Spring reproductive success influences autumnal malarial load in a passerine bird

Romain PIGEAULT[1,2]*, Camille-Sophie COZZAROLO[2,3]*, Jérôme WASSEF[2], Jérémy GREMION[2], Marc BASTARDOT[2], Olivier GLAIZOT[2,4] & Philippe CHRISTE[2]

1 Laboratoire EBI, Equipe EES, UMR CNRS 7267, 86000 Poitiers, France

2 Department of Ecology and Evolution, Université de Lausanne, Biophore, 1015 Lausanne, Switzerland

3 Biogéosciences, UMR 6282 CNRS, université de Bourgogne, 6 boulevard Gabriel, 21000 Dijon, France

4 Muséum cantonal des sciences naturelles - Département de zoologie, Palais de Rumine, Place de la Riponne 6, 1005 Lausanne, Switzerland

* Co-first

Corresponding author : romain.pigeault@univ-poitiers.fr

romain.pigeault@univ-poitiers.fr - 0000-0002-8011-4600

camille-sophie.cozzarolo@ik.me - 0000-0002-9056-8622

jerome.wassef@gmail.com

jeremy.gremion@vogelwarte.ch

marcbastardot@hotmail.com

olivier.glaizot@unil.ch - 0000-0001-9116-3355

philippe.christe@unil.ch - 0000-0002-8605-7002

Abstract

Although avian haemosporidian parasites are widely used as model organisms to study
fundamental questions in evolutionary and behavorial ecology of host-parasite interactions, some of
their basic characteristics, such as seasonal variations in within-host density, are still mostly unknown.
In addition, their interplay with host reproductive success in the wild seems to depend on the
interaction of many factors, starting with host and parasite species and the temporal scale under
study. Here, we monitored the intensities of infection by two haemosporidian parasites – *Plasmodium*
relictum and *P. homonucleophilum* – in two wild populations of great tits (*Parus major*) in Switzerland
over three years, to characterize their dynamics. We also collected data on birds' reproductive output
– laying date, clutch size, fledging success – to determine whether they were associated with infection
intensity before (winter), during (spring) and after (autumn) breeding season. Both parasite species
dramatically increased their within-host density in spring, in a way that was correlated to their
parasitaemia in winter. Infection intensity before and during breeding season did not explain
reproductive success. However, the birds which fledged the more chicks had higher parasite burdens
in autumn, which were not associated with their parasite burden in previous spring. Our results tend
to indicate that high haemosporidian parasite loads do not impair reproduction in great tits, but high
resource allocation into reproduction can leave birds less able to maintain low parasitaemia over the
following months.

Keywords: avian malaria, annual variations, relapses, recurences, parasitaemia, life history traits, bird

Introduction

The assumed impact of parasitic infections on animal fitness is at the basis of several evolutionary

theories such as the Hamilton-Zuk (Hamilton & Zuk, 1982) or the terminal investment (Stearns, 1989)

hypotheses. A recent meta-analysis highlighted the overall negative cost of parasites on reproductive

success (Hasik & Siepielski, 2022). This meta-analysis focussed on the infection status as a binary

variable (parasitized versus non-parasitized), as is the case of many studies. However, the parasite

load might better represent the host's ability to control the infection (resistance) and might be a finer

correlate of its physiological or energetical costs (Stjernman et al., 2008; Risely et al., 2018; Sánchez

et al., 2018; Methling et al., 2019). A negative association between infection intensity and

reproductive success was shown in several host-parasite systems (Madsen et al., 2005; Asghar et al.,

2011; Gooderham & Schulte-Hostedde, 2011; Hicks et al., 2019; Schoepf et al., 2022) but some studies

also show an absence or a positive correlation (Siikamäki et al., 1997; Edler et al., 2004; Raveh et al.,

2011; Kulma et al., 2014; Delefortrie et al., 2022).

The different results obtained from studies sometimes involving similar host-parasite pairs could

be explained by a number of factors. For instance, the results can vary with the host's sex. Several

studies have reported a negative relationship between parasite burden and reproductive success only

in males (Sundberg, 1995; Dawson & Bortolotti, 2001; Gooderham & Schulte-Hostedde, 2011; but see

Hicks et al., 2019). The age of the infected individuals could also partly blur the signal, since it has been

shown that in some species, individuals nearing the end of their lives may invest heavily in

reproduction (i.e. terminal investment hypothesis, Velando et al., 2006; Duffield et al., 2018), and in

others, that the age-reproductive success relationship follows a bell-shaped curve (Lecomte et al.,

2010; Saraux & Chiaradia, 2022). The fluctuation of parasite burdens over time may also pose a

significant challenge. A large diversity of parasites, ranging from viruses to eukaryotic organisms,

exhibit highly dynamic patterns of replication rate, leading to temporal fluctuations of within-host

parasite burden (Hasker et al., 2013; Pigeault, Caudron, et al., 2018; Colangeli et al., 2020). These

fluctuations can occur on short-term scales, such as daily variations, as well as on long-term scales,

spanning months or even years (Martinez-Bakker & Helm, 2015; Prior et al., 2020). Although the

parasite density experienced by the host during the breeding period is most likely to have a direct (e.g.

pathogenicity, resource exploitation) influence on host's reproductive success (e.g. Madsen et al.,

2005; Asghar et al., 2011; Gooderham & Schulte-Hostedde, 2011; Hicks et al., 2019; Schoepf et al.,

2022), the within-host infection intensity before the breeding season may also have indirect effects,

by influencing for instance premating trade-offs in resource allocation (i.e. carry over effect, Harrison

et al., 2011). On the other hand, as life-history theory assumes that components of reproductive effort

are costly, investment in reproduction may have longer-term consequences for the host's ability to
clear or at least to control parasite replication rate (Nordling et al., 1998; Leivesley et al., 2019).

In this study, we used a longitudinal approach to assess whether the intensity of parasite infection,
quantified a few months before and during the breeding season, could explain host reproductive
output, and whether reproductive effort could predict infection intensity later in the year. To do this,
we used avian malaria as a biological system. For over a century, this vector borne disease has been
used in studies of host-parasite interactions (Rivero & Gandon, 2018) and provides an excellent model
for exploring the influence of parasitic infections on host life-history traits (e.g. Oppliger et al., 1997;
Asghar et al., 2015; Pigeault, Cozzarolo, et al., 2018). To date, the vast majority of studies that
investigated the impact of malaria infection on bird reproduction used the infection status (i.e.
parasitized versus non-parasitized) as a predictor of diverse reproductive parameters (e.g. Sanz et al.,
2001; Norte et al., 2009; Podmokła et al., 2014; Zylberberg et al., 2015; Pigeault, Cozzarolo, et al.,
2018). The few non-interventional studies that have examined the influence of parasite load on the
reproductive success of birds have reported contrasting results (e.g., Siikamäki et al., 1997; Edler et
al., 2004; Asghar et al., 2011; Kulma et al., 2014). On the other hand, drug-induced reduction of
parasitaemia usually results in higher reproductive success (e.g., Merino et al., 2000; Marzal et al.,
2005; Knowles et al., 2010; Schoepf et al., 2022). A possible explanation for the difference in results
between the non-interventional studies and those in which the parasite load was experimentally
reduced is that only experimental reduction of the parasite load can ensure that treated birds have a
permanently lower parasitaemia than control individuals. Indeed, avian malaria is characterized by
radical temporal variations in the intensity of parasite infection in the peripheral blood of the bird.
After transmission of the parasite by a mosquito, the number of parasites in the blood compartment
increases rapidly and reaches a maximum in 10-20 days (Pigeault, Caudron, et al., 2018; Palinauskas
et al., 2018). Activation of the host immune system then controls avian malaria parasite replication
rate, but in most cases it is not able to eliminate it completely, leading to the establishment of the
chronic phase of the infection (Valkiūnas, 2005; Asghar et al., 2012) during which parasites persist at
low densities for several months or years. However, this chronic phase can be regularly interrupted
by recrudescence events when the parasite load in the host's peripheral blood increases significantly
over a short period of time (three to seven days, Cornet et al., 2014; Pigeault et al., 2023).
Consequently, the results of non-interventional studies investigating the relationship between the
intensity of avian malaria parasite infection and host reproductive success will be highly dependent
on the timing of parasite load measurements.

Our non-interventional study was carried out over a period of three years, during which field
sessions were organized to capture and recapture great tits in order to (i) monitor annual infection

dynamics, (ii) study the link between infection intensity and reproductive success, and (iii) investigate the influence of investment in reproduction on parasite load measured later in the year. Although commonly mentioned in the literature, but rarely reported, we predict a significant increase in parasite load in spring, illustrating spring reccurences (or spring relapses, Applegate, 1971). In light of the studies which showed a trade-off between activation of the great tit's immune system and their reproduction (e.g., Ots & Hõrak, 1998; Grzędzicka, 2017; Kubacka & Cichoń, 2020), we predict a negative relationship between infection intensity and reproductive success. Finally, in view of the energy expenditure associated with reproduction (Visser & Lessells, 2001; Nilsson & Råberg, 2001), we also predict that infected individuals who invest substantially in reproduction will later experience higher parasite loads (Hanssen et al., 2005).

Material and Methods

Study area and host species

The study was carried out between 2017 and 2019 in two populations of great tits (*Parus major*) breeding in nest-boxes located near the University of Lausanne (Dorigny Forest) and 15 km apart in the Marais des Monod, both in Switzerland (Pigeault, Cozzarolo, et al., 2018). At each breeding season, we trapped adults in the nest boxes to determine whether they were infected by haemosporidian parasites and, if so, we molecularly identified the parasites involved in the infection (see Appendix 1). We also took morphometric measurements (mass and tarsus length) to calculate the scaled mass index of each individual as a proxy for body condition (SMI, Peig & Green, 2009) and we collected several breeding parameters (i.e., laying date, clutch size and fledging success). To correct for between-year differences in average breeding time, laying date was standardized by using the first year of the study as a reference (2017, 1 April = day 1). The differences in laying date between 2017 and each subsequent year were subtracted from the actual laying date of that particular year (Allander & Bennett, 1995).

In addition to the spring monitoring, birds were also captured with mist nets during winter (between January and end of March) and autumn (between mid-October and mid-December). Three capture sessions were done in winter and in autumn in the Dorigny Forest, while for logistic reasons it was only possible to perform two sessions per season in the Marais des Monod. We also took blood samples and morphometric measurements on all caught individuals. No capture session was carried out in summer because we know from previous attempts that it is very difficult to capture great tits during this period in our study area.

Molecular analyses

Only birds captured at least twice in a focal year and diagnosed as infected by *Plasmodium* parasite

on at least one of their capture dates were retained for analyses (see **Appendix 1, section 1**). The

quantification of infection intensity in all the blood samples collected was carried out using qPCR (see

Appendix 1, section 2). Parasite load was calculated with relative quantification values (RQ). RQ can

be interpreted as the fold-amount of target gene (*Plasmodium* 18S rDNA) with respect to the amount

of the reference gene (avian 18S rDNA) and is calculated as $2^{-(Ct\_\textit{Plasmodium} - Ct\_bird)}$. For convenience, RQ

values were standardized by $\times 10^4$ factor. Since some individuals were trapped and blood sampled

several times per season, we calculated the average parasitemia of individuals for each season when

we wanted to compare the infection intensity between seasons. To note, in this manuscript, we

indiscriminately use the words "parasite load", "parasite burden", "density", "infection intensity" and

"parasitaemia" to refer to the relative quantity of *Plasmodium* sp. in birds' blood. When using these

words to discuss other studies, they might refer to other measurements of parasite quantities in hosts

or hosts' tissues.

Statistical analysis

We used a mixed model procedure with a normal error structure and bird fitted as a random factor

to test for an effect of month of capture on change in infection intensity by Plasmodium parasite

('lme4' package, Bates et al., 2014). Sex, age, Plasmodium lineage and year were added as fixed factors

into the model. As infection intensity could be expected to be a non-linear function of month of

capture, because of spring recurrences, the quadratic term month of capture was added to assess if it

significantly improved the model fit. In order to study the influence of the infection intensity measured

in winter and spring on the reproductive parameters and the impact of reproductive parameters on

the infection intensity measured in autumn, we calculated the average parasitemia of each bird in

each season. Indeed, for the same year a certain number of individuals were captured several times

in winter or in autumn. Then, for a focal year, we used generalized linear models (glm), with an error

structure appropriate to each response variable (see **Appendix 1, Table S1**), to study the impact of

winter and spring parasitemia on bird's reproductive parameters. We also used a glm to measure the

influence of the birds' reproductive parameters on their parasitemia measured in the autumn. Finally,

we used a glm to study the impact of spring recurrence intensity on the set of reproductive

parameters. Spring recurrence intensity was defined by the difference in infection intensity between

winter and spring. Sex, age, *Plasmodium* lineage, SMI and year were added as explanatory variables

into the models. All statistical analyses were carried out using the R statistical software (v. 4.1.3). The

raw data and the R script used for the analyses and to produce the figures are available on the figshare

website (10.6084/m9.figshare.23695422).

 Ethics statement

     This study was approved by the Ethical Committee of the Vaud Canton veterinary authority

 (authorization number is 1730.4).

Results

Variation of host parasite load

We followed the annual infection dynamics in 70 individuals (31 females and 39 males). We

detected five lineages of *Plasmodium* parasites (Parus1: 1, BT7: 2, Turdus1: 4, SW2: 20, SGS1: 43) but

we focused our subsequent analyses on the two most prevalent lineages (i.e., *P. homonucleophilum*

SW2 and *P. relictum* SGS1). The parasite communities were very different in both sites. The SW2

lineage was only observed in the Marais des Monod while hosts infected by SGS1 were only captured

in the Dorigny forest.

A significant influence of month of capture on the infection intensity was observed but only when

month was added as a quadratic term in the model (model 1, month: $\chi2 = 62.17$, $p < 0.0001$, month²:

$\chi2 = 63.205$, $p < 0.0001$, Figure 1). We indeed observed that the parasite load followed a bell-shaped

function: peaking during spring and decreasing thereafter (**Figure 1**). When spring captures were

removed from our analyses, we no longer observed an influence of month on host parasite load

(model 2, month: $\chi2 = 1.21$, $p = 0.271$, month²: $\chi2 = 1.09$, $p = 0.315$). Although the general shape of

infection for both lineages are similar, we observed that host infected by SW2 showed a higher

parasite load than those infected by SGS1 (model 1, $\chi2 = 31.69$, $p < 0.0001$, mean RQ10 ± se, SW2 =

294 ± 99, SGS1 = 148 ±87, Figure 1). Infection intensity differed between years (model 1, $\chi2 = 9.44$, p

= 0.002, mean RQ10 ± se, 2017 = 174 ± 69, 2018 = 83 ± 31, 2019 = 247 ± 145) but we did not observe

any difference between host sex and age (model 1, sex: $\chi2 = 1.41$, $p = 0.235$, age: $\chi2 = 0.002$, $p = 0.967$).

After calculating the average parasite load of hosts in each season, we observed a significant

relationship between winter and spring parasite loads (model 3, $\chi2 = 6.16$, p = 0.010). The most

infected hosts in winter exhibited higher parasite loads in spring (**Figure 2**). Conversely, we did not

observe any relationship between winter and autumn parasite loads or between spring and autumn

(model 4-5, $\chi2 = 0.62$, p = 0.445, $\chi2 = 0.04$, p = 0.837, respectively, **Figure 2**). As observed previously

in the analysis of the annual dynamics of *Plasmodium* infection, in each season the infection intensity

was higher in individuals infected by SW2 than in birds infected by SGS1 (model 3: $\chi2 = 8.40$, p = 0.006,

model 4: $\chi2 = 4.77$, p = 0.049, model 5: $\chi2 = 6.55$, p = 0.016).

Figure 2 – Relationship between parasite loads measured at different season. (A) Relationship
between spring and winter parasite load (Log(RQ10+1)), (B) between autumn and winter parasite
load and (C) between autumn and spring parasite load. Blue dots correspond to individuals infected
with *P. relictum SGS1*, green dots correspond to individuals infected with *P. homonucleophilum* SW2.

Relationship between host parasite load and reproductive parameters

The parasite load recorded during the winter period did not influence the laying date (model 6, $\chi2$

= 0.244, p = 0.623), the investment in the reproduction or the reproductive success of the hosts (model

7, clutch size: $\chi2 = 0.103$, p = 0.748, model 8, number of fledged chicks: $\chi2 = 0.479$, p = 0.489). We also

did not observe any relationship between parasite load measured during the reproductive period and

any reproductive parameters (model 9-11, laying date: χ^2 = 0.037, p = 0.847, clutch size: χ^2 = 0.141, p = 0.707, number of fledged chicks: χ^2 = 0.001, p = 0.975). However, the number of fledged chicks was higher in hosts infected by SW2 than in birds infected by SGS1 (model 11, χ^2 = 7.60, p = 0.006, mean ± se, SW2 = 6.22 ± 0.46, SGS1 = 4.20 ± 0.32). Finally, the clutch size did not impact the parasite load measured later in autumn (model 12, χ^2 = 1.340, p = 0.247, **Figure 3A**), but we observed a significant effect of the number of fledged chicks (model 12, χ^2 = 5.91, p = 0.015). Individuals with high fledging success also trended toward higher parasite loads next autumn (**Figure 3B**). We observed a negative relationship between individual SMI and autumn parasite load (model 12, χ^2 = 4.66, p = 0.031). The birds with the lowest body condition were those that had the highest parasite burden (**Figure S1**). However, there was no relationship between the breeding success (i.e. the number of fledged chicks) of birds in spring and their body condition a few months later in autumn (model 13, χ^2 = 0.55, p = 0.457).

Figure 3 – Influence of reproductive parameters monitored in spring and host's parasite load measured a few months later in autumn. Relationship between (**A**) the number of eggs laid or (**B**) the number of chicks fledged and bird's parasite load (Log RQ10+1). Blue dots correspond to individuals infected with *P. relictum SGS1*, green dots correspond to individuals infected with *P. homonucleophilum* SW2.

Discussion

Using data collected from two wild populations of great tits infected by two different malaria parasite lineages, we highlighted that parasite load increased drastically between winter and spring,

and we found that increased reproductive allocation in spring is associated with higher *Plasmodium*
parasite load in autumn.

Numerous studies have attempted to characterize the yearly fluctuations of infection prevalence
in different bird populations. They either found no seasonality in probability of infection (Himalayan
bird community; Ishtiaq et al., 2017), a link with migration (Hellgren et al., 2013; Pulgarín-R et al.,
2019), or seasonality patterns that vary across *Plasmodium* lineages (Neto et al., 2020) and
geographical regions (Cosgrove et al., 2008; Lynton-Jenkins et al., 2020; Neto et al., 2020). On the
other hand, little is known about annual within-host variation in infection intensity. In a wild bird
community in Slovakian woodlands, *Plasmodium* sp. parasitaemia peaked in summer (Šujanová et al.,
2021) while *Plasmodium relictum* parasitaemia in captive house sparrows (*Passer domesticus*) in Spain
varied monthly without clear seasonal pattern (Garcia-Longoria et al., 2022). In our study we reported
that for two *Plasmodium* lineages the parasitaemia followed a bell-shaped function. The number of
infected erythrocytes increased more than tenfold between winter and spring, and then decreased in
autumn to winter levels. Our study illustrates a phenomenon described by Applegate & Beaudoin in
the 70s but has been rarely documented since: the spring recurrences (or spring relapses Applegate,
1970). Spring recurrences can be due to several factors likely acting on bird's physiology or immunity,
such as resource availability (Cornet, Bichet, et al., 2014), environmental stressors (Becker et al., 2020;
Pigeault et al., 2023), immune challenges, co-infections with other parasites (Palinauskas et al., 2011;
Reinoso-Pérez et al., 2020; Garcia-Longoria et al., 2022) and energy allocation to other functions such
as reproduction (Christe et al., 2012).

Although various factors may be responsible for triggering recurrences of infection, we have
demonstrated that the spring parasitaemia of birds was positively correlated to their parasite load
recorded in winter. However, contrary to our predictions we did not observe any effect of both winter
and spring parasite burden on bird reproductive parameters. Indeed, individuals with extremely high
parasite burdens (RQ > 1000) did not lay more eggs nor fledged more chicks than those with very low
parasite burdens (RQ < 1). This result is not consistent with a premating trade-offs in resource
allocation (i.e., carry-over effect) or with a reallocation of host resources towards immunity during the
mating period (Stearns, 1989; Harrison et al., 2011; Stahlschmidt et al., 2013; Albery et al., 2020).

Interestingly, while the infection intensity in winter is a significant predictor of the intensity of the
spring recurrence, we did not observe any relationship between the parasite load measured in winter
or in spring and the infection intensity recorded the following autumn. This result suggests that
between spring and autumn, biotic and/or abiotic parameters may have modified the interactions
between *Plasmodium* and its host. Infections by new parasites during spring or summer periods could
directly or indirectly influence the within-host infection dynamics of *Plasmodium* (Cellier-Holzem et

al., 2010). Our study was not designed to test this hypothesis, but we noted that the birds recaptured
in the autumn were all infected with the same lineage of *Plasmodium* that was identified in the spring.
Further, we did not record any new haemosporidian parasite infection. However, we cannot exclude
the possibility that, during the breeding season, the birds were infected by other parasites (e.g.
gastrointestinal nematodes, viruses, Leucocytozoon).

Inter-individual variability in reproductive investment may also explain why there was no
relationship between the parasite load measured at the beginning of the reproductive period and that
measured several months later in autumn. Because immunity and reproduction compete for host
resources, in resource-limited environments, hosts that reproduce should have fewer resources to
allocate to immune defense which may ultimately influence within-host infection dynamics. Increased
allocation to reproduction was found to be associated with increased burden of gastrointestinal
nematodes in wild Soay sheep during both late gestation and early lactation (Leivesley et al., 2019).
Brood manipulation studies on birds showed that increased allocation to reproduction was associated
with greater parasite burdens and less effective immune responses at the end of the breeding period
(Richner et al., 1995, Oppliger et al., 1997, Knowles et al., 2009; Christe et al., 2012). Here, we observed
that birds with the highest reproductive success tended to be those with the highest parasite load 6-
8 months later. Although supported by a small sample size (n = 36), this result suggests a long-term
effect of investment in reproduction on the ability of hosts to control the replication rate of blood
parasites.

Finally, in addition to seasonal variations, we observed a significant difference in the intensity of
infection in birds, depending on the *Plasmodium* lineage involved in the infection. Hosts infected with
P. homonucleophilum had a higher parasite load than hosts infected with *P. relictum*, irrespective of
the time of year when the blood samples were taken. This result is consistent with that of an earlier
study conducted on the same great tits populations in 2009-2011 (Rooyen et al., 2013). Interestingly,
we also found that birds infected by *P. homonucleophilum* showed higher reproductive success than
those infected by *P. relictum.* However, it is not possible to determine whether the differences
observed between hosts infected by these two *Plasmodium* species are the result of differences
inherent to the two blood parasites, or an effect of the vertebrate host genetic background, the living
environment or interactions between all these factors. Indeed, the individuals infected with *P.*
homonucleophilum all originated from the Monod marshlands, whereas the birds infected with *P.*
relictum came from the Dorigny forest. Although these two populations are only 15km apart, we have
never observed birds migrating between the two populations and the habitats are very different (large
marshy forest massif *versus* periurban forest patches on the campus of the University of Lausanne,
respectively).

In conclusion, our monitoring of great tit malarial load across seasons evidenced a spring

recurrence, the triggers of which are still to be determined. Further, our study supported the idea that

strong allocation in reproduction incurs costs later in life, without evidence that higher parasite

burdens prior to and during breeding season reduce reproductive success.

Appendices

317 Appendix 1 includes:
318 Section 1: Detection of hemosporidian parasites
319 Section 2: Quantification of infection intensity
320 Table S1: Description of the statistical models presented in the main text.
321 Figure S1: Relationship between scale mass index and intensity of infection in birds captured in autumn.

Funding

323 This project was funded by the Swiss National Science Foundation (SNSF), grants 31003A_179378 to PC.

Conflict of interest disclosure

325 The authors declare that they comply with the PCI rule of having no conflicts of interest in relation to the
326 content of the article.

Data, scripts, code, and supplementary information availability

328 Data and scripts are available online: https://doi.org/10.6084/m9.figshare.23695422

References

Albery GF, Watt KA, Keith R, Morris S, Morris A, Kenyon F, Nussey DH, Pemberton JM (2020)

Reproduction has different costs for immunity and parasitism in a wild mammal. *Functional*

Ecology, **34**, 229–239. https://doi.org/10.1111/1365-2435.13475

Allander K, Bennett GF (1995) Retardation of breeding onset in great tits (*Parus major*) by blood

parasites. *Functional Ecology*, **9**, 677. https://doi.org/10.2307/2390160

Applegate JE (1970) Population changes in latent avian malaria infections associated with season and

corticosterone treatment. *The Journal of Parasitology*, **56**, 439. https://doi.org/10.2307/3277599

Applegate JE (1971) Spring relapse of *Plasmodium relictum* infections in an experimental field

population of English sparrows (*Passer domesticus*). *Journal of Wildlife Diseases*, **7**, 37–42.

https://doi.org/10.7589/0090-3558-7.1.37

Applegate JE, Beaudoin RL (1970) Mechanism of spring relapse in avian malaria: Effect of gonadotropin
and corticosterone. *Journal of Wildlife Diseases*, **6**, 443–447. https://doi.org/10.7589/0090-3558-
6.4.443

Asghar M, Hasselquist D, Bensch S (2011) Are chronic avian haemosporidian infections costly in wild
birds? *Journal of Avian Biology*, **42**, 530–537. https://doi.org/10.1111/j.1600-048X.2011.05281.x

Asghar M, Hasselquist D, Hansson B, Zehtindjiev P, Westerdahl H, Bensch S (2015) Hidden costs of
infection: Chronic malaria accelerates telomere degradation and senescence in wild birds. *Science*,
347, 436–438. https://doi.org/10.1126/science.1261121

Asghar M, Westerdahl H, Zehtindjiev P, Ilieva M, Hasselquist D, Bensch S (2012) Primary peak and
chronic malaria infection levels are correlated in experimentally infected great reed warblers.
Parasitology, **139**, 1246–1252. https://doi.org/10.1017/S0031182012000510

Bates D, Machler M, Bolker B, Walker S (2015). Fitting linear mixed-effects models using lme4. J. Statis.
Soft. **67** (1), 1–48. https://doi.org/10.18637/jss.v067.i01

Becker DJ, Singh D, Pan Q, Montoure JD, Talbott KM, Wanamaker SM, Ketterson ED (2020) Artificial
light at night amplifies seasonal relapse of haemosporidian parasites in a widespread songbird.
Proceedings of the Royal Society B: Biological Sciences, **287**, 20201831.
https://doi.org/10.1098/rspb.2020.1831

Cellier-Holzem E, Esparza-Salas R, Garnier S, Sorci G (2010) Effect of repeated exposure to *Plasmodium
relictum* (lineage SGS1) on infection dynamics in domestic canaries. *International Journal for
Parasitology*, **40**, 1447–1453. https://doi.org/10.1016/j.ijpara.2010.04.014

Christe P, Glaizot O, Strepparava N, Devevey G, Fumagalli L (2012) Twofold cost of reproduction: An
increase in parental effort leads to higher malarial parasitaemia and to a decrease in resistance to
oxidative stress. *Proceedings of the Royal Society B: Biological Sciences*, **279**, 1142–1149.
https://doi.org/10.1098/rspb.2011.1546

Colangeli R, Gupta A, Vinhas SA, Chippada Venkata UD, Kim S, Grady C, Jones-López EC, Soteropoulos
P, Palaci M, Marques-Rodrigues P, Salgame P, Ellner JJ, Dietze R, Alland D (2020) *Mycobacterium
tuberculosis* progresses through two phases of latent infection in humans. *Nature
Communications*, **11**, 4870. https://doi.org/10.1038/s41467-020-18699-9

Cornet S, Bichet C, Larcombe S, Faivre B, Sorci G (2014) Impact of host nutritional status on infection
dynamics and parasite virulence in a bird-malaria system. *Journal of Animal Ecology*, **83**, 256–265.
https://doi.org/10.1111/1365-2656.12113

Cornet S, Nicot A, Rivero A, Gandon S (2014) Evolution of plastic transmission strategies in avian
malaria. *PLOS Pathogens*, **10**, e1004308. https://doi.org/10.1371/journal.ppat.1004308

Cosgrove CL, Wood MJ, Day KP, Sheldon BC (2008) Seasonal variation in *Plasmodium* prevalence in a population of blue tits *Cyanistes caeruleus*. *Journal of Animal Ecology*, **77**, 540–548. https://doi.org/10.1111/j.1365-2656.2008.01370.x

Dawson R, Bortolotti G (2001) Sex-specific associations between reproductive output and hematozoan parasites of American kestrels. *Oecologia*, **126**, 193–200. https://doi.org/10.1007/s004420000506

Delefortrie Z, Gante HF, Gordo O, Schwab KR, Gonser RA (2022) Effects of haemosporidian co-infection and parasitemia on reproductive strategies in a polymorphic species. *bioRxiv*, 2022.03.28.486032. https://doi.org/10.1101/2022.03.28.486032

Duffield KR, Hampton KJ, Houslay TM, Hunt J, Rapkin J, Sakaluk SK, Sadd BM (2018) Age-dependent variation in the terminal investment threshold in male crickets. *Evolution*, **72**, 578–589. https://doi.org/10.1111/evo.13443

Edler R, Klump GM, Friedl TWP (2004) Do blood parasites affect reproductive performance in male red bishops (*Euplectes orix*)? A test of the Hamilton-Zuk hypothesis. *Ethology Ecology and Evolution*, **16**, 315–328. https://doi.org/10.1080/08927014.2004.9522623

Garcia-Longoria L, Magallanes S, Huang X, Drews A, Råberg L, Marzal A, Bensch S, Westerdahl H (2022) Reciprocal positive effects on parasitemia between coinfecting haemosporidian parasites in house sparrows. *BMC Ecology and Evolution*, **22**, 73. https://doi.org/10.1186/s12862-022-02026-5

Gooderham K, Schulte-Hostedde A (2011) Macroparasitism influences reproductive success in red squirrels (*Tamiasciurus hudsonicus*). *Behavioral Ecology*, **22**, 1195–1200. https://doi.org/10.1093/beheco/arr112

Grzędzicka E (2017) Immune challenge of female great tits at nests affects provisioning and body conditions of their offspring. *Acta Ethologica*, **20**, 223–233. https://doi.org/10.1007/s10211-017-0265-4

Hamilton WD, Zuk M (1982) Heritable true fitness and bright birds: A role for parasites? *Science, New Series*, **218**, 384–387.

Hanssen SA, Hasselquist D, Folstad I, Erikstad KE (2005) Cost of reproduction in a long-lived bird: incubation effort reduces immune function and future reproduction. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1039–1046. https://doi.org/10.1098/rspb.2005.3057

Harrison XA, Blount JD, Inger R, Norris DR, Bearhop S (2011) Carry-over effects as drivers of fitness differences in animals: Carry-over effects in animal populations. *Journal of Animal Ecology*, **80**, 4–18. https://doi.org/10.1111/j.1365-2656.2010.01740.x

Hasik AZ, Siepielski AM (2022) Parasitism shapes selection by drastically reducing host fitness and increasing host fitness variation. *Biology letters*, **18**, 20220323. https://doi.org/10.1098/rsbl.2022.0323

Hasker E, Kansal S, Malaviya P, Gidwani K, Picado A, Singh RP, Chourasia A, Singh AK, Shankar R,

Menten J, Wilson ME, Boelaert M, Sundar S (2013) Latent Infection with *Leishmania donovani* in

Highly Endemic Villages in Bihar, India. *PLOS Neglected Tropical Diseases*, **7**, e2053.

https://doi.org/10.1371/journal.pntd.0002053

Hellgren O, Wood MJ, Waldenström J, Hasselquist D, Ottosson U, Stervander M, Bensch S (2013)

Circannual variation in blood parasitism in a sub-Saharan migrant passerine bird, the garden

warbler. *Journal of Evolutionary Biology*, **26**, 1047–1059. https://doi.org/10.1111/jeb.12129

Hicks O, Green JA, Daunt F, Cunningham EJA, Newell M, Butler A, Burthe SJ (2019) Sublethal effects of

natural parasitism act through maternal, but not paternal, reproductive success in a wild

population. *Ecology*, **100**, e02772. https://doi.org/10.1002/ecy.2772

Ishtiaq F, Bowden CGR, Jhala YV (2017) Seasonal dynamics in mosquito abundance and temperature

do not influence avian malaria prevalence in the Himalayan foothills. *Ecology and Evolution*, **7**,

8040–8057. https://doi.org/10.1002/ece3.3319

Knowles SCL, Nakagawa S, Sheldon BC (2009) Elevated reproductive effort increases blood

parasitaemia and decreases immune function in birds: A meta-regression approach. *Functional

Ecology*, **23**, 405–415. https://doi.org/10.1111/j.1365-2435.2008.01507.x

Knowles SCL, Palinauskas V, Sheldon BC (2010) Chronic malaria infections increase family inequalities

and reduce parental fitness: Experimental evidence from a wild bird population. *Journal of

Evolutionary Biology*, **23**, 557–569. https://doi.org/10.1111/j.1420-9101.2009.01920.x

Kubacka J, Cichoń M (2020) An immune challenge of female great tits decreases offspring survival and

has sex-specific effects on offspring body size. *acta ethologica*, **23**, 173–181.

https://doi.org/10.1007/s10211-020-00351-w

Kulma K, Low M, Bensch S, Qvarnström A (2014) Malaria-infected female collared flycatchers (Ficedula

albicollis) do not pay the cost of late breeding. *PLoS ONE*, **9**.

https://doi.org/10.1371/journal.pone.0085822

Lecomte VJ, Sorci G, Cornet S, Jaeger A, Faivre B, Arnoux E, Gaillard M, Trouvé C, Besson D, Chastel O,

Weimerskirch H (2010) Patterns of aging in the long-lived wandering albatross. *Proceedings of the

National Academy of Sciences*, **107**, 6370–6375. https://doi.org/10.1073/pnas.0911181107

Leivesley JA, Bussière LF, Pemberton JM, Pilkington JG, Wilson K, Hayward AD (2019) Survival costs of

reproduction are mediated by parasite infection in wild Soay sheep. *Ecology Letters*, **22**, 1203–

1213. https://doi.org/10.1111/ele.13275

Lynton-Jenkins JG, Bründl AC, Cauchoix M, Lejeune LA, Sallé L, Thiney AC, Russell AF, Chaine AS,

Bonneaud C (2020) Contrasting the seasonal and elevational prevalence of generalist avian

haemosporidia in co-occurring host species. *Ecology and Evolution*, **10**, 6097–6111.
https://doi.org/10.1002/ece3.6355

Madsen T, Ujvari B, Olsson M (2005) Old pythons stay fit; effects of haematozoan infections on life
history traits of a large tropical predator. *OECOLOGIA*, **142**, 407–412.
https://doi.org/10.1007/s00442-004-1742-9

Martinez-Bakker M, Helm B (2015) The influence of biological rhythms on host–parasite interactions.
Trends in Ecology & Evolution, **30**, 314–326. https://doi.org/10.1016/j.tree.2015.03.012

Marzal A, Lope FD, Navarro C, Møller AP (2005) Malarial parasites decrease reproductive success: An
experimental study in a passerine bird. *Oecologia*, **142**, 541–545. https://doi.org/10.1007/s00442-
004-1757-2

Merino S, Moreno J, Sanz JJ, Arriero E (2000) Are avian blood parasites pathogenic in the wild? A
medication experiment in blue tits (Parus caeruleus). *Proceedings of the Royal Society B: Biological*
Sciences, **267**, 2507–2510. https://doi.org/10.1098/rspb.2000.1312

Methling C, Douda K, Reichard M (2019) Intensity-dependent energetic costs in a reciprocal parasitic
relationship. *Oecologia*, **191**, 285–294. https://doi.org/10.1007/s00442-019-04504-y

Neto JM, Mellinger S, Halupka L, Marzal A, Zehtindjiev P, Westerdahl H (2020) Seasonal dynamics of
haemosporidian (Apicomplexa, Haemosporida) parasites in house sparrows *Passer domesticus* at
four European sites: comparison between lineages and the importance of screening methods.
International Journal for Parasitology, **50**, 523–532. https://doi.org/10.1016/j.ijpara.2020.03.008

Nilsson J-Å, Råberg L (2001) The resting metabolic cost of egg laying and nestling feeding in great tits.
Oecologia, **128**, 187–192. https://doi.org/10.1007/s004420100653

Nordling D, Andersson M, Zohari S, Lars G (1998) Reproductive effort reduces specific immune
response and parasite resistance. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences, **265**, 1291–1298. https://doi.org/10.1098/rspb.1998.0432

Norte AC, Araújo PM, Sampaio HL, Sousa JP, Ramos JA (2009) Haematozoa infections in a Great Tit
Parus major population in Central Portugal: relationships with breeding effort and health:
Haematozoa infections in Great Tits from Central Portugal. *Ibis*, **151**, 677–688.
https://doi.org/10.1111/j.1474-919X.2009.00960.x

Oppliger A, Christe P, Richner H (1997) Clutch size and malarial parasites in female great tits.
Behavioral Ecology, **8**, 148–152.

Ots I, Hõrak P (1998) Health impact of blood parasites in breeding great tits. *Oecologia*, **116**, 441–448.
https://doi.org/10.1007/s004420050608

Palinauskas V, Valkiūnas G, Bolshakov CV, Bensch S (2011) *Plasmodium relictum* (lineage SGS1) and
Plasmodium ashfordi (lineage GRW2): The effects of the co-infection on experimentally infected

passerine birds. *Experimental Parasitology*, **127**, 527–533. https://doi.org/10.1016/j.exppara.2010.10.007

Palinauskas V, Žiegytė R, Šengaut J, Bernotienė R (2018) Different paths – the same virulence: experimental study on avian single and co-infections with Plasmodium relictum and Plasmodium elongatum. *International Journal for Parasitology*, **48**, 1089–1096. https://doi.org/10.1016/j.ijpara.2018.08.003

Peig J, Green AJ (2009) New perspectives for estimating body condition from mass/length data: The scaled mass index as an alternative method. *Oikos*, **118**, 1883–1891. https://doi.org/10.1111/j.1600-0706.2009.17643.x

Pigeault R, Caudron Q, Nicot A, Rivero A, Gandon S (2018) Timing malaria transmission with mosquito fluctuations. *Evolution Letters*, **2**, 378–389. https://doi.org/10.1002/evl3.61

Pigeault R, Cozzarolo C-S, Choquet R, Strehler M, Jenkins T, Delhaye J, Bovet L, Wassef J, Glaizot O, Christe P (2018) Haemosporidian infection and co-infection affect host survival and reproduction in wild populations of great tits. *International Journal for Parasitology*, **48**, 1079–1087. https://doi.org/10.1016/j.ijpara.2018.06.007

Pigeault R, Ruiz De Paz A, Baur M, Isaïa J, Glaizot O, Christe P (2023) Impact of host stress on the replication rate of *Plasmodium*: take it easy to avoid malaria recurrences. *Frontiers in Ecology and Evolution*, **11**.

Podmokła E, Dubiec A, Drobniak SM, Arct A, Gustafsson L, Cichoń M (2014) Avian malaria is associated with increased reproductive investment in the blue tit. *Journal of Avian Biology*, **45**, 219–224. https://doi.org/10.1111/j.1600-048X.2013.00284.x

Prior KF, Rijo-Ferreira F, Assis PA, Hirako IC, Weaver DR, Gazzinelli RT, Reece SE (2020) Periodic parasites and daily host rhythms. *Cell Host & Microbe*, **27**, 176–187. https://doi.org/10.1016/j.chom.2020.01.005

Pulgarín-R PC, Gómez C, Bayly NJ, Bensch S, FitzGerald AM, Starkloff N, Kirchman JJ, González-Prieto AM, Hobson KA, Ungvari-Martin J, Skeen H, Castaño MI, Cadena CD (2019) Migratory birds as vehicles for parasite dispersal? Infection by avian haemosporidians over the year and throughout the range of a long-distance migrant. *Journal of Biogeography*, **46**, 83–96. https://doi.org/10.1111/jbi.13453

Raveh S, Heg D, Dobson FS, Coltman DW, Gorrell JC, Balmer A, Röösli S, Neuhaus P (2011) No experimental effects of parasite load on male mating behaviour and reproductive success. *Animal Behaviour*, **82**, 673–682. https://doi.org/10.1016/j.anbehav.2011.06.018

Reinoso-Pérez MT, Dhondt KV, Sydenstricker AV, Heylen D, Dhondt AA (2020) Complex interactions between bacteria and haemosporidia in coinfected hosts: An experiment. *Ecology and Evolution*, **10**, 5801–5814. https://doi.org/10.1002/ece3.6318

Richner H, Christe P, Oppliger A (1995) Paternal investment affects prevalence of malaria. *Proceedings*
of the National Academy of Sciences, **92**, 1192–1194. https://doi.org/10.1038/381565a0

Risely A, Klaassen M, Hoye BJ (2018) Migratory animals feel the cost of getting sick: A meta-analysis
across species. *Journal of Animal Ecology*, **87**, 301–314. https://doi.org/10.1111/1365-2656.12766

Rivero A, Gandon S (2018) Evolutionary ecology of avian malaria: Past to present. *Trends in*
Parasitology, **34**, 712–726. https://doi.org/10.1016/j.pt.2018.06.002

Sánchez CA, Becker DJ, Teitelbaum CS, Barriga P, Brown LM, Majewska AA, Hall RJ, Altizer S (2018) On
the relationship between body condition and parasite infection in wildlife: a review and meta-
analysis (J Davies, Ed,). *Ecology Letters*, **21**, 1869–1884. https://doi.org/10.1111/ele.13160

Sanz JJ, Arriero E, Moreno J, Merino S (2001) Interactions between hemoparasite status and female
age in the primary reproductive output of pied flycatchers. *Oecologia*, **126**, 339–344.
https://doi.org/10.1007/s004420000530

Saraux C, Chiaradia A (2022) Age-related breeding success in little penguins: a result of selection and
ontogenetic changes in foraging and phenology. *Ecological Monographs*, **92**, 10.1111/evo.13443.
https://doi.org/10.1002/ecm.1495

Schoepf I, Olson S, Moore IT, Bonier F (2022) Experimental reduction of haemosporidian infection
affects maternal reproductive investment, parental behaviour and offspring condition.
Proceedings of the Royal Society B: Biological Sciences, **289**, 20221978.
https://doi.org/10.1098/rspb.2022.1978

Siikamäki P, Rätti O, Hovi M, Bennett GF (1997) Association between haematozoan infections and
reproduction in the Pied Flycatcher. *Functional Ecology*, **11**, 176–183.
https://doi.org/10.1046/j.1365-2435.1997.00075.x

Stahlschmidt ZR, Rollinson N, Acker M, Adamo SA (2013) Are all eggs created equal? Food availability
and the fitness trade-off between reproduction and immunity. *Functional Ecology*, **27**, 800–806.
https://doi.org/10.1111/1365-2435.12071

Stearns SC (1989) Trade-offs in life-history evolution. *Ecology*, **3**, 259–268.

Stjernman M, Råberg L, Nilsson J-Å (2008) Maximum host survival at intermediate parasite infection
intensities. *PLOS ONE*, **3**, e2463. https://doi.org/10.1371/journal.pone.0002463

Šujanová A, Špitalská E, Václav R (2021) Seasonal dynamics and diversity of haemosporidians in a
natural woodland bird community in Slovakia. *Diversity*, **13**, 439.
https://doi.org/10.3390/d13090439

Sundberg J (1995) Parasites, plumage coloration and reproductive success in the yellowhammer,
Emberiza citrinella. *Oikos*, **74**, 331. https://doi.org/10.2307/3545664

Van Rooyen J, Lalubin F, Glaizot O, Christe P (2013). Avian haemosporidian persistence and co-

infection in great tits at the individual level. Malaria Journal **12**, 40. . https://doi.org/10.1186/1475-2875-12-40

Valkiūnas G (2005) *Avian malaria parasites and other haemosporidia*. CRC Press, Boca Raton.

Velando A, Drummond H, Torres R (2006) Senescent birds redouble reproductive effort when ill:

confirmation of the terminal investment hypothesis. *Proceedings of the Royal Society B: Biological*

Sciences, **273**, 1443–1448. https://doi.org/10.1098/rspb.2006.3480

Visser ME, Lessells CM (2001) The costs of egg production and incubation in great tits (*Parus major*).

Proceedings of the Royal Society of London. Series B: Biological Sciences, **268**, 1271–1277.

https://doi.org/10.1098/rspb.2001.1661

Zylberberg M, Derryberry EP, Breuner CW, Macdougall-Shackleton EA, Cornelius JM, Hahn TP (2015)

Haemoproteus infected birds have increased lifetime reproductive success. *Parasitology*, **142**,

1033–1043. https://doi.org/10.1017/S0031182015000256