Structure and function of bacterial YeeE-YeeD complex in thiosulfate
uptake pathway

Authors
Mai Ikei[1,†], Ryoji Miyazaki[1,†], Keigo Monden[1], Yusuke Naito[1], Azusa Takeuchi[1],
Yutaro S. Takahashi[1], Yoshiki Tanaka[1], Keina Murata[2], Takaharu Mori[2], Muneyoshi
Ichikawa[3,*], Tomoya Tsukazaki[1,*]

Affiliations
[1]Division of Biological Science, Graduate School of Science and Technology, Nara
Institute of Science and Technology, Ikoma, Nara 630-0192, Japan.
[2]Department of Chemistry, Faculty of Science, Tokyo University of Science, Shinjuku-
ku, Tokyo 162-8601, Japan
[3]State Key Laboratory of Genetic Engineering, Department of Biochemistry and
Biophysics, School of Life Sciences, Fudan University, Shanghai 200438, China.
*Corresponding author. Email: ichikawa_muneyoshi@fudan.edu.cn (M. Ichikawa);
ttsukaza@bs.naist.jp (T.T.)
†These authors contributed equally to this work.

Abstract
Uptake of thiosulfate ions as an inorganic sulfur source from the environment is important
for bacterial sulfur assimilation. Recently, a selective thiosulfate uptake pathway
involving a membrane protein YeeE (TsuA) in *Escherichia coli* was characterized. YeeE-
like proteins are conserved in some bacteria, archaea, and eukaryotes. However, the
precise function of YeeE, along with its potential partner protein in the thiosulfate ion
uptake pathway, remained unclear. Here, we assessed selective thiosulfate transport via
Spirochaeta thermophila YeeE in vitro and characterized *E. coli* YeeD (TsuB) as an
adjacent and essential protein for YeeE-mediated thiosulfate uptake in vivo. We further
showed that *S. thermophila* YeeD possesses thiosulfate decomposition activity and that a
conserved cysteine in YeeD was modified to several forms in the presence of thiosulfate.
Finally, the crystal structures of *S. thermophila* YeeE-YeeD fusion proteins at 3.34-Å and
2.60-Å resolutions revealed their interactions. The association was evaluated by a binding

assay using purified *S. thermophila* YeeE and YeeD. Based on these results, a model of

the sophisticated uptake of thiosulfate ions by YeeE and YeeD is proposed.

125-character sentence

The structural study of the YeeE-YeeD membrane protein complex provides insight into

the thiosulfate uptake and degradation mechanism.

INTRODUCTION

From bacteria to eukaryotes, sulfur is a vital element for cellular activities. For example, sulfur-containing biomolecules, such as L-cysteine, L-methionine, thiamine, glutathione, and biotin, play a variety of essential roles in cells (*1*). Bacteria and plants can utilize L-cysteine as a source of sulfate, but they also have sulfur assimilation pathways to synthesize L-cysteine from inorganic sulfur compounds. In bacteria, L-cysteine is important not only as a component for protein synthesis but also as a reducing agent against oxidative stress (*2*). There are two pathways for bacterial L-cysteine synthesis, which use O-acetylserine as a precursor, called the sulfate and thiosulfate pathways (*3*). In the sulfate pathway, sulfate ion is first decomposed into sulfide ion through phosphorylation by two molecules of ATP and reduction by four molecules of NADPH. Subsequently, L-cysteine is synthesized from sulfide ion and O-acetylserine by O-acetylserine sulfhydrylase-A (CysK). In the thiosulfate pathway, S-sulfocysteine is synthesized from thiosulfate ion and O-acetylserine by O-acetylserine sulfhydrylase-B (CysM); S-sulfocysteine is then reduced by one NADPH molecule, and L-cysteine is synthesized. The sulfate and thiosulfate ions used in these pathways are taken up from the environment by transporters on the cytoplasmic membrane. Sulfate and thiosulfate ions are trapped by periplasmic proteins Sbp (*4*) and CysP (*5*), respectively. Both sulfate and thiosulfate ions are then passed on to a complex formed by the inner membrane proteins CysU and CysW and the cytosolic protein CysA (CysUWA) (also called CysTWA) (*3*) and transported into the cytoplasm by the CysUWA complex using the energy from ATP hydrolysis.

A bacterial membrane protein, YeeE, has been identified as mediating thiosulfate uptake based on growth complementation assay of *Escherichia coli* cells, and the structure of *Spirochaeta thermophila* YeeE (*St*YeeE) has been determined by X-ray crystallography (*6*). YeeE was proposed to be almost entirely buried in the membrane and shows a unique hourglass-like structure. An electron density near a conserved cysteine residue on the outside surface was assigned to a thiosulfate ion, which seemed to be connected to the sulfur atom of the cysteine via a S─H─S hydrogen bond (*6*). In addition, the binding of thiosulfate ions to purified YeeE protein was shown by isothermal titration calorimetry experiments. In the center of YeeE, three conserved, functionally important cysteines, including the thiosulfate-interacting one, are arranged side by side and perpendicular to the membrane. Based on these structural features and in vivo functional

analysis using a series of mutants, thiosulfate ion was proposed to be transported from the environment to the cytoplasm while transiently interacting with the three conserved cysteine residues by relays mediated via hydrogen bonds (*6*), independently of the thiosulfate pathway described above. However, it has not been elucidated whether YeeE alone transports thiosulfate ions. Moreover, no YeeE-associated protein has been clearly defined, although one candidate gene is *yeeD*, which resides within the same operon that includes *yeeE* (*7*), and the regulatory mechanism of the alternative, sophisticated thiosulfate uptake by YeeE remains unclear.

Recently, in addition to the *E. coli yeeE* gene, *yeeD* was shown to be involved in thiosulfate uptake and the two were named *tsuA* and *tsuB*, respectively (*8*). YeeD is a cytoplasmic protein that belongs to the TusA (tRNA 2-thiouridine synthesizing protein A) protein family (*9*). TusA plays various roles in sulfate transfer activities in cells, such as thiomodification of tRNA (*10*), molybdenum cofactor biosynthesis (*11*), and dissimilatory sulfur and tetrathionate oxidation (*12*). The cysteine residue in a Cys-Pro-X-Pro (CPxP) motif of TusA receives activated sulfur from the L-cysteine desulfurase IscS (*13*). Although YeeD possesses the CPxP motif, it cannot complement TusA function (*11*). Therefore, YeeD is thought to have a sulfate-related yet distinct function from TusA in bacteria. In some bacteria, such as Gram-positive *Corynebacterium* species YeeE and YeeD are encoded as one polypeptide, implying that they function together in the thiosulfate uptake pathway. However, the enzymatic activity of YeeD and the functional cooperativity of YeeE and YeeD have not been well characterized.

In this study, first, we measured thiosulfate uptake activity using *St*YeeE-reconstituted liposomes to clarify YeeE function. Second, according to an *E. coli* growth complementation assay, we found that *E. coli* YeeD (*Ec*YeeD) plays an essential role in YeeE function in vivo. Third, we demonstrated a thiosulfate decomposition activity in purified *S. thermophila* YeeD (*St*YeeD). Fourth, direct interaction between *St*YeeE and *St*YeeD proteins was detected, and substrate thiosulfate ions weakened the binding of these proteins. Fifth, the crystal structures of *St*YeeE–YeeD complex were determined at 3.34-Å and 2.60-Å resolutions. Critical residues of YeeD for both its activity and interaction with YeeE were defined. Based on these results, detailed mechanisms for the functional cooperation between YeeE and YeeD in thiosulfate uptake are discussed.

RESULTS

YeeE-mediated thiosulfate transport

Although *E. coli* growth complementation tests showed that YeeE is involved in the uptake of thiosulfate ions (*6*), thiosulfate transport via YeeE has not yet been demonstrated. To detect thiosulfate uptake activity using a purified reconstituted system, we adapted solid-supported membrane (SSM)-based electrophysiology (*14*). First, *St*YeeE-reconstituted liposomes were applied to the SSM, and charge displacement was measured in the presence of sulfate ions or thiosulfate ions (Fig. 1A). A change in the potential toward the negative direction was observed in the presence of thiosulfate ions but not sulfate ions, meaning that negatively charged thiosulfate ions were selectively transported into the proteoliposomes. The slight change in the potential toward the positive direction in the presence of sulfate ions was probably due to buffer shock since there was no counter ion present in these measurements. Next, because *St*YeeE-reconstituted liposomes showed no transport activity for sulfate ions, we measured YeeE activity using sulfate as a counter ion. As a result, a clear peak toward the negative direction (-0.62 nA) was observed only in the presence of *St*YeeE (Fig. 1B). The negative peak area increased when higher concentrations of thiosulfate ions were used (Fig. 1C). These experiments verified that YeeE transports thiosulfate ions specifically but not sulfate ions.

YeeD is essential for the YeeE-mediated thiosulfate pathway

According to recent reports (*7, 8*), YeeD may function in combination with YeeE in *E. coli*. YeeD is composed of about 80 amino acid residues and possesses one or two cysteines, which are highly conserved among bacteria. In some organisms, such as *E. coli*, YeeD has two conserved cysteine residues (C13 and C39 in *Ec*YeeD), while other organisms like *S. thermophila* have one conserved cysteine residue (C17 in *St*YeeD) (Fig. 2A). The invariant cysteine residue in the N-terminal region is part of a CPxP motif that is conserved among TusA family proteins (Fig. 2A and fig. S1). The predicted *Ec*YeeD structure in the AlphaFold2 database (*15, 16*) shows a globular shape with two α-helices and a β-sheet composed of four β-strands (Fig. 2B), similar to other TusA protein structures (*17, 18*). To evaluate the influence of YeeD on YeeE activity in cells, a growth complementation assay using *E. coli* MG1655 *ΔcysPUWA ΔyeeE::kan* (DE3) cells was performed, as described previously (*6*). These *E. coli* cells, which lack sulfate and

thiosulfate ion-uptake pathways, cannot grow in the presence of thiosulfate as a sole

sulfur source (Fig. 2C and fig. S2, A and B), but can when they carry a plasmid containing

the *E. coli yeeE* operon, which includes *yeeE* and *yeeD* (*6*) (Fig. 2C). When the *yeeD*

region was deleted, growth was similar to the negative control, which was conducted

using a vector without *yeeE* and *yeeD* regions (Fig. 2C), revealing that YeeD is essential

for thiosulfate uptake via YeeE in vivo. Further, we prepared a series of alanine mutations

in *Ec*YeeD, including the conserved cysteine mutations C13A and C39A, that are located

on the surface of the AlphaFold2 model (Fig. 2B). Subsequent growth complementation

assays using these YeeD mutations demonstrated the significance of C13 and C39, but

not of the other mutated residues (E19, K21, E26, D31, E32, and D38), in *Ec*YeeD

function (Fig. 2C).

Thiosulfate decomposition activity of YeeD and its catalytic center residue

To unveil the enzymatic function of YeeD itself, we purified *St*YeeD and its mutant

proteins and then conducted their biochemical analyses in vitro. The gel filtration profile

of purified His-tagged *St*YeeD(WT) (11.3 kDa) appears to be a mixture of dimers and

monomers as compared to gel filtration markers (fig. S2C), similar to TusA (*12*). Non-

reducing and reducing SDS-PAGE after the addition of iodoacetamide (IAA) or N-

Ethylmaleimide (NEM), which block the sidechain of Cys residues, showed that almost

all *St*YeeD(WT) lacks disulfide bond formation: the dimeric *St*YeeD apparent in gel

filtration profiles is not due to disulfide bond formation (fig. S2D). In the range of 10–15

kDa on SDS-PAGE gels, purified *St*YeeD(WT) exhibited main and faint minor bands,

whereas *St*YeeD(C17A), a putative catalytic center mutant, exhibited only the main band

(Fig. 2, D and E). Since this slightly upshifted band did not disappear even in reducing

SDS-PAGE (fig. S2D), the -SH group might be modified to an irreversible oxidation state.

Indeed, oxidation of WT *St*YeeD with hydrogen peroxide significantly increased the

reductant-resistant band (fig. S2E), presumably due to irreversible formation of sulfinic

acid ($-SO_2^-$) or sulfonic acid ($-SO_3^-$) (*19*). To examine *St*YeeD modification at C17, we

performed matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass

spectrometry (MS) of YeeD in several conditions. The samples for MS were prepared

under aerobic conditions. The spectrum of purified YeeE in the absence of thiosulfate

revealed three major peaks with m/z of 11317.3, 11349.7, and 11394.3 (Fig. 2D). The

first peak (11317.3 m/z) was assigned as unmodified *St*YeeD (-SH at C17) as its

theoretical m/z value is 11315.7. The second peak (11349.7 m/z) might correspond to

sulfinic acid (-S-O_2^- at C17) and/or persulfide modification (-S-S^- at C17) as it increased

by 32.4 (close to the value of 2 oxygen atoms, 32.0, or 1 sulfur atom, 32.1). Although the

formation of sulfinic acid (see above) may be an artifact due to oxidation in the

purification process, the persulfide modification was also found by MS in purified

Pseudomonas aeruginosa PA1006, an ortholog of *E. coli* TusA (*20*), which has one

conserved cysteine residue corresponding to C17 of *St*YeeD. The third 77.0 m/z-

increased peak (11394.3 m/z) may be attributable to perthiosulfonic acid (-S-SO_3^- at C17).

In contrast, *St*YeeD(C17A) showed only one prominent peak (11283.4 m/z) as its

theoretical unmodified value (11283.6 m/z) (Fig. 2E). These results suggest that the C17

of *St*YeeD is highly reactive to sulfur-related molecules and susceptible to sulfur(S_γ)-

related modifications. Next, we performed MS analysis of the *St*YeeD protein in the

presence of DL-Dithiothreitol (DTT), a reductant. The third peak entirely disappeared

after the addition of DTT (Fig. 2F). By contrast, the second peak remained even in the

presence of DTT. Persulfide modification (-S-S^-) can be reduced by DTT (*20*) but sulfinic

acid (-S-O_2^-), an overoxidation product, is irreversible as previously shown (*19, 21*).

Therefore, in the DTT condition, the second peak is thought to be residual sulfinic acid

product. Using sample treated with DTT, to gain insights into catalytic intermediates, MS

analysis was performed in the presence of thiosulfate (Fig. 2G). In the condition with

thiosulfate (500 μM), a peak with m/z value of 11431.0 appeared. The peak with an m/z

value of 11431.0 may correspond to -S-$S_2O_3^-$ because the m/z increments were equivalent

to the addition of S_2O_3 to StYeeD(-SH). The previously reported crystal structure of

TsdBA (PDB ID 5LO9) included the same modification of a cysteine residue's thiol (-

SH) group to -S-$S_2O_3^-$, indicating that this modification is one of the stable forms (*22*). In

the case of StYeeD(C17A), the addition of thiosulfate did not increase the m/z (Fig. 2E).

In addition, we performed MS using *St*YeeD sample treated with DTT and then hydrogen

peroxide (Fig. 2H). In this condition, the peak corresponding to *St*YeeD(-SH) completely

disappeared, but peaks corresponding to sulfinic acid (-S-O_2^-), sulfonic acid (-S-O_3^-), and

perthiosulfonic acid (-S-SO_3^-) were observed. Given these results, although

perthiosulfonic acid (-S-SO_3^-) detected in the purified StYeeD is thought to be due to

oxidization during the purification procedure, *St*YeeD may temporarily take the -S-SO_3^-

form in the process of thiosulfate decomposition.

The additional binding of S_2O_3 to the C17 side chain, in form of -S-$S_2O_3^-$, and

the existence of persulfide modification (-S-S^- at C17) and perthiosulfonic acid (-S-SO_3^-

at C17), as indicated by the MS results, raise the possibility that YeeD decomposes

thiosulfate ion ($S_2O_3^{2-}$) and releases sulfide ion (S^{2-}). To test this, we monitored hydrogen

sulfide (H_2S) derived from thiosulfate-decomposed sulfide ions using a fluorescent probe

for H_2S, HSip-1. A significant increase in fluorescence intensity was observed in the

presence of *St*YeeD compared with an irrelevant protein, BSA, and with no protein (Fig.

2I and fig. S2G), suggesting that YeeD catalyzes the decomposition of thiosulfate as a

substrate. We also tested the thiosulfate decomposition activity of several *St*YeeD

mutants. Among these, only C17A did not show a significant increase in fluorescence;

NEM-modified *St*YeeD also did not. These results indicated that the conserved cysteine

residue of YeeD (C17 in *St*YeeD) is the catalytic center residue for thiosulfate

decomposition.

Crystal structure of the YeeE-YeeD complex

We succeeded in crystallizing a fusion protein of *St*YeeE and *St*YeeD, designed based on

the amino acid sequence of *Corynebacterium pollutisoli* YeeED, which is expressed as a

single protein. First, because a comparison of *St*YeeE/D and several YeeEDs' amino acid

sequences revealed that *C. pollutisoli* YeeED possesses the shortest linker, of 40 amino

acids, between its YeeE and YeeD regions, we prepared the fusion protein by introducing

the 40 residues between the C-terminus of *St*YeeE and the N-terminus of *St*YeeD. Second,

after attempting but failing to crystallize the fusion protein, *St*YeeE-*St*YeeD, we tried

crystallization using alanine mutants of the conserved cysteines in *St*YeeE-*St*YeeD and

succeeded in crystallizing *St*YeeE(C22A)-*St*YeeD. Using the X-ray diffraction dataset of

*St*YeeE(C22A)-*St*YeeD, we determined the crystal structure of the *St*YeeE-*St*YeeD

complex at 3.34 Å (fig. S3A,B, and Table 1 8J4C). However, the resolution was

insufficient to reveal the details of the interactions between YeeE and YeeD. Introducing

a mutation in YeeD (L45A) strengthened the interaction, as found in binding experiments

using purified YeeE and YeeD mutants (described later). The diffraction dataset of

crystals of *St*YeeE(C22A)-*St*YeeD(L45A) was improved to 2.60 Å (Fig. 3A and Table 1

8K1R), and the resulting 2.60 Å crystal structure enabled us to discuss the detailed

interactions between YeeE and YeeD.

Each crystal lattice in the datasets was the same, except for having slightly
different unit cell dimensions (Table 1). The asymmetric unit contains two molecules of
*St*YeeED, Mol A and Mol B. In the crystal packing, *St*YeeD in Mol A is involved in
interacting with an adjacent YeeD region, but YeeD in Mol B is not (fig S3A). When the
two *St*YeeE-*St*YeeD structures (Mol A and Mol B), showing different packing
interactions, were superimposed, the crystal structures were almost identical with an
RSMD of 0.433 Å for 8J4C or 0.155 Å for 8K1R for Cα atoms (fig. S3A). Although the
orientation of YeeD relative to YeeE is slightly different, the RMSD between 8J4C and
8K1R for Cα atoms is 0.896 (fig. S3C) and the active site of YeeD is similar. Therefore,
in this paper, we introduce Mol A of the higher resolution structure, which shows the best
electron density map, to explain YeeE and YeeD interactions.

The overall structure of YeeE in the crystal structure of *St*YeeE-*St*YeeD
resembles the previously reported crystal structure of *St*YeeE (PDB ID 6LEP) (*6*), with
an RMSD of 0.596 Å for Cα atoms (fig. S3D). The cytoplasmic indentation of *St*YeeE
associates with *St*YeeD, which consists of two α-helices and four β-sheets. The
architecture of *St*YeeD is similar in shape to the sequence-homologous *E. coli* TusA (22.5%
identical) (PDB ID 3LVK) (fig. S3E) (*18*). Residues involved in the YeeE-YeeD
interaction are shown in Fig. 3B. C22 of *St*YeeE is located at the interface, implying that
C22A affects the interaction. Residues R28, R98, E101, and E261 on the cytoplasmic
surface, conserved in YeeEs (*6*), were found to be involved in the interaction with YeeD.
While the important, conserved residues C293, C91, and C22, termed the first, second,
and third cysteines, respectively, are arranged perpendicular to the membrane at 8.6- and
6.7-Å intervals, C17 of *St*YeeD, one of the YeeE-interacting residues, is the putative
enzymatic active site and is located adjacent to C22 of *St*YeeE as the "fourth cysteine".
This positioning means that a thiosulfate ion transported through the membrane via the
center region of YeeE interacts with the first-to-third cysteines of YeeE in succession and
then interacts with the fourth cysteine of YeeD. The distance between the third and the
fourth cysteines, 10.8 Å, is wider than that between the other cysteines, and this space
may be needed for transient binding of the thiosulfate ion rather than its transport (Fig.
3C). A water molecule is located at position III, a predicted thiosulfate-binding site (6).
A position where another water molecule is located below position III corresponds to the
position of the side chain of C22 (6). The relay of a thiosulfate ion from position III to
C17 is likely facilitated by the conserved positively charged residues R28 and R98 (Fig.

3C). The CPxP motif forms a cis peptide between C17 and P18 and is located at the
beginning of the α-helix (Fig. 3D), similar to a previous report (*17*). In addition, the
electron density of the peptide bonds between V16 and C17 is faint, which may be closely
related to C17 being the active center.
271

Interaction between YeeE and YeeD.
To confirm that the interactions between YeeE and YeeD revealed in the crystal structures
of *St*YeeE-YeeD fusion protein also occur in solution, we performed a binding assay
based on the bio-layer interferometry (BLI) method (*23*) using purified *St*YeeE and His-
tagged *St*YeeD. Clear association and dissociation of *St*YeeE with an *St*YeeD-solidified
Ni-NTA sensor were observed as real-time wavelength shifts (Fig. 4A, blue). Based on
the *St*YeeE binding-dependent wavelength changes, the K_D between *St*YeeD and *St*YeeE
was estimated as 1.6 ± 0.20 μM. When a buffer containing sulfate ion was used, the
binding was similar to that in a buffer without a sulfur source (fig. S4, A and B). In
contrast, *St*YeeE showed significantly decreased binding to *St*YeeD in a buffer containing
thiosulfate ions (fig. S4, A and B). We next sought the important residues for the
interaction using a series of *St*YeeD mutants (fig. S2F). The mutants E15A, V16A, C17A,
D42A, Y43A, E48A, and R49A severely impaired *St*YeeD's affinity for *St*YeeE. while
the mutant L45A enhanced the affinity (Fig. 4B). This enhanced affinity could stabilize
the crystal packing of *St*YeeE-*St*YeeD, leading to its higher-resolution structure. In the
crystal structure, most of these influential residues for affinity were located at the border
region between YeeE and YeeD (Fig. 4C). On the other hand, Ala substitutions on the
opposite side maintained affinity (Fig. 4). The calculated K_a, K_d and K_D values for YeeD
mutant proteins are summarized in Table 2. Notably, the conserved cysteine residue (C17)
of *St*YeeD proved to be pivotal for not only the enzyme activity but also the interaction
with *St*YeeE. These BLI results in solution are consistent with the crystal structure.
To further assess the important residues of YeeE for the interaction with YeeD,
we used a growth complementation assay. *Ec*YeeE mutations of residues at the
interaction surface (R21A, R101A, E104A, E271A, R275A) could not complement the
growth of *E. coli* MG1655 *ΔcysPUWA ΔyeeE::kan* (DE3) cells (fig. S4, C and D). These
results emphasize that the interaction of YeeE and YeeD is essential for in vivo function
of thiosulfate uptake.
299

MD simulation of YeeE/YeeD complex.

In the presence of thiosulfate ions, *St*YeeD was easily modified (Fig. 2G) and the binding between *St*YeeE and *St*YeeD decreased (fig. S4A, B), leading to the hypothesis that the modified *St*YeeD dissociates from *St*YeeE. Here, to address how the *St*YeeE-YeeD complex undergoes structural changes when *St*YeeD is in the YeeD-SH or YeeD-S-$S_2O_3^-$ states, we conducted molecular dynamics (MD) simulation analysis (fig. S5). In the case of *St*YeeD-SH, no significant structural changes were observed during the 1-μs MD simulation (fig. S5A): the simulation structure remained similar to the crystal structure of the *St*YeeE-YeeD complex (PDB ID 8K1R) (fig. S5C). On the other hand, in the case of *St*YeeD-S-$S_2O_3^-$, the *St*YeeD root mean square deviations (RMSD) for C-alpha atoms increased from around 400 ns; in other words, *St*YeeD moved away from the cytoplasmic concave surface of *St*YeeE (fig. S5A–C). The calculation results are consistent with the idea that the interaction between YeeE and YeeD weakens upon Cys-modifications of YeeD. The dissociation of YeeD from YeeE may facilitate the release of thiosulfate ions' decomposition products from the Cys positions of YeeD into the cytoplasm.

DISCUSSION

By the growth complementation assay, we have demonstrated that YeeD is a required enzyme for growth when thiosulfate is solely imported via YeeE. YeeD's substrate was identified as the thiosulfate ion, and the uptake activity for thiosulfate ions was measured using YeeE-reconstituted liposomes. In addition, the interaction between YeeE and YeeD was revealed by both binding assay and X-ray crystallography. By mutational analysis of YeeD, a conserved cysteine residue was found to be critical for YeeE's activity in vivo as well as thiosulfate ion decomposition activity.

Compared with other TusA family proteins, YeeD shows some similarities and differences. Although conserved cysteine residues in other TusA family proteins, corresponding to C17 in *St*YeeD, are important for their own activities (*10, 20*), TusA could not decompose thiosulfate ions (*24*). *Metallosphaera cuprina* TusA was reported to exist as both monomer and dimer forms (*12*). Similarly, *St*YeeD appeared to be monomers and dimers in gel filtration (fig. S2C). *St*YeeD may exploit this feature to transiently form a dimer in the thiosulfate decomposition reaction (discussed below). The conserved cysteine residues in YeeD and TusA are important for binding to their respective partners (Fig. 4) (*12, 18*), which seems to be a common feature in TusA family

proteins. Meanwhile, YeeD is unique in that its interaction partner, YeeE, is a membrane protein, unlike other TusA family proteins. Previously, the conserved cysteine residue of PA1006 was reported to be persulfide-modified (*20*). Furthermore, *St*YeeD has several different sulfur-related modification statuses, including persulfide modification, perthiosulfonic acid, and thiosulfonate (Fig. 2, D, F–H). In the case of DsrE3 and TusA, the conserved cysteines can possess thiosulfonate, but the modifications were detected only with their substrate tetrathionate (*12*). Along with the fact that YeeE only allows thiosulfate to pass through, YeeD also seems to be specialized for thiosulfate reactivity. Both can contribute to the efficient uptake of thiosulfate. Since YeeE alone can transport thiosulfate ions but was unable to rescue the growth deficiency in vivo, cooperation between YeeE and YeeD is crucial. In this study, we have unveiled a sophisticated pathway for thiosulfate uptake in the bacterial membrane. An open question is whether there are other proteins that directly receive sulfur compounds from YeeD in sulfur assimilation pathways.

Based on our findings, we propose detailed mechanisms for thiosulfate ion uptake by YeeE and YeeD (Fig. 5A). From the previous crystal structure of *St*YeeE, the transportation of thiosulfate ions across the membrane was proposed to be relayed by the first-to-third cysteine residues (*6*). Our crystal structure of the *St*YeeE and *St*YeeD complex now further shows that YeeD is positioned to interact with the YeeE cytoplasmic side, which we propose to be the exit of the thiosulfate pathway on YeeE; the conserved cysteine residue of YeeD can function as a "fourth cysteine". Therefore, the interactions between the two can facilitate the delivery of a thiosulfate ion from YeeE to YeeD: the conserved cysteine residue of YeeD can directly capture a thiosulfate ion passed through YeeE. This transport scheme of thiosulfate ions by YeeE and YeeD recalls a group-translocation originally proposed by Mitchell and Moyle (*25*). From our X-ray crystal structure, YeeE and YeeD appear to work as a 1:1 complex. The fact that YeeE and YeeD are encoded as a single polypeptide in some species supports this idea. After receiving the thiosulfate ion, YeeD's conserved cysteine would be thiosulfonated. Our BLI method analysis in the presence of thiosulfate showed that the interaction between *St*YeeD and *St*YeeE was weakened, possibly due to thiosulfonation on YeeD. The thiosulfonated YeeD may therefore dissociate from YeeE and be released into the cytoplasm. This dissociation model agrees well with our MD simulation results. Since YeeD can decompose thiosulfate by itself, it would do so after dissociating from YeeE. Of

additional note, since the existence of StYeeD dimers in solution was suggested by gel
filtration chromatography, StYeeD may form a transient dimer after being released from
StYeeE.

We suggest a plausible mechanism for decomposition of thiosulfate ion by
YeeD (Fig. 5B). While the reducing power for this series of reactions is thought to be
supplied from the cytoplasm, a reducing environment, the electron route remains
unknown and needs to be elucidated in a future study. In the presence of thiosulfate ion,
StYeeD incorporated thiosulfate, but StYeeD(C17A) did not (Fig. 2E and G), suggesting
that the thiosulfate ion directly binds to the cysteine residue of YeeD. Our MS results also
suggest that there are StYeeD molecules with perthiosulfonic acid (-S-SO_3^-) and
persulfide modifications (-S-S^-). Taking this together with the detection of H$_2$S in the
presence of StYeeD and thiosulfate ion (Fig. 2H), it would be reasonable for sulfide ion
(S^{2-}) to be released from a persulfide-modified version of StYeeD (-S-S^-) (Fig. 5B).
Although sulfonic acid (-SO_3^-) formation by the oxidization of a cysteine residue is an
irreversible process, perthiosulfonic oxidization of a cysteine residue (-S-SO_3^-) is
reversible as shown by Doka et al. (*19*). Therefore, even if -S-SO_3^- is formed, sulfite ion
(SO$_3^{2-}$) will be released. In our experimental condition for the thiosulfate decomposition
assay, reductant β-ME is present to facilitate these reactions. In the cell, there could be
other acceptor proteins which take S or SO$_3$ atoms from YeeD protein. Such proteins
should be identified in a future study. While the significance of the StYeeD dimer in this
study has not been demonstrated, if it were involved in this series of reactions, the scheme
depicted in the Fig. 5 box would likely occur. Because StYeeD does not form disulfide
bonds in its dimer (fig. S2C and D), one StYeeD molecule in a dimer can be
thiosulfonated and the other can have an active -SH group, which can take the SO_3 from
thiosulfonated YeeD (-S-$S_2O_3^-$). Through this process, a persulfide-modified form (-S-S^-)
can be generated. In some bacteria, including *E. coli*, YeeD has two cysteine residues. In
species having YeeD with two cysteine residues, the above-mentioned process could be
carried out by one YeeD molecule using these two cysteine residues. Another possibility
is that the non-conserved cysteine (C39) of EcYeeD may not serve as the decomposition
active center, but may interact with YeeE and/or enhance its activity. In agreement with
these ideas, our growth complementation assay showed that both cysteines are important
for the growth of *E. coli* cells.

In the 2.60-Å structure of *St*YeeE-*St*YeeD, the electron density is faint on the
N-terminal side of the CPxP motif, raising the possibility that some electrons may be
attracted to the CPxP motif (Fig. 3D). If that happens, the CPxP motif could make the
region electron-rich and exhibit high reducing power. Previously, the CPxP architecture
was proposed to stabilize an α-helix-1 by capping it (*17*). In the TusA family proteins,
the formation of CPxP may also provide high reducing power, conferring on them the
unique property of selective decomposition of their substrates.

In conclusion, we revealed that YeeE and YeeD cooperatively contribute to the
thiosulfate uptake pathway and possess unprecedented regulation mechanisms. Our
findings also deepen the understanding of the functions of TusA family proteins.

MATERIALS AND METHODS

Strains and plasmids

For the growth complementation assay, we used plasmids derived from pAZ061, which
is based on pET-16b-TEV (*26*) and possesses *E. coli yeeE* and *yeeD* genes between
BamHI and XhoI sites amplified from *E. coli* genomic DNA (JCM 20135, RIKEN BRC)
using   primer   set   5′-
AAATTTATATTTTCAAGGATCCCATATGTTTTCAATGATATTAAG-3′ and 5′-
GGCTTTGTTAGCAGCCCTCGAGTCAGGCTTTTTGAACGG-3′ (fig. S2A), and the
E. coli MG1655 *ΔcysPUWA ΔyeeE::kan* (DE3) strain, which cannot grow on a minimum
medium with thiosulfate ion as the single sulfur source (*6*). pAZ061 derivatives having
mutations in the *yeeD* region were prepared by site-directed mutagenesis. The DNA
sequence encoding *Ec*YeeD was deleted from pAZ061 using primers 5′-
GTTTGGGTTAGGCATCGCTTCCCCAACGGCC-3′   and   5′-
GGCCGTTGGGGAAGCGATGCCTAACCCAAAC-3′.

The plasmid used to express *St*YeeE (1-328aa, UniProt ID: G0GAP6) with C-
terminal GSSGENLYFQGEDVE-His$_8$ sequence (pKK550) was prepared as in (*6*). For
the expression of the *St*YeeE-*St*YeeD fusion protein, we modified pKK550; the resulting
plasmid,   pAZ150,   encodes   *St*YeeE   (1–330   aa)-
AATPTPVAEAAPSSAEDRVLPFQVATGAVALQTAPRVKKA-*St*YeeD (1–80 aa,
UniProt ID: G0GAP7)-GSSGENLYFQGEDVE-His$_6$. The mutations in the *St*YeeE and
*St*YeeD regions were introduced by site-directed mutagenesis.

For the expression of *St*YeeD, the DNA sequence encoding *St*YeeD (1–80 aa)
was inserted into a modified pCGFP-BC plasmid. The resulting plasmid, pNT015,
expresses *St*YeeD (1–80 aa)- GSSGENLYFQGEDVE-His$_6$. The pNT015 derivatives
were prepared by Gene Synthesis and Mutagenesis (SC1441, GenScript) or site-directed
mutagenesis.
435

Growth complementation tests of *E. coli* cells

E. coli MG1655 *ΔcysPUWA ΔyeeE::kan* (DE3) cells harboring pAZ061 or its derivatives
were cultured in LB medium containing 50 μg/ml ampicillin and 25 μg/ml kanamycin for
16 h at 37 °C. The culture was diluted 100-fold and cultured for another 8 h at 37 °C,
after which the cells were precipitated by centrifugation and washed twice with S-free
medium (42 mM Na$_2$HPO$_4$, 22 mM KH$_2$PO$_4$, 8.6 mM NaCl, 19 mM NH$_4$Cl, 1 mM MgCl$_2$,
0.2% (w/v) glucose, 0.01% (w/v) thiamine hydrochloride, and 0.1 mM CaCl$_2$).
Subsequent passage cultures in S-free medium containing 500 μM Na$_2$S$_2$O$_3$ were started
at OD$_{600}$ = 0.2. The ΔOD$_{600}$ of the culture was measured every 30 min with OD-Monitor
C&T (TAITEC) for 24 h. Measurement was performed three times for each transformant.
446

Protein expression and purification

*St*YeeE and *St*YeeE(C22A)-*St*YeeD proteins were purified as follows. *E. coli* C41(DE3)
cells transformed with plasmids expressing these proteins were cultured in 2.5 l of LB
medium containing 50 μg/ml ampicillin until the OD$_{600}$ reached 0.4. Isopropyl β-D-
thiogalactopyranoside (IPTG) was then added to 1 mM, and the cells were cultured at
30 °C for 17 h. Subsequent procedures were performed at 4 °C or on ice. The cells were
collected by centrifugation as pellets, suspended in a buffer (10 mM Tris-HCl pH 8.0,
300 mM NaCl, 1 mM EDTA-Na pH 8.0, 2 mM Na$_2$S$_2$O$_3$, 0.1 mM phenylmethylsulfonyl
fluoride (PMSF), and 1 mM β–mercaptoethanol (β-ME)), and disrupted with a
Microfluidizer Processor M-110EH (Microfluidics International). The suspension was
centrifuged (10,000 rpm for 20 min, himac R13A rotor), and the collected supernatant
was further ultracentrifuged (40,000 rpm for 60 min, Beckman 45Ti rotor) to obtain the
membrane fraction, which was flash-frozen in liquid nitrogen and stored at -80 °C until
purification. The membrane fraction was resuspended in solubilization buffer (20 mM
Tris-HCl pH 8.0, 300 mM NaCl, 2 mM Na$_2$S$_2$O$_3$, 10 mM imidazole-HCl pH 8.0, 5% (v/v)
glycerol, 1 mM β-ME, 0.1 mM PMSF) containing 1% (w/v) n-dodecyl β-maltoside

(DDM) and stirred at 4 °C for 60 min. The insoluble fraction was removed by

ultracentrifugation (45,000 rpm for 30 min, Beckman 70Ti rotor), and the supernatant

was mixed with 5 ml of Ni-NTA agarose (QIAGEN) pre-equilibrated with solubilization

buffer for 60 min. Then, 100 ml of wash buffer (20 mM Tris-HCl pH 8.0, 300 mM NaCl,

2 mM $Na_2S_2O_3$, 50 mM imidazole-HCl pH 8.0, 5% (v/v) glycerol, 1 mM β-ME, 0.1 mM

PMSF, 0.1% (w/v) DDM) was added to the column. Next, 5 ml of elution buffer (20 mM

Tris-HCl pH 8.0, 300 mM NaCl, 2 mM $Na_2S_2O_3$, 200 mM imidazole-HCl pH 8.0, 5%

(v/v) glycerol, 1 mM β-ME, 0.1 mM PMSF, and 0.1% (w/v) DDM) was added to the

column six times, and the fractions with target proteins were pooled. To remove the His-

tag, TEV(S219V) protease (*27*) was added with a protein weight ratio of 10 (proteins):1

(TEV) and the protein solution was dialyzed overnight against dialysis buffer (20 mM

Tris-HCl pH 8.0, 300 mM NaCl, 2 mM $Na_2S_2O_3$, 0.1% (w/v) DDM, 1 mM β-ME). Ni-

NTA agarose (2.5 ml) pre-equilibrated with dialysis buffer was added to the sample and

stirred for 60 min to remove TEV proteases. The flow-through fraction was collected,

concentrated with an Amicon Ultra 50K NMWL (Merck Millipore), and ultracentrifuged

(45,000 rpm for 30 min, himac S55A2 rotor). The sample was then applied to a Superdex

200 increase 10/300 GL column (Cytiva) equilibrated with a buffer (20 mM Tris-HCl pH

8.0, 300 mM NaCl, 2 mM $Na_2S_2O_3$, 0.1% (w/v) DDM, 1 mM β-ME). The fractions with

target proteins were collected and pooled. *St*YeeE(C22A)-*St*YeeD(L45A) protein was

purified as for *St*YeeE and *St*YeeE(C22A)-YeeD except that the buffer lacked $Na_2S_2O_3$.

*St*YeeD purification was performed as follows. *E. coli* C41(DE3) cells containing

*St*YeeD-encoding plasmids were inoculated into 25 ml of LB medium containing 50

μg/ml ampicillin and cultured overnight at 37 °C. The culture was then added to 0.5 l of

LB medium containing 50 μg/ml ampicillin and allowed to grow until $OD_{600} = 0.4$. After

the addition of IPTG to 1 mM, the cells were cultured at 30 °C for 17 h. The cells were

collected by centrifugation (4,500 rpm for 10 min, himac R9A2), flash-frozen in liquid

nitrogen, and stored at -80 °C until use. The next procedures were performed at 4 °C or

on ice. The frozen pellets were suspended in 25 ml of sonication buffer (20 mM Tris-HCl

pH 8.0, 300 mM NaCl, 10 mM imidazole-HCl pH 8.0, 1 mM β-ME, 0.1 mM PMSF),

sonicated for 10 min on ice with a Q500 sonicator (QSONICA), and centrifuged (15,000

rpm for 30 min, himac R13A). Ni-NTA agarose (2.5 ml) pre-equilibrated with the

sonication buffer was added to the supernatant and rotated for 1 h. After the flow-through

fraction was removed, the resin was washed with 125 ml of a buffer (20 mM Tris-HCl

pH 8.0, 300 mM NaCl, 20 mM imidazole-HCl pH 8.0, 1 mM β-ME, 0.1 mM PMSF).

Next, 5 ml of a buffer (20 mM Tris-HCl pH 8.0, 300 mM NaCl, 200 mM imidazole-HCl

pH 8.0, 1 mM β-ME, 0.1 mM PMSF) was added, and the elution fraction was collected.

The elution step was repeated six times. The fractions with target proteins were pooled

and concentrated using an Amicon Ultra 3K NMWL (Merck Millipore). The sample was

ultracentrifuged (45,000 rpm x 30 min, himac S55A2 rotor) and loaded onto a Superdex

200 increase 10/300 GL column pre-equilibrated with YeeD gel filtration buffer (10 mM

Tris-HCl pH 8.0 and 300 mM NaCl). The fractions with *St*YeeD proteins were

concentrated using an Amicon Ultra 3K NMWL.

MALDI-TOF MS

For *St*YeeD(WT) sample preparation for MS (Fig. 2D), *St*YeeD(WT) protein was mixed

with gel filtration buffer, so that the final concentration of *St*YeeD(WT) was 48.2 μM and

the total volume was 200 μl, and incubated for 2.5 h at 37 °C. The sample was then

ultracentrifuged, and the buffer was exchanged with MS buffer (10 mM Tris-HCl pH 8.0)

using a NAP-5 Column (Cytiva). The final concentration of *St*YeeD(WT) for MS analysis

was 17.5 μM.

For *St*YeeD(C17A) sample preparation for MS (Fig. 2E), 48.2 μM

*St*YeeD(C17A) in gel filtration buffer was incubated with or without 500 μM thiosulfate

for 2.5 h at 37 °C in a total volume of 200 μl. After the incubation, the sample was

ultracentrifuged, and the buffer was exchanged with MS buffer using a NAP-5 Column.

The final concentrations of *St*YeeD(C17A) were 12.3 μM for without thiosulfate and 16.6

μM for with thiosulfate.

For the DTT condition of *St*YeeD(WT) (Fig.2F), DTT was added to

*St*YeeD(WT) to give final concentrations of 401 μM and 16.7 mM, respectively, and a

total volume of 288 μl. The sample was then incubated for 10 min at 37 °C and

ultracentrifuged. A 48-μl aliquot was taken and the buffer was exchanged with MS buffer

using a NAP-5 Column. The concentration of resultant DTT-treated *St*YeeD(WT) was

10.96 μM. For the thiosulfate or hydrogen peroxide condition of *St*YeeD(WT), the DTT-

treated *St*YeeD(WT) sample was processed as follows. First, to remove DTT, the sample

was applied to a Superdex 200 Increase 10/300 GL column (Cytiva) pre-equilibrated with

gel filtration buffer. Next, the sample after gel filtration was concentrated using Amicon

ultra 3K NMWL. For the thiosulfate condition (Fig. 2G), the sample was then mixed with

thiosulfate so that the final concentrations of *St*YeeD(WT) and thiosulfate were 48.2 μM

and 500 μM, respectively, and the total volume was 200 μl. After incubating for 2.5 h at

37 °C, the sample was ultracentrifuged, and the buffer was exchanged with MS buffer

using a NAP-5 Column. The final concentration of *St*YeeD(WT) for MS analysis was

14.0 μM. For the hydrogen peroxide condition (Fig. 2H), the sample after gel filtration

and concentration was mixed with hydrogen peroxide so that the final concentrations of

*St*YeeD(WT) and hydrogen peroxide were 48.2 μM and 0.1% (w/v), respectively, and the

total volume was 200 μl. After incubating for 10 min at room temperature, the sample

was ultracentrifuged, and the buffer was exchanged with MS buffer using a NAP-5

Column. The final concentration of *St*YeeD(WT) for MS analysis was 23.8 μM.

Samples for MALDI-TOF analysis were prepared by the sinapinic acid (SA)

double layer method (*28*). In brief, 1 μl matrix solution A (saturated solution of SA in

ethanol) was deposited onto the MALDI target and allowed to dry. Matrix solution B

(saturated solution of SA in TA30 solvent (1:2 (v/v) acetonitrile:0.1% (v/v)

trifluoroacetic acid in water)) and protein solution were mixed at a ratio of 1:24 (v/v).

Matrix solution B/protein mixture was then deposited onto the matrix spot and allowed

to dry. The dried samples were analyzed by Autoflex-II (Bruker Daltonics) with a linear

positive mode and acquisition mass range of 3,000–20,000 Da. Theoretical m/z values

were calculated with the equation of $m/z = (M + n)/n$, where M is molecular mass and n

is the charge (n = 1 was adopted).

Measurement of enzyme activity of *St*YeeD

The release of H_2S due to chemical decomposition of thiosulfate ions by *St*YeeD was

monitored by HSip-1 (Dojindo, SB21-10). The reaction solution contained 19.6 μM

HSip-1, 24.1 μM YeeD, and 70 μM β-ME in the YeeD gel filtration buffer. As negative

controls, 24.1 μM BSA was used instead of YeeD, or no protein was added to the solution.

The reaction was started by adding 5 μl of 100 mM $Na_2S_2O_3$ at a 1:200 [v/v] ratio (final,

500 μM $Na_2S_2O_3$), and the solution was incubated at 37 °C. The fluorescence of HSip-1

was measured using a fluorescence spectrophotometer (F-7000, Hitachi), at an excitation

wavelength of 491 nm and an emission wavelength of 521 nm, every 30 min until 150

min.

Crystallization

Purified *St*YeeE(C22A)-*St*YeeD fusion protein was concentrated to 21.5 mg/ml by

Amicon Ultra 50K NMWL and crystallized using the lipidic cubic phase (LCP) method,

as previously performed in the case of purified *St*YeeE (*6*). Fifteen microliters of 21.5

mg/ml *St*YeeE(C22A)-*St*YeeD was mixed with 4.3 μl of a buffer (20 mM Tris-HCl pH

8.0, 300 mM $Na_2S_2O_3$, 0.1% (w/v) DDM, 1 mM β-ME) and incubated on ice for 20 min.

Then, 16 μl of the sample was mixed with 24 μl of monoolein (M-239, Funakoshi) in an

LCP syringe (Art Robbins Instruments) for 10 min. After 30 min of incubation at 20°C,

30 nl of the mixed samples were spotted onto MRC under oil crystallization plates

(Hampton Research) using the Crystal Gryphon protein crystallization aliquot system

(Art Robbins Instruments) with 3 μl of buffers (18 to 24% (v/v) pentaerythritol-

propoxylate (5/4 PO/OH), 100 mM 2-morpholinoethanesulfonic acid (MES)-NaOH pH

7.0, and 300 mM NaCl) covering them. The plate was incubated at 20 °C for 7 days. For

*St*YeeE(C22A)-*St*YeeD(L45A) fusion protein, the purified protein was concentrated to

16.5 mg/ml and crystallized using the LCP method. The crystals appeared using 0.35 M

ammonium formate, 0.1 M Tris-HCl (pH 8.0 to 9.0), and 22 to 42% (v/v) 1,4-Butanediol

as covering solutions. The plate was incubated at 20 °C for 14 days. The crystals that

appeared were harvested by Crystal Mounts and Loops (MiTegen) without using

cryoprotectant, directly flash-frozen in liquid nitrogen, and stored in liquid nitrogen until

X-ray diffraction experiments.

Data collection and structural determination

X-ray diffraction experiments of *St*YeeE(C22A)-*St*YeeD and *St*YeeE(C22A)-

*St*YeeD(L45A) were performed at beamline BL32XU of SPring-8. Data were collected

with the automated data collection system ZOO (*29*). The complete data sets were

obtained by combining multiple small wedge datasets from hundreds of tiny crystals with

sizes of approximately 20 μm. The principle of this method has been described in detail

(*30*). Data processing was performed with KAMO (*30*) using XDS (*31*) programs. For

*St*YeeE(C22A)-*St*YeeD dataset, the initial phase was determined by molecular

replacement using the *St*YeeE crystal structure (PDB ID: 6LEO) as a template by

PHASER (*32*), and a *St*YeeD structure predicted by AlphaFold2 (*15, 16*) was manually

fitted to the density map using COOT (*33*). Structure refinement was performed using

COOT (*33*) and PHENIX (*34*) iteratively until R_{work}/R_{free} reached 0.265/0.310 at 3.34 Å

resolution. For *St*YeeE(C22A)-*St*YeeD(L45A) dataset, the initial phase was determined

by molecular replacement using the *St*YeeE(C22A)-*St*YeeD structure as a template by

MOLREP (*35*). Refinement of the structure was performed using COOT (*33*) and

PHENIX (*34*) in a iterative way until R_{work}/R_{free} reached 0.206/0.255 at 2.60 Å resolution.

Figures of the structures were prepared using PyMOL (https://pymol.org/2/) and Chimera

(*36*).

Interaction analysis between *St*YeeE and *St*YeeD by the BLI method

BLI method (*23*) was performed to analyze the interaction between *St*YeeE and *St*YeeD-

His$_6$ proteins using the Octet N1 System (Sartorius) at room temperature. A Ni-NTA

biosensor was hydrated with Octet buffer (20 mM Tris-HCl pH 8.0, 300 mM NaCl, 0.1%

(w/v) DDM, 1 mM β-ME) for 10 min and mounted in the Octet N1 System. The biosensor

was first dipped in Octet buffer, and the initial baseline was measured for 30 sec. Next,

the biosensor was dipped in 4.82 μM *St*YeeD-His$_6$ protein solution in Octet buffer for

120 sec for loading. The solution was then changed to Octet buffer for 30 sec to measure

the baseline, after which, the biosensor was submerged in 11.05 μM *St*YeeE solution in

Octet buffer for 120 sec to measure the association of *St*YeeE. Finally, the buffer was

exchanged to Octet buffer, and the dissociation of *St*YeeE from the biosensor was

measured for 120 sec. Biosensors without *St*YeeD-His$_6$ proteins were measured with the

same procedure and used as references. After the reference data were subtracted from

*St*YeeD-His$_6$ protein data, the association rate constant (k_a), dissociation rate constant (k_d),

and affinity constant (K_D) were calculated by a local fitting method. Measurements were

performed three times for each protein.

Reconstitution of proteoliposomes

A mixture of 0.8 mg/ml *St*YeeE and 4 mg/ml *E. coli* total lipid extract (Avanti Polar

Lipids) in a buffer (20 mM HEPES-HCl pH 7.0, 300 mM NaCl, 5% (v/v) glycerol, and

0.1% (w/v) DDM) was rotated at 4 °C for 1 h. Next, the detergent was removed using

SM2-beads (Bio-Rad). The resulting solution was ultracentrifuged, and the precipitates

were suspended in a buffer (25 mM HEPES-NaOH pH 7.5 and 100 mM NaCl). The

reconstituted proteoliposome samples were flash-frozen in liquid nitrogen and stored at -

80 °C until measurements.

Measurement of ion transport activity

The SSM method was used to detect thiosulfate uptake activity of the *St*YeeE-
reconstituted proteoliposomes. Frozen proteoliposomes were thawed on ice and sonicated
using a bath sonicator (VELVO-CLEAR, VS-50R) for 10 s three times before use.
Measurement was performed using SURFE^{2}R N1 (Nanion Technologies) as described
(*14*). First, 50 μl of 0.5 mM 1-octadecanethiol (dissolved in isopropanol) was applied to
N1 Single Sensors (Nanion Technologies, Nr. 2-03-35002-000) and incubated for 30 min
at room temperature. Next, the sensors were washed twice with isopropanol, washed
twice with distilled water, and dried. Then, 1.5 μl of 7.5 μg/μl 1,2-diphytanoyl-sn-
glycero-3-phosphocholin (dissolved in n-decane) was applied to the sensor, followed by
50 μl of non-activating buffer (B buffer: 140 mM NaCl, 4 mM MgCl$_2$, 25 mM HEPES,
25 mM MES, KOH pH 6.7, with or without Na$_2$SO$_4$). Proteoliposomes were pipetted
toward the sensor beneath the B buffer surface, and the sensor was centrifuged (2,000*g*
for 30 min, 25 °C) to adsorb the liposomes onto the sensor surface. The resultant sensors
were mounted in SURFE^{2}R N1, and the sensors were rinsed with buffer B before each
measurement. The current change on the sensor was monitored while B buffer and A
buffer (140 mM NaCl, 4 mM MgCl$_2$, 25 mM HEPES, 25 mM MES, KOH pH 6.7, with
Na$_2$S$_2$O$_3$) were exchanged. For each condition, measurement was performed four times,
and the three results with least noise were analyzed.
646
MD simulations of *St*YeeE-*St*YeeD complex
All-atom MD simulations of the StYeeE-StYeeD complex were performed in
POPE/POPG mixed lipid bilayers and 150 mM NaCl solution, where two intermediate
states, *St*YeeE-*St*YeeD-SH and *St*YeeE-*St*YeeD-S-$S_2O_3^-$, were examined. The initial
structure of the protein–membrane complex was constructed using the CHARMM-GUI
membrane builder (*37*). The system size is 100 Å × 100 Å × 120 Å, and the total number
of atoms is ~111,600. The CHARMM C36m force field parameters were used for proteins
and lipids (*38*), and the topology and parameters for S-$S_2O_3^-$-modified Cys were derived
from the CGenFF parameters (*39*). The system was equilibrated using 2,000-step energy
minimization, followed by equilibration in the *NVT* and *NPT* ensembles for 3.5 ns using
the positional restraints on the proteins and lipids. The production run was then carried
out for 1 μs in the *NPT* ensemble at $T = 310$ K and $P = 1$ atm with the time step of 3.5 fs
using the RESPA integrator with hydrogen-mass repartitioning and bond-length
constraining (*40*) and the Bussi thermostat and barostat for temperature and pressure

control (*41*). The particle-mesh Ewald method was used for the calculation of long-range

electrostatic interaction (*42*). All simulations were performed using GENESIS (*43, 44*).

REFERENCES AND NOTES

1. A. Francioso, A. Baseggio Conrado, L. Mosca, M. Fontana, Chemistry and Biochemistry of Sulfur Natural Compounds: Key Intermediates of Metabolism and Redox Biology. *Oxid Med Cell Longev* **2020**, 8294158 (2020).

2. G. D. Westrop, G. Goodall, J. C. Mottram, G. H. Coombs, Cysteine biosynthesis in Trichomonas vaginalis involves cysteine synthase utilizing O-phosphoserine. *J Biol Chem* **281**, 25062-25075 (2006).

3. T. Nakatani *et al.*, Enhancement of thioredoxin/glutaredoxin-mediated L-cysteine synthesis from S-sulfocysteine increases L-cysteine production in Escherichia coli. *Microb Cell Fact* **11**, 62 (2012).

4. C. T. Pereira, C. Roesler, J. N. Faria, M. R. Fessel, A. Balan, Sulfate-Binding Protein (Sbp) from Xanthomonas citri: Structure and Functional Insights. *Mol Plant Microbe Interact* **30**, 578-588 (2017).

5. A. Sirko, M. Hryniewicz, D. Hulanicka, A. Bock, Sulfate and thiosulfate transport in Escherichia coli K-12: nucleotide sequence and expression of the cysTWAM gene cluster. *J Bacteriol* **172**, 3351-3357 (1990).

6. Y. Tanaka *et al.*, Crystal structure of a YeeE/YedE family protein engaged in thiosulfate uptake. *Sci Adv* **6**, eaba7637 (2020).

7. B. K. Mohanty, S. R. Kushner, Inactivation of RNase P in Escherichia coli significantly changes post-transcriptional RNA metabolism. *Mol Microbiol* **117**, 121-142 (2022).

8. S. Morigasaki, A. Umeyama, Y. Kawano, Y. Aizawa, I. Ohtsu, Defect of RNA pyrophosphohydrolase RppH enhances fermentative production of L-cysteine in Escherichia coli. *J Gen Appl Microbiol* **66**, 307-314 (2021).

9. T. Yamashino, M. Isomura, C. Ueguchi, T. Mizuno, The yhhP gene encoding a small ubiquitous protein is fundamental for normal cell growth of Escherichia coli. *J Bacteriol* **180**, 2257-2261 (1998).

10. Y. Ikeuchi, N. Shigi, J. Kato, A. Nishimura, T. Suzuki, Mechanistic insights into sulfur relay by multiple sulfur mediators involved in thiouridine biosynthesis at tRNA wobble positions. *Mol Cell* **21**, 97-108 (2006).

11.   J. U. Dahl *et al.*, The sulfur carrier protein TusA has a pleiotropic role in Escherichia coli that
also affects molybdenum cofactor biosynthesis. *J Biol Chem* **288**, 5426-5442 (2013).

12.   L. J. Liu *et al.*, Thiosulfate transfer mediated by DsrE/TusA homologs from acidothermophilic
sulfur-oxidizing archaeon Metallosphaera cuprina. *J Biol Chem* **289**, 26949-26959 (2014).

13.   T. S. Tanabe, S. Leimkuhler, C. Dahl, The functional diversity of the prokaryotic sulfur carrier
protein TusA. *Adv Microb Physiol* **75**, 233-277 (2019).

14.   A. Bazzone, M. Barthmes, K. Fendler, SSM-Based Electrophysiology for Transporter Research.
Methods Enzymol **594**, 31-83 (2017).

15.   M. Varadi *et al.*, AlphaFold Protein Structure Database: massively expanding the structural
coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439-
D444 (2022).

16.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-
589 (2021).

17.   E. Katoh *et al.*, High precision NMR structure of YhhP, a novel Escherichia coli protein
implicated in cell division. *J Mol Biol* **304**, 219-229 (2000).

18.   R. Shi *et al.*, Structural basis for Fe-S cluster assembly and tRNA thiolation mediated by IscS
protein-protein interactions. *PLoS Biol* **8**, e1000354 (2010).

19.   E. Doka *et al.*, Control of protein function through oxidation and reduction of persulfidated
states. *Sci Adv* **6**, eaax8358 (2020).

20.   G. Tombline *et al.*, Pseudomonas aeruginosa PA1006 is a persulfide-modified protein that is
critical for molybdenum homeostasis. *PLoS One* **8**, e55593 (2013).

21.   B. Ezraty, A. Gennaris, F. Barras, J. F. Collet, Oxidative stress, protein damage and repair in
bacteria. *Nat Rev Microbiol* **15**, 385-396 (2017).

22.   J. M. Kurth *et al.*, Electron Accepting Units of the Diheme Cytochrome c TsdA, a Bifunctional
Thiosulfate Dehydrogenase/Tetrathionate Reductase. *J Biol Chem* **291**, 24804-24818 (2016).

23.   Y. Abdiche, D. Malashock, A. Pinkerton, J. Pons, Determining kinetics and affinities of protein
interactions using a parallel real-time label-free biosensor, the Octet. *Anal Biochem* **377**, 209-
217 (2008).

24.   Y. Stockdreher *et al.*, New proteins involved in sulfur trafficking in the cytoplasm of
Allochromatium vinosum. *J Biol Chem* **289**, 12390-12403 (2014).

25.   P. Mitchell, J. Moyle, Group-translocation: a consequence of enzyme-catalysed group-transfer.
Nature **182**, 372-373 (1958).

26. Y. Daimon *et al.*, The TPR domain of BepA is required for productive interaction with substrate

proteins and the beta-barrel assembly machinery complex. *Mol Microbiol* **106**, 760-776 (2017).

27. R. B. Kapust *et al.*, Tobacco etch virus protease: mechanism of autolysis and rational design of

stable mutants with wild-type catalytic proficiency. *Protein Eng* **14**, 993-1000 (2001).

28. Y. Dai, R. M. Whittal, L. Li, Two-layer sample preparation: a method for MALDI-MS analysis

of complex peptide and protein mixtures. *Anal Chem* **71**, 1087-1091 (1999).

29. K. Hirata *et al.*, ZOO: an automatic data-collection system for high-throughput structure analysis

in protein microcrystallography. *Acta Crystallogr D Struct Biol* **75**, 138-150 (2019).

30. K. Yamashita, K. Hirata, M. Yamamoto, KAMO: towards automated data processing for

microcrystals. *Acta Crystallogr D Struct Biol* **74**, 441-449 (2018).

31. W. Kabsch, Xds. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132 (2010).

32. A. J. McCoy *et al.*, Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674 (2007).

33. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta

Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

34. P. V. Afonine *et al.*, Towards automated crystallographic structure refinement with

phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352-367 (2012).

35. A. Vagin, A. Teplyakov, Molecular replacement with MOLREP. *Acta Crystallogr D Biol

Crystallogr* **66**, 22-25 (2010).

36. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and

analysis. *J Comput Chem* **25**, 1605-1612 (2004).

37. E. L. Wu *et al.*, CHARMM-GUI Membrane Builder toward realistic biological membrane

simulations. *J Comput Chem* **35**, 1997-2004 (2014).

38. J. Huang *et al.*, CHARMM36m: an improved force field for folded and intrinsically disordered

proteins. *Nat Methods* **14**, 71-73 (2017).

39. K. Vanommeslaeghe *et al.*, CHARMM general force field: A force field for drug-like molecules

compatible with the CHARMM all-atom additive biological force fields. *J Comput Chem* **31**,

671-690 (2010).

40. J. Jung *et al.*, Optimized Hydrogen Mass Repartitioning Scheme Combined with Accurate

Temperature/Pressure Evaluations for Thermodynamic and Kinetic Properties of Biological

Systems. *J Chem Theory Comput* **17**, 5312-5321 (2021).

41. G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling. *J Chem

Phys* **126**, 014101 (2007).

42.   U. Essmann *et al.*, A smooth particle mesh Ewald method. *The Journal of Chemical Physics*
103, 8577-8593 (1995).

43.   J. Jung *et al.*, GENESIS: a hybrid-parallel and multi-scale molecular dynamics simulator with
enhanced sampling algorithms for biomolecular and cellular simulations. *Wiley Interdiscip Rev*
Comput Mol Sci **5**, 310-323 (2015).

44.   C. Kobayashi *et al.*, GENESIS 1.1: A hybrid-parallel molecular dynamics simulator with
enhanced sampling algorithms on multiple computational platforms. *J Comput Chem* **38**, 2193-
2206 (2017).

45.   X. Robert, P. Gouet, Deciphering key features in protein structures with the new ENDscript
server. *Nucleic Acids Res* **42**, W320-324 (2014).

Acknowledgments: We thank Kayo Abe for secretarial assistance, Akira Sasaki for
liposome preparation, Rie Kurata for performing mass spectrometry, Naoki Sakai for
helping with the data analysis, the beamline scientists at BL32XU of SPring-8 (Hyogo,
Japan) for helping with data collection, and Nanion Technologies for assistance with
initial data collection. The synchrotron radiation experiments were performed at BL32XU
of SPring-8 with the approval of the Japan Synchrotron Radiation Research Institute
(JASRI) (Proposal Nos. 2020A2564, 2021A2745, 2022A2738, 2023A2727). **Funding:**
This work was supported by JSPS/MEXT KAKENHI (Grant No. 21H05157 to T.M.,
Grant Nos. JP22K15075, JP20K15733 to M. Ichikawa, Grant No. JP22K15061,
JP22H05567 to R.M., and Grant Nos. JP22H02567, JP22H02586, JP21H05155,
JP21H05153, JP21K19226, JP21KK0125 to T.T.), MEXT as "Program for Promoting
Researches on the Supercomputer Fugaku" (Development and application of large-scale
simulation-based inferences for biomolecules JPMXP1020230119) and HPCI project
(hp200064 and hp230209) to T.M., PRESTO (JPMJPR20E1 to M. Ichikawa) from the
Japan Science and Technology Agency (JST), and private research foundations (the
Chemo-Sero-Therapeutic Research Institute, Naito Foundation, Takeda Science
Foundation, G-7 Scholarship Foundation, the Sumitomo Foundation, the Institute for
Fermentation (Osaka), Yamada Science Foundation, KOSÉ Cosmetology Research
Foundation, and Japan Foundation for Applied Enzymology) to T.T. *E. coli* JCM20135
was provided by RIKEN BRC, which is participating in the National Bio-Resources
Project of MEXT, Japan. **Author Contributions:** Conceptualization, A.T., M. Ichikawa,
and T.T.; Methodology and investigation, M. Ikei, R.M., K. Monden, Y.N., A.T., Y.S.T.,

Y.T., K. Murata, T.M., M. Ichikawa, and T.T.; Writing – original draft, M. Ichikawa and

T.T.; Writing, review & editing, M. Ichikawa and T.T.; Supervision, M. Ichikawa and

T.T. **Competing interests:** The authors declare no competing interests. **Data and**

materials availability: Coordinates and structure factors have been deposited in the

Protein Data Bank under accession number 8J4C and 8K1R. All data needed to evaluate

the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Additional data related to this paper may be requested from the authors.

FIGURES AND TABLES
Table 1. Data collection and refinement
Statistics for the highest-resolution shell are shown in parentheses.

PDB ID		8J4C	8K1R
Crystal		*St*YeeE(C22A)-*St*YeeD	*St*YeeE(C22A)-*St*YeeD(L45A)
Data Collection			
X-ray source		SPring-8 BL32XU	SPring-8 BL32XU
Wavelength (Å)		1.00	1.00
Space group		$P2_122_1$	$P2_122_1$
a, b, c (Å)		73.98, 102.44, 186.52	73.58, 101.75, 182.76
Resolution range (Å)		47.60–3.34 (3.46–3.34)	49.01–2.60 (2.69–2.60)
Total reflections		432750 (38159)	13803448 (1256177)
Unique reflections		21249 (2011)	42977 (4206)
Multiplicity		20.4 (18.4)	321.2 (297.1)
Completeness (%)		99.04 (96.87)	99.52 (99.32)
$I/\sigma(I)$		4.12 (0.83)	15.25 (2.10)
Wilson B-factor		49.19	31.95
R-pim		0.2207 (1.017)	0.0478 (0.2706)
$CC_{1/2}$		0.963 (0.578)	0.996 (0.861)
Refinement			
Reflections used in refinement		21070 (2011)	42846 (4206)
Reflections used for R-free		1990 (190)	1993 (196)
R_{work}		0.2654 (0.3596)	0.2064 (0.2705)
R_{free}		0.3095 (0.4113)	0.2547 (0.3108)
Number of non-hydrogen atoms		6675	6900
protein		6238	6247
thiosulfate		10	0
monoolein		427	545
water		0	108
RMS derivations	bond length (Å)	0.003	0.009
	bond angles (°)	0.56	1.16
MolProbity Validation	overall score	1.95	1.80
	clashscore	11.91	7.67
	rotamer outliers (%)	0.00	1.72
Ramachandran	favoured (%)	94.80	96.77
	allowed (%)	4.95	2.73
	outliers (%)	0.25	0.50
Average B-factor		66.46	39.54
protein		67.97	38.61
thiosulfate		73.25	-
monoolein		44.20	51.12
water		-	34.03

804 **Table 2. k_a, k_d, and K_D values of the interactions between YeeE and YeeD proteins**

	k_a (1/μM · s)	k_d (1/s)	K_D (μM)	
WT	0.017 ± 0.0029	0.027 ± 0.0029	1.6 ± 0.20	806
K8A	0.0089 ± 0.0011	0.037 ± 0.0019	4.3 ± 0.61	807
E9A	0.0060 ± 0.00058	0.027 ± 0.0021	4.6 ± 0.22	808
E15A	0.0075 ± 0.0013	0.071 ± 0.033	9.5 ± 4.2	809
V16A	0.017 ± 0.0016	0.093 ± 0.023	5.6 ± 0.86	810
C17A	*	*	*	
L38A	0.018 ± 0.0013	0.032 ± 0.0030	1.7 ± 0.055	811
V39A	0.0058 ± 0.00033	0.032 ± 0.00058	5.4 ± 0.30	812
D42A	**	**	**	813
Y43A	0.0082 ± 0.00095	0.16 ± 0.017	20 ± 2.1	
L45A	0.029 ± 0.0016	0.035 ± 0.0040	1.2 ± 0.10	814
E48A	0.017 ± 0.010	0.14 ± 0.20	4.5 ± 6.2	815
R49A	0.0090 ± 0.0014	0.12 ± 0.16	17 ± 23	
E63A	0.018 ± 0.00026	0.030 ± 0.0013	1.7 ± 0.074	816

818 *: The affinity of YeeD C17A derivative for YeeE was too low, and the value could not be

819 calculated accurately.

820 **: The YeeD D42A sample contained a significant amount of intermolecular S-S dimers (fig.

821 S2F); therefore, the value was omitted here.

Fig. 1. Thiosulfate ion transport activity of *St*YeeE-reconstituted liposomes.

(**A**) Selective uptake of thiosulfate ion by *St*YeeE. *St*YeeE-reconstituted liposomes were

used for the measurement using the SSM method. The non-activation buffer, lacking

any sulfur-related ions, was exchanged with buffer containing either 20 mM $Na_2S_2O_3$ or

20 mM Na_2SO_4 at 0.0 s. (**B**) YeeE-dependent negative current changes. Liposomes with

and without *St*YeeE were used. The non-activation buffer containing 20 mM Na_2SO_4

was exchanged with buffer containing 20 mM $Na_2S_2O_3$ instead of Na_2SO_4 at 0.0 s. The

current values from three measurements were standardized based on the average values

from -0.05 to 0.0 s, and then the mean values from the three measurements were

calculated. Error bars indicate the SD (standard deviation). (**C**) Thiosulfate

concentration dependency of the uptake activity. The current changes were monitored in

the same manner as in (B), except for the concentration of thiosulfate and sulfate ions

(1.25, 2.5, 5, 10, 20, and 40 mM). The negative peak area values (nA・s) from three

measurements were calculated and plotted with the SD. The plot was fitted with

Michaelis–Menten equation, and V_{max} was estimated as 0.01184 nA・s and K_m as 0.6261

mM.

Fig. 2. Characterization of YeeD functions in vivo and in vitro.

(**A**) Sequence alignment of *Ec*YeeD and *St*YeeD. The regions of α-helices and β-strands

predicted by AlphaFold2 in (B) are highlighted with colors. Red text indicates amino acid

residues that were mutated in this study. The conserved CPxP motif is enclosed in a red

rectangle. (**B**) Cartoon model of AlphaFold2-predicted structure of *Ec*YeeD. α-helices

and β-strands are colored as in (A). The side chains of amino acid residues mutated for

growth complementation analysis in (C) are shown as stick models. Corresponding amino

acid residues from *St*YeeD are also indicated in parentheses. (**C**) Growth

complementation assay of Δ*cysPUWA* Δ*yeeE* (DE3) cells, depending on *Ec*YeeE and

*Ec*YeeD expression from plasmids. The mean values of increased OD_{600} (ΔOD_{600}) after

24 h are shown. Error bars represent the SD from three measurements. Statistical

significance compared with the Δ*cysPUWA* Δ*yeeE* (DE3) cells possessing an empty

vector was determined using one-way analysis of variance (ANOVA) followed by

Dunnett's multiple comparisons tests (****, $p < 0.0001$; ns, not significant). (**D**) CBB-

stained SDS-PAGE gel and mass spectrometry of purified *St*YeeD(WT). Red and black

arrowheads indicate the major and minor bands of purified WT *St*YeeD, respectively. In

mass spectrometry, three major peaks corresponding to -SH, -S-O_2^-/-S-S^-, and -S-SO_3^-

were detected. (**E**) CBB-stained SDS-PAGE gel and mass spectrometry of purified

*St*YeeD(C17A) (upper) and 500 μM thiosulfate-treated *St*YeeD(C17A) (lower). In mass

spectrometry, a single peak was observed. (**F**) Mass spectrometry of DTT-reduced

*St*YeeD(WT). (**G**) Mass spectrometry of 500 μM thiosulfate-treated *St*YeeD after DTT-

reduction. (**H**) Mass spectrometry of *St*YeeD oxidized by hydrogen peroxide after DTT-

reduction. (**I**) Measurement of in vitro enzymatic activity of *St*YeeD. Fluorescence

intensity changes derived from an H_2S-detectable fluorescent probe, HSip-1, in the

presence of thiosulfate (500 μM) were measured using a series of purified *St*YeeD

derivatives in 10 mM Tris-HCl pH 8.0, 300 mM NaCl, and 70 μM β-ME. Mean values

of three experiments are shown. Error bars represent SD. Statistical significance

compared with samples containing no protein was determined using one-way ANOVA

followed by Dunnett's multiple comparisons tests (***, $p < 0.001$; ****, $p < 0.0001$; ns,

not significant). Dashed lines indicate the mean values of no-protein samples and

*St*YeeD(WT).

Fig. 3. Crystal structure of *St*YeeE-*St*YeeD complex (PDB ID 8K1R).
(A) Cartoon model of *St*YeeE(C22A)-YeeD(L45A). The side chains of amino acid
residues on the contact surfaces between *St*YeeE and *St*YeeD are shown as stick
models. The side chains of the conserved cysteine positions are shown in the spheres
model. The thiosulfate ion in the *St*YeeE(C22A)-YeeD model (PDB ID 8J4C) was
superimposed as spheres model. (**B**) Close-up view of the contact region. (**C**) 2Fo-Fc
electron density map at 1.5 δ of the border region around C22A in *St*YeeE and C17 in
*St*YeeD. Position III is a previously proposed thiosulfate-binding site. The red stars
indicate water molecules. (**D**) 2Fo-Fc electron density map at 1.3 δ around the CPxP
motif. The red arrowhead indicates where electron density map is faint.

Fig. 4. Interaction between *St*YeeE and *St*YeeD.

(**A**) Real-time detection by the BLI method of association and dissociation of *St*YeeE with/from solidified *St*YeeD and its derivatives with Octet buffer (20 mM Tris-HCl pH 8.0, 300 mM NaCl, 0.1% (w/v) DDM, 1 mM β-ME) Each line shows the mean value of three measurements. The dashed line indicates the 120-s point where *St*YeeD-solidified biosensors were submerged in the YeeE-free buffer. (**B**) Mean values of *St*YeeE binding to solidified *St*YeeD at 120 s. Error bars show the SD of three measurements. The dashed line indicates the mean value of *St*YeeD(WT). (**C**) Mutation site mapping on the crystal structure of *St*YeeE-YeeD complex (PDB ID 8J4C). The side chains of mutated amino acid residues in (A) are shown as stick models. Magenta, blue, and red residues indicate that those Ala-substituted mutants showed increased, similar, and reduced binding affinity, respectively, compared to WT.

Fig. 5. Model of uptake and decomposition of thiosulfate ion by YeeE and YeeD.

(**A**) Uptake of a thiosulfate ion by YeeE and YeeD. The thiosulfate ion is relayed by conserved cysteine residues of YeeE and binds to the conserved cysteine residue of YeeD. After that, the association of YeeD with YeeE is destabilized. YeeD then dissociates from YeeE and decomposes the thiosulfate. (**B**) Thiosulfate decomposition by YeeD. First, thiosulfate binds directly to the conserved cysteine residue of YeeD. Next, sulfide ion (SO_3^{2-}) is released. Finally, sulfite ion (S^{2-}) is released. In the blue box, a putative dimerization model of *St*YeeD is shown.