Dynamic simulations of feeding and respiration of the early Cambrian periderm-bearing cnidarian polyps

Yiheng Zhang[1,2,4], Xing Wang[3,4,*], Jian Han[2,*] Juyue Xiao[1], Yuanyuan Yong[2], Chiyang Yu[2], Ning Yue[2], Jie Sun[2], Kaiyue He[2], Wenjing Hao[2], Tao Zhang[1,2,*], Bin Wang[1], Deng Wang[2], *and* Xiaoguang Yang[2]

1. *School of Information Science & Technology, Northwest University, Xi'an* 710069, *China.*
2. *State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life and Environments, Department of Geology, Northwest University, Xi'an* 710069, *China.*
3. *College of Life Science, Linyi University, Linyi* 276000, *China.*
4. *These authors contributed equally to this work.*
*Corresponding authors: Jian Han (elihanj@nwu.edu.cn); Tao Zhang (zhangtao129@nwu.edu.cn) and Xing Wang (wx5432183@126.com)

Impact statement

To the best of our knowledge, our study is the first to simulate the feeding and respiration patterns of *Quadrapyrgites* by introducing a dynamic fluid-structure coupling method, revealing the relationship between the ancient medusozoans and modern jellyfish.

Abstract

Although fossil evidence suggests the existence of an early muscular system in the ancient cnidarian jellyfish from the early Cambrian Kuanchuanpu biota (*ca.* 535 Ma), south China, the mechanisms underlying the feeding and respiration of the early jellyfish are conjectural. Recently, the polyp inside the periderm of olivooids was demonstrated to be a calyx-like structure, most likely bearing short tentacles and bundles of coronal muscles at the edge of the calyx, thus presumably contributing to feeding and respiration. Here, we simulate the contraction and expansion of the microscopic periderm-bearing olivooid *Quadrapyrgites* via the fluid-structure interaction computational fluid dynamics (CFD) method to investigate their feeding and respiratory activities. The simulations show that the rate of water inhalation by the polyp subumbrella is positively correlated with the rate of contraction and expansion of the coronal muscles, consistent with the previous feeding and respiration hypothesis. The dynamic simulations also show that the frequent inhalation/exhalation of water through the periderm polyp expansion/contraction conducted by the muscular system of *Quadrapyrgites* most likely represents the ancestral feeding and respiration patterns of Cambrian sedentary medusozoans that predated the rhythmic jet-propelled swimming of the modern jellyfish. Most importantly for these Cambrian microscopic sedentary medusozoans, the increase of body size and stronger capacity of muscle contraction may have been indispensable in the stepwise evolution of active feeding and subsequent swimming in a higher flow (or higher Reynolds number) environment.

Introduction

Introduction

Cnidarians, such as medusozoans, corals, sea fans, and hydromedusae, are generally considered to
be a sister group of bilateral animals that live predominantly in the ocean. In general,
medusozoans have a two-stage life cycle, consisting of a swimming medusoid stage and a
sedentary polypoid stage. Swimming jellyfish rely on rhythmic contraction of the coronal muscles
and expansion of the mesoglea at the umbrella rim to swim in a "jet-like" manner in the water
column (Arai 1997; Brusca *et al.* 2016; Leclère & Röttinger 2016; Zapata *et al.* 2015). Because
both the ectoderm and endoderm of jellyfish are in direct contact with seawater, no specialised
respiratory organs are required to meet aerobic metabolic needs. Sedentary polyps rely on free,
extensible tentacles and can feed actively or passively (Ishi 2008).

Although the origin of the common ancestor of medusozoans was dated using the molecular
clock technique to the Cryogenian (Erwin *et al.* 2011), the earliest known medusozoan fossil
records were found in the Ediacaran. *Haootia quadriformis* from the Ediacaran Fermeuse
Formation (*ca.* 560 Ma), was suggested to be a stalked jellyfish based on external morphological
evidence and possible coronal muscles on its surface (Liu *et al.* 2014). By comparing periderm
morphology, several taxa of tetradial conulariids in the Ediacaran were also proposed to be more
closely related to modern scyphozoan polyps (Van Iten *et al.* 2006; Leme *et al.* 2022). All these
fossil types were suggested to be sedentary forms, with no definitive evidence of a free-swimming
lifestyle. The earliest known swimming jellyfish, *Yunnanoascus haikouensis* Hu *et al.* (2007),
from the early Cambrian Chengjiang biota (Stage 3, *ca.* 519 Ma) exhibits a typical tetraradial
symmetry; eight sensory rods distributed around the umbrella rim, 16 pairs of elongated
retractable tentacles evenly spaced with rhopalias, and a less pronounced manubrium. This
configuration allows direct comparison with modern scyphozoans (Han *et al.* 2007; 2016) and
suggests that the origin of swimming jellyfish may have occurred much earlier.

Phosphatised microfossil medusozoans from the early Cambrian Kuanchuanpu biota (*ca.* 535
Ma) provide critical clues for investigating the origin and evolution of cnidarians and swimming
medusae. At least four families have been identified in the Kuanchuanpu biota, involving
Hexangulaconulariidae, Anabaritidae, Carinachitidae, and Olivooidae (Han *et al.* 2020). Of these,
the family Olivooidae includes the tetraradial symmetrical *Quadrapyrgites* as well as multiple
pentaradial forms, such as *Olivooides*, *Sinaster*, *Hanagyroia*, and other undetermined taxa (Li *et al.*
2007; Liu *et al.* 2014; Han *et al.* 2013, 2016; Dong *et al.* 2013, 2016; Wang *et al.* 2017, 2020).

As revealed by scanning electron microscopy (SEM), the hemi-globular shaped embryo of
Olivooides (Figure 1), enclosed by an egg membrane, is equipped with very complex internal
structures at the prehatched embryonic stage, such as a manubrium in a relatively deep
subumbrella cavity, short tentacles, bundles of coronal muscles, paired gonad-like lamellae at
either side of the interradial septa, and many other sheet-like lamellae (Han *et al.* 2013; 2016;
Dong *et al.* 2013; Wang *et al.* 2017; 2020). Remarkably, the regular distribution of ring-like
fibrous structures on the surface of the umbrella of their prehatched embryos, which are densely
packed in bundles at the edge of the subumbrella and gradually become sparse towards the aboral
side, allows for a comparison with the coronal muscles of modern jellyfish (Han *et al.* 2020; Wang
et al. 2022).

In the hatched stages of development, the soft tissue of millimetre-scale olivooids consists of
an upper calyx and basal stalk (Figure 1G), in a torch-like shape similar to that of the extant

medusozoan polyps (Wang *et al.* 2020; Steiner *et al.* 2014). Unfortunately, the internal structure of

polyps remains ambiguous. Considering their cubomedusa-type anatomy in the prehatched stage,

the hatched torch shaped olivooids appear to be a type of periderm-bearing polyp-shaped medusa

(Wang *et al.* 2020). As mentioned above, the rhythmic contraction and expansion of the coronal

muscles aided by the mesoglea leads to the consequent inhalation and discharge of water,

propelling modern jellyfish to swim through the water column, which in turn facilitates more

efficient tentacle feeding (Brusca *et al.* 2016). The presence of coronal muscles in early Cambrian

embryonic olivooids suggests that early medusozoans in the hatched/juvenile polypoid stage

probably used a similar muscular system to control the opening and closing of the subumbrella to

drive the water flow in and out, and to assist in feeding (Wang *et al.* 2022); however, the early

Cambrian polyp-shaped medusa most likely could not swim freely due to the external periderm.

The compromise of these two seemingly conflicting conditions led to the hypothesis on the

benthic origin of medusa swimming and feeding (Han *et al.* 2020; Wang *et al.* 2022). Specifically,

there may be a series of intermediate types (*i.e.*, stalked jellyfish) between sedentary polyps and

free-swimming medusae. These transitional types may have evolved divergently from sedentary

forms to free-swimming medusae through a series of morphological and structural innovations in

evolution, such as rhythmic contraction of the coronal muscles, loss or degradation of the

periderm, and increased thickness of the mesogleal layer (Han *et al.* 2020; Wang *et al.* 2022).

Quadrapyrgites (Figure 2A, B) are one of the most recognisable taxa in the olivooids from

the Kuanchuanpu biota (Fortunian Stage, early Cambrian). It has drawn much attention from

biologists and palaeontologists, given its tetra-radial symmetry comparable to that of modern

jellyfish (Liu *et al.* 2014; Dzik *et al.* 2017). The pagoda-shaped, thin, flexible periderm was

divisible into basal and an abapical sections. The abapical section showed an increasing number of

annular ridges as it grew. The surface of an annular ridge exhibits many irregular longitudinal

folds and fine striations (Yong *et al.* 2022). Similar to the type of pentaradial forms found in

olivooids (Steiner *et al.* 2014, Figures 10.3, 11.6, and 11.13), the twelve longitudinal apertural

lobes of *Quadrapyrgites* converge towards the central axis of the periderm and then extend

downwards, leaving a narrow, star-shaped but contractile channel, which is called the periderm

aperture (Figure 2B). The upper side of the polyp calyx was bound to the periderm aperture

(Figure 2C). The manubrium in the subumbrella cavity was conceived with a mouth at the top. A

ring of four pairs of short tentacles was possibly located close to the subumbrella margin, as

inferred from other contemporaneous tetraradial olivooid embryos (Figure 2D, Han *et al.* 2016*b*).

The varying heights and expansions of the peridermal aperture of *Quadrapyrgites* with the twelve

centripetal lobes (Figure 2A–C) indicate the peridermal aperture could move up and down along

the body axis, and expand centrifugally or contract centripetally, a behaviour that was undoubtedly

triggered by the interaction of circular and longitudinal muscles and the mesoglea of the polyp

inside the periderm. To date, there is no evidence to support that the tentacles of *Quadrapyrgites*

could protrude from the periderm to feed in the same way as modern scyphopolyps.

Modelling the living environment of macrofossils to verify their morphological and

functional roles is one of the most recent advances in paleobiology (Dynowski *et al.* 2016;

Darroch *et al.* 2017; Waters *et al.* 2017; Gibson *et al.* 2019; Rahman *et al.* 2020; Song *et al.* 2021).

For example, computational fluid dynamics (CFD) methods were used to simulate oral feeding in

the Ediacaran *Tribrachidium heraldicum*, demonstrating that its oral morphology was more

oriented towards suspension filter feeding, providing evidence for late Ediacaran ecosystem

complexity (Rahman *et al.* 2015). Our recent findings suggest that fluid simulation tools can also
be used for microfossil morphological and functional studies (Liu *et al.* 2022). Additionally,
compared with macrofossil fluid simulations, the boundary layer conditions should be considered
(Zhang *et al.* 2022). Although the swimming mechanism of modern jellyfish has long been studied
by biological modelling and the use of fluid simulations (Sahin *et.al.* 2009; Gemmell *et al.* 2013;
Gemmell *et al.* 2018), to the best of our knowledge, such methodologies have hardly been applied
to modelling the dynamic pattern of Cambrian sedentary polyps.
In the present study, we attempted to simulate both the contraction of the coronoid muscle of
the subumbrella and the expansion of the mesoglea layer of structurally simplified polyps of
Quadrapyrgites using a fluid-structure interaction CFD (Figure 2). Thus, we were able to
reconstruct and investigate the active dynamic pattern of *Quadrapyrgites*. It is also possible to
further probe into the autecology of more microscopic Cambrian sedentary periderm-bearing
polyps.
144
Results
Flow velocity
The velocity line profiles for the simulations with different expansion/contraction time ratios show
that the velocities in the contraction phase of the four sets of simulations had almost the same
trend with time (Table supplement 3). During the expansion phase, the maximum values of the
mouth flow velocity for all four sets of simulations increased as the expansion velocity increased,
with the maximum values occurring to the right of the centre of the time axis (Figure 4A–D). By
comparing the maximum values of the flow velocities at the sampling cut points in each
simulation (Figure 5), the accelerated expansion velocity lead to a more remarkable change in
flow velocity within the region of z from 2.05 to 2.15 mm than that within the region of z>2.15
mm. Since the flow velocity dropped to below 0.001 m/s, suggesting that the polyp subumbrella
had a reduced capacity to take in food from this region.
In all simulations, the general trend of the flow changed with different expansion/contraction
time ratios; however, only the maximum values of the flow velocities differed. We considered the
results of the simulation with an expansion/contraction time ratio of 3:1 as an example. The
subumbrella inside the polyp was in a resting state at 0 s (Figure 6A, B), the flow velocity in the
flow field was close to 0 m/s, and the opening of the subumbrella was in an expanded state. After
0 s, the subumbrella started to expand, and the external water flow was sucked in. Then, the
expansion velocity of the subumbrella gradually decreased with time until around 3 s, when the
flow velocity near the opening of the subumbrella became 0 m/s and the mouth shrank to a
minimum (Figure 6E, F). After 3 s, the subumbrella started to contract, accelerating the
contraction, and the opening of the subumbrella began to be restored. At approximately 3.5 s, the
contraction velocity of the mesoglea layer reached a maximum, and the flow velocity near the
opening of the subumbrella also reached a maximum during the contraction phase (Figure 6G, H).
At 4 s, the subumbrella stopped contracting, the opening of the subumbrella was restored to its
original state, and the flow velocity decreased to a minimum. At this point, the polyp completed
the expansion/contraction cycle.
172
Vortex visualisation
The visualisation of the intensity of the dimensionless vorticity around *Quadrapyrgites* during its

expansion and contraction phases is shown in Figure 7-animation supplement 1 (dynamic 2D
visualisation and 3D vortex reconstruction in Animation supplement 1 and 2, respectively). The
magnitude of vorticity of the colour scale bar was set to [–0.001,0.001] so that the magnitude was
appropriate for visualising vortex formation near the periderm. At approximately 0.1 s (Figure 7A),
the main vortex started to form near the peridermal aperture, and a secondary vortex, which
flowed in the opposite direction to the main vortex, also started to form in the middle portion of
the periderm. Both the main vortex and secondary vortex enlarged gradually over time. At
approximately 1 s, the lower secondary vortex was in contact with the bottom surface (Figure 7B),
and its morphology changed, manifesting itself as a flow from the middle of the periderm towards
the bottom surface. At approximately 2 s, the main vortex developed to the maximum visualisation
range (Figure 7C), at which time, due to an increase in the velocity of the water, a partial
microflow in the opposite direction of the vortex was also formed on the surface of the periderm.
After 2 s, the secondary vortex began to move up along the surface of the periderm to a position
close to the aperture. From 2.87–2.88 s (Figure 7D, E), the main vortex separated from the
periderm, and the secondary vortex moved to the original position of the main vortex at the
peridermal aperture, replacing it with the main vortex for the next stage of contraction at the end
of the expansion movement of the subumbrella at 3 s (Figure 7F).

After the onset of contraction, the newly formed main vortex pushed the expansion phase
vortex away from the peridermal aperture. The vortex development process in the contraction
phase follows a pattern similar to that in the expansion phase, with the vortex eventually
separating from the peridermal aperture at 3.9–4.0 s (Figure 7G, H), at which point the
expansion/contraction cycle was completed.

Because the contraction process took less time and the vortex enlarged faster than that in the
expansion phase, the shapes and maximum sizes of the newly formed main vortex and secondary
vortex differed, but the overall trend of alternating main vortex and secondary vortex formation
was maintained during the expansion-contraction-expansion movement of the subumbrella.

Discussion

Expansion/contraction frequency and feeding and respiration efficiencies
The simulations demonstrated the dynamic pattern of *Quadrapyrgites* during one cycle of
expansion and contraction and the visualisation of ambient water flow in its vicinity. A faster
expansion rate (*i.e.*, a shorter expansion-contraction cycle) leads to a relatively greater water
exchange and flow velocity formed near the subumbrella aperture. Subsequently, the tentacles will
have more opportunities to make contact with suspended food particles in fresh water inputted
from outside the periderm per unit of time. In this regard, compared with the stagnant condition,
the increased velocity of the water flowing into the periderm and then the subumbrella cavity will
improve the efficiency of food intake. In the subsequent contraction phase, the polyp expels water
from the subumbrella cavity at a high rate of movement.

During the contraction/expansion movement of the polyp, the vortices formed around the
periderm could slowly bring food particles close to the periderm aperture, where small food
particles were more likely to gather owing to the viscous force of the peridermal surface instead of
being transported away by the current. This combination of active and passive feeding could also
improve the feeding and gas exchange efficiencies of polyps. Notably, although the deeply

concave subumbrellar cavity of olivooids is equipped with short tentacles, it is unlikely that they
can protrude their short tentacles out of the periderm. Therefore, except for the random passive
flow of microscopic food particles, their main mode of food intake is likely suspension feeding
through active contraction by the ring muscles (Wang *et al.* 2022). The relatively high rate of
contraction of *Quadrapyrgites/*olivooids, if contracted frequently or rhythmically, helps the
four/five pairs of short tentacles capture relatively larger quantities of food particles in a short
period of time. In this regard, olivooids can be functionally considered as active suspension
feeders rather than conventional predators.

The feeding and excretory activities of marine benthic organisms are closely related to their
surrounding water as an appropriate flow velocity can increase feeding efficiency (Pratt 2008).
Active suspension feeders with the ability to move can actively avoid areas where currents are
unsuitable for survival and choose areas where food is abundant (Labarbera 1984). In contrast, in
sedentary forms, olivooids were unable to displace directionally and actively; therefore, their
feeding was more dependent on the surrounding environments. A suitable current environment is
conducive to the formation of eddies around the periderm to inhale food particles efficiently from
the space above the subumbrella aperture, whereas excessive current velocities may take away
large amounts of food particles.

Potential influence of current velocity on feeding patterns
It has been suggested that microbenthos mostly inhabited in the low flow region above the sea

floor, also called the bottom boundary layer (Trowbridge & Lentz 2018). The flow velocity above

the *Quadrapyrgites* peridermal aperture showed that the maximum velocity ranged from 0.005–
0.0155 m/s depending on the rate of contraction and expansion (Figure 5). Such a flow velocity
magnitude indicated that polyps may live in relative low flow environment protected by the
viscous boundary layer, which enable them to maintain a relatively stable posture for feeding
activities in a lower ambient flow speed environment (Liu *et al.* 2022). Otherwise, inhaling food
from the ambient environment with a current velocity much higher than the inhaling velocity is
more difficult.

However, the protective ability of the viscous boundary layer is varied because of the
fluctuations of flow caused by oscillating current or uneven seafloor (Zhang *et al.* 2022).
Simultaneously, the current velocity (at a height of 2–2.5 mm) in the vicinity of the peridermal
aperture within the viscous boundary layer might be close to 0.03 m/s (Caldwell & Chriss 1979;
Wengrove & Foster 2014), about twice the maximum inhalation velocity of the polyp in the
simulations, especially in areas with frequent intertidal current velocity changes (Wengrove &
Foster 2014). Under these relatively high current velocities, food particles will be rapidly
transported, making it difficult for polyps to inhale them efficiently. Thus, it can be hypothesised
that the physiological activity of *Quadrapyrgites* relies on other potential environmental factors in
addition to the protectiveness of the viscous boundary layer. For example, the greater viscous
effect of rougher sediment surfaces or the enrichment of organisms in the area results in more
turbulence among the community, increasing the thickness of the boundary layer (Grant &
Madsen 1986), and simultaneously reducing the current velocity. A turbulent environment can
enhance the mixing of nutrients, thereby increasing the feeding efficiency (Denny 2014), but the
effectiveness requires further investigation. In addition, this type of inhalation feeding is

profoundly dependent on the surrounding environment and may be an important constraint in
survival.
The pattern of alternating ring-shaped main and secondary vortices formed during the
expansion and contraction of polyps (Figure 7) shares some similarities with those in modern
swimming medusae. Modern medusae swimming patterns can be divided into jet propulsion (jet
propulsive force is generated by the contraction of the circular muscle fibres) and jet-paddling
propulsion (the edge of the subumbrella can act as "paddles" to assist jet propulsion) (Dabiri *et al.*
2007). The dye visualisation on *Aurelia aurita* showed a ring-shaped starting vortex generated
during the swimming stroke (contraction), followed by a stopping vortex of opposite rotational
sense during the recovery stroke (expansion) (Dabiri *et al.* 2005). The vortex formed was
suggested to be related to both feeding and propulsion. Sahin *et al.* (2009) recovered the
swimming mode of hydromedusae *Sarsia tubulosa* (jet propulsion) and *Aequorea victoria* (jet-
paddling propulsion) using a CFD method and visualised the ring-shaped starting and stopping
vortices. The flow pattern of a vortex can be analysed to estimate the efficiency at which a jet
produces thrust (Sahin *et al.* 2009). Dabiri *et al.* (2005) suggested that the interaction between the
starting and stopping vortex functions to reduce the kinetic energy was lost during medusa
swimming, while Sahin *et al.* (2009) suggested that the formation of toroidal vortex rings in the
wake of medusa swimming does that. Gemmell *et al.* also suggested that the starting and stopping
vortices may be related to the passive energy recapture mechanism, which is supported by the fact
that *Aurelia aurita* is one of the most energetically efficient propulsors (Gemmell *et al.* 2013;
2017). Although the dynamic process of sedentary *Quadrapyrgites* is more or less similar to the jet
propulsion swimming of medusae, they may not stroke to move or swim concerning the external
periderm. The formed main and secondary vortices are presumed to improve the efficiency of
water flow in and out of the polyp cavity, hence influencing the feeding and respiration efficiency.
Furthermore, the contraction (or changeable diameter) of the subumbrella opening can bolster the
kinetic energy of expelled water and the vortex (Mohseni & Gharib 1998; Mohseni *et al.* 2001),
which may also help them gain extra feeding and respiration efficiency. However, jet propulsion is
significantly less efficient than jet-paddling propulsion swimming patterns, although the former
can produce relatively higher thrust and kinetic energy in most cases. Constrained by peridermal
structure, the *Quadrapyrgites* polyps were unable to expand and contract to a large magnitude as
modern medusae or hydrae do, thus their dynamic efficiency is considered to be rather limited,
although this cannot be calculated by using formulas for modern swimming animals (*e.g.*, net cost
of transport analysis proposed in Gemmell *et al.* 2013) due to their sedentary life and the
ambiguity of certain truly essential parameters (*e.g.*, net mass of polyps).
296
Functions of muscular system in post-embryonic olivooids
Extant anthozoans, and possibly, medusozoan polyps (Arai, 1997), are exclusively passive
opportunists in feeding (Shick, 1991), minimising the energetic cost of obtaining food. In contrast,
jellyfish are undoubtedly active feeders (Arai, 1997). The earliest Cambrian medusozoan
olivooids, although encased in the periderm, may be able to switch between active and passive
feeding modes as modern benthic organisms. In an energetic tidal setting with a high
concentration of oxygen, despite the low flow due to the bottom boundary layer, olivooids can
presumably stay inside the periderm, waiting for the suspension of food particles transported by
the bimodal current or absorbing dissolved organic matter. Morpho-anatomical studies suggest

that the coronal muscles of the pentaradial symmetrical medusozoan embryos from the early
Cambrian Kuanchuanpu biota during the pre-hatching stage are perfectly comparable to those of
modern medusae, and it was further hypothesised that the contraction of the coronal muscles of
the soft tissue and the expansion of the mesoglea layer guided the opening of the periderm,
thereby facilitating animal feeding (Wang et al. 2022). Taken together with CFD simulation, it
suggested that Cambrian olivooids could also be active feeders and that they may have developed
some of the behavioural capabilities of swimming medusae.
313
Body size, Reynolds number, and swimming
Previous studies have revealed that some Cambrian echinoderms may adopt a more favourable
posture in relation to the current to reduce drag or create a suitable recirculation environment to
improve feeding efficiency (Rahman *et al.* 2020). Olivooids were unable to do so because of their
sedentary life within a tetraradial symmetrical periderm. However, their periderm aperture will
stand upright or tilt at various angles. As polyps and their periderm became larger or taller, due to
growth or evolution, the upper portion of the periderm left the bottom boundary layer region
(Zhang *et al.* 2022) and encountered higher velocity currents. Therefore, larger polyps have to
develop a series of novel strategies to adhere to the seafloor by a holdfast and to access more
suspended food particles per unit time transported by higher velocity currents (Shick, 1991). One
of the strategies for periderm-dwelling medusozoans is to develop stronger coronal muscles to
achieve a greater contraction or expansion ability (*i.e.*, higher rate or larger magnitude for
contraction and expansion), so as to change the direction of flow to a greater extent. Alternatively,
they may have evolved many longer, extensible tentacles protruding from the periderm, which
should result in a larger periderm aperture even without a cover, similar to living scyphopolyps
(Jarms *et al.* 2002). Third, if the periderm largely degenerated, such as in living sea anemone or
cubozoan polyps (Straehler-Pohl 2017), the polyps could change their adhesive position.
Otherwise, they could develop a streamlined body to reduce drag and allow for more favourable
feeding gestures (Liu *et al.* 2022).
Furthermore, from the aspect of biological fluid mechanics, a low Reynolds number (*Re*)
swimming strategy can lead to a lower propulsion efficiency in modern propulsive swimming
animals, such as swimming medusae (Sahin *et al.* 2009). In certain cases, animals are unable to
move forward if the swimming *Re* is too low; this is called "the scallop theorem" (Robertson *et al.*
2019). Specifically, this implies the medusae may evolve to much larger sizes and still be able to
move through the surrounding fluid (Sahin *et al.* 2009). For modern scyphomedusae, such as
Aurelia aurita, an increase in body size (or bell diameter) allowed them to swim in a higher *Re*
manner (Feitl *et al.* 2009). Simultaneously, for millimetre-scale olivooids inhabiting the viscous
boundary layer region, both the dynamic *Re* (approximately 1; calculated from the results in this
study, see "Computational fluid dynamics") and the environmental *Re* could be very low (due to
the higher fluid viscosity caused by enhanced mixing of sediments and lower flow), the limitation
to the efficiency of food inhalation and respiration through contraction and expansion may also be
significant. Thus, it is reasonable that both larger body size and stronger capacity of body
contraction of Cambrian polyps may have been indispensable toward the stepwise evolution of
active feeding, enabling them to live and subsequently swim in a higher flow (or higher *Re*)
environment.
Notably, from the earliest Cambrian to Cambrian Stage 3, the trend of increasing body size

was remarkable on a global scale (Zhuravlev & Wood 2020), although with some exceptions, *i.e.*,
the large, skeletonised conulariid-like *Paraconularia* found in the terminal Ediacaran Tamengo
Formation of Brazil (Leme *et al.* 2022). For example, in South China, millimetre-scale, sedentary
medusozoan polyps from the Cambrian Fortunian Stage to Stage 2, except for the tubulous
Anabaritidae without tube closure (Liu *et al.* 2017; Guo *et al.* 2020a, b; 2021), have small
peridermal apertures. For polyps living inside the periderm, access to food is relatively inefficient,
and food particles are limited in size by virtue of contraction and expansion of the peridermal
aperture and the coronal muscle-mesoglea layer. In contrast, centimetric polyp-type sedentary
tube-dwelling cnidarian fossils such as *Sphenothallus* (Li *et al.* 2004), *Cambrorhytium* (Conway
Morris *et al.* 2015), and *Byronia* (Zhu *et al.* 2000; Chang *et al.* 2018) from the mid- to late
Cambrian allow their tentacles to protrude completely out of the cone-shaped closureless tube. In
addition, the tube surface was much smoother, capable of reducing the drag-to-water flow. This
body structure, together with protruded tentacles, allowed for larger amounts and sizes of food to
be obtained, thus establishing the evolutionary foundation for the rise of polyp strobilation, the
emergence of saucer-like planula larvae, and the origin of swimming medusae. Changes in the
periderm/exoskeleton of Cambrian benthic medusozoans were also consistent with our hypothesis
regarding the interaction between animal body size and current velocity (or Reynolds number).

In summary, the simulation results illustrate that the rate of water intake near the periderm
aperture is directly related to the expansion rate of the mesoglea layer. Increasing the strength of
the expansion-contraction requires a highly concentrated and well-developed coronal muscle and
thicker mesoglea layer, which inevitably reduces the density of the polyp body. Hence, olivooid-
type feeding was most likely one of the prerequisite transitional forms for the rise of the jet-
propelled swimming style; in other words, rhythmic jet-propelled swimming is most likely a by-
product of occasional/frequent olivooid-type feeding of periderm-bearing sedentary medusozoans.
These inferences fit well with the appearance of centimetre-scale predatory swimming medusae
with rhopalias at the beginning of the Cambrian Stage 3 (Cartwright et al. 2007; Han et al. 2016a).
376

Perspectives for future work and improvements
Previous palaeontological CFD simulations applied to extinct fossils have mostly used static
models of organisms and have mainly focused on the hydrodynamic efficiency of organisms in
water flowing at different velocities (Gutarra & Rahman 2022). This type of simulation has
potential for testing hypotheses in terms of an organism's functional and morphological
performance (Rahman 2017), providing assessment of and deep insights into the adaptability of
organisms to their environment from hydrodynamic perspectives. However, they did not involve
the dynamic effects of organisms on the ambient environment. Most previously simulated
palaeontological organisms were on the scale of millimetres to centimetres (Waters *et al.* 2017;
Gibson *et al.* 2019; Rahman *et al.* 2020; Song *et al.* 2021). The smaller size of *Quadrapyrgites*
indicated that they lived in a different current environment compared with larger or taller
organisms. For instance, the velocity gradient in the bottom boundary layer flow regime can lead
to higher ambient flow speeds with the increasing height of benthic organisms (Gibson *et al.*
2021). Therefore, lack of an ambient current environment may have an impact on the results.
However, adding an ambient current to the simulations can introduce more technical issues that
are too problematic to be addressed at this stage, such as poorer simulation convergency. In
addition, the dynamic effect of *Quadrapyrgites* on ambient water, which was the main focus of the

present study, needs to be investigated and visualised without the interference of current (as many
CFD simulations for modern jellyfish, *e.g.*, McHenry & Jed 2003; Gemmell *et al.* 2013; Sahin *et*
al. 2009 were conducted under hydrostatic conditions). Considering the two important points
stated above, the simulations here were conducted under a hydrostatic environment. We
emphasised that members of *Quadrapyrgites* were not configured to live in stagnant water
naturally, as discussed in a study by Liu *et al.* (2022). However, further investigations of the
impact of ambient currents on the feeding abilities of organisms may need to be performed by
designing a set of new insightful simulations.

Furthermore, it remains obscure whether *Quadrapyrgites* lived primarily in solitary or
gregarious modes. A benthic community with variable organism density can affect ambient water
conditions or the feeding capability of a single organism among it (Gibson *et al.* 2019; Liu *et al.*
2021). Remarkably, simulations of large gregarious communities are restricted here by the
computational resources and complexity of the model. Although simulating the gregarious benthic
communities generally involves modelling a multiple of organisms, which is beyond the present
computational capability, the results and data collected here could be used for further simulations
to achieve a better understanding of gregarious active feeding and respiration behaviours of the
co-existing small shelly fossils. One promising approach is building a simplified active feeding
model by reconstructing the flow velocity profile collected accordingly. Thus, it is possible to
further investigate the ecological characteristics of gregarious organisms and the effect of
community size on the adaptability of organisms living in the ambient environments.

It is also noteworthy that we omitted the possible effects of the polyp stalk of *Quadrapyrgites*
and other internal structures of the calyx on the water flowing in and out of the peridermal
aperture. The interaction between these internal structures and subumbrella contraction remains
unclear. To this end, we did not model the internal structure of the polyp. In addition, the
manubrium in the subumbrella cavity as well as the tentacles could act as a barrier to water flow in
and out. The true thickness of the mesoglea is also unknown because of diagenesis, which may
influence the exact magnitude of contraction and expansion. However, based on the fossil record,
the mesoglea would have been much thinner than that in modern medusae (Han *et al.* 2016).
Although the primary dynamic pattern of *Quadrapyrgites* could be much more subtle, the
simplified model required less computational resources and adequate restoration of the polyp body
plan. Our study will shed new light on the autecology of Cambrian micro-benthos using numerical
computational methods.
426
Conclusions
Our simulations of *Quadrapyrgites* show that the accelerated expansion of the polyp body can
improve active feeding efficiency and increase the range in the upper flow above the peridermal
aperture height. The contraction/expansion pattern of the polyp body and rough peridermal surface
helps the polyp to access food particles in the ambient environment of the periderm, thereby
enhancing the polyp's feeding and gas exchange efficiencies under relatively low flow velocity
conditions. Eventually, as body size and height of the Cambrian benthic medusozoans increase,
this mode of feeding will be replaced by more efficient feeding methods (*e.g.*, relying on free
tentacles). Our study has implications for understanding the feeding and respiration of olivooids
and other sedentary medusozoans completely dwelling in their periderm. Furthermore, our
findings provide valuable insights into the interactions between the evolution of animal body size,

an increased incidence of swimming behaviour in medusa, and the ambient environment during

the Cambrian explosion. This is also the first time a dynamic numerical simulation method has

been applied to a microfossil, demonstrating the further possibilities for utilising this approach in

palaeontological research.

Material and methods

Geological setting and fossil pre-treatment

Rock samples of olivooids were collected from the Shizhongguo and Zhangjiagou sections of the

Kuanchuanpu Formation, Shaanxi Province, China. The Shizhongguo section of the Kuanchuanpu

Formation in is an interbedded set of cherts, flint, and phosphatic tuffs and is approximately 60 m

thick. Overlying the Guojiaba Formation is a black carbonaceous shale and siltstone,

approximately 8 m thick, dominated by detrital dolomite. The Zhangjiagou in the Kuanchuanpu

Formation in section is a thickly bedded set of phosphorus limestone, approximately 22 m thick,

whereas the underlying Dengying Formation is dominated by massive, thickly-bedded, black

dolomite. Small shell fossil specimens were obtained from the *Anabarites-Protohertzina-*

*Arthrochitie*s biozone, which corresponds to the Cambrian Fortunian Stage (Qian 1977; 1999).

The rocks were smashed to a width of 2–3 cm, immersed in a 7–10% acetic acid solution to

decompose, and the residue was air-dried before the fossil samples were manually picked out

under a binocular microscope (Leica M20 stereoscopic microscope). SEM (FEI Quanta 400 FEG

scanning electron microscope) was used for the scanning photography.

The collected fossil data were modelled using Dragonfly 4.0 and high-resolution 3D models

were generated using Autodesk Maya 2018 and saved in the '.stl' format. All fossil specimens and

model files were stored at the Shaanxi Key Laboratory of Early Life and Environments and

Department of Geology, Northwestern University.

Three-dimensional modelling

Simulations were carried out for *Quadrapyrgites* (Figure supplement 1A). The 3D model of

Quadrapyrgites (height: 2.0 mm, length: 0.6 mm, width: 0.6 mm) consists of two parts: an outer

pagoda-shaped periderm and a hollow bowl-shaped polyp (Figure 3A). The latter is functionally a

proxy of the polyp subumbrella without the manubrium, internal tissue of polyps, and basal stalk

(Figure 3B). The dynamic process of the simulated polyp was modelled by rhythmic contraction

and expansion of the subumbrella, and it was divided into contraction, expansion, and normal

resting phases. In the initial state, the subumbrella was about 0.7 mm in height and 0.1 mm in

diameter at its widest point. At this point, the umbrella muscle and mesoglea layer contracted, and

the size of the subumbrella opening was minimal. Subsequently, the umbrella expanded and

increased in size. Finally, the polyp subumbrella shrank and returned to its original shape.

Computational fluid dynamics

We used COMSOL Multiphysics v. 5.6 (https://cn.comsol.com) to carry out 3D simulations of

Quadrapyrgites. The computational domain consisted of a cuboid with a length, width, and height

of 20, 10, and 10 mm, respectively. The *Quadrapyrgites* model was placed at the centre of the

lower boundary of the cuboid domain in the direction of the peridermal aperture-apex from top to

bottom and was inserted vertically into the lower boundary approximately 0.18 mm (Figure

supplement 1B). The part below the lower boundary was then removed by Boolean operations

(Figure 3C). The mesh mainly consisted of the computational domain mesh and boundary layer
mesh applied to water-solid interacted boundaries (with a layer number of 2 and stretching factor
of 1.2) (Figure 3A). A free tetrahedral mesh was used to cover the entire simulation domain to
capture as much detail of the model as possible (Figure 3C, D). Each subdomain of the whole
simulation domain was meshed with specific levels of refinement (*i.e.*, varied maximum and
minimum element sizes were applied according to sizes of subdomains) to ensure a balance
between the accuracy and computational cost. Sensitivity tests were performed to determine the
optimal settings for the subsequent simulations (see "Mesh sensitivity analysis").

The simulations used the hyper-elastic material model in the membrane node of solid
mechanics to define the structure of the umbrella surface of the *Quadrapyrgites* polyp inside the
periderm. The stalk and other internal structures of the calyx, such as the manubrium and tentacles
in the subumbrella cavity, were ignored in the simplified model of the *Quadrapyrgites* polyp in
this study. The subumbrella, with a circular muscle bundle and mesoglea, was replaced with an
elastic membrane. Considering that the physical parameters of the polyp umbrella were difficult to
obtain directly from preserved fossil material, we used the physical parameters of elastic rubber
instead (Odgen material model (Holzapfel 2002), specific parameters are listed in Table
supplement 1). As the dynamics of the polyp subumbrella were determined by a displacement
function and not by the material elasticity of the subumbrella itself, this alternative setting would
not have a significant effect on the locomotion of the polyp subumbrella. The boundary of the
simulated polyp apex was set as a fixed constraint boundary that supported the simulated polyp
subumbrella.

To define the contraction and expansion motion of the umbrella of the polyp using the
prescribed displacement method, a columnar coordinate system was established in advance with
parameters a, φ, and r, where a is the axis of the centre of rotation of the polyp, φ is the angle of
rotation, and r is the distance between the polyp surface and origin. To simplify the physics setup
and mathematical model, our simulation reduced the motion of the umbrella inside the polyp to
motion in the r-direction only, without considering its motion in the a-direction. For this reason,
the displacement of the contraction and expansion motion of the umbrella inside the simulated
polyp in the r-direction was defined by an interpolation function with height z as a variable (Table
supplement 1), and the function was fitted using the cubic spline method to ensure the smoothness
of the displacement process inside the simulated polyp.

The ratio of contraction to expansion times varies between different species and even within
the same individual in modern cnidarians, with generally short contraction times and relatively
long expansion times, as demonstrated by *Aurelia aurita* (McHenry & Jed 2003). Considering that
the muscle contraction capacity of the polyp subumbrella may differ from that of modern jellyfish,
the contraction time of the polyp in the simulation was fixed at 1 s. Four sets of experiments with
different contraction time duty cycles, or expansion/contraction time ratios, were conducted to
simulate the expansion and contraction of the polyp subumbrella: 1 s:1 s, 2 s:1 s, 3 s:1 s, and 4 s:1
s, respectively. All sets of simulated movements were implemented separately using the
corresponding smoothing functions. As the displacements of the subumbrella of the simulated
polyp only occurred in the r-direction, the displacement of the subumbrella of the simulated polyp
in the a-direction and φ-direction were prescribed as 0 to avoid unexpected twisting of the mesh
and to ensure convergence of the simulations.

In the fluid domain, the Reynolds number (approximately 0.96, less than 1) was calculated by

using the diameter of the opening of the simulated subumbrella at 0.003 mm within the polyp as

the characteristic size. In this case, viscous forces dominated the fluid domain, and the influence

of inertial forces was negligible (*i.e.*, the inertial term in the Navier-Stokes equations was equal to

zero); thus, the peristaltic flow model was chosen for all simulations. We assumed that the

tetraradial pagoda-shaped *Quadrapyrgites* lived in environments with relatively low flow

velocities. To better visualise the effect of the dynamics of *Quadrapyrgites* on the surrounding

environment, the simulated fluid domain was set as a hydrostatic environment. The top and

perimeter of the simulated domain were set as open boundaries to allow the flow of water in and

out without restrictions. In addition, the bottom boundary, umbrella surface, and peridermal

surface of the polyp in contact with the fluid were set as the no-slip boundaries. The density and

dynamic viscosity of water in the simulation were set to 1,000 kg/m^3 and 0.001 Pa·s, respectively.

The simulations used a unidirectional structure to the fluid coupling method (*i.e.*, velocity

transmission to fluid) to couple fluid and solid fields. In addition, the moving mesh feature was

enabled, allowing for the subumbrella model inside the periderm to be deformed. Considering the

complexity and computational cost of the physical fields in the simulation domain, the governing

equations were solved using a time-dependent, non-linear, and fully coupled solver with a relative

tolerance of 0.005 and a time step of T=0.01 s for the final output.

To obtain flow velocity data above the peridermal aperture and determine the effective range

of the active expansion/contraction of *Quadrapyrgites*, ten sampling cut points (Cut Point 3D)

were set above the peridermal aperture, ranging from z = 2.05 mm to z = 2.5 mm at the

coordinates of the simulated domain, with 0.05 mm between each cut point (Figure supplement

1B). The results were visualised as 2D cross sections along the centre of the rectangular

simulation domain. The simulation files were saved as '.mph' files.

Mesh sensitivity analysis

For the sensitivity analysis, a mesh model (Figure 3) with an expansion/contraction time ratio of

3:1 was used as the test model. Meshes with different levels of refinement were tested, with the

number of elements ranging from 75,781 to 1,009,782 (Table supplement 2). The flow velocity

values obtained from the cut points with different meshes on each timestep were compared, and

the average difference of each timestep was calculated. Results were considered mesh independent

when the difference ranged from 5–10% between those obtained with the current mesh and the

next most fine mesh. Finally, the mesh selected for subsequent simulations contained 670,654

elements in total (see Table supplement 3 for specific mesh parameters). The average error

between the results obtained with such meshes and those with a finer mesh was approximately 5.5%

(Table supplement 2).

Acknowledgements

We thank H.J. Gong, X. Liu, and M.R. Cheng (State Key Laboratory for Continental Dynamics,

Northwest University, Xi'an, China) for their assistance in both field and lab work.

Data Availability Statement

Supplementary data of this manuscript have been deposited in the general data repository figshare

(https://doi.org/10.6084/m9.figshare.23282627.v2). Simulation files have been deposited to Dryad

(https://datadryad.org/stash/share/QGDSqLh8HOll7ofl6JWVrqM57Rp62ZPjvZU0AQQHwTY).

Funding

Funder	Grant reference number	Author
Natural Science Foundation of China	Nos. 42372012, 41720104002, 41911530236	Jian Han
Strategic Priority Research Program of Chinese Academy of Sciences	No. XDB26000000	Jian Han
Ministry of Science and 111 project of Ministry of Education of China	No. D17013	Jian Han
the Most Special Fund from the State Key Laboratory of Continental Dynamics, Northwest University, China	BJ11060	Jian Han
the High Level Talent Research Initiation Fund (Technology), Linyi University, China	Z6122059	Xing Wang
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication		

Author contributions. JH and XW designed the study work; XW and JH provided and interpreted the fossil samples; YZ conducted the simulations; YZ, JH, XW and TZ completed the main text with input from the other authors. All authors read and approved the final manuscript.

Author ORCIDs

Yiheng Zhang http://orcid.org/0009-0002-2426-0838

Xing Wang http://orcid.org/0000-0002-1777-864X

Jian Han http://orcid.org/0000-0002-2134-4078

Juyue Xiao http://orcid.org/0009-0000-4014-5763

Yuanyuan Yong http://orcid.org/0009-0003-9718-9953

Chiyang Yu http://orcid.org/0009-0002-9620-0500

Ning Yue http://orcid.org/0000-0003-0775-9936

Jie Sun http://orcid.org/0009-0005-4870-9670

Kaiyue He http://orcid.org/0000-0002-1816-5298

Wenjing Hao http://orcid.org/0000-0002-0622-369X

Tao Zhang http://orcid.org/0000-0002-5622-0227

Bin Wang http://orcid.org/0000-0002-1589-8939

Deng Wang http://orcid.org/0000-0002-4464-9632

Xiaoguang Yang http://orcid.org/0000-0002-5941-9899

References

Arai MN. 1997. A Functional Biology of Scyphozoa. Chapman & Hall, London

Caldwell DR, Chriss TM. 1979. The viscous sublayer at the sea floor. Science 205:1131–1132. doi:10.1126/science.205.4411.1131

Cartwright P, Halgedahl SL, Hendricks JR, Jarrard RD, Marques AC, Collins AG, Lieberman BS.

2007. Exceptionally preserved jellyfishes from the Middle Cambrian. PloS one 2:e1121. doi:
10.1371/journal.pone.0001121

Chang S, Clausen S, Zhang L, Feng Q, Steiner M, Bottjer DJ, Zhang Y, Shi M. 2018. New
probable cnidarian fossils from the Lower Cambrian of the Three Gorges area, South China,
and their ecological implications. Palaeogeography, Palaeoclimatology, Palaeoecology
505:150–166. doi:10.1016/j.palaeo.2018.05.039

Dabiri JO, Colin SP, Costello JH. 2007. Morphological diversity of medusan lineages constrained
by animal-fluid interactions. Journal of Experimental Biology 210:1868–1873.
doi:10.1242/jeb.003772

Dabiri JO, Colin SP, Costello JH, Gharib M. 2005. Flow patterns generated by oblate medusan
jellyfish: field measurements and laboratory analyses. Journal of Experimental Biology
208:1257–1265. doi:10.1242/jeb.01519

Darroch SA, Rahman IA, Gibson B, Racicot RA, Laflamme M. 2017. Inference of facultative
mobility in the enigmatic Ediacaran organism *Parvancorina*. Biology Letters 13:20170033.
doi:10.1098/rsbl.2017.0033

Denny M. 1988. Biology and the Mechanics of the Wave-swept Environment. Princeton
University Press.

Dong X, Cunningham JA, Bengtson S, Thomas CW, Liu J, Stampanoni M, Donoghue PC. 2013.
Embryos, polyps and medusae of the Early Cambrian scyphozoan *Olivooides*. Proceedings of
the Royal Society B: Biological Sciences 280:20130071. doi:10.1098/rspb.2013.0071

Dong X, Vargas K, Cunningham JA, Zhang H, Liu T, Chen F, Liu J, Bengtson S, Donoghue PC.
2016. Developmental biology of the early Cambrian cnidarian *Olivooides*. Palaeontology
59:387–407. doi:10.1111/pala.12231

Dynowski JF, Nebelsick JH, Klein A, Roth-Nebelsick A. 2016. Computational fluid dynamics
analysis of the fossil crinoid *Encrinus liliiformis* (Echinodermata: Crinoidea). PLoS One
11:e0156408. doi:10.1371/journal.pone.0156408

Dzik J, Baliński A, Sun Y. 2017. The origin of tetraradial symmetry in cnidarians. Lethaia 50:306–
321. doi:10.1111/let.12199

Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D, Peterson KJ. 2011. The Cambrian
conundrum: early divergence and later ecological success in the early history of animals.
Science 334:1091–1097. doi:10.1126/science.1206375

Feitl KE, Millett AF, Colin SP, Dabiri JO, Costello JH. 2009. Functional morphology and fluid
interactions during early development of the scyphomedusa *Aurelia aurita*. The Biological
Bulletin 217:283–291. https://doi.org/10.1086/BBLv217n3p283

Gemmell BJ, Colin SP, Costello JH. 2018. Widespread utilization of passive energy recapture in
swimming medusae. Journal of Experimental Biology 221:jeb168575.
doi:10.1242/jeb.168575

Gemmell BJ, Costello JH, Colin SP, Stewart CJ, Dabiri JO, Tafti D, Priya S. 2013. Passive energy
recapture in jellyfish contributes to propulsive advantage over other metazoans. Proceedings
of the National Academy of Sciences 110:17904–17909. doi:10.1073/pnas.1306983110

Gibson BM, Furbish DJ, Rahman IA, Schmeeckle MW, Laflamme M, Darroch SA. 2021. Ancient
life and moving fluids. Biological Reviews 96:129–152. doi:10.1111/brv.12649

Gibson BM, Rahman IA, Maloney KM, Racicot RA, Mocke H, Laflamme M, Darroch SA. 2019.
Gregarious suspension feeding in a modular Ediacaran organism. Science Advances

5:eaaw0260. doi:10.1126/sciadv.aaw0260

Grant WD, Madsen OS. 1986. The continental-shelf bottom boundary layer. Annual Review of
Fluid Mechanics 18:265–305. doi:10.1146/annurev.fl.18.010186.001405

Guo J, Han J, Van Iten H, Song Z, Qiang Y, Wang W, Zhang Z, Li G. 2021. A ten-faced
hexangulaconulariid from Cambrian Stage 2 of South China. Journal of Paleontology
95:957–964. doi:doi.org/10.1017/jpa.2021.25

Guo J, Han J, Van Iten H, Song Z, Qiang Y, Wang W, Zhang Z, Li G, Sun Y, Sun J. 2020a. A new
tetraradial olivooid (Medusozoa) from the Lower Cambrian (Stage 2) Yanjiahe Formation,
South China. Journal of Paleontology 94:457–466. doi:10.1017/jpa.2019.101

Guo J, Han J, Van Iten H, Wang X, Qiang Y, Song Z, Wang W, Zhang Z, Li G. 2020b. A fourteen-
faced hexangulaconulariid from the early Cambrian (Stage 2) Yanjiahe Formation, South
China. Journal of Paleontology 94:45–55. doi:10.1017/jpa.2019.56

Gutarra S, Rahman IA. 2022. The locomotion of extinct secondarily aquatic tetrapods. Biological
Reviews 97:67–98. doi:10.1111/brv.12790

Han J, Hu S, Cartwright P, Zhao F, Ou Q, Kubota S, Wang X, Yang X. 2016a. The earliest pelagic
jellyfish with rhopalia from Cambrian Chengjiang Lagerstätte. Palaeogeography,
Palaeoclimatology, Palaeoecology 449:166–173. doi:10.1016/j.palaeo.2016.02.025

Han J, Kubota S, Li G, Ou Q, Wang X, Yao X, Shu D, Li Y, Uesugi K, Hoshino M. 2016b.
Divergent evolution of medusozoan symmetric patterns: Evidence from the microanatomy of
Cambrian tetramerous cubozoans from South China. Gondwana Research 31:150–163.
doi:10.1016/j.gr.2015.01.003

Han J, Kubota S, Li G, Yao X, Yang X, Shu D, Li Y, Kinoshita S, Sasaki O, Komiya T. 2013.
Early Cambrian pentamerous cubozoan embryos from South China. PloS one 8:e70741.
doi:10.1371/journal.pone.0070741

Han J, Guo J, Ou Q, Song Z, Liu P, Hao W, Sun J, Wang X. 2020. Evolutionary framework of
early Cambrian cnidarians from South China. Earth Science Frontiers 27:67.
doi:10.13745/j.esf.sf.2020.6.3

Holzapfel GA. 2002. Nonlinear Solid Mechanics: A Continuum Approach for Engineering Science.
Kluwer Academic Publishers Dordrecht.

Hu S, Steiner M, Zhu M, Erdtmann B-D, Luo H, Chen L, Weber B. 2007. Diverse pelagic
predators from the Chengjiang Lagerstätte and the establishment of modern-style pelagic
ecosystems in the early Cambrian. Palaeogeography, Palaeoclimatology, Palaeoecology
254:307–316. doi:10.1016/j.palaeo.2007.03.044

Jarms G, Morandini A, da Silveira F. 2002. Cultivation of polyps and medusae of Coronatae
(Cnidaria, Scyphozoa) with a brief review of important characters. Helgoland Marine
Research 56:203–210. doi:10.1007/s10152-002-0113-3

LaBarbera M. 1984. Feeding currents and particle capture mechanisms in suspension feeding
animals. American Zoologist 24:71–84. doi:10.1093/icb/24.1.71

Leclère L, Röttinger E. 2017. Diversity of cnidarian muscles: function, anatomy, development and
regeneration. Frontiers in Cell and Developmental Biology 4:157. doi:
10.3389/fcell.2016.00157

Leme JM, Van Iten H, Simões MG. 2022. A new conulariid (Cnidaria, Scyphozoa) from the
terminal Ediacaran of Brazil. Frontiers in Earth Science 10:777746.
doi:10.3389/feart.2022.777746

Li G, Zhu M, Van Iten H, Li C 2004. Occurrence of the earliest known Sphenothallus Hall in the Lower Cambrian of southern Shaanxi Province, China. Geobios 37:229–237. doi:10.1016/j.geobios.2003.04.002

Li P, Hua H, Zhang L, Zhang D, Jin X, Liu Z. 2007. Lower Cambrian phosphatized *Punctatus* from southern Shaanxi and their ontogeny sequence. Chinese Science Bulletin 52:2820–2828. doi:10.1007/s11434-007-0447-y

Liu P, Yang X, Tang Q, He K, Zhang Y. 2021. Computational fluid dynamics analysis of the gregarious behavior of *Anabarites trisulcatus* from the early Cambrian. Acta Micropalaeontologica Sinica 38:234–240. doi:10.16087/j.cnki.1000-0674.20210906.002

Liu P, Zhang Y, Yang X, Wang B, Zhang T, Sun J, Tang Q, He K, Hao W, Yue N. 2022. Hydrodynamic simulations of millimeter-scale Cambrian sedentary medusozoans. Journal of Geophysical Research: Biogeosciences 127:e2022JG006854. doi:10.1029/2022JG006854

Liu Y, Li Y, Shao T, Zhang H, Wang Q, Qiao J. 2014. *Quadrapyrgites* from the Lower Cambrian of South China: growth pattern, post-embryonic development, and affinity. Chinese Science Bulletin 59:4086–4095. doi:10.1007/s11434-014-0481-5

Liu Y, Shao T, Zhang H, Wang Q, Zhang Y, Chen C, Liang Y, Xue J. 2017. A new scyphozoan from the Cambrian Fortunian Stage of South China. Palaeontology 60:511–518. doi:doi.org/10.1111/pala.12306

McHenry MJ, Jed J. 2003. The ontogenetic scaling of hydrodynamics and swimming performance in jellyfish (*Aurelia aurita*). Journal of Experimental Biology 206:4125–4137. doi:10.1242/jeb.00649

Mohseni K, Gharib M. 1998. A model for universal time scale of vortex ring formation. Physics of Fluids 10:2436–2438. doi:10.1063/1.869785

Mohseni K, Ran H, Colonius T. 2001. Numerical experiments on vortex ring formation. Journal of Fluid Mechanics 430:267–282. doi:0.1017/S0022112000003025

Conway Morris S, Selden PA, Gunther G, Jamison PG, Robison RA. 2015. New records of Burgess Shale-type taxa from the middle Cambrian of Utah. Journal of Paleontology 89:411–423. doi:10.1017/jpa.2015.26

Pratt MC. 2008. Living where the flow is right: How flow affects feeding in bryozoans. Integrative and Comparative Biology 48:808–822. doi:10.1093/icb/icn052

Qian Y. 1977. Hyolitha and some problematica from the Lower Cambrian Meishucun Stage in central and SW China. Acta Palaeontologica Sinica 16:255–275.

Rahman IA. 2017. Computational fluid dynamics as a tool for testing functional and ecological hypotheses in fossil taxa. Palaeontology 60:451–459. doi:10.1111/pala.12295

Rahman IA, Darroch SA, Racicot RA, Laflamme M. 2015. Suspension feeding in the enigmatic Ediacaran organism *Tribrachidium* demonstrates complexity of Neoproterozoic ecosystems. Science Advances 1:e1500800. doi:10.1126/sciadv.1500800

Rahman IA, O'Shea J, Lautenschlager S, Zamora S. 2020. Potential evolutionary trade‑off between feeding and stability in Cambrian cinctan echinoderms. Palaeontology 63:689–701. doi:10.1111/pala.12495

Robertson MA, Efremov F, Paik J. 2019. RoboScallop: A bivalve inspired swimming robot. IEEE Robotics and Automation Letters 4:2078–2085. doi:10.1109/LRA.2019.2897144

Sahin M, Mohseni K, Colin SP. 2009. The numerical comparison of flow patterns and propulsive
performances for the hydromedusae *Sarsia tubulosa* and *Aequorea victoria*. Journal of
Experimental Biology 212:2656–2667. doi:10.1242/jeb.025536

Shick, J.M., 1991. A Functional Biology of Sea Anemones. Springer Dordrecht.

Song Hanchen, Song Haijun, Rahman IA, Chu D. 2021. Computational fluid dynamics confirms
drag reduction associated with trilobite queuing behaviour. Palaeontology 64:597–608.
doi:10.1111/pala.12562

Steiner M, Qian Y, Li G, Hagadorn JW, Zhu M. 2014. The developmental cycles of early
Cambrian Olivooidae fam. nov.(? Cycloneuralia) from the Yangtze Platform (China).
Palaeogeography,   Palaeoclimatology,   Palaeoecology   398:97–124.
doi:10.1016/j.palaeo.2013.08.016

Straehler-Pohl I. 2017. Cubozoa and Scyphozoa: The results of 20 years of scyphozoan life cycle
research with new results on cubozoan life cycles to suggest a new nomenclature referring to
both classes. Frontiers in Ecological Studies of Jellyfish (eds Toyokawa M, Miyake H,
Nishikawa, J), Seibutsu Kenkyu Sha Co Ltd (Organisms Research Co Ltd), Tokyo 17–29.

Trowbridge JH, Lentz SJ. 2018. The bottom boundary layer. Annual Review of Marine Science
10:397–420. doi:10.1146/annurev-marine-121916-063351

Van Iten H, de Moraes Leme J, Simoes MG, Marques AC, Collins AG. 2006. Reassessment of the

phylogenetic position of conulariids (? Ediacaran-Triassic) within the subphylum medusozoa

(Phylum   Cnidaria).   Journal   of   Systematic   Palaeontology   4:109   –   118.

doi:10.1017/S1477201905001793

Wang X, Han J, Vannier J, Ou Q, Yang X, Uesugi K, Sasaki O, Komiya T. 2017. Anatomy and

affinities of a new 535‑million‑year‑old medusozoan from the Kuanchuanpu Formation,

South China. Palaeontology 60:853–867. doi:10.1111/pala.12320

Wang X, Vannier J, Yang X, Kubota S, Ou Q, Yao X, Uesugi K, Sasaki O, Komiya T, Han J. 2020.
An intermediate type of medusa from the early Cambrian Kuanchuanpu Formation, South
China. Palaeontology 63:775–789. doi:10.1111/pala.12483

Wang X, Vannier J, Yang X, Leclère L, Ou Q, Song X, Komiya T, Han J. 2022. Muscle systems
and motility of early animals highlighted by cnidarians from the basal Cambrian. eLife
11:e74716. doi:10.7554/eLife.74716

Waters JA, White LE, Sumrall CD, Nguyen BK. 2017. A new model of respiration in blastoid
(Echinodermata) hydrospires based on computational fluid dynamic simulations of virtual 3D
models. Journal of Paleontology 91:662–671. doi:10.1017/jpa.2017.1

Wengrove ME, Foster DL. 2014. Field evidence of the viscous sublayer in a tidally forced
developing   boundary   layer.   Geophysical   Research   Letters   41:5084–5090.
doi:10.1002/2014GL060709

Yong Y, Peng J, Wang X, Yu C, Hao, Sun J, Guo J, Han J. 2022. A new species of *Octapyrgites*
(Cnidaria) from the Lower Cambrian Kuanchuanpu formation, Southern Shaanxi. Acta
Micropalaeontologica Sinica 39:223–234.

Zhang Y, Zhang T, Yong Y, Yu C, Xiao J, He K, Wang D, Wang X, Wang B, Yang X, Han J. 2022.
Effect of boundary layer on simulation of microbenthic fossils in coastal and shallow seas.

770   Earth Science Frontiers. doi:10.13745/j.esf.sf.2023.5.32

771 Zhu M, Van Iten H, Cox RS, Zhao Y, Erdtmann B-D. 2000. Occurrence of *Byronia* Matthew and

772   *Sphenothallus* Hall in the Lower Cambrian of China. PalZ 74:227–238.

773   doi:10.1007/BF02988098

774 Zhuravlev AY, Wood R. 2020. Dynamic and synchronous changes in metazoan body size during

775   the Cambrian Explosion. Scientific Reports 10:1–8. doi:10.1038/s41598-020-63774-2

777
778
779 **Figure 1** *Olivooides mirabilis* from the Cambrian Fortunian Stage Kuanchuanpu Formation, in
780 Shizhonggou section, Ningqiang County, Shaanxi Province, China. A, juvenile stage; B,
781 prehatched development stage; C, peridermal apertural view; D–G, possible internal calyx-like
782 polyp. Solid white dots indicate the pentaradial symmetry. Abbreviations: c, calyx; st, stalk; pe,
783 periderm. Scale bars: A, 500 μm; B, 200 μm; C–F, 300 μm; G, 400 μm.
784

785
786
Figure 2. *Quadrapyrgites quadratacris* of the Cambrian Fortunian Stage Kuanchuanpu Formation, in

the Zhangjiagou section, Xixiang County. A, polyp stage with uplifted peridermal aperture; B, apertural

view showing concaved peridermal aperture; C, possible calyx-like polyp with a stalk. D, a 2D

structure of *Quadrapyrgites* with simplified internal anatomy. Solid white dots indicate the pentaradial

symmetry. Abbreviations: eu, exumbrella; su, subumbrella; sc, subumbrellar cavity; te, tentacle; m,

mouth; pe, periderm; pa, periderm aperture; st, stalk; ca, calyx. Scale bar: 200 μm.

Figure 3. A, a 3D model of *Quadrapyrgites*; B, a reduced model of the polyp subumbrella; C, meshed

computational domain and boundary conditions (dashed box in C marks the position of D).

799
800
801   **Figure 4.** A, B, C, and D are the velocity variations with time collected by sampling cut points in all
802   simulations with expansion/contraction time ratios of 1:1, 2:1, 3:1, and 4:1, respectively.
803

804
805
806   **Figure 5.** Maximum flow velocity data collected by sampling cut points in all simulations with
807   different expansion/contraction time ratios.
808

Figure 6. A, C, E, and G show the status of the polyp subumbrella at different moments. B, D, F, and H
show the corresponding enlarged views of the polyp, with the white arrows representing the flow
direction and velocity magnitude (the size of the arrows is proportional to the natural logarithm of the
flow velocity magnitude with a range quotient of 1,000) of water flow. A, B, t = 0 s, the polyp is at rest,
and the subumbrella opening is in its maximum state; C, D, t = 1.8 s, the subumbrella is in the process
of expansion, and the flow velocity near the peridermal aperture has reached its maximum; E, F, t = 3 s,
the subumbrella is in its maximum state, and the subumbrella opening is in its minimum state; G, H, t =
3.5 s, the subumbrella is in the process of contraction, and the flow velocity near the peridermal
aperture has reached its maximum value.

Figure 7. Vortex visualisation of the dynamic process of *Quadrapyrgite* (the length of arrows was normalised to represent the orientation of velocity). A, t = 0.1 s, the upper main vortex, and the lower secondary vortex begin to form. B, t = 1s, contact between the secondary vortex and the lower boundary C, t = 2 s, the main and secondary vortex is developed to the maximum visualisation range of vorticity. D, E, t = 2.87–2.88s, separation of the main vortex occurs. F, t = 3s, the new main vortex of the contraction process formed, the expansion process ended, and the contraction process began. G, H, t = 3.9–4s, separation of the main vortex occurred.

Animation supplement 1. 2D visualisation of vortex (the length of arrows was normalised to represent the orientation of flow velocity).

832   **Figure supplement 1.** A, height of *Quadrapyrgites*. B, positions of *Quadrapyrgites* and cut points

833   in computational domain.

835