Chromatin activity of IκBα mediates the exit from naïve pluripotency

Luis G. Palma[1,2,3,δ], Daniel Álvarez-Villanueva[1,4,δ], María Maqueda[1,2,3], Mercedes Barrero[5], Arnau Iglesias[1,2,3], Joan Bertran[6], Damiana Álvarez-Errico[2], Carlos A. García-Prieto[2], Cecilia Ballaré[5], Virginia Rodriguez-Cortez[2], Clara Bueno[2], August Vidal[4], Alberto Villanueva[7], Pablo Menéndez[2,3,8,9,10], Gregoire Stik[2], Luciano Di Croce[5,10,11], Bernhard Payer[5,11], Manel Esteller[2,3,10,12], Lluís Espinosa[1,3,*] & Anna Bigas[1,2,3,*]

[1]Program in Cancer Research. Hospital del Mar Research Institute, Barcelona, Spain.

[2]Josep Carreras Leukemia Research Institute, Barcelona Spain.

[3]Centro de Investigación Biomédica en Red Cancer (CIBERONC), Madrid, Spain.

[4]Institut Investigació Biomèdica de Bellvitge (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

[5]Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, Spain.

[6]Faculty of Sciences, Technology and Engineering, Universitat de Vic - Universitat Central de Catalunya, Vic 08500 Spain

[7]Catalan Institute of Oncology (ICO/IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

[8]Spanish Network for Advanced Therapies (RICORS-TERAV). Carlos III Health Institute (ISCIII), Spain.

[9]Department of Biomedicine. University of Barcelona, Spain.

[10]Institucio Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

[11] Universitat Pompeu Fabra, Barcelona, Spain.

[12]Physiological Sciences Department, School of Medicine and Health Sciences, University of Barcelona (UB), Barcelona, Catalonia, Spain.

[δ]These authors contributed equally

Keywords: IκBα, NFKBIA, NF-κB, DNA Methylation, Polycomb, Embryonic Stem Cells, Pluripotency Exit, Germ Layer Specification,

*Corresponding authors:

Anna Bigas and Lluis Espinosa

Cancer Research Program. CIBERONC.

Hospital del Mar Research Institute

Doctor Aiguader 88, 08003, Spain.

Tel: +34 933160440; email: abigas@researchmar.net; lespinosa@researchmar.net

Summary

Maintenance of pluripotency is a multifactorial process in which NF-κB is a negative regulator. Our

previous work identified a chromatin role for IκBα, the master regulator of NF-κB signaling, that is

critical for the proper regulation of various tissue stem cells. Here, we found that IκBα accumulates

specifically in the chromatin fraction of pluripotent embryonic stem cells. IκBα depletion does not

affect NF-kB-dependent transcription, but causes a profound epigenetic rewiring in pluripotent stem

cells, including alterations in H3K27me3, a histone mark catalyzed by Polycomb repression complex

2. Chromatin changes induced by IκBα depletion affect a subset of pluripotency genes and are

associated with altered gene transcription. At the cellular level, IκBα-deficient embryonic stem cells

are arrested in a naive pluripotency state when cultured in serum/LIF conditions and fail to exit

pluripotency under differentiation conditions. By constructing separation-of-function mutants, we

show that the effects of IκBα in regulating stem cell pluripotency are NF-κB-independent, but mainly

rely on its chromatin-related function. Taken together, our results reveal a novel mechanism by which

IκBα participates in the regulation of the pluripotent state of embryonic stem cells and shed light on

the interplay between inflammatory signals and the regulation of pluripotency.

INTRODUCTION

Embryonic Stem Cells (ESCs) pluripotency is characterized by the capacity to generate all somatic and germline lineages both *in vitro* and *in vivo*, depending on the culture conditions. When cultured in serum plus Leukemia Inhibitory Factor (LIF) (Serum/LIF), murine ESCs (mESCs) include cells interconverting to different metastable states [1] that either resemble preimplantation blastocyst (naïve pluripotency) or post-implantation (primed pluripotency) embryonic stages [2–5]. The heterogeneity of mESCs in Serum/LIF is reduced by culturing them in the presence of GSK3β and MEK1/2 inhibitors (CHIR99021 and PD0325901, respectively) along with LIF (2i/LIF). Under these conditions, mESCs enter a ground state of naïve pluripotency characterized by self-renewal activity while suppressing any pro-differentiating signals [6,7]. This pluripotent state closely resembles the inner cell mass of the E4.0 mouse preimplantation blastocyst [8]. After implantation, pluripotent capability of the mouse embryo (E5.5) becomes restricted to Epiblast Stem Cells (EpiSCs), which represent a primed pluripotent state, characterized by coexpression of pluripotency and lineage specification markers [9,10]. EpiSCs can give rise to cells from the three germ layers (endoderm, mesoderm and ectoderm), albeit their contribution to chimeras or to germ cells is greatly compromised [4,11]. EpiSCs can be derived *in vitro* from naïve pluripotent cells when cultured in a medium supplemented with Nodal and fibroblast growth factor (FGF) [12]. Different epigenetic features distinguish naïve from primed pluripotent states [13], such as changes in the X chromosome inactivation in female cells [12,14], histone post-translational modifications [10] or a global DNA hypomethylation profile observed in naïve pluripotent stem cells [15–17]. These epigenetic changes are a prerequisite for the subsequent activation of gene circuits associated with pluripotency exit and germ layer specification [9,10]. Crucial regulatory elements for the transition from naïve to primed pluripotency state are located in distal DNA regions or enhancers to facilitate the resolution toward the primed state [18,19]. Therefore, identifying novel players to fine-tune the equilibrium between ground and primed states is crucial for preserving both pluripotency stability and differentiation capability in mESCs.

There is now strong evidence that inflammatory signals are critical for stem cell development [20–22], with NF-κB being the main effector. Different strategies to inhibit or attenuate NF-κB in ESCs lead to increase pluripotency gene expression and impair cell differentiation [23,24], or it facilitates reprogramming to induced pluripotent stem cells (iPSC) [25]. Canonical NF-κB signaling is triggered by the IKK kinase complex, which induce the phosphorylation and subsequent degradation of the NF-κB inhibitor, IκBα, leading to nuclear translocation of the NF-κB factors (e.g. p50/p65) [26]. However, we have previously identified a nuclear IκBα function, which is critical for tissue stem cell homeostasis and the proper differentiation of epidermal, intestinal and hematopoietic stem cells [27–30]. IκBα chromatin function is mediated by the interaction with histone deacetylases (HDACs), core elements of the Polycomb Repressor Complex 2 (PRC2) and histone H2A and H4 [27,29,31].

We have now investigated the role of IκBα in the context of pluripotent stem cells. Whereas
cytoplasmic IκBα is detected in the different states and its levels decrease after
differentiation, in agreement with NF-κB activity as a promoter of differentiation, we
specifically detected chromatin-bound IκBα in the naïve pluripotent stem cells. Importantly,
IκBα depletion in ESCs did not affect NF-κB activity but it stabilizes the naïve (2i/LIF-like) state
of pluripotency under Serum/LIF conditions, displaying an epigenetic rewiring, including
changes in the PRC2-dependent H3K27me3 histone mark and impacting in proximal and distal
regulation of pluripotent genes. Using a new Separation-Of-Function (SOF) IκBα mutants[32],
we have now definitively demonstrated that chromatin-bound IκBα is required for naïve
pluripotency exit of mESCs in an NF-κB-independent manner. Overall, we identified IκBα as a
novel key player in the regulation of the exit from naïve-to-primed pluripotency and essential
for the activation of gene programs involved in germ layer specification.

 RESULTS

 IκBα and NF-κB have opposite expression dynamics in naïve pluripotent cells

Due to the prominent role of NF-κB signaling in mESCs regulation, we analyzed the expression
pattern of the different NF-κB members including the IκB inhibitors in the three defined
pluripotent stages: ground state of naïve pluripotency (2i/LIF), naïve pluripotency
(Serum/LIF), and primed pluripotency (Epiblast Stem Cells or EpiSCs) [5] (Figure 1A). In
agreement with NF-κB being a pro-differentiation factor, canonical NF-κB inhibitors including
IκBα (*Nfkbia*), IκBβ (*Nfkbib*) and IκBε (*Nfkbie*) were all expressed in the naïve pluripotency
state (both 2i/LIF and Serum/LIF), with *Nfkbia* levels gradually reduced upon naïve
pluripotency exit towards EpiSCs (Figure 1B). In contrast, *Rela, Nfkb1 and Nfkb2* genes,
codifying for the NF-κB factors p65/RelA, p105 and p100, respectively, showed opposite
expression dynamics increasing the levels in EpiSCs (Figure 1C). We further characterized the
distribution of IκBα in the cytoplasm, nucleus and chromatin fractions of the three states of
pluripotency (Figure 1D). Total IκBα protein levels correlated with the detected RNA levels in
the three different cellular states being highest in the 2i/LIF condition (Figure 1B), with a large
accumulation in the cytoplasm. A fraction of IκBα protein was detected in the chromatin of
naïve pluripotent cells, being highest in cells cultured in Serum/LIF conditions compared to
2i/LIF and absent in the EpiSCs. (Figure 1D). The dynamics of IκBα RNA and protein levels from
naïve pluripotent ESCs to EpiSCs was further corroborated by analysis of an additional
database that includes both naïve and primed pluripotent states[33] (Figure 1E).

Next, we studied IκBα expression in differentiated embryoid bodies (EBs) from mESCs
(Serum/LIF) (see materials and methods) and analyzed the expression levels of different
pluripotency, differentiation-linked, and NF-κB genes by RNA sequencing. *Nfkbia* (IκBα gene)
was expressed at higher levels specifically in mESCs (Serum/LIF) whereas cells undergoing
differentiation (48h and 96h) displayed a reduction in its expression levels (Figure 1F). The
expression pattern of *Nfkbia* was similar to the pluripotency genes whereas most of the NF-
κB genes were upregulated upon differentiation, clustering together with developmental
genes, as previously described [24]. Notably, several polycomb elements were highly expressed
in mESCs, consistent with the possibility that chromatin-bound IκBα modulates PRC2 activity
[27] in the pluripotent state.

Taken together, these data show that, unlike the other NF-kB inhibitors and target genes, the
expression of *Nfkbia* is high in the naive pluripotent state, with IκBα protein present in the
cytoplasm and chromatin of naive pluripotent cells. Moreover, whereas canonical NF-κB
subunits are primarily absent, we detected several Polycomb elements expressed in mESCs.

 IκBα is required in mESCs to exit the naïve pluripotency state

To investigate the functional implication of IκBα in naïve pluripotency, we knocked-out (KO)
IκBα protein in mESCs by CRISPR-Cas9 (IκBα-KO mESCs) (Figure S1A and materials and

methods). Notably, IκBα KO mESCs have an impaired ability to undergo differentiation and
cannot properly switch off the pluripotency program after 216h (9 days) of pro-differentiation
signals, as it is demonstrated by high levels of pluripotency markers OCT3/4 and NANOG (Fig
1A and Fig S1C), higher number of alkaline phosphatase cells (Fig 1B and Fig S1) and higher
percentage of SSEA-1$^+$ cells (Fig S1B) within the 216h EBs. These results were also supported
by the detection of increased expression level of the naïve pluripotency genes *Pou5f1 (Oct3/4*
gene), Gbx2, Klf2, Sox2, Zfp42 (Rex1) and *Nanog* in IκBα KO EBs after 216h of differentiation
(Figure S1B).

To shed light into the molecular mechanisms governing the differentiation impairment in IκBα
KO EBs, we performed RNA-seq of IκBα WT and IκBα KO mESCs in the first stages of the
embryoid body differentiation. Analysis of the data indicated that IκBα KO EBs cannot
successfully evolve towards differentiation trajectories, as shown by principal component
analysis (PCA) of IκBα-WT and IκBα-KO EBs at 48h and 96h in differentiation media (Figure
2C). Analysis of naïve pluripotency markers at 96h (*Gab1, Sox2, Esrrb, Zfp42, Dppa4, Klf2,*
Prdm14, Klf4, Tfcp2l1) also showed higher expression levels in IκBα KO compared to their IκBα
WT counterparts (Figures 2D and S1F). Defective differentiation capacity of IκBα KO EBs
involved programs associated with specification of all 3 germ layers (endoderm, mesoderm,
and ectoderm) (Figures 2E-G). Moreover, IκBα KO EBs were smaller in size (Figure S1E), in
agreement with the reduced developmental potential of IκBα-depleted mESCs [34]. To further
study the requirement of IκBα in pluripotency exit *in vivo*, we established teratomas by
subcutaneous injection of IκBα-WT and IκBα-KO mESCs (Figure 2H and materials and
methods) into NSG (NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ) mice. IκBα-KO teratomas showed a
decreased differentiation potential, as indicated by higher number of OCT4$^+$ cells 6 weeks
after injection (Figures 2H-J). These observations further confirm the essential role of IκBα in
the exit from the pluripotency state not only *in vitro* but also under *in vivo* differentiation
signals. To understand the conditions in which IκBα is required for pluripotency exit, we
differentiated IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF (naïve pluripotency)
towards EpiSCs (primed pluripotency) (Figure 3A, upper panel). IκBα-KO cells were not
committed towards the primed pluripotency stage and maintained an elevated expression
levels of naïve pluripotency genes (*Dppa3, Nanog, Sox2, Rex1, Klf2, Klf4, Gbx2, Tbx3*) (Figure
3A, bottom panel). Additionally, IκBα-KO cells retained a naïve pluripotent morphology,
forming tight clusters that resemble cells cultured in 2i/LIF medium (Figure S2A).

mESCs cultured in serum/LIF are highly heterogeneous, comprising a mixture of cell states
that resemble developmental transitions from preimplantation (naïve pluripotency) to
postimplantation (primed pluripotency) embryo [2,3]. To investigate how IκBα deficiency was
affecting the exit of mESCs from naïve pluripotency, we performed a Gene Set Enrichment
Analysis (GSEA) using RNA-seq data from IκBα-KO and IκBα-WT mESCs cultured in Serum/LIF
against gene signatures for 2i/LIF and Serum/LIF pluripotency [35]. GSEA results revealed a
significant enrichment in Serum/LIF IκBα-KO mESCs transcriptome for genes specifically
expressed in the naïve ground state (2i/LIF), while they negatively correlate with mESCs

cultured in Serum/LIF (Figure 3B) [35]. These results agree with the fact that IκBα-KO mESCs
cultured in Serum/LIF resemble 2i/LIF morphology, with homogeneous tight clusters of cells,
whereas IκBα-WT mESCs retain the colony heterogeneity (Figure S2B). Interestingly, the
higher levels of ground-state-related genes *Zfp42* (*Rex1*), *Klf2* and *Tbx3* in IκBα-KO mESCs
were partially obtained when IκBα-WT mESCs were cultured in 2i/LIF for two consecutive
passages (Figure 3C).

One of the key features of the naïve pluripotency state is its global DNA hypomethylation
pattern [15,16]. We assessed the DNA methylation status of IκBα-WT and IκBα-KO mESCs
cultured in Serum/LIF conditions by analysis of 5-methyl-cytosine (5mC) mark and using DNA
methylation arrays [36]. We found that IκBα-KO mESCs contained much lower levels of 5mC
compared to IκBα-WT cells (Figure 3D). Moreover, DNA methylation arrays confirmed a global
pattern of DNA hypomethylation in IκBα-KO mESCs, which was maintained in 96h EBs (Figure
3E). Interestingly, the DNA hypomethylation status occurred in a genome-wide fashion
affecting all chromosomes (Figure S2C), in accordance with a higher pluripotent state of IκBα-
KO mESCs.

Overall, these data demonstrate that IκBα is required for the exit of the naïve state of
pluripotency and its deficiency results in impaired differentiation of mESCs *in vivo* and *in vitro*.
Accordingly, IκBα-KO mESCs cultured in Serum/LIF showed genome-wide pattern of DNA
hypomethylation.

Lack of IκBα causes an epigenetic rewiring in pluripotent stem cells to resemble 2i/LIF naïve
pluripotency in Serum/LIF culture

ChIP Enrichment Analysis (ChEA) of genes identified in our RNA-seq data of IκBα-KO and WT
mESCs revealed that differentially expressed genes were putative targets of chromatin
regulators including the PRC2 subunit SUZ12 and MTF2 (Figure 4A). We previously described
that IκBα is important for the proper deposition of the H3K27me3 mark in different types of
tissue stem cells by associating with elements of the PRC2 complex [27,28]. Since 2i/LIF and
Serum/LIF states are characterized by important differences in epigenetic marks, we carried
out an epigenetic profiling of IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF. We
performed chromatin immunoprecipitation followed by sequencing (ChIP-seq) of histone
modifications associated with gene activation (H3K4me3) and gene repression (H3K27me3)
(Figures 4B and S3A-B). Overall, Serum/LIF IκBα-KO mESCs exhibited a general increased
H3K4me3 status and redistribution of the PRC2-catalyzed H3K27me3 mark (Figure 4B).
Notably, gain of the H3K4me3 mark was associated with a reduction of H3K27me3 levels in
genes associated with naïve pluripotency function (Figures 4B-C and S3A-B). In particular, 41
genes with concomitant gain of H3K4me3 and loss of H3K27me3 were found to be enriched
in mechanisms associated with pluripotency function (Figure 4C), including Tbx3 and Gbx2
(Figure 4D-4E). A second set of pluripotency-related genes showed an increase in H3K4me3

without changes in H3K27me3 (Zfp42 or Klf2). Both sets of pluripotency genes showed
increased expression in the absence of IκBα (Figure 4H).

We next aimed to study whether the lack of IκBα was also affecting distal regulatory regions
by analyzing enhancer-associated histone marks in IκBα-WT and IκBα-KO mESCs. We
quantified the total amount of H3K27Ac (active enhancer) and H3K4me1 (poised enhancer)
by ChIP-seq experiments conducted in both IκBα-WT and IκBα-KO mESCs. Overall, IκBα-KO
mESCs showed a reduction in poised enhancers at genes (or related genomic regions)
associated with differentiation processes (Figures 4F and S3C) and an increase in active
enhancers at genes associated with pluripotency (Figure 4F and S3D), such as Tbx3, Tfcp2l1
and Zfp42 (Rex1) (Figure 4G), which positively favors their higher expression in IκBα-KO mESCs
(Figure 4H). These results further support that mESCs lacking IκBα are epigenetically
remodeled to favor the ground state of naive pluripotency, which may negatively impact on
their differentiation potential, similar to that described for other models with stabilized naïve
pluripotent state[37,38].

The chromatin function of IκBα in pluripotency exit is independent of classical NF-κB
activity.

Although previous results from C. elegans [39] and Drosophila [27] support the functional
relevance of chromatin-related IκBα function, the investigation of this alternative IκBα
activity remains challenging due to the predominant role of this protein in NF-κB regulation.
However, and similar to that found in other models, we did not detect major changes in
canonical NF-κB target genes upon IκBα depletion neither in Serum/LIF mESCs, nor 48h and
96h EB-differentiating conditions (Figure 5A). Thus, either NF-κB regulation does not require
IκBα at this stage or it might be compensated by the other IκBs (IκBβ, IκBε or p100), as
previously demonstrated [40,41]. This observation suggests that either NF-κB signaling is not
active in naïve pluripotency, which is supported by the low expression levels of NF-κB
subunits (see Figures 1B-C and 1E-F), or IκBα function on NF-κB is compensated by the other
IκBs (IκBβ, IκBε or p100), as previously demonstrated[40,41].

To further investigate the relative impact of both IκBα functions in mESCs, we took advantage
of recent results from our group that led to the identification of the specific IκBα protein
residues that define NF-κB- or chromatin-binding through H2A/H4. Mutation of these
residues allowed the generation of doxycycline-inducible (i) separation-of-function IκBα
mutants that are specifically deficient in one or the other function, at least in the intestinal
epithelial cells[32]. We called i-IκBα[ΔChromatin] the i-SOF IκBα mutant that interacts with NF-κB
but is deficient for H2A and H4 binding and i-IκBα[ΔNF-κB] the one that interact with histones but
is deficient for NF-κB subunits binding (Figure 5B and [32]). The three different IκBα forms (i-
IκBα[WT], i-IκBα[ΔNF-κB] and i-IκBα[ΔChromatin]) were stably transfected in the IκBα KO mESCs and
induced by 16h of doxycycline treatment (Figure S4A). We used these cells to investigate the
biochemical properties and functional impact of SOF IκBα mutants in mESCs. By pull-down

experiments, we found that i-IκBα$^{\Delta NF-\kappa B}$ expressed in ESCs was able to interact with histone
H2A but not with the NF-κB member p50. In contrast, i-IκBα$^{\Delta Chromatin}$ interacts with p50 but
not with H2A, while i-IκBα^{WT} preserves both binding capacities (Figure 5C). In concordance
with these results, i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ were found in both the cytoplasmic and chromatin
fractions of mESCs, whereas IκBα$^{\Delta Cromatin}$ was retained in the cytoplasm (Figures 5D-E). In
multiple experiments, we noticed a significant reduction in protein levels of IκBα$^{\Delta chromatin}$
despite the mRNA expression levels of the different IκBα forms being comparable (Figures 5C-
D and S4A).

Then, we studied whether i-IκBα^{WT} and/or the i-SOF IκBα mutants (IκBα$^{\Delta NF-\kappa B}$ or IκBα$^{\Delta Chromatin}$)
were able to reverse any of the phenotypes observed in the IκBα-KO mESCs (Figure S4B).
Induction of i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ reduced the expression levels of the naïve pluripotent
genes *Zfp42, Klf2, Sox2* and *Tbx3*, which were increased by i-IκBα$^{\Delta Chromatin}$ (Figure 5F). In
addition, the ground state-associated DNA hypomethylation was specifically reverted in i-
IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ mESCs but not in i-IκBα$^{\Delta Chromatin}$ mESCs (Figure 5G). Finally, we
addressed whether i-IκBα^{WT}, i-IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta Chromatin}$ mESCs were capable to revert the
differentiation blockage of IκBα-KO ESCs (Figure S4B). Induction of IκBα^{WT} and IκBα$^{\Delta NF-\kappa B}$ was
enough to restore the differentiation potential of IκBα-KO mESCs into EBs, as indicated the
increase in the number of differentiated cells (based on percentage of SSEA-1^{neg} cells) (Figures
5H and S4C) and the reduced number of AP+ colonies (in the 216h IκBα-KO reconstituted EB
assays), which was similar to the number of colonies obtained from IκBαWT EBs (Figures 5I).
Lastly, the number of remaining pluripotent-like cells (defined by OCT3/4 NANOG staining)
was also reduced in i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ 216h EBs, and only small and few regions of
undifferentiated cells was observed. In contrast, induction of i-IκBα$^{\Delta Chromatin}$ did not restore
the differentiation potential of IκBα-KO mESCs (Figures 5J and S4D).

These results demonstrate that the chromatin-dependent IκBα function, but not its NF-κB-
related activity, is necessary for the proper regulation of chromatin marks at specific genomic
regions that are linked to transcriptional changes that are likely at the base of the capacity of
ESCs to exit the ground state of naïve pluripotency and differentiate.

DISCUSSION

Here we have identified a new and unexpected function for the IκBα protein in the regulation of the naïve pluripotency exit. Using mESCs to investigate the role of IκBα in pluripotency maintenance and differentiation, we discovered that the absence of IκBα severely impairs the exit of pluripotency. Furthermore, IκBα-KO mESCs stabilizes the ground state of naïve pluripotency under Serum/LIF conditions, shedding light on its function as a critical regulator of the transition from naïve to primed pluripotency. While IκBα is commonly known for its role as an inhibitor of the NF-κB signaling pathway [26], our laboratory has previously furnished compelling evidence of its alternative role through its interaction with histones and other chromatin components [27,29,31]. We have demonstrated that chromatin-bound IκBα plays a crucial role in regulating skin and intestinal stem cells [27,28], hematopoietic stem cell development [42], and it influences the regenerative capacity of these tissues, and in some cases, their susceptibility to neoplastic transformation. Disentangling the chromatin-related function of IκBα from its canonical role as an inhibitor of NF-κB has been challenging, making it difficult to determine its truly biological significance. To tackle this challenge, we engineered separation-of-function (SOF) mutants of IκBα, as detailed in [43], by identifying specific residues that are essential for binding either NF-κB elements or histones. Interestingly, we have shown that only the histone-binding proficient IκBα mutant (IκBα$^{\Delta NF-\kappa B}$), which lacks the ability to bind to NF-κB, has the capability to reverse the stabilization of the ground state observed in IκBα-KO mESCs. This is sufficient to facilitate their exit from the state of naïve pluripotency, restoring their full differentiation potential (Figure 5). The involvement of NF-κB in pluripotency has been a subject of prior investigation. Studies have revealed that pluripotent cells display a dampened inflammatory signaling pathway, which is induced upon pluripotency dissolution and activation of differentiation [24,44]. However, the precise molecular mechanisms underlying the attenuated status of NF-κB during this stage remain incompletely understood. A recent work has shed light on a potential regulatory axis involving ATG5-β-TrCP1-NF-κB, proposing it as a mechanism that diminishes NF-κB activity by stabilizing the IκBα protein in mESCs [23]. In any case, since modification of canonical NF-κB activity directly impacts on the levels IκBα expression, the previous NF-κB-associated phenotypes should be reevaluated considering this chromatin role of IκBα. Furthermore, the expression of IκBα$^{\Delta Chromatin}$, which retains the inhibitory function of NF-κB, failed to mitigate the ground state stabilization observed in IκBα-KO mESCs (Figure 5). This finding suggests that the observed role of IκBα in regulating the exit from naïve pluripotency is not due to NF-κB inhibition. In fact, there is evidence that the chromatin function of IκBα is an ancestral function preceding its role as NF-κB inhibitor. In Caenorhabditis Elegans (*C.elegans*), which lacks NF-κB orthologs but possesses two homologs of IκBα, the absence of IκBα orthologs has severe multi-organ differentiation defects [39]. This observation further supports the notion that the chromatin-related function of IκBα plays a fundamental role in developmental processes, independent of its canonical role in NF-κB signaling.

We have observed that IκBα-KO mESCs exhibit epigenetic and transcriptomic profiles
resembling the ground state of naïve pluripotency under Serum/LIF culture (Figure 3). In this
regard, the fact that IκBα is preferentially located in the chromatin of mESCs cultured in
Serum/LIF, when they exhibit a high degree of heterogeneity, suggests that IκBα might act as
a regulator of this heterogeneity. One potential explanation might be that IκBα could be
affecting the stability of fluctuating transcription factors, which is crucial to mediate the
pluripotent-to-differentiation balance [3]. Thus, the lack of IκBα might stabilize those
transcription factors that would favor the ground-state observed in IκBα-KO mESCs cultured
in Serum/LIF. Further experiments are required to demonstrate this hypothesis. Moreover,
the differences in murine and human pluripotency [11] point out the necessity to address the
real impact of IκBα modulation in the stabilization of the ground state in human pluripotent
stem cells.

In summary, our work establishes that IκBα mediates the exit from naïve pluripotency by
modulating the activation of naïve pluripotency genes. Importantly, this newly identified role
of IκBα operates independently of its canonical function in inhibiting the NF-κB pathway. Our
findings underscore the intricate interplay between inflammation and pluripotency status,
revealing a previously unrecognized complexity in their interaction.

METHODS

Cell culture

mESC line ES-E14TG2a (ATCC; Cat #CRL-1821) was cultured on plastic dishes precoated with 0.1% (w/v) Gelatin (Sigma; Cat #G2500-100G) in Serum/LIF medium. Serum/LIF is composed by DMEM basal medium (Sigma-Aldrich; Cat #D5796) supplemented with 15% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 1X NEAA (Gibco; Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070), 1000U/mL Leukemia Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-mercaptoethanol (Gibco; Cat#31350010). Medium was changed every day, and cells were splitted every other day using TrypLE Express (Gibco; Cat #12605010) for harvesting. Cells were maintained in a 5% CO2 incubator at 37°C.

For inducing the ground-state of naïve pluripotency, mESCs were cultured in 2i/LIF medium for 2 consecutive passages. 2i/LIF medium is composed by NDiff 227 medium (TAKARA, Cat #Y40002) supplemented with 1000U/mL LIF, 0.4 µM PD032591 (Selleck Chemicals, Cat #S1036) and 3 µM CHIR99021 (Merck, Cat #SML1046).

Epiblast stem cell differentiation

Differentiation from mESCs towards Epiblast Stem Cells (EpiSCs) was performed as previously described [45]. $3x10^4$ mESCs were seeded in 6-well plastic dishes, and they were cultured in Serum/LIF medium for 24h. Medium was then switched to NDiff 227 medium (TAKARA, Cat #Y40002) supplemented with 20 ng/ml activin A (Cat. #120-14E; PeproTech) and 10 ng/ml Fgf basic (R&D Systems, Cat #233-FB-025/CF). Cells were submitted to daily media changes till day 5 (120h), when they were further analyzed.

Embryoid bodies differentiation from mESCs

Embryoid bodies differentiation was established as described in [46]. Briefly, mESCs were splitted twice in Serum/LIF medium before inducing differentiation. Once mESCs are 80% confluent, cells were collected using TrypLE Express. One well of 6-well plate was splitted in an entire 6-well plate in IMDM-ES medium for 48h. IMDM-ES medium is composed of Iscove's Modified Dulbecco's Medium (IMDM) (Cytiva, Cat #16SH30259.01) supplemented with 20% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 1X NEAA (Gibco; Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070), 10ng/mL Leukemia Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-mercaptoethanol (Gibco; Cat#31350010). For embryoid body induction, mESCs were harvested, and they were rinsed twice with DPBS (Gibco; Cat #14190144). Cells were very well disaggregated into single-cells, and 1.2×10^4 cells/mL embryoid body differentiation (EB^{diff}) medium were resuspended to a total volume of 25mL of EB^{diff} medium. EB^{diff} medium is composed by IMDM supplemented with 15% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 50

µg/mL ascorbic acid (Sigma; Cat #A-4544), 180 µg/mL Transferrin (Roche; Cat #10652202001)

and 0.45 mM alpha-monothioglycerol (MTG) (Sigma; Cat #M6145). EBs were formed in

suspension for 5 days (120h). At day 5, in order to elongate the differentiation up to 216h

(day 9), EB^{diff} medium was refreshed by harvesting EBs with 10mL serological pipette,

centrifuge at 200 g for 3 min and EBs were resuspended in fresh EB^{diff} medium for 4 more

days.

CRISPR/Cas9 gene editing and cell lines generation in mESCs

The two guide RNAs (gRNAs) targeting the Nfkbia locus were designed using the CRISPR design

tool from MIT (http://crispr.mit.edu). The best 2 gRNAs (based on on-target and off-target

scores) that were targeting the exon 1 of Nfkbia (IκBα gene) were selected. After annealing,

one of the gRNAs was cloned into SpCas9(BB)-2A-GFP (px458) plasmid (Addgene; Cat #48138),

and the other gRNA was cloned into the px330-mCherry plasmid (modified from Addgene;

Cat #98750 to incorporate mCherry reporter). mESCs were co-transfected with the two

plasmids (px458-gRNA1 and px330-mCherry-gRNA2). 3×10^5 cells were seeded a day before

per well of 6-well plate. At day of transfection, cells were washed once with 1X DPBS (Gibco;

Cat #14190144), and 2mL Opti-MEM medium (Gibco; Cat #31985070) was added into cells.

1.25 µg of each plasmid (2.5 µg of total DNA) was incubated with 10 µL of Lipofectamine 2000

(Invitrogen;   Cat   #11668019)   in   240   µL   of   Opti-MEM   for   20   min.

OptiMEM::Lipofectamine::DNA mixture was added into mESCs, and cells were incubated for

5 hours. Cells were then washed with DPBS, and medium was replaced by Serum/LIF. 48h

after transfection, GFP$^+$ mCherry$^+$ single cells were sorted by FACS using BD FACSAria II Cell

Sorter (BD Bioscience). Mutant clones were screened through PCR and western blotting to

identify single clones with no IκBα protein expression. Non-targeting scrambled gRNAs were

cloned into px458 and px330-mCherry plasmids as wildtype clones. Three independent clones

from each genotype (IκBα-WT and IκBα-KO) were selected for further experiments.

Doxycycline-inducible IκBα (IκBα^{WT}, IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta H2A/H4}$) mESCs were generated by

cloning the three different IκBα versions into a PiggyBac transposon system [47]. IκBα protein-

coding cDNAs were PCR-amplified, and NheI and SalI restriction sites were placed at the 5'

and 3' ends, respectively. T2A-EGFP fragment was amplified from PX458 plasmid, and SalI and

AgeI restriction sites were introduced at its 5' and 3' ends. PB-TRE backbone (Addgene, Cat#

63800) was digested with NheI and AgeI enzymes, and mNfkbia and T2A-EGFP fragments

were ligated into the PB-TRE digested vector. To generate mESCs containing the inducible

vector, IκBα-KO mESCs were co-transfected with PB-TRE-mNfkbia-T2A-EGFP and PiggyBac

transposase plasmids, and pool of transfected cells was selected by Hygromycin (100µg/mL)

for 7 days. Cells were doubly screened by FACS sorting green fluorescent protein after 24h of

doxycycline (1µg/mL) treatment. Pool of cells were used for further experiments. Oligos

sequences are included in supplementary table 1.

Teratoma formation assay

IκBα-WT and IκBα-KO mESCs grown in Serum/LIF (5×10^5) were injected intramuscularly into
severe combined immunodeficient mice (NSG). In order to favor 3D aggregation of cells and
teratoma formation, cells were resuspended in Matrigel Matrix (BD; Cat #356234) prior to
injection. Six weeks later, mice with tumors were euthanized, and tumors were fixed in
formaldehyde, embedded in paraffin, sectioned and stained with hematoxylin and eosin for
histological analysis.

Immunofluorescence staining

For immunostaining of mESCs, cells were seeded on 0.1% gelatin-coated coverslips on 6-well
plates; for EB immunofluorescence, they were collected directly from the plate. Samples were
washed twice with 1X PBS, and they were fixed at 4ºC for 30 minutes with 4% PFA (Electron
Microscopy Sciences; Cat #15713S). For immunofluorescence of 5-Methylcytosine (5mC),
fixed cells were incubated with 2M Hydrochloric Acid (Sigma-Aldrich, Cat #H1758) for 20
minutes at room temperature. Samples were then washed twice with 1X Tris-buffered saline
(TBS; 50 mM Tris-Cl, pH 8) and permeabilized and blocked with 1X TBS supplemented with 1%
Triton-X100 (MERCK; Cat #9036-19-5) and 6% FBS (Biological Industries; Cat #04-001-1A) for
2h at 4ºC. Samples were washed twice with 1X TBS supplemented with 6% FBS. Primary
antibody incubation was performed in 1X TBS plus 6% FBS and 0.3% or 0.5% Triton-X100 (0.3%
for 2D culture and 0.5% for EBs) overnight at 4ºC. The following primary antibodies were used:
OCT3/4 (1:250; Santa Cruz; Cat #sc-5279), NANOG (1:250; Novus Biologicals; Cat #NB100-
588), 5-Methylcytosine (1:500, Invitrogen, Cat #MA5-24694), H3K27Ac (1:2000, Abcam, Cat
#AB4729). Samples were then washed 4 times with 1X TBS, 5 min each washing. Secondary
antibody incubation was performed in TBS 1X plus 1% BSA (Sigma-Aldrich; Cat #9048-46-8)
for 2h at room temperature. The following antibodies were used: Alexa Fluor 488 donkey anti-
mouse antibody (1:1000; Invitrogen; Cat #A-21202), Alexa Fluor 647 donkey anti-mouse
antibody (1:1000; Invitrogen; Cat #A-31571), Alexa Fluor 594 donkey anti-rabbit antibody
(1:1000; Life Technologies; Cat #A-21207), Alexa Fluor 647 donkey anti-rabbit
antibody(1:1000, Invitrogen; Cat #A-31573). Samples were washed three times with 1X TBS
for 5 min at room temperature each rinse. Samples were mounted using DAPI Fluoromount-
G (Southern Biotech; Cat #0100-20).

Microscopy and image acquisition

Fluorescence images were acquired using Confocal Leica TCS SP5 (Leica Microsystems), and
Leica application software LAS AF (Leica Microsystems) was used to visualize the images.
Images of teratoma haematoxylin and eosin and immunohistochemistry stainings were
acquired using BX61 Olympus Microscope (Olympus), and PRECiV™ 2D Image and
Measurement Software (Olympus) was used to visualize the images. The Image J (version
1.15) was used for further analysis [48].

Flow cytometry sample preparation and analysis

Cells were collected and they were purified based on FACS using BD FACSAria II Cell Sorter
(BD Bioscience). Cells were sorted at 3,500 events/second, and at maximum flow rate of 4
and 85 μm nozzle. 5 μg/ml of DAPI (Biotium; Cat #BT-40043) was used as a viability dye. For
flow cytometry analysis of pluripotency exit, cells were disaggregated using TrypLE Express,
and they were incubated at 4ºC for 20 minutes with SSEA1-eFluor 660 Monoclonal Antibody
(1:200, Invitrogene, Cat #50-8813-42). BD LSRFortessa Cell Analyzer or BD LSR II Flow
Cytometer (BD Bioscience) were used for Flow Cytometry Analysis experiments. Flow
cytometry data were analyzed using FlowJo X v10.0.7 (BD Biosciences).

RNA isolation, cDNA synthesis and quantitative RT-PCR

Total RNA isolation from cells was performed using the RNeasy Plus Mini Kit (Qiagen; Cat
#74136) or RNeasy Micro Kit (Qiagen; Cat #74004) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
2 μg of total RNA was retro-transcribed using Transcriptor First Strand cDNA Synthesis Kit
(Roche; Cat #04897030001) following the manufacturer's instructions.

Quantitative RT-PCR was performed in triplicates for each sample, and SYBR Green I Master
Kit (Roche; Cat #04887352001) was used to carry out the reaction. qRT-PCR was performed
using the LightCycler 480 system (Roche). Relative expression levels were calculated as $2^{-\Delta CT}$
normalized with the average CT of the housekeeping gene Tbp or Gapdh. Oligos sequences
are found in supplementary table 1.

Chromatin Immunoprecipitation (ChIP)

$4\text{-}6\text{x}10^7$ mESCs were cross-linked by incubating them in DPBS (Gibco, Cat #14190094)
supplemented with 1% formaldehyde (Sigma-Aldrich; Cat #252549) for 10 min rocking at
room temperature. Cross-linking was stopped using 125 mM Glycine (Sigma-Aldrich, Cat
#G8790) rocking for 5 minutes at room temperature. Fixed cells were then washed twice with
ice-cold DPBS, and scrapped and collected using DPBS supplemented with
protease/phosphatase inhibitor cocktail composed by 1X cOmplete™, EDTA-free Protease
Inhibitor Cocktail (Roche, Cat #11873580001), 1 mM PMSF, 1mM Sodium Ortovanadate and
20 mM β-Glycerol phosphate. Cells were centrifuged at 3200g for 5 minutes at 4ºC. Cells were
lysed by resuspending cell pellet in ice-cold ChIP buffer (1 volume of SDS buffer [100mM NaCl,
50mM Tris-HCl pH 8.1, 5mM EDTA pH 8, 0.5% SDS] and 0.5 volume of triton dilution buffer
[100mM Tris-HCl pH 8.6, 100mM NaCl, 5mM EDTA pH 8, 5% Triton X-100]) supplemented with
protease/phosphatase inhibitor cocktail, and samples were sonicated using sonication beads
(Diagenode; Cat #C01020031) and the Bioruptor Pico Sonicator (Diagenode; Cat #B01060010)
for 20 cycles (each cycle 30 seconds on/30 seconds off) or till DNA fragments have 100-300bp
size. Samples were centrifuged at 16000g for 20 minutes at 4ºC. Chromatin was quantified,
and 30 μg of chromatin was incubated with every 5μg of antibody rotating for 16 hours (or
overnight) at 4ºC. The following antibodies were used: H3K4me3 (Abcam, Cat #ab8580),

H3K27me3 (Millipore, Cat ##07-449), H3K27Ac (Abcam, Cat #ab4729), H3K4me1 (Abcam, Cat
#ab8895). Samples were pulled down by incubation with Protein A-Sepharose™ CL-4B
(previously hydrated and blocked with 0.05% BSA) for 3h at 4ºC in rotation. Sample was
washed with a low salt washing buffer (50mM HEPES pH 7.5, 140mM NaCl and 1% Triton)
three times, one wash with a high salt wash buffer (50mM HEPES pH 7.5, 500mM NaCl and
1% Triton) and one wash with TE pH 8 (10mM Tris-HCl pH8, 1mM EDTA). Samples were then
eluted in 1% SDS and 100mM $NaHCO_3$. Samples were descrosslinked by incubating for 16
hours (or overnight) at 65ºC shaking at 450 rpm, and were treated with Proteinase K for 1h at
45ºC. DNA was purified using QIAquick PCR Purification Kit (Qiagen, Cat #28106), following
the manufacturer's instructions.

For ChIP followed by sequencing (ChIP-seq) Purified DNA concentration and integrity were
determined using Agilent Bioanalyzer (Agilent Technologies; Cat #G2939BA). Libraries were
prepared using standard protocols. Chromatin was sequenced using Illumina HiSeq platform
(Illumina, Inc.) (50bp single-end reads). Samples sequencing depth range was the following:
(i) H3K4me3: $2.56\text{-}2.74 \times 10^7$ reads, (ii) H3K27me3: $6.19\text{-}7.2 \times 10^7$ reads, (iii) H3K27Ac: 9.11-
10.65×10^7 reads and (iv) H3K4me1: $9.76\text{-}10.58 \times 10^7$ reads.

ChIP-seq data analysis

Quality control was performed on raw data with FASTQC tool. Raw reads were trimmed to
remove adapters presence with Trimgalore (v0.6.6) [49]. Default parameters were used except
for a minimum quality of 15 (Phred score) and an adapter removal stringency of 3bp overlap.
For H3K4me3, H3K27me3 and H3K4me1 data, trimmed reads were aligned to the reference
genome with Bowtie2 (v2.4.4) which was executed with default parameters [50]. Required
genome index was built with corresponding GRCm38 fasta file retrieved from Ensembl
(http://ftp.ensembl.org/pub/release-102/). Multimapped reads and those exhibiting MAPQ <
20 were removed. Randomly placed multi mappers were removed from the mapped reads.
For all cases, duplicated reads were marked with SAMtools (v1.15) [51]. Drosophila
Melanogaster spike-ins, present in H3K27me3, were discarded. NSC and RSC quality metrics
were computed with PhantomPeakQualTools (v1.2) [52]. ENCODE blacklisted regions (mm10
v2) were removed prior to peak calling. BigWig files were individually generated using
deepTools (v3.5.1) bamCoverage with -ignoreDuplicates -binSize 10 -smoothLength 30 -
effectiveGenomeSize   2308125349   -normalizeUsing   RPGC   and   -extendReads
Fragment_Length options [53]. The effective genome size was particularized to a read length of
50bp   and   directly   retrieved   from   deepTools   web   site
(https://deeptools.readthedocs.io/en/develop/content/feature/effectiveGenomeSize.html).
Fragment_Length was retrieved from PhantomPeakQualTools results. For all histone marks
except for H3K4me3, peak calling was conducted by means of epic2 (v0.0.52) with -effective-
genome-fraction 0.8452 -fragment-size *Fragment_Length* options and chromosome sizes
only referring to canonical chromosomes [54]. For H3K4me3, MACS2 (v2.2.7.1) was used to
identify peaks with -nomodel -extsize *Fragment_Length* -g 2308125349 options [55]. The

corresponding input sample was used in all peak calling computations. For the histone marks,
peaks were called with adjusted p-values (FDR) < 0.05.

Peak annotation was performed with ChIPseeker R (v4.2.1) package (v1.34.1) considering a
TSS region range between 5000 bp upstream and 100 bp downstream (Yu et al., 2015).
Required TxDb object was generated from the Ensembl GRCm38 gtf file retrieved
(http://ftp.ensembl.org/pub/release-102/). A consensus peakset was obtained from the
three biological replicates per histone mark and condition (IκBα-WT or IκBα-KO) by identifying
overlapping peaks in at least two out of the three replicates.

Differential binding analysis (DBA) was conducted with DiffBind (v3.8.4) per histone mark [56].
Default parameters were used except for the summits parameter which was set to 500bp for
all histone marks except for H3K4me3 (150bp) in order to consider, for testing, intervals of
1000bp or 300bp respectively. Summit values were selected to have interval widths between
the minimum and first quartile peak width values for each histone mark. EdgeR was the
statistical method used for all the three analyses. Differentially bound regions (DBRs)
between IκBα-KO and IκBα-WT samples were called with adjusted p-values (FDR) < 0.05.
Identified DBRs with any annotated gene were plotted with EnrichedHeatmap R package
(v1.18.1) [57]. To summarize replicates per condition, normalizeToMatrix function was used
based on corresponding BigWig files in w0 mean mode in 50bp windows. Tracks visualization
were obtained by means of Integrative Genomics Browser (IGV) tool [58].

Enhancers activity identification

Consensus peaksets derived from the three H3K27ac and H3K4me1 IκBα-WT mESCs replicates
were used to identify poised and active enhancers in an IκBα-WT scenario. Putative poised
enhancers were defined as those regions with H3K4me1 but no H3K27ac peaks. Putative
active enhancers were defined as those regions with H3K27ac and presence/absence
H3K4me1 peaks. To assess any differential enhancers activity in IκBα-KO vs IκBα-WT mESCs
cultured in Serum/LIF, differentially bound regions and consensus peaksets (IκBα-KO)
identified in H3K4me1 and H3K27ac were required. Putative gained or lost poised/active
enhancers in IκBα $^{-/-}$ were obtained with following criteria: (i) for gained/lost poised enhancers
in IκBα $^{-/-}$: a differential H3K4me1 increase/decrease and absence of H3K27ac peaks in IκBα $^{-/-}$
consensus peakset, (ii) for gained/lost active enhancers (without H3K4me1) in IκBα-KO: a
differential H3K27ac increase/decrease and absence of H3K4me1 in IκBα-KO consensus
peakset and (iii) for gained/lost active enhancers (with H3K4me1) in IκBα-KO: a differential
H3K27ac increase/decrease and presence of H3K4me1 in IκBα $^{-/-}$ consensus peakset.

A maximum gap of 1000bp was allowed for checking the overlap between two different
histone marks. For this purpose, the 'subsetByOverlaps' function from the IRanges R package
was used (v.2.34.1).

RNA-seq experiments

Total RNA from three independent clones from mESCs, 48h EBs and 96h EBs was isolated
using the RNeasy Plus Mini Kit (Qiagen; Cat #74136) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
RNA integrity was addressed by agarose gel and Agilent Bioanalyzer (Agilent Technologies;
Cat #G2939BA). Libraries sequenced using Illumina HiSeq 2500 (Illumina, Inc.) (125bp paired-
end reads). Samples sequencing depth ranged between 35M and 52M reads (average 41M
reads) per sample.

RNA-seq data analysis

Quality control was performed on raw data with the FASTQC tool (v0.11.9). Raw reads were
trimmed to remove adapter presence with Trimgalore (v0.6.6) [49]. Default parameters were
used except for a minimum quality of 15 (Phred score) and an adapter removal stringency of
3bp overlap. Trimmed reads were aligned to reference the genome with STAR aligner tool
(v2.7.8). STAR was executed with default parameters except for the number of allowed
mismatches which was set to 1. Required genome index was built with corresponding
GRCm38 gtf and fasta files retrieved from Ensembl (http://ftp.ensembl.org/pub/release-
102/). Obtained BAM files with uniquely mapped reads were considered for further analysis.
Raw gene expression was quantified using featureCounts tool from subRead software (v2.0.1)
with exon as feature [59]. The raw counts matrix was imported into the R Statistical Software
environment (v4.2.1) for downstream analysis. Raw expression matrix included 55,487 genes
per 18 samples in total. Experimental design considered three timepoints: mouse ESCs, EBs
at 48h and EBs at 96h. Each time point included 6 samples distributed in two conditions: 3
IκBα-WT and 3 IκBα-KO. Prior to statistical analysis, those genes with less than 10 raw counts
across the 6 samples under test were removed. After pre-filtering, 21,843 genes (mESCs),
21,424 genes (EBs 48h) or 21,664 genes (EBs 96h) were available for testing. For visualization
purposes, counts were normalized by the variance-stabilizing transformation method as
implemented in DESeq2 R package [60] (v1.38.3). Differential expression analysis (DEA) was
conducted with DESeq2. Each time point was independently analyzed. Fitted statistical model
included sample conditions as covariable with IκBα-WT as the reference. Obtained log2 fold
change values were shrunken with apeglm shrinkage estimator R package (v1.20.0) [61]. Raw p-
values were adjusted for multiple testing using the Benjamini-Hochberg False Discovery Rate
(FDR) [62]. Differentially Expressed Genes (DEGs) between IκBα-KO and IκBα-WT samples were
called with adjusted p-values (FDR) < 0.05 and absolute shrunken log2 Fold change > 1. Data
visualization was performed with the ggplot2 (v3.4.1).

Genome-wide DNA methylation samples preparation

DNA from frozen mESCs pellets was extracted using DNeasy Blood and Tissue Kit (Qiagen
GmbH, Hilden, Germany). Purified genomic DNA was quantified with Qubit (Invitrogen,
Carlsbad, CA, USA) according to manufacturer's instructions. Infinium Mouse Methylation
BeadChip (Illumina, Inc., San Diego, CA, USA) arrays were used to profile DNA methylation

genome wide. This platform allows over 285,000 methylation sites per sample to be
interrogated at single-nucleotide resolution. The samples were bisulfite converted using EZ
DNA Methylation-Gold™ Kit (Zymo Research, CA, USA) and were hybridized in the array
following the manufacturer's instructions.

DNA methylation data analysis

The DNA methylation profile of the studied samples was assessed using the Infinium Mouse
Methylation BeadChip Array (~285,000 methylation sites) as previously described [63]. Briefly,
raw signal intensities were obtained with GenomeStudio Software 2011.1 (Illumina) and DNA
methylation beta values were computed from raw IDAT files using GenomeStudio default
normalization with control probes and background subtraction. Quality control steps to
remove erratic probe signals were performed within the R statistical environment (v4.0.3).
We removed probes with detection p value > 0.01, genotyping probes and manufacturing
flagged (MFG) probes described in the Illumina manifest file
(https://support.illumina.com/downloads/infinium-mouse-methylation-manifest-file.html).
The differentially methylated probes (DMPs) between IκBα-WT and IκBα-KO samples were
computed separately for each time point (mESCs and EBs at 96h) by deriving a linear model
with the limma R package (v3.46.0). Each condition included 3 samples per time point. DMPs
with adjusted p-value (FDR) <0.05 and absolute mean methylation beta value difference
between conditions >0.3 were considered significant. DNA methylation analysis was
performed using the mm10 mouse genome reference build and the complete annotation was
downloaded from the annotated manifest file
(http://zwdzwd.github.io/InfiniumAnnotation#mouse) [36].

Functional analysis

Overrepresentation analysis was applied over lists of selected genes derived from RNA-seq
data (DEGs) or from ChIP-seq data (differentially bound regions). The Gene Ontology
(Biological Process ontology, GO BP terms), KEGG PATHWAY and WikiPathways databases for
Mus Musculus [64–66] were interrogated by means of clusterProfiler R package (v4.6.2) (Wu et
al., 2021). Corresponding Entrez identifiers were used. Benjamini-Hochberg procedure was
used to obtain adjusted p-values. Obtained GO BP terms were simplified using the simplify
function from clusterProfiler with default parameters. Overrepresented terms or processes
were called with adjusted p-values (FDR) < 0.05.

Gene Set Enrichment Analysis (GSEA) was performed for mESCs against the three cell states
signatures defined by other authors [67]. Briefly, each signature was composed of 200 genes
uniquely expressed in each state obtained from bulk RNA-seq performed per state. For this
purpose, the complete list of genes from mESCs samples (21,843 genes) was ranked based on
the shrunken log2 Fold Change obtained from DEA. GSEA was conducted through the

fgseaMultilevel function from fgsea R package (v1.24.0) [68] with default parameters.

Enrichment plots were generated with the same package.

Additionally, the testEnrichment function from SeSAMe R package (v1.14.2) [69] was used for

conducting functional analysis with default parameters over the list of DMPs. Probe design,

transcription factor binding site and histone modifications consensus database sets, included

in the same package, were interrogated.

Gene Set Variation Analysis (GSVA) R package (v1.46.0) with default parameters were used to

obtain Z-score values for genes annotated to endoderm (GO:0001706), mesoderm

(GO:0001707) and ectoderm (GO:0001705) GO BP terms [70].

Statistical analysis

Statistical analysis was performed with GraphPad Prism v.8.0.1. (GraphPad Software, Inc.). Unless specified, the comparison between two groups was performed with unpaired two-sided t-test. A p-value < 0.05 was considered significant.

DATA AVAILABILITY

All sequencing data is deposited at GEO under a SuperSeries with the accession number GSE239565)[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE239565]. Individual SubSeries can be found at: GSE239563 (RNA-seq data), GSE239564 (ChIP-seq data) and GSE239562 (Methylation array data).

CODE AVAILABILITY

Scripts used to process the generated bulk RNA-seq and ChIP-seq data are available in Github repository: https://github.com/BigaSpinosaLab/PAPER_IkBa_Naive_Pluripotency

ACKNOWLEDGEMENTS

We would like to acknowledge all members of the Bigas/Espinosa labs for helpful discussions. We are grateful for technical support to CRG/UPF Flow Cytometry Unit, CRG Genomics and Advanced Light Microscopy Units. This work has been supported by Spanish Ministry of Science and Innovation (PID2019-104695RB-I00, PLEC2021-007518, PDC2021-120817-I00), Generalitat de Catalunya (2021SGR 39) and Departament de Salut (SLT002/16/00299) to AB; the Spanish Ministry of Science and Innovation (PID2019-108322GB-100 and PID2022-142679NB-I00) to LDC; the Spanish Ministry of Science and Innovation (PID2021-123383NB-I00.) and the Agencia de Gestió d'Ajuts Universitaris i de Recerca (2021 SGR 01222) to BP. L.G.P has been a recipient of FI AGAUR fellowship (2019 FI-B 00151/2020 FI_B1 00130) from Generalitat de Catalunya. D.A-V was funded by the FIS fellowship (FI20/00130) from Instituto Carlos III. M.B. received funding from the Ramón Areces Foundation. M.M. is a recipient of a grant from the Instituto Carlos III, grant number CA22/00011 (co-funded by the European Social Fund Plus, ESF+ and by the European Union).

AUTHOR CONTRIBUTION

L.G.P, L.E and A.B designed the study. L.G.P, D.A-V, M.B, A.I, J.G, J.B, D.A.E, C.A.G.P, C.B, V.R.C and A.V performed experiments. L.G.P, M.M, M.B and C.A.G.P performed bioinformatics analysis. L.G and A.B wrote the manuscript with input from all authors. C.B, A.V, A.V, P.M,

	FIGURE LEGENDS

Figure 1. IκBα is highly expressed in pluripotent cells. A. Schematic of pluripotent states in
mouse embryonic stem cells (mESCs). mESCs were maintained regularly in naïve pluripotency
by culturing them in serum/LIF medium, or they were polarized towards the ground state of
naïve pluripotency by culturing them for two passages in LIF plus Gsk3β (CHIR99021 or Chiron)
and MEK (PD0325091) inhibitors (2i/LIF). mESCs were further differentiated towards the
primed pluripotent state (epiblast-like cells) by culturing mESCs in a medium containing
20ng/mL Activin A and 10ng/mL Fgf2 for 120h. **B-C.** Relative mRNA levels (based on qPCR
experiments) of the canonical IκB genes (*Nfkbia, Nfkbib, Nfkbie*) (**B**) or Nf-κB effector genes
(*Rela, c-Rel, Relb, Nfkb1, Nfkb2*) (**C**) across the different states of pluripotency. Expression was
normalized using the house-keeping gene *Tbp* relative to 2i/LIF. Data from three independent
experiments. Dots indicate mean values and error bars refer to ± standard deviation (SD). **D.**
Western blot analysis of IκBα protein distribution in cytoplasm, nucleoplasm (Nucleus) or
chromatin in cells at the ground-state of naïve pluripotency (2i/L), naïve pluripotency (S/L) or
primed pluripotency (Epi). **E.** Transcriptome and proteome differences across naïve (2i/LIF)
and primed (Epiblast-like cells or EpiSCs) pluripotent states. Enriched transcripts and proteins
in naïve state are highlighted in red (Log$_2$ fold-change < -1). Enriched transcripts and proteins
in the primed state are marked in dark gray (Log$_2$ fold-change > 1). Data from [33]. **F.** Heatmap
showing the expression levels of pluripotency (*Dppa4, Pou5f1, Sox2, Prdm14, Klf4, Zfp42,*
Nanog), differentiation (*Mesp1, Gata4, Gata6, Lhx1, Wnt3a, Fgf5, Nes, Axin2, Wnt3, Eomes,*
Foxa2, T, Mixl1, Lef1) NF-κB (*Nfkbia, Nfkbib, Nfkbie, Nfkbiz, Rela, c-Rel, Relb*) and Polycomb
(*Ezh1, Ezh2, Eed, Epop, Jarid2, Suz12, Mtf2, Aebp2*) genes from the RNAseq samples at mESCs
(Serum/LIF) and differentiating cells (embryoid bodies) at 48h and 96h of IκBα-WT cells.
Normalized counts based on z-score are represented.

Figure 2. The absence of IκBα hinders the exit from the pluripotent state and prevents the
activation of differentiation programs. A. Representative immunofluorescence of 216h EBs
stained with the pluripotency markers NANOG and OCT3/4. **B.** Quantification of total number
of alkaline phosphatase (AP) colonies after 216h EB differentiation and additional culture of
96h in Serum/LIF (S/L) medium. Dots represent values from two independent experiments
from three different IκBα-WT or IκBα-KO clones. Unpaired two-sided t-test applied.
Representative images from AP colonies are included in the upper panel. **C.** PC1 and PC2 from
Principal Component Analysis (PCA) of normalized RNA-seq data from 3 different time points
(mESCs, 48h EBs and 96h EBs) of IκBα-WT and IκBα-KO cells (three independent clones from
each genotype). **D.** Heatmap showing the expression levels of naïve pluripotency genes
(*Gab1, Sox2, Esrrb, Zfp42, Dppa4, Klf2, Prdm14, Klf4, Tfcp2l1*) from the RNAseq samples of
IκBα-WT and IκBα-KO EBs at 96h. Normalized counts based on z-score are represented. **E-G.**
Z-score values for gene sets referring to Endoderm (E), Mesoderm (F) and Ectoderm (G)
formation from The Gene Ontology database (GO:0001706, GO:0001707 and GO:0001705
terms, respectively) and obtained from   corresponding expression levels at mESCs, 48h and
96h EBs IκBα-WT or IκBα-KO cells (three independent clones of each genotype). Loess curves

are also represented with 95% confidence intervals (shadowed area surrounding the curves).
H. Schematic of teratoma formation assay. 5×10^5 IκBα-WT or IκBα-KO mESCs cultured in
Serum/LIF were injected intramuscularly in the leg of immunocompromised NSG mice. 6
weeks after the transplant, teratomas were formed, and mice were euthanized for further
teratoma analysis. **I.** Representative images of immunohistochemistry for OCT3/4 in IκBα-WT
(left panel) or IκBα-KO (right panel) teratomas. Teratomas were derived from three
independent clones of mESCs of both genotypes. **J.** Percentage of OCT3/4$^+$ cells in teratomas.
Positive cells were counted from five different microscope fields from each IκBα-WT and IκBα-
KO clone. Unpaired two-sided t-test was performed. Bars indicate mean values and error bars
refer to ± SD.

Figure 3. IκBα-KO mESCs retain the ground-state of naïve pluripotency under Serum/LIF
culture. A. (Upper panel) Schematics of mESCs differentiation towards Epiblast Stem Cells
(EpiSCs). mESCs cultured in Serum/LIF were then seeded in N2B27 medium supplemented
with 10ng/mL Fgf2 and 20ng/mL Activin A for 120h (see Materials and Methods section for
further information). (Bottom panel) Relative RNA levels (based on qPCR experiments) of
naïve pluripotency genes (*Nanog, Sox2, Rex1, Klf4, Gbx2, Tbx3, Dppa3*) in IκBα-WT or IκBα-KO
EpiSCs at 120h. RNA levels were normalized based on *Tbp* expression relative to IκBα-WT.
Each dot represents an independent clone either from IκBα-WT or IκBα-KO genotypes.
Unpaired two-sided t-test applied. Horizontal bars indicate mean values and error bars refer
to ± SD. **B.** GSEA results comparing IκBα-KO vs IκBα-WT mESCs cultured in Serum/LIF against
ground (2i/LIF) and naïve (Serum/LIF) state pluripotency signatures retrieved from [35].
Adjusted p-value by Benjamini-Hochberg procedure and Normalized Enrichment Score (NES)
indicated. **C.** Relative RNA levels of the naïve pluripotency genes *Rex1 (Zfp42), Klf2* and *Tbx3*
upon culture of IκBα-WT either in Serum/LIF or two passages in 2i/LIF. 2i/LIF IκBα-WT mESCs
were compared with IκBα-KO mESCs cultured in Serum/LIF. Unpaired two-sided t-test was
applied to calculate statistical significance. Each dot represents an independent clone from
each of the two genotypes. **D.** Representative immunofluorescence images of DNA-
methylation-related mark 5-methylcytosine (5mC) in IκBα-WT (left) and IκBα-KO (right)
mESCs cultured in Serum/LIF. **E.** DNA methylation profile (using DNA methylation arrays [36]) in
IκBα-WT vs IκBα-KO mESCs (upper) and 96h EBs (bottom panel) showing the distribution
density of mean ß-values from all 261,220 CpGs under test per condition.

Figure 4. Absence of IκBα reinforces the pluripotency program at the epigenetic and
transcriptional levels. A. ChIP-seq Enrichment Analysis (ChEA) of differentially expressed
genes (DEGs; adjusted p-value<0.05) in IκBα-KO mESCs. P-value was calculated using Fisher
exact test. **B.** Number of differentially bound regions (adjusted p-value <0.05, FDR) for
H3K4me3 and H3K27me3 histone marks in IκBα $^{-/-}$ vs IκBα $^{+/+}$ mESCs cultured in Serum/LIF.
Enriched or reduced regions in IκBα $^{-/-}$ mESCs are distinguished (log$_2$ Fold-change >0 or <0,
respectively). Overrepresented WikiPathways, for (left) H3K4me3 and (right) H3K27me3 are
indicated. Annotated genes to differential regions showing H3K4me3 enrichment or
H3K27me3 reduction were considered. A one-sided hypergeometric test was conducted. **C.**

Venn Diagram of genes contained in regions with either increased H3K4me3 or reduced levels of H3K27me3. Statistical significance of the overlapping was calculated using the chi-squared method. Overrepresented pathways for genes containing increased H3K4me3 and reduced H3K27me3 levels are indicated (only pathways with an adjusted p-value<0.05 were considered). **D-E.** Representative genomic regions of naïve pluripotency genes having either differential enriched H3K4me3-only (D) or reduced H3K27me3-only (E) levels in IκBα-KO vs IκBα-WT mESCs cultured in Serum/LIF. Shadowed regions highlight the differential levels of the histone marks. **F.** Number of regions with differential enhancer activity in IκBα-KO vs IκBα-WT mESCs cultured in Serum/LIF. Increased/Reduced activity for (left) poised enhancers based on the H3K4me1 differential binding and absence of H3K27ac overlapping peaks in IκBα-KO mESCs and, (right) active enhancers based on the differential binding of H3K27ac and presence of H3K4me1 overlapping peaks in IκBα-KO. Differential binding based on adjusted p-value <0.05 (FDR). Increased or reduced regions in IκBα-KO are distinguished by $\log_2$ Fold-change >0 or <0, respectively, compared to IκBα-WT mESCs. Overrepresented WikiPathways for annotated genes to (left) poised enhancers with reduced activity and (right) active enhancers with increased activity in IκBα-KO vs IκBα-WT mESCs are indicated (adjusted p-value < 0.05, FDR). A one-sided hypergeometric test was conducted. **G.** Representative genomic regions of naïve pluripotency genes having an increase in active enhancer status (H3K27Ac enrichment) in IκBα-KO mESCs cultured in Serum/LIF. Shadowed regions highlight the differential levels of the histone mark. **H.** Heatmap showing the expression levels of the naïve pluripotency genes from the RNAseq samples of IκBα-KO and IκBα-WT mESCs cultured in Serum/LIF. Normalized counts (z-score) from genes are represented.

Figure 5. IκBα prevents the hyperactivation of naïve pluripotency program in an NF-κB-independent manner. A. Boxplots, violin plots and individual values representing the normalized gene expression levels in IκBα-WT or IκBα-KO mESCs, EBs 48h and EBs 96h. Considered genes are those annotated to NF-κB signaling pathway (KEGG PATHWAY mmu04064). Box Plots: center line, median; lower and upper hinges, first and third quartiles; whiskers, 1.5x interquartile range (IQR) extended to the largest/smallest value within 1.5xIQR; no outliers present. Unpaired two-sided Wilcoxon test applied. **B.** Schematic of chromatin-specific/NF-κB-deficient (IκBα$^{\Delta NF-κB}$) or NF-κB-specific/chromatin-deficient (IκBα$^{\Delta Chrom}$) IκBα mutants (Separation-Of-Function or SOF mutants). **C.** Pull-down of GST-H2A and GST-P50 with cell lysates from different SOF IκBα mutant mESCs. **D.** Western blot analysis of cytoplasmic protein extracts of doxycycline-inducible IκBα forms (i-IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-κB}$, i-IκBα$^{\Delta Chrom}$) after 16h of doxycycline induction. **E.** Western blot analysis of chromatin protein extracts of doxycycline-inducible IκBα forms (i-IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-κB}$, IκBα$^{\Delta Chrom}$) after 16h of doxycycline induction. **F.** Relative RNA levels of naïve pluripotency genes (*Zfp42, Klf2, Sox2, Tbx3*) in i-IκBα^{WT}, i-IκBα$^{\Delta NF-κB}$ and i-IκBα$^{\Delta Chrom}$ mESCs after doxycycline induction. Unpaired two-sided t-test was performed. Bars indicate mean values and error bars refer to ± SEM. **G.** Representative immunofluorescence images of DNA-methylation-related mark 5-

methylcytosine (5mC) in i-IκBα$^{\Delta WT}$ (upper panel), i-IκBα$^{\Delta NF-\kappa B}$ (medium panel), IκBα$^{\Delta Chrom}$
(bottom panel) mESCs cultured in Serum/LIF. **H.** Flow cytometry analysis of the pluripotency
surface marker SSEA-1 at 216h EBs in i-IκBα^{WT}, i-IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta Chrom}$ cells. **I.**
Quantification of Alkaline Phosphatase (AP) staining in 216h EBs upon reconstitution with the
different IκBα forms. Each dot represents an independent experiment. One-way ANOVA was
applied. Bars indicate mean values and error bars refer to ± SEM. Representative images of
Alkaline Phosphatase (AP) staining in 216h EBs are included in the upper panel. **J.**
Representative immunofluorescence for NANOG and OCT3/4 at 216h EBs in i-IκBα^{WT} (upper
panel), i-IκBα$^{\Delta NF-\kappa B}$ (medium panel) and i-IκBα$^{\Delta Chrom}$ (bottom panel) cells.

Figure S1. Depletion of IκBα in mESCs compromises the exit from pluripotency. Related to
Figure 2. A. Schematic of the approach followed to deplete IκBα protein in mESCs (upper
panel) and western blot analysis of three independent IκBα-WT and IκBα-KO mESC clones
(bottom panel). **B.** Flow cytometry plots of SSEA-1 from IκBα-WT or IκBα-KO EBs 216h. **C.**
Relative RNA levels (based on qPCR experiments) of naïve pluripotency genes (*Pou5f1, Gbx2,*
Klf2, Sox2, Rex1, Nanog) in IκBα-WT or IκBα-KO 216h EBs. RNA levels were normalized based
on *Tbp* expression relative to IκBα-WT. Each dot represents an independent clone either from
IκBα-WT or IκBα-KO genotypes. Unpaired two-sided t-test was applied. **D.** Schematic of
Alkaline Phosphatase (AP) staining carried out in EBs. IκBα-WT or IκBα-KO 216h EBs were
dissociated and seeded in Serum/LIF medium for another 96h. Attached cells were screened
for AP$^+$ total cell number. **E.** Violin plot and individual values showing the quantification of
relative size of 168h EBs. Each dot represents an embryoid body. Unpaired two-sided t-test
was applied. **F.** Volcano plot from differential expression analysis between IκBα-KO vs IκBα-
WT 96h EBs. Dashed vertical lines represent an absolute shrunken log$_2$ Fold Change = 1 and
dashed horizontal line represents an adjusted p-value = 0.05 (FDR).

Figure S2. Absence of IκBα favors the ground-state of naïve pluripotency in Serum/LIF
conditions. Related to Figure 3. A. Brightfield images of IκBα-WT or IκBα-KO EpiSCs at 120h.
Each picture corresponds to an independent clone from either IκBα-WT or IκBα-KO
genotypes. **B.** Bright-field microscopy images of IκBα-WT and IκBα-KO mESCs cultured in
Serum/LIF conditions. Two independent clones from each genotype are shown. **C.** DNA
methylation levels (mean ß-values) across chromosomes in IκBα-WT vs IκBα-KO mESCs.

Figure S3. Epigenetic rewiring characterization in IκBα-KO mESCs cultured in Serum/LIF.
Related to Figure 4. A-D. Volcano plots from differential binding analysis of H3K4me3 (A),
H3K27me3 (B), H3K4me1 (C) and H3K27Ac (D) in IκBα-WT vs IκBα-KO mESCs cultured in
Serum/LIF. Representative genes with enriched or reduced levels of H3K4me3 (A), H3K27me3
(B), H3K4me1 (C) and H3K27Ac (D) marks in IκBα-KO mESCs are highlighted.

Figure S4. Reversion of IκBα-KO-related hyper pluripotent status is NF-κB independent.
Related to Figure 5. A. Relative expression level of Nfkbia gene after overnight doxycycline
induction in i-IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-\kappa B}$ and IκBα$^{\Delta Chrom}$ mESCs. **B.** Schematic representation of

reconstitution of IκBα-KO mESCs with either of the two doxycycline-inducible SOF mutants (i-
IκBα$^{\Delta NF-κB}$ or IκBα$^{\Delta Chrom}$). mESCs were then kept in Serum/LIF medium supplemented with
1μg/mL Doxycycline for 4 consecutive passages. mESCs were characterized at the pluripotent
level and then their potential to be further differentiated was addressed. **C.** Percentage of
differentiated cells (determined as SSEA-1.) at 216h EBs 216h derived from i-IκBα$^{\Delta WT}$, i-
IκBα$^{\Delta NF-κB}$ and IκBα$^{\Delta Chrom}$ mESCs. **D.** Quantification of EB differentiation status (percentage of
pluripotent cells) at 216h after induction of SOF IκBα mutants. Data from three independent
experiments.

REFERENCES

1. Festuccia, N., Osorno, R., Wilson, V., and Chambers, I. (2013). The role of pluripotency gene regulatory network components in mediating transitions between pluripotent cell states. Curr. Opin. Genet. Dev. *23*, 504–511.

2. Graf, T., and Stadtfeld, M. (2008). Heterogeneity of embryonic and adult stem cells. Cell Stem Cell *3*, 480–483.

3. Martinez Arias, A., and Brickman, J.M. (2011). Gene expression heterogeneities in embryonic stem cell populations: origin and function. Curr. Opin. Cell Biol. *23*, 650–656.

4. Pera, M.F., and Rossant, J. (2021). The exploration of pluripotency space: Charting cell state transitions in peri-implantation development. Cell Stem Cell *28*, 1896–1906.

5. Wang, X., and Wu, Q. (2022). The Divergent Pluripotent States in Mouse and Human Cells. Genes *13*. 10.3390/genes13081459.

6. Ying, Q.-L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P., and Smith, A. (2008). The ground state of embryonic stem cell self-renewal. Nature *453*, 519–523.

7. Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H., Nichols, J., Kranz, A., Stewart, A.F., Smith, A., et al. (2012). The transcriptional and epigenomic foundations of ground state pluripotency. Cell *149*, 590–604.

8. Boroviak, T., Loos, R., Lombard, P., Okahara, J., Behr, R., Sasaki, E., Nichols, J., Smith, A., and Bertone, P. (2015). Lineage-Specific Profiling Delineates the Emergence and Progression of Naive Pluripotency in Mammalian Embryogenesis. Dev. Cell *35*, 366–382.

9. Brons, I.G.M., Smithers, L.E., Trotter, M.W.B., Rugg-Gunn, P., Sun, B., Chuva de Sousa Lopes, S.M., Howlett, S.K., Clarkson, A., Ahrlund-Richter, L., Pedersen, R.A., et al. (2007). Derivation of pluripotent epiblast stem cells from mammalian embryos. Nature *448*, 191–195.

10. Tesar, P.J., Chenoweth, J.G., Brook, F.A., Davies, T.J., Evans, E.P., Mack, D.L., Gardner, R.L., and McKay, R.D.G. (2007). New cell lines from mouse epiblast share defining features with human embryonic stem cells. Nature *448*, 196–199.

11. Weinberger, L., Ayyash, M., Novershtern, N., and Hanna, J.H. (2016). Dynamic stem cell states: naive to primed pluripotency in rodents and humans. Nat. Rev. Mol. Cell Biol. *17*, 155–169.

12. Guo, G., Yang, J., Nichols, J., Hall, J.S., Eyres, I., Mansfield, W., and Smith, A. (2009). Klf4 reverts developmentally programmed restriction of ground state pluripotency. Development *136*, 1063–1069.

13. Takahashi, S., Kobayashi, S., and Hiratani, I. (2018). Epigenetic differences between

naïve and primed pluripotent stem cells. Cell. Mol. Life Sci. *75*, 1191–1203.

14. Bao, S., Tang, F., Li, X., Hayashi, K., Gillich, A., Lao, K., and Surani, M.A. (2009).
Epigenetic reversion of post-implantation epiblast to pluripotent embryonic stem cells.
Nature *461*, 1292–1295.

15. Habibi, E., Brinkman, A.B., Arand, J., Kroeze, L.I., Kerstens, H.H.D., Matarese, F.,
Lepikhov, K., Gut, M., Brun-Heath, I., Hubner, N.C., et al. (2013). Whole-genome
bisulfite sequencing of two distinct interconvertible DNA methylomes of mouse
embryonic stem cells. Cell Stem Cell *13*, 360–369.

16. Leitch, H.G., McEwen, K.R., Turp, A., Encheva, V., Carroll, T., Grabole, N., Mansfield, W.,
Nashun, B., Knezovich, J.G., Smith, A., et al. (2013). Naive pluripotency is associated
with global DNA hypomethylation. Nat. Struct. Mol. Biol. *20*, 311–316.

17. Singer, Z.S., Yong, J., Tischler, J., Hackett, J.A., Altinok, A., Surani, M.A., Cai, L., and
Elowitz, M.B. (2014). Dynamic heterogeneity and DNA methylation in embryonic stem
cells. Mol. Cell *55*, 319–331.

18. Factor, D.C., Corradin, O., Zentner, G.E., Saiakhova, A., Song, L., Chenoweth, J.G.,
McKay, R.D., Crawford, G.E., Scacheri, P.C., and Tesar, P.J. (2014). Epigenomic
comparison reveals activation of "seed" enhancers during transition from naive to
primed pluripotency. Cell Stem Cell *14*, 854–863.

19. Buecker, C., Srinivasan, R., Wu, Z., Calo, E., Acampora, D., Faial, T., Simeone, A., Tan,
M., Swigut, T., and Wysocka, J. (2014). Reorganization of enhancer patterns in
transition from naive to primed pluripotency. Cell Stem Cell *14*, 838–853.

20. Gerondakis, S., Grumont, R., Gugasyan, R., Wong, L., Isomura, I., Ho, W., and Banerjee,
A. (2006). Unravelling the complexities of the NF-kappaB signalling pathway using
mouse knockout and transgenic models. Oncogene *25*, 6781–6799.

21. Kaltschmidt, C., Greiner, J.F.W., and Kaltschmidt, B. (2021). The Transcription Factor
NF-κB in Stem Cells and Development. Cells *10*. 10.3390/cells10082042.

22. Espín-Palazón, R., and Traver, D. (2016). The NF-κB family: Key players during
embryonic development and HSC emergence. Exp. Hematol. *44*, 519–527.

23. Li, S., Sun, J., Zhang, B.-W., Yang, L., Wan, Y.-C., Chen, B.-B., Xu, N., Xu, Q.-R., Fan, J.,
Shang, J.-N., et al. (2024). ATG5 attenuates inflammatory signaling in mouse embryonic
stem cells to control differentiation. Dev. Cell. 10.1016/j.devcel.2024.01.026.

24. Torres, J., and Watt, F.M. (2008). Nanog maintains pluripotency of mouse embryonic
stem cells by inhibiting NFkappaB and cooperating with Stat3. Nat. Cell Biol. *10*, 194–
201.

25. Dutta, D., Ray, S., Home, P., Larson, M., Wolfe, M.W., and Paul, S. (2011). Self-renewal
versus lineage commitment of embryonic stem cells: protein kinase C signaling shifts
the balance. Stem Cells *29*, 618–628.

26. Zhang, Q., Lenardo, M.J., and Baltimore, D. (2017). 30 Years of NF-κB: A Blossoming of
Relevance to Human Pathobiology. Cell *168*, 37–57.

27. Mulero, M.C., Ferres-Marco, D., Islam, A., Margalef, P., Pecoraro, M., Toll, A., Drechsel,
N., Charneco, C., Davis, S., Bellora, N., et al. (2013). Chromatin-bound IκBα regulates a
subset of polycomb target genes in differentiation and cancer. Cancer Cell *24*, 151–166.

28. Marruecos, L., Bertran, J., Guillén, Y., González, J., Batlle, R., López-Arribillaga, E.,
Garrido, M., Ruiz-Herguido, C., Lisiero, D., González-Farré, M., et al. (2020). IκBα
deficiency imposes a fetal phenotype to intestinal stem cells. EMBO Rep. *21*, e49708.

29. Marruecos, L., Bertran, J., Álvarez-Villanueva, D., Mulero, M.C., Guillén, Y., Palma, L.G.,
Floor, M., Vert, A., Arce-Gallego, S., Pecharroman, I., et al. (2021). Dynamic chromatin
association of IκBα is regulated by acetylation and cleavage of histone H4. EMBO Rep.
22, e52649.

30. Thambyrajah, R., Maqueda, M., Fadlullah, M.Z., Proffitt, M., Neo, W.H., Guillén, Y.,
Casado-Pelaez, M., Herrero-Molinero, P., Brujas, C., Castelluccio, N., et al. (2024). IκBα
controls dormancy in hematopoietic stem cells via retinoic acid during embryonic
development. Nat. Commun. *15*, 4673.

31. Aguilera, C., Hoya-Arias, R., Haegeman, G., Espinosa, L., and Bigas, A. (2004).
Recruitment of IκBα to the *hes1* promoter is associated with transcriptional repression.
Proceedings of the National Academy of Sciences *101*, 16537–16542.

32. Alvarez-Villanueva, D., Galan, L., Bertran, J., and Floor, M. (2023). Separation-of-
function mutants reveal the NF-κB-independent involvement of IκBα in the regulation
of stem cell and oncogenic programs. bioRxiv.

33. Atlasi, Y., Jafarnejad, S.M., Gkogkas, C.G., Vermeulen, M., Sonenberg, N., and
Stunnenberg, H.G. (2020). The translational landscape of ground state pluripotency.
Nat. Commun. *11*, 1617.

34. Tian, T.V., Di Stefano, B., Stik, G., Vila-Casadesús, M., Sardina, J.L., Vidal, E., Dasti, A.,
Segura-Morales, C., De Andrés-Aguayo, L., Gómez, A., et al. (2019). Whsc1 links
pluripotency exit with mesendoderm specification. Nat. Cell Biol. *21*, 824–834.

35. Ghimire, S., Van der Jeught, M., Neupane, J., Roost, M.S., Anckaert, J., Popovic, M., Van
Nieuwerburgh, F., Mestdagh, P., Vandesompele, J., Deforce, D., et al. (2018).
Comparative analysis of naive, primed and ground state pluripotency in mouse
embryonic stem cells originating from the same genetic background. Sci. Rep. *8*, 5884.

36. Zhou, W., Hinoue, T., Barnes, B., Mitchell, O., Iqbal, W., Lee, S.M., Foy, K.K., Lee, K.-H.,
Moyer, E.J., VanderArk, A., et al. (2022). DNA methylation dynamics and dysregulation
delineated by high-throughput profiling in the mouse. Cell Genom *2*.
10.1016/j.xgen.2022.100144.

37. Choi, J., Huebner, A.J., Clement, K., Walsh, R.M., Savol, A., Lin, K., Gu, H., Di Stefano, B.,
Brumbaugh, J., Kim, S.-Y., et al. (2017). Prolonged Mek1/2 suppression impairs the

developmental potential of embryonic stem cells. Nature *548*, 219–223.

38. Di Stefano, B., Luo, E.-C., Haggerty, C., Aigner, S., Charlton, J., Brumbaugh, J., Ji, F.,
Rabano Jiménez, I., Clowers, K.J., Huebner, A.J., et al. (2019). The RNA Helicase DDX6
Controls Cellular Plasticity by Modulating P-Body Homeostasis. Cell Stem Cell *25*, 622–
638.e13.

39. Brena, D., Bertran, J., Porta-de-la-Riva, M., Guillén, Y., Cornes, E., Kukhtar, D., Campos-
Vicens, L., Fernández, L., Pecharroman, I., García-López, A., et al. (2020). Ancestral
function of Inhibitors-of-kappaB regulates Caenorhabditis elegans development. Sci.
Rep. *10*, 16153.

40. Hayden, M.S., and Ghosh, S. (2008). Shared principles in NF-kappaB signaling. Cell *132*,
344–362.

41. Basak, S., Kim, H., Kearns, J.D., Tergaonkar, V., O'Dea, E., Werner, S.L., Benedict, C.A.,
Ware, C.F., Ghosh, G., Verma, I.M., et al. (2007). A fourth IkappaB protein within the
NF-kappaB signaling module. Cell *128*, 369–381.

42. Thambyrajah, R., Fadlullah, Z., Proffitt, M., Neo, W.H., Guillén, Y., Casado-Pelaez, M.,
Herrero-Molinero, P., Brujas, C., Castelluccio, N., González, J., et al. (2022). IκBα
controls dormancy induction in Hematopoietic stem cell development via retinoic acid.
bioRxiv, 2022.11.17.516971. 10.1101/2022.11.17.516971.

43. Álvarez-Villanueva, D., Galán-Palma, L., Bertran, J., Floor, M., Solé, L., Lobo-Jarne, T.,
Maqueda, M., Rajbhandari, R., Marruecos, L., Villà-Freixa, J., et al. (2023). Separation-
of-function mutants reveal the NF-κB-independent involvement of IκBα in the
regulation of stem cell and oncogenic programs. bioRxiv, 2023.06.21.545928.
10.1101/2023.06.21.545928.

44. Guo, Y.-L. (2019). The underdeveloped innate immunity in embryonic stem cells: The
molecular basis and biological perspectives from early embryogenesis. Am. J. Reprod.
Immunol. *81*, e13089.

45. Pantier, R., Tatar, T., Colby, D., and Chambers, I. (2019). Endogenous epitope-tagging of
Tet1, Tet2 and Tet3 identifies TET2 as a naïve pluripotency marker. Life Sci Alliance *2*.
10.26508/lsa.201900516.

46. Sroczynska, P., Lancrin, C., Pearson, S., Kouskoff, V., and Lacaud, G. (2009). In vitro
differentiation of mouse embryonic stem cells as a model of early hematopoietic
development. Methods Mol. Biol. *538*, 317–334.

47. Chen, Q., Luo, W., Veach, R.A., Hickman, A.B., Wilson, M.H., and Dyda, F. (2020).
Structural basis of seamless excision and specific targeting by piggyBac transposase.
Nat. Commun. *11*, 3446.

48. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years
of image analysis. Nat. Methods *9*, 671–675.

49. Ewels, P., Afyounian, E., and Schuster-Boeckler, B.F.K. (2012). TrimGalore: v0. 6.7-DOI
via Zenodo.(Zenodo, 2021). doi: 10.5281/zenodo. 5127899. 2. Langmead, B. & Salzberg,
SL Fast gapped-read alignment with …. Nat. Methods.

50. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357–359.

51. Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham,
A., Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and
BCFtools. Gigascience *10*. 10.1093/gigascience/giab008.

52. Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S.,
Bernstein, B.E., Bickel, P., Brown, J.B., Cayting, P., et al. (2012). ChIP-seq guidelines and
practices of the ENCODE and modENCODE consortia. Genome Res. *22*, 1813–1831.

53. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res. *44*, W160–W165.

54. Stovner, E.B., and Sætrom, P. (2019). epic2 efficiently finds diffuse domains in ChIP-seq
data. Bioinformatics *35*, 4392–4393.

55. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq
(MACS). Genome Biol. *9*, R137.

56. Stark, R., Brown, G., and Others (2011). DiffBind: differential binding analysis of ChIP-
Seq peak data. R package version *100*.

57. Gu, Z., Eils, R., Schlesner, M., and Ishaque, N. (2018). EnrichedHeatmap: an
R/Bioconductor package for comprehensive visualization of genomic signal
associations. BMC Genomics *19*, 234.

58. Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G.,
and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

59. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–
930.

60. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

61. Zhu, A., Ibrahim, J.G., and Love, M.I. (2019). Heavy-tailed prior distributions for
sequence count data: removing the noise and preserving large differences.
Bioinformatics *35*, 2084–2092.

62. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Series B Stat. Methodol. *57*,

289–300.

63.   Garcia-Prieto, C.A., Álvarez-Errico, D., Musulen, E., Bueno-Costa, A., N. Vazquez, B.,
1027   Vaquero, A., and Esteller, M. (2022). Validation of a DNA methylation microarray for
1028   285,000 CpG sites in the mouse genome. Epigenetics *17*, 1677–1685.

1029   64.   Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P.,
1030   Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene ontology: tool for the
1031   unification of biology. The Gene Ontology Consortium. Nat. Genet. *25*, 25–29.

1032   65.   Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes.
1033   Nucleic Acids Res. *28*, 27–30.

1034   66.   Martens, M., Ammar, A., Riutta, A., Waagmeester, A., Slenter, D.N., Hanspers, K., A
1035   Miller, R., Digles, D., Lopes, E.N., Ehrhart, F., et al. (2021). WikiPathways: connecting
1036   communities. Nucleic Acids Res. *49*, D613–D621.

1037   67.   Hu, S., Metcalf, E., Mahat, D.B., Chan, L., Sohal, N., Chakraborty, M., Hamilton, M.,
1038   Singh, A., Singh, A., Lees, J.A., et al. (2022). Transcription factor antagonism regulates
1039   heterogeneity in embryonic stem cell states. Mol. Cell *82*, 4410–4427.e12.

1040   68.   Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N., and Sergushichev, A.
1041   (2021). Fast gene set enrichment analysis. bioRxiv, 060012. 10.1101/060012.

1042   69.   Zhou, W., Triche, T.J., Jr, Laird, P.W., and Shen, H. (2018). SeSAMe: reducing artifactual
1043   detection of DNA methylation by Infinium BeadChips in genomic deletions. Nucleic
1044   Acids Res. *46*, e123.

1045   70.   Hänzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis
1046   for microarray and RNA-seq data. BMC Bioinformatics *14*, 7.

Figure 1

Figure 2

A EBs 216h

IκBα-WT   IκBα-KO

DAPI/NANOG/OCT4

80 μm   80 μm

B EBs (216h)

IκBα-WT

IκBα-KO

p=0.0001

AP+ colonies (number)

IκBα-WT   IκBα-KO

C Transcriptome dynamics during mESC differentiation

○ mESCs   ■ IκBα-WT
△ 48h   ■ IκBα-KO
□ 96h

PC2: 17.5% variance

PC1: 30.7% variance

D EBs 96h

Gab1
Sox2
Esrrb
Zfp42
Dppa4   Pluripotency
Klf2
Prdm14
Klf4
Tfcp2l1

clone #1 | clone #2 | clone #3 | clone #1 | clone #2 | clone #3

IκBα WT   IκBα KO

Row Z-Score
-2 -1 0 1 2

E Endoderm score

Z-score

mESCs   EBs 48h   EBs 96h

F Mesoderm score

mESCs   EBs 48h   EBs 96h

G Ectoderm score

mESCs   EBs 48h   EBs 96h

■ IκBα-WT
■ IκBα-KO

H

mESCs (IκBα WT or KO)

5x10⁵ cells

NSG mice

1.5 months (6 weeks)

Teratoma formation

I Teratomas - 1.5 months (OCT3/4+ cells)

IκBα-WT #1 | IκBα-KO #1
IκBα-WT #2 | IκBα-KO #2
IκBα-WT #3 | IκBα-KO #3

200μm

J Teratomas (6weeks)

p=0.0025

% OCT3/4+ cells

IκBα-WT   IκBα-KO

Figure 3

A

mESCs (IκBα-WT/IκBα-KO) → 120h, Activin A + Fgf2 → EpiSCs (IκBα-WT/IκBα-KO)

Naïve pluripotency in EpiSCs (120h)

Relative RNA expression

Nanog — p=0.0004
Sox2 — p=0.0004
Rex1 — p<0.0001
Klf4 — p=0.1422
Gbx2 — p=0.0003
Tbx3 — p=0.0002
Dppa3 — p=0.0039

0h (mESCs) — 120h (EpiSCs)

IκBα-WT
IκBα-KO

B mESCs in Serum/LIF

2i/LIF signature — KO > WT / KO < WT — NES: 2.25, p-adj: 2.0116E-36

Serum/LIF signature — KO > WT / KO < WT — NES: -1.49, p-adj: 0.00011934

Enrichment score / Rank

C Relative RNA expression

Rex1 — Serum/LIF, 2i/LIF, Serum/LIF — p<0.0001
Klf2 — Serum/LIF, 2i/LIF, Serum/LIF — p=0.25
Tbx3 — Serum/LIF, 2i/LIF, Serum/LIF — p=0.33

IκBα-WT
IκBα-KO

D mESCs (Serum/LIF)

IκBα-WT / IκBα-KO — DAPI / 5mC — 10μm

E DNA Methylation profile

mESCs / EBs 96h

Density / Methylation value (β-score)

IκBα-WT
IκBα-KO

Figure 4

A
ChEA analysis
(DEGs in IκBα-KO mESCs)

A scatter/volcano plot with y-axis "-Log₁₀ (adjusted p-value)" ($-\text{Log}_{10}$ (adjusted p-value)) and x-axis "Odds Ratio". Labeled points: MBD3, MTF2, TRIM28, SUZ12.

B
mESCs IκBα-KO (Serum/LIF)

Increased H3K4me3 | Reduced H3K27me3

Left panel (H3K4me3): 295 (blue), 1162 (807 genes) (red). Inset: "Mechanisms associated with pluripotency" vs $-\text{Log}_{10}$ (adjusted p-value).

Right panel (H3K27me3): 612 (417 genes) (blue), 1070 (red). Inset bars: "Integrin-mediated cell adhesion", "Mechanisms associated with pluripotency", "IL-3 signaling pathway" vs $-\text{Log}_{10}$ (adjusted p-value).

Y-axis: Percentage of regions

C
Genes with increased H3K4me3 | **Genes with reduced H3K27me3**

Venn diagram: 766 | 41 | 376
p=0.001625

Genes with both increased H3K4me3 and reduced H3K27me3

Bars: Focal Adhesion; Mechanisms associated with pluripotency; White fat cell differentiation; Integrin-mediated Cell Adhesion

x-axis: $-\text{Log}10$ (adjusted p-value)

D
Increased H3K4me3

Tbx3, Gbx2, Zfp42, Klf2 (track plots, IκBα-KO vs IκBα-WT)

E
Reduced H3K27me3

Tbx3, Gbx2, Zfp42, Klf2 (track plots)

— IκBα-KO — IκBα-WT

G
Increased H3K27Ac

Tbx3, Tfcp2l1, Zfp42 (track plots)

— IκBα-KO — IκBα-WT

H
mESCs (Serum/LIF)

Heatmap. Rows: IκBα WT clone #1, clone #2, clone #3; IκBα KO clone #1, clone #2, clone #3. Columns: Klf2, Nanog, Pou5f1, Prdm14, Gbx2, Tbx3, Zfp42, Klf4.
Row Z-Score scale: −1 to 1.

F
Enhancer status in IκBα-KO mESCs
(Serum/LIF)

Left panel (H3K4me1): 2764 (blue), 810 (red). Poised Enhancers. Inset GO terms: regulation of membrane potential, axon guidance, limb morphogenesis, appendage morphogenesis, telencephalon development, kidney development, renal system development, cell-cell adhesion, axonogenesis, forebrain development. x-axis: $-\text{Log}_{10}$ (adjusted p-value)

Right panel (H3K4me1 H3K27Ac): 1550 (blue), 2468 (red). Active Enhancers. Inset: "Mechanisms associated with pluripotency". x-axis: $-\text{Log}_{10}$ (adjusted p-value)

■ Reduced   ■ Increased

Figure 5

A NF-kB genes (KEGG 2019 Mouse)

mESCs — p = 0.96 — IκBα-WT / IκBα-KO
EBs 48h — p = 0.82 — IκBα-WT / IκBα-KO
EBs 96h — p = 0.93 — IκBα-WT / IκBα-KO

Normalized Expression

B

IκBα^ΔH2A/H4 (IκBα^84-86A)
IκBα^ΔNF-κB (IκBα^N108A)

C Input / PD. GST / PD. GST-p50 / PD. GST / PD. GST-H2A
Dox, i-IκBα^WT, i-IκBα^ΔNF-κB, i-IκBα^ΔChrom
IκBα, p50, Comassie

D Cytoplasm protein extracts
Doxy (16h): − +
i-IκBα^WT, i-IκBα^ΔNF-kb, i-IκBα^ΔChromatin
IκBα, Tubulin

E Chromatin protein extracts
Doxy (16h): − +
i-IκBα^WT, i-IκBα^ΔNF-kb, i-IκBα^ΔChromatin
IκBα, H3

F Naïve pluripotency markers (Serum/LIF)

Zfp42 (Rex1): p=0.0018, p<0.0001, p=0.6273
Sox2: p<0.0001, p=0.0531, p=0.5490
Klf2: p=0.0028, p=0.0010, p=0.5841
Tbx3: p=0.0003, p=0.0008, p<0.0001

Relative RNA expression
i-IκBα^WT, i-IκBα^ΔNF-κB, i-IκBα^ΔChrom
■ No Doxy ■ Doxy

G Serum/LIF — 5mC
i-IκBα^WT, i-IκBα^ΔNF-kB, i-IκBα^ΔChrom

H EBs 216h
IκBα-WT 58.9%
IκBα-KO 17.8%
i-IκBα^WT 33.4%
i-IκBα^ΔNF-kB 28.2%
i-IκBα^ΔChrom 17.5%
SSEA-1

I EBs 216h
IκBα-WT, IκBα-KO
i-IκBα^WT, i-IκBα^ΔNF-kB, i-IκBα^ΔChrom

EBs 216h
AP colonies (Total number)
p=0.0038
p=0.9847
p=0.9367
p=0.0043
IκBα-WT, IκBα-KO, i-IκBα^WT, i-IκBα^ΔNF-kB, i-IκBα^ΔChrom

J EBs 216h
i-IκBα^WT, i-IκBα^ΔNF-kB, i-IκBα^ΔChrom
100 μm, 100 μm, 20 μm
DAPI/NANOG/OCT4

Figure S1 Related to Figure 2

A Nfkbia chr12:55,536,195-55,539,432

gRNA 1 px548 (GFP)
gRNA 2 px330 (mCherry)

IκBα-WT   IκBα-KO
clone #   1 2 3   1 2 3
IκBα
Tubulin

B SSEA-1 (EBs 216h)

clone #:
IκBα-WT
38.8%   1
41.3%   2
24.3%   3
IκBα-KO
86.2%   1
85.2%   2
76.3%   3

Counts

SSEA-1

C Pluripotency (EBs 216h)

Pou5f1   p=0.0002
Gbx2   p=0.0004
Klf2   p<0.0001
Sox2   p<0.0001
Rex1   p<0.0001
Nanog   p<0.0001

Relative RNA expression

IκBα-WT
IκBα-KO

D Alkaline Phosphatase (AP) staining

EBs in non-adherent dish → 216h → Dissociated EBs plated in adherent dith with Serum/LIF → 96h → Count AP+ colonies

E EBs 168h

p=0.0085

Relative size

IκBα-WT   IκBα-KO

F EBs 96h IκBα-WT vs IκBα-KO

1121 genes   1263 genes

$-\log_{10}$ (FDR)

T
Foxa2
Mixl1
Eomes
Klf2
Zfp42
Esrrb
Prdm14

FDR <0.05 |LFc| > 1

Log₂ Fold-Change
(IκBα-WT vs IκBα-KO)

Figure S2 related to figure 3

Figure S3 related to figure 4

Figure S4 related to figure 5

A Nfkbia

Fold-activation (y-axis): 0.1, 1, 10, 100, 1000

x-axis: i-IκBα^{WT}, i-IκBα$^{ΔNF-kB}$, i-IκBα^{ΔChrom}

○ No dox.
● Dox. (16h)

B Doxy-inducible IκBα$^{ΔNF-kB}$ or IκBα^{ΔChrom}

IκBα, IκBα

IκBα-KO mESCs → +Doxy (4x passages) Serum/LIF → i-IκBα$^{ΔNF-kB}$ mESCs, i-IκBα^{ΔChrom} mESCs

Pluripotent status

Differentiation potential

216h → Embryoid Bodies

C Differentiated cells
(% SSEA-1⁻ in EBs 216h)

Percentage (y-axis): 0, 10, 20, 30, 40

x-axis: i-IκBα^{WT}, i-IκBα$^{ΔNF-kB}$, i-IκBα$^{ΔH2A/H4}$

D Reconstituted EBs (216h)

%EBs (y-axis): 0, 50, 100

x-axis: IκBα-WT, IκBα-KO, IκBα^{WT}, IκBα$^{ΔNF-kB}$, IκBα$^{ΔH2A/H4}$

■ Totally Pluripotent EBs
■ Partially Differentiated EBs
■ Totally Differentiated EBs