Chromatin enhancer activity of IκBα mediates the exit from naïve pluripotency

Luis G. Palma[1,2,3,δ], Daniel Álvarez-Villanueva[1,3,4,δ], María Maqueda[1,2,3], Mercedes Barrero[5],

Arnau Iglesias[1,2,3], Joan Bertran[6], Damiana Álvarez-Errico[2], Carlos A. García-Prieto[2], Cecilia

Ballaré[5], Virginia Rodriguez-Cortez[2], Clara Bueno[2], August Vidal[4], Alberto Villanueva[7], Pablo

Menéndez[2,3,8,9,10], Gregoire Stik[2], Luciano Di Croce[5,10,11], Bernhard Payer[5,11], Manel

Esteller[2,3,10,12], Lluís Espinosa[1,3,*] & Anna Bigas[1,2,3,*]

[1]Program in Cancer Research. Hospital del Mar Research Institute, Barcelona, Spain.

[2]Josep Carreras Leukemia Research Institute, Barcelona Spain.

[3]Centro de Investigación Biomédica en Red Cancer (CIBERONC), Madrid, Spain.

[4]Institut Investigació Biomèdica de Bellvitge (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

[5]Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, Spain.

[6]Faculty of Sciences, Technology and Engineering, Universitat de Vic - Universitat Central de Catalunya, Vic 08500 Spain

[7]Chemoresistance and Predictive Factors Group, Program Against Cancer Therapeutic Resistance (ProCURE), Catalan Institute of Oncology (ICO), Oncobell Program, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

[8]Spanish Network for Advanced Therapies (RICORS-TERAV). Carlos III Health Institute (ISCIII), Spain.

[9]Department of Biomedicine. University of Barcelona, Spain.

[10]Institucio Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

[11]Universitat Pompeu Fabra, Barcelona, Spain.

[12]Physiological Sciences Department, School of Medicine and Health Sciences, University of Barcelona (UB), Barcelona, Catalonia, Spain.

[δ]These authors contributed equally

Keywords: IκBα, NFKBIA, NF-κB, DNA Methylation, Polycomb, Embryonic Stem Cells, Pluripotency Exit, Germ Layer Specification,

*Corresponding authors:

Anna Bigas and Lluis Espinosa

Cancer Research Program. CIBERONC.

Hospital del Mar Research Institute

Doctor Aiguader 88, 08003, Spain.

Tel: +34 933160440; email: abigas@researchmar.net; lespinosa@researchmar.net

36   Summary

Inflammatory signals are key in development and cell differentiation, yet their precise involvement in
the transition from naïve to primed pluripotency remains elusive. One of the main sensors of
inflammation in cells is the IκBα protein which exerts its function through activation of NF-κB. Our
previous work identified a chromatin role of IκBα that is crucial for the proper regulation of different
tissue stem cells. Here, we demonstrate that IκBα depletion causes profound epigenetic rewiring in
pluripotent stem cells characterized by altered histone post-translational modifications and DNA
methylation, which is associated with changes in transcriptional regulation. These changes led to
stabilization of cells in a ground state of naïve pluripotency when cultured in Serum/LIF conditions and
differentiation impairment. Mechanistically, IκBα binds to intergenic chromatin regions in the naïve
pluripotency state, which result in altered activation status of a subset of enhancers, including several
associated to pluripotency-related genes. Through the engineering of separation-of-function mutants,
we demonstrate that the effects of IκBα in the regulation of stem cell pluripotency are NF-κB-
independent, but mainly relies on its chromatin-related function. Altogether, our results highlight a
novel mechanism by which IκBα participates in regulating the pluripotent states of embryonic stem
cells, shedding light on the interplay between inflammatory signals and pluripotency regulation.

INTRODUCTION

Embryonic Stem Cells (ESCs) pluripotency is characterized by the capacity to generate all somatic and germline lineages both *in vitro* and *in vivo*, depending on the culture conditions. When cultured in serum plus Leukemia Inhibitory Factor (LIF) (Serum/LIF), murine ESCs (mESCs) include cells interconverting to different metastable states [1] that either resemble preimplantation blastocyst (naïve pluripotency) or post-implantation (primed pluripotency) embryonic stages [2–5]. The heterogeneity of mESCs in Serum/LIF is reduced by culturing them in the presence of GSK3β and MEK1/2 inhibitors (CHIR99021 and PD0325901, respectively) along with LIF (2i/LIF). Under these conditions, mESCs enter a ground state of naïve pluripotency characterized by self-renewal activity while suppressing any pro-differentiating signals [6,7]. This pluripotent state closely resembles the inner cell mass of the E4.0 mouse preimplantation blastocyst [8]. After implantation, pluripotent capability of the mouse embryo (E5.5) becomes restricted to Epiblast Stem Cells (EpiSCs), which represent a primed pluripotent state, characterized by coexpression of pluripotency and lineage specification markers [9,10]. EpiSCs can give rise to cells of the three different germ layers (endoderm, mesoderm and ectoderm), albeit their contribution to chimeras or to germ cells is greatly compromised [4,11]. EpiSCs can be derived *in vitro* from naïve pluripotent cells cultured in a medium supplemented with Nodal and fibroblast growth factor (FGF) [12]. Different epigenetic features distinguish naïve from primed pluripotent states [13], such as changes in the X chromosome inactivation in female cells [12,14], histone post-translational modifications [10] or a global DNA hypomethylation profile observed in naïve pluripotent stem cells [15–17]. These epigenetic changes are a prerequisite for the subsequent activation of gene circuits associated with pluripotency exit and germ layer specification [9,10]. Crucial regulatory elements for the transition from naïve to primed pluripotency state are located in distal DNA regions or enhancers to facilitate the resolution toward the primed state [18,19]. Therefore, identifying novel players to fine-tune the equilibrium between ground and primed states is crucial for preserving both pluripotency stability and differentiation capability in mESCs.

Inflammation is a defense mechanism mostly activated in response to harmful endogenous or foreign stimuli, which involves different signaling pathways including NF-κB. Canonical NF-κB signaling is triggered by the IKK kinase complex, which induce the phosphorylation and subsequent degradation of the NF-κB inhibitor, IκBα, leading to nuclear translocation of the NF-κB factors (e.g. p50/p65) [20]. We have previously identified an alternative nuclear function of IκBα involving the interaction with histone deacetylases (HDACs), core elements of the Polycomb Repressor Complex 2 (PRC2) and histones [21–23]. This nuclear IκBα function is crucial for tissue stem cell homeostasis and the proper differentiation of epidermal, intestinal and hematopoietic cells [22–25].

There is now robust evidence that inflammatory signals are crucial in stem cell development [26,27], and it was recently demonstrated that attenuation of inflammatory signals by ATG5 facilitates mESCs differentiation [28]. Unexpectedly, being NF-κB pathway a pivotal regulator of

inflammation, classical NF-κB factors including p50 and RelA/p65 are expressed at very low
levels in mESCs. Overexpression of NF-κB proteins leads to cell differentiation while
expression of the IκBα super-repressor (S32-36A mutant) increases expression of
pluripotency markers [29]. Similarly, inhibition of Protein Kinase C (PKC) activity in mESCs, which
in turn restrains NF-κB signaling, allows keeping mESCs in a ground pluripotency state, thus
facilitating the generation of induced pluripotent stem cells (iPSCs) [30]. Together these results
indicate that NF-κB inhibition is required for maintaining the pluripotent state of murine ESCs.
Given the importance of NF-κB in the multiple aspects of stem cell biology [31] and the pivotal
role of IκBα in NF-κB regulation, the possible impact of the IκBα chromatin function in cell
fate has remained challenging.

We have now investigated the role of IκBα in a context of pluripotent stem cells.
Unexpectedly, and in contrast with the pro-differentiation effects of NF-κB activation,
depletion of the NF-κB inhibitor IκBα precludes the exit of mESCs from naïve pluripotency
associated with an epigenetic rewiring. By using the recently described Separation-Of-
Function (SOF) IκBα mutants [32], we have now demonstrated that pluripotency regulation by
IκBα is mainly exerted through its chromatin-related function but independent on its capacity
to negatively regulate NF-κB.

     RESULTS

     IκBα accumulates in the chromatin of naïve pluripotent cells

Because of the prominent role of NF-κB signaling in mESCs regulation, we analyzed the
expression pattern of the different NF-κB members including the IκB inhibitors in the three
defined pluripotent stages: ground state of naïve pluripotency (2i/LIF), naïve pluripotency
(Serum/LIF), and primed pluripotency (Epiblast Stem Cells or EpiSCs) (Figure 1A). In
agreement with NF-κB being a pro-differentiation factor, canonical NF-κB inhibitors including
IκBα (*Nfkbia*), IκBβ (*Nfkbib*) and IκBε (*Nfkbie*) were all expressed in the naïve pluripotency
state (both ground and Serum/LIF), with *Nfkbia* levels significantly reduced upon naïve
pluripotency exit towards EpiSCs (Figure 1B). In contrast, *Rela, Nfkb1 and Nfkb2 genes*
codifying for the NF-κB factors p65/RelA, p105 and p100, respectively, showed an opposite
pattern and either increased or maintained their levels in EpiSCs (Figure 1C). We further
characterized the distribution of IκBα in the cytoplasm, nucleus and chromatin fractions of
the three states of pluripotency (Figure 1D). IκBα protein levels in the cytoplasm and the
nucleus correlated with the detected RNA levels in the three different cellular states being
highest in 2i/LIF condition (Figure 1B). Interestingly, a notable accumulation of IκBα was
observed in the chromatin of naïve pluripotent cells cultured in Serum/LIF conditions
compared to 2i/LIF. In contrast, IκBα was entirely absent from the chromatin of EpiSCs. (Figure
1D). The abundance of IκBα RNA and protein in the context of naïve pluripotency compared
to EpiSCs was further corroborated through analysis of an additional database encompassing
both naïve and primed pluripotent states [33] (Figure 1E).

Next, we studied IκBα expression in differentiated embryoid bodies (EBs) from mESCs
(Serum/LIF) (see materials and methods) and analyzed the expression levels of different
pluripotency (*Dppa4, Pou5f1, Sox2, Prdm14, Klf4, Zfp42 [Rex1], Nanog*), differentiation-linked
(*Mesp1, Gata6, Lhx1, Gata4, Wnt3a, Fgf5, Nes, Axin2, Wnt3, Eomes, T, Foxa2, Mixl1, Lef1*),
and NF-κB genes (*Nfkbid, Nfkbia, Nfkbie, Nfkbib, Nfkbiz, Rel, Rela, Relb*) by RNA sequencing.
Nfkbia (IκBα gene) was expressed at higher levels specifically in mESCs (Serum/LIF) whereas
cells undergoing differentiation (48h and 96h) displayed a reduction in its expression levels
(Figure 1F). The expression pattern of *Nfkbia* was similar to the pluripotency genes, whereas
most of the NF-κB genes were upregulated upon differentiation, clustering together with
developmental genes, as previously described [29].

Taken together, these data show that the expression of *Nfkbia*, like the other *Nfkbi*, is high in
the naive pluripotent state, in contrast to the NF-κB genes, which cluster together with
differentiation markers. In addition, IκBα accumulates in the chromatin of naive pluripotent
cells, more so in cells cultured under serum/LIF.

Lack of IκBα impairs the exit from the naïve pluripotency state

To investigate the functional implication of the chromatin accumulation of IκBα in naïve
pluripotency, we knocked-out IκBα protein in mESCs by CRISPR-Cas9 (IκBα-KO mESCs) (Figure
1G and materials and methods). We did not detect major changes in NF-κB target genes upon
IκBα depletion neither in mESCs, nor 48h and 96h EBs (Figure 1H), suggesting that NF-κB
regulation did not require IκBα at this stage or it might be compensated by the other canonical
or non-canonical IκBs (IκBβ, IκBε or p100), as previously demonstrated [34,35]. Paradoxically,
IκBα-KO mESCs have an impaired capability to undergo differentiation, since IκBα-KO EBs
cannot properly evolve towards differentiation trajectories (Figure 2A). This observation was
confirmed by analysis of naïve pluripotency markers at 96h (*Gab1, Sox2, Esrrb, Zfp42, Dppa4,*
Klf2, Prdm14, Klf4, Tfcp2l1), which displayed higher expression levels in IκBα-KO compared
to their wild-type (IκBα-WT) counterparts (Figures 2B and S1A). Differentiation of IκBα-KO EBs
was deficient in programs associated with all germ layer specification (endoderm, mesoderm,
and ectoderm) (Figures 2C-E). Moreover, IκBα-KO EBs were smaller in size (Figure S1B), and
maintained higher expression of the pluripotency program even at 216h (Figures 2F and S1C).
We also detected a higher number of Alkaline Phosphatase-expressing cells (Figures 2G and
S1D) and higher expression of the pluripotency marker SSEA-1 (Figure S1E) at 216h IκBα-KO
EBs. To further study the requirement of IκBα in pluripotency exit *in vivo*, we established
teratomas by subcutaneous injection of IκBα-WT and IκBα-KO mESCs (Figure S1F and
materials and methods) in NSG (NOD.Cg-Prkdc^scid Il2rg^tm1Wjl/SzJ) mice. IκBα-KO teratomas
showed a decreased differentiation potential, as indicated by higher number of OCT4[+] cells
(Figures S1G-H). These observations underscore the essential role of IκBα in the exit from the
pluripotency state in mouse embryonic stem cells (mESCs).

To determine the specific time-point in which IκBα is required for pluripotency exit, we
differentiated IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF (naïve pluripotency)
towards EpiSCs (primed pluripotency) for 120h (Figure 2H and materials and methods). IκBα-
KO cells were not committed towards the primed pluripotency stage and maintained an
elevated expression levels of naïve pluripotency genes (*Dppa3, Nanog, Sox2, Rex1, Klf2, Klf4,*
Gbx2, Tbx3) (Figure 2I). Additionally, IκBα-KO cells retained a naïve pluripotent morphology,
forming tight clusters and resembling cells cultured in 2i/LIF medium (Figure S1I). These data
demonstrate that IκBα is required for the exit of the naïve state of pluripotency and its
deficiency results in impaired differentiation of mESC *in vivo* and *in vitro*, without imposing
major changes in the regulation of canonical NF-κB.

IκBα-KO mESCs resemble the ground-state of naïve pluripotency under Serum/LIF
conditions

As mentioned, mESCs cultured in serum/LIF are highly heterogeneous, containing a mixture
of cell states that resemble developmental transitions from preimplantation (naïve
pluripotency) to postimplantation (primed pluripotency) embryo [2,3]. To investigate how IκBα
deficiency was affecting the exit of mESCs from naïve pluripotency, we performed a Gene Set
Enrichment Analysis (GSEA) using RNA-seq data from IκBα-KO and IκBα-WT mESCs cultured

in Serum/LIF against gene signatures for ground (2i/LIF) and naïve (Serum/LIF) pluripotency
[36]. GSEA results revealed a significant enrichment in Serum/LIF IκBα-KO mESCs transcriptome
for genes specifically expressed in the ground state (2i/LIF), while they negatively correlate
with mESCs cultured in Serum/LIF (Figure 3A) [36]. Specifically, IκBα-KO mESCs expressed higher
levels of the ground state pluripotency genes *Klf2, Nanog, Pou5f1, Prdm14, Gbx2, Tbx3,*
Zfp42[Rex1] and *Klf4* (Figure 3B), similar to IκBα-WT mESCs cultured in 2i/LIF (Figure S2A).
Since ground (2i/LIF) and naïve (Serum/LIF) states are characterized by important differences
in epigenetic marks, we carried out an epigenetic profiling of IκBα-WT and IκBα-KO mESCs
cultured in Serum/LIF. We performed chromatin immunoprecipitation followed by
sequencing (ChIP-seq) of histone modifications associated to gene activation (H3K4me3),
gene repression (H3K27me3) and constitutive heterochromatin (H3K9me3) (Figures 3C-D and
S2B-D). Overall, Serum/LIF IκBα-KO mESCs exhibited a more permissive chromatin status as
evidenced by increased H3K4me3 and decreased H3K9me3 levels, while H3K27me3 was
redistributed in both directions (Figure 3C). Notably, the gain in H3K4me3 mark was
associated with a reduction of H3K27me3 levels in genes associated with naïve pluripotency
(Figures 3C-D and S2B-C), in agreement with the higher pluripotent state observed in IκBα-
KO mESCs. As mentioned, detection of the H3K9me3 mark was significantly reduced in IκBα-
KO mESCs (Figures 3C and S2D). These regions correspond to genes involved in DNA
methylation and chromatin edition functions (Figure 3C). By analyzing the DNA methylation
status under Serum/LIF conditions by immunofluorescence of 5-methyl-cytosine (5mC) we
found a global DNA hypomethylation in IκBα-KO mESCs (Figure 3E), which is one of the key
features of the ground state of naïve pluripotency (2i/LIF) [15,37]. To further characterize the
DNA methylation status in IκBα-KO cells, we performed DNA methylation arrays [38] in IκBα-
KO and IκBα-WT mESCs (Serum/LIF) and 96h after EB differentiation. We observed a strong
reduction of the DNA methylation level in IκBα-KO mESCs in a genome-wide fashion (Figure
3F). Specifically, IκBα-KO mESCs contain lower DNA methylation levels in promoters, gene
bodies, enhancers and imprinted regions (Figure 3G), and this DNA hypomethylation status
was not associated with any particular chromosome (Figure S2E). DNA hypomethylation was
maintained upon 96h in IκBα-KO EBs (Figures 3F-G and S2F).

These results further indicate that IκBα is required for the transition from ground to primed
pluripotent states and for the proper regulation of the histone post-translational
modifications (PTMs) and DNA methylation changes associated with this process. We propose
that the enhanced pluripotent state found in IκBα-KO mESCs under basal conditions
(Serum/LIF) may negatively impact on their differentiation potential, similar to that described
for other models with stabilized naïve pluripotent state [39,40].

IκBα binds to intergenic regions to attenuate the activation status of chromatin linked to
naïve pluripotency genes

To gain insight into the mechanism by which IκBα regulates the transition from naïve to
primed pluripotent states, we knocked-in a 3xFLAG-tag into the IκBα locus and used these

transgenic cells for ChIP-seq assay with the anti-FLAG antibody (see Materials and Methods).
We identified 817 bona-fide IκBα-bound chromatin regions which were consistent between
three independent replicates (Figure S3A). Among all the genome regions, IκBα was mainly
bound to distal intergenic and intronic regions (38.5% and 42.2% of total IκBα peaks,
respectively) (Figure 4A). Interestingly, IκBα-bound regions coincide with those that gain
enhancer activation-related H3K27Ac signals in EpiSCs (Figures S3B-C). The fact that IκBα was
released from chromatin at the stage of primed pluripotency (Figure 1D) suggested that IκBα
might be involved in fine-tuning these regions at the exit from naive pluripotency. To address
this question, we quantified the total amount of H3K27Ac (active enhancer) by western blot
analysis of chromatin extracts of IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF. We
observed an increase of this activation mark in IκBα-KO mESCs, whereas we did not observe
global changes in H3K4me1 levels, a marker of poised enhancers (Figures S3D-E). These
results suggested that IκBα depletion in naïve pluripotency might lead to an increase in the
global levels of active enhancers, which are characterized by a gain of H3K27Ac without
changes in H3K4me1 levels [41]. The increase in H3K27Ac was further validated by ChIP-seq
experiments conducted in both IκBα-WT and IκBα-KO mESCs. Notably, depletion of IκBα
protein under Serum/LIF conditions led to an increase in the deposition of H3K27Ac at regions
where we detected IκBα binding (Figure 4B), while levels of H3K4me1 were not generally
altered. Since IκBα deficiency was affecting enhancer-related histone marks, we further
profiled the enhancer status of IκBα-KO mESCs by analyzing the differential distribution of
poised enhancers (showing changes in H3K4me1 levels without affecting H3K27Ac) and active
enhancers (chromatin regions decorated with H3K4me1 that also display H3K27ac binding)
(Figures 4D). Overall, IκBα-KO mESCs showed a reduction in poised enhancers at genes (or
related genomic regions) associated to differentiation processes (Figures 4D and S3F),
whereas active enhancers of genes associated to pluripotency were significantly enriched
(Figures 4D-E and S3G). These results further support that mESCs lacking IκBα are
epigenetically reinforced in the ground state of naïve pluripotency. Since the majority of IκBα
chromatin peaks were located in distal intergenic and intronic regions (Figure 4A), and we
found that IκBα deficiency led to changes in the enhancer status (Figure 4D), we took
advantage of publicly available Hi-C data to identify topologically associating domains (TADs)
at putative IκBα target genes in Serum/LIF mESCs [42]. It was previously demonstrated that TADs
are organized in distinct chromatin compartments defined as active (Compartment A) or
inactive (Compartment B) [43]. Importantly, chromatin regions bound by IκBα mainly
correspond to TADs of the active compartment (81%), whereas a small proportion of IκBα
peaks were found in inactive chromatin (19%) (Figure S3H). Furthermore, TADs harboring IκBα
binding exhibited a greater proportion of enhancers differentially regulated in IκBα-KO cells
when compared to all TADs (92% vs. 74%, respectively) (Figure S3H). Additionally, genes
located within TADs bound by IκBα are predominantly associated with mechanisms linked to
pluripotency identity (data not shown). To address the functional relevance of this
observation, we investigated the expression pattern of pluripotency-related genes in IκBα-
WT and IκBα-KO mESCs at naïve pluripotency (cultured in Serum/LIF) and differentiation (48h

EBs) stages (Figure 4F). Most of the genes analyzed were significantly upregulated in IκBα-KO
mESCs, and this difference was further accentuated in IκBα-KO 48h EBs. These results suggest
that IκBα modulates the dinamyc expression of pluripotency genes through a distal regulatory
mechanism involving enhancer activity and 3D chromatin architecture. To address this
hypothesis, we crossed Hi-C data with ChIP-seq of IκBα and enhancer-related histone marks
(H3K27Ac and H3K4me1), together with RNA-seq data. In this meta-analysis, we identified a
TAD region encompassing the Tbx3 locus with IκBα binding sites (Figure 4G). Interestingly, we
observed that the IκBα-bound regions interact with the region containing Tbx3 gene (Figure
4G; interactions highlighted with an arrow in the contact map). Moreover, that region harbors
an active enhancer marked by H3K27Ac/K4me1, whose activation status is heightened in
IκBα-KO mESCs, likely participating in increased expression of the Tbx3 gene (Figure 4G,
bottom panel) and other interactions are detected between the differentially activated
enhancer and the Tbx3 locus in this TAD (Figure 4G, interaction highlighted with a gray arrow
in the contact map).

We then determined whether differential enhancer activation status observed in IκBα-KO
mESCs (evidenced by increased H3K27Ac) was reversible. We engineered a doxycycline-
inducible IκBα construct and introduced it into IκBα-KO mESCs. Treatment of these cells with
doxycycline for several passages under Serum/LIF conditions (i-IκBα^{WT} mESCs, Figure 4H) led
to sustained IκBα expression and a significant reduction of global levels of H3K27Ac (Figures
4I and S3I). Ectopic expression of i-IκBα^{WT} in mESCs was also sufficient to attenuate the
expression of the naïve pluripotency genes *Zfp42, Klf2, Sox2* and *Tbx3 in* IκBα-KO mESCs under
Serum/LIF conditions (Figure S3L).

IκBα attenuates the activation of intergenic regions in an NF-κB independent manner

Uncovering the actual impact of chromatin-related IκBα function is challenging due to the
predominant role of this protein on NF-κB regulation. Recent work from our group has
identified the residues in the IκBα protein critical for NF-κB-binding or chromatin-binding
through H2A/H4. Mutation of these residues, allows to specifically discriminate between
either of the two functions of IκBα [32]. We generated doxycycline-inducible constructs to
express separation-of-function (SOF) IκBα mutants: i) exclusively interacts with NF-κB and is
deficient for chromatin binding (i-IκBα^{ΔChromatin}) or ii) interacts with histones but is deficient
for NF-κB interaction and inhibition (i-IκBα$^{ΔNF-κB}$) (Figure 5A). We introduced one of the SOF
IκBα mutants into IκBα-KO mESCs and confirmed their functionality by pull-down
experiments. We found that i-IκBα$^{ΔNF-κB}$ was able to interact with histone H2A but not with
the NF-κB member p50 and, i-IκBα^{ΔChromatin} could interact with p50 but not with H2A, while i-
IκBα^{WT} preserved all interactions (Figures 5B-C). Importantly, when we induced the three
different IκBα forms (i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and i-IκBα^{ΔChromatin}, Figure S4A), only i-IκBα^{WT} and i-
IκBα$^{ΔNF-κB}$ were found in the chromatin fraction of mESCs, whereas IκBα^{ΔCromatin} was retained
in the cytoplasm (Figures 5D and S4B). To address whether IκBα was participating in the
stabilization of the ground-state of naïve pluripotency in an NF-κB-independent manner, we

studied which of the SOF mutants (IκBα$^{\Delta NF-\kappa B}$ or IκBα$^{\Delta Chromatin}$) was able to reverse the
phenotype of IκBα-KO mESCs (Figure S4C). We observed the reduction of the global levels of
H3K27Ac upon induction of i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$, but not of IκBα$^{\Delta Chromatin}$ (Figures 5E and
S4D). Similarly, induction of IκBα$^{\Delta NF-\kappa B}$ was enough to reduce the H3K27Ac amount in
enhancers with increased H3K27Ac in IκBα-KO mESCs, but not when IκBα$^{\Delta Chromatin}$ was
expressed (Figure 5F). Interestingly, H3K4me1 levels followed a similar pattern as H3K27Ac in
those specific loci, and a significant reduction was achieved upon IκBα$^{\Delta NF-\kappa B}$ overexpression,
whereas IκBα$^{\Delta Chromatin}$ was mediating an opposite effect (Figure 5G). In addition, the ground
state-associated DNA hypomethylation was reverted in i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ mESCs but
not in i-IκBα$^{\Delta Chromatin}$ mESCs (Figure 5H). Furthermore, the naïve pluripotent gene expression
(*Zfp42, Klf2, Sox2, Tbx3*) was reduced upon inducing i-IκBα$^{\Delta NF-\kappa B}$, while induction of i-
IκBα$^{\Delta Chromatin}$ resulted in their increase (Figure S4E).

To address whether reverting to the basal epigenetic status is enough to rescue the
differentiation capability, i-IκBα^{WT}, i-IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta Chromatin}$ mESCs were differentiated
into EBs (Figure S4C). Induction of IκBα^{WT} and IκBα$^{\Delta NF-\kappa B}$ was enough to restore the
differentiation potential of IκBα-KO mESCs, since the number of differentiated cells (based on
percentage of SSEA-1^{neg} cells) increased (Figures 5I and S4F). Moreover, we found a reduced
number of AP+ colonies from 216h IκBα-KO EBs after induction of i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$,
similar to the number of colonies obtained from IκBα-WT EBs (Figures 5J and S4G). Lastly,
the number of remaining pluripotent-like cells (defined by OCT3/4 NANOG staining) was also
reduced in i-IκBα^{WT} and i-IκBα$^{\Delta NF-\kappa B}$ 216h EBs, and only small and few regions of
undifferentiated cells was observed (Figures 5K and S4H). On the other hand, induction of i-
IκBα$^{\Delta Chromatin}$ did not restore the differentiation potential of IκBα-KO mESCs, and phenocopied
the differentiation defects observed in IκBα-depleted mESCs.

These results demonstrate that the chromatin-dependent IκBα function, but not its NF-κB-
related activity, is necessary and sufficient to revert the stabilization in the ground state of
naïve pluripotency correlating with higher DNA methylation and lower H3K27Ac levels. Our
results demonstrate that the chromatin function of IκBα alters the activation status of
enhancers, which impacts the expression of genes involved in the maintenance of the naïve
pluripotency. This function is independent of the NF-κB-inhibitory activity of IκBα.

DISCUSSION

Here we have identified a new and unexpected function for the IκBα protein in the regulation of the naïve pluripotency exit. Using mESCs to investigate the role of IκBα in pluripotency maintenance and differentiation, we discovered that the absence of IκBα severely impairs the exit of pluripotency. Furthermore, IκBα-KO mESCs stabilizes the ground state of naïve pluripotency under Serum/LIF conditions, shedding light on its function as a critical regulator of the transition from naïve to primed pluripotency. While IκBα is commonly known for its role as an inhibitor of the NF-κB signaling pathway [20], our laboratory has previously furnished compelling evidence of its alternative role through its interaction with histones and other chromatin components [21–23]. We have demonstrated that chromatin-bound IκBα plays a crucial role in regulating skin and intestinal stem cells [22,24], hematopoietic stem cell development [25], and it influences the regenerative capacity of these tissues, and in some cases, their susceptibility to neoplastic transformation. Disentangling the chromatin-related function of IκBα from its canonical role as an inhibitor of NF-κB has been challenging, making it difficult to determine its truly biological significance. To tackle this challenge, we engineered separation-of-function (SOF) mutants of IκBα, as detailed in [44], by identifying specific residues that are essential for binding either NF-κB elements or histones. Interestingly, we have shown that only the histone-binding proficient IκBα mutant (IκBα$^{\Delta NF-\kappa B}$), which lacks the ability to bind to NF-κB, has the capability to reverse the stabilization of the ground state observed in IκBα-KO mESCs. This is sufficient to facilitate their exit from the state of naïve pluripotency, restoring their full differentiation potential (Figure 5). The involvement of NF-κB in pluripotency has been a subject of prior investigation. Studies have revealed that pluripotent cells display a dampened inflammatory signaling pathway, which is induced upon pluripotency dissolution and activation of differentiation [29,45]. However, the precise molecular mechanisms underlying the attenuated status of NF-κB during this stage remain incompletely understood. A recent work has shed light on a potential regulatory axis involving ATG5-β-TrCP1-NF-κB, proposing it as a mechanism that diminishes NF-κB activity by stabilizing the IκBα protein in mESCs [28]. In any case, since modification of canonical NF-κB activity directly impacts on the levels IκBα expression, the previous NF-κB-associated phenotypes should be reevaluated considering this chromatin role of IκBα. Furthermore, the expression of IκBα$^{\Delta Chromatin}$, which retains the inhibitory function of NF-κB, failed to mitigate the ground state stabilization observed in IκBα-KO mESCs (Figure 5). This finding suggests that the observed role of IκBα in regulating the exit from naïve pluripotency is not due to NF-κB inhibition. In fact, there is evidence that the chromatin function of IκBα is an ancestral function preceding its role as NF-κB inhibitor. In Caenorhabditis Elegans (*C.elegans*), which lacks NF-κB orthologs but possesses two homologs of IκBα, the absence of IκBα orthologs has severe multi-organ differentiation defects [46]. This observation further supports the notion that the chromatin-related function of IκBα plays a fundamental role in developmental processes, independent of its canonical role in NF-κB signaling.

We have observed that IκBα-KO mESCs exhibit epigenetic and transcriptomic profiles
resembling the ground state of naïve pluripotency under Serum/LIF culture (Figure 3). In this
regard, the fact that IκBα is preferentially located in the chromatin of mESCs cultured in
Serum/LIF, when they exhibit a high degree of heterogeneity, suggests that IκBα might act as
a regulator of this heterogeneity. One potential explanation might be that IκBα could be
affecting the stability of fluctuating transcription factors, which is crucial to mediate the
pluripotent-to-differentiation balance [3]. Thus, the lack of IκBα might stabilize those
transcription factors that would favor the ground-state observed in IκBα-KO mESCs cultured
in Serum/LIF. Further experiments are required to demonstrate this hypothesis. Moreover,
the differences in murine and human pluripotency [11] point out the necessity to address the
real impact of IκBα modulation in the stabilization of the ground state in human pluripotent
stem cells. Our investigation has also revealed that IκBα accumulates in the intergenic
chromatin regions of naïve pluripotent cells (Figures 1D and 4A). Mechanistically, this binding
of IκBα serves to safeguard enhancers associated with pluripotency from excessive activation.
This is evidenced by the higher abundance of active enhancers observed in IκBα-depleted
mESCs, particularly in regions linked to pluripotency located within TADs where IκBα is bound.
A recent study highlights the significance of maintaining a proper 3D chromatin architecture
during the resolution of the naïve pluripotent state [47]. Our findings suggest that IκBα may
potentially influence the 3D conformation of regions relevant to pluripotency, thereby fine-
tuning the expression of genes crucial for sustaining naïve pluripotency (Figure 6).
Nevertheless, the precise mechanisms by which the absence of IκBα impacts 3D chromatin
architecture remain to be elucidated.

In summary, our work establishes that IκBα mediates the exit from naïve pluripotency by
modulating the activation of enhancers associated with pluripotency genes. Importantly, this
newly identified role of IκBα operates independently of its canonical function in inhibiting the
NF-κB pathway. Our findings underscore the intricate interplay between inflammation and
pluripotency status, revealing a previously unrecognized complexity in their interaction.

METHODS

Cell culture

mESC line ES-E14TG2a (ATCC; Cat #CRL-1821) was cultured on plastic dishes precoated with 0.1% (w/v) Gelatin (Sigma; Cat #G2500-100G) in Serum/LIF medium. Serum/LIF is composed by DMEM basal medium (Sigma-Aldrich; Cat #D5796) supplemented with 15% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 1X NEAA (Gibco; Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070), 1000U/mL Leukemia Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-mercaptoethanol (Gibco; Cat#31350010). Medium was changed every day, and cells were splitted every other day using TrypLE Express (Gibco; Cat #12605010) for harvesting. Cells were maintained in a 5% CO_2 incubator at 37°C.

For inducing the ground-state of naïve pluripotency, mESCs were cultured in 2i/LIF medium for 2 consecutive passages. 2i/LIF medium is composed by NDiff 227 medium (TAKARA, Cat #Y40002) supplemented with 1000U/mL LIF, 0.4 µM PD032591 (Selleck Chemicals, Cat #S1036) and 3 µM CHIR99021 (Merck, Cat #SML1046).

Epiblast stem cell differentiation

Differentiation from mESCs towards Epiblast Stem Cells (EpiSCs) was performed as previously described [48]. 3×10^4 mESCs were seeded in 6-well plastic dishes, and they were cultured in Serum/LIF medium for 24h. Medium was then switched to NDiff 227 medium (TAKARA, Cat #Y40002) supplemented with 20 ng/ml activin A (Cat. #120-14E; PeproTech) and 10 ng/ml Fgf basic (R&D Systems, Cat #233-FB-025/CF). Cells were submitted to daily media changes till day 5 (120h), when they were further analyzed.

Embryoid bodies differentiation from mESCs

Embryoid bodies differentiation was established as described in [49]. Briefly, mESCs were splitted twice in Serum/LIF medium before inducing differentiation. Once mESCs are 80% confluent, cells were collected using TrypLE Express. One well of 6-well plate was splitted in an entire 6-well plate in IMDM-ES medium for 48h. IMDM-ES medium is composed of Iscove's Modified Dulbecco's Medium (IMDM) (Cytiva, Cat #16SH30259.01) supplemented with 20% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 1X NEAA (Gibco; Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070), 10ng/mL Leukemia Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-mercaptoethanol (Gibco; Cat#31350010). For embryoid body induction, mESCs were harvested, and they were rinsed twice with DPBS (Gibco; Cat #14190144). Cells were very well disaggregated into single-cells, and 1.2×10^4 cells/mL embryoid body differentiation (EB^{diff}) medium were resuspended to a total volume of 25mL of EB^{diff} medium. EB^{diff} medium is composed by IMDM supplemented with 15% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 50

µg/mL ascorbic acid (Sigma; Cat #A-4544), 180 µg/mL Transferrin (Roche; Cat #10652202001)
and 0.45 mM alpha-monothioglycerol (MTG) (Sigma; Cat #M6145). EBs were formed in
suspension for 5 days (120h). At day 5, in order to elongate the differentiation up to 216h
(day 9), EB^{diff} medium was refreshed by harvesting EBs with 10mL serological pipette,
centrifuge at 200 g for 3 min and EBs were resuspended in fresh EB^{diff} medium for 4 more
days.

CRISPR/Cas9 gene editing and cell lines generation in mESCs

The two guide RNAs (gRNAs) targeting the Nfkbia locus were designed using the CRISPR design
tool from MIT (http://crispr.mit.edu). The best 2 gRNAs (based on on-target and off-target
scores) that were targeting the exon 1 of Nfkbia (IκBα gene) were selected. After annealing,
one of the gRNAs was cloned into SpCas9(BB)-2A-GFP (px458) plasmid (Addgene; Cat #48138),
and the other gRNA was cloned into the px330-mCherry plasmid (modified from Addgene;
Cat #98750 to incorporate mCherry reporter). mESCs were co-transfected with the two
plasmids (px458-gRNA1 and px330-mCherry-gRNA2). 3x10^5 cells were seeded a day before
per well of 6-well plate. At day of transfection, cells were washed once with 1X DPBS (Gibco;
Cat #14190144), and 2mL Opti-MEM medium (Gibco; Cat #31985070) was added into cells.
1.25 µg of each plasmid (2.5 µg of total DNA) was incubated with 10 µL of Lipofectamine 2000
(Invitrogen;   Cat   #11668019)   in   240   µL   of   Opti-MEM   for   20   min.
OptiMEM::Lipofectamine::DNA mixture was added into mESCs, and cells were incubated for
5 hours. Cells were then washed with DPBS, and medium was replaced by Serum/LIF. 48h
after transfection, GFP$^+$ mCherry$^+$ single cells were sorted by FACS using BD FACSAria II Cell
Sorter (BD Bioscience). Mutant clones were screened through PCR and western blotting to
identify single clones with no IκBα protein expression. Non-targeting scrambled gRNAs were
cloned into px458 and px330-mCherry plasmids as wildtype clones. Three independent clones
from each genotype (IκBα-WT and IκBα-KO) were selected for further experiments.

To increase IκBα specific detection, the IκBα endogenous gene (Nfkbia) was knocked-in with
3XFLAG sequence (DYKDHDGDYKDHDIDYKDDDDK) at the 3' end. To generate IκBα-3XFLAG
mESCs, 3XFLAG sequence was cloned into a plasmid containing Nfkbia left and right homology
arms flanking the endogenous stop codon; a spacer sequence of three glycines (GGCGGTGGG)
was also introduced to link the IκBα endogenous protein with the 3XFLAG (donor plasmid).
To induce the knock-in of the 3XFLAG, a gRNA complementary to the region surrounding the
endogenous stop codon was cloned into the PX458 plasmid (PX458-gRNA-mNfkbia-Cterm).
mESCs were co-transfected with donor and PX458-gRNA-mNfkbia-Cterm plasmids, cells were
sorted by FACS and individual clones were screened for IκBα-3XFLAG homozygous clones.
gRNAs and oligos sequences are included in supplementary table 1.

Doxycycline-inducible IκBα (IκBα^{WT}, IκBα$^{ΔNF-κB}$ and i-IκBα$^{ΔH2A/H4}$) mESCs were generated by
cloning the three different IκBα versions into a PiggyBac transposon system [50]. IκBα protein-
coding cDNAs were PCR-amplified, and NheI and SalI restriction sites were placed at the 5'

and 3' ends, respectively. T2A-EGFP fragment was amplified from PX458 plasmid, and SalI and AgeI restriction sites were introduced at its 5' and 3' ends. PB-TRE backbone (Addgene, Cat# 63800) was digested with NheI and AgeI enzymes, and mNfkbia and T2A-EGFP fragments were ligated into the PB-TRE digested vector. To generate mESCs containing the inducible vector, IκBα-KO mESCs were co-transfected with PB-TRE-mNfkbia-T2A-EGFP and PiggyBac transposase plasmids, and pool of transfected cells was selected by Hygromycin (100μg/mL) for 7 days. Cells were doubly screened by FACS sorting green fluorescent protein after 24h of doxycycline (1μg/mL) treatment. Pool of cells were used for further experiments. Oligos sequences are included in supplementary table 1.

Teratoma formation assay

IκBα-WT and IκBα-KO mESCs grown in Serum/LIF (5×10^5) were injected intramuscularly into severe combined immunodeficient mice (NSG). In order to favor 3D aggregation of cells and teratoma formation, cells were resuspended in Matrigel Matrix (BD; Cat #356234) prior to injection. Six weeks later, mice with tumors were euthanized, and tumors were fixed in formaldehyde, embedded in paraffin, sectioned and stained with hematoxylin and eosin for histological analysis.

Immunofluorescence staining

For immunostaining of mESCs, cells were seeded on 0.1% gelatin-coated coverslips on 6-well plates; for EB immunofluorescence, they were collected directly from the plate. Samples were washed twice with 1X PBS, and they were fixed at 4ºC for 30 minutes with 4% PFA (Electron Microscopy Sciences; Cat #15713S). For immunofluorescence of 5-Methylcytosine (5mC), fixed cells were incubated with 2M Hydrochloric Acid (Sigma-Aldrich, Cat #H1758) for 20 minutes at room temperature. Samples were then washed twice with 1X Tris-buffered saline (TBS; 50 mM Tris-Cl, pH 8) and permeabilized and blocked with 1X TBS supplemented with 1% Triton-X100 (MERCK; Cat #9036-19-5) and 6% FBS (Biological Industries; Cat #04-001-1A) for 2h at 4ºC. Samples were washed twice with 1X TBS supplemented with 6% FBS. Primary antibody incubation was performed in 1X TBS plus 6% FBS and 0.3% or 0.5% Triton-X100 (0.3% for 2D culture and 0.5% for EBs) overnight at 4ºC. The following primary antibodies were used: OCT3/4 (1:250; Santa Cruz; Cat #sc-5279), NANOG (1:250; Novus Biologicals; Cat #NB100-588), 5-Methylcytosine (1:500, Invitrogen, Cat #MA5-24694), H3K27Ac (1:2000, Abcam, Cat #AB4729). Samples were then washed 4 times with 1X TBS, 5 min each washing. Secondary antibody incubation was performed in TBS 1X plus 1% BSA (Sigma-Aldrich; Cat #9048-46-8) for 2h at room temperature. The following antibodies were used: Alexa Fluor 488 donkey anti-mouse antibody (1:1000; Invitrogen; Cat #A-21202), Alexa Fluor 647 donkey anti-mouse antibody (1:1000; Invitrogen; Cat #A-31571), Alexa Fluor 594 donkey anti-rabbit antibody (1:1000; Life Technologies; Cat #A-21207), Alexa Fluor 647 donkey anti-rabbit antibody (1:1000, Invitrogen; Cat #A-31573). Samples were washed three times with 1X TBS for 5 min

at room temperature each rinse. Samples were mounted using DAPI Fluoromount-G
(Southern Biotech; Cat #0100-20).

Microscopy and image acquisition

Fluorescence images were acquired using Confocal Leica TCS SP5 (Leica Microsystems), and
Leica application software LAS AF (Leica Microsystems) was used to visualize the images.
Images of teratoma haematoxylin and eosin and immunohistochemistry stainings were
acquired using BX61 Olympus Microscope (Olympus), and PRECiV™ 2D Image and
Measurement Software (Olympus) was used to visualize the images. The Image J (version
1.15) was used for further analysis [51].

Flow cytometry sample preparation and analysis

Cells were collected and they were purified based on FACS using BD FACSAria II Cell Sorter
(BD Bioscience). Cells were sorted at 3,500 events/second, and at maximum flow rate of 4
and 85 µm nozzle. 5 µg/ml of DAPI (Biotium; Cat #BT-40043) was used as a viability dye. For
flow cytometry analysis of pluripotency exit, cells were disaggregated using TrypLE Express,
and they were incubated at 4ºC for 20 minutes with SSEA1-eFluor 660 Monoclonal Antibody
(1:200, Invitrogene, Cat #50-8813-42). BD LSRFortessa Cell Analyzer or BD LSR II Flow
Cytometer (BD Bioscience) were used for Flow Cytometry Analysis experiments. Flow
cytometry data were analyzed using FlowJo X v10.0.7 (BD Biosciences).

RNA isolation, cDNA synthesis and quantitative RT-PCR

Total RNA isolation from cells was performed using the RNeasy Plus Mini Kit (Qiagen; Cat
#74136) or RNeasy Micro Kit (Qiagen; Cat #74004) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
2 µg of total RNA was retro-transcribed using Transcriptor First Strand cDNA Synthesis Kit
(Roche; Cat #04897030001) following the manufacturer's instructions.

Quantitative RT-PCR was performed in triplicates for each sample, and SYBR Green I Master
Kit (Roche; Cat #04887352001) was used to carry out the reaction. qRT-PCR was performed
using the LightCycler 480 system (Roche). Relative expression levels were calculated as $2^{-\Delta CT}$
normalized with the average CT of the housekeeping gene Tbp or Gapdh. Oligos sequences
are found in supplementary table 1.

Chromatin Immunoprecipitation (ChIP)

4-6x10^7 mESCs were cross-linked by incubating them in DPBS (Gibco, Cat #14190094)
supplemented with 1% formaldehyde (Sigma-Aldrich; Cat #252549) for 10 min rocking at
room temperature. Cross-linking was stopped using 125 mM Glycine (Sigma-Aldrich, Cat
#G8790) rocking for 5 minutes at room temperature. Fixed cells were then washed twice with
ice-cold DPBS, and scrapped and collected using DPBS supplemented with

protease/phosphatase inhibitor cocktail composed by 1X cOmplete™, EDTA-free Protease
Inhibitor Cocktail (Roche, Cat #11873580001), 1 mM PMSF, 1mM Sodium Ortovanadate and
20 mM β-Glycerol phosphate. Cells were centrifuged at 3200g for 5 minutes at 4ºC. Cells were
lysed by resuspending cell pellet in ice-cold ChIP buffer (1 volume of SDS buffer [100mM NaCl,
50mM Tris-HCl pH 8.1, 5mM EDTA pH 8, 0.5% SDS] and 0.5 volume of triton dilution buffer
[100mM Tris-HCl pH 8.6, 100mM NaCl, 5mM EDTA pH 8, 5% Triton X-100]) supplemented with
protease/phosphatase inhibitor cocktail, and samples were sonicated using sonication beads
(Diagenode; Cat #C01020031) and the Bioruptor Pico Sonicator (Diagenode; Cat #B01060010)
for 20 cycles (each cycle 30 seconds on/30 seconds off) or till DNA fragments have 100-300bp
size. Samples were centrifuged at 16000g for 20 minutes at 4ºC. Chromatin was quantified,
and 30 µg of chromatin was incubated with every 5µg of antibody rotating for 16 hours (or
overnight) at 4ºC. The following antibodies were used: H3K4me3 (Abcam, Cat #ab8580),
H3K27me3 (Millipore, Cat ##07-449), H3K9me3 (Abcam, Cat #ab8898), H3K27Ac (Abcam, Cat
#ab4729), H3K4me1 (Abcam, Cat #ab8895). Samples were pulled down by incubation with
Protein A-Sepharose™ CL-4B (previously hydrated and blocked with 0.05% BSA) for 3h at 4ºC
in rotation. Sample was washed with a low salt washing buffer (50mM HEPES pH 7.5, 140mM
NaCl and 1% Triton) three times, one wash with a high salt wash buffer (50mM HEPES pH 7.5,
500mM NaCl and 1% Triton) and one wash with TE pH 8 (10mM Tris-HCl pH8, 1mM EDTA).
Samples were then eluted in 1% SDS and 100mM NaHCO$_3$. Samples were descrosslinked by
incubating for 16 hours (or overnight) at 65ºC shaking at 450 rpm, and were treated with
Proteinase K for 1h at 45ºC. DNA was purified using QIAquick PCR Purification Kit (Qiagen, Cat
#28106), following the manufacturer's instructions.

For ChIP IκBα-3XFLAG, a previous cross-linking step using Gold (Diagenode, Cat #C01019027)
was performed following manufacturer's instructions, and cells (IκBα$^{3XFLAG/3XFLAG}$ and IκBα-WT
mESCs) were later treated with 1% formaldehyde as mentioned above. Cell nuclei were
isolated by incubating cell pellets in a buffer containing 5mM Pipes pH 8, 85mM KCl and 0.5%
NP40, and later incubated on ice for 10 minutes followed by centrifuge at 4000 rpm for 5
minutes at 4ºC. Nucleus were lysed in a buffer containing 1% SDS, 10mM EDTA and 50 mM
Tris-HCl pH 8.1 and incubated on ice for 10 min. Samples were sonicated as mentioned above
and SDS was removed by incubating the sonicated sample on ice for 1 hour and later
centrifuged for 15 at 4ºC and 16000g. 60µg of chromatin was incubated with 10 µg of ANTI-
FLAG® M2 (Sigma-Aldrich, Cat #F1804). Samples were pulled down by a 50/50 mix of
Dynabeads A (Invitrogen, Cat #10002D) and G (Invitrogen, Cat #10003D). Samples were
washed 4 times in 0.1% SDS, 1% Triton-X, 2mM EDTA, 20mM Tris-HCl pH 8 and 150mM NaCl;
followed by 1 wash in TE pH 8.

For ChIP followed by PCR, 1 µL of ChIP sample was normalized by its input and enrichment
was calculated based on IgG isotype control. Oligos used are found in supplementary table 1.

For ChIP followed by sequencing (ChIP-seq) Purified DNA concentration and integrity were
determined using Agilent Bioanalyzer (Agilent Technologies; Cat #G2939BA). Libraries were

prepared using standard protocols. Chromatin was sequenced using Illumina HiSeq platform
(Illumina, Inc.) (50bp single-end reads). Samples sequencing depth range was the following:
(i) H3K4me3: 2.56-2.74x10^7 reads, (ii) H3K27me3: 6.19-7.2x10^7 reads, (iii) H3K9me3: 7.12-
8.92x10^7 reads, (iv) IκBα-3XFLAG: 9.81-10.86x10^7 reads, (v) H3K27Ac: 9.11-10.65x10^7 reads
and (vi) H3K4me1: 9.76-10.58x10^7 reads.

ChIP-seq data analysis

Quality control was performed on raw data with FASTQC tool. Raw reads were trimmed to
remove adapters presence with Trimgalore (v0.6.6) [52]. Default parameters were used except
for a minimum quality of 15 (Phred score) and an adapter removal stringency of 3bp overlap.
For H3K4me3, H3K27me3, H3K4me1 and IκBα-3XFLAG or   IκBα-WT data, trimmed reads
were aligned to reference genome with Bowtie2 (v2.4.4) which was executed with default
parameters [53]. Required genome index was built with corresponding GRCm38 fasta file
retrieved from Ensembl (http://ftp.ensembl.org/pub/release-102/). Multimapped reads and
those exhibiting MAPQ < 20 were removed. For H3K9me3 data, trimmed reads were aligned
to reference genome (GRCm38 release 102 from Ensembl) with ShortStack (v3.8.5) [54].
Randomly placed multimappers were removed from the mapped reads. For all cases,
duplicated reads were marked with SAMtools (v1.15) [55]. Drosophila Melanogaster spike-ins,
present in H3K27me3, were discarded. NSC and RSC quality metrics were computed with
PhantomPeakQualTools (v1.2) [56]. ENCODE blacklisted regions (mm10 v2) were removed prior
to peak calling except for H3K9me3 (Amemiya et al., 2019). BigWig files were individually
generated using deepTools (v3.5.1) bamCoverage with -ignoreDuplicates -binSize 10 -
smoothLength 30 -effectiveGenomeSize 2308125349 -normalizeUsing RPGC and -
extendReads *Fragment_Length* options [57]. The effective genome size was particularized to a
read   length   of   50bp   and   directly   retrieved   from   deepTools   web   site
(https://deeptools.readthedocs.io/en/develop/content/feature/effectiveGenomeSize.html).
Fragment_Length was retrieved from PhantomPeakQualTools results except for H3K9me3
where 150bp value was considered. Duplicates were kept in IκBα-3XFLAG for BigWig files
generation. For all histone marks except for H3K4me3, peak calling was conducted by means
of epic2 (v0.0.52) with -effective-genome-fraction 0.8452 -fragment-size *Fragment_Length*
options and chromosome sizes only referring to canonical chromosomes [58]. For H3K4me3 and
IκBα-3XFLAG/WT, MACS2 (v2.2.7.1) was used to identify peaks with -nomodel -extsize
Fragment_Length -g 2308125349 options [59]. The corresponding input sample was used in all
peak calling computations. For the histone marks, peaks were called with adjusted p-values
(FDR) < 0.05. In case of IκBα-3XFLAG/WT samples, peaks were called with adjusted p-values
(FDR) < 0.05 and a minimum 4-fold enrichment.

Peak annotation was performed with ChIPseeker R (v4.2.1) package (v1.34.1) considering a
TSS region range between 5000 bp upstream and 100 bp downstream (Yu et al., 2015).
Required TxDb object was generated from the Ensembl GRCm38 gtf file retrieved
(http://ftp.ensembl.org/pub/release-102/). A consensus peakset was obtained from the

three biological replicates per histone mark and condition (IκBα-WT or IκBα-KO) by identifying
overlapping peaks in at least two out of the three replicates. In case of IκBα ChIP-seq data,
peaks from consensus peakset identified from the -3XFLAG samples not overlapping with
IκBα-WT were considered the high-fidelity peaks for this experiment. For this purpose,
findOverlapsOfPeaks function from ChIPpeakAnno R package (v3.32.0) was used with default
parameters [60].

Differential binding analysis (DBA) was conducted with DiffBind (v3.8.4) per histone mark [61].
Default parameters were used except for the summits parameter which was set to 500bp for
all histone marks except for H3K4me3 (150bp) in order to consider, for testing, intervals of
1000bp or 300bp respectively. Summit values were selected to have interval widths between
the minimum and first quartile peak width values for each histone mark. EdgeR was the
statistical method used for all the three analyses.　Differentially bound regions (DBRs)
between IκBα-KO and IκBα-WT samples were called with adjusted p-values (FDR) < 0.05.
Identified DBRs with any annotated gene were plotted with EnrichedHeatmap R package
(v1.18.1) [62]. To summarize replicates per condition, normalizeToMatrix function was used
based on corresponding BigWig files in w0 mean mode in 50bp windows. Tracks visualization
were obtained by means of Integrative Genomics Browser (IGV) tool [63].

Enhancers activity identification

Consensus peaksets derived from the three H3K27ac and H3K4me1 IκBα-WT mESCs replicates
were used to identify poised and active enhancers in an IκBα-WT scenario. Putative poised
enhancers were defined as those regions with H3K4me1 but no H3K27ac peaks. Putative
active enhancers were defined as those regions with H3K27ac and presence/absence
H3K4me1 peaks. To assess any differential enhancers activity in IκBα-KO vs IκBα-WT mESCs
cultured in Serum/LIF, differentially bound regions and consensus peaksets (IκBα-KO)
identified in H3K4me1 and H3K27ac were required. Putative gained or lost poised/active
enhancers in IκBα $^{-/-}$ were obtained with following criteria: (i) for gained/lost poised enhancers
in IκBα $^{-/-}$: a differential H3K4me1 increase/decrease and absence of H3K27ac peaks in IκBα $^{-/-}$
consensus peakset, (ii) for gained/lost active enhancers (without H3K4me1) in IκBα-KO: a
differential H3K27ac increase/decrease and absence of H3K4me1 in IκBα-KO consensus
peakset and (iii) for gained/lost active enhancers (with H3K4me1) in IκBα-KO: a differential
H3K27ac increase/decrease and presence of H3K4me1 in IκBα $^{-/-}$ consensus peakset.

A maximum gap of 1000bp was allowed for checking the overlap between two different
histone marks. For this purpose, the 'subsetByOverlaps' function from the IRanges R package
was used (v.2.34.1).

Hi-C data visualization and TADs annotations

Juicebox v1.9.8 was run locally to visualize Hi-C data from [42], specifically in the mESCs
(Serum/LIF) dataset. Corresponding maps from this dataset were already included in the tool.

To visualize contact maps, balanced normalization was considered and applied color range up
to 3 in the selected regions. Additionally, the list of 2,712 TADs with their corresponding
genome coordinates was retrieved and used to locate the identified IκBα binding peaks,
enhancers in the IκBα-WT scenario and showing differential activity together with
differentially expressed genes from in-house RNAseq data.

RNA-seq experiments

Total RNA from three independent clones from mESCs, 48h EBs and 96h EBs was isolated
using the RNeasy Plus Mini Kit (Qiagen; Cat #74136) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
RNA integrity was addressed by agarose gel and Agilent Bioanalyzer (Agilent Technologies;
Cat #G2939BA). Libraries sequenced using Illumina HiSeq 2500 (Illumina, Inc.) (125bp paired-
end reads). Samples sequencing depth ranged between 35M and 52M reads (average 41M
reads) per sample.

RNA-seq data analysis

Quality control was performed on raw data with the FASTQC tool (v0.11.9). Raw reads were
trimmed to remove adapter presence with Trimgalore (v0.6.6) [52]. Default parameters were
used except for a minimum quality of 15 (Phred score) and an adapter removal stringency of
3bp overlap. Trimmed reads were aligned to reference the genome with STAR aligner tool
(v2.7.8). STAR was executed with default parameters except for the number of allowed
mismatches which was set to 1. Required genome index was built with corresponding
GRCm38 gtf and fasta files retrieved from Ensembl (http://ftp.ensembl.org/pub/release-
102/). Obtained BAM files with uniquely mapped reads were considered for further analysis.
Raw gene expression was quantified using featureCounts tool from subRead software (v2.0.1)
with exon as feature [64]. The raw counts matrix was imported into the R Statistical Software
environment (v4.2.1) for downstream analysis. Raw expression matrix included 55,487 genes
per 18 samples in total. Experimental design considered three timepoints: mouse ESCs, EBs
at 48h and EBs at 96h. Each time point included 6 samples distributed in two conditions: 3
IκBα-WT and 3 IκBα-KO. Prior to statistical analysis, those genes with less than 10 raw counts
across the 6 samples under test were removed. After pre-filtering, 21,843 genes (mESCs),
21,424 genes (EBs 48h) or 21,664 genes (EBs 96h) were available for testing. For visualization
purposes, counts were normalized by the variance-stabilizing transformation method as
implemented in DESeq2 R package [65] (v1.38.3). Differential expression analysis (DEA) was
conducted with DESeq2. Each time point was independently analyzed. Fitted statistical model
included sample conditions as covariable with IκBα-WT as the reference. Obtained log2 fold
change values were shrunken with apeglm shrinkage estimator R package (v1.20.0) [66]. Raw p-
values were adjusted for multiple testing using the Benjamini-Hochberg False Discovery Rate
(FDR) [67]. Differentially Expressed Genes (DEGs) between IκBα-KO and IκBα-WT samples were

called with adjusted p-values (FDR) < 0.05 and absolute shrunken log2 Fold change > 1. Data
visualization was performed with the ggplot2 (v3.4.1).

Genome-wide DNA methylation samples preparation

DNA from frozen mESCs pellets was extracted using DNeasy Blood and Tissue Kit (Qiagen
GmbH, Hilden, Germany). Purified genomic DNA was quantified with Qubit (Invitrogen,
Carlsbad, CA, USA) according to manufacturer's instructions. Infinium Mouse Methylation
BeadChip (Illumina, Inc., San Diego, CA, USA) arrays were used to profile DNA methylation
genome wide. This platform allows over 285,000 methylation sites per sample to be
interrogated at single-nucleotide resolution. The samples were bisulfite converted using EZ
DNA Methylation-Gold™ Kit (Zymo Research, CA, USA) and were hybridized in the array
following the manufacturer's instructions.

DNA methylation data analysis

The DNA methylation profile of the studied samples was assessed using the Infinium Mouse
Methylation BeadChip Array (~285,000 methylation sites) as previously described [68]. Briefly,
raw signal intensities were obtained with GenomeStudio Software 2011.1 (Illumina) and DNA
methylation beta values were computed from raw IDAT files using GenomeStudio default
normalization with control probes and background subtraction. Quality control steps to
remove erratic probe signals were performed within the R statistical environment (v4.0.3).
We removed probes with detection p value > 0.01, genotyping probes and manufacturing
flagged   (MFG)   probes   described   in   the   Illumina   manifest   file
(https://support.illumina.com/downloads/infinium-mouse-methylation-manifest-file.html).
The differentially methylated probes (DMPs) between IκBα-WT and IκBα-KO samples were
computed separately for each time point (mESCs and EBs at 96h) by deriving a linear model
with the limma R package (v3.46.0). Each condition included 3 samples per time point. DMPs
with adjusted p-value (FDR) <0.05 and absolute mean methylation beta value difference
between conditions >0.3 were considered significant. DNA methylation analysis was
performed using the mm10 mouse genome reference build and the complete annotation was
downloaded   from   the   annotated   manifest   file
(http://zwdzwd.github.io/InfiniumAnnotation#mouse) [38].

Functional analysis

Overrepresentation analysis was applied over lists of selected genes derived from RNA-seq
data (DEGs) or from ChIP-seq data (differentially bound regions). The Gene Ontology
(Biological Process ontology, GO BP terms), KEGG PATHWAY and WikiPathways databases for
Mus Musculus [69–71] were interrogated by means of clusterProfiler R package (v4.6.2) (Wu et
al., 2021). Corresponding Entrez identifiers were used. Benjamini-Hochberg procedure was
used to obtain adjusted p-values. Obtained GO BP terms were simplified using the simplify

function from clusterProfiler with default parameters. Overrepresented terms or processes

were called with adjusted p-values (FDR) < 0.05.

Gene Set Enrichment Analysis (GSEA) was performed for mESCs against the three cell states

signatures defined by other authors [72]. Briefly, each signature was composed of 200 genes

uniquely expressed in each state obtained from bulk RNA-seq performed per state. For this

purpose, the complete list of genes from mESCs samples (21,843 genes) was ranked based on

the shrunken log2 Fold Change obtained from DEA. GSEA was conducted through the

fgseaMultilevel function from fgsea R package (v1.24.0) [73] with default parameters.

Enrichment plots were generated with the same package.

Additionally, the testEnrichment function from SeSAMe R package (v1.14.2) [74] was used for

conducting functional analysis with default parameters over the list of DMPs. Probe design,

transcription factor binding site and histone modifications consensus database sets, included

in the same package, were interrogated.

Gene Set Variation Analysis (GSVA) R package (v1.46.0) with default parameters were used to

obtain Z-score values for genes annotated to endoderm (GO:0001706), mesoderm

(GO:0001707) and ectoderm (GO:0001705) GO BP terms [75].

Statistical analysis

Statistical analysis was performed with GraphPad Prism v.8.0.1. (GraphPad Software, Inc.). Unless specified, the comparison between two groups was performed with unpaired two-sided t-test. A p-value < 0.05 was considered significant.

DATA AVAILABILITY

All sequencing data is deposited at GEO under a SuperSeries with the accession number GSE239565)[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE239565]. Individual SubSeries can be found at: GSE239563 (RNA-seq data), GSE239564 (ChIP-seq data) and GSE239562 (Methylation array data).

CODE AVAILABILITY

Scripts used to process the generated bulk RNA-seq and ChIP-seq data are available in Github repository: https://github.com/BigaSpinosaLab/IkBa_Naive_Pluripotency

ACKNOWLEDGEMENTS

We would like to acknowledge all members of the Bigas/Espinosa labs for helpful discussions. We are grateful for technical support to CRG/UPF Flow Cytometry Unit, CRG Genomics and Advanced Light Microscopy Units. This work has been supported by Spanish Ministry of Science and Innovation (PID2019-104695RB-I00, PLEC2021-007518, PDC2021-120817-I00), Generalitat de Catalunya (2021SGR 39) and Departament de Salut (SLT002/16/00299) to AB; the Spanish Ministry of Science and Innovation (PID2019-108322GB-100 and PID2022-142679NB-I00) to LDC; the Spanish Ministry of Science and Innovation (PID2021-123383NB-I00.) and the Agencia de Gestió d'Ajuts Universitaris i de Recerca (2021 SGR 01222) to BP. L.G.P has been a recipient of FI AGAUR fellowship (2019 FI-B 00151/2020 FI_B1 00130) from Generalitat de Catalunya. D.A-V was funded by the FIS fellowship (FI20/00130) from Instituto Carlos III. M.B. received funding from the Ramón Areces Foundation. M.M. is a recipient of a grant from the Instituto Carlos III, grant number CA22/00011 (co-funded by the European Social Fund Plus, ESF+ and by the European Union). A.V was a recipient of PI22-00548 funded by the Instituto de Salud Carlos III (ISCIII) and co-funded by the European Union.

AUTHOR CONTRIBUTION

Figure 1. IκBα specifically accumulates in the chromatin of the naïve pluripotent state. A.
Schematic of pluripotent states in mouse embryonic stem cells (mESCs). mESCs were
maintained regularly in naïve pluripotency by culturing them in serum/LIF medium, or they
were polarized towards the ground state of naïve pluripotency by culturing them for two
passages in LIF plus Gsk3β (CHIR99021 or Chiron) and MEK (PD0325091) inhibitors (2i/LIF).
mESCs were further differentiated towards the primed pluripotent state (epiblast-like cells)
by culturing mESCs in a medium containing 20ng/mL Activin A and 10ng/mL Fgf2 for 120h. **B.**
C. Relative mRNA levels of the canonical IκB genes (*Nfkbia, Nfkbib, Nfkbie*) (**B**) or Nf-κB
effector genes (*Rela, c-Rel, Relb, Nfkb1, Nfkb2*) (**C**) across the different states of pluripotency.
Expression was normalized using the house-keeping gene *Tbp* relative to 2i/LIF. Data from
three independent experiments. Dots indicate mean values and error bars refer to ± standard
deviation (SD). **D.** Western blot analysis of IκBα protein distribution in cytoplasm,
nucleoplasm (Nucleus) or chromatin in cells at the ground-state of naïve pluripotency (2i/L),
naïve pluripotency (S/L) or primed pluripotency (Epi). **E.** Transcriptome and proteome
differences across naïve (2i/LIF) and primed (Epiblast-like cells or EpiSCs) pluripotent states.
Enriched transcripts and proteins in naïve state are highlighted in red (Log$_2$ fold-change < -1).
Enriched transcripts and proteins in the primed state are marked in dark gray (Log$_2$ fold-
change > 1). Data from [33]. **F.** Heatmap showing the expression levels of pluripotency (*Dppa4,*
Pou5f1, Sox2, Prdm14, Klf4, Zfp42, Nanog), differentiation (*Mesp1, Gata4, Gata6, Lhx1,*
Wnt3a, Fgf5, Nes, Axin2, Wnt3, Eomes, Foxa2, T, Mixl1, Lef1) and NF-κB (*Nfkbia, Nfkbib,*
Nfkbie, Nfkbiz, Rela, c-Rel, Relb) genes from the RNAseq samples at mESCs (Serum/LIF) and
differentiating cells (embryoid bodies) at 48h and 96h of IκBα-WT cells. . **G.** Schematic of the
approach followed to deplete IκBα protein in mESCs (upper panel) and western blot analysis
of three independent IκBα-WT and IκBα-KO mESC clones (bottom panel). **H.** Boxplots, violin
plots and individual values representing the normalized gene expression levels in IκBα-WT or
IκBα-KO mESCs, EBs 48h and EBs 96h. Considered genes are those annotated to NF-κB
signaling pathway (KEGG PATHWAY mmu04064). Box Plots: center line, median; lower and
upper hinges, first and third quartiles; whiskers, 1.5x interquartile range (IQR) extended to
the largest/smallest value within 1.5xIQR; no outliers present. Unpaired two-sided Wilcoxon
test applied.

Figure 2. Lack of IκBα impairs the exit from naïve pluripotency. A. PC1 and PC2 from Principal
Component Analysis (PCA) of normalized RNA-seq data from 3 different time points (mESCs,
48h EBs and 96h EBs) of IκBα-WT and IκBα-KO cells (three independent clones from each
genotype). **B.** Heatmap showing the expression levels of naïve pluripotency genes (*Gab1,*
Sox2, Esrrb, Zfp42, Dppa4, Klf2, Prdm14, Klf4, Tfcp2l1) from the RNAseq samples of IκBα-WT
and IκBα-KO EBs at 96h of **C, D, E**. Z-score values for gene sets referring to Endoderm (C),
Mesoderm (D) and Ectoderm (E) formation from The Gene Ontology database (GO:0001706,
GO:0001707 and GO:0001705 terms, respectively) and obtained from   corresponding
expression levels at mESCs, 48h and 96h EBs IκBα-WT or IκBα-KO cells (three independent

clones of each genotype). Loess curves are also represented with 95% confidence intervals.
F. Representative immunofluorescence of 216h EBs stained with the pluripotency markers
NANOG and OCT3/4. **G.** Quantification of total number of alkaline phosphatase (AP) colonies
after 216h EB differentiation and additional culture of 96h in Serum/LIF (S/L) medium. Dots
represent values from two independent experiments from three different IκBα-WT or IκBα-
KO clones. Unpaired two-sided t-test applied. **H.** Schematics of mESCs differentiation towards
Epiblast Stem Cells (EpiSCs). mESCs cultured in Serum/LIF were then seeded in N2B27 medium
supplemented with 10ng/mL Fgf2 and 20ng/mL Activin A for 120h (see Materials and
Methods section for further information). **I.** Relative RNA levels of naïve pluripotency genes
(*Dppa3, Nanog, Sox2, Rex1, Klf2, Klf4, Gbx2, Tbx3*) in IκBα-WT or IκBα-KO EpiSCs at 120h. RNA
levels were normalized based on *Tbp* expression relative to IκBα-WT. Each dot represents an
independent clone either from IκBα-WT or IκBα-KO genotypes. Unpaired two-sided t-test
applied. Horizontal bars indicate mean values and error bars refer to ± SD.

Figure 3. IκBα-KO mESCs preserve the ground-state of naïve pluripotency under Serum/LIF
culture. A. GSEA results comparing IκBα-KO vs IκBα-WT mESCs cultured in Serum/LIF against
ground (2i/LIF) and naïve (Serum/LIF) state pluripotency signatures retrieved from [36].
Adjusted p-value by Benjamini-Hochberg procedure and Normalized Enrichment Score (NES)
indicated. **B.** Heatmap showing the expression levels of the naïve pluripotency genes from
the RNAseq samples of IκBα-KO and IκBα-WT mESCs cultured in Serum/LIF. **C.** Number of
differentially bound regions (adjusted p-value <0.05, FDR) for H3K4me3, H3K27me3 and
H3K9me3 histone marks in IκBα $^{-/-}$ vs IκBα $^{+/+}$ mESCs cultured in Serum/LIF. Enriched or
reduced regions in IκBα $^{-/-}$ mESCs are distinguished (log$_2$ Fold-change >0 or <0, respectively).
Overrepresented WikiPathways, for (left) H3K4me3 and (center) H3K27me3, or The Gene
Ontology terms, for (right) H3K9me3 (adjusted p-value < 0.05, FDR) are indicated per mark.
Annotated genes to differential regions showing H3K4me3 enrichment, H3K27me3 reduction
or H3K9me3 reduction in IκBα-KOmESCs were considered. A one-sided hypergeometric test
was conducted. **D.** Representative naïve pluripotency genes having differential H3K4me3
(enriched) and H3K27me3 (reduced) levels in IκBα $^{-/-}$ vs IκBα $^{+/+}$ mESCs cultured in Serum/LIF.
Shadowed regions highlight the differential levels of the histone marks. **E.** Representative
immunofluorescence images of DNA-methylation-related mark 5-methylcytosine (5mC) in
IκBα-WT (left) and IκBα-KO (right) mESCs cultured in Serum/LIF. **F.** DNA methylation profile in
IκBα-WT vs IκBα-KO mESCs (left) and 96h EBs (right) showing the distribution density of mean
ß-values from all 261,220 CpGs under test per condition. **G.** DNA methylation levels in
different genomic regions in either IκBα-WT vs IκBα-KO mESCs or 96h EBs: promoters, gene
bodies, enhancers and imprinted regions. Number of CpGs analyzed in each category is
indicated between brackets. Unpaired two-sided t-test was applied.

Figure 4. IκBα binds to enhancer regions to prevent their hyperactivation in the naïve
pluripotency state. A. Distribution of genomic regions annotated to the 817 identified IκBα
chromatin binding peaks. **B, C.** H3K27Ac (B) and H3K4me1 (C) distribution profiles in IκBα-WT
vs IκBα-KO mESCs cultured in Serum/LIF across IκBα-bound chromatin regions. Histone marks

data generated in this study. **D.** Number of regions with differential enhancer activity in IκBα-
KO vs IκBα-WT mESCs cultured in Serum/LIF. Increased/Reduced activity for (left) poised
enhancers based on the H3K4me1 differential binding and absence of H3K27ac overlapping
peaks in IκBα-KO mESCs and, (right) active enhancers based on the differential binding of
H3K27ac and presence of H3K4me1 overlapping peaks in IκBα-KO. Differential binding based
on adjusted p-value <0.05 (FDR). Increased or reduced regions in IκBα-KO are distinguished
by log$_2$ Fold-change >0 or <0, respectively, compared to IκBα-WT mESCs. Overrepresented
WikiPathways for annotated genes to (left) poised enhancers with reduced activity and (right)
active enhancers with increased activity in IκBα-KO vs IκBα-WT mESCs are indicated
(adjusted p-value < 0.05, FDR). A one-sided hypergeometric test was conducted. **E.**
Representative naïve pluripotency genes having an increase in active enhancer status
(H3K27Ac enrichment without changes in H3K4me1) in IκBα-KO mESCs cultured in Serum/LIF.
Shadowed regions highlight the differential levels of the histone mark. **F.** Heatmap showing
the expression levels of selected naïve pluripotency genes contained in TADs with active
enhancer marks (H3K27Ac and H3K4me1) and IκBα chromatin binding from RNAseq samples
of IκBα-WT and IκBα-KO mESCs and 48h EBs. **G.** (Upper panel) Contact map (Hi-C data from
[42]) of the TAD (chr5:118,785,000-119,707,000) which contains the Tbx3 locus. An extended
region is shown (chr5:118660246-119843912). Arrows indicate the interaction between the
IκBα-bound region and TSS of Tbx3 gene (upper arrow) or differential active enhancer
(bottom arrow). (Center panel) Genome browser showing ChIP-seq data from H3K27Ac and
H3K4me1 in IκBα-WT and IκBα-KO mESCs, and identified IκBα chromatin peaks. Shadowed
areas indicate differential active enhancer (light gray), the TSS of the gene of interest (darker
gray) or IκBα peaks (dark green). (Bottom panel) *Tbx3* expression (RNA-seq data) in IκBα-WT
and IκBα-KO mESCs and 48h EBs from two independent clones of each genotype. **H.**
Schematic representation of IκBα reconstitution of IκBα-KO mESCs. IκBα-KO mESCs were
reconstituted by introducing a doxycycline-inducible IκBα construct (i-IκBα^{WT}). mESCs were
then kept in Serum/LIF medium supplemented with 1μg/mL Doxycycline for 4 consecutive
passages. **I.** Western blot analysis of H3K27Ac levels in i-IκBα^{WT} mESCs (left panel). Two
independent experiments were performed to quantify H3K27Ac levels (right panel).

Figure 5. IκBα safeguards enhancer hyperactivation in naïve pluripotency in an NF-κB-
independent manner. A. Schematic of chromatin-specific/NF-κB-deficient (IκBα$^{ΔNF-κB}$) or NF-
κB-specific/chromatin-deficient (IκBα^{ΔChrom}) IκBα mutants (Separation-Of-Function or SOF
mutants). **B.** Pull-down of GST-H2A with cell lysates from different SOF IκBα mutant mESCs.
C. Pull-down of GST-p50 with lysates from SOF IκBα mutant mESCs. **D.** Western blot analysis
of chromatin protein extracts of doxycycline-inducible IκBα forms (i-IκBα^{ΔWT}, i-IκBα$^{ΔNF-κB}$,
IκBα^{ΔChrom}) after 16h of doxycycline induction. **E.** Western blot analysis of H3K27Ac levels in i-
IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and IκBα^{ΔChrom} mESCs (upper panel). H3K27Ac levels were quantified
(bottom panel). **F, G.** ChIP followed by PCR of H3K27Ac (F) and H3K4me1 (G) marks within
genomic regions bound by IκBα in i-IκBα$^{ΔNF-κB}$ and IκBα^{ΔChrom} mESCs cultured in Serum/LIF.
Unpaired two-sided t-test was performed. Bars indicate mean values and error bars refer to

± standard error of the mean (SEM). **H.** Representative immunofluorescence images of DNA-methylation-related mark 5-methylcytosine (5mC) in i-IκBα$^{\Delta WT}$ (upper panel), i-IκBα$^{\Delta NF-\kappa B}$ (medium panel), IκBα$^{\Delta Chrom}$ (bottom panel) mESCs cultured in Serum/LIF. **I.** Flow cytometry analysis of the pluripotency surface marker SSEA-1 at 216h EBs in i-IκBα^{WT}, i-IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta Chrom}$ cells. **J.** Quantification of Alkaline Phosphatase (AP) staining in 216h EBs upon reconstitution with the different IκBα forms. Each dot represents an independent experiment. One-way ANOVA was applied. Bars indicate mean values and error bars refer to ± SEM. **K.** Representative immunofluorescence for NANOG and OCT3/4 at 216h EBs in i-IκBα^{WT} (upper panel), i-IκBα$^{\Delta NF-\kappa B}$ (medium panel) and i-IκBα$^{\Delta Chrom}$ (bottom panel) cells.

Figure 6. Proposed model: IκBα attenuates enhancer activity in the naïve-to-primed pluripotency transition. In mESCs cultured in Serum/LIF, IκBα binds to distal regions to maintain a moderate pluripotency gene expression by avoiding enhancer hyperactivation. This allows naïve pluripotency exit and differentiation upon LIF withdrawal. In the absence of IκBα, enhancer regions acquire a higher activation status, resulting in an aberrantly increased expression of pluripotency genes. This leads to an impaired pluripotency exit and differentiation.

Figure S1. Depletion of IκBα in mESCs compromises the exit from naïve pluripotency. Related to Figure 2. A. Volcano plot from differential expression analysis between IκBα-KO vs IκBα-WT 96h EBs. Dashed vertical lines represent an absolute shrunken log$_2$ Fold Change = 1 and dashed horizontal line represents an adjusted p-value = 0.05 (FDR). **B.** Boxplots and individual values showing the quantification of relative size of 168h EBs. Each dot represents an embryoid body. Unpaired two-sided t-test was applied. **C.** Relative RNA levels of naïve pluripotency genes (*Pou5f1, Gbx2, Klf2, Sox2, Rex1, Nanog*) in IκBα-WT or IκBα-KO 216h EBs. RNA levels were normalized based on *Tbp* expression relative to IκBα-WT . Each dot represents an independent clone either from IκBα-WT or IκBα-KO genotypes. Unpaired two-sided t-test was applied. **D.** Schematic of Alkaline Phosphatase (AP) staining carried out in EBs. IκBα-WT or IκBα-KO 216h EBs were dissociated and seeded in Serum/LIF medium for another 96h. Attached cells were screened for AP$^+$ total cell number. **E.** Flow cytometry plots of SSEA-1 from IκBα-WT or IκBα-KO EBs 216h. **F.** Schematic of teratoma formation assay. 5x10^5 IκBα-WT or IκBα-KO mESCs cultured in Serum/LIF were injected intramuscularly in the leg of immunocompromised NSG mice. 6 weeks after the transplant, teratomas were formed, and mice were euthanized for further teratoma analysis. **G.** Representative images of immunohistochemistry for OCT3/4 in IκBα-WT (left panel) or IκBα-KO (right panel) teratomas. Teratomas were derived from three independent clones of mESCs of both genotypes. **H.** Percentage of OCT3/4+ cells in teratomas. Positive cells were counted from five different microscope fields from each IκBα-WT and IκBα-KO clone. Unpaired two-sided t-test was performed. Bars indicate mean values and error bars refer to ± SD. **I.** Brightfield images of IκBα-WT or IκBα-KO EpiSCs at 120h. Each picture corresponds to an independent clone from either IκBα-WT or IκBα-KO genotypes.

Figure S2. Absence of IκBα favors the ground-state of naïve pluripotency in Serum/LIF conditions. Related to Figure 3. A. Relative RNA levels of the naïve pluripotency genes *Rex1 (Zfp42), Klf2* and *Tbx3* upon culture of IκBα-WT either in Serum/LIF or two passages in 2i/LIF. 2i/LIF IκBα-WT mESCs were compared with IκBα-KO mESCs cultured in Serum/LIF. Unpaired two-sided t-test was applied to calculate statistical significance. Each dot represents an independent clone from each of the two genotypes. **B, C, D.** Volcano plots from differential binding analysis of H3K4me3 (B), H3K27me3 (C) and H3K9me3 (D) in IκBα-WT vs IκBα-KO mESCs cultured in Serum/LIF. Representative genes with enriched or reduced levels of H3K4me3 (B), H3K27me3 (C) and H3K9me3 (D) marks in IκBα-KO mESCs are highlighted. **E, F.** DNA methylation levels (mean ß-values) across chromosomes in IκBα-WT vs IκBα-KO mESCs (E) or 96h EBs (F).

Figure S3. IκBα binds to enhancer regions to preserve their proper activation level. Related to Figure 4. A. Heatmap of the chromatin binding profile of IκBα in mESCs cultured in Serum/LIF. High-fidelity IκBα chromatin peaks (n=817) were retrieved by considering common peaks in at least two out of the three different replicates in IκBα$^{3XFLAG/3XFLAG}$ mESCs that were not identified in IκBα-WT mESCs. ChIP of 3XFLAG was performed in either of the two conditions (IκBα$^{3XFLAG/3XFLAG}$ or IκBα-WT mESCs). **B, C .** H3K27Ac (B) or H3K4me1 (C) distribution profile in naïve (Serum/LIF) or primed (EpiSCs) pluripotency state across IκBα-bound chromatin regions. Histone marks data retrieved from [76]. **D.** Western blot analysis of H3K27Ac (Active enhancer) and H3K4me1 (total enhancer) in IκBα-WT vs IκBα-KO mESCs cultured in Serum/LIF. Two independent IκBα-KO mESC clones were analyzed. **E.** Representative immunofluorescence images of H3K27Ac in mESCs cultured in Serum/LIF. **F, G.** Volcano plots from differential binding analysis of H3K4me1 (F) and H3K27Ac (G) in IκBα-WT vs IκBα-KO mESCs cultured in Serum/LIF. Representative genes annotated to regions with enriched or reduced levels of histones marks in IκBα-KO mESCs are highlighted. **H.** Distribution of Topologically Associated Domains (TADs) into A (active) and B (inactive) compartments in a genome-wide fashion (left) or only for those including at least one IκBα peaks (right). Respective proportion of differential enhancers (poised or active as defined in panel B) are shown. Among the 74% of genome-wide TADs with differential enhancers, 69% of them were located within A compartment. Among the 92% of TADs with IκBα peak with differential enhancers, 81% of them were located within A compartment. TADs were retrieved from [42], mESCs dataset. **I.** Representative immunofluorescence images of H3K27Ac in i-IκBα^{WT} (bottom panel) mESCs vs IκBα-KO mESCs (upper panel) cultured in Serum/LIF. **J.** Relative RNA levels of naïve pluripotency genes (*Zfp42, Klf2, Sox2, Tbx3*) in i-IκBα^{WT} mESCs after doxycycline induction. Relative RNA levels were calculated based on *Tbp* expression. Each dot represents an independent experiment. Unpaired two-sided t-test was performed. Bars indicate mean values and error bars refer to ± SD.

Figure S4. Generation of separation-of-function (SOF) IκBα mutants demonstrates that IκBα safeguards hyperactivation of enhancers in an NF-κB-independent manner. Related to Figure 5. A. Relative expression level of Nfkbia gene after overnight doxycycline induction in

i-IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-\kappa B}$ and IκBα$^{\Delta Chrom}$ mESCs. **B.** Western blot analysis of cytoplasmic protein

extracts of doxycycline-inducible IκBα forms (i-IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-\kappa B}$, i-IκBα$^{\Delta Chrom}$) after 16h of

doxycycline induction. **C.** Schematic representation of reconstitution of IκBα-KO mESCs with

either of the two doxycycline-inducible SOF mutants (i-IκBα$^{\Delta NF-\kappa B}$ or IκBα$^{\Delta Chrom}$). mESCs were

then kept in Serum/LIF medium supplemented with 1μg/mL Doxycycline for 4 consecutive

passages. mESCs were characterized at the epigenetic level and then their potential to be

further differentiated was addressed. **D.** Representative immunofluorescence images of

H3K27Ac in i-IκBα$^{\Delta NF-\kappa B}$ (upper panel) mESCs vs i-IκBα$^{\Delta Chrom}$ mESCs (bottom panel) cultured in

Serum/LIF**. E.** Relative RNA levels of naïve pluripotency genes (*Zfp42, Klf2, Sox2, Tbx3*) in i-

IκBα$^{\Delta NF-\kappa B}$ and i-IκBα$^{\Delta Chrom}$ mESCs after doxycycline induction. Unpaired two-sided t-test was

performed.Bars indicate mean values and error bars refer to +/- standard deviation. **F.**

Percentage of differentiated cells (determined as SSEA-1.) at 216h EBs 216h derived from i-

IκBα$^{\Delta WT}$, i-IκBα$^{\Delta NF-\kappa B}$ and IκBα$^{\Delta Chrom}$ mESCs. **G.** Representative images of Alkaline Phosphatase

(AP) staining in 216h EBs upon reconstitution with the different IκBα forms. **H.** Quantification

of EB differentiation status (percentage of pluripotent cells) at 216h after induction of SOF

IκBα mutants. Data from two independent experiments.

REFERENCES

1. Festuccia, N., Osorno, R., Wilson, V., and Chambers, I. (2013). The role of pluripotency gene regulatory network components in mediating transitions between pluripotent cell states. Curr. Opin. Genet. Dev. *23*, 504–511.

2. Graf, T., and Stadtfeld, M. (2008). Heterogeneity of embryonic and adult stem cells. Cell Stem Cell *3*, 480–483.

3. Martinez Arias, A., and Brickman, J.M. (2011). Gene expression heterogeneities in embryonic stem cell populations: origin and function. Curr. Opin. Cell Biol. *23*, 650–656.

4. Pera, M.F., and Rossant, J. (2021). The exploration of pluripotency space: Charting cell state transitions in peri-implantation development. Cell Stem Cell *28*, 1896–1906.

5. Wang, X., and Wu, Q. (2022). The Divergent Pluripotent States in Mouse and Human Cells. Genes *13*. 10.3390/genes13081459.

6. Ying, Q.-L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P., and Smith, A. (2008). The ground state of embryonic stem cell self-renewal. Nature *453*, 519–523.

7. Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H., Nichols, J., Kranz, A., Stewart, A.F., Smith, A., et al. (2012). The transcriptional and epigenomic foundations of ground state pluripotency. Cell *149*, 590–604.

8. Boroviak, T., Loos, R., Lombard, P., Okahara, J., Behr, R., Sasaki, E., Nichols, J., Smith, A., and Bertone, P. (2015). Lineage-Specific Profiling Delineates the Emergence and Progression of Naive Pluripotency in Mammalian Embryogenesis. Dev. Cell *35*, 366–382.

9. Brons, I.G.M., Smithers, L.E., Trotter, M.W.B., Rugg-Gunn, P., Sun, B., Chuva de Sousa Lopes, S.M., Howlett, S.K., Clarkson, A., Ahrlund-Richter, L., Pedersen, R.A., et al. (2007). Derivation of pluripotent epiblast stem cells from mammalian embryos. Nature *448*, 191–195.

10. Tesar, P.J., Chenoweth, J.G., Brook, F.A., Davies, T.J., Evans, E.P., Mack, D.L., Gardner, R.L., and McKay, R.D.G. (2007). New cell lines from mouse epiblast share defining features with human embryonic stem cells. Nature *448*, 196–199.

11. Weinberger, L., Ayyash, M., Novershtern, N., and Hanna, J.H. (2016). Dynamic stem cell states: naive to primed pluripotency in rodents and humans. Nat. Rev. Mol. Cell Biol. *17*, 155–169.

12. Guo, G., Yang, J., Nichols, J., Hall, J.S., Eyres, I., Mansfield, W., and Smith, A. (2009). Klf4 reverts developmentally programmed restriction of ground state pluripotency. Development *136*, 1063–1069.

13. Takahashi, S., Kobayashi, S., and Hiratani, I. (2018). Epigenetic differences between

naïve and primed pluripotent stem cells. Cell. Mol. Life Sci. *75*, 1191–1203.

14. Bao, S., Tang, F., Li, X., Hayashi, K., Gillich, A., Lao, K., and Surani, M.A. (2009).
Epigenetic reversion of post-implantation epiblast to pluripotent embryonic stem cells.
Nature *461*, 1292–1295.

15. Habibi, E., Brinkman, A.B., Arand, J., Kroeze, L.I., Kerstens, H.H.D., Matarese, F.,
Lepikhov, K., Gut, M., Brun-Heath, I., Hubner, N.C., et al. (2013). Whole-genome
bisulfite sequencing of two distinct interconvertible DNA methylomes of mouse
embryonic stem cells. Cell Stem Cell *13*, 360–369.

16. Leitch, H.G., McEwen, K.R., Turp, A., Encheva, V., Carroll, T., Grabole, N., Mansfield, W.,
Nashun, B., Knezovich, J.G., Smith, A., et al. (2013). Naive pluripotency is associated
with global DNA hypomethylation. Nat. Struct. Mol. Biol. *20*, 311–316.

17. Singer, Z.S., Yong, J., Tischler, J., Hackett, J.A., Altinok, A., Surani, M.A., Cai, L., and
Elowitz, M.B. (2014). Dynamic heterogeneity and DNA methylation in embryonic stem
cells. Mol. Cell *55*, 319–331.

18. Factor, D.C., Corradin, O., Zentner, G.E., Saiakhova, A., Song, L., Chenoweth, J.G.,
McKay, R.D., Crawford, G.E., Scacheri, P.C., and Tesar, P.J. (2014). Epigenomic
comparison reveals activation of "seed" enhancers during transition from naive to
primed pluripotency. Cell Stem Cell *14*, 854–863.

19. Buecker, C., Srinivasan, R., Wu, Z., Calo, E., Acampora, D., Faial, T., Simeone, A., Tan,
M., Swigut, T., and Wysocka, J. (2014). Reorganization of enhancer patterns in
transition from naive to primed pluripotency. Cell Stem Cell *14*, 838–853.

20. Zhang, Q., Lenardo, M.J., and Baltimore, D. (2017). 30 Years of NF-κB: A Blossoming of
Relevance to Human Pathobiology. Cell *168*, 37–57.

21. Aguilera, C., Hoya-Arias, R., Haegeman, G., Espinosa, L., and Bigas, A. (2004).
Recruitment of IκBα to the *hes1* promoter is associated with transcriptional repression.
Proceedings of the National Academy of Sciences *101*, 16537–16542.

22. Mulero, M.C., Ferres-Marco, D., Islam, A., Margalef, P., Pecoraro, M., Toll, A., Drechsel,
N., Charneco, C., Davis, S., Bellora, N., et al. (2013). Chromatin-bound IκBα regulates a
subset of polycomb target genes in differentiation and cancer. Cancer Cell *24*, 151–166.

23. Marruecos, L., Bertran, J., Álvarez-Villanueva, D., Mulero, M.C., Guillén, Y., Palma, L.G.,
Floor, M., Vert, A., Arce-Gallego, S., Pecharroman, I., et al. (2021). Dynamic chromatin
association of IκBα is regulated by acetylation and cleavage of histone H4. EMBO Rep.
22, e52649.

24. Marruecos, L., Bertran, J., Guillén, Y., González, J., Batlle, R., López-Arribillaga, E.,
Garrido, M., Ruiz-Herguido, C., Lisiero, D., González-Farré, M., et al. (2020). IκBα
deficiency imposes a fetal phenotype to intestinal stem cells. EMBO Rep. *21*, e49708.

25. Thambyrajah, R., Fadlullah, Z., Proffitt, M., Neo, W.H., Guillén, Y., Casado-Pelaez, M.,

Herrero-Molinero, P., Brujas, C., Castelluccio, N., González, J., et al. (2022). IκBα
controls dormancy induction in Hematopoietic stem cell development via retinoic acid.
bioRxiv, 2022.11.17.516971. 10.1101/2022.11.17.516971.

26. Gerondakis, S., Grumont, R., Gugasyan, R., Wong, L., Isomura, I., Ho, W., and Banerjee,
A. (2006). Unravelling the complexities of the NF-kappaB signalling pathway using
mouse knockout and transgenic models. Oncogene *25*, 6781–6799.

27. Kaltschmidt, C., Greiner, J.F.W., and Kaltschmidt, B. (2021). The Transcription Factor
NF-κB in Stem Cells and Development. Cells *10*. 10.3390/cells10082042.

28. Li, S., Sun, J., Zhang, B.-W., Yang, L., Wan, Y.-C., Chen, B.-B., Xu, N., Xu, Q.-R., Fan, J.,
Shang, J.-N., et al. (2024). ATG5 attenuates inflammatory signaling in mouse embryonic
stem cells to control differentiation. Dev. Cell. 10.1016/j.devcel.2024.01.026.

29. Torres, J., and Watt, F.M. (2008). Nanog maintains pluripotency of mouse embryonic
stem cells by inhibiting NFkappaB and cooperating with Stat3. Nat. Cell Biol. *10*, 194–
201.

30. Dutta, D., Ray, S., Home, P., Larson, M., Wolfe, M.W., and Paul, S. (2011). Self-renewal
versus lineage commitment of embryonic stem cells: protein kinase C signaling shifts
the balance. Stem Cells *29*, 618–628.

31. Espín-Palazón, R., and Traver, D. (2016). The NF-κB family: Key players during
embryonic development and HSC emergence. Exp. Hematol. *44*, 519–527.

32. Alvarez-Villanueva, D., Galan, L., Bertran, J., and Floor, M. (2023). Separation-of-
function mutants reveal the NF-κB-independent involvement of IκBα in the regulation
of stem cell and oncogenic programs. bioRxiv.

33. Atlasi, Y., Jafarnejad, S.M., Gkogkas, C.G., Vermeulen, M., Sonenberg, N., and
Stunnenberg, H.G. (2020). The translational landscape of ground state pluripotency.
Nat. Commun. *11*, 1617.

34. Hayden, M.S., and Ghosh, S. (2008). Shared principles in NF-kappaB signaling. Cell *132*,
344–362.

35. Basak, S., Kim, H., Kearns, J.D., Tergaonkar, V., O'Dea, E., Werner, S.L., Benedict, C.A.,
Ware, C.F., Ghosh, G., Verma, I.M., et al. (2007). A fourth IkappaB protein within the
NF-kappaB signaling module. Cell *128*, 369–381.

36. Ghimire, S., Van der Jeught, M., Neupane, J., Roost, M.S., Anckaert, J., Popovic, M., Van
Nieuwerburgh, F., Mestdagh, P., Vandesompele, J., Deforce, D., et al. (2018).
Comparative analysis of naive, primed and ground state pluripotency in mouse
embryonic stem cells originating from the same genetic background. Sci. Rep. *8*, 5884.

37. Yagi, M., Kishigami, S., Tanaka, A., Semi, K., Mizutani, E., Wakayama, S., Wakayama, T.,
Yamamoto, T., and Yamada, Y. (2017). Derivation of ground-state female ES cells
maintaining gamete-derived DNA methylation. Nature *548*, 224–227.

38.   Zhou, W., Hinoue, T., Barnes, B., Mitchell, O., Iqbal, W., Lee, S.M., Foy, K.K., Lee, K.-H.,
Moyer, E.J., VanderArk, A., et al. (2022). DNA methylation dynamics and dysregulation
delineated by high-throughput profiling in the mouse. Cell Genom *2*.
10.1016/j.xgen.2022.100144.

39.   Choi, J., Huebner, A.J., Clement, K., Walsh, R.M., Savol, A., Lin, K., Gu, H., Di Stefano, B.,
Brumbaugh, J., Kim, S.-Y., et al. (2017). Prolonged Mek1/2 suppression impairs the
developmental potential of embryonic stem cells. Nature *548*, 219–223.

40.   Di Stefano, B., Luo, E.-C., Haggerty, C., Aigner, S., Charlton, J., Brumbaugh, J., Ji, F.,
Rabano Jiménez, I., Clowers, K.J., Huebner, A.J., et al. (2019). The RNA Helicase DDX6
Controls Cellular Plasticity by Modulating P-Body Homeostasis. Cell Stem Cell *25*, 622–
638.e13.

41.   Wu, X., Wu, X., and Xie, W. (2023). Activation, decommissioning, and dememorization:
enhancers in a life cycle. Trends Biochem. Sci. *48*, 673–688.

42.   Bonev, B., Mendelson Cohen, N., Szabo, Q., Fritsch, L., Papadopoulos, G.L., Lubling, Y.,
Xu, X., Lv, X., Hugnot, J.-P., Tanay, A., et al. (2017). Multiscale 3D Genome Rewiring
during Mouse Neural Development. Cell *171*, 557–572.e24.

43.   Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling,
A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive
mapping of long-range interactions reveals folding principles of the human genome.
Science *326*, 289–293.

44.   Álvarez-Villanueva, D., Galán-Palma, L., Bertran, J., Floor, M., Solé, L., Lobo-Jarne, T.,
Maqueda, M., Rajbhandari, R., Marruecos, L., Villà-Freixa, J., et al. (2023). Separation-
of-function mutants reveal the NF-κB-independent involvement of IκBα in the
regulation of stem cell and oncogenic programs. bioRxiv, 2023.06.21.545928.
10.1101/2023.06.21.545928.

45.   Guo, Y.-L. (2019). The underdeveloped innate immunity in embryonic stem cells: The
molecular basis and biological perspectives from early embryogenesis. Am. J. Reprod.
Immunol. *81*, e13089.

46.   Brena, D., Bertran, J., Porta-de-la-Riva, M., Guillén, Y., Cornes, E., Kukhtar, D., Campos-
Vicens, L., Fernández, L., Pecharroman, I., García-López, A., et al. (2020). Ancestral
function of Inhibitors-of-kappaB regulates Caenorhabditis elegans development. Sci.
Rep. *10*, 16153.

47.   Lando, D., Ma, X., Cao, Y., Jartseva, A., Stevens, T.J., Boucher, W., Reynolds, N.,
Montibus, B., Hall, D., Lackner, A., et al. (2024). Enhancer-promoter interactions are
reconfigured through the formation of long-range multiway hubs as mouse ES cells exit
pluripotency. Mol. Cell *84*, 1406–1421.e8.

48.   Pantier, R., Tatar, T., Colby, D., and Chambers, I. (2019). Endogenous epitope-tagging of
Tet1, Tet2 and Tet3 identifies TET2 as a naïve pluripotency marker. Life Sci Alliance *2*.
10.26508/lsa.201900516.

49. Sroczynska, P., Lancrin, C., Pearson, S., Kouskoff, V., and Lacaud, G. (2009). In vitro
differentiation of mouse embryonic stem cells as a model of early hematopoietic
development. Methods Mol. Biol. *538*, 317–334.

50. Chen, Q., Luo, W., Veach, R.A., Hickman, A.B., Wilson, M.H., and Dyda, F. (2020).
Structural basis of seamless excision and specific targeting by piggyBac transposase.
Nat. Commun. *11*, 3446.

51. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years
of image analysis. Nat. Methods *9*, 671–675.

52. Ewels, P., Afyounian, E., and Schuster-Boeckler, B.F.K. (2012). TrimGalore: v0. 6.7-DOI
via Zenodo.(Zenodo, 2021). doi: 10.5281/zenodo. 5127899. 2. Langmead, B. & Salzberg,
SL Fast gapped-read alignment with …. Nat. Methods.

53. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357–359.

54. Axtell, M.J. (2013). ShortStack: comprehensive annotation and quantification of small
RNA genes. RNA *19*, 740–751.

55. Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham,
A., Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and
BCFtools. Gigascience *10*. 10.1093/gigascience/giab008.

56. Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S.,
Bernstein, B.E., Bickel, P., Brown, J.B., Cayting, P., et al. (2012). ChIP-seq guidelines and
practices of the ENCODE and modENCODE consortia. Genome Res. *22*, 1813–1831.

57. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res. *44*, W160–W165.

58. Stovner, E.B., and Sætrom, P. (2019). epic2 efficiently finds diffuse domains in ChIP-seq
data. Bioinformatics *35*, 4392–4393.

59. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq
(MACS). Genome Biol. *9*, R137.

60. Zhu, L.J., Gazin, C., Lawson, N.D., Pagès, H., Lin, S.M., Lapointe, D.S., and Green, M.R.
(2010). ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip
data. BMC Bioinformatics *11*, 237.

61. Stark, R., Brown, G., and Others (2011). DiffBind: differential binding analysis of ChIP-
Seq peak data. R package version *100*.

62. Gu, Z., Eils, R., Schlesner, M., and Ishaque, N. (2018). EnrichedHeatmap: an
R/Bioconductor package for comprehensive visualization of genomic signal

associations. BMC Genomics *19*, 234.

63. Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G.,
and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

64. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–
930.

65. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

66. Zhu, A., Ibrahim, J.G., and Love, M.I. (2019). Heavy-tailed prior distributions for
sequence count data: removing the noise and preserving large differences.
Bioinformatics *35*, 2084–2092.

67. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Series B Stat. Methodol. *57*,
289–300.

68. Garcia-Prieto, C.A., Álvarez-Errico, D., Musulen, E., Bueno-Costa, A., N. Vazquez, B.,
Vaquero, A., and Esteller, M. (2022). Validation of a DNA methylation microarray for
285,000 CpG sites in the mouse genome. Epigenetics *17*, 1677–1685.

69. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P.,
Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene ontology: tool for the
unification of biology. The Gene Ontology Consortium. Nat. Genet. *25*, 25–29.

70. Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res. *28*, 27–30.

71. Martens, M., Ammar, A., Riutta, A., Waagmeester, A., Slenter, D.N., Hanspers, K., A
Miller, R., Digles, D., Lopes, E.N., Ehrhart, F., et al. (2021). WikiPathways: connecting
communities. Nucleic Acids Res. *49*, D613–D621.

72. Hu, S., Metcalf, E., Mahat, D.B., Chan, L., Sohal, N., Chakraborty, M., Hamilton, M.,
Singh, A., Singh, A., Lees, J.A., et al. (2022). Transcription factor antagonism regulates
heterogeneity in embryonic stem cell states. Mol. Cell *82*, 4410–4427.e12.

73. Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N., and Sergushichev, A.
(2021). Fast gene set enrichment analysis. bioRxiv, 060012. 10.1101/060012.

74. Zhou, W., Triche, T.J., Jr, Laird, P.W., and Shen, H. (2018). SeSAMe: reducing artifactual
detection of DNA methylation by Infinium BeadChips in genomic deletions. Nucleic
Acids Res. *46*, e123.

75. Hänzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis
for microarray and RNA-seq data. BMC Bioinformatics *14*, 7.

76. Bleckwehl, T., Crispatzu, G., Schaaf, K., Respuela, P., Bartusel, M., Benson, L., Clark, S.J.,

1221 Dorighi, K.M., Barral, A., Laugsch, M., et al. (2021). Enhancer-associated H3K4
1222 methylation safeguards in vitro germline competence. Nat. Commun. *12*, 5771.

Figure 1

Figure 2

Figure 3

Figure 4

A IκBα binding

B H3K27Ac (Serum/LIF)

C H3K4me1 (Serum/LIF)

D Enhancer status in IκBα-KO mESCs (Serum/LIF)

E Tbx3 · Tfcp2l1 · Klf4 · Zfp42

F mESCs · EBs 48h

G chr5:118,660,246-119,843,912

H Doxy-inducible IκBα

I i-IκBα^WT mESCs (Serum/LIF) · H3K27Ac Intensity

Figure 5

Figure 6