Epigenetic modifications driving ground state pluripotency exit require an NF-κB-independent chromatin IκBα function

Luís G. Palma[1,2,δ], Daniel Álvarez-Villanueva[1,10,δ], María Maqueda[1,2], Mercedes Barrero[7], Arnau Iglesias[1,2], Joan Bertran[9], Damiana Alvarez-Errico[2], Carlos A. García-Prieto[2], Cecilia Ballaré[7], Virginia Rodriguez-Cortez[2], Clara Bueno[2], August Vidal[10], Alberto Villanueva[10], Pablo Menéndez[4,5,11,12,13], Luciano Di Croce[5,7,8], Bernhard Payer[7,8], Manel Esteller[2,4,5,6], Lluís Espinosa[1,*] & Anna Bigas[1,2,*]

[1]Program in Cancer Research. Hospital del Mar Research Institute, CIBERONC, Barcelona, Spain

[2]Josep Carreras Leukemia Research Institute, Barcelona Spain.

[4]Centro de Investigación Biomédica en Red Cancer (CIBERONC), Madrid, Spain.

[5]Institucio Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

[6]Physiological Sciences Department, School of Medicine and Health Sciences, University of Barcelona (UB), Barcelona, Catalonia, Spain.

[7]Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, Spain.

[8]Universitat Pompeu Fabra, Barcelona, Spain.

[9]Faculty of Sciences, Technology and Engineering, Universitat de Vic - Universitat Central de Catalunya, Vic 08500 Spain

[10]Institut Investigació Biomèdica de Bellvitge (IDIBELL), Hospitalet, Barcelona, Spain

[11]Stem Cell Biology, Immunotherapy and Developmental Leukemia Laboratory. Josep Carreras Leukemia Research Institute

[12]Spanish Network for Advanced Therapies (RICORS-TERAV). Carlos III Health Institute (ISCIII).

[13]Department of Biomedicine. University of Barcelona.

[δ] These authors contributed equally

Keywords: IκBα, NFKBIA, NF-κB, DNA Methylation, Polycomb, Embryonic Stem Cells, Pluripotency Exit, Germ Layer Specification,

*Corresponding authors:

Anna Bigas and Lluis Espinosa

Cancer Research Program. CIBERONC.

Hospital del Mar Research Institute

Doctor Aiguader 88, 08003, Spain.

Tel: +34 933160440; email: abigas@imim.es; lespinosa@imim.es

Summary

Inflammatory signals are key in development and cell differentiation but their orchestration

with pluripotency and stemness signals is poorly understood. Our previous work identified a

chromatin function of IκBα, the NF-κB inhibitor, that is crucial for differentiation in different

types of somatic stem cells. Here we demonstrate that deficiency of IκBα imposes a profound

chromatin rewiring defect that impacts on DNA methylation, histone post-translational

modifications and transcriptional regulation, stabilizing mouse embryonic stem cells (ESCs) in

a ground state of pluripotency while preventing them from pluripotency exit and

differentiation. By engineering separation-of-function mutants of IκBα with specific binding

to either NF-κB or histones, we demonstrate that regulation of pluripotency state by IκBα is

independent of NF-κB but requires the chromatin-related IκBα function.

INTRODUCTION

The pluripotent capability of Embryonic Stem Cells (ESCs) allows for the generation of all the somatic and germline lineages both *in vitro* and *in vivo*. However, murine ESCs (mESCs) exhibit high heterogeneity when cultured in serum plus Leukemia Inhibitory Factor (LIF) (S/L medium), with cells interconverting to different metastable states (Festuccia et al. 2013). Some cells retain a ground state pluripotency program, while others are primed for differentiation (Graf and Stadtfeld 2008; Martinez Arias and Brickman 2011). Fluctuations in pluripotency-related transcription factors (Kumar et al. 2014; Filipczyk et al. 2015), microRNA biogenesis (Wang et al. 2007; Kumar et al. 2014) and antagonistic effects on distal regulatory regions (Hu et al. 2022) have been proposed to contribute to this disparity in mESCs. However, the upstream mechanisms governing these interconvertible states remain to be uncovered.

The heterogeneity of mESCs can be abolished by culturing the cells with inhibitors of ERK and GS3K signaling, in combination with the use of LIF (2i/L) (Ying et al. 2008). Under both inhibitors plus LIF (2i/LIF) conditions, mESCs enter a ground state of naïve pluripotency promoting homogenous self-renewal while blocking any residual pro-differentiating activity (Ying et al. 2008). Changes in different histone post-translational modifications (PTMs) have been shown to activate pluripotency genes and silence pro-differentiation genes in the ground state (Marks et al. 2012). One of the major epigenetic differences between S/L and 2i/L conditions is the global DNA hypomethylation profile observed in mESCs cultured in 2i/L medium (Habibi et al. 2013; Leitch et al. 2013; Singer et al. 2014). However, long-term stabilization of the ground state of pluripotency is detrimental as it can limit the developmental potential of mESCs (Choi et al. 2017). Therefore, identifying novel players to fine-tune the equilibrium between ground and primed states is crucial for preserving both pluripotency stability and differentiation capability in mESCs.

Inflammation is an immune response to pathogens that involves different signaling molecules. One of the main effectors of the inflammatory cascade is NF-κB, whose canonical activation involves the phosphorylation of the IKK kinase complex, the phosphorylation and degradation of the NF-κB inhibitor, IκBα, and the nuclear translocation of the NF-κB heterodimer p50/p65 (Zhang, Lenardo, and Baltimore 2017). There is now robust evidence that inflammatory signals are crucial in stem cell development (Gerondakis et al. 2006; Kaltschmidt, Greiner, and Kaltschmidt 2021), though the integration of the inflammatory signals at the pluripotency state is still poorly understood. In this regard, classical NF-κB pathway elements are very lowly expressed in mESCs and overexpression of NF-κB effector proteins leads to loss of pluripotency. In contrast, overexpression of the IκBα super-repressor (S32-36A mutant) has been shown to increase expression of pluripotency markers (Torres and Watt 2008). Similarly, inhibition of Protein Kinase C (PKC) activity in mESCs, which in turn restrains NF-κB signaling, allows keeping mESCs in a ground pluripotency state, thus facilitating the generation of induced pluripotent stem cells (iPSCs) (Dutta et al. 2011).

However, whether NF-κB components affect mESCs function and heterogeneity specifically
through regulation of NF-κB signaling or by additional mechanisms remains unknown.

We previously have identified an alternative nuclear function of IκBα involving the interaction
with different transcriptional regulators, histone deacetylases (HDACs), core elements of the
Polycomb Repressor Complex 2 (PRC2) as well as histones H2A and H4 (Aguilera et al. 2004;
Mulero et al. 2013; Marruecos et al. 2021). This nuclear IκBα function is crucial to efficiently
silence sets of developmental genes and ensure a proper cell fate specification of somatic
stem cells in the epidermis, intestine and hematopoietic stem cells (Mulero et al. 2013;
Marruecos et al. 2020, 2021; Thambyrajah et al. 2022). We have now investigated the role of
IκBα in the pluripotency of mESCs by using deletion mutants and characterizing their capacity
to self-renew and differentiate. We found that lack of IκBα forces a ground chromatin state
which involves changes in H3K9me3, H3K4me3 and H3K27me3 levels, and it is coupled with
a global DNA hypomethylation in S/L conditions, resembling the ground state of pluripotency.
This stabilization of the ground state imposes a block in the pluripotency exit and a general
impairment of differentiation. Expression of Separation-of-function (SOF) IκBα mutants that
either bind to histones or to NF-κB demonstrate that chromatin-associated IκBα controls
heterogeneity in mESCs by an epigenetic rewiring independent of NF-κB function.

RESULTS

IκBα is present in the nucleus of pluripotent embryonic stem cells but not after
differentiation

NF-κB-dependent IκBα function takes place mainly in the cytoplasm, whereas chromatin-
associated IκBα function occurs in the nucleus (Zhang, Lenardo, and Baltimore 2017). We first
investigated IκBα localization in mESCs by immunofluorescence (IF) and western blot (WB).
We observed an accumulation of total IκBα in the nucleus and chromatin fractions in
pluripotent mESCs (Figures 1A upper panel and 1B). We previously found that nuclear IκBα
mainly corresponds to phosphorylated Ser32/36 IκBα (p-IκBα), which is preserved from
degradation because of K21 SUMOylation (Mulero et al. 2013). In line with this, our IF analysis
showed p-IκBα accumulation in the nucleus of mESCs (Figure 1A bottom panel). In the skin
and intestine, nuclear IκBα and p-IκBα are restricted to the stem cell compartments and lost
upon differentiation (Mulero et al. 2013; Marruecos et al. 2020). To explore whether a similar
distribution happens upon pluripotency exit, we next investigated p-IκBα dynamics in
embryoid body (EB) differentiating conditions (Figure 1C upper panel). At 216 hours of
culture, most of the cells in EBs were differentiated as detected by the absence of NANOG,
and only few cells maintained NANOG expression (Figure 1C bottom panel and 1D). In these
conditions, nuclear p-IκBα was restricted to pluripotent cells, co-localizing with NANOG
(Figures 1C bottom panel and 1D). These results indicate that p-IκBα is accumulated in the
nucleus of pluripotent cells, which is lost upon pluripotency exit. Supporting this observation,
RNAseq analysis of mESCs (0h) and early differentiation stages of EBs (48h and 96h), showed
that pluripotent cells express higher levels of IκBα gene (*Nfkbia*), which decreases upon
differentiation (Figure 1E). Importantly, *Nfkbia* expression follows similar expression
dynamics as pluripotency genes (*Dppa4, Zfp42, Prdm14, Klf4, Sox2 and Nanog*), whereas NF-
κB members (*Rela, Relb* and *Rel*) and other IκB genes (*Nfkbib, Nfkbiz* and *Nfkbie*) showed
expression patterns similar to differentiation genes (Figure 1E).

Taken together, these data indicate that p-IκBα protein is specifically present in the nucleus
in the pluripotency state of mESCs when *Nfkbia* gene expression is highest.

IκBα is required for pluripotency silencing and the *in vitro* and *in vivo* differentiation of
mESCs

To investigate the role of IκBα in pluripotency, we generated IκBα$^{-/-}$ mESC clones by CRISPR-
Cas9 with guide RNAs (gRNAs) targeting the first exon of *Nfkbia* (Figure S1A and see Methods
section). Initial characterization of three independent IκBα$^{-/-}$ and IκBα$^{+/+}$ mESCs clones
demonstrated that lack of IκBα was not affecting expression of the pluripotency marker SSEA-
1 (Figure S1B). Cell cycle analysis showed increased number of proliferating cells in IκBα$^{-/-}$
compared to IκBα$^{+/+}$ mESCs (G$_2$ phase: 15.5%±3.5 in IκBα$^{+/+}$/20.1%±3.5 in IκBα$^{-/-}$; M phase:
2.5%±0.2 in IκBα$^{+/+}$/4.1%±0.7 in IκBα$^{-/-}$) (Figure S1C). Next, we assessed the ability of IκBα$^{-/-}$

and IκBα[+/+] mESCs to differentiate into embryoid bodies (EBs) (Figure S2A) and detected
higher numbers of pluripotent cells (NANOG[+]/OCT3/4[+]) after 120h of differentiation in IκBα[-]
[/-] compared to IκBα[+/+] EBs (Figure S2B). The majority of cells in IκBα[-/-] EBs maintained a
pluripotency state after 216h, when IκBα[+/+] EBs had only a few small clusters of pluripotent
cells remaining (Figure 2A). In addition to the presence of OCT3/4 and NANOG, the higher
pluripotency status was confirmed by the presence of Alkaline Phosphatase (AP) assay
(Figures 2B and S2C), SSEA-1 (Figures 2C and S2E) as well as the expression of *Pou5f1* (*Oct4*)*,*
Nanog, Zfp42 (Rex1) and Sox2 (Figure S2F). We also observed smaller sizes in EBs from IκBα[-/-]
cells (Figure S2D), a feature previously related to lower differentiation potential (Tian et al.
2019). To further study the requirement of IκBα in mESC differentiation *in vivo*, we established
teratomas by subcutaneous injection of IκBα[+/+] and IκBα[-/-] mESCs (Figure S2G) in NSG
(NOD.Cg-Prkdc[scid] Il2rg[tm1Wjl]/SzJ) mice. IκBα[-/-] teratomas show a decreased differentiation
potential, as indicated by the presence of less differentiated structures (Figure S2H) and the
higher number of OCT4[+] cells (Figures 2D and 2E).

The lack of differentiation capability of IκBα[-/-] mESCs was confirmed at the transcriptomic
level by RNAseq of mESCs and EBs at 48h and 96h in differentiation medium (Figure 2F and
supplementary table 1). Moreover, IκBα[-/-] 96h EBs failed to properly silence naïve
pluripotency genes (Figures 2G, 2K, S3B-D) while they cannot activate differentiation
programs of germ layer (endoderm, mesoderm and ectoderm) specification (Figures 2H-K).
These results support the pluripotency fate retention in IκBα[-/-] mESCs upon differentiation
conditions. Notably, we did not detect major changes in the expression levels of NF-κB
signaling pathway genes neither at IκBα[-/-] mESCs nor in IκBα[-/-] EBs 48h and IκBα[-/-] EBs 96h
(Figure S1D), suggesting that the phenotype imposed by IκBα deficiency is not due to aberrant
NF-κB signaling. This could be due to the fact that there is no NF-κB activity in mESCs that
need to be repressed by IκBα, or the NF-κB-related IκBα function might be compensated by
other IκBs, as previously demonstrated (Hayden and Ghosh 2008).

Altogether, our results demonstrate that the lack of IκBα in mESCs imposes a blockade at the
pluripotency state which inhibits differentiation in both *in vivo* and *in vitro* conditions without
affecting canonical NF-κB signaling.

Absence of IκBα stabilizes the ground state pluripotency through epigenetic rewiring in
mESCs

In vitro cell states for mESCs are highly heterogeneous mimicking different embryonic states
and range from a highly pluripotent (ground) state to primed for differentiation (Graf and
Stadtfeld 2008; Martinez Arias and Brickman 2011). To investigate the role of IκBα in
pluripotency, we performed a Gene Set Enrichment Analysis (GSEA) over IκBα[+/+] and IκBα[-/-]
mESCs comparing gene signatures for ground and primed states of mouse ESC
pluripotency(Hu et al. 2022). We found that IκBα[-/-] mESCs transcriptome is significantly
enriched for genes specifically expressed in the ground state, while negatively correlating with

the primed state (Figure 3A). These results suggested that IκBα is required for the transition
from ground to primed states, which is characterized by an important epigenetic rewiring that
involves histone post-translational modifications and DNA methylation. To further investigate
the state of IκBα$^{-/-}$ mESCs, we performed genome-wide profiling of H3K4me3, H3K27me3 and
H3K9me3 by Chromatin Immunoprecipitation followed by sequencing (ChIP-seq). In general,
IκBα$^{-/-}$ mESCs showed a more permissive chromatin status with a predominant enrichment of
H3K4me3, loss of H3K9me3 and redistribution of H3K27me3 (Figures 3B, 3C, 3E, 3G, 3I and
Figures S3E, S3F, S3H, S3I, and supplementary table 2). Gain of H3K4me3 was found in genes
associated to pluripotency such as *Klf2, Zfp42 (Rex1), Sall4* and *Esrrb* (Figures 3C-E and S3E),
and it directly correlates with their increased expression in IκBα$^{-/-}$ mESCs (Figure 3F). Analysis
of the H3K27me3 mark in IκBα$^{-/-}$ mESCs showed a slight increase of the number of regions
with higher H3K27me3 (1070 regions) (Figures 3B, 3G and S3F). Notably, the subset of genes
with loss of the H3K27me3 mark (612 regions) were associated with mechanisms of
pluripotency (Figure 3H), including those genes associated to the ground state of
pluripotency, such as *Tbx3, Klf4* and *Gbx2*, whose transcription is upregulated in IκBα$^{-/-}$ mESCs
(Figures 3I and S3G).

Analysis of the H3K9me3 mark showed a significant reduction in the detection of regions
enriched for this mark in IκBα$^{-/-}$ mESCs (87 regions gained and 450 regions lost) (Figures 3B,
S3H, S3I). Among regions with decreased levels of H3K9me3, we found that they mainly
correspond to DNA-methylated and imprinted regions (Figures 3J and S3J), which gain
H3K4me3 decoration and gene expression (Figures 3J and S3K). H3K9me3 is associated with
heterochromatin and DNA methylation (Allis and Jenuwein 2016). We evaluated the DNA
methylation levels in IκBα$^{+/+}$ and IκBα$^{-/-}$ mESCs by immunofluorescence. We found a strong
decrease in the 5-methylCytosine (5mC) signal in the absence of IκBα (Figure 3K), suggesting
a hypomethylation phenotype. To further investigate the specific changes in DNA
methylation, we performed a DNA methylation array (Zhou et al. 2022) in IκBα$^{+/+}$ and IκBα$^{-/-}$
mESCs and 96h EBs. IκBα$^{-/-}$ mESCs showed a global DNA hypomethylation profile which
persists upon induction of differentiation at 96h compared to the IκBα$^{+/+}$ counterparts
(Figures 3L, S4A-C). Analysis of differentially methylated regions (DMPs; Δβ < 0.3) identified
9940 hypomethylated regions specifically in IκBα$^{-/-}$ mESCs, 1497 in IκBα$^{-/-}$ 96h EBs and 518
common regions among both timepoints (Figure 3M and supplementary table 3). Specifically,
DNA-hypomethylated positions exclusively found in IκBα$^{-/-}$ mESCs correspond to regions of
heterochromatin typically decorated by H3K9me2 (Figure S4D). Motif enrichment analysis of
IκBα$^{-/-}$ 96h-EBs hypomethylated regions identified target enhancers for pluripotency-related
factors such as PRDM14, ESRRB, NANOG (Figure S4E). Moreover, positions that were
commonly hypomethylated in mESCs and 96h EBs IκBα$^{-/-}$ corresponded to imprinted regions
enriched in H3K9me3 (also confirmed by H3K9me3 ChIP-seq in mESCs (Figure 3J, Figure S3H-
J)) and generally bound by the SETDB1/TRIM28 complex (Figure S4F).

These results suggest that IκBα in mESCs regulates different chromatin-modifying activities,
including histones PTMs and DNA methylation, which specifically affects pluripotency genes.

Moreover, our data supports previous studies indicating that long-term stabilization of the
ground state of pluripotency limits the developmental potential of mESCs (Choi et al. 2017).

218 Chromatin function of IκBα induces changes in ground state pluripotency and pluripotency
219 exit in an NF-κB independent manner

Uncovering the actual impact of chromatin-related IκBα functions is challenging due to the
predominant role of this protein on NF-κB regulation. Recent work from our group has
identified the residues in the IκBα protein which are responsible to specifically interact with
NF-κB subunits or with histones, and we have generated separation-of-function (SOF) IκBα
mutants which retain either NF-κB or chromatin-dependent functions (Álvarez-Villanueva et
al. 2023). Taking advantage of IκBα$^{-/-}$ mESCs, we generated different doxycycline-inducible
IκBα (i-IκBα) expressing cell lines containing either a wildtype IκBα protein (i-IκBα^{WT}) or one
of the SOF IκBα mutants: the one that interacts with NF-κB and it is deficient for H2A/H4
binding (IκBα$^{ΔH2A/H4}$) or the one that interacts with H2A/H4 but not with NF-κB (IκBα$^{ΔNF-κB}$)
(Figures 4A, S5A). By pull down experiments, we confirmed that IκBα$^{ΔNF-κB}$ was able to interact
with histone H2A but not with the NF-κB member p50 (Figures S5C and S5D), although in our
conditions it can still interact with p65 (Figure S5D). On the other hand, IκBα$^{ΔH2A/H4}$ can
interact with the classical NF-κB heterodimer p50-p65, but it cannot bind H2A (Figures S5C
and S5D). As expected, IκBα^{WT} is able to bind to both H2A and NF-κB. To address how IκBα is
mediating the stabilization of the ground state in mESCs, we induced expression of i-IκBα^{WT},
i-IκBα$^{ΔNF-κB}$ and i-IκBα$^{ΔH2A/H4}$ in IκBα$^{-/-}$ mESCs by overnight (16h) treatment of doxycycline.
Temporary induction of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ was enough to decrease expression of
pluripotency genes *Tbx3, Gbx2, Klf4, Zfp42* and *Nanog*, and to induce differentiation markers
(Figures 4B and 4C). Notably, induction of i-IκBα$^{ΔH2A/H4}$ represses most of the differentiation
genes and it causes upregulation of the naïve pluripotency genes *Zfp42* and *Klf2* (Figures 4B
and 4C). In addition, the ground state-associated DNA hypomethylation was reverted upon
expression of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ but not upon i-IκBα$^{ΔH2A/H4}$ induction (Figure 4D). These
results demonstrate that the chromatin-dependent IκBα function, but not its NF-κB-related
activity, is necessary and sufficient to revert the ground state of pluripotency with higher DNA
methylation levels and lower pluripotency gene expression.

In order to address which IκBα activities were required for full mESC differentiation, we
induced EB differentiation in i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and i-IκBα$^{ΔH2A/H4}$ lines (Figure S5E).
Induction of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ in EBs restored the differentiation potential in IκBα$^{-/-}$
mESCs (Figures 4E-G and S5E-G). In both conditions, we observed that the number of
pluripotent-like cells (NANOG$^+$ and OCT3/4$^+$) remaining at 216h EBs was similar to the ones
observed at IκBα$^{+/+}$ EBs (Figures 4E, S5F and 2A), and only small and few patches of
undifferentiated cells are observed. On the other hand, induction of i-IκBα$^{ΔH2A/H4}$ did not
restore the differentiation potential of IκBα$^{-/-}$ and most of the EB cells were still expressing
high levels of NANOG and OCT3/4 at 216h. To further define the differentiation potential of
i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and i-IκBα$^{ΔH2A/H4}$ EBs after 216h, we performed AP staining as well as

analysis of SSEA-1. We found that expression of i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ reduced the number of
undifferentiated AP$^+$ colonies obtained from the EBs (Figure 4F). In agreement with this
observation, the number and proportion of cells lacking expression of the pluripotency
marker SSEA-1 in the EBs was partially restored upon induction of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ but
not i-IκBα$^{ΔH2A/H4}$ (Figures 4G and S5G).

Taken together, these results indicate that chromatin-dependent IκBα function controls the
balance between ground and primed-to-differentiate states of pluripotency in mESCs in an
NF-κB independent manner, and it is required for cells to exit pluripotency and activate the
differentiation programs.

DISCUSSION

We here identified a new and unexpected function for the IκBα protein in the regulation of
pluripotency and differentiation. Using mESCs to investigate the role of IκBα in differentiation,
we discovered that the absence of IκBα severely impairs the exit of pluripotency.
Furthermore, IκBα$^{-/-}$ mESCs acquire a ground state of pluripotency, indicating the role of IκBα
as a regulator of the transition between ground and primed states. While IκBα is primarily
known as an inhibitor of the NF-κB signaling pathway, our lab has previously provided strong
evidence of its alternative role by binding to histones and other chromatin elements. We have
demonstrated that chromatin-bound IκBα has a crucial role in regulating skin and intestinal
stem cells (Mulero et al. 2013; Marruecos et al. 2020), as well as affecting the regenerative
capacity of these tissues and the neoplastic transformation. Differentiating this chromatin
function of IκBα from its NF-κB inhibitory function has been challenging, making it difficult to
determine its significance. To address this, we have engineered separation-of-function (SOF)
mutants of IκBα (Álvarez-Villanueva et al. 2023) by identifying specific residues that are
essential to bind to NF-κB elements or histones. By reintroducing these mutants into IκBα-
deficient mESCs, we have shown that only the histone-binding proficient IκBα mutant
(IκBα$^{ΔNF-κB}$) with deficient NF-κB binding can restore the differentiation capacity of mESCs.
Similarly, we found that the same IκBα$^{ΔNF-κB}$ mutant is able to restore differentiation capacity
in intestinal cells (Álvarez-Villanueva et al. 2023). The role of NF-κB in mESC differentiation
and pluripotency has been previously explored, and while several reports suggest that NF-κB
activity correlates with differentiation and loss of pluripotency (Torres and Watt 2008), others
claim that NF-κB inhibition correlates with a decrease in pluripotency gene expression
(Armstrong et al. 2006). In any case, since modification of canonical NF-κB activity directly
impacts on the levels IκBα expression, the previous NF-kB associated phenotypes should be
reevaluated considering this chromatin role of IκBα.

Our results support the observation that expression of NF-κB elements is low in
undifferentiated mESCs and increases with differentiation (Figure 1E). Moreover, the
expression of IκBα$^{ΔH2A/H4}$, which retains the inhibitory function of NF-κB, cannot reconstitute
differentiation capacity in cells that lack IκBα (Figures 4 and S5) indicating that the observed
IκBα function in controlling pluripotency exit is not due to NF-κB inhibition. In fact, there is
evidence that the chromatin function of IκBα is an ancestral function preceding its role as NF-
κB inhibitor. In Caenorhabditis elegans (*C.elegans*), which lacks NF-κB orthologs but possesses
two IκBα homolog genes, the absence of IκBα orthologs has severe multi-organ
differentiation defects (Brena et al. 2020). In mice, IκBα-deficiency results in perinatal death
without major developmental defects in the early stages of embryogenesis, which suggest
that other mechanisms may compensate for the lack of IκBα *in vivo*. Alternatively, we
speculate that in vitro maintenance of pluripotency is differentially regulated compared to
blastocyst development.

We found that IκBα⁻/⁻ mESCs exhibit an epigenetic and transcriptomic profile that is
compatible with a ground state of pluripotency, strongly suggesting that temporary inhibition
of IκBα in mESCs or iPSCs could be used to modulate ground versus primed pluripotency. Our
results show that IκBα-deficient mESCs exhibit general DNA hypomethylation, loss of
heterochromatin mark H3K9me3 in imprinted genes and a general gain of H3K4me3
activation mark (Figure 3). These results suggest IκBα to be a crucial member of a complex
which indirectly regulates different histones PTMs and it allows a permissive chromatin status
to successfully allow the exit from pluripotency. Nevertheless, the chromatin target regions
of IκBα as well as its chromatin partners need to be identified.

Additionally, H3K27me3 is modified in several genomic regions, although there is no net gain
or loss of this mark. We have previously demonstrated that IκBα interacts with various
elements of the Polycomb Repressor Complex 2 (PRC2) in epidermal and intestinal stem cells
(Mulero et al. 2013; Marruecos et al. 2020), an interaction that is conserved in *Drosophila*
(Mulero et al. 2013) and *C.Elegans* (Brena et al. 2020). While we have not observed this
physical interaction in mESCs, it is likely that some of these changes result from IκBα
modifying the activity of the PRC2 complex.

Overall, our work demonstrates that a chromatin activity of IκBα controls pluripotency states
in mESCs through an epigenetic rewiring which permits pluripotency exit in an NF-κB-
independent manner.

METHODS

Cell lines and culture mediums

HEK293T and NIH/3T3 cell lines were obtained from American Type Culture Collection (ATCC;
HEK293T: Cat #CRL-3216; NIH/3T3: Cat #CRL-1658). They were cultured in Dulbecco's
modified Eagle's medium (DMEM; Invitrogen Cat #11965092) supplemented with 10% fetal
bovine serum (FBS; Biological Industries Cat #04-001-1A), 4.5 g/L glucose (Life Technologies
Cat #A2494001), 2 mM L-glutamine (Biological Industries Cat #03-020-1A), 56 U/mL penicillin
and 56 µg/mL streptomycin (Biological Industries Cat #03-031-1B).

mESC line ES-E14TG2a (ATCC; Cat #CRL-1821) was cultured on plastic dishes precoated with
0.1% (w/v) Gelatin (Sigma; G2500-100G) with DMEM-ES medium. DMEM-ES is composed by
DMEM basal medium (Sigma-Aldrich; Cat #D5796) supplemented with 15% FBS (ESC-
qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat #35050061), 1X NEAA (Gibco;
Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070), 103U/mL Leukemia
Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-mercaptoethanol (Gibco;
Cat#31350010).

Medium was changed every day, and cells were splitted every other day using TrypLE Express
(Gibco; Cat #12605010) for harvesting. Cells were maintained in a 5% CO2 incubator at 37°C.

Embryoid bodies differentiation from mESCs

Embryoid bodies differentiation was established as described in (Sroczynska et al. 2009).
Briefly, mESCs were thawed and passed twice in DMEM-ES before inducing differentiation.
After the second passage, and once mESCs are 80% confluent, cells are trypsinized with TrypLE
Express. One well of 6-well plate was splitted in an entire 6-well plate in IMDM-ES medium
for 48h. IMDM-ES medium is composed of Iscove's Modified Dulbecco's Medium (IMDM)
supplemented with 20% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat
#35050061), 1X NEAA (Gibco; Cat#11140050), 1mM Sodium Pyruvate (Gibco; Cat#11360070),
10ng/mL Leukemia Inhibitory Factor (LIF) (Millipore, Cat #ESG1107) and 0.125mM 2-
mercaptoethanol (Gibco; Cat#31350010).

For embryoid body induction, mESCs were harvested by trypsinization with TrypLE, and they
were rinsed twice with 1X PBS to remove any traces of LIF, since this one can inhibit a proper
EB differentiation. Cells were very well disaggregated into single-cells, and 1.2 x 104 cells/mL
embryoid body differentiation (EBdiff) medium were resuspended to a total volume of 25mL
of EBdiff medium. Embryoid body differentiation medium is composed by IMDM
supplemented with 15% FBS (ESC-qualified; Gibco Cat #26140079), 1X Glutamax (Gibco; Cat
#35050061), 50 µg/mL ascorbic acid (Sigma; Cat #A-4544), 180 µg/mL Transferrin (Roche; Cat
#10652202001) and 0.45 mM alpha-monothioglycerol (MTG) (Sigma; Cat #M6145). EBs were
formed in suspension for 5 days (120h). At day 5, in order to elongate the differentiation up

to 216h (day 9), EBdiff medium was refreshed by harvesting EBs with 10mL serological
pipette, centrifuge at 200 g for 3 min and EBs were resuspended in fresh EBdiff medium for 4
days longer.

CRISPR/Cas9 gene editing in mESCs

The two guide RNAs (gRNAs) targeting the Nfkbia locus were designed using the CRISPR design
tool from MIT (http://crispr.mit.edu.sare.upf.edu). The best 2 gRNAs (based on on-target and
off-target scores) that were targeting the exon 1 of Nfkbia (IκBα gene) were selected. After
annealing, one of the gRNAs was cloned into SpCas9(BB)-2A-GFP (px458) plasmid (Addgene;
Cat #48138), and the other gRNA was cloned into the px330-mCherry plasmid (Addgene; Cat
#98750). mESCs were transfected with the two plasmids (px458-gRNA1 and px330-mCherry-
gRNA2) at the same time, following the DECKO (Double Excision CRISPR Knockout) approach
(Aparicio-Prat et al. 2015). Briefly, 3×10^5 cells were seeded a day before. At day of
transfection, cells were washed once with 1X DPBS (Gibco; Cat #14190144), and Opti-MEM
medium (Gibco; Cat #31985070) was added. 1.25 µg of each plasmid (2.5 µg of total DNA)
was incubated with 10 µL of Lipofectamine 2000 (Invitrogen; Cat #11668019) in 240 µL of
Opti-MEM for 20 min. OptiMEM::Lipofectamine::DNA mixture was added into mESCs, and
cells were incubated for 5 hours with the mixture. After that time, cells were washed with
PBS, and DMEM-ES was added into the well. 48h after transfection, GFP+ mCherry+ single
cells were sorted by FACS using BD FACSAria II Cell Sorter (BD Bioscience). Mutant clones were
screened through PCR and western blotting to identify single clones with no IκBα protein
expression. Non-targeting scrambled gRNAs were cloned into px458 and px330-mCherry
plasmids to generate control clones. Three independent clones from each genotype (wild-
type and mutant) were selected for further experiments. gRNAs sequences are included in
supplementary table 4.

Teratoma formation assay

IκBα+/+ and IκBα-/- mESCs grown in DMEM-ES (5 × 105) were injected intramuscularly into
severe combined immunodeficient mice (NSG). In order to favor 3D aggregation of cells and
teratoma formation, cells were resuspended in Matrigel Matrix (BD; Cat #356234) prior to
injection. Six weeks later, mice with tumors were euthanized, and tumors were fixed in
formaldehyde, embedded in paraffin, sectioned and stained with hematoxylin and eosin for
histological analysis.

Immunofluorescence staining

389 For immunostaining of mESCs, cells were seeded on 0.1% gelatin-coated coverslips on 6-well
390 plates; for EB immunofluorescence, they were collected directly from the plate. Samples were
391 washed twice with 1X PBS, and they were fixed at RT for 10 minutes with 4% PFA (Electron
392 Microscopy Sciences; Cat #15713S). After fixation, samples were washed twice with 1X Tris-
393 buffered saline (TBS; 50 mM Tris-Cl, pH 8) and permeabilized and blocked with 1X TBS

supplemented with 1% Triton-X100 (MERCK; Cat #9036-19-5) and 6% FBS (Biological Industries; Cat #04-001-1A) for 2h at 4ºC. Samples were washed twice with 1X TBS supplemented with 6% FBS. Primary antibody incubation was performed in 1X TBS plus 6% FBS and 0.5% Triton-X100 for overnight at 4ºC. The following antibodies were used: phospho (Ser32/36) IκBα (1:250; Cell Signaling; Cat #9246), OCT3/4 (1:250; Santa Cruz; Cat #sc-5279), NANOG (1:250; Novus Biologicals; Cat #NB100-588), BRACHYURY (1:200; Invitrogen; Cat #PA5-78422). Samples were then washed 4 times with 1X TBS, 5 min each washing. Secondary antibody incubation was performed in TBS 1X plus 1% BSA (Sigma-Aldrich; Cat #9048-46-8) for 4 h at RT. The following antibodies were used: Alexa Fluor 488 donkey anti-mouse antibody (1:1000; Invitrogen; Cat #A-21202), Alexa Fluor 647 donkey anti-mouse antibody (1:1000; Invitrogen; Cat #A-31571), Alexa fluor 594 goat anti-mouse antibody (1:1000; Invitrogen; Cat #A-11032), Alexa Fluor 488 donkey anti-rabbit antibody (1:1000; Invitrogen; Cat #A-21206), Alexa Fluor 594 donkey anti-rabbit antibody (1:1000; Life Technologies; Cat #A-21207). Samples were washed three times with 1X TBS for 5 min at RT each rinse. Samples were mounted using DAPI Fluoromount-G (Southern Biotech; Cat #0100-20).

Microscopy and image acquisition

Fluorescence images were acquired using Confocal Leica TCS SP5 (Leica Microsystems), and Leica application software LAS AF (Leica Microsystems) was used to visualize the images. Images of teratoma haematoxylin and eosin and immunohistochemistry stainings were acquired using BX61 Olympus Microscope (Olympus), and PRECiV™ 2D Image and Measurement Software (Olympus) was used to visualize the images. The Image J (version 1.15) was used for further analysis (Schneider, Rasband, and Eliceiri 2012).

Cell cycle analysis

mESCs were detached from the plates using TrypLE Express, and they were fixed and permeabilized with a FIX & PERM Cell permeabilization kit (Invitrogen, Cat #GAS004) following manufacturer's instructions. Cells were stained with Alexa Fluor 647 Mouse anti-Ki-67 antibody (BD Biosciences, Cat #561126) (1:100), and DAPI (5 µg/ml) (Biotium, Cat #BT-40043). Staining with DAPI was done for 30 min, prior to flow cytometry analysis using a BD LSR II Flow Cytometer (BD Bioscience).

Flow cytometry analysis

Cells were purified based on FACS using BD FACSAria II Cell Sorter (BD Bioscience). Cells were sorted at around 3,500 events/second, and at maximum flow rate of 4 and with a 85 µm nozzle. BD LSRFortessa Cell Analyzer or BD LSR II Flow Cytometer (BD Bioscience) were used for Flow Cytometry Analysis experiments. 5 µg/ml of DAPI (Biotium; Cat #BT-40043) was used as a viability dye. Flow cytometry analysis was performed

RNA isolation, cDNA synthesis and quantitative RT-PCR

Total RNA isolation from cells was performed using the RNeasy Plus Mini Kit (Qiagen; Cat
#74136) or RNeasy Micro Kit (Qiagen; Cat #74004) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
2 µg of total RNA was retro-transcribed using Transcriptor First Strand cDNA Synthesis Kit
(Roche; Cat #04897030001) following the manufacturer's instructions.

Quantitative RT-PCR was performed in triplicates for each sample, and SYBR Green I Master
Kit (Roche; Cat #04887352001) was used to carry out the reaction. qRT-PCR reaction was done
using the LightCycler 480 system (Roche). Gene expression levels were calculated as $2^{-\Delta CT}$
normalized with the average CT of the housekeeping gene Tbp or Gapdh. Primer sequences
are found in supplementary table 4.

ChIP-seq experiments

mESCs were cross-linked by incubating them in a cross-link solution (50 mM HEPES pH 8.0, 10
mM NaCl, 1 mM EDTA, 0.5 mM EGTA (Egtazic acid)) supplemented with 0.5% formaldehyde
(Sigma-Aldrich; Cat #252549). Samples were Incubated for 10 min rocking at RT. Cross-linking
was stopped by adding 1.25 M Glycine, 10 mM Tris-HCl pH 8.0 into the sample and incubating
for 5 min rocking at RT. Cells were then lysed with 1 mL of lysis buffer (10 mM Tris-HCl pH 8.0,
0.25% Triton X-100, 10 mM EDTA, 0.5 mM EGTA, 10 mM Na-butyrate, 20 mM β-glycerol-
phosphate, 0.1 mM Naorthovanadate and cOmplete™ Protease Inhibitor Cocktail) for 30 min
at 4ºC. Samples were then resuspended in sonication buffer (10 mM Tris-HCl pH 8.0, 0.1 M
NaCl, 1 mM EDTA, 0.5 mM EGTA, 10 mM Na-butyrate, 20 mM β-glycerol-phosphate, 0.1 mM
Na Orthovanadate and 1% SDS), and they were sonicated using sonication beads (Diagenode;
Cat #C01020031) and the Bioruptor Pico Sonicator (Diagenode; Cat #B01060010) for 7 min,
to fragment DNA into 500 bp peaces. Fragmented DNA was concentrated using Vivaspin 20
columns (Sartorius; Cat #VS2032), and unspecific binding was blocked by incubating the
sample with 1% BSA, 1µg salmon sperm DNA, unspecific pre-immune IgG for 2h rocking. To
immunoprecipitate, samples were incubated with 5 µg of antibody at 4ºC overnight.
Chromatin-IgG complexes were pull down using protein A/G sepharose beads (GE Healthcare;
Cat #17-0618-01 and #17-0780-01) for 2h rocking at 4ºC. Cross-linked DNA-protein was
reverted by incubating samples at 65 ºC overnight. DNA was purified using GFX Kit (GE
Healthcare Life Sciences; Cat #28-9034-71), following the manufacturer's instructions.

Purified DNA concentration and integrity were determined using Agilent Bioanalyzer (Agilent
Technologies; Cat #G2939BA). Libraries were prepared using standard protocols. Chromatin
was sequenced using Illumina HiSeq platform (Illumina, Inc.) (50bp single-end reads). Samples
sequencing depth ranged between, per histone mark: (i) H3K4me3: 26M and 27M reads
(average 27M reads), (ii) H3K27me3: 62M and 72M reads (average 67M reads) and (iii)
H3K9me3: 71M and 89M reads (average 80M reads).

ChIP-seq data analysis

Quality control was performed on raw data with FASTQC tool. Raw reads were trimmed to
remove adapters presence with Trimgalore (v0.6.6) (Krueger et al., 2020). Default parameters
were used except for a minimum quality of 15 (Phred score) and an adapter removal
stringency of 3bp overlap. For H3K4me3 and H3K27me3 data, trimmed reads were aligned to
reference genome with Bowtie2 (v2.4.4) which was executed with default parameters
(Langmead & Salzberg, 2012). Required genome index was built with corresponding GRCm38
fasta file retrieved from Ensembl (http://ftp.ensembl.org/pub/release-102/). Multimapped
reads and those exhibiting MAPQ < 20 were removed. For H3K9me3 data, trimmed reads
were aligned to reference genome (GRCm38 release 102 from Ensembl) with ShortStack
(v3.8.5) (Axtell, 2013). Randomly placed multimappers were removed from the mapped
reads. For all three histone marks, duplicated reads were marked with SAMtools (v1.15)
(Danecek et al., 2021). Drosophila Melanogaster spike-ins, present in H3K27me3, were
discarded. NSC and RSC quality metrics were computed with PhantomPeakQualTools (v1.2)
(Landt et al., 2012). ENCODE blacklisted regions (mm10 v2) were removed prior to peak calling
except for H3K9me3 (Amemiya et al., 2019). BigWig files were individually generated using
deepTools (v3.5.1) bamCoverage with -ignoreDuplicates -binSize 10 -smoothLength 30 -
effectiveGenomeSize   2308125349   -normalizeUsing   RPGC   and   -extendReads
Fragment_Length options. (Ramírez et al., 2016). The effective genome size was
particularized to a read length of 50bp and directly retrieved from deepTools web site
(https://deeptools.readthedocs.io/en/develop/content/feature/effectiveGenomeSize.html).
Fragment_Length was retrieved from PhantomPeakQualTools results except for H3K9me3
where 150bp value was considered. For H3K9me3 and H3K27me3 data, peak calling was
conducted by means of epic2 (v0.0.52) with -effective-genome-fraction 0.8452 -fragment-
size *Fragment_Length* options and chromosome sizes only referring to canonical
chromosomes (Stovner & Sætrom, 2019). For H3K4me3 narrow profile, MACS2 (v2.2.7.1) was
used to identify peaks with -nomodel -extsize *Fragment_Length* -g 2308125349 options
(Zhang et al., 2008). For the three histone marks, peaks were called with adjusted p-values
(FDR) < 0.05. Peak annotation was performed with ChIPseeker R (v4.2.1) package (v1.34.1)
considering a TSS region range between 5000 bp upstream and 100 bp downstream (Yu et
al., 2015). Required TxDb object was generated from the Ensembl GRCm38 gtf file retrieved
(http://ftp.ensembl.org/pub/release-102/). A consensus peakset was obtained from the
three biological replicates per histone mark and condition (WT or KO). For this purpose,
findOverlapsOfPeaks function from ChIPpeakAnno R package (v3.32.0) was used with default
parameters (L. J. Zhu et al., 2010).

Differential binding analysis (DBA) was conducted with DiffBind (v3.8.4) per histone mark
(Stark & Brown, 2011). Default parameters were used except for the summits parameter
which was set to 150bp for H3K4me3 and 500bp for H3K27me3 and H3K9me3 in order to
consider, for testing, intervals of 300bp and 1000bp respectively. Summit values were
selected to have interval widths between the minimum and first quartile peak width values
for each histone mark. EdgeR was the statistical method used for all the three analyses.

Differentially bound regions (DBRs) between KO and WT samples were called with adjusted
p-values (FDR) < 0.05. Identified DBRs with any annotated gene were plotted with
EnrichedHeatmap R package (v1.18.1) (Gu et al., 2018). To summarize replicates per
condition, normalizeToMatrix function was used based on corresponding BigWig files in w0
mean mode in 50bp windows. Tracks visualization were obtained by means of Integrative
Genomics Browser (IGV) tool (Robinson et al., 2011).

RNA-seq experiments

Total RNA from three independent clones from mESCs, 48h EBs and 96h EBs was isolated
using the RNeasy Plus Mini Kit (Qiagen; Cat #74136) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
RNA integrity was addressed by agarose gel and Agilent Bioanalyzer (Agilent Technologies;
Cat #G2939BA). Libraries sequenced using Illumina HiSeq 2500 (Illumina, Inc.) (125bp paired-
end reads). Samples sequencing depth ranged between 35M and 52M reads (average 41M
reads) per sample.

RNA-seq data analysis

Quality control was performed on raw data with the FASTQC tool (v0.11.9). Raw reads were
trimmed to remove adapter presence with Trimgalore (v0.6.6) (Krueger et al., 2020). Default
parameters were used except for a minimum quality of 15 (Phred score) and an adapter
removal stringency of 3bp overlap. Trimmed reads were aligned to reference the genome
with STAR aligner tool (v2.7.8). STAR was executed with default parameters except for the
number of allowed mismatches which was set to 1. Required genome index was built with
corresponding GRCm38 gtf and fasta files retrieved from Ensembl
(http://ftp.ensembl.org/pub/release-102/). Obtained BAM files with uniquely mapped reads
were considered for further analysis. Raw gene expression was quantified using
featureCounts tool from subRead software (v2.0.1) with exon as feature (Liao et al., 2014).
The raw counts matrix was imported into the R Statistical Software environment (v4.2.1) for
downstream analysis. Raw expression matrix included 55,487 genes per 18 samples in total.
Experimental design considered three timepoints: mouse ESCs, EBs at 48h and EBs at 96h.
Each time point included 6 samples distributed in two conditions: 3xWT and 3xKO. Prior to
statistical analysis, those genes with less than 10 raw counts across the 6 samples under test
were removed. After pre-filtering, 21,843 genes (mESCs), 21,424 genes (EBs 48h) or 21,664
genes (EBs 96h) were available for testing. For visualization purposes, counts were normalized
by the variance-stabilizing transformation method as implemented in DESeq2 R package (Love
et al., 2014) (v1.38.3). Differential expression analysis (DEA) was conducted with DESeq2.
Each time point was independently analyzed. Fitted statistical model included sample
conditions as covariable with WT as the reference. Obtained log2 fold change values were
shrunken with apeglm shrinkage estimator R package (v1.20.0) (A. Zhu et al., 2019). Raw p-
values were adjusted for multiple testing using the Benjamini-Hochberg False Discovery Rate

(FDR) (Benjamini & Hochberg, 1995). Differentially Expressed Genes (DEGs) between KO and
WT samples were called with adjusted p-values (FDR) < 0.05 and absolute shrunken log2 Fold
change > 1. Data visualization was performed with the ggplot2 (v3.4.1).

Genome-wide DNA methylation samples preparation

DNA from frozen mESCs pellets was extracted using DNeasy Blood and Tissue Kit (Qiagen
GmbH, Hilden, Germany). Purified genomic DNA was quantified with Qubit (Invitrogen,
Carlsbad, CA, USA) according to manufacturer's instructions. Infinium Mouse Methylation
BeadChip (Illumina, Inc., San Diego, CA, USA) arrays were used to profile DNA methylation
genome wide. This platform allows over 285,000 methylation sites per sample to be
interrogated at single-nucleotide resolution. The samples were bisulfite converted using EZ
DNA Methylation-Gold™ Kit (Zymo Research, CA, USA) and were hybridized in the array
following the manufacturer's instructions.

DNA methylation data analysis

The DNA methylation profile of the studied samples was assessed using the Infinium Mouse
Methylation BeadChip Array (~285,000 methylation sites) as previously described (Garcia-
Prieto et al., 2022). Briefly, raw signal intensities were obtained with GenomeStudio Software
2011.1 (Illumina) and DNA methylation beta values were computed from raw IDAT files using
GenomeStudio default normalization with control probes and background subtraction.
Quality control steps to remove erratic probe signals were performed within the R statistical
environment (v4.0.3). We removed probes with detection p value > 0.01, genotyping probes
and manufacturing flagged (MFG) probes described in the Illumina manifest file
(https://support.illumina.com/downloads/infinium-mouse-methylation-manifest-file.html).
The differentially methylated probes (DMPs) between WT and KO samples were computed
separately for each time point (mESCs and EBs at 96h) by deriving a linear model with the
limma R package (v3.46.0). Each condition included 3 samples per time point. DMPs with
adjusted p-value (FDR) <0.05 and absolute mean methylation beta value difference between
conditions >0.3 were considered significant. DNA methylation analysis was performed using
the mm10 mouse genome reference build and the complete annotation was downloaded
from the annotated manifest file (http://zwdzwd.github.io/InfiniumAnnotation#mouse)
(Zhou et al., 2022).

Functional analysis

Overrepresentation analysis was applied over lists of selected genes derived from RNA-seq
data (DEGs) or from ChIP-seq data (differentially bound regions). The Gene Ontology
(Biological Process ontology, GO BP terms), KEGG PATHWAY and WikiPathways databases for
Mus Musculus (Ashburner et al., 2000; Kanehisa & Goto, 2000; Martens et al., 2021) were
interrogated by means of clusterProfiler R package (v4.6.2) (Wu et al., 2021). Corresponding
Entrez identifiers were used. Benjamini-Hochberg procedure was used to obtain adjusted p-

values. Obtained GO BP terms were simplified using the simplify function from clusterProfiler
with default parameters. Overrepresented terms or processes were called with adjusted p-
values (FDR) < 0.05.

Gene Set Enrichment Analysis (GSEA) was performed for mESCs against the three cell states
signatures defined by other authors (Hu et al., 2022). Briefly, each signature was composed
of 200 genes uniquely expressed in each state obtained from bulk RNA-seq performed per
state. For this purpose, the complete list of genes from mESCs samples (21,843 genes) was
ranked based on the shrunken log2 Fold Change obtained from DEA. GSEA was conducted
thru fgseaMultilevel function from fgsea R package (v1.24.0) (Korotkevich et al., 2021) with
default parameters. Enrichment plots were generated with the same package.

Additionally, the testEnrichment function from SeSAMe R package (v1.14.2) (Zhou et al.,
2018) was used for conducting functional analysis with default parameters over the list of
DMPs. Probe design, transcription factor binding site and histone modifications consensus
database sets, included in the same package, were interrogated.

Gene Set Variation Analysis (GSVA) R package (v1.46.0) with default parameters were used to
obtain Z-score values for a particular gene set (Hänzelmann et al., 2013).

DATA AVAILABILITY

All sequencing data is deposited at GEO under a superseries (accession number GSEXXXXX).
Individual datasets can be found at: <mark>GSEXXXX</mark> (RNA-seq data), <mark>GSEXXX</mark> (ChIP-seq data) and
<mark>GSEXXX</mark> (Methylation array data).

ACKNOWLEDGEMENTS

L.G.P has been a recipient of FI AGAUR fellowship (2019 FI-B 00151/2020 FI_B1 00130) from
Generalitat de Catalunya. D.A-V was funded by the FIS fellowship (FI20/00130) from Instituto
Carlos III. This work has been supported by Spanish Ministry of Science and Innovation
(PID2019-104695RB-I00, PLEC2021-007518, PDC2021-120817-I00), Generalitat de Catalunya
(2021SGR 39) and Departament de Salut (SLT002/16/00299) to AB; the Spanish Ministry of
Science and Innovation (PID2019-108322GB-100 and PID2022-142679NB-I00) to LDC; the
Spanish Ministry of Science and Innovation (PID2021-123383NB-I00.) and the Agencia de
Gestió d'Ajuts Universitaris i de Recerca (2021 SGR 01222) to BP. M.B. received funding from
the Ramón Areces Foundation. M.M. is a recipient of a grant from the Instituto Carlos III, grant
number CA22/00011 (cofunded by the European Social Fund Plus, ESF+ and by the European
Union).

AUTHOR CONTRIBUTION

L.G.P, L.E and A.B designed the study. L.G.P, D.A-V, M.B, A.I, J.G, J.B, D.A.E, C.A.G.P, C.B, V.R.C
and A.V performed experiments. L.G, M.M and M.B performed bioinformatics analysis. L.G
and A.B wrote the manuscript with input from all authors. C.B, A.V, A.V, P.M, L.D.C, B.P, M.E
and L.E provided input on experimental design. L.E and A.B provided resources and supervised
the research.

 FIGURE LEGENDS

Figure 1. p-IκBα specifically accumulates in the nucleus of pluripotent cells. A.
Immunofluorescence of mESCs stained for total IκBα and OCT3/4 (upper panel), and
phosphorylated Se 32/36 IκBα (p-IκBα) co-stained with NANOG (bottom panel). **B.** Western
blot analysis of cytoplasm (Cyt), nucleus (Nuc) and chromatin (Chr) from mESCs. Red arrow:
specific band for total IκBα. *: non-specific band. **C.** Schematic of EBs differentiation protocol
(upper panel). Representative image of immunofluorescence for NANOG and p-IκBα on 216h
EBs (lower panel). **D.** Percentage of cells displaying p-IκBα and NANOG protein expression at
216h EBs. Each dot represents the average quantification of % cells in five EBs in each
independent experiment. **E.** Heatmap representing the normalized and scaled expression of
genes related to pluripotency, differentiation and NF-κB pathways at mESCs, 48h embryoid
bodies (EBs) and 96h EBs. Three independent IκBα$^{+/+}$ clones are analyzed.

Figure 2. Lack of IκBα prevents pluripotency exit upon induction of *in vitro* and *in vivo*
differentiation. Representative image of immunofluorescence of 216h EBs stained for
NANOG and OCT3/4 from IκBα$^{+/+}$ (upper panel) and IκBα$^{-/-}$ (lower panel) cells. **B.**
Quantification of total number of alkaline phosphatase (AP) colonies after 216h EB
differentiation and additional culture of 96h in S/L medium. Dots represent independent
experiments from three different IκBα$^{+/+}$ or IκBα$^{-/-}$ clones **C.** Percentage of SSEA-1 cells at EBs
216h analyzed by flow cytometry. Each dot represents an independent clone. **D.**
Representative images of immunohistochemistry for OCT3/4 in IκBα$^{+/+}$ (left panel) or IκBα$^{-/-}$
(right panel) teratomas. Teratomas were derived from three independent clones of mESCs of
both genotypes. **E.** Percentage of OCT3/4$^+$ cells in teratomas. Positive cells were counted by
from five different microscope fields from each IκBα$^{+/+}$ and IκBα$^{-/-}$ clone. **F.** Principal
component analysis (PCA) based on normalized RNA-seq data from 3 different time points
(mESCs, 48h EBs and 96h EBs) of IκBα$^{+/+}$ and IκBα$^{-/-}$ cells. **G, H, I, J.** Z-score from normalized
counts retrieved from mESCs, 48h and 96h EBs IκBα$^{+/+}$ or IκBα$^{-/-}$ cells. Pluripotency score (G)
refers to the subset of genes from WikiPathways database ('Mechanisms associated with
pluripotency', WP1763) which are differentially expressed at 96h EBs. Genes from the
Endoderm (H), Mesoderm (I) and Ectoderm (J) scores are retrieved from corresponding Gene
Ontology BP formation terms (GO:0001706, GO:0001707 or GO:0001705, respectively). **K.**
Network of selected overrepresented biological pathways (adjusted p-val <0.05, FDR) from
WikiPathways database. Pathways are connected through their annotated differentially
expressed genes (DEGs) in IκBα$^{+/+}$ vs IκBα$^{-/-}$ comparison at 96h EBs. Pathway circle size is
proportional to the number of annotated genes. Gene node color represents shrunken Log$_2$
Fold-change obtained in differential expression analysis.

Figure 3. IκBα controls ground state pluripotency exit through epigenetic rewiring. A. GSEA
results from IκBα$^{-/-}$ vs IκBα$^{+/+}$ mESCs samples against ground and primed state pluripotency
signatures retrieved from Hu et al., 2022. **B.** Number of differentially binded regions (adjusted
p-val <0.05, FDR) for H3K9me3 (K9me3), H3K4me3 (K4me3) and H3K27me3 (K27me3) histone

marks in IκBα $^{-/-}$ compared to IκBα $^{+/+}$ mESCs. Gained or lost regions in IκBα $^{-/-}$ samples are distinguished (log$_2$ Fold-change >0 or <0, respectively) . **C.** Heatmap of differentially gained and lost H3K4me3 regions in IκBα $^{-/-}$ and IκBα $^{+/+}$ mESCs (adjusted p-val < 0.05, FDR). Only those regions with any associated gene are considered. Highlighted genes are annotated to 'Mechanisms associated to pluripotency' pathway from WikiPathways database (WP1763). Regions are sorted per their adjusted p-value obtained from differential binding analysis (DBA). Heatmaps are centered in the region center and extended 3Kb upstream and downstream. Fold Change values are retrieved from DBA. **D.** Functional overrepresentation analysis results of those genes with differential gain of H3K4me3 mark in IκBα $^{-/-}$ mESCs using WikiPathways database. **E.** Tracks showing H3K4me3 distribution for ground state pluripotency genes associated to differential regions in IκBα $^{-/-}$ compared to IκBα $^{+/+}$ mESCs. Shadowed regions highlight the differential levels of the H3K4me3. **F.** Expression analysis (qPCR) of those genes shown in previous panel. **G.** Heatmap of differentially gained and lost H3K27me3 regions in IκBα $^{-/-}$ and IκBα $^{+/+}$ mESCs (adjusted p-val < 0.05, FDR). Only those regions with any associated gene are considered. Highlighted genes are annotated to 'Mechanisms associated to pluripotency' pathway from WikiPathways database (WP1763). Regions are sorted per their adjusted p-value obtained from differential binding analysis (DBA). Heatmaps are centered in the region center and extended +/- 3kb. Fold Changes values are retrieved from DBA. **H.** Functional overrepresentation analysis results of those genes with differential lost H3K27me3 mark in IκBα $^{-/-}$ mESCs against WikiPathways database. **I.** Tracks showing H3K27me3 distribution for ground state pluripotency genes associated with differential regions in IκBα $^{-/-}$ mESCs compared to IκBα $^{+/+}$ mESCs. Shadowed regions highlight the differential levels of the H3K27me3. **J.** Tracks showing H3K9me3 and H3K4me3 distribution for imprinted genes associated to differential regions in IκBα $^{-/-}$ mESCs compared to IκBα $^{+/+}$ mESCs. Shadowed regions highlight the differential levels of the H3K9me3. **K.** Immunofluorescence of DNA methylation (5mC) in IκBα $^{+/+}$ mESCs (upper) and IκBα $^{-/-}$ (lower) mESCs. **N.** DNA methylation profile in IκBα $^{+/+}$ and IκBα $^{-/-}$ mESCs showing the distribution density of all 261,220 CpGs under test from mean ß-values per condition. **M.** Venn diagram representing differentially DNA hypomethylated regions in either IκBα $^{-/-}$ mESCs or 96h EBs.

Figure 4. Chromatin-associated IκBα controls pluripotency exit in an NF-κB-independent manner. A. Schematic of chromatin-specific/NF-κB-deficient (IκBα$^{\Delta NF-κB}$) or NF-κB-specific/chromatin-deficient (IκBα$^{\Delta H2A/H4}$) IκBα mutant functions. **B.** Expression analysis of pluripotency markers after inducing IκBα^{WT}, IκBα$^{\Delta NF-κB}$ and IκBα$^{\Delta H2A/H4}$ in IκBα$^{-/-}$ mESCs. Each dot represents an independent experiment. **C.** Expression analysis of differentiation markers after inducing IκBα^{WT}, IκBα$^{\Delta NF-κB}$ and IκBα$^{\Delta H2A/H4}$. Each dot represents an independent experiment. **D.** Immunofluorescence of DNA methylation (measured by 5mC) after re-expressing IκBα^{WT}, IκBα$^{\Delta NF-κB}$ and IκBα$^{\Delta H2A/H4}$ **E.** Immunofluorescence for NANOG and OCT3/4 at EBs 216h after induction of IκBα^{WT}, IκBα$^{\Delta NF-κB}$ and IκBα$^{\Delta H2A/H4}$. **F.** Alkaline phosphatase staining (upper) and quantification (lower) of reconstitution experiments. **G.** Flow cytometry

analysis of the pluripotency surface marker SSEA-1 at 216h EBs after IκBα^{WT}, IκBα$^{ΔNF-κB}$ and

IκBα$^{ΔH2A/H4}$ induction.

Figure S1. Depletion of IκBα does not compromise neither pluripotency self-renewal nor

NF-κB activity. A. Upper panel. Strategy to delete IκBα gene in mESCs using CRISPR/Cas9.

Three independent clones from each genotype (IκBα$^{+/+}$ or IκBα$^{-/-}$) were chosen for further

experiments. **Lower panel.** Western Blotting for IκBα between IκBα$^{+/+}$ and IκBα$^{-/-}$ clones. **B.**

Flow cytometry for the pluripotent surface marker SSEA-1 in IκBα$^{+/+}$ and IκBα$^{-/-}$ mESCs

cultured in S/L medium. **C. i)** Representative plots of the cell cycle profile in IκBα$^{+/+}$ and IκBα$^{-/-}$

mESCs by flow cytometry. **ii)** Quantification of cell cycle distribution in three independent

mESCs clones from each genotype. **D.** Boxplots and violin plots representing the normalized

gene expression levels in IκBα$^{+/+}$ or IκBα$^{-/-}$ mESCs, EBs 48h and EBs 96h. Considered genes are

those annotated to NF-κB signaling pathway (KEGG PATHWAY mmu04064). Box Plots: center

line, median; box limits, first and third quartiles; whiskers, 1.5× interquartile; points, outliers.

Violin plots show the corresponding probability density function. Two-sided Wilcoxon test

applied.

Figure S2. Absence of IκBα blocks pluripotency exit upon EB and teratoma formation. A.

Schematic of EB differentiation with brightfield images of different time points of

differentiation. **B.** Representative immunofluorescence images of 120h EBs stained for

NANOG and OCT3/4 from IκBα$^{+/+}$ (upper panel) and IκBα$^{-/-}$ (bottom panel) cells. **C. i)** Schematic

of AP staining experiment. **ii)** Brightfield images from AP colonies from three independent

clones from IκBα$^{+/+}$ or IκBα$^{-/-}$. **D.** Quantification of relative size of 168h EBs. Each dot

represents one EB. **E.** Flow cytometry plots of SSEA-1 from IκBα$^{+/+}$ or IκBα$^{-/-}$ EBs 96h. **F.**

Expression dynamics of different pluripotency markers *Pou5f1 (Oct4), Zfp42, Nanog* and *Sox2*

measured across EB differentiation. RNA expression levels were relative to time point 0

(mESCs). **G.** Schematic of the protocol to induce teratoma formation in NSG mice. **H.**

Hematoxylin & Eosin staining representative images from teratomas formed from

independent clones of IκBα$^{+/+}$ or IκBα$^{-/-}$ mESCs. Asterisks denote the presence of immature

structures.

Figure S3. Transcriptomic and epigenetic differences in IκBα$^{+/+}$ or IκBα$^{-/-}$ mESCs. A, B, C.

Volcano plots from IκBα$^{-/-}$ against IκBα$^{+/+}$ mESCs (A), 48h EBs (B) and 96h EBs (C) differential

expression analysis. Dashed vertical lines represent an absolute shrunkend log2 Fold Change

= 1 and dashed horizontal lines represented and adjusted p-value = 0.05 (FDR) **D.** Heatmap of

96h EBs in which normalized counts (z-score) of pluripotency or differentiation (gastrulation-

related) genes are represented. **E.** Volcano plot of H3K4me3 in IκBα$^{+/+}$ vs IκBα$^{-/-}$ mESCs

differential binding analysis. Representative genes with increased levels of H3K4me3 in IκBα$^{-/-}$

mESCs are highlighted. **F.** Volcano plot of H3K27me3 in IκBα$^{+/+}$ vs IκBα$^{-/-}$ mESCs differential

binding analysis. Representative genes with decreased levels of H3K27me3 in IκBα$^{-/-}$ mESCs

are highlighted. **G.** qPCR analysis of pluripotency genes with decreased levels of H3K27me3

in IκBα$^{-/-}$ mESCs. Expression analysis was performed in mESCs. **H.** Heatmap of differentially

gained and lost H3K9me3 regions in IκBα$^{-/-}$ and IκBα$^{+/+}$ mESCs (adjusted p-val < 0.05, FDR).

Only those regions with any associated gene are considered. Highlighted genes are imprinted

genes. Regions are sorted per their adjusted p-value obtained from differential binding

analysis (DBA). Heatmaps are centered in the region center and extended +/- 3kb. Fold

Change values are retrieved from DBA. **I.** Volcano plot of H3K9me3 in IκBα$^{+/+}$ vs IκBα$^{-/-}$ mESCs

differential binding analysis. Representative imprinted genes with decreased levels of

H3K9me3 in IκBα$^{-/-}$ mESCs are highlighted. **J.** Functional overrepresentation analysis results of

those genes with differential gain of H3K9me3 mark in IκBα$^{-/-}$ mESCs against GO Biological

Processes database. **K.** qPCR analysis of imprinted genes with differential loss of H3K9me3 in

mESCs.

Figure S4. IκBα$^{-/-}$ mESCs and 96h EBs display a global DNA hypomethylation status. A, B.

DNA methylation levels (mean ß-values) across chromosomes in mESCs (A) or 96h EBs (B). **C.**

DNA methylation levels in different genomic regions: promoters, gene bodies, enhancers and

imprinted regions (Unpaired T-Test). **D, E, F.** Functional analysis of differentially methylated

regions exclusively found in IκBα$^{-/-}$ mESCs (D), 96h EBs (E) or common between time points

(F).

Figure S5. Generation of separation-of-function (SOF) IκBα mutants to study how

chromatin-related IκBα controls pluripotency exit. A. IκBα$^{-/-}$ mESCs after overnight induction

(1µg/mL doxycycline) of the different SOF IκBα mutants. **B.** Quantification of *Nfkbia* mRNA

activation upon doxycycline induction in reconstituted IκBα$^{-/-}$ mESCs. **C.** Pull-down of GST-H2A

with cell lysates from different SOF IκBα mutant mESCs. **D.** Pull-down of GST-p65 (left) or GST-

p50 (right) with lysates from SOF IκBα mutant mESCs. **E.** Western blot analysis of IκBα protein

during EB differentiation after 48h of induction. **F.** Quantification of EB differentiation status

(percentage of pluripotent cells) at 216h after induction of SOF IκBα mutants. Data from two

independent experiments. **G.** Percentage of differentiated cells (determined as SSEA-1

negative) at EBs 216h after SOF IκBα induction.

REFERENCES

Aguilera, Cristina, Ruben Hoya-Arias, Guy Haegeman, Lluís Espinosa, and Anna Bigas. 2004. "Recruitment of IκBα to the *hes1* Promoter Is Associated with Transcriptional Repression." *Proceedings of the National Academy of Sciences* 101 (47): 16537–42.

Allis, C. David, and Thomas Jenuwein. 2016. "The Molecular Hallmarks of Epigenetic Control." *Nature Reviews. Genetics* 17 (8): 487–500.

Álvarez-Villanueva, Daniel, Luis Galán-Palma, Joan Bertran, Martin Floor, Laura Solé, Teresa Lobo-Jarne, María Maqueda, et al. 2023. "Separation-of-Function Mutants Reveal the NF-κB-Independent Involvement of IκBα in the Regulation of Stem Cell and Oncogenic Programs." *bioRxiv*. https://doi.org/10.1101/2023.06.21.545928.

Aparicio-Prat, Estel, Carme Arnan, Ilaria Sala, Núria Bosch, Roderic Guigó, and Rory Johnson. 2015. "DECKO: Single-Oligo, Dual-CRISPR Deletion of Genomic Elements Including Long Non-Coding RNAs." *BMC Genomics* 16 (October): 846.

Armstrong, Lyle, Owen Hughes, Sun Yung, Louise Hyslop, Rebecca Stewart, Ilka Wappler, Heiko Peters, et al. 2006. "The Role of PI3K/AKT, MAPK/ERK and NFkappabeta Signalling in the Maintenance of Human Embryonic Stem Cell Pluripotency and Viability Highlighted by Transcriptional Profiling and Functional Analysis." *Human Molecular Genetics* 15 (11): 1894–1913.

Brena, David, Joan Bertran, Montserrat Porta-de-la-Riva, Yolanda Guillén, Eric Cornes, Dmytro Kukhtar, Lluís Campos-Vicens, et al. 2020. "Ancestral Function of Inhibitors-of-kappaB Regulates Caenorhabditis Elegans Development." *Scientific Reports* 10 (1): 16153.

Choi, Jiho, Aaron J. Huebner, Kendell Clement, Ryan M. Walsh, Andrej Savol, Kaixuan Lin, Hongcang Gu, et al. 2017. "Prolonged Mek1/2 Suppression Impairs the Developmental Potential of Embryonic Stem Cells." *Nature* 548 (7666): 219–23.

Dutta, Debasree, Soma Ray, Pratik Home, Melissa Larson, Michael W. Wolfe, and Soumen Paul. 2011. "Self-Renewal versus Lineage Commitment of Embryonic Stem Cells: Protein Kinase C Signaling Shifts the Balance." *Stem Cells* 29 (4): 618–28.

Festuccia, Nicola, Rodrigo Osorno, Valerie Wilson, and Ian Chambers. 2013. "The Role of Pluripotency Gene Regulatory Network Components in Mediating Transitions between Pluripotent Cell States." *Current Opinion in Genetics & Development* 23 (5): 504–11.

Filipczyk, Adam, Carsten Marr, Simon Hastreiter, Justin Feigelman, Michael Schwarzfischer, Philipp S. Hoppe, Dirk Loeffler, et al. 2015. "Network Plasticity of Pluripotency Transcription Factors in Embryonic Stem Cells." *Nature Cell Biology* 17 (10): 1235–46.

Gerondakis, S., R. Grumont, R. Gugasyan, L. Wong, I. Isomura, W. Ho, and A. Banerjee. 2006. "Unravelling the Complexities of the NF-kappaB Signalling Pathway Using Mouse Knockout and Transgenic Models." *Oncogene* 25 (51): 6781–99.

Graf, Thomas, and Matthias Stadtfeld. 2008. "Heterogeneity of Embryonic and Adult Stem Cells." *Cell Stem Cell* 3 (5): 480–83.

Habibi, Ehsan, Arie B. Brinkman, Julia Arand, Leonie I. Kroeze, Hindrik H. D. Kerstens, Filomena Matarese, Konstantin Lepikhov, et al. 2013. "Whole-Genome Bisulfite Sequencing of Two Distinct Interconvertible DNA Methylomes of Mouse Embryonic Stem Cells." *Cell Stem Cell* 13 (3): 360–69.

Hayden, Matthew S., and Sankar Ghosh. 2008. "Shared Principles in NF-kappaB Signaling." *Cell* 132 (3): 344–62.

Hu, Sofia, Emily Metcalf, Dig Bijay Mahat, Lynette Chan, Noor Sohal, Meenakshi

Chakraborty, Maxwell Hamilton, et al. 2022. "Transcription Factor Antagonism Regulates Heterogeneity in Embryonic Stem Cell States." *Molecular Cell* 82 (23): 4410–27.e12.

Kaltschmidt, Christian, Johannes F. W. Greiner, and Barbara Kaltschmidt. 2021. "The Transcription Factor NF-κB in Stem Cells and Development." *Cells* 10 (8). https://doi.org/10.3390/cells10082042.

Kumar, Roshan M., Patrick Cahan, Alex K. Shalek, Rahul Satija, Ajay DaleyKeyser, Hu Li, Jin Zhang, et al. 2014. "Deconstructing Transcriptional Heterogeneity in Pluripotent Stem Cells." *Nature* 516 (7529): 56–61.

Leitch, Harry G., Kirsten R. McEwen, Aleksandra Turp, Vesela Encheva, Tom Carroll, Nils Grabole, William Mansfield, et al. 2013. "Naive Pluripotency Is Associated with Global DNA Hypomethylation." *Nature Structural & Molecular Biology* 20 (3): 311–16.

Marks, Hendrik, Tüzer Kalkan, Roberta Menafra, Sergey Denissov, Kenneth Jones, Helmut Hofemeister, Jennifer Nichols, et al. 2012. "The Transcriptional and Epigenomic Foundations of Ground State Pluripotency." *Cell* 149 (3): 590–604.

Marruecos, Laura, Joan Bertran, Daniel Álvarez-Villanueva, María Carmen Mulero, Yolanda Guillén, Luis G. Palma, Martin Floor, et al. 2021. "Dynamic Chromatin Association of IκBα Is Regulated by Acetylation and Cleavage of Histone H4." *EMBO Reports* 22 (8): e52649.

Marruecos, Laura, Joan Bertran, Yolanda Guillén, Jéssica González, Raquel Batlle, Erika López-Arribillaga, Marta Garrido, et al. 2020. "IκBα Deficiency Imposes a Fetal Phenotype to Intestinal Stem Cells." *EMBO Reports* 21 (6): e49708.

Martinez Arias, Alfonso, and Joshua M. Brickman. 2011. "Gene Expression Heterogeneities in Embryonic Stem Cell Populations: Origin and Function." *Current Opinion in Cell Biology* 23 (6): 650–56.

Mulero, María Carmen, Dolors Ferres-Marco, Abul Islam, Pol Margalef, Matteo Pecoraro, Agustí Toll, Nils Drechsel, et al. 2013. "Chromatin-Bound IκBα Regulates a Subset of Polycomb Target Genes in Differentiation and Cancer." *Cancer Cell* 24 (2): 151–66.

Singer, Zakary S., John Yong, Julia Tischler, Jamie A. Hackett, Alphan Altinok, M. Azim Surani, Long Cai, and Michael B. Elowitz. 2014. "Dynamic Heterogeneity and DNA Methylation in Embryonic Stem Cells." *Molecular Cell* 55 (2): 319–31.

Thambyrajah, Roshana, Zaki Fadlullah, Martin Proffitt, Wen Hao Neo, Yolanda Guillén, Marta Casado-Pelaez, Patricia Herrero-Molinero, et al. 2022. "IκBα Controls Dormancy Induction in Hematopoietic Stem Cell Development via Retinoic Acid." *bioRxiv*. https://doi.org/10.1101/2022.11.17.516971.

Tian, Tian V., Bruno Di Stefano, Grégoire Stik, Maria Vila-Casadesús, José Luis Sardina, Enrique Vidal, Alessandro Dasti, et al. 2019. "Whsc1 Links Pluripotency Exit with Mesendoderm Specification." *Nature Cell Biology* 21 (7): 824–34.

Torres, Josema, and Fiona M. Watt. 2008. "Nanog Maintains Pluripotency of Mouse Embryonic Stem Cells by Inhibiting NFkappaB and Cooperating with Stat3." *Nature Cell Biology* 10 (2): 194–201.

Wang, Yangming, Rostislav Medvid, Collin Melton, Rudolf Jaenisch, and Robert Blelloch. 2007. "DGCR8 Is Essential for microRNA Biogenesis and Silencing of Embryonic Stem Cell Self-Renewal." *Nature Genetics* 39 (3): 380–85.

Ying, Qi-Long, Jason Wray, Jennifer Nichols, Laura Batlle-Morera, Bradley Doble, James Woodgett, Philip Cohen, and Austin Smith. 2008. "The Ground State of Embryonic Stem Cell Self-Renewal." *Nature* 453 (7194): 519–23.

Zhang, Qian, Michael J. Lenardo, and David Baltimore. 2017. "30 Years of NF-κB: A
Blossoming of Relevance to Human Pathobiology." *Cell* 168 (1-2): 37–57.
Zhou, Wanding, Toshinori Hinoue, Bret Barnes, Owen Mitchell, Waleed Iqbal, Sol Moe Lee,
Kelly K. Foy, et al. 2022. "DNA Methylation Dynamics and Dysregulation Delineated by
High-Throughput Profiling in the Mouse." *Cell Genomics* 2 (7).
https://doi.org/10.1016/j.xgen.2022.100144.

Figure 1

Figure 2

Figure 3

Figure 4