Chromatin enhancer activity of IκBα mediates the exit from naïve pluripotency

Luis G. Palma[1,2,3,δ], Daniel Álvarez-Villanueva[1,3,4,δ], María Maqueda[1,2,3], Mercedes Barrero[5],

Arnau Iglesias[1,2,3], Joan Bertran[6], Damiana Álvarez-Errico[2], Carlos A. García-Prieto[2], Cecilia

Ballaré[5], Virginia Rodriguez-Cortez[2], Clara Bueno[2], August Vidal[4], Alberto Villanueva[7], Pablo

Menéndez[2,3,8,9,10], Gregoire Stik[2], Luciano Di Croce[5,10,11], Bernhard Payer[5,11], Manel

Esteller[2,3,10,12], Lluís Espinosa[1,3,*] & Anna Bigas[1,2,3,*]

[1]Program in Cancer Research. Hospital del Mar Research Institute, Barcelona, Spain.

[2]Josep Carreras Leukemia Research Institute, Barcelona Spain.

[3]Centro de Investigación Biomédica en Red Cancer (CIBERONC), Madrid, Spain.

[4]Institut Investigació Biomèdica de Bellvitge (IDIBELL), L'Hospitalet de Llobregat, Barcelona,
Spain.

[5]Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology,
Barcelona, Spain.

[6]Faculty of Sciences, Technology and Engineering, Universitat de Vic - Universitat Central de
Catalunya, Vic 08500 Spain

[7]Chemoresistance and Predictive Factors Group, Program Against Cancer Therapeutic
Resistance (ProCURE), Catalan Institute of Oncology (ICO), Oncobell Program, Bellvitge
Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain.

[8]Spanish Network for Advanced Therapies (RICORS-TERAV). Carlos III Health Institute (ISCIII),
Spain.

[9]Department of Biomedicine. University of Barcelona, Spain.

[10]Institucio Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

[11] Universitat Pompeu Fabra, Barcelona, Spain.

[12]Physiological Sciences Department, School of Medicine and Health Sciences, University of
Barcelona (UB), Barcelona, Catalonia, Spain.

[δ]These authors contributed equally

Keywords: IκBα, NFKBIA, NF-κB, DNA Methylation, Polycomb, Embryonic Stem Cells,
Pluripotency Exit, Germ Layer Specification,

*Corresponding authors:

Anna Bigas and Lluis Espinosa

Cancer Research Program. CIBERONC.

Hospital del Mar Research Institute

Doctor Aiguader 88, 08003, Spain.

Tel: +34 933160440; email: abigas@researchmar.net; lespinosa@researchmar.net

Running title: IκBα function in pluripotency exit

Keywords: Embryonic Stem Cells, IkappaB, NfkappaB, Chromatin Regulation, Enhancer,
Inflammation, Naïve pluripotency, Pluripotency

Abstract

Inflammation is crucial in development and cell differentiation, albeit their involvement in the
transition from naïve to primed pluripotency remains elusive. One of the main sensors of inflammation
is IκBα, which exerts its function through activation of NF-κB. Our previous work identified a chromatin
role of IκBα that is crucial for the proper regulation of adult stem cells. Here, we demonstrate that
IκBα depletion causes profound epigenetic rewiring in pluripotent stem cells characterized by altered
histone post-translational modifications and DNA methylation, which is associated with changes in
transcriptional regulation. These changes led to stabilization of cells in a ground state of naïve
pluripotency when cultured in Serum/LIF conditions and differentiation impairment. Mechanistically,
IκBα binds to intergenic chromatin regions in the naïve pluripotency state, which result in altered
activation status of a subset of enhancers, including several associated to pluripotency-related genes.
Through the engineering of separation-of-function mutants, we demonstrate that the effects of IκBα
in the regulation of stem cell pluripotency are NF-κB-independent, but mainly relies on its chromatin-
related function. Altogether, our results highlight a novel mechanism by which IκBα participates in
regulating the pluripotent states of embryonic stem cells, shedding light on the interplay between
inflammatory signals and pluripotency regulation.

INTRODUCTION

Embryonic Stem Cells (ESCs) pluripotency is characterized by the capacity to generate all somatic and germline lineages both *in vitro* and *in vivo*, depending on the culture conditions. When cultured in serum plus Leukemia Inhibitory Factor (LIF) (Serum/LIF), murine ESCs (mESCs) include cells interconverting to different metastable states (Festuccia et al. 2013) that either resemble preimplantation blastocyst (naïve pluripotency) or post-implantation (primed pluripotency) embryonic stages (Graf and Stadtfeld 2008; Martinez Arias and Brickman 2011; Pera and Rossant 2021; Wang and Wu 2022). The heterogeneity of mESCs in Serum/LIF is reduced by culturing them in the presence of GSK3β and MEK1/2 inhibitors (CHIR99021 and PD0325901, respectively) along with LIF (2i/LIF). Under these conditions, mESCs enter a ground state of naïve pluripotency characterized by self-renewal activity while suppressing any pro-differentiating signals (Ying et al. 2008; Marks et al. 2012). This pluripotent state closely resembles the inner cell mass of the E4.0 mouse preimplantation blastocyst (Boroviak et al. 2015). After implantation, pluripotent capability of the mouse embryo (E5.5) becomes restricted to Epiblast Stem Cells (EpiSCs), which represent a primed pluripotent state, characterized by coexpression of pluripotency and lineage specification markers (Brons et al. 2007; Tesar et al. 2007). EpiSCs can give rise to cells of the three different germ layers (endoderm, mesoderm and ectoderm), albeit their contribution to chimeras or to germ cells is greatly compromised (Weinberger et al. 2016; Pera and Rossant 2021). EpiSCs can be derived *in vitro* from naïve pluripotent cells cultured in a medium supplemented with Nodal and fibroblast growth factor (FGF) (Guo et al. 2009). Different epigenetic features distinguish naïve from primed pluripotent states (Takahashi et al. 2018), such as changes in the X chromosome inactivation in female cells (Bao et al. 2009; Guo et al. 2009), histone post-translational modifications (Tesar et al. 2007) or a global DNA hypomethylation profile observed in naïve pluripotent stem cells (Habibi et al. 2013; Leitch et al. 2013; Singer et al. 2014). These epigenetic changes are a prerequisite for the subsequent activation of gene circuits associated with pluripotency exit and germ layer specification (Brons et al. 2007; Tesar et al. 2007). Crucial regulatory elements for the transition from naïve to primed pluripotency state are located in distal DNA regions or enhancers to facilitate the resolution toward the primed state (Factor et al. 2014; Buecker et al. 2014). Therefore, identifying novel players to fine-tune the equilibrium between ground and primed states is crucial for preserving both pluripotency stability and differentiation capability in mESCs.

Inflammation is a defense mechanism mostly activated in response to harmful endogenous or foreign stimuli, which involves different signaling pathways including NF-κB. Canonical NF-κB signaling is triggered by the IKK kinase complex, which induce the phosphorylation and subsequent degradation of the NF-κB inhibitor, IκBα, leading to nuclear translocation of the NF-κB factors (e.g. p50/p65) (Zhang et al. 2017). We have previously identified an alternative nuclear function of IκBα involving the interaction with histone deacetylases (HDACs), core elements of the Polycomb Repressor Complex 2 (PRC2) and histones (Aguilera et al. 2004;

Mulero et al. 2013; Marruecos et al. 2021). This nuclear IκBα function is crucial for tissue stem
cell homeostasis and the proper differentiation of epidermal, intestinal and hematopoietic
cells (Mulero et al. 2013; Marruecos et al. 2020, 2021; Thambyrajah et al. 2022).

There is now robust evidence that inflammatory signals are crucial in stem cell development
(Gerondakis et al. 2006; Kaltschmidt et al. 2021), and it was recently demonstrated that
attenuation of inflammatory signals by ATG5 facilitates mESCs differentiation (Li et al. 2024).
Unexpectedly, being NF-κB pathway a pivotal regulator of inflammation, classical NF-κB
factors including p50 and RelA/p65 are expressed at very low levels in mESCs. Overexpression
of NF-κB proteins leads to cell differentiation while expression of the IκBα super-repressor
(S32-36A mutant) increases expression of pluripotency markers (Torres and Watt 2008).
Similarly, inhibition of Protein Kinase C (PKC) activity in mESCs, which in turn restrains NF-κB
signaling, allows keeping mESCs in a ground pluripotency state, thus facilitating the
generation of induced pluripotent stem cells (iPSCs) (Dutta et al. 2011). Together these results
indicate that NF-κB inhibition is required for maintaining the pluripotent state of murine ESCs.
Given the importance of NF-κB in the multiple aspects of stem cell biology (Espín-Palazón and
Traver 2016) and the pivotal role of IκBα in NF-κB regulation, the possible impact of the IκBα
chromatin function in cell fate has remained challenging.

We have now investigated the role of IκBα in a context of pluripotent stem cells.
Unexpectedly, and in contrast with the pro-differentiation effects of NF-κB activation,
depletion of the NF-κB inhibitor IκBα precludes the exit of mESCs from naïve pluripotency
associated with an epigenetic rewiring. By using the recently described Separation-Of-
Function (SOF) IκBα mutants (Alvarez-Villanueva et al. 2023), we have now demonstrated that
pluripotency regulation by IκBα is mainly exerted through its chromatin-related function but
independent on its capacity to negatively regulate NF-κB.

RESULTS

IκBα accumulates in the chromatin of naïve pluripotent cells

Because of the prominent role of NF-κB signaling in mESCs regulation, we analyzed the expression pattern of the different NF-κB members including the IκB inhibitors in the three defined pluripotent stages: ground state of naïve pluripotency (2i/LIF), naïve pluripotency (Serum/LIF), and primed pluripotency (Epiblast Stem Cells or EpiSCs) (Figure 1A). In agreement with NF-κB being a pro-differentiation factor, canonical NF-κB inhibitors including IκBα (*Nfkbia*), IκBβ (*Nfkbib*) and IκBε (*Nfkbie*) were all expressed in the naïve pluripotency state (both ground and Serum/LIF), with *Nfkbia* levels significantly reduced upon naïve pluripotency exit towards EpiSCs (Figure 1B). In contrast, *Rela, Nfkb1 and Nfkb2 genes* codifying for the NF-κB factors p65/RelA, p105 and p100, respectively, showed an opposite pattern and either increased or maintained their levels in EpiSCs (Figure 1C). We further characterized the distribution of IκBα in the cytoplasm, nucleus and chromatin fractions of the three states of pluripotency (Figure 1D). IκBα protein levels in the cytoplasm and the nucleus correlated with the detected RNA levels in the three different cellular states being highest in 2i/LIF condition (Figure 1B). Interestingly, a notable accumulation of IκBα was observed in the chromatin of naïve pluripotent cells cultured in Serum/LIF conditions compared to 2i/LIF. In contrast, IκBα was entirely absent from the chromatin of EpiSCs. (Figure 1D). The abundance of IκBα RNA and protein in the context of naïve pluripotency compared to EpiSCs was further corroborated through analysis of an additional database encompassing both naïve and primed pluripotent states (Atlasi et al. 2020) (Figure 1E).

Next, we studied IκBα expression in differentiated embryoid bodies (EBs) from mESCs (Serum/LIF) (see materials and methods) and analyzed the expression levels of different pluripotency (*Dppa4, Pou5f1, Sox2, Prdm14, Klf4, Zfp42 [Rex1], Nanog*), differentiation-linked (*Mesp1, Gata6, Lhx1, Gata4, Wnt3a, Fgf5, Nes, Axin2, Wnt3, Eomes, T, Foxa2, Mixl1, Lef1*), and NF-κB genes (*Nfkbid, Nfkbia, Nfkbie, Nfkbib, Nfkbiz, Rel, Rela, Relb*) by RNA sequencing. *Nfkbia* (IκBα gene) was expressed at higher levels specifically in mESCs (Serum/LIF) whereas cells undergoing differentiation (48h and 96h) displayed a reduction in its expression levels (Figure 1F). The expression pattern of *Nfkbia* was similar to the pluripotency genes, whereas most of the NF-κB genes were upregulated upon differentiation, clustering together with developmental genes, as previously described (Torres and Watt 2008).

Taken together, these data show that the expression of *Nfkbia*, like the other *Nfkbi*, is high in the naive pluripotent state, in contrast to the NF-κB genes, which cluster together with differentiation markers. In addition, IκBα accumulates in the chromatin of naive pluripotent cells, more so in cells cultured under serum/LIF.

Lack of IκBα impairs the exit from the naïve pluripotency state

To investigate the functional implication of the chromatin accumulation of IκBα in naïve
pluripotency, we knocked-out IκBα protein in mESCs by CRISPR-Cas9 (IκBα-KO mESCs) (Figure
1G and materials and methods). We did not detect major changes in NF-κB target genes upon
IκBα depletion neither in mESCs, nor 48h and 96h EBs (Figure 1H), suggesting that NF-κB
regulation did not require IκBα at this stage or it might be compensated by the other canonical
or non-canonical IκBs (IκBβ, IκBε or p100), as previously demonstrated (Hayden and Ghosh
2008; Basak et al. 2007). Paradoxically, IκBα-KO mESCs have an impaired capability to undergo
differentiation, since IκBα-KO EBs cannot properly evolve towards differentiation trajectories
(Figure 2A). This observation was confirmed by analysis of naïve pluripotency markers at 96h
(*Gab1, Sox2, Esrrb, Zfp42, Dppa4, Klf2, Prdm14, Klf4, Tfcp2l1*), which displayed higher
expression levels in IκBα-KO compared to their wild-type (IκBα-WT) counterparts (Figures 2B
and S1A). Differentiation of IκBα-KO EBs was deficient in programs associated with all germ
layer specification (endoderm, mesoderm, and ectoderm) (Figures 2C-E). Moreover, IκBα-KO
EBs were smaller in size (Figure S1B), and maintained higher expression of the pluripotency
program even at 216h (Figures 2F and S1C). We also detected a higher number of Alkaline
Phosphatase-expressing cells (Figures 2G and S1D) and higher expression of the pluripotency
marker SSEA-1 (Figure S1E) at 216h IκBα-KO EBs. To further study the requirement of IκBα in
pluripotency exit *in vivo*, we established teratomas by subcutaneous injection of IκBα-WT and
IκBα-KO mESCs (Figure S1F and materials and methods) in NSG (NOD.Cg-Prkdc[scid]
Il2rg[tm1Wjl]/SzJ) mice. IκBα-KO teratomas showed a decreased differentiation potential, as
indicated by higher number of OCT4[+] cells (Figures S1G-H). These observations underscore
the essential role of IκBα in the exit from the pluripotency state in mouse embryonic stem
cells (mESCs).

To determine the specific time-point in which IκBα is required for pluripotency exit, we
differentiated IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF (naïve pluripotency)
towards EpiSCs (primed pluripotency) for 120h (Figure 2H and materials and methods). IκBα-
KO cells were not committed towards the primed pluripotency stage and maintained an
elevated expression levels of naïve pluripotency genes (*Dppa3, Nanog, Sox2, Rex1, Klf2, Klf4,
Gbx2, Tbx3*) (Figure 2I). Additionally, IκBα-KO cells retained a naïve pluripotent morphology,
forming tight clusters and resembling cells cultured in 2i/LIF medium (Figure S1I). These data
demonstrate that IκBα is required for the exit of the naïve state of pluripotency and its
deficiency results in impaired differentiation of mESC *in vivo* and *in vitro*, without imposing
major changes in the regulation of canonical NF-κB.

IκBα-KO mESCs resemble the ground-state of naïve pluripotency under Serum/LIF
conditions

As mentioned, mESCs cultured in serum/LIF are highly heterogeneous, containing a mixture
of cell states that resemble developmental transitions from preimplantation (naïve
pluripotency) to postimplantation (primed pluripotency) embryo (Graf and Stadtfeld 2008;
Martinez Arias and Brickman 2011). To investigate how IκBα deficiency was affecting the exit

of mESCs from naïve pluripotency, we performed a Gene Set Enrichment Analysis (GSEA)
using RNA-seq data from IκBα-KO and IκBα-WT mESCs cultured in Serum/LIF against gene
signatures for ground (2i/LIF) and naïve (Serum/LIF) pluripotency (Ghimire et al. 2018). GSEA
results revealed a significant enrichment in Serum/LIF IκBα-KO mESCs transcriptome for
genes specifically expressed in the ground state (2i/LIF), while they negatively correlate with
mESCs cultured in Serum/LIF (Figure 3A) (Ghimire et al. 2018). Specifically, IκBα-KO mESCs
expressed higher levels of the ground state pluripotency genes *Klf2, Nanog, Pou5f1, Prdm14,*
Gbx2, Tbx3, Zfp42[Rex1] and *Klf4* (Figure 3B), similar to IκBα-WT mESCs cultured in 2i/LIF
(Figure S2A). Since ground (2i/LIF) and naïve (Serum/LIF) states are characterized by
important differences in epigenetic marks, we carried out an epigenetic profiling of IκBα-WT
and IκBα-KO mESCs cultured in Serum/LIF. We performed chromatin immunoprecipitation
followed by sequencing (ChIP-seq) of histone modifications associated to gene activation
(H3K4me3), gene repression (H3K27me3) and constitutive heterochromatin (H3K9me3)
(Figures 3C-D and S2B-D). Overall, Serum/LIF IκBα-KO mESCs exhibited a more permissive
chromatin status as evidenced by increased H3K4me3 and decreased H3K9me3 levels, while
H3K27me3 was redistributed in both directions (Figure 3C). Notably, the gain in H3K4me3
mark was associated with a reduction of H3K27me3 levels in genes associated with naïve
pluripotency (Figures 3C-D and S2B-C), in agreement with the higher pluripotent state
observed in IκBα-KO mESCs. As mentioned, detection of the H3K9me3 mark was significantly
reduced in IκBα-KO mESCs (Figures 3C and S2D). These regions correspond to genes involved
in DNA methylation and chromatin edition functions (Figure 3C). By analyzing the DNA
methylation status under Serum/LIF conditions by immunofluorescence of 5-methyl-cytosine
(5mC) we found a global DNA hypomethylation in IκBα-KO mESCs (Figure 3E), which is one of
the key features of the ground state of naïve pluripotency (2i/LIF) (Habibi et al. 2013; Yagi et
al. 2017). To further characterize the DNA methylation status in IκBα-KO cells, we performed
DNA methylation arrays (Zhou et al. 2022) in IκBα-KO and IκBα-WT mESCs (Serum/LIF) and
96h after EB differentiation. We observed a strong reduction of the DNA methylation level in
IκBα-KO mESCs in a genome-wide fashion (Figure 3F). Specifically, IκBα-KO mESCs contain
lower DNA methylation levels in promoters, gene bodies, enhancers and imprinted regions
(Figure 3G), and this DNA hypomethylation status was not associated with any particular
chromosome (Figure S2E). DNA hypomethylation was maintained upon 96h in IκBα-KO EBs
(Figures 3F-G and S2F).

These results further indicate that IκBα is required for the transition from ground to primed
pluripotent states and for the proper regulation of the histone post-translational
modifications (PTMs) and DNA methylation changes associated with this process. We propose
that the enhanced pluripotent state found in IκBα-KO mESCs under basal conditions
(Serum/LIF) may negatively impact on their differentiation potential, similar to that described
for other models with stabilized naïve pluripotent state (Choi et al. 2017; Di Stefano et al.
2019).

IκBα binds to intergenic regions to attenuate the activation status of chromatin linked to naïve pluripotency genes

To gain insight into the mechanism by which IκBα regulates the transition from naïve to primed pluripotent states, we knocked-in a 3xFLAG-tag into the IκBα locus and used these transgenic cells for ChIP-seq assay with the anti-FLAG antibody (see Materials and Methods). We identified 817 bona-fide IκBα-bound chromatin regions which were consistent between three independent replicates (Figure S3A). Among all the genome regions, IκBα was mainly bound to distal intergenic and intronic regions (38.5% and 42.2% of total IκBα peaks, respectively) (Figure 4A). Interestingly, IκBα-bound regions coincide with those that gain enhancer activation-related H3K27Ac signals in EpiSCs (Figures S3B-C). The fact that IκBα was released from chromatin at the stage of primed pluripotency (Figure 1D) suggested that IκBα might be involved in fine-tuning these regions at the exit from naive pluripotency. To address this question, we quantified the total amount of H3K27Ac (active enhancer) by western blot analysis of chromatin extracts of IκBα-WT and IκBα-KO mESCs cultured in Serum/LIF. We observed an increase of this activation mark in IκBα-KO mESCs, whereas we did not observe global changes in H3K4me1 levels, a marker of poised enhancers (Figures S3D-E). These results suggested that IκBα depletion in naïve pluripotency might lead to an increase in the global levels of active enhancers, which are characterized by a gain of H3K27Ac without changes in H3K4me1 levels (Wu et al. 2023). The increase in H3K27Ac was further validated by ChIP-seq experiments conducted in both IκBα-WT and IκBα-KO mESCs. Notably, depletion of IκBα protein under Serum/LIF conditions led to an increase in the deposition of H3K27Ac at regions where we detected IκBα binding (Figure 4B), while levels of H3K4me1 were not generally altered. Since IκBα deficiency was affecting enhancer-related histone marks, we further profiled the enhancer status of IκBα-KO mESCs by analyzing the differential distribution of poised enhancers (showing changes in H3K4me1 levels without affecting H3K27Ac) and active enhancers (chromatin regions decorated with H3K4me1 that also display H3K27ac binding) (Figures 4D). Overall, IκBα-KO mESCs showed a reduction in poised enhancers at genes (or related genomic regions) associated to differentiation processes (Figures 4D and S3F), whereas active enhancers of genes associated to pluripotency were significantly enriched (Figures 4D-E and S3G). These results further support that mESCs lacking IκBα are epigenetically reinforced in the ground state of naïve pluripotency. Since the majority of IκBα chromatin peaks were located in distal intergenic and intronic regions (Figure 4A), and we found that IκBα deficiency led to changes in the enhancer status (Figure 4D), we took advantage of publicly available Hi-C data to identify topologically associating domains (TADs) at putative IκBα target genes in Serum/LIF mESCs (Bonev et al. 2017). It was previously demonstrated that TADs are organized in distinct chromatin compartments defined as active (Compartment A) or inactive (Compartment B) (Lieberman-Aiden et al. 2009). Importantly, chromatin regions bound by IκBα mainly correspond to TADs of the active compartment (81%), whereas a small proportion of IκBα peaks were found in inactive chromatin (19%) (Figure S3H). Furthermore, TADs harboring IκBα binding exhibited a greater proportion of

enhancers differentially regulated in IκBα-KO cells when compared to all TADs (92% vs. 74%,
respectively) (Figure S3H). Additionally, genes located within TADs bound by IκBα are
predominantly associated with mechanisms linked to pluripotency identity (data not shown).
To address the functional relevance of this observation, we investigated the expression
pattern of pluripotency-related genes in IκBα-WT and IκBα-KO mESCs at naïve pluripotency
(cultured in Serum/LIF) and differentiation (48h EBs) stages (Figure 4F). Most of the genes
analyzed were significantly upregulated in IκBα-KO mESCs, and this difference was further
accentuated in IκBα-KO 48h EBs. These results suggest that IκBα modulates the dynamic
expression of pluripotency genes through a distal regulatory mechanism involving enhancer
activity and 3D chromatin architecture. To address this hypothesis, we crossed Hi-C data with
ChIP-seq of IκBα and enhancer-related histone marks (H3K27Ac and H3K4me1), together with
RNA-seq data. In this meta-analysis, we identified a TAD region encompassing the Tbx3 locus
with IκBα binding sites (Figure 4G). Interestingly, we observed that the IκBα-bound regions
interact with the region containing Tbx3 gene (Figure 4G; interactions highlighted with an
arrow in the contact map). Moreover, that region harbors an active enhancer marked by
H3K27Ac/K4me1, whose activation status is heightened in IκBα-KO mESCs, likely participating
in increased expression of the Tbx3 gene (Figure 4G, bottom panel) and other interactions are
detected between the differentially activated enhancer and the Tbx3 locus in this TAD (Figure
4G, interaction highlighted with a gray arrow in the contact map).

We then determined whether differential enhancer activation status observed in IκBα-KO
mESCs (evidenced by increased H3K27Ac) was reversible. We engineered a doxycycline-
inducible IκBα construct and introduced it into IκBα-KO mESCs. Treatment of these cells with
doxycycline for several passages under Serum/LIF conditions (i-IκBα^{WT} mESCs, Figure 4H) led
to sustained IκBα expression and a significant reduction of global levels of H3K27Ac (Figures
4I and S3I). Ectopic expression of i-IκBα^{WT} in mESCs was also sufficient to attenuate the
expression of the naïve pluripotency genes *Zfp42, Klf2, Sox2* and *Tbx3 in* IκBα-KO mESCs under
Serum/LIF conditions (Figure S3L).

IκBα attenuates the activation of intergenic regions in an NF-κB independent manner

Uncovering the actual impact of chromatin-related IκBα function is challenging due to the
predominant role of this protein on NF-κB regulation. Recent work from our group has
identified the residues in the IκBα protein critical for NF-κB-binding or chromatin-binding
through H2A/H4. Mutation of these residues, allows to specifically discriminate between
either of the two functions of IκBα (Alvarez-Villanueva et al. 2023). We generated doxycycline-
inducible constructs to express separation-of-function (SOF) IκBα mutants: i) exclusively
interacts with NF-κB and is deficient for chromatin binding (i-IκBα^{ΔChromatin}) or ii) interacts with
histones but is deficient for NF-κB interaction and inhibition (i-IκBα$^{ΔNF-κB}$) (Figure 5A). We
introduced one of the SOF IκBα mutants into IκBα-KO mESCs and confirmed their functionality
by pull-down experiments. We found that i-IκBα$^{ΔNF-κB}$ was able to interact with histone H2A
but not with the NF-κB member p50 and, i-IκBα^{ΔChromatin} could interact with p50 but not with

H2A, while i-IκBα^{WT} preserved all interactions (Figures 5B-C). Importantly, when we induced
the three different IκBα forms (i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and i-IκBα^{ΔChromatin}, Figure S4A), only i-
IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ were found in the chromatin fraction of mESCs, whereas IκBα^{ΔCromatin}
was retained in the cytoplasm (Figures 5D and S4B). To address whether IκBα was
participating in the stabilization of the ground-state of naïve pluripotency in an NF-κB-
independent manner, we studied which of the SOF mutants (IκBα$^{ΔNF-κB}$ or IκBα^{ΔChromatin}) was
able to reverse the phenotype of IκBα-KO mESCs (Figure S4C). We observed the reduction of
the global levels of H3K27Ac upon induction of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$, but not of
IκBα^{ΔChromatin} (Figures 5E and S4D). Similarly, induction of IκBα$^{ΔNF-κB}$ was enough to reduce the
H3K27Ac amount in enhancers with increased H3K27Ac in IκBα-KO mESCs, but not when
IκBα^{ΔChromatin} was expressed (Figure 5F). Interestingly, H3K4me1 levels followed a similar
pattern as H3K27Ac in those specific loci, and a significant reduction was achieved upon
IκBα$^{ΔNF-κB}$ overexpression, whereas IκBα^{ΔChromatin} was mediating an opposite effect (Figure 5G).
In addition, the ground state-associated DNA hypomethylation was reverted in i-IκBα^{WT} and
i-IκBα$^{ΔNF-κB}$ mESCs but not in i-IκBα^{ΔChromatin} mESCs (Figure 5H). Furthermore, the naïve
pluripotent gene expression (*Zfp42, Klf2, Sox2*, *Tbx3*) was reduced upon inducing i-IκBα$^{ΔNF-κB}$,
while induction of i-IκBα^{ΔChromatin} resulted in their increase (Figure S4E).

To address whether reverting to the basal epigenetic status is enough to rescue the
differentiation capability, i-IκBα^{WT}, i-IκBα$^{ΔNF-κB}$ and i-IκBα^{ΔChromatin} mESCs were differentiated
into EBs (Figure S4C). Induction of IκBα^{WT} and IκBα$^{ΔNF-κB}$ was enough to restore the
differentiation potential of IκBα-KO mESCs, since the number of differentiated cells (based on
percentage of SSEA-1^{neg} cells) increased (Figures 5I and S4F). Moreover, we found a reduced
number of AP+ colonies from 216h IκBα-KO EBs after induction of i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$,
similar to the number of colonies obtained from IκBα-WT EBs (Figures 5J and S4G). Lastly, the
number of remaining pluripotent-like cells (defined by OCT3/4 NANOG staining) was also
reduced in i-IκBα^{WT} and i-IκBα$^{ΔNF-κB}$ 216h EBs, and only small and few regions of
undifferentiated cells was observed (Figures 5K and S4H). On the other hand, induction of i-
IκBα^{ΔChromatin} did not restore the differentiation potential of IκBα-KO mESCs, and phenocopied
the differentiation defects observed in IκBα-depleted mESCs.

These results demonstrate that the chromatin-dependent IκBα function, but not its NF-κB-
related activity, is necessary and sufficient to revert the stabilization in the ground state of
naïve pluripotency correlating with higher DNA methylation and lower H3K27Ac levels. Our
results demonstrate that the chromatin function of IκBα alters the activation status of
enhancers, which impacts the expression of genes involved in the maintenance of the naïve
pluripotency. This function is independent of the NF-κB-inhibitory activity of IκBα.

DISCUSSION

Here we have identified a new and unexpected function for the IκBα protein in the regulation of the naïve pluripotency exit. Using mESCs to investigate the role of IκBα in pluripotency maintenance and differentiation, we discovered that the absence of IκBα severely impairs the exit of pluripotency. Furthermore, IκBα-KO mESCs stabilizes the ground state of naïve pluripotency under Serum/LIF conditions, shedding light on its function as a critical regulator of the transition from naïve to primed pluripotency. While IκBα is commonly known for its role as an inhibitor of the NF-κB signaling pathway (Zhang et al. 2017), our laboratory has previously furnished compelling evidence of its alternative role through its interaction with histones and other chromatin components (Aguilera et al. 2004; Mulero et al. 2013; Marruecos et al. 2021). We have demonstrated that chromatin-bound IκBα plays a crucial role in regulating skin and intestinal stem cells (Mulero et al. 2013; Marruecos et al. 2020), hematopoietic stem cell development (Thambyrajah et al. 2022), and it influences the regenerative capacity of these tissues, and in some cases, their susceptibility to neoplastic transformation. Disentangling the chromatin-related function of IκBα from its canonical role as an inhibitor of NF-κB has been challenging, making it difficult to determine its truly biological significance. To tackle this challenge, we engineered separation-of-function (SOF) mutants of IκBα, as detailed in (Álvarez-Villanueva et al. 2023), by identifying specific residues that are essential for binding either NF-κB elements or histones. Interestingly, we have shown that only the histone-binding proficient IκBα mutant (IκBα$^{\Delta NF-\kappa B}$), which lacks the ability to bind to NF-κB, has the capability to reverse the stabilization of the ground state observed in IκBα-KO mESCs. This is sufficient to facilitate their exit from the state of naïve pluripotency, restoring their full differentiation potential (Figure 5). The involvement of NF-κB in pluripotency has been a subject of prior investigation. Studies have revealed that pluripotent cells display a dampened inflammatory signaling pathway, which is induced upon pluripotency dissolution and activation of differentiation (Torres and Watt 2008; Guo 2019). However, the precise molecular mechanisms underlying the attenuated status of NF-κB during this stage remain incompletely understood. A recent work has shed light on a potential regulatory axis involving ATG5-β-TrCP1-NF-κB, proposing it as a mechanism that diminishes NF-κB activity by stabilizing the IκBα protein in mESCs (Li et al. 2024). In any case, since modification of canonical NF-κB activity directly impacts on the levels IκBα expression, the previous NF-κB-associated phenotypes should be reevaluated considering this chromatin role of IκBα. Furthermore, the expression of IκBα$^{\Delta Chromatin}$, which retains the inhibitory function of NF-κB, failed to mitigate the ground state stabilization observed in IκBα-KO mESCs (Figure 5). This finding suggests that the observed role of IκBα in regulating the exit from naïve pluripotency is not due to NF-κB inhibition. In fact, there is evidence that the chromatin function of IκBα is an ancestral function preceding its role as NF-κB inhibitor. In Caenorhabditis Elegans (*C.elegans*), which lacks NF-κB orthologs but possesses two homologs of IκBα, the absence of IκBα orthologs has severe multi-organ differentiation defects (Brena et al. 2020). This observation further supports the notion that the chromatin-related function

of IκBα plays a fundamental role in developmental processes, independent of its canonical
role in NF-κB signaling.

We have observed that IκBα-KO mESCs exhibit epigenetic and transcriptomic profiles
resembling the ground state of naïve pluripotency under Serum/LIF culture (Figure 3). In this
regard, the fact that IκBα is preferentially located in the chromatin of mESCs cultured in
Serum/LIF, when they exhibit a high degree of heterogeneity, suggests that IκBα might act as
a regulator of this heterogeneity. One potential explanation might be that IκBα could be
affecting the stability of fluctuating transcription factors, which is crucial to mediate the
pluripotent-to-differentiation balance (Martinez Arias and Brickman 2011). Thus, the lack of
IκBα might stabilize those transcription factors that would favor the ground-state observed in
IκBα-KO mESCs cultured in Serum/LIF. Further experiments are required to demonstrate this
hypothesis. Moreover, the differences in murine and human pluripotency (Weinberger et al.
2016) point out the necessity to address the real impact of IκBα modulation in the stabilization
of the ground state in human pluripotent stem cells. Our investigation has also revealed that
IκBα accumulates in the intergenic chromatin regions of naïve pluripotent cells (Figures 1D
and 4A). Mechanistically, this binding of IκBα serves to safeguard enhancers associated with
pluripotency from excessive activation. This is evidenced by the higher abundance of active
enhancers observed in IκBα-depleted mESCs, particularly in regions linked to pluripotency
located within TADs where IκBα is bound. A recent study highlights the significance of
maintaining a proper 3D chromatin architecture during the resolution of the naïve pluripotent
state (Lando et al. 2024). Our findings suggest that IκBα may potentially influence the 3D
conformation of regions relevant to pluripotency, thereby fine-tuning the expression of genes
crucial for sustaining naïve pluripotency (Figure 6). Nevertheless, the precise mechanisms by
which the absence of IκBα impacts 3D chromatin architecture remain to be elucidated.

In summary, our work establishes that IκBα mediates the exit from naïve pluripotency by
modulating the activation of enhancers associated with pluripotency genes. Importantly, this
newly identified role of IκBα operates independently of its canonical function in inhibiting the
NF-κB pathway. Our findings underscore the intricate interplay between inflammation and
pluripotency status, revealing a previously unrecognized complexity in their interaction.

MATERIALS & METHODS

Cell culture

mESC line ES-E14TG2a (ATCC; Cat #CRL-1821) was cultured on plastic dishes precoated with 0.1% (w/v) Gelatin (Sigma; Cat #G2500-100G) in Serum/LIF medium. Medium was changed every day, and cells were passaged every other day using TrypLE Express (Gibco; Cat #12605010) for harvesting. Cells were maintained in a 5% CO2 incubator at 37°C. For inducing the ground-state of naïve pluripotency, mESCs were cultured in 2i/LIF medium for 2 consecutive passages.

Epiblast stem cell differentiation

Differentiation from mESCs towards Epiblast Stem Cells (EpiSCs) was performed as previously described (Pantier et al. 2019). 3×10^4 mESCs were seeded in 6-well plastic dishes, and they were cultured in Serum/LIF medium for 24h. Medium was then switched to EpiSCs medium. Cells were submitted to daily media changes till day 5 (120h), when they were further analyzed.

Embryoid bodies differentiation from mESCs

Embryoid bodies (EBs) differentiation was established as described in (Sroczynska et al. 2009). Briefly, mESCs were splitted twice in Serum/LIF medium before inducing differentiation. One well of 6-well plate was splitted in an entire 6-well plate in IMDM-ES medium for 48h. mESCs were then harvested, and they were rinsed twice with DPBS (Gibco; Cat #14190144). Cells were very well disaggregated into single-cells, and 1.2×10^4 cells/mL embryoid body differentiation (EB^{diff}) medium were resuspended to a total volume of 25mL of EB^{diff} medium. EBs were formed in suspension for 5 days (120h). At day 5, in order to elongate the differentiation up to 216h (day 9), EB^{diff} medium was refreshed by harvesting EBs with 10mL serological pipette, centrifuge at 200 g for 3 min and EBs were resuspended in fresh EB^{diff} medium for 4 more days.

Culture mediums composition are found in supplementary table 3.

CRISPR/Cas9 gene editing and cell lines generation in mESCs

The two guide RNAs (gRNAs) targeting the Nfkbia locus were designed using the CRISPR design tool from MIT (http://crispr.mit.edu). The best 2 gRNAs (based on on-target and off-target scores) that were targeting the exon 1 of Nfkbia (IκBα gene) were selected. After annealing, one of the gRNAs was cloned into SpCas9(BB)-2A-GFP (px458) plasmid (Addgene; Cat #48138), and the other gRNA was cloned into the px330-mCherry plasmid (modified from Addgene; Cat #98750 to incorporate mCherry reporter). mESCs were co-transfected with the two plasmids (px458-gRNA1 and px330-mCherry-gRNA2) by the use of Lipofectamine 200 following manufacturer's instructions (Invitrogen; Cat #11668019). GFP$^+$/mCherry$^+$ single cells

were FACS-purified. Mutant clones were screened through PCR and western blotting. Non-
targeting scrambled gRNAs were used as controls. Three independent clones from each
genotype (IκBα-WT and IκBα-KO) were selected for further experiments.

To increase IκBα specific detection, the IκBα endogenous gene (Nfkbia) was knocked-in with
3XFLAG sequence (DYKDHDGDYKDHDIDYKDDDDK) at the 3' end. To generate IκBα-3XFLAG
mESCs, 3XFLAG sequence was cloned into a plasmid containing Nfkbia left and right homology
arms flanking the endogenous stop codon; a spacer sequence of three glycines (GGCGGTGGG)
was also introduced. gRNAs and oligos sequences are included in supplementary table 1.

Doxycycline-inducible IκBα (IκBα^{WT}, IκBα$^{ΔNF-κB}$ and i-IκBα$^{ΔH2A/H4}$) mESCs were generated by
cloning the three different IκBα versions into a PiggyBac transposon system (Chen et al. 2020).
IκBα protein-coding cDNAs were PCR-amplified, and the T2A-EGFP fragment was amplified
from PX458 and incorporated into the 3' end. PB-TRE was used as backbone (Addgene, Cat#
63800). Pool of transfected IκBα-KO mESCs was selected by Hygromycin (100μg/mL) for 7
days. Cells were doubly screened by FACS sorting green fluorescent protein after 24h of
doxycycline (1μg/mL) treatment. Pool of cells were used for further experiments. Oligos
sequences are included in supplementary table 1.

Teratoma formation assay

IκBα-WT and IκBα-KO mESCs grown in Serum/LIF (5×10^5) were injected intramuscularly into
severe combined immunodeficient mice (NSG). In order to favor 3D aggregation of cells and
teratoma formation, cells were resuspended in Matrigel Matrix (BD; Cat #356234) prior to
injection. Six weeks later, mice with tumors were euthanized, and tumors were fixed in
formaldehyde, embedded in paraffin, sectioned and stained with hematoxylin and eosin for
histological analysis.

Immunofluorescence staining

For immunostaining of mESCs, cells were seeded on 0.1% gelatin-coated coverslips on 6-well
plates; for EB immunofluorescence, they were collected directly from the plate. Samples were
washed twice with 1X PBS, and they were fixed at 4ºC for 30 minutes with 4% PFA (Electron
Microscopy Sciences; Cat #15713S). For immunofluorescence of 5-Methylcytosine (5mC),
fixed cells were incubated with 2M Hydrochloric Acid (Sigma-Aldrich, Cat #H1758) for 20
minutes at room temperature. Samples were then washed twice with 1X Tris-buffered saline
(TBS; 50 mM Tris-Cl, pH 8) and permeabilized and blocked with 1X TBS supplemented with 1%
Triton-X100 (MERCK; Cat #9036-19-5) and 6% FBS (Biological Industries; Cat #04-001-1A) for
2h at 4ºC. Samples were washed twice with 1X TBS supplemented with 6% FBS. Primary
antibody incubation was performed in 1X TBS plus 6% FBS and 0.3% or 0.5% Triton-X100 (0.3%
for 2D culture and 0.5% for EBs) overnight at 4ºC. Samples were then washed 4 times with 1X
TBS, 5 min each washing. Secondary antibody incubation was performed in TBS 1X plus 1%
BSA (Sigma-Aldrich; Cat #9048-46-8) for 2h at room temperature. Samples were washed three

times with 1X TBS for 5 min at room temperature each rinse. Samples were mounted using
DAPI Fluoromount-G (Southern Biotech; Cat #0100-20). List of antibodies used is found in
Supplementary table 2.

Microscopy and image acquisition

491   Fluorescence images were acquired using Confocal Leica TCS SP5 (Leica Microsystems), and
492   Leica application software LAS AF (Leica Microsystems) was used to visualize the images. The
493   Image J (version 1.15) was used for further analysis (Schneider et al. 2012).

Flow cytometry sample preparation and analysis

Cells were collected and purified based on FACS using BD FACSAria II Cell Sorter (BD
Bioscience). Cells were sorted at 3,500 events/second, and at maximum flow rate of 4 and 85
μm nozzle. 5 μg/ml of DAPI (Biotium; Cat #BT-40043) was used as a viability dye. For flow
cytometry analysis of pluripotency exit, cells were disaggregated using TrypLE Express, and
they were incubated at 4ºC for 20 minutes with the antibody. BD LSRFortessa Cell Analyzer or
BD LSR II Flow Cytometer (BD Bioscience) were used for Flow Cytometry Analysis experiments.
Flow cytometry data were analyzed using FlowJo X v10.0.7 (BD Biosciences).

RNA isolation, cDNA synthesis and quantitative RT-PCR

Total RNA isolation from cells was performed using the RNeasy Plus Mini Kit (Qiagen; Cat
#74136) or RNeasy Micro Kit (Qiagen; Cat #74004) following manufacturer's instructions.
Amount of RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and
2 μg of total RNA was retro-transcribed using Transcriptor First Strand cDNA Synthesis Kit
(Roche; Cat #04897030001) following the manufacturer's instructions.

Quantitative RT-PCR was performed in triplicates for each sample, and SYBR Green I Master
Kit (Roche; Cat #04887352001) was used to carry out the reaction. qRT-PCR was performed
using the LightCycler 480 system (Roche). Relative expression levels were calculated as $2^{-\Delta CT}$
normalized with the average CT of the housekeeping gene *Tbp* or *Gapdh*. Oligos sequences
are found in supplementary table 1.

Chromatin Immunoprecipitation (ChIP)

$4\text{-}6\text{x}10^7$ mESCs were cross-linked by incubating them in DPBS (Gibco, Cat #14190094)
supplemented with 1% formaldehyde (Sigma-Aldrich; Cat #252549) for 10 min rocking at
room temperature. Cross-linking was stopped using 125 mM Glycine (Sigma-Aldrich, Cat
#G8790) rocking for 5 minutes at room temperature. Fixed cells were then washed twice with
ice-cold DPBS, and scrapped and collected using DPBS supplemented with
protease/phosphatase inhibitor cocktail composed by 1X cOmplete™, EDTA-free Protease
Inhibitor Cocktail (Roche, Cat #11873580001), 1 mM PMSF, 1mM Sodium Ortovanadate and
20 mM β-Glycerol phosphate. Cells were centrifuged at 3200g for 5 minutes at 4ºC. Cells were

lysed by resuspending cell pellet in ice-cold ChIP buffer (1 volume of SDS buffer [100mM NaCl, 50mM Tris-HCl pH 8.1, 5mM EDTA pH 8, 0.5% SDS] and 0.5 volume of triton dilution buffer [100mM Tris-HCl pH 8.6, 100mM NaCl, 5mM EDTA pH 8, 5% Triton X-100]) supplemented with protease/phosphatase inhibitor cocktail, and samples were sonicated using sonication beads (Diagenode; Cat #C01020031) and the Bioruptor Pico Sonicator (Diagenode; Cat #B01060010) for 20 cycles (each cycle 30 seconds on/30 seconds off) or till DNA fragments have 100-300bp size. Samples were centrifuged at 16000g for 20 minutes at 4ºC. Chromatin was quantified, and 30 µg of chromatin was incubated with every 5µg of antibody rotating for 16 hours (or overnight) at 4ºC. Samples were pulled down by incubation with Protein A-Sepharose™ CL-4B (previously hydrated and blocked with 0.05% BSA) for 3h at 4ºC in rotation. Sample was washed with a low salt washing buffer (50mM HEPES pH 7.5, 140mM NaCl and 1% Triton) three times, one wash with a high salt wash buffer (50mM HEPES pH 7.5, 500mM NaCl and 1% Triton) and one wash with TE pH 8 (10mM Tris-HCl pH8, 1mM EDTA). Samples were then eluted in 1% SDS and 100mM NaHCO$_3$. Samples were de-crosslinked by incubating for 16 hours (or overnight) at 65ºC shaking at 450 rpm, and were treated with Proteinase K for 1h at 45ºC. DNA was purified using QIAquick PCR Purification Kit (Qiagen, Cat #28106), following the manufacturer's instructions.

For ChIP IκBα-3XFLAG, a previous cross-linking step using Gold (Diagenode, Cat #C01019027) was performed following manufacturer's instructions, and cells (IκBα$^{3XFLAG/3XFLAG}$ and IκBα-WT mESCs) were later treated with 1% formaldehyde as mentioned above. Cell nuclei were isolated by incubating cell pellets in a buffer containing 5mM Pipes pH 8, 85mM KCl and 0.5% NP40, and later incubated on ice for 10 minutes followed by centrifuge at 4000 rpm for 5 minutes at 4ºC. Nucleus were lysed in a buffer containing 1% SDS, 10mM EDTA and 50 mM Tris-HCl pH 8.1 and incubated on ice for 10 min. Samples were sonicated as mentioned above and SDS was removed by incubating the sonicated sample on ice for 1 hour and later centrifuged for 15 at 4ºC and 16000g. 60µg of chromatin was incubated with 10 µg of antibody. Samples were pulled down by a 50/50 mix of Dynabeads A (Invitrogen, Cat #10002D) and G (Invitrogen, Cat #10003D). Samples were washed 4 times in 0.1% SDS, 1% Triton-X, 2mM EDTA, 20mM Tris-HCl pH 8 and 150mM NaCl; followed by 1 wash in TE pH 8. For ChIP followed by PCR, 1 µL of ChIP sample was normalized by its input. Oligos used are found in supplementary table 1. For ChIP followed by sequencing (ChIP-seq) Purified DNA concentration and integrity were determined using Agilent Bioanalyzer (Agilent Technologies; Cat #G2939BA). Libraries were prepared using standard protocols. Chromatin was sequenced using Illumina HiSeq platform (Illumina, Inc.) (50bp single-end reads). List of antibodies used is found in Supplementary table 2.

ChIP-seq data analysis

Quality control was performed on raw data with FASTQC tool. Raw reads were trimmed to remove adapters presence with Trimgalore (v0.6.6) (Ewels et al. 2012). Default parameters were used except for a minimum quality of 15 (Phred score) and an adapter removal

stringency of 3bp overlap. For H3K4me3, H3K27me3, H3K4me1 and IκBα-3XFLAG or IκBα-WT data, trimmed reads were aligned to reference genome with Bowtie2 (v2.4.4) which was executed with default parameters (Langmead and Salzberg 2012). Required genome index was built with corresponding GRCm38 fasta file retrieved from Ensembl (http://ftp.ensembl.org/pub/release-102/). Multimapped reads and those exhibiting MAPQ < 20 were removed. For H3K9me3 data, trimmed reads were aligned to reference genome (GRCm38 release 102 from Ensembl) with ShortStack (v3.8.5) (Axtell 2013). Randomly placed multimappers were removed from the mapped reads. For all cases, duplicated reads were marked with SAMtools (v1.15) (Danecek et al. 2021). Drosophila Melanogaster spike-ins, present in H3K27me3, were discarded. NSC and RSC quality metrics were computed with PhantomPeakQualTools (v1.2) (Landt et al. 2012). ENCODE blacklisted regions (mm10 v2) were removed prior to peak calling except for H3K9me3 (Amemiya et al., 2019). BigWig files were individually generated using deepTools (v3.5.1) bamCoverage with -ignoreDuplicates -binSize 10 -smoothLength 30 -effectiveGenomeSize 2308125349 -normalizeUsing RPGC and -extendReads *Fragment_Length* options (Ramírez et al. 2016). The effective genome size was particularized to a read length of 50bp and directly retrieved from deepTools web site (https://deeptools.readthedocs.io/en/develop/content/feature/effectiveGenomeSize.html). *Fragment_Length* was retrieved from PhantomPeakQualTools results except for H3K9me3 where 150bp value was considered. Duplicates were kept in IκBα-3XFLAG for BigWig files generation. For all histone marks except for H3K4me3, peak calling was conducted by means of epic2 (v0.0.52) with -effective-genome-fraction 0.8452 -fragment-size *Fragment_Length* options and chromosome sizes only referring to canonical chromosomes (Stovner and Sætrom 2019). For H3K4me3 and IκBα-3XFLAG/WT, MACS2 (v2.2.7.1) was used to identify peaks with -nomodel -extsize *Fragment_Length* -g 2308125349 options (Zhang et al. 2008). The corresponding input sample was used in all peak calling computations. For the histone marks, peaks were called with adjusted p-values (FDR) < 0.05. In case of IκBα-3XFLAG/WT samples, peaks were called with adjusted p-values (FDR) < 0.05 and a minimum 4-fold enrichment.

Peak annotation was performed with ChIPseeker R (v4.2.1) package (v1.34.1) considering a TSS region range between 5000 bp upstream and 100 bp downstream (Yu et al., 2015). Required TxDb object was generated from the Ensembl GRCm38 gtf file retrieved (http://ftp.ensembl.org/pub/release-102/). A consensus peakset was obtained from the three biological replicates per histone mark and condition (IκBα-WT or IκBα-KO) by identifying overlapping peaks in at least two out of the three replicates. In case of IκBα ChIP-seq data, peaks from consensus peakset identified from the -3XFLAG samples not overlapping with IκBα-WT were considered the high-fidelity peaks for this experiment. For this purpose, findOverlapsOfPeaks function from ChIPpeakAnno R package (v3.32.0) was used with default parameters (Zhu et al. 2010).

Differential binding analysis (DBA) was conducted with DiffBind (v3.8.4) per histone mark (Stark et al. 2011). Default parameters were used except for the summits parameter which

was set to 500bp for all histone marks except for H3K4me3 (150bp) in order to consider, for
testing, intervals of 1000bp or 300bp respectively. Summit values were selected to have
interval widths between the minimum and first quartile peak width values for each histone
mark. EdgeR was the statistical method used for all the three analyses. Differentially bound
regions (DBRs) between IκBα-KO and IκBα-WT samples were called with adjusted p-values
(FDR) < 0.05. Identified DBRs with any annotated gene were plotted with EnrichedHeatmap R
package (v1.18.1) (Gu et al. 2018). To summarize replicates per condition, normalizeToMatrix
function was used based on corresponding BigWig files in w0 mean mode in 50bp windows.
Tracks visualization were obtained by means of Integrative Genomics Browser (IGV) tool
(Robinson et al. 2011).

Enhancers activity identification

Consensus peaksets derived from the three H3K27ac and H3K4me1 IκBα-WT mESCs replicates
were used to identify poised and active enhancers in an IκBα-WT scenario. Putative poised
enhancers were defined as those regions with H3K4me1 but no H3K27ac peaks. Putative
active enhancers were defined as those regions with H3K27ac and presence/absence
H3K4me1 peaks. Putative gained or lost poised/active enhancers in IκBα-KO were obtained
with following criteria: (i) for gained/lost poised enhancers: a differential H3K4me1
increase/decrease and absence of H3K27ac peaks in IκBα-KO consensus peakset, (ii) for
gained/lost active enhancers (without H3K4me1): a differential H3K27ac increase/decrease
and absence of H3K4me1 in IκBα-KO consensus peakset and (iii) for gained/lost active
enhancers (with H3K4me1: a differential H3K27ac increase/decrease and presence of
H3K4me1 in IκBα-KO consensus peakset.

A maximum gap of 1000bp was allowed for checking the overlap between two different
histone marks. For this purpose, the 'subsetByOverlaps' function from the IRanges R package
was used (v.2.34.1).

Hi-C data visualization and TADs annotations

Juicebox v1.9.8 was run locally to visualize Hi-C data from (Bonev et al. 2017), specifically in
the mESCs (Serum/LIF) dataset. Corresponding maps from this dataset were already included
in the tool. To visualize contact maps, balanced normalization was considered and applied
color range up to 3 in the selected regions. Additionally, the list of 2,712 TADs with their
corresponding genome coordinates was retrieved and used to locate the IκBα peaks,
enhancers and showing differential activity together with differentially expressed genes from
in-house RNAseq data.

RNA-seq experiments

635 Total RNA from three independent clones from mESCs, 48h EBs and 96h EBs was isolated
636 using the RNeasy Plus Mini Kit (Qiagen; Cat #74136) following manufacturer's instructions.

RNA was quantified with Nanodrop (Thermo Fisher; Cat #ND2000CLAPTOP), and RNA integrity was addressed by Agilent Bioanalyzer (Agilent Technologies; Cat #G2939BA). Libraries sequenced using Illumina HiSeq 2500 (Illumina, Inc.) (125bp paired-end reads). Samples sequencing depth ranged between 35M and 52M reads (average 41M reads) per sample.

RNA-seq data analysis

Quality control was performed using FASTQC tool (v0.11.9). Raw reads were trimmed with Trimgalore (v0.6.6) (Ewels et al. 2012). Default parameters were used except for a minimum quality of 15 (Phred score) and an adapter removal stringency of 3bp overlap. Trimmed reads were aligned to reference the genome with STAR aligner tool (v2.7.8). STAR was executed with default parameters except for the number of allowed mismatches which was set to 1. Genome index was built with corresponding GRCm38 gtf and fasta files retrieved from Ensembl (http://ftp.ensembl.org/pub/release-102/). Obtained BAM files with uniquely mapped reads were considered for further analysis. Raw gene expression was quantified using featureCounts tool from subRead software (v2.0.1) with exon as feature (Liao et al. 2014). The raw counts matrix was imported into the R Statistical Software environment (v4.2.1) for downstream analysis. Experimental design considered three timepoints: mouse ESCs, EBs at 48h and EBs at 96h. Each time point included 6 samples distributed in two conditions: 3 IκBα-WT and 3 IκBα-KO. Prior to statistical analysis, those genes with less than 10 raw counts across the 6 samples under test were removed. After pre-filtering, 21,843 genes (mESCs), 21,424 genes (EBs 48h) or 21,664 genes (EBs 96h) were available for testing. For visualization purposes, counts were normalized by the variance-stabilizing transformation method as implemented in DESeq2 R package (Love et al. 2014) (v1.38.3). Differential expression analysis (DEA) was conducted with DESeq2. Each time point was independently analyzed. Fitted statistical model included sample conditions as covariable with IκBα-WT as the reference. Obtained $\log_2$ fold change values were shrunken with apeglm shrinkage estimator R package (v1.20.0) (Zhu et al. 2019). Raw p-values were adjusted for multiple testing using the Benjamini-Hochberg False Discovery Rate (FDR) (Benjamini and Hochberg 1995). Differentially Expressed Genes (DEGs) between IκBα-KO and IκBα-WT samples were called with adjusted p-values (FDR) < 0.05 and absolute shrunken log2 Fold change > 1. Data visualization was performed with the ggplot2 (v3.4.1).

Genome-wide DNA methylation samples preparation

DNA from frozen mESCs pellets was extracted using DNeasy Blood and Tissue Kit (Qiagen GmbH, Hilden, Germany). Purified genomic DNA was quantified with Qubit (Invitrogen, Carlsbad, CA, USA) according to manufacturer's instructions. Infinium Mouse Methylation BeadChip (Illumina, Inc., San Diego, CA, USA) arrays were used to profile DNA methylation genome wide. Samples were bisulfite converted using EZ DNA Methylation-Gold™ Kit (Zymo

674 Research, CA, USA) and were hybridized in the array following the manufacturer's
675 instructions.

DNA methylation data analysis

The DNA methylation profile of the studied samples was assessed using the Infinium Mouse
Methylation BeadChip Array (~285,000 methylation sites) as previously described (Garcia-
Prieto et al. 2022). Quality control steps to remove erratic probe signals were performed
within the R statistical environment (v4.0.3). We removed probes with detection p value >
0.01, genotyping probes and manufacturing flagged (MFG) probes described in the Illumina
manifest file (https://support.illumina.com/downloads/infinium-mouse-methylation-
manifest-file.html). The differentially methylated probes (DMPs) between IκBα-WT and IκBα-
KO samples were computed by deriving a linear model with the limma R package (v3.46.0).
Each condition included 3 samples per time point. DMPs with adjusted p-value (FDR) <0.05
and absolute mean methylation beta value difference between conditions >0.3 were
considered significant. DNA methylation analysis was performed using the mm10 mouse
genome reference build and the complete annotation was downloaded from the annotated
manifest file (http://zwdzwd.github.io/InfiniumAnnotation#mouse) (Zhou et al. 2022).

Functional analysis

Overrepresentation analysis was applied over lists of selected genes derived from RNA-seq
data (DEGs) or from ChIP-seq data (differentially bound regions). The Gene Ontology
(Biological Process ontology, GO BP terms), KEGG PATHWAY and WikiPathways databases for
Mus Musculus (Ashburner et al. 2000; Kanehisa and Goto 2000; Martens et al. 2021) were
interrogated by means of clusterProfiler R package (v4.6.2) (Wu et al., 2021). Corresponding
Entrez identifiers were used. Benjamini-Hochberg procedure was used to obtain adjusted p-
values. Obtained GO BP terms were simplified using the simplify function from clusterProfiler
with default parameters. Overrepresented terms or processes were called with adjusted p-
values (FDR) < 0.05.

Gene Set Enrichment Analysis (GSEA) was performed for mESCs against the three cell states
signatures defined by (Ghimire et al. 2018). Complete list of genes from mESCs samples
(21,843 genes) was ranked based on the shrunken log2 Fold Change obtained from DEA. GSEA
was conducted through the fgseaMultilevel function from fgsea R package (v1.24.0)
(Korotkevich et al. 2021) with default parameters. Enrichment plots were generated with the
same package.

Gene Set Variation Analysis (GSVA) R package (v1.46.0) with default parameters were used to
obtain Z-score values for genes annotated to endoderm (GO:0001706), mesoderm
(GO:0001707) and ectoderm (GO:0001705) GO BP terms (Hänzelmann et al. 2013).
709

Statistical analysis

Statistical analysis was performed with GraphPad Prism v.8.0.1. (GraphPad Software, Inc.). Unless specified, the comparison between two groups was performed with unpaired two-sided t-test. A p-value < 0.05 was considered significant.

DATA AVAILABILITY

Data is deposited at GEO under a SuperSeries with the accession number GSE239565)[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE239565]. Individual SubSeries can be found at: GSE239563 (RNA-seq data), GSE239564 (ChIP-seq data) and GSE239562 (Methylation array data).

CODE AVAILABILITY

Scripts used to process the generated bulk RNA-seq and ChIP-seq data are available in Github repository: https://github.com/BigaSpinosaLab/IkBa_Naive_Pluripotency

COMPETING INTEREST STATEMENT

Authors declared no potential conflicts of interest.

ACKNOWLEDGEMENTS

We would like to acknowledge all members of the Bigas/Espinosa labs for helpful discussions. We are grateful for technical support to CRG/UPF Flow Cytometry Unit, CRG Genomics and Advanced Light Microscopy Units. This work has been supported by Spanish Ministry of Science and Innovation (PID2019-104695RB-I00, PLEC2021-007518, PDC2021-120817-I00), Generalitat de Catalunya (2021SGR/39) and Departament de Salut (SLT002/16/00299) to AB; the Spanish Ministry of Science and Innovation (PID2019-108322GB-100 and PID2022-142679NB-I00) to LDC; the Spanish Ministry of Science and Innovation (PID2021-123383NB-I00.) and the Agencia de Gestió d'Ajuts Universitaris i de Recerca (2021 SGR 01222) to BP. L.G.P has been a recipient of FI AGAUR fellowship (2019 FI-B 00151/2020 FI_B1 00130) from Generalitat de Catalunya. D.A-V was funded by the FIS fellowship (FI20/00130) from Instituto Carlos III. M.B. received funding from the Ramón Areces Foundation. M.M. is a recipient of a grant from the Instituto Carlos III, grant number CA22/00011 (co-funded by the European Social Fund Plus, ESF+ and by the European Union).

Author contributions

L.G.P, L.E and A.B designed the study. L.G.P, D.A-V, M.B, A.I, J.G, J.B, D.A.E, C.A.G.P, C.B, V.R.C and A.V performed experiments. L.G.P, M.M, M.B and C.A.G.P performed bioinformatics analysis. L.G and A.B wrote the manuscript with input from all authors. C.B, A.V, A.V, P.M,

742 L.D.C, G.S, B.P, M.E and L.E provided input on experimental design. L.E and A.B provided
743 resources and supervised the research.

REFERENCES

Aguilera C, Hoya-Arias R, Haegeman G, Espinosa L, Bigas A. 2004. Recruitment of IκBα
to the *hes1* promoter is associated with transcriptional repression. *Proceedings of the*
National Academy of Sciences **101**: 16537–16542.

Alvarez-Villanueva D, Galan L, Bertran J, Floor M. 2023. Separation-of-function mutants
reveal the NF-κB-independent involvement of IκBα in the regulation of stem cell and
oncogenic programs. *bioRxiv*.
https://www.biorxiv.org/content/10.1101/2023.06.21.545928.abstract.

Álvarez-Villanueva D, Galán-Palma L, Bertran J, Floor M, Solé L, Lobo-Jarne T, Maqueda
M, Rajbhandari R, Marruecos L, Villà-Freixa J, et al. 2023. Separation-of-function
mutants reveal the NF-κB-independent involvement of IκBα in the regulation of stem
cell and oncogenic programs. *bioRxiv* 2023.06.21.545928.
https://www.biorxiv.org/content/10.1101/2023.06.21.545928v1 (Accessed July 17,
2023).

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K,
Dwight SS, Eppig JT, et al. 2000. Gene ontology: tool for the unification of biology. The
Gene Ontology Consortium. *Nat Genet* **25**: 25–29.

Atlasi Y, Jafarnejad SM, Gkogkas CG, Vermeulen M, Sonenberg N, Stunnenberg HG.
2020. The translational landscape of ground state pluripotency. *Nat Commun* **11**: 1617.

Axtell MJ. 2013. ShortStack: comprehensive annotation and quantification of small RNA
genes. *RNA* **19**: 740–751.

Bao S, Tang F, Li X, Hayashi K, Gillich A, Lao K, Surani MA. 2009. Epigenetic reversion of
post-implantation epiblast to pluripotent embryonic stem cells. *Nature* **461**: 1292–
1295.

Basak S, Kim H, Kearns JD, Tergaonkar V, O'Dea E, Werner SL, Benedict CA, Ware CF,
Ghosh G, Verma IM, et al. 2007. A fourth IkappaB protein within the NF-kappaB
signaling module. *Cell* **128**: 369–381.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J R Stat Soc Series B Stat Methodol* **57**: 289–
300.

Bleckwehl T, Crispatzu G, Schaaf K, Respuela P, Bartusel M, Benson L, Clark SJ, Dorighi
KM, Barral A, Laugsch M, et al. 2021. Enhancer-associated H3K4 methylation
safeguards in vitro germline competence. *Nat Commun* **12**: 5771.

Bonev B, Mendelson Cohen N, Szabo Q, Fritsch L, Papadopoulos GL, Lubling Y, Xu X, Lv
X, Hugnot J-P, Tanay A, et al. 2017. Multiscale 3D Genome Rewiring during Mouse
Neural Development. *Cell* **171**: 557–572.e24.

Boroviak T, Loos R, Lombard P, Okahara J, Behr R, Sasaki E, Nichols J, Smith A, Bertone

P. 2015. Lineage-Specific Profiling Delineates the Emergence and Progression of Naive
Pluripotency in Mammalian Embryogenesis. *Dev Cell* **35**: 366–382.

Brena D, Bertran J, Porta-de-la-Riva M, Guillén Y, Cornes E, Kukhtar D, Campos-Vicens L,
Fernández L, Pecharroman I, García-López A, et al. 2020. Ancestral function of
Inhibitors-of-kappaB regulates Caenorhabditis elegans development. *Sci Rep* **10**: 16153.

Brons IGM, Smithers LE, Trotter MWB, Rugg-Gunn P, Sun B, Chuva de Sousa Lopes SM,
Howlett SK, Clarkson A, Ahrlund-Richter L, Pedersen RA, et al. 2007. Derivation of
pluripotent epiblast stem cells from mammalian embryos. *Nature* **448**: 191–195.

Buecker C, Srinivasan R, Wu Z, Calo E, Acampora D, Faial T, Simeone A, Tan M, Swigut T,
Wysocka J. 2014. Reorganization of enhancer patterns in transition from naive to
primed pluripotency. *Cell Stem Cell* **14**: 838–853.

Chen Q, Luo W, Veach RA, Hickman AB, Wilson MH, Dyda F. 2020. Structural basis of
seamless excision and specific targeting by piggyBac transposase. *Nat Commun* **11**:
3446.

Choi J, Huebner AJ, Clement K, Walsh RM, Savol A, Lin K, Gu H, Di Stefano B,
Brumbaugh J, Kim S-Y, et al. 2017. Prolonged Mek1/2 suppression impairs the
developmental potential of embryonic stem cells. *Nature* **548**: 219–223.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools.
Gigascience **10**. http://dx.doi.org/10.1093/gigascience/giab008.

Di Stefano B, Luo E-C, Haggerty C, Aigner S, Charlton J, Brumbaugh J, Ji F, Rabano
Jiménez I, Clowers KJ, Huebner AJ, et al. 2019. The RNA Helicase DDX6 Controls Cellular
Plasticity by Modulating P-Body Homeostasis. *Cell Stem Cell* **25**: 622–638.e13.

Dutta D, Ray S, Home P, Larson M, Wolfe MW, Paul S. 2011. Self-renewal versus lineage
commitment of embryonic stem cells: protein kinase C signaling shifts the balance.
Stem Cells **29**: 618–628.

Espín-Palazón R, Traver D. 2016. The NF-κB family: Key players during embryonic
development and HSC emergence. *Exp Hematol* **44**: 519–527.

Ewels P, Afyounian E, Schuster-Boeckler BFK. 2012. TrimGalore: v0. 6.7-DOI via
Zenodo.(Zenodo, 2021). doi: 10.5281/zenodo. 5127899. 2. Langmead, B. & Salzberg, SL
Fast gapped-read alignment with …. *Nat Methods*.

Factor DC, Corradin O, Zentner GE, Saiakhova A, Song L, Chenoweth JG, McKay RD,
Crawford GE, Scacheri PC, Tesar PJ. 2014. Epigenomic comparison reveals activation of
"seed" enhancers during transition from naive to primed pluripotency. *Cell Stem Cell*
14: 854–863.

Festuccia N, Osorno R, Wilson V, Chambers I. 2013. The role of pluripotency gene
regulatory network components in mediating transitions between pluripotent cell

states. *Curr Opin Genet Dev* **23**: 504–511.

Garcia-Prieto CA, Álvarez-Errico D, Musulen E, Bueno-Costa A, N. Vazquez B, Vaquero
A, Esteller M. 2022. Validation of a DNA methylation microarray for 285,000 CpG sites
in the mouse genome. *Epigenetics* **17**: 1677–1685.

Gerondakis S, Grumont R, Gugasyan R, Wong L, Isomura I, Ho W, Banerjee A. 2006.
Unravelling the complexities of the NF-kappaB signalling pathway using mouse
knockout and transgenic models. *Oncogene* **25**: 6781–6799.

Ghimire S, Van der Jeught M, Neupane J, Roost MS, Anckaert J, Popovic M, Van
Nieuwerburgh F, Mestdagh P, Vandesompele J, Deforce D, et al. 2018. Comparative
analysis of naive, primed and ground state pluripotency in mouse embryonic stem cells
originating from the same genetic background. *Sci Rep* **8**: 5884.

Graf T, Stadtfeld M. 2008. Heterogeneity of embryonic and adult stem cells. *Cell Stem*
Cell **3**: 480–483.

Guo G, Yang J, Nichols J, Hall JS, Eyres I, Mansfield W, Smith A. 2009. Klf4 reverts
developmentally programmed restriction of ground state pluripotency. *Development*
136: 1063–1069.

Guo Y-L. 2019. The underdeveloped innate immunity in embryonic stem cells: The
molecular basis and biological perspectives from early embryogenesis. *Am J Reprod*
Immunol **81**: e13089.

Gu Z, Eils R, Schlesner M, Ishaque N. 2018. EnrichedHeatmap: an R/Bioconductor
package for comprehensive visualization of genomic signal associations. *BMC Genomics*
19: 234.

Habibi E, Brinkman AB, Arand J, Kroeze LI, Kerstens HHD, Matarese F, Lepikhov K, Gut
M, Brun-Heath I, Hubner NC, et al. 2013. Whole-genome bisulfite sequencing of two
distinct interconvertible DNA methylomes of mouse embryonic stem cells. *Cell Stem*
Cell **13**: 360–369.

Hänzelmann S, Castelo R, Guinney J. 2013. GSVA: gene set variation analysis for
microarray and RNA-seq data. *BMC Bioinformatics* **14**: 7.

Hayden MS, Ghosh S. 2008. Shared principles in NF-kappaB signaling. *Cell* **132**: 344–
362.

Hu S, Metcalf E, Mahat DB, Chan L, Sohal N, Chakraborty M, Hamilton M, Singh A, Singh
A, Lees JA, et al. 2022. Transcription factor antagonism regulates heterogeneity in
embryonic stem cell states. *Mol Cell* **82**: 4410–4427.e12.

Kaltschmidt C, Greiner JFW, Kaltschmidt B. 2021. The Transcription Factor NF-κB in
Stem Cells and Development. *Cells* **10**. http://dx.doi.org/10.3390/cells10082042.

Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*

Acids Res **28**: 27–30.

Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. 2021. Fast
gene set enrichment analysis. *bioRxiv* 060012.
https://www.biorxiv.org/content/10.1101/060012v3 (Accessed March 11, 2024).

Lando D, Ma X, Cao Y, Jartseva A, Stevens TJ, Boucher W, Reynolds N, Montibus B, Hall
D, Lackner A, et al. 2024. Enhancer-promoter interactions are reconfigured through the
formation of long-range multiway hubs as mouse ES cells exit pluripotency. *Mol Cell* **84**:
1406–1421.e8.

Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein BE,
Bickel P, Brown JB, Cayting P, et al. 2012. ChIP-seq guidelines and practices of the
ENCODE and modENCODE consortia. *Genome Res* **22**: 1813–1831.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**: 357–359.

Leitch HG, McEwen KR, Turp A, Encheva V, Carroll T, Grabole N, Mansfield W, Nashun
B, Knezovich JG, Smith A, et al. 2013. Naive pluripotency is associated with global DNA
hypomethylation. *Nat Struct Mol Biol* **20**: 311–316.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–930.

Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I,
Lajoie BR, Sabo PJ, Dorschner MO, et al. 2009. Comprehensive mapping of long-range
interactions reveals folding principles of the human genome. *Science* **326**: 289–293.

Li S, Sun J, Zhang B-W, Yang L, Wan Y-C, Chen B-B, Xu N, Xu Q-R, Fan J, Shang J-N, et al.
2024. ATG5 attenuates inflammatory signaling in mouse embryonic stem cells to
control differentiation. *Dev Cell*. http://dx.doi.org/10.1016/j.devcel.2024.01.026.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Marks H, Kalkan T, Menafra R, Denissov S, Jones K, Hofemeister H, Nichols J, Kranz A,
Stewart AF, Smith A, et al. 2012. The transcriptional and epigenomic foundations of
ground state pluripotency. *Cell* **149**: 590–604.

Marruecos L, Bertran J, Álvarez-Villanueva D, Mulero MC, Guillén Y, Palma LG, Floor M,
Vert A, Arce-Gallego S, Pecharroman I, et al. 2021. Dynamic chromatin association of
IκBα is regulated by acetylation and cleavage of histone H4. *EMBO Rep* **22**: e52649.

Marruecos L, Bertran J, Guillén Y, González J, Batlle R, López-Arribillaga E, Garrido M,
Ruiz-Herguido C, Lisiero D, González-Farré M, et al. 2020. IκBα deficiency imposes a
fetal phenotype to intestinal stem cells. *EMBO Rep* **21**: e49708.

Martens M, Ammar A, Riutta A, Waagmeester A, Slenter DN, Hanspers K, A Miller R,

Digles D, Lopes EN, Ehrhart F, et al. 2021. WikiPathways: connecting communities.
Nucleic Acids Res **49**: D613–D621.

Martinez Arias A, Brickman JM. 2011. Gene expression heterogeneities in embryonic
stem cell populations: origin and function. *Curr Opin Cell Biol* **23**: 650–656.

Mulero MC, Ferres-Marco D, Islam A, Margalef P, Pecoraro M, Toll A, Drechsel N,
Charneco C, Davis S, Bellora N, et al. 2013. Chromatin-bound IκBα regulates a subset of
polycomb target genes in differentiation and cancer. *Cancer Cell* **24**: 151–166.

Pantier R, Tatar T, Colby D, Chambers I. 2019. Endogenous epitope-tagging of Tet1,
Tet2 and Tet3 identifies TET2 as a naïve pluripotency marker. *Life Sci Alliance* **2**.
http://dx.doi.org/10.26508/lsa.201900516.

Pera MF, Rossant J. 2021. The exploration of pluripotency space: Charting cell state
transitions in peri-implantation development. *Cell Stem Cell* **28**: 1896–1906.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F,
Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**: W160–5.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.
2011. Integrative genomics viewer. *Nat Biotechnol* **29**: 24–26.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**: 671–675.

Singer ZS, Yong J, Tischler J, Hackett JA, Altinok A, Surani MA, Cai L, Elowitz MB. 2014.
Dynamic heterogeneity and DNA methylation in embryonic stem cells. *Mol Cell* **55**:
319–331.

Sroczynska P, Lancrin C, Pearson S, Kouskoff V, Lacaud G. 2009. In vitro differentiation
of mouse embryonic stem cells as a model of early hematopoietic development.
Methods Mol Biol **538**: 317–334.

Stark R, Brown G, Others. 2011. DiffBind: differential binding analysis of ChIP-Seq peak
data. *R package version* **100**.
https://www.andersvercelli.com/packages/2.13/bioc/vignettes/DiffBind/inst/doc/DiffB
ind.pdf.

Stovner EB, Sætrom P. 2019. epic2 efficiently finds diffuse domains in ChIP-seq data.
Bioinformatics **35**: 4392–4393.

Takahashi S, Kobayashi S, Hiratani I. 2018. Epigenetic differences between naïve and
primed pluripotent stem cells. *Cell Mol Life Sci* **75**: 1191–1203.

Tesar PJ, Chenoweth JG, Brook FA, Davies TJ, Evans EP, Mack DL, Gardner RL, McKay
RDG. 2007. New cell lines from mouse epiblast share defining features with human
embryonic stem cells. *Nature* **448**: 196–199.

Thambyrajah R, Fadlullah Z, Proffitt M, Neo WH, Guillén Y, Casado-Pelaez M, Herrero-
Molinero P, Brujas C, Castelluccio N, González J, et al. 2022. IκBα controls dormancy
induction in Hematopoietic stem cell development via retinoic acid. *bioRxiv*
2022.11.17.516971. https://www.biorxiv.org/content/10.1101/2022.11.17.516971v1
(Accessed July 11, 2023).

Torres J, Watt FM. 2008. Nanog maintains pluripotency of mouse embryonic stem cells
by inhibiting NFkappaB and cooperating with Stat3. *Nat Cell Biol* **10**: 194–201.

Wang X, Wu Q. 2022. The Divergent Pluripotent States in Mouse and Human Cells.
Genes **13**. http://dx.doi.org/10.3390/genes13081459.

Weinberger L, Ayyash M, Novershtern N, Hanna JH. 2016. Dynamic stem cell states:
naive to primed pluripotency in rodents and humans. *Nat Rev Mol Cell Biol* **17**: 155–
169.

Wu X, Wu X, Xie W. 2023. Activation, decommissioning, and dememorization:
enhancers in a life cycle. *Trends Biochem Sci* **48**: 673–688.

Yagi M, Kishigami S, Tanaka A, Semi K, Mizutani E, Wakayama S, Wakayama T,
Yamamoto T, Yamada Y. 2017. Derivation of ground-state female ES cells maintaining
gamete-derived DNA methylation. *Nature* **548**: 224–227.

Ying Q-L, Wray J, Nichols J, Batlle-Morera L, Doble B, Woodgett J, Cohen P, Smith A.
2008. The ground state of embryonic stem cell self-renewal. *Nature* **453**: 519–523.

Zhang Q, Lenardo MJ, Baltimore D. 2017. 30 Years of NF-κB: A Blossoming of Relevance
to Human Pathobiology. *Cell* **168**: 37–57.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers
RM, Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome*
Biol **9**: R137.

Zhou W, Hinoue T, Barnes B, Mitchell O, Iqbal W, Lee SM, Foy KK, Lee K-H, Moyer EJ,
VanderArk A, et al. 2022. DNA methylation dynamics and dysregulation delineated by
high-throughput profiling in the mouse. *Cell Genom* **2**.
http://dx.doi.org/10.1016/j.xgen.2022.100144.

Zhou W, Triche TJ Jr, Laird PW, Shen H. 2018. SeSAMe: reducing artifactual detection of
DNA methylation by Infinium BeadChips in genomic deletions. *Nucleic Acids Res* **46**:
e123.

Zhu A, Ibrahim JG, Love MI. 2019. Heavy-tailed prior distributions for sequence count
data: removing the noise and preserving large differences. *Bioinformatics* **35**: 2084–
2092.

Zhu LJ, Gazin C, Lawson ND, Pagès H, Lin SM, Lapointe DS, Green MR. 2010.
ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. *BMC*
Bioinformatics **11**: 237.

Figure 1. IκBα specifically accumulates in the chromatin of the naïve pluripotent state. A. Schematic of pluripotent states in mouse embryonic stem cells (mESCs). **B. C.** Relative mRNA levels of the canonical IκB genes (**B**) or Nf-κB effector genes (**C**). Expression was normalized using the house-keeping gene *Tbp* relative to 2i/LIF. Data from three independent experiments. Dots indicate mean values and error bars refer to ± standard deviation (SD). **D.** Western blot analysis of IκBα protein distribution in cytoplasm, nucleoplasm (Nucleus) or chromatin in cells at different pluripotency states. **E.** Transcriptome and proteome differences between 2i/LIF and EpiSCs. Enriched transcripts and proteins in naïve or primed states are highlighted in red or dark gray, respectively (Log$_2$ fold-change < -1 or >1, respectively). Data from (Atlasi et al. 2020). **F.** Heatmap showing the expression levels of pluripotency (*Dppa4, Pou5f1, Sox2, Prdm14, Klf4, Zfp42, Nanog*), differentiation (*Mesp1, Gata4, Gata6, Lhx1, Wnt3a, Fgf5, Nes, Axin2, Wnt3, Eomes, Foxa2, T, Mixl1, Lef1*) and NF-κB (*Nfkbia, Nfkbib, Nfkbie, Nfkbiz, Rela, c-Rel, Relb*) genes from the RNAseq samples in mESCs and differentiating cells (embryoid bodies) at 48h and 96h of IκBα-WT cells. **G.** Schematic to deplete IκBα protein in mESCs (upper panel) and western blot analysis of three independent IκBα-WT and IκBα-KO mESC clones (bottom panel). **H.** Boxplots, violin plots and individual values representing the normalized gene expression levels in IκBα-WT or IκBα-KO mESCs, EBs 48h and EBs 96h. Considered genes are those annotated to NF-κB signaling pathway (KEGG). Unpaired two-sided Wilcoxon test applied.

987   **Figure 2. Lack of IκBα impairs the exit from naïve pluripotency. A.** PC1 and PC2 from Principal
988   Component Analysis (PCA) of normalized RNA-seq data from mESCs, 48h EBs and 96h EBs of
989   IκBα-WT and IκBα-KO cells (three independent clones from each genotype). **B.** Expression

level of naïve pluripotency genes in IκBα-WT and IκBα-KO 96h EBs. **C, D, E**. Expression of endoderm (C), mesoderm (D) and ectoderm (E) genes from Gene Ontology (GO) database at mESCs, 48h and 96h EBs IκBα-WT or IκBα-KO cells (three independent clones). Loess curves are also represented with 95% confidence intervals. **F.** Representative immunofluorescence of 216h EBs stained with NANOG and OCT3/4. **G.** Quantification of total number of alkaline phosphatase (AP) colonies after 216h EB differentiation and additional culture of 96h in Serum/LIF (S/L) medium. Dots represent values from two independent experiments from three different IκBα-WT or IκBα-KO clones. Unpaired two-sided t-test applied. **H.** Schematics of mESCs differentiation to EpiSCs (Materials and Methods section for further information). **I.** Relative RNA levels of naïve pluripotency genes in IκBα-WT or IκBα-KO EpiSCs at 120h. RNA levels were normalized based on *Tbp* expression relative to IκBα-WT. Each dot represents an independent clone. Unpaired two-sided t-test applied. Error bars refer to ± SD.

Figure 3. IκBα-KO mESCs preserve the ground-state of naïve pluripotency under Serum/LIF culture. A. GSEA results comparing IκBα-KO vs IκBα-WT mESCs cultured in Serum/LIF against ground (2i/LIF) and naïve (Serum/LIF) state pluripotency signatures from (Ghimire et al. 2018). Adjusted p-value by Benjamini-Hochberg procedure and Normalized Enrichment Score (NES) indicated. **B.** Expression of the naïve pluripotency genes from the RNAseq samples of IκBα-KO and IκBα-WT mESCs cultured in Serum/LIF. **C.** Number of differentially bound regions (adjusted p-value <0.05, FDR) for H3K4me3, H3K27me3 and H3K9me3 in IκBα$^{-/-}$ vs IκBα$^{+/+}$ mESCs cultured in Serum/LIF. Enriched or reduced regions in IκBα $^{-/-}$ mESCs are distinguished (log$_2$ Fold-change >0 or <0, respectively). Overrepresented WikiPathways for (left) H3K4me3 and (center) H3K27me3, or GO terms for (right) H3K9me3 (adjusted p-value < 0.05, FDR) are indicated. A one-sided hypergeometric test was conducted. **D.** Representative naïve pluripotency genes having differential H3K4me3 (enriched) and H3K27me3 (reduced) levels

in IκBα $^{-/-}$ vs IκBα $^{+/+}$ mESCs cultured in Serum/LIF. Shadowed regions highlight the differential

levels of the histone marks. **E.** Representative immunofluorescence images of DNA-

methylation-related mark 5-methylcytosine (5mC) in IκBα-WT (left) and IκBα-KO (right)

mESCs cultured in Serum/LIF. **F.** DNA methylation profile in IκBα-WT vs IκBα-KO mESCs (left)

and 96h EBs (right) showing the distribution density of mean ß-values from all 261,220 CpGs.

G. DNA methylation levels in different genomic regions in either IκBα-WT vs IκBα-KO mESCs

or 96h EBs. Unpaired two-sided t-test was applied.

1023

1024

Figure 4. IκBα binds to enhancer regions to prevent their hyperactivation in the naïve pluripotency state. A. Distribution of IκBα chromatin binding peaks. **B, C.** H3K27Ac (B) and

H3K4me1 (C) distribution profiles in IκBα-WT vs IκBα-KO mESCs cultured in Serum/LIF in IκBα-
bound chromatin regions. **D.** Number of regions with differential enhancer activity in IκBα-KO
vs IκBα-WT mESCs cultured in Serum/LIF. Increased/Reduced activity for (left) poised
enhancers based on the H3K4me1 differential binding and absence of H3K27ac overlapping
peaks in IκBα-KO mESCs and, (right) active enhancers based on the differential binding of
H3K27ac and presence of H3K4me1 overlapping peaks in IκBα-KO. Differential binding based
on adjusted p-value <0.05 (FDR). Increased or reduced regions in IκBα-KO are distinguished
by $\log_2$ Fold-change >0 or <0, respectively. Overrepresented WikiPathways for annotated
genes with (left) reduced poised enhancers and (right) increased active enhancers in IκBα-KO
mESCs are indicated (adjusted p-value < 0.05, FDR). **E.** Representative naïve pluripotency
genes having an increase in active enhancer status in IκBα-KO mESCs. Shadowed regions
highlight the differential levels of the histone mark. **F.** Expression of selected naïve
pluripotency genes contained in TADs with active enhancer marks (H3K27Ac and H3K4me1)
and IκBα binding in IκBα-WT and IκBα-KO mESCs and 48h EBs. **G.** (Upper panel) Contact map
(Hi-C data from (Bonev et al. 2017)) of the TAD (chr5:118,785,000-119,707,000) which
contains the Tbx3 locus. Arrows indicate the interaction between the IκBα-bound region and
TSS of Tbx3 gene (upper arrow) or differential active enhancer (bottom arrow). (Center panel)
Genome browser with H3K27Ac and H3K4me1 data in IκBα-WT and IκBα-KO mESCs.
Shadowed areas indicate differential active enhancer (light gray), the TSS of the gene of
interest (darker gray) or IκBα peaks (dark green). (Bottom panel) *Tbx3* expression (RNA-seq
data) in IκBα-WT and IκBα-KO mESCs and 48h EBs from two independent clones. **H.** Schematic
representation of IκBα reconstitution of IκBα-KO mESCs. **I.** Western blot analysis of H3K27Ac
levels in i-IκBα^{WT} mESCs (left panel). Two independent experiments were performed to
quantify H3K27Ac levels (right panel).

Figure 5. IκBα safeguards enhancer hyperactivation in naïve pluripotency in an NF-κB-independent manner. A. Schematic of chromatin-specific/NF-κB-deficient (IκBα^{ΔNF-κB}) or NF-κB-specific/chromatin-deficient (IκBα^{ΔChrom}) IκBα mutants. **B.** Pull-down of GST-H2A with cell lysates from different SOF IκBα mutant mESCs. **C.** Pull-down of GST-p50 with lysates from SOF IκBα mutant mESCs. **D.** Western blot analysis of chromatin protein extracts of doxycycline-

inducible IκBα forms after 16h of doxycycline induction. **E.** Western blot analysis of H3K27Ac levels in i-IκBα^{WT}, i-IκBα$^{\Delta NF\text{-}\kappa B}$ and IκBα$^{\Delta Chrom}$ mESCs (upper panel). Quantification of H3K27Ac levels (bottom panel). **F, G.** ChIP-PCR of H3K27Ac (F) and H3K4me1 (G) marks within genomic regions bound by IκBα in i-IκBα$^{\Delta NF\text{-}\kappa B}$ and IκBα$^{\Delta Chrom}$ mESCs cultured in Serum/LIF. Unpaired two-sided t-test was performed. Bars indicate mean values and error bars refer to ± standard error of the mean (SEM). **H.** Representative immunofluorescence images of 5-methylcytosine (5mC) in i-IκBα$^{\Delta WT}$ (upper panel), i-IκBα$^{\Delta NF\text{-}\kappa B}$ (medium panel), IκBα$^{\Delta Chrom}$ (bottom panel) mESCs cultured in Serum/LIF. **I.** Flow cytometry of SSEA-1 at 216h EBs in i-IκBα^{WT}, i-IκBα$^{\Delta NF\text{-}\kappa B}$ and i-IκBα$^{\Delta Chrom}$ cells. **J.** Quantification of AP staining in 216h EBs upon reconstitution with the different IκBα forms. Each dot represents an independent experiment. One-way ANOVA was applied. Bars indicate mean values and error bars refer to ± SEM. **K.** Representative immunofluorescence for NANOG and OCT3/4 at 216h EBs in i-IκBα^{WT} (upper panel), i-IκBα$^{\Delta NF\text{-}\kappa B}$ (medium panel) and i-IκBα$^{\Delta Chrom}$ (bottom panel) cells.

1073

| Ac H3K27Ac | ■ Enhancer | ■ Pluripotency gene |

Figure 6. Proposed model: IκBα attenuates enhancer activity in the naïve-to-primed
pluripotency transition. In mESCs cultured in Serum/LIF, IκBα binds to distal regions to
maintain a moderate pluripotency gene expression by avoiding enhancer hyperactivation.
This allows naïve pluripotency exit and differentiation upon LIF withdrawal. In the absence of
IκBα, enhancer regions acquire a higher activation status, resulting in an aberrantly increased
expression of pluripotency genes. This leads to an impaired pluripotency exit and
differentiation.