Compositional editing of extracellular matrices by CRISPR/Cas9 engineering of human mesenchymal

stem cell lines

Sujeethkumar Prithiviraj [1,2,3], Alejandro Garcia Garcia [1,2,3], Karin Linderfalk [1,2,3], Bai Yiguang [4], Sonia

Ferveur [1,2,3], Ludvig Nilsén Falck [5], Agatheeswaran Subramaniam [5], Sofie Mohlin [6], David Hidalgo [1, 2,3],

Steven J Dupard[1, 2,3], Deepak Bushan Raina [7], Paul E Bourgine [1,2,3].

[1]Cell, Tissue & Organ engineering laboratory, BMC B11, 22184, Department of Clinical Sciences, Lund Stem

Cell Centre, Lund University, Lund, Sweden

[2]Wallenberg Centre for Molecular Medicine, Lund University, Lund, Sweden

[3]Lund University Cancer Centre, Lund University, Lund, Sweden

[4]Department of Orthopaedics, Nanchong central hospital, The second clinical Institute of North Sichuan

Medical college Nanchong, Sichuan, China

[5]Division of Molecular Medicine and Gene Therapy, Lund Stem Cell Centre, Lund University, 22184 Lund,

Sweden

[6]Division of Pediatrics, Clinical Sciences, Translational Cancer Research, Lund University Cancer Center at

Medicon Village, Lund, 22363, Sweden

[7]The Faculty of Medicine, Department of Clinical Sciences Lund, Orthopedics, Lund 221 84, Sweden

18
Abstract

Tissue engineering strategies predominantly rely on the production of living substitutes, whereby

implanted cells actively participate in the regenerative process. Beyond cost and delayed graft availability,

the patient-specific performance of engineered tissues poses serious concerns on their clinical translation

ability. A more exciting paradigm consist in exploiting cell-laid, engineered extracellular matrices (eECM),

which can be used as off-the-shelf materials. Here, the regenerative capacity solely relies on the

preservation of the eECM structure and embedded signals to instruct an endogenous repair. We recently

described the possibility to exploit custom human stem cell lines for eECM manufacturing. In addition to

the conferred standardization, the availability of such cell lines opened avenues for the design of tailored

eECMs by applying dedicated genetic tools. In this study, we demonstrated the exploitation of

CRISPR/Cas9 as a high precision system for editing the composition and function of eECMs. Human

mesenchymal stromal/stem cell (hMSC) lines were modified to knockout VEGF and RUNX2 and assessed

for their capacity to generate osteoinductive cartilage matrices. We report the successful editing of

hMSCs, subsequently leading to targeted VEGF and RUNX2-knockout cartilage eECMs. Despite the

absence of VEGF, eECMs retained full capacity to instruct ectopic endochondral ossification. Conversely,

RUNX2-edited eECMs exhibited impaired hypertrophy, reduced ectopic ossification and superior cartilage

repair in a rat osteochondral defect. In summary, our approach can be harnessed to identify the necessary

eECM factors driving endogenous repair. Our work paves the road towards the compositional eECMs

editing and their exploitation in broad regenerative contexts.

Introduction

Extracellular matrices (ECMs) are complex networks of proteins providing tissue structural and mechanical

support, but also acting as growth factor storing and presenting entities[1]. As such, ECMs are receiving

increasing attention in tissue engineering as templates capable of guiding homeostasis, remodeling, and

regenerative processes[2,3,4].

ECMs can be derived from native tissue or organs (native ECMs, nECMs) by applying a decellularization

step which effectively removes the cellular fraction. The resulting nECMs are commonly used as

biomaterials in regenerative medicine[5,6], offering a natural biocompatible structure. However, they suffer

from substantial batch-to-batch variation with their properties being largely affected by the tissue source

and the decellularization process[5,7]. Importantly, while displaying a high level of biological complexity and

fidelity, the composition of nECMs cannot be tailored to specific needs.

A valuable option emerged from recent advances in bioengineering which led to the development of

synthetic ECMs (sECMs). These tunable materials are predominantly composed of biopolymers such as

poly (lactic acid) (PLA), poly (glycolic acid) (PGA), poly (lactic co-glycolic acid) (PLGA) and Poly (ethylene

glycol) (PEG), which can be functionalized with bioactive substances such as peptides, growth factors or

enzymes[8,9,10,]. The possible modulation of sECM's composition and chemistry offers precise control over

their mechanical properties, degradability, and temporal release of instructive molecules towards

improved tissue repair. Despite holding great promises, yet their structure and function remain simplified

compared to their native counterpart [11,12].

With the aim of combining high biological fidelity and design flexibility, engineered ECMs (eECMs) have

become a credible alternative. Those result from the exploitation of stem/progenitor populations capable

of depositing ECMs under specific in vitro culture conditions[13,14,15,16], which can be subsequently isolated

from the cellular fraction and exploited as a cell-laid product. Early studies were conducted using eECM

derived from primary cells, exhibiting variable performance associated with the inter-donor variability[17,18].

Most recently, we demonstrated the possibility to standardize the production of eECMs through the

engineering of dedicated human mesenchymal stromal/stem cell (hMSC) lines[19]. Precisely, the

Mesenchymal Sword of Damocles – BMP2 (MSOD-B) line offered the generation of eECMs in the form of

human cartilage templates[20]. Those cartilage templates can be further devitalized, lyophilized, and stored

as an *off-the-shelf* tissue, retaining remarkable bone formation capacity by instructing endochondral

ossification[21,22].

The exploitation of cell lines as an unlimited cell source for eECM generation offers unprecedented

standardization in graft production and performance. Meanwhile, it also consists of a robust cellular

platform facilitating their further genetic engineering, to achieve the production of eECMs tailored in

composition and function. Towards this objective, CRISPR/Cas9 appears as a versatile and efficient gene-

editing tool. While this technology has been extensively used for genetic screening or disease

modeling[23,24,25], the possibility to edit hMSCs and study the compositional impact on deposited eECMs has

not been investigated so far.

In this study, we aim at demonstrating that CRISPR/Cas9 can be applied as an editing tool for the

engineering of custom eECMs. As proof-of-principle, our strategy relies on the CRISPR/Cas9-guided editing

of the MSOD-B line for knocking out Vascular endothelial growth factor (VEGF) and Runt related

transcription factor 2 (RUNX2), respectively a key pro-angiogenic protein and transcription factor critically

involved in the endochondral ossification pathway. The resulting hMSC lines will be evaluated for their

capacity to generate hypertrophic cartilage in vitro and instruct tailored skeletal tissue formation in vivo

as cell-free and lyophilized templates. If successful, our study is expected to validate the use of

CRISPR/Cas9 for eECMs edited in composition, while providing a novel platform towards decoding the

necessary molecular signals driving effective tissue repair.

Results

CRISPR/Cas9 editing of hMSC lines lead to efficient VEGF knockout in cartilage tissues.

VEGF is a known master regulator of angiogenesis[26] and a key mediator of endochondral ossification.

When reaching hypertrophy, chondrocytes highly express VEGF, prompting vasculature invasion and

subsequent osteoprogenitor recruitment. This was proven to be essential for cartilage template

remodeling into bone and bone marrow[27]. Here, we thus first aimed at engineering a hMSC line

CRISPR/Cas9-edited for VEGF knockout and evaluate the corresponding impact on cartilage and

endochondral bone formation (Figure 1A). To this end, specific guide RNAs (gRNAs) targeting different

regions of the VEGF gene but conserved across all isoforms of VEGF were designed based on a previously

established protocol[28]. Three gRNAs targeting exon 1 (VEGF_1.1, VEGF_1,2, VEGF1.3), one targeting exon

2 (VEGF_2.1), and one targeting exon 8 (VEGF_8.1) were designed (Figure 1B, Supplementary Figure 1A).

These gRNAs were cloned into the pU6-(BbsI)_CBh-Cas9-T2A-mCherry vector encoding the Streptococcus

pyogenes Cas9 (SpCas9) machinery. Following transfection in MSOD-B cells, single cell clones were sorted

based on the transient mCherry expression and expanded for characterization (Figure 1A). Out of 163

single clones, 14 could be successfully expanded (8.5 %). From these clones, five were retrieved from the

gRNA 1.1, six from gRNA 1.3 and three from gRNA 8.1. In order to assess a successful editing and directly

correlate it to a knockout of VEGF, an ELISA assay was performed to measure the concentration of VEGF

in the supernatant of expanded MSOD-B clones (Figure 1C). In non-edited cells (MSOD-B, control), 5000

pg/mL were secreted and detected by ELISA. From the successfully expanded clones, most of them

showed reduced but non-abolished secretion of VEGF. However, two clones exhibited undetectable levels

of VEGF, both resulting from the gRNA targeting the Exon1 of the VEGF gene. These clones were defined

as MSOD-BΔV1 and MSOD-BΔV2 and further characterized for their capacity to form cartilage.

To this end, clones were expanded and seeded on collagen scaffold and induced for three weeks towards

chondrogenic differentiation. As anticipated with the MSOD-B tissues, histological assessment revealed

the formation of a collagenous matrix (Masson's Trichrome) with cartilage features (Safranin-O, Figure

1D). Similarly, a successful chondrogenic differentiation and deposition of glycosaminoglycans could be

observed in the MSOD-BΔV1 (Figure 1D) constructs. A quantitative assessment (blyscan assay) confirmed

the comparable content in glycosaminoglycans in MSOD-B and VEGF-edited clones (Figure 1E).

Importantly, VEGF quantification in corresponding pellets validated the successful knockout of the

protein, barely detectable in the MSOD-BΔV1 and ΔV2 tissues (47.64 pg/pellet in edited clones versus

552.7 pg/pellet in MSOD-B, Figure 1F).

To further examine the potential impact of VEGF-editing on tissue formation, we performed

immunostaining analysis of the engineered cartilages. Confocal microscopy revealed strong deposition of

Collagen Type I (COL1) and Collagen Type X (COLX) in all samples, characteristic of mature hypertrophic

cartilage tissues (Figure 1G). However, while the immunostaining revealed the VEGF deposition in the

MSOD-B ECM, the protein could not be detected in the MSOD-BΔV1 samples (Figure 1G)

Taken together, this data indicates the successful generation of MSOD-B lines knocked-out in VEGF. The

MSOD-BΔV1 clones exhibited frank cartilage formation capacity with minimal level of VEGF. This validates

the use of CRISPR/Cas9 as a precision tool to edit the composition of cartilage tissue.

VEGF knockout cartilage tissues retain bone remodeling capacity despite reduced early-stage vascularization.

To evaluate the functional impact of VEGF knockout in engineered constructs, we assessed both the angiogenic and bone formation assays. Since the two clones exhibited similar in vitro tissue formation capacity, only the MSOD-BΔV1 was selected for further performance assessment.

We first performed the Chorioallantoic Membrane (CAM) assay as ex vivo assessment of angiogenic potential (Figure 2A). This assay provides insights into the biological activity and neovascularization potential of each construct, through quantitative evaluation of vascular density around engineered grafts After 4 days, we observed an extensive vessel formation in the periphery of both MSOD-B and MSOD-BΔV1 eECMs (Figure 2B). Quantification of vascular densities [29] suggested a reduced vessel formation in MSOD-BΔV1 samples but without reaching significance (Figure 2C and Supplementary Figure 2A). Thus, the CAM assay indicated that both MSOD-B and MSOD-BΔV1 cartilages retain angiogenic potentials.

We next investigate in vivo the impact of the VEGF knockout on the graft capacity to undergo endochondral ossification, by implanting cartilage grafts subcutaneously in an immunodeficient mouse model. Importantly, prior to implantation, tissues were lyophilized using a pre-established protocol[21]. This allowed us to assess the performance of the generated tissue itself, in line with the idea of developing off-the-shelf substitutes. Samples were extracted after two and six weeks as early and late development timepoints (Figure 2D). After two weeks in vivo, we first evaluated the angiogenic potential of tissues by quantitative confocal microscopy imaging of 100 μm thick sections stained for CD31, a well-defined vascular marker. This allowed us to evidence a dense vascular network in MSOD-B constructs, covering the entirety of the grafts (Figure 2E). Instead, MSOD-BΔV1 displayed a more limited vascularization, pre-dominantly at the tissue periphery (Figure 2E). Quantification confirmed these observations with MSOD-B exhibiting a significantly higher volume of vessels per total section volume (0.523 μm^3 versus 0.231 μm^3 for MSOD-B and MSOD-BΔV1 respectively, Figure 2F).

Using microtomography, we further assessed the amount of mineralized tissue formed in a temporal fashion. At two weeks, the formation of a cortical ring could already be observed in both MSOD-B and MSOD-BΔV1 tissues (Figure 2G). Quantifications revealed a similar bone/mineralized volume normalized to the total volume of the graft (BV/TV, Figure 2H, supplementary figure 2C and 2D) with 17 % and 16 % in MSOD-B and MSOD-BΔV1 samples respectively. At the six-week timepoint, trabecular structures could

be observed in reconstructed 3D scans from both groups (Figure 2G). Remarkably, histological analysis

confirmed the similar development of the tissues with frank bone formation (Masson´s trichrome) already

at two weeks post-implantation and minimal remnants of cartilage (Safranin O) (Figure 2I). At six weeks

both samples remodeled into fully mature bone organs, characterized by presence of cortical and

trabecular structures as well as a bone marrow tissue filling the cavity (Figure 2I).

These results indicate that the absence of VEGF in cartilage tissue can delay the early vascularization of

MSOD-BΔV samples. However, this did not prevent nor impact the remodeling of the lyophilized grafts

into bone and bone marrow tissues indicating that VEGF is non-essential in order to efficiently instruct

endochondral ossification.

RUNX2 knockdown does not prevent chondrogenic differentiation but impairs hypertrophy.

We next investigated whether the composition and thus function of a graft could be modified by editing

transcriptional factors involved in mesenchymal cell differentiation. Using CRISPR/Cas9, we thus targeted

the RUNX2, a known master regulator particularly important for chondrocyte differentiation and

hypertrophy[30,31]. We designed gRNAs targeting the coding regions involved in cell signaling integration,

activation, and inhibition domain which are conserved in both isoforms of RUNX2 (Figure 3A,

Supplementary Figure 3A).

One gRNA was designed for targeting exon 2 (RUNX2_2.1), one targeting exon 5 (RUNX2_5.1), two

targeting exon 6 (RUNX2_6.1 and RUNX2_6.2) and one targeting exon 8 (RUNX2_8.1). Similar to the VEGF

set-up, gRNAs were transfected in MSOD-B cells together with a pU6-(BbsI)_CBh-Cas9-T2A-mCherry

plasmid to ensure transfection efficiency and single cell sorting of positive clones. Out of 385 single clones,

62 could be successfully expanded, corresponding to a 16.1% efficiency. From these clones, five were

derived from the RUNX2_5.1 gRNA, 18 from RUNX2_6.1, 15 from RUNX26.2 and 24 from RUNX2_8.1.

To identify successfully edited clones, we first screened for those exhibiting a decrease in their RUNX2

protein expression using intra cellular flow cytometry. Among the 62 clones, 17 displayed a reduced

RUNX2 pattern as compared to the MSOD-B control (Figure 3B and Supplementary Figure 3B) and were

further sent for sequencing analysis. This led to the identification of two successfully edited clones,

demonstrating a point mutation in the exon 6 of RUNX2 gene (Supplementary Figure 3C). To confirm the

knockout impact on the transcription factor structure, a Western Blot analysis was conducted on cellular

extracts of in vitro cultured cells. MSOD-B cells exhibited intact RUNX2 proteins of 52 – 62 kDa[32,33](Figure

3C). Instead, the RUNX2-edited cells displayed truncated versions of 32 − 42 kDa, consistent with the

expected mRNA shortening.

These two clones were defined as MSOD-BΔR1 and MSOD-BΔR2 and further assessed for their ability to

form cartilage in vitro. Following 3D chondrogenic differentiation, samples were processed for histological

analysis. The presence of cartilage structures embedded in a rich collagenous matrix could be observed in

all MSOD-B, MSOD-BΔR2 (Figure 3D) and MSOD-BΔR1 tissues (Supplementary Figure 3D). Quantification

of glycosaminoglycans revealed a similar content in all the groups (Figure 3E, Supplementary Figure 3E).

Using immunostaining combined with quantitative confocal microscopy, the possible impact of RUNX2-

editing on tissue hypertrophy was further investigated. First, Collagen Type II (COL2) staining confirmed

the similar distribution of cartilage matrix across MSOD-B, MSOD-BΔR1 and MSOD-BΔR2 samples (Figure

3F and Supplementary Figure 3F). However, we observed a clear distinction in the Collagen Type X (COLX)

expression pattern, a specific marker of hypertrophy which was hardly detectable in MSOD-BΔR tissues

(Figure 3F and Supplementary Figure 3F). Subsequent quantification confirmed a significant reduction of

COLX volume in MSOD-BΔR1 (0.236 μm^3) and MSOD-BΔR2 (0.182 μm^3) compared to MSOD-B (0.81 μm^3)

(Figure 3G, Figure 3H, Supplementary figure 3G-H-I).

To further characterize the functional impact of the RUNX2 knock-down, the MSOD-B and MSOD-BΔR1

osteogenic differentiation capacity was assessed in vitro. After 3 weeks of culture in osteogenic medium

(or expansion medium, control), Alizarin Red staining revealed a s a marked absence of mineralization in

MSOD-BΔR1 culture, while in stark contrast with MSOD-B cells (Figure 3I). Quantitative PCR analysis

indicated a lower expression of COLX and ALPL in MSOD-BΔR1 (Figure 3J), key osteogenic-associated

genes, of interest, the expression level of RUNX2 was only partially decreased in MSOD-BΔR1, in line with

the truncated mRNA reducing but not abrogating qPCR primers binding probability.

In summary, we here report the successful RUNX2 knockout in MSOD-B lines using CRISPR/Cas9. RUNX2

did not impair chondrogenesis but prevented the tissue hypertrophy, as well as the osteogenic potential

of the cells.

RUNX2 knockout in cartilage tissues disrupts effective ectopic bone formation.

To assess the corresponding impact of RUNX2-edited cartilages on bone formation, MSOD-B tissues (as

control) and MSOD-BΔR1 were lyophilized and implanted subcutaneously in immunodeficient mice for

two to six weeks. From microtomography image reconstructions, we observed an evident reduction of

mineralization in MSOD-BΔR1 compared to MSOD-B (Figure 4A) already after two weeks. This qualitative

and visual difference in mineralization persisted after six weeks in vivo (Figure 4A). Subsequent quantifications confirmed these observations with a clear reduction in BV/TV as early as week 2 (16.46 % in MSOD-B and 7.01 % in MSOD-BΔR1) and persisting at week 6 (20.03% in MSOD-B and 4.25 % in MSOD-BΔR1) (Figure 4B). Histological analysis revealed an early bone formation at two weeks in both samples, but to a lower extent in the MSOD-BΔR1 group (Figure 4C). As anticipated, MSOD-B tissues underwent full remodeling after six weeks in vivo. In sharp contrast, the MSOD-BΔR1 only displayed a partial maturation with limited presence of bone and bone marrow.

Altogether, this indicates an incomplete remodeling of RUNX2 edited samples with significantly delayed cortical and trabecular structure formation. This correlates with the impaired hypertrophic phenotype in the corresponding in vitro samples.

RUNX2 knockout in cartilage tissues leads to better cartilage regeneration in a rat osteochondral defect.

In order to assess the performance of CRISPR-Cas9 edited eECMs in a relevant skeletal regenerative context, a proof-of-concept study in the rat osteochondral defect was performed. MSOD-B and MSOD-BΔR1 tissues were lyophilized and placed in sub-chondral defect in the distal femur of a rat. The defect consisted in a drill hole of 1mm diameter and 2mm depth (Figure 5A and supplementary figure 5A). Positive controls consisted of untreated healthy rats.

Following explantation, histological stainings and micro-tomography were performed on all sample groups. Hematoxylin and Eosin (H&E) and Masson's Trichrome indicated de novo bone formation in the defect area for both the MSOD-B and MSOD-BΔR1 groups (Figure 5B and supplementary figure 5B). The presence of fibrotic tissue as well as a reduced marrow compartment suggested an incomplete remodeling at that timepoint (Figure 5B). Upon Tartrate-resistant acid phosphatase (TRAP) staining, we observed elevated level osteoclastic activities in MSOD-B samples compared to MSOD-BΔR1 indicating reduced rate of active bone formation in MSOD-BΔR1 (Figure 5C). The micro-CT quantification revealed a 30% repair of the damaged bone in the defect for both MSOD-B and MSOD-BΔR1 groups (Figure 5D, Supplementary figure 5C). Of interest, a significantly higher trabecular separation was observed in MSOD-B and MSOD-BΔR1, confirming an ongoing bone remodeling process (Figure 5E).

While no statistical differences in bone formation could be identified between the two groups, the MSOD-BΔR1 tissue led to a superior regeneration of the cartilage area, as revealed by Safranin-O staining (Figure 5F) in the chondral zone (Figure 5F, a,b and c magnification). Interestingly, the MSOD-BΔR1 group exhibited a higher remnants of GAGs in the subchondral area (Figure 5F, d,e and f magnification).

Quantification of cartilage tissue within condyle surface area confirmed a poor-to-no repair in the MSOD-
B group (1.05%, Figure 5G). In sharp contrast, the MSOD-BΔR1 implanted eECMs initiated a chondral
regeneration reaching approximately 20.67% of the total healthy cartilage area. The integration and
regeneration potential of engineered constructs was further evaluated by performing a semi-quantitative
histological assessment following a pre-existing grading system[34] (Table 1 and Supplementary table 1).
This approach allows for the systematic evaluation of critical repair tissue parameters, offering a
comparative measure of the regenerative efficacy of the engineered constructs against healthy tissue
benchmarks. In the assessment of stained femur condyle sections, cellular morphology and matrix staining
of the MSOD-BΔR1 group (66% and 50%, respectively) were significantly better compared to the MSOD-
B one (33% and 16%, respectively). Although both constructs exhibited reduced cartilage thickness and
subchondral bone regeneration, MSOD-BΔR1 consistently outperformed the MSOD-B grafts (33.33%
versus 8.33% and 25% versus 16.66%). Surface regularity and integration with adjacent cartilage also
revealed better outcomes in MSOD-BΔR1 (41% and 50%) as opposed to MSOD-B (50% and 33%), further
indicating an enhanced regenerative potential of the MSOD-BΔR1 constructs.

In conclusion, while both grafts yielded comparable bone regeneration within the osteochondral defects,
the RUNX2 deletion in cartilage grafts led to a superior cartilage regeneration.

Discussion

We here report the possibility to edit the composition and function of eECM by CRISPR/Cas9 engineering
of human mesenchymal lines. VEGF knockout led to successful in vitro formation of cartilage with targeted
protein depletion. Upon implantation, this impacted the graft vascularization onset but not the
endogenous instruction of the endochondral program. In turn, RUNX2 knockout prevented cartilage
hypertrophy in vitro, significantly delaying ectopic bone and bone marrow formation in vivo. This strategy
was validated in a functional osteochondral defect model, whereby prevention of cartilage hypertrophy
by RUNX2 deletion led to superior cartilage regeneration.
A clear interdependency of angiogenesis and osteogenesis occurs upon bone formation[35,36]. For this
reason, VEGF-enrichment has been naturally proposed as a strategy to accelerate or increase bone graft
vascularization[37,38]. Our study demonstrates that VEGF knocked-out cartilage templates retained full
osteoinductive potential in a challenging ectopic environment. Those results are in sharp contrast with
intramembranous strategies[39,40], including our previous work[19], whereby grafts remain insufficient in

promoting complete bone remodeling even upon VEGF-enrichment. This suggests that the first stage of

host progenitor recruitment by our eECM is not VEGF dependent, although subsequent tissue

vascularization and remodeling may be orchestrated in a second step by endogenous cell secretion. Other

pro-angiogenic proteins embedded in the matrix may also have compensated for the lack of VEGF, such

as BMP-2[20] known to stimulate endothelial cell proliferation, migration, and differentiation[41,42].

Conversely, RUNX2 deletion was shown to fully prevent cartilage hypertrophy, in line with published

mouse models also reporting a lack of endochondral ossification and lethality at birth upon RUNX2

knockout[43,44]. Here, MSOD-BΔR cartilage templates still exhibited frank bone formation capacity, but with

cortical and trabecular structures largely reduced as compared to controls. Future work may clarify if this

strictly results from the absence of COLX, or if an additional specific cartilage matrix component (e.g.

matrix metalloproteinases) impaired the instruction of endochondral ossification.

The performance in a regenerative context was further evaluated in a rat osteochondral defect model[45].

We defined an early timepoint of 3-week as providing the opportunity to assess the early contribution of

the grafts to both the cartilage and bone tissue regeneration. Strikingly, only the MSOD-BΔR could

contribute to a superior chondrogenesis. We hypothesize that the absence of hypertrophic features

prevented the template degradation and favored its integration to native cartilage. Despite the repair

performance being limited to an approximate 20% of native cartilage restoration, this is remarkable in

light of the BMP-2 content in our ECM prompting endochondral ossification[46,47,48]. In fact, preventing the

in vivo remodeling of engineered cartilage templates remains challenging in the skeletal regeneration

field, with bone marrow MSCs (BM-MSCs) cartilage systematically reaching hypertrophy and subsequent

ossification. Our results warrant further studies comprising additional timepoints to decipher the repair

dynamic, as well as the long-term stability of newly formed tissue. It also prompts a comparison with other

recently proposed *off-the-shelf* strategies, based on ECM[49] or synthetic scaffold materials[50], in order to

comprehend the feasibility to exploit our grafts for stable cartilage / joint repair[51,52]. Taken together, while

the relevance of CRISPR/Cas9 eECMs in cartilage repair remains to be further validated, our study

demonstrates the relevance of genetically-edited ECMs in regenerative contexts.

CRISPR/Cas9 editing of human mesenchymal cells has previously been explored for gene and cell therapy

applications[53,54], for tailoring the immunomodulatory and/or differentiation capacity of engineered cells.

However, no studies have described the exploitation of editing strategies for the generation of eECMs,

whereby modified cells are absent from the final grafting product. Our work illustrate that this can be

achieved by direct targeting of secreted factors typically embedded upon ECM deposition, as

demonstrated with VEGF. Alternatively, we also propose the knockout of key transcription factors as a

strategy for custom eECM generation by impacting their tissue developmental/maturation stages.

A clear implication of our work lies in the possibility to decipher the necessary factors capable of

instructing de novo tissue formation. Those findings will be of high relevance for the design of eECMs

tailored in composition. Beyond bone repair, our study also bears high relevance in other regenerative

contexts. In fact, an exciting opportunity also lies in harnessing CRISPR/Cas9 for editing eECMs and tuned

their immunogenicity. Key inflammatory components could be turned down, leading to improved efficacy

of repair in line with the immuno-engineering principles[55,56] .

Importantly, our study describes the exploitation of eECMs in a lyophilized form, thus conferring an off-

the-shelf storage solution. Together with the standardization of production conferred by stable cell

sources, our concept offers exciting translational opportunities. In fact, instrumental to this work is the

use of dedicated hMSC lines. The genetic modification of primary cells is laborious, and their limited

lifespan ex vivo challenges their selection, characterization, and exploitation for tissue engineering

applications[57,58]. Here, the MSOD-B was harnessed as an unlimited cell source with robust differentiation

potential. Nonetheless, the editing of eECMs remains tedious in part due to the limited CRISPR/Cas9

efficiency and potential off-target effects, calling for a systematic clonal selection. In addition, while large

CRISPR/Cas9 screening can be performed in other stem cell systems[59,60], the critical cell mass required for

effective cartilage formation affects parallelization and leads to limited throughput. Nonetheless, after

the identification and characterization process, the resulting cell lines can be banked and used for

unlimited tissue manufacturing.

The MSOD-B line remains the only human cell source capable of priming endochondral ossification by

engineering living or cell-free grafts. This was demonstrated to be driven by the combined low dose of

BMP-2 embedded in the tissue (~40 ng/tissue) together with glycosaminoglycans and other thousands of

identified ECM proteins[20]. The BMP-2 amount is thus far below the typical amount used in sECMs

approaches for effective osteoinduction, falling in the microgram to milligram range[21]. This is one key

advantage of eECM graft, exhibiting a biological complexity that so far has not been matched by sECMs.

Ideally, the two approaches are complementary as the editing of eECMs could inform on the necessary

but sufficient factors driving effective regeneration. Those would in turn be embedded in a sECMs strategy

and offer a fast and possibly cost-effective solution.

To conclude, the present work offers a platform for decoding factors involved in tissue regeneration and

generate tailored eECM. Here, illustrated in the context of skeletal repair, our study may offer similar

opportunities in other regenerative situations.

Materials and Methods:

Cell expansion

MSOD-B cells and their modified progeny were cultured in a humidified incubator at 37 °C and 5% CO_2 using complete medium consisting of α-minimum essential medium (αMEM) supplemented with 10% fetal bovine serum, 1% HEPES, 1% sodium pyruvate, 1% penicillin-streptomycin-glutamine solution, and 5 ng/ml of fibroblast growth factor-2 (FGF-2) (all from Gibco). Cells were seeded at a density of 3200 cells/cm^2 until they reached 90% confluency. The medium was replaced twice a week.

Chondrogenic and osteogenic differentiation

MSOD-B cells and their modified progeny were harvested from culture flasks by adding Trypsin-EDTA (0.25%) (Gibco) and subsequently seeded on cylindrical Collagen Type I scaffold (Avitene™ Ultrafoam™ Collagen Sponge, Davol) of 6 mm in diameter and 3 mm in thickness at a density of 2×10^6 cells per scaffold in 12 well plates coated with 1 % agarose (Sigma) for chondrogenic differentiation and in a 12-well plate for osteogenic differentiation. Tissue constructs were cultured for three weeks in chondrogenic medium (DMEM supplemented with 1% penicillin-streptomycin-glutamine, 1% HEPES (1M), 1% sodium pyruvate (100 mM), 1% ITS (100x) (Insulin, Transferrin, Selenium) (all from Gibco), 0,47 mg/ml linoleic acid (Sigma), 0.12% bovine serum albumin (25 mg/mL) (BSA) (Sigma), 0.1 mM ascorbic acid (Sigma), 10^{-7} M dexamethasone (Sigma) and 10 ng/ml TGF-β3 (Novartis)). The cells in the 12 well plates were supplemented for three weeks with osteogenic (or expansion medium, control) differentiation medium (α-minimum essential medium (αMEM) with 10% fetal bovine serum, 1% HEPES (1M), 1% sodium pyruvate (100×10^{-3}M) and 1% penicillin-streptomycin-glutamine solution (100X), supplemented with 0.01M β-dexamethasone and 0.1M ascorbic acid). Media were replaced twice a week.

Lyophilization

After three weeks of chondrogenic differentiation, the tissue constructs were rinsed twice with PBS 7.2 (without calcium/magnesium Gibco), snap frozen in liquid nitrogen for five minutes and then lyophilized using a freeze dryer (Labconco) (-80 °C and 0.05 mbar) overnight. Thereafter, the lyophilized tissue constructs were stored at 4 °C.

Transfection of MSOD-B cells with gRNAs

MSOD-B cells were seeded at a density of 400,000 cells per well in 12 well plates to reach a minimum of 80% confluency the following day. The medium was replaced before transfection. The transfection was performed with a / ratio of 2 μL lipofectamine to 1 μg DNA. A DNA mix was prepared for each of the five gRNAs and the two plasmid controls. DNA mixtures were composed of 1 μg of plasmid, 2 μL of P3000

reagent and 50 µL of OptiMEM (Thermo-Fischer). A lipofectamine cocktail was prepared for all DNA

mixtures consisting of 24 µL of lipofectamine 3000 in 400 µL of OptiMEM. Lipofectamine and the DNA

mixtures were added at a 1:1 ratio. After 48h, cells were analyzed and mCherry positive clones were FACS

sorted as single cells in 96-well plates using ARIAIII (BD Biosciences). Successfully expanded clones were

then further characterized.

ELISA

VEGF protein content was measured in supernatant collected from cells seeded at 570 cells/cm^2 in T175

flask and cultured for 3 days. Content from engineered cartilage tissues was assessed following digestion

in RIPA buffer. The Quantikine ELISA kit for Human VEGF-A Immunoassay from the R&D System was used

according to the manufacturer instructions to determine protein concentration.

Intracellular flow cytometry

MSOD-B and MSOD-B ΔRUNX2 cells were trypsinized, fixed and permeabilized using the

Fixation/Permeabilization Kit (BD Biosciences). Following a blocking step in 10% normal donkey serum

(Sigma), cells were stained with a primary antibody against RUNX2 (Rabbit anti-human, Thermo-Fischer

PA5-82787) for one hour at room temperature. After primary antibody incubation, cells were washed and

stained with an Allophycocyanin (APC) labeled secondary antibody (Donkey anti-rabbit IgG DyLight™ 649,

Biolegend 406406) for 30-45 minutes at room temperature in 2% normal donkey serum. The Control

samples consisted of unstained cells (negative control) and cells incubated only with the secondary

antibody (secondary Ab control). Data were recorded on a BD LSRFortessa™ Cell Analyzer (BD

Biosciences). FCS files were analyzed using the FlowJo™ software (FlowJo LLC, 10.5.3, BD Biosciences).

Western blotting

MSOD-B and MSOD-B ΔRUNX2 cells were trypsinized and washed with ice cold PBS twice and lysed on ice

for 10 min in RIPA buffer (#10017003, Thermo Fisher Scientific) supplemented with 1× proteinase and

phosphatase inhibitor cocktail (#78440, Thermo Fisher Scientific). The lysates were centrifuged at 16,

000*g* for 15 min +4 °C, and the supernatants were collected. Sample buffer (Laemmli buffer (#161-0737,

BioRad) supplemented with 5% 2-mercaptoethanol, 1× proteinase and phosphatase inhibitor cocktail

(#78440, Thermo Fisher Scientific) was added to the supernatant at 1:1 ratio. Samples were boiled at 95

°C for 5 minutes and stored at −80 °C or kept on ice until gel loading. Proteins were separated using Bolt

gels according to manufacturer's protocol (#NW04122, #B0002, Thermo Fisher Scientific). iBlot2 system

was used to transfer the proteins on polyvinylidene fluoride membrane (PVDF) membrane according to

manufacturer's protocol (#IB24001, Thermo Fisher Scientific). PVDF membrane was washed once in 1×

PBST buffer (#28352, Thermo Fisher Scientific) and blocked in 2% blocking solution (#10156414, Thermo

Fisher Scientific) for 1 h at room temperature. Membranes were incubated overnight at +4 °C with primary

antibodies at recommended concentrations in 1% blocking solution. Membranes were washed three

times (5 min for each wash) with 1× PBST buffer, and secondary HRP-conjugated antibodies in 1% blocking

solution were added to the membranes at 1:5000 concentration for 1 h incubation at room temperature.

Membranes were washed three times with 1× PBST and proteins were detected by chemiluminescence

according to manufacturer's protocol (#RPN2232, Thermo Fisher Scientific). The following antibodies

were used. primary antibodies: RunX– (D1L7F - 12556) from Cell Signaling Technology and Actin (612656)

is from Becton Dickinson. The secondary antibodies were anti-Mouse (GENA931) from Sigma-Aldrich, anti-

Rabbit (NA9340V) from Thermo Fisher Scientific.

Biochemical analysis (GAG analysis)

Lyophilized tissue constructs were digested by overnight incubation in 0.5 ml of Proteinase K solution (1

mg/mL Proteinase K, Sigma; 10 ug/ml pepstatin A, Sigma; 1 mM EDTA, Sigma; 100 mM Iodoacetamide;

50mM Tris) at pH 7.6 and 56 °C. The GAG content of digested samples was analyzed using

Glycosaminoglycan Assay Blyscan kit (Biocolor) following the manufacturer instruction.

Mice

FoxN1 KO BALB/C (nude mice) 6–8-weeks old, were obtained from Charles River Laboratories. All mouse

experiments and animal care were performed in accordance with the Lund University Animal Ethical

Committee guidelines (ethical permit #M15485–18). Mice were housed at a 12-hour light cycle in

individually ventilated cages at a positive air pressure and constant temperature. Mice were fed with

autoclaved diet and water ad libitum. During the implantation procedure, anaesthesia was performed

with 2-3 % Isoflurane (Attane). The mice were kept on a heating pad during the procedure to avoid the

heat loss.

Micro-CT scanning

In vivo samples were explanted and fixed overnight with 4% formaldehyde before ex vivo micro-CT

analysis using a U-CT system (MILABS, Netherland) equipped with a tungsten x-ray source at 50 kV and

0.21 mA. Volumes were reconstituted at 10 µm isotropic voxel size. For total volume (TV) analysis, each

sample was assessed with Blender (v2.82a, Netherland). Briefly, a mesh was created surrounding the 3D

reconstruction of each sample and the volume occupied was then quantified. For bone volume (BV)

analysis, the highly mineralized tissue volume was quantified using Seg3D (v2.2.1, NIH, NCRR, Science

Computing and Imaging Institute (SCI))

Sample preparation for histological analysis.

In vitro samples were directly fixed prior to sectioning. In vivo samples were fixed and subsequently

decalcified with a 10% EDTA solution, pH = 8, at 4 °C for two weeks prior to tissue embedding.

Paraffin embedding was performed on samples fixed in 4% formalin (Solveca AB). Tissues were

dehydrated by immersion in consecutive solutions of 35%, 70%, 95%, and 99.5% graded ethanol solution

(Solveco AB). Immersion in a 99.5% ethanol/xylene solution (1:1, Fisher Scientific) was then performed

for ten minutes followed by two rinsing in xylene (Fisher Scientific) for 20 minutes. Subsequently,

tissues were embedded in paraffin at 56 °C overnight, before sectioning with a microtome (Microm HM

355 Rotary Microto–e) in 5 - 10 µm sections. The sections were then dried overnight at 37 °C before

staining. Prior to staining, sections were deparaffinized by two washes in xylene for 7 minutes and once

in 99.5% ethanol/xylene solution (1:1) for 3 minutes. Afterwards, sections were hydrated twice in

consecutive solutions of 99.5%, 95%, 70%, and 35% ethanol, for 7 minutes each.

Agarose embedding was performed on samples fixed overnight with 4% paraformaldehyde (Thermo

Scientific), using 4% low-melting agarose (Sigma). Sections of 100 µm thickness were obtained using a

7000smz vibratome (Campden) with stainless steel or ceramic blades.

Safranin O Staining

Paraffin-embedded sections were stained using Mayer's hematoxylin solution (Sigma-Aldrich) for 10

minutes. Samples were then placed under running distilled water to remove superfluous staining from

the sections. Subsequently, sections were stained with 0.01% fast green solution (Fisher Scientific) for five

minutes and rinsed with 1% acetic acid solution (glacial, Fisher Scientific) for 15 seconds. After that, the

slides were stained with 0.1% safranin O (Fisher Scientific) solution for 5 minutes. Dehydration and

clearing were performed by immersion in 95%, 99.5% ethanol, 99.5% ethanol/xylene solution (1:1) and

xylene twice successively for 2 minutes. Finally, the stained slides were mounted with glass slides using

PERTEX mounting medium (PERTEX, HistoLab).'

Masson's trichrome staining

Masson's trichrome staining was performed using the trichrome staining kit (Sigma-Aldrich Sweden AB)

according to the manufacturer's guidelines. Briefly, tissue sections were deparaffinized and immersed in

cold running deionized water for 3 minutes. Then the sections were kept in Bouin's solution at room

temperature overnight or at 56 °C for 15 minutes. The slides were washed by running tap water and

stained using working Weigert's iron hematoxylin solution (Sigma) for five minutes for nuclei detection

(in black). After washing the slides, the cytoplasm was stained in red with Biebrich Scarlet-Acid fuchsin for

five minutes followed by clearing the slides by immersion in working phosphotungstic/ phosphomolybdic

acid solution for five minutes. Collagen stained blue by immersion in aniline blue solution for five minutes

followed by clearing in 1% acetic acid (glacial, Fisher Scientific) solution diluted in distilled water (glacial,

Fisher Scientific) for two minutes and washing with running deionized water. Finally, the sections were

dehydrated in graded ethanol solutions (95% once, 100% twice) for two minutes each before washing

with xylene twice for two minutes and mounted with PERTEX mounting medium.

Hematoxylin and Eosin staining

Paraffin-embedded sections were stained using Mayer's hematoxylin solution (Sigma-Aldrich) for 10

minutes. Samples were then placed under running distilled water to remove superfluous staining from

the sections. The slides were immersed precisely in 1X phosphate-buffered saline (PBS) for 1 minute to

intensify the blue nuclei staining while preserving tissue integrity. Following this, a thorough rinse with 3

changes of distilled water effectively removed any residual PBS, preparing the sections for the subsequent

counterstaining step. Subsequently, Alcoholic-Eosin (Sigma-Aldrich) was applied for 1 minute as the

counterstain without any rinsing post-application, ensuring optimal interaction of the staining solution

with the tis. Dehydration and clearing were performed by immersion in 95%, 99.5% ethanol, 99.5%

ethanol/xylene solution (1:1) and xylene twice successively for 2 minutes. Finally, the stained slides were

mounted with glass coverslips using PERTEX mounting medium (PERTEX, HistoLab).

Quantitative PCR

Total RNA isolation was conducted from in vitro engineered tissues utilizing the Quick-RNA extraction kit

(Zymo Research, R1055) following the manufacturer's protocol. Subsequently, cDNA extraction was

carried out using cDNA synthesis kit (Invitrogen 11917020). The quantitative real-time polymerase chain

reaction (qRT-PCR) was performed, utilizing the assay on demand from Applied Biosystems to assess the

expression of specific genes. These genes include glyceraldehyde 3-phosphate dehydrogenase (GAPDH,

HS02786624_G1), runt-related transcription factor 2 (Runx2, Hs00231692_m1), alkaline phosphatase

(ALP, Hs01029144_m1), and collagen type X (ColX, Hs00166657_m1).

Rat osteochondral defect model

Ten- to 12-week-old male Sprague Dawley rats (n=11) were purchased from JANVIER LABS

(France). After 7 days of acclimation the rats were anesthetized using 3% isoflurane. Then,

animals were placed on a 37°C warm heating pad in prone position. Once anesthetized, isoflurane

was lowered to 2 to 2.5%, and buprenorphine (Temgesic, 30 µg/kg; Indivior Europe Ltd., Dublin,

Ireland) was injected subcutaneously for analgesia. The right hind limb of the animal was shaved

and carefully disinfected, and an incision was made along the skin and soft tissue to expose the

right knee. After cleaning the knee laterally from soft tissue, patella was displaced laterally to

expose the distal femur. An articular cartilage defect with 2mm depth and 1mm diameter was

created at the trochlear groove of femur. The defect was then push-fitted with the grafts, and

not secured with any sutures. The wound was closed in a layered fashion using resorbable sutures

(Vicryl 45-0, Ethicon, Somerville, USA) by closing the joint capsule, followed by the muscle tissue

(continuous interlaced suture), followed by closing the skin (Donati suture). Animals started load

bearing immediately after surgery. After 3 weeks (n=11) animals were anesthetized by isoflurane

inhalation (3%), followed by CO_2 asphyxiation, and right-femur were harvested prior to

subsequent micro-CT and histological characterization.

Histology grading method

Histological sections from rat femur condyles were stained with Safranin-O, Masson's trichrome, and H&E

staining. A semi-quantitative analysis of the repaired tissue was performed by utilizing a customized

adaptation of the histological grading system initially outlined by Wakitani et al[61]. This modified scale

encompasses six distinct parameters: cellular morphology, matrix staining, surface regularity, cartilage

thickness, regenerated subchondral bone formation, and integration with neighbouring cartilage. Each

parameter was graded on a numerical scale ranging from 0 to 4 points by independent expert observers,

where a maximum total score of 16 indicates the presence of tissue displaying characteristics akin to

entirely healthy and normal tissue (Table 1).

Cartilage quantification method

An evaluation of three distinct sites within the condyle defect was conducted to ascertain the proportion

of cartilage in each knee. Utilizing the measured femoral condyle thickness (200μm) and defect length

(1mm), a standardized Region of Interest (ROI) was defined using ImageJ. The percentage of cartilage-

positive regions within each defect was calculated in relation to the total ROI size, offering a quantitative

assessment of cartilage regeneration. This identical approach was applied to evaluate the percentage of

cartilage in the healthy control samples.

Trabecular thickness (Tb.Th) and trabecular separation (Tb.Sp)

Image analysis was performed by ImageJ. To evaluate the defect space, a rectangular interface region

(2 × 1 mm) was defined as the Region of Interest . Trabecular thickness (Tb.Th) and trabecular separation

(Tb.Sp),was calculated inside the ROI using BoneJ plugin.

Vascular density quantification

To assess vascular density, the Quantitative Vascular Analysis Tool (Q-VAT)[29] was utilized. The samples

were placed inside the chicken egg for CAM assay, and the images were acquired using a digital

microscope. These images were then segmented into tiles for detailed analysis. The binary vascular masks

were generated using the Q-VAT's automated capabilities, to distinguish vascular structures within the

tissue. The tool then facilitated the separation of vessel measurements based on their diameters, allowing

for a differentiated quantification of macro- and microvasculature. These binary masks were then

processed to calculate the vascular density, which refers to the proportion of the tissue area occupied by

vessels. This quantitative metric was evaluated for each tile, and the mean vascular density was computed

across all tiles for each tissue sample. As part of the analysis, we applied Q-VAT to double or triple-stained

slides to quantify the overlapping percentage of vessels, comparing various time points to observe

changes over the course of the experiment.

Immunofluorescence

Agarose-embedded sections were treated with 0.5% Triton X-100 (Sigma) in PBS supplemented with 20%

donkey serum (Jackson ImmunoResearch) for one hour at room temperature (RT) for blocking and

permeabilization. After blocking/permeabilization, sections were stained with primary antibodies (mouse

anti-Collagen II (Invitrogen, MA137493), Anti-Collagen 1 (Abcam, ab138492), VEGF polyclonal antibody

(Bioss Antibodies, bs-0279R),   rabbit anti-Collagen Type X (ColX) (abbexa, abx101469), Goat anti

mouse/rat CD31 (R&D system, YZU0119121), VEGF at 4 °C overnight. Sections were washed three times

using cold PBS with 0.1% Triton X-100, 20 minutes each, followed by staining with secondary antibodies

(CF568 donkey anti-rabbit (Biotium, 20098), CF567 anti-goat (Invitrogen, A11057), CF633 donkey anti-

mouse (Sigma-Aldrich, SAB4600131 in 2% donkey serum for three hours at room temperature. Sections

were washed three times using cold PBS with 0.1% Triton X-100, 20 minutes each once more.

All sections were mounted using the Vectashield antifade mounting medium containing DAPI (Vector

Laboratories, H1200). A LSM780 confocal microscope (Zeiss) with a 10x/20x objective was used to capture

images and image series of whole tissue sections (z-step = 2 µm). The IMARIS 9.5 software package (Oxford

Instruments) was used to analyze the data.

Acknowledgements

We express our gratitude to the following institutions and individuals for their invaluable support and

contributions to this study: Lund University, Medical Faculty, FACS Facility (Anna Hammarberg), Cell and

Gene Therapy core facility (Pia Johansson) and Imaging Facility at Lund Stem Cell Centre, as well as the

Lund University Bioimaging Centre, for providing access to the X-ray CT system. The funding for this

research was made possible by a Knut and Alice Wallenberg grant, the Medical Faculty at Lund University,

the Swedish Research Council, the European Research Council (Starting grant hOssicle #948588 to P.E.B),

and Region Skåne. Figures were generated using BioRender.com. Special thanks to Ani Grigoryan and

Dimitra Zacharaki for their expertise and valuable insights regarding FACS experiments, and to Steven

Dupard for his advice and guidance on the design of gRNAs and molecular techniques. Christian Hansen

deserves our appreciation for providing the mcherry plasmid and offering valuable guidance and

knowledge related to CRISPR-Cas9.

References

1.   Bonnans C, Chou J, Werb Z. Remodelling the extracellular matrix in development and disease.
Nat Rev Mol Cell Biol. 2014;15(12):786-801. doi:10.1038/nrm3904

2.   Hussey GS, Dziki JL, Badylak SF. Extracellular matrix-based materials for regenerative medicine.
Nat Rev Mater. 2018;3(7):159-173. doi:10.1038/s41578-018-0023-x

3.   Fattahi R, Chamkhorami FM, Taghipour N, Keshel SH. The effect of extracellular matrix
remodeling on material-based strategies for bone regeneration: Review article. *Tissue Cell*.
2022;76(January):101748. doi:10.1016/j.tice.2022.101748

4.   Mouw JK, Ou G, Weaver VM. Extracellular matrix assembly: A multiscale deconstruction. *Nat Rev
Mol Cell Biol*. 2014;15(12):771-785. doi:10.1038/nrm3902

5.   Assunção M, Dehghan-Baniani D, Yiu CHK, Später T, Beyer S, Blocki A. Cell-Derived Extracellular
Matrix for Tissue Engineering and Regenerative Medicine. *Front Bioeng Biotechnol*.
2020;8(December):1-10. doi:10.3389/fbioe.2020.602009

6.   Hinderer S, Layland SL, Schenke-Layland K. ECM and ECM-like materials - Biomaterials for
applications in regenerative medicine and cancer therapy. *Adv Drug Deliv Rev*. 2016;97:260-269.
doi:10.1016/j.addr.2015.11.019

7.   Hinderer S, Layland SL, Schenke-Layland K. ECM and ECM-like materials - Biomaterials for
applications in regenerative medicine and cancer therapy. *Adv Drug Deliv Rev*. 2016;97:260-269.
doi:10.1016/j.addr.2015.11.019

8.   Fattahi R, Chamkhorami FM, Taghipour N, Keshel SH. The effect of extracellular matrix
remodeling on material-based strategies for bone regeneration: Review article. *Tissue Cell*.
2022;76(January):101748. doi:10.1016/j.tice.2022.101748

9.   Tang G, Liu Z, Liu Y, et al. Recent Trends in the Development of Bone Regenerative Biomaterials.
Front Cell Dev Biol. 2021;9(May):1-18. doi:10.3389/fcell.2021.665813

10.   Gao C, Peng S, Feng P, Shuai C. Bone biomaterials and interactions with stem cells. *Bone Res*.
2017;5(May):1-33. doi:10.1038/boneres.2017.59

11.   Filippi M, Born G, Chaaban M, Scherberich A. Natural Polymeric Scaffolds in Bone Regeneration.
Front Bioeng Biotechnol. 2020;8(May). doi:10.3389/fbioe.2020.00474

12.   Reddy MSB, Ponnamma D, Choudhary R, Sadasivuni KK. A comparative review of natural and
synthetic biopolymer composite scaffolds. *Polymers (Basel)*. 2021;13(7).
doi:10.3390/polym13071105

13.   Decaris ML, Leach JK. Design of experiments approach to engineer cell-secreted matrices for
directing osteogenic differentiation. *Ann Biomed Eng*. 2011;39(4):1174-1185.
doi:10.1007/s10439-010-0217-x

14.   Harvestine JN, Orbay H, Chen JY, Sahar DE, Leach JK. Cell-secreted extracellular matrix,
independent of cell source, promotes the osteogenic differentiation of human stromal vascular
fraction. *J Mater Chem B*. 2018;6(24):4104-4115. doi:10.1039/c7tb02787g

15.   Hoch AI, Mittal V, Mitra D, Vollmer N, Zikry CA, Leach JK. Cell-secreted matrices perpetuate the
bone-forming phenotype of differentiated mesenchymal stem cells. *Biomaterials*. 2016;74:178-
187. doi:10.1016/j.biomaterials.2015.10.003

16.   Haumer A, Bourgine PE, Occhetta P, Born G, Tasso R, Martin I. Delivery of cellular factors to
regulate bone healing. *Adv Drug Deliv Rev*. 2018;129:285-294. doi:10.1016/j.addr.2018.01.010

17.   Sadr N, Pippenger BE, Scherberich A, et al. Enhancing the biological performance of synthetic
polymeric materials by decoration with engineered, decellularized extracellular matrix.
Biomaterials. 2012;33(20):5085-5093. doi:10.1016/j.biomaterials.2012.03.082

18.   Prewitz MC, Seib FP, von Bonin M, et al. Tightly anchored tissue-mimetic matrices as instructive
stem cell microenvironments. *Nat Methods*. 2013;10(8):788-794. doi:10.1038/nmeth.2523

19.   Bourgine PE, Gaudiello E, Pippenger B, et al. Engineered Extracellular Matrices as Biomaterials of
Tunable Composition and Function. *Adv Funct Mater*. Published online 2017.
doi:10.1002/adfm.201605486

20.   Pigeot S, Klein T, Gullotta F, et al. Manufacturing of Human Tissues as off-the-Shelf Grafts
Programmed to Induce Regeneration. *Advanced Materials*. 2021;33(43).
doi:10.1002/adma.202103737

21.   Pigeot S, Klein T, Gullotta F, et al. Manufacturing of Human Tissues as off-the-Shelf Grafts
Programmed to Induce Regeneration. *Advanced Materials*. 2021;33(43).
doi:10.1002/adma.202103737

22.   Grigoryan A, Zacharaki D, Balhuizen A, et al. Engineering human mini-bones for the standardized
modeling of healthy hematopoiesis, leukemia, and solid tumor metastasis. *Sci Transl Med*.
2022;14(666):1-16. doi:10.1126/scitranslmed.abm6391

23.   Golchin A, Shams F, Karami F. Advancing Mesenchymal Stem Cell Therapy with CRISPR/Cas9 for
630   Clinical Trial Studies. In: *Cell Biology and Translational Medicine,*. Vol 8. ; 2019:89-100.
631   doi:10.1007/5584_2019_459

632   24.   Hazrati A, Malekpour K, Soudi S, Hashemi SM. CRISPR/Cas9-engineered mesenchymal
633   stromal/stem cells and their extracellular vesicles: A new approach to overcoming cell therapy
634   limitations. *Biomedicine & Pharmacotherapy*. 2022;156:113943.
635   doi:10.1016/j.biopha.2022.113943

636   25.   Geurts MH, Clevers H. CRISPR engineering in adult stem cell-derived organoids for gene repair
637   and disease modelling. 2023;1(January):32-45. doi:10.1038/s44222-022-00013-5

638   26.   Hu K, Olsen BR. The roles of vascular endothelial growth factor in bone repair and regeneration.
639   *Bone*. 2016;91:30-38. doi:10.1016/j.bone.2016.06.013

640   27.   Duda GN, Geissler S, Checa S, Tsitsilonis S, Petersen A, Schmidt-Bleek K. The decisive early phase
641   of bone regeneration. *Nat Rev Rheumatol*. 2023;19(February):78-95. doi:10.1038/s41584-022-
642   00887-0

643   28.   Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-
644   Cas9 system. *Nat Protoc*. 2013;8(11):2281-2308. doi:10.1038/nprot.2013.143

645   29.   Callewaert B, Gsell W, Himmelreich U, Jones EAV. Q-VAT: Quantitative Vascular Analysis Tool.
646   *Front Cardiovasc Med*. 2023;10. doi:10.3389/fcvm.2023.1147462

647   30.   Otto F, Thornell AP, Crompton T, et al. Cbfa1, a candidate gene for cleidocranial dysplasia
648   syndrome, is essential for osteoblast differentiation and bone development. *Cell*. 1997;89(5):765-
649   771. doi:10.1016/S0092-8674(00)80259-7

650   31.   Komori T, Yagi * H., Nomura S, Yamaguchi A. Targeted disruption of. *Proc Natl Acad Sci U S A*.
651   1998;95(15):8692-8697.

652   32.   Kim HJ, Kim WJ, Ryoo HM. Post-Translational Regulations of Transcriptional Activity of RUNX2.
653   *Mol Cells*. 2020;43(2):160-167. doi:10.14348/molcells.2019.0247

654   33.   Hattori EY, Masuda T, Mineharu Y, et al. A RUNX-targeted gene switch-off approach modulates
655   the BIRC5/PIF1-p21 pathway and reduces glioblastoma growth in mice. *Commun Biol*.
656   2022;5(1):1-11. doi:10.1038/s42003-022-03917-5

657   34.   Maehara H, Sotome S, Yoshii T, et al. Repair of large osteochondral defects in rabbits using
658   porous hydroxyapatite/collagen (HAp/Col) and fibroblast growth factor-2 (FGF-2). *Journal of
659   Orthopaedic Research*. 2010;28(5):677-686. doi:10.1002/jor.21032

660   35.   Kusumbe AP, Ramasamy SK, Adams RH. Coupling of angiogenesis and osteogenesis by a specific
661   vessel subtype in bone. *Nature*. 2014;507(7492):323-328. doi:10.1038/nature13145

662   36.   Tuckermann J, Adams RH. The endothelium–bone axis in development, homeostasis and bone
663   and joint disease. *Nat Rev Rheumatol*. 2021;17(10):608-620. doi:10.1038/s41584-021-00682-3

37.　Largo RD, Burger MG, Harschnitz O, et al. VEGF Over-Expression by Engineered BMSC Accelerates
Functional Perfusion, Improving Tissue Density and In-Growth in Clinical-Size Osteogenic Grafts.
Front Bioeng Biotechnol. 2020;8(July):1-13. doi:10.3389/fbioe.2020.00755

38.　Yang YQ, Tan YY, Wong R, Wenden A, Zhang LK, Rabie ABM. The role of vascular endothelial
growth factor in ossification. *Int J Oral Sci*. 2012;4(2):64-68. doi:10.1038/ijos.2012.33

39.　Burger MG, Grosso A, Briquez PS, et al. Robust coupling of angiogenesis and osteogenesis by
VEGF-decorated matrices for bone regeneration. *Acta Biomater*. 2022;149:111-125.
doi:10.1016/j.actbio.2022.07.014

40.　Behr B, Tang C, Germann G, Longaker MT, Quarto N. Locally applied vascular endothelial growth
factor A increases the osteogenic healing capacity of human adipose-derived stem cells by
promoting osteogenic and endothelial differentiation. *Stem Cells*. 2011;29(2):286-296.
doi:10.1002/stem.581

41.　Zuo WH, Zeng P, Chen X, Lu YJ, Li A, Wu J Bin. Promotive effects of bone morphogenetic protein 2
on angiogenesis in hepatocarcinoma via multiple signal pathways. *Sci Rep*. 2016;6(July):1-9.
doi:10.1038/srep37499

42.　Lowery JW, de Caestecker MP. BMP signaling in vascular development and disease. *Cytokine
Growth Factor Rev*. 2010;21(4):287-298. doi:10.1016/j.cytogfr.2010.06.001

43.　Kamekura S, Kawasaki Y, Hoshi K, et al. Contribution of runt-related transcription factor 2 to the
pathogenesis of osteoarthritis in mice after induction of knee joint instability. *Arthritis Rheum*.
2006;54(8):2462-2470. doi:10.1002/art.22041

44.　Komori T, Yagi * H., Nomura S, Yamaguchi A. Targeted disruption of. *Proc Natl Acad Sci U S A*.
1998;95(15):8692-8697.

45.　Meng X, Ziadlou R, Grad S, et al. Animal Models of Osteochondral Defect for Testing Biomaterials.
Biochem Res Int. 2020;2020. doi:10.1155/2020/9659412

46.　Sekiya I, Vuoristo JT, Larson BL, Prockop DJ. In vitro cartilage formation by human adult stem cells
from bone marrow stroma defines the sequence of cellular and molecular events during
chondrogenesis. *Proc Natl Acad Sci U S A*. 2002;99(7):4397-4402. doi:10.1073/pnas.052716199

47.　Pelttari K, Winter A, Steck E, et al. Premature induction of hypertrophy during in vitro
chondrogenesis of human mesenchymal stem cells correlates with calcification and vascular
invasion after ectopic transplantation in SCID mice. *Arthritis Rheum*. 2006;54(10):3254-3266.
doi:10.1002/art.22136

48.　Mueller MB, Tuan RS. Functional characterization of hypertrophy in chondrogenesis of human
mesenchymal stem cells. *Arthritis Rheum*. 2008;58(5):1377-1388. doi:10.1002/art.23370

49.　Browe DC, Burdis R, Díaz-Payno PJ, et al. Promoting endogenous articular cartilage regeneration
using extracellular matrix scaffolds. *Mater Today Bio*. 2022;16. doi:10.1016/j.mtbio.2022.100343

50.   Steele JAM, Moore AC, St-Pierre JP, et al. In vitro and in vivo investigation of a zonal
700   microstructured scaffold for osteochondral defect repair. *Biomaterials*. 2022;286.
701   doi:10.1016/j.biomaterials.2022.121548

702   51.   Park DY, Min BH, Park SR, et al. Engineered cartilage utilizing fetal cartilage-derived progenitor
703   cells for cartilage repair. *Sci Rep*. 2020;10(1):1-13. doi:10.1038/s41598-020-62580-0

704   52.   Zeng J, Huang L, Xiong H, et al. Injectable decellularized cartilage matrix hydrogel encapsulating
705   urine-derived stem cells for immunomodulatory and cartilage defect regeneration. *NPJ Regen*
706   *Med*. 2022;7(1):1-12. doi:10.1038/s41536-022-00269-w

707   53.   Golchin A, Shams F, Karami F. Advancing Mesenchymal Stem Cell Therapy with CRISPR/Cas9 for
708   Clinical Trial Studies. In: *Cell Biology and Translational Medicine,*. Vol 8. ; 2019:89-100.
709   doi:10.1007/5584_2019_459

710   54.   Hazrati A, Malekpour K, Soudi S, Hashemi SM. CRISPR/Cas9-engineered mesenchymal
711   stromal/stem cells and their extracellular vesicles: A new approach to overcoming cell therapy
712   limitations. *Biomedicine & Pharmacotherapy*. 2022;156:113943.
713   doi:10.1016/j.biopha.2022.113943

714   55.   Chen Z, Klein T, Murray RZ, et al. Osteoimmunomodulation for the development of advanced
715   bone biomaterials. *Materials Today*. 2016;19(6):304-321. doi:10.1016/j.mattod.2015.11.004

716   56.   Xie Y, Hu C, Feng Y, et al. Osteoimmunomodulatory effects of biomaterial modification strategies
717   on macrophage polarization and bone regeneration. *Regen Biomater*. 2020;7(3):233-245.
718   doi:10.1093/rb/rbaa006

719   57.   Aamodt JM, Grainger DW. Extracellular matrix-based biomaterial scaffolds and the host
720   response. *Biomaterials*. 2016;86:68-82. doi:10.1016/j.biomaterials.2016.02.003

721   58.   Ramakrishnaiah Siddappa, Ruud Licht, Clemens van Blitterswijk J de B. Donor variation and loss of
722   multipotency during in vitro expansion of human mesenchymal stem cells for bone tissue
723   engineering. *Journal of Orthopaedic Research*. 2007;25(8).

724   59.   LaFleur MW, Nguyen TH, Coxe MA, et al. A CRISPR-Cas9 delivery system for in vivo screening of
725   genes in the immune system. *Nat Commun*. 2019;10(1):1-10. doi:10.1038/s41467-019-09656-2

726   60.   Bock C, Datlinger P, Chardon F, et al. High-content CRISPR screening. *Nature Reviews Methods*
727   *Primers*. 2022;2(1). doi:10.1038/s43586-021-00093-4

728   61.   Maehara H, Sotome S, Yoshii T, et al. Repair of large osteochondral defects in rabbits using
729   porous hydroxyapatite/collagen (HAp/Col) and fibroblast growth factor-2 (FGF-2). *Journal of*
730   *Orthopaedic Research*. 2010;28(5):677-686. doi:10.1002/jor.21032

Figure 1. CRISPR/Cas9 editing of mesenchymal cell lines lead to efficient VEGF knockout in cartilage tissues. A) Experimental scheme depicting the generation of CRISPR/Cas9-edited MSOD-B lines, and the subsequent in vitro and in vivo tissue formation assessment. B) Overview of the native human VEGF coding sequence composed of eight exons. Designed gRNAs and their targeted binding sites are illustrated, as well as the corresponding expected impact on the coding sequence. gRNAs targeting exon 1 (orange) disrupt translation initiation and inhibit protein expression, gRNA targeting exon 2 (blue) disrupts VEGF receptor binding, and gRNA targeting exon 8 (green) alters the C-Terminal sequence and repress activation of protein. C) ELISA-based quantitative analysis of VEGF protein content in cell culture supernatant from expanded single cell colonies. From all clones, only two had no detectable level of VEGF (1.3_3 and 1.3_6). These clones were subsequently defined as MSOD-BΔV1 and MSOD-BΔV2. D) Histological assessment of in vitro differentiated constructs using Safranin O (top panel) and Masson's trichrome (bottom panel) stains. Both the MSOD-B and MSOD-BΔV1 displayed glycosaminoglycans (GAG) (orange to reddish in Safranin O) and collagen content (Blueish in Masson's trichrome), indicating successful cartilage formation (Scale bars = 200 μm). Left bottom inserts show the whole tissue section. E) Quantitative assessment of the total GAG content in MSOD-B and MSOD-BΔV1 in vitro differentiated constructs. Unpaired t-test, n = 3 biological replicates, n.s. = not significant. F) ELISA-based quantitative assessment of VEGF protein in in vitro differentiated constructs. Unpaired t-test, n = 3-4 biological replicates, ****p < 0.0001. G) Immunofluorescence images of MSOD-B and MSOD-BΔV1 tissues. Displayed images consist of 3D-stacks from 80-100 μm thick sections, stained for VEGF (yellow), Collagen Type X (COLX, red) and Collagen Type I (COL1, grey). A clear reduction in the VEGF signal could be observed in the MSOD-BΔV1 tissues, indicating a successful VEGF knockdown (Scale bars = 80 μm).

A

Day -6 — Day -3 — Day 0 — Day 4

Started incubation At 37.5 °C — Eggs windowing — Implantation of samples — Termination of assay

B

MSOD-B | MSOD-BΔV1

Day 0

Day 4

C

Vascular Density after 4 days %

MSOD-B | MSOD-BΔV1

ns

D

Subcutaneous Implantation of engineered constructs in mice and in vivo analysis, 2 – 6 weeks

E

DAPI | CD31 | Merged

MSOD-B

MSOD-B ΔV1

a | b
c | d

F

CD31+ volume (μm³) Normalized to total volume

MSOD-B | MSOD-BΔV1

*

G

2 weeks | 6 weeks

MSOD-B

MSOD-B ΔV1

H

BV/TV %

2 weeks | 6 weeks

ns | ns

● MSOD-B
○ MSOD-BΔV1

I

2 weeks | 6 weeks

Safranin O | Masson's Trichrome | Safranin O | Masson's Trichrome

MSOD-B

CB BM TB | CB TB BM

MSOD-BΔV1

BM CB | CB BM TB

Figure 2. VEGF knockout cartilage tissues retain bone remodeling capacity despite reduced early-stage vascularization

A) Experimental scheme of the Chorioallantoic Membrane (CAM) assay, for ex vivo evaluation of angiogenic potential. B) Macroscopic comparison of in vitro MSOD-B and MSOD-BΔV1 constructs at day 0 (day of implantation) and day 4 (4 days post-implantation), illustrating robust de novo vessel formation perfusing the tissues (Scale bars at 1 mm). C) Quantitative Analysis of Vascular Density in MSOD-B and MSOD-BΔV1. Vascular densities were quantified from macroscopic images obtained on day 4 using ImageJ. Unpaired t-test, n = 3-4 biological replicates, n.s. = not significant. D) overview of the subcutaneous implantation procedure in mice and subsequent in vivo evaluation of vascularization from 2 to 6 weeks post-implantation. E) Immunofluorescence images of MSOD-B and MSOD-BΔV1 tissues two weeks post-in vivo implantation. Displayed images consist of 3D-stacks from 80-100 μm thick sections, vessels stained with mouse CD31 (red) and nuclei with DAPI (cyan) (Scale bars at 500 μm except for magnified white inserts at 80 μm). Box "a" and "c" display the periphery whereas Box "b" and "d" show the central region of MSOD-B and MSOD-BΔV1 constructs respectively. A reduction in tissue vascularization is observed in MSOD-BΔV1 samples. F) Quantitative analysis of the CD31 signal using an isosurface-based strategy (IMARIS software). Unpaired t-test, n = 3-4 biological replicates, $*p < 0.05$. G) Representative macroscopic and microtomography images of in vivo constructs retrieved at two and six weeks post-implantation. H) Microtomography-based quantification of the sample's bone/ mineralized volume over their total volume (ratio). No significant differences between MSOD-B and MSOD-BΔV1 could be observed. (BV: Bone Volume, TV: Total Volume). Ordinary One-way ANOVA, n=8 biological replicates, n.s.=not significant I) Histological analysis of in vivo tissues using Safranin O and Masson's trichrome stains, at two (2W) and six weeks (6W) post-implantation. Both sample types underwent full remodeling into a bone organ after 6 weeks, with presence of bone structures and a bone marrow compartment (Scale bars = 200 μm). CB – cortical bone; BM – bone marrow. TB – trabecular bone.

A Native coding sequence

RUNX2_6.2
RUNX2_6.1

5′ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 3′ 528 aa

RUNX2_2.1   RUNX2_5.1   RUNX2_8.1

Truncated coding sequences

5′ | 1 | 2 | 90 - 100 aa
5′ | 1 | 2 | 3 | 4 | 5 | 250 - 260 aa
5′ | 1 | 2 | 3 | 4 | 5 | 6 | 340 - 350 aa
5′ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 410 - 420 aa

B

Count

RUNX2

Isotype ctrl
MSOD-BΔR1
MSOD-BΔR2
MSOD-B ctrl

C

(kDa)

60
50

40 — RUNX2

40 — Actin

MSOD-B / MSOD-BΔR1 / MSOD-BΔR2

D

MSOD-B | MSOD-BΔR1

Safranin O

Masson's Trichrome

E

GAG (ug/pellet)

ns

MSOD-B / MSOD-BΔR1

F

DAPI | COL2 | COLX | Merged

MSOD-B

MSOD-BΔR1

G

Col2 volume (µm^3)
Normalized to total volume

ns

MSOD-B / MSOD-BΔR1

H

ColX volume (µm^3)
Normalized to total volume

***

MSOD-B / MSOD-BΔR1

I

Expansion medium | Osteogenic medium

MSOD-B | MSOD-BΔR1

J

Gene expression normalised to GAPDH (log scale)

RUNX2   COLX   ALPL

n.d

□ Expanded cells
■ MSOD-B
▨ MSOD-BΔR1

Figure 3. RUNX2 knockout does not prevent chondrogenic differentiation but impairs hypertrophy A) Overview of the human RUNX2 coding sequence comprising 8 exons. gRNAs and their corresponding expected protein structure. The gRNA targeting exon 2 (orange) disrupts the DNA binding domain, gRNA targeting exon 5 (violet) disrupts nuclear translocation, gRNAs targeting exon 6 (blue) disrupt the transcriptional activation domain and gRNAs targeting exon 8 (green) disrupt the nuclear matrix targeting signal and repress the protein function. B) Intracellular flow cytometry for RUNX2 detection in MSOD-B and RUNX2-edited clones. A clear protein reduction could be observed in the 6.1_1 and 6.1_23 clones. C) Western blot analysis of RUNX2 in cultured MSOD-B and RUNX2-edited cells. The genetic editing of RUNX2 is confirmed by the detection of the truncated proteins. Actin is used as a control to normalize the protein content. D) Histological analysis of in vitro constructs using Safranin O (top) and Masson's trichrome (bottom) stains, indicating the presence of cartilage matrices (Scale bars = 200 μm). E) Quantitative assessment of the total GAG content in corresponding in vitro generated tissues. Unpaired t-test, n=3 biological replicates, n.s.=not significant. F) Immunofluorescence images of MSOD-B and MSOD-BΔR1 tissues. Displayed images consist of 3D-stacks from 80-100 μm thick sections, stained for DAPI (blue), Collagen Type II (COL2, yellow) and Collagen Type X (COLX, red). A clear reduction in the COLX signal could be observed in the MSOD-BΔR1 tissues, indicating impaired hypertrophy. (Scale bars = 500 μm) G) Isosurface-based quantification of the COL2 immunofluorescent signal using the IMARIS software. No significant difference between groups confirm the retention of cartilage formation in RUNX2-edited constructs. Unpaired t-test, n=3 biological replicates, n.s.=not significant. H) Isosurface-based quantification of the COLX immunofluorescent signal using the IMARIS software, confirming the disruption of hypertrophy in the MSOD-BΔR1 constructs. Unpaired t-test, n=3, ***$p < 0.001$, n.s. = not significant. I) Alizarin Red staining evidencing a lack of mineralization in the MSOD-BΔR1 culture compared to the MSOD-B. Two way ANOVA test, n=3, **$p < 0.01$, n.s. = not significant. J) Quantitative PCR analysis displaying the relative expression levels of osteogenesis-related genes: RUNX2, COL1, and ALPL. The expression is normalized to GAPDH as housekeeping gene. n.d= not detected.

Figure 4. RUNX2 knockout in cartilage tissues disrupts effective ectopic bone formation A) Representative macroscopic and microtomography images of in vivo constructs retrieved at two and six weeks post-implantation. B) Microtomography-based quantification of the sample's bone/mineralized volume over their total volume (ratio). (BV: Bone Volume, TV: Total Volume). A marked difference is observed as early as two weeks, with a clear lower mineral content in MSOD-BΔR1 samples. Ordinary one way ANOVA, n=3 biological replicates, *p < 0.05, ***p < 0.001. C) Histological analysis of in vivo constructs using Safranin O and Masson's trichrome stains. After two weeks (2W), a higher bone formation is already evident in the MSOD-B control group. The MSOD-BΔR1 samples explanted after 6 weeks (6W) displayed presence of cortical and trabecular bone, but also large amount of fibrous tissue indicating an incomplete remodeling. (Scale bars = 200 μm).

Figure 5. RUNX2 knockout in cartilage tissues leads to better cartilage regeneration and maintenance in osteochondral defect in rats A) Experimental scheme for the regenerative potential assessment of MSOD-B & MSOD-BΔR1 cartilage tissues in a rat osteochondral defect. B) Histological analysis of the osteochondral defects for each group using H&E and Masson's trichrome stains, at three weeks post-implantation. The dash-line marks the defect area. (Scale bars = 500 μm) C) Histological analysis the osteochondral defects using TRAP staining reporting osteoclastic activity, at three weeks post-implantation. The dash-line marks the defect area. (Scale bars = 500 μm and 100 μm for magnified areas) D) Microtomography-based quantification of the sample's total bone/mineralized volume normalized to the healthy control in percentage. Unpaired t-test, n=3 biological replicates, n.s.=not significant. E) Image J based quantification of Trabecular separation (Tb.Sp). One way ANOVA test, n=3 biological replicates, *p < 0.05 F) Histological analysis of the osteochondral defects using Safranin O staining. After three weeks, a higher regeneration of the surface cartilage is evident in the MSOD-BΔR1 group (Scale bars = 500 μm and 100 μm for magnified areas). The magnified regions of the subchondral area (d,e,f) show higher cartilage remnants in the MSOD-BΔR1 group. G) Quantitative analysis of cartilage regeneration in the osteochondral surface as compare to the healthy control (100%). Unpaired t-test, n=3 biological replicates, *p < 0.05.

Table 1: Scoring rate for individual categories
Semi-quantitative analysis utilizing a modified histological grading system, including parameters such as cellular morphology, matrix staining, surface regularity, thickness of cartilage, subchondral bone formation and integration of adjacent cartilage , resulting in a comprehensive assessment of tissue regeneration and integration

	Scoring Rate (%) (SD)					
	Cell Morphology	Matrix Staining	Surface Regularity	Thickness of Cartilage	Regenerated Subchondral Bone	Integration with Adjacent Cartilage
Healthy	100 (0)	100 (0)	100 (0)	100 (0)	100 (0)	100 (0)
MSOD-B	33,33 (28,86)	16,66 (14,43)	50 (0)	8,33 (14,33)	16,66 (14,33)	33,33 (14,33)
MSOD-B ΔR1	66,66 (14,33)	50 (0)	41,66 (14,33)	33,33 (28,86)	25 (25)	50 (0)

A

Supplementary figure 1. CRISPR/Cas9 editing of mesenchymal cell lines lead to efficient VEGF knockout in cartilage tissues A) Overview of the human VEGF exon structure, nucleotide sequences of targeting gRNAs and their corresponding amino acid sequences.

A

B

C

- ● MSOD-B
- ○ MSOD-BΔV1

Supplementary figure 2. VEGF knockout cartilage tissues retain bone remodeling capacity despite reduced early-stage vascularization

A) Sequential processing of a vascular tissue section using Q-VAT, depicting the original image, binary mask creation, and vessel segmentation for quantitative vascular density analysis. B) Microtomography analysis (Bone/mineralized volume) showed significant differences between in vivo constructs at both two weeks and six weeks time points (One way ANOVA, n=3, **p < 0.01). C) Microtomography analysis (total volume) did not show significant difference between in vivo constructs at either two weeks or six weeks time points (One way ANOVA, n=3, n.s.= not significant) (BV: Bone Volume, TV: Total Volume).

A

| | 245 | 248 | 252 |
| | R L | S D L | G R I |

RUNX2_5.1 ←

g gag tca cta aat ccc gcg **tccac**

---- cgc ctc agt gat tta ggg cgc att ----
---- gcg gag tca cta aat ccc gcg taa ----

| | 406 | 409 | 412 |
| | S G | M S L | G M |

RUNX2_8.1 ←

gt ccg tac agg gag cca tac**ccac**

---- tca ggc atg tcc ctc ggt atg----
---- agt ccg tac agg gag cca tac----

QA repeat   RUNT domain and DNA binding site   Cell signal integration, activation and inhibition domain

5'   1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 528   3'

NLS   NMTS   VWRPY

| | 92 | 95 | 99 |
| | R L | R P P | H D N |

RUNX2_ 2.1 ←

gcc aac gcc ggc ggg gtg ct**ccac**

---- cgg ttg cgg ccg ccc cac gac aac----
---- gcc aac gcc ggc ggg gtg ctg ttg ---

| | 332 | 335 | 338 |
| | I T | D V P | R R |

RUNX2_6.1 ←

ag tgg cta cac gga tcc gcg**ccac**

---- gcc atc acc gat gtg cct agg cgc ----
---- cgg tag tgg cta cac gga tcc gcg ----

cgg tag tgg cta cac gga tc**ccac**

RUNX2_6.2 ←

B

(Panel B: flow cytometry histograms for RunX2)

Top left — legend:
- Isotype ctrl
- 5.1_11
- 5.1_14
- 5.1_16
- 5.1_17
- MSOD-B ctrl

Top middle — legend:
- Isotype ctrl
- 6.1_3
- 6.1_5
- 6.1_7
- 6.1_9
- 6.1_10
- 6.1_11
- 6.1_13
- MSOD-B ctrl

Top right — legend:
- Isotype ctrl
- 6.1_14
- 6.1_17
- 6.1_18
- 6.1_19
- 6.1_21
- 6.1_24
- 6.1_25
- 6.1_26
- 6.1_27
- MSOD-B ctrl

Bottom left — legend:
- Isotype ctrl
- 6.2_1
- 6.2_2
- 6.2_3
- 6.2_4
- 6.2_5
- 6.2_6
- 6.2_7
- MSOD-B ctrl

Bottom middle — legend:
- Isotype ctrl
- 6.2_8
- 6.2_9
- 6.2_10
- 6.2_11
- 6.2_12
- 6.2_13
- 6.2_14
- 6.2_16
- MSOD-B ctrl

Bottom right — legend:
- 8.1_1
- 8.1_2
- 8.1_4
- 8.1_7
- 8.1_8
- 8.1_9
- 8.1_10
- MSOD-B ctrl

C

D

E

F

G

Col2 volume (μm³) / Normalised to total volume

0.45
0.40
0.35
0.30
0.25
0.20

*

MSOD-B   MSOD-BΔR2

H

ColX volume (μm³) / Normalised to total volume

1.0
0.8
0.6
0.4
0.2
0.0

**

MSOD-B   MSOD-BΔR2

I

Supplementary figure 3: **RUNX2 knockout does not prevent chondrogenic differentiation but impairs hypertrophy** A) Overview of human RUNX2 exon structure, nucleotide sequences of targeting gRNAs and its corresponding amino acid sequence. B) Intracellular flow cytometry for RUNX2 detection in MSOD-B and RUNX2-edited clones. Clones whose peaks fell left of the dotted line were selected for sequencing. C) Sanger sequencing results of DNA extracted from MSOD-BΔR samples and the modifications compared to MSOD-B DNA sequence. D) Histological analysis of in vitro constructs from using Safranin O and Masson trichrome staining displayed the presence of glycosaminoglycans (GAG) and collagen content respectively, indicating the presence of cartilage formation in MSOD-BΔR2 (Scale bars = 200 μm). E) Quantitative assessment of the total GAG content (unpaired t-test, n=3, n.s. = not significant) confirms the presence of cartilage formation in both constructs thus confirming RUNX2 knockout in hMSCs (MSOD-BΔR2) retain cartilage formation. (unpaired t-test, n=3, n.s. = not significant). F) Reduction of RUNX2 expression in cartilage tissues lead to reduced hypertrophy. Representative immunofluorescence staining images of COL2, COLX and DAPI tissue section of in vitro cartilage constructs shows reduction of COLX expression in knockout construct (MSOD-BΔR2) indicating reduction of hypertrophy. (Scale bars = 500 μm). G) Quantitative analysis of the immunofluorescent staining sections using IMARIS software displayed significant difference in the expression of COL2, demonstrating reduction of cartilage formation in knockout constructs (MSOD-BΔR2) (unpaired t-test, n=3, *p < 0.05). H) Quantitative analysis of the immunofluorescent staining sections using IMARIS software displayed significant reduction in expression of COLX, confirming the disruption of hypertrophy in knockout constructs (unpaired t-test, n=3, n.s. = not significant). I) Representative isosurface images of immunofluorescent staining sections constructed using IMARIS software.

Supplementary figure 4. RUNX2 knockout in cartilage tissues disrupts effective bone formation A) Microtomography analysis (Bone/mineralized volume) showed very high significant difference between in vivo constructs at both two weeks and six weeks time points (One was ANOVA, n=3, *p < 0.05, ***p < 0.001). B) Microtomography analysis (total volume) showed significant difference between in vivo constructs at two weeks time point but no significant difference at six weeks time point (One was ANOVA, n=3, n.s. = not significant) (BV: Bone Volume, TV: Total Volume).

Supplementary figure 5. RUNX2 knockout in cartilage tissues exhibit better cartilage regeneration and maintenance in osteochondral defect in rats A) Macroscopic images osteochondral defect model in rats depicting in of empty defect, defect filled with MSOD-B and defect filled with MSOD-BΔR1 respectively . B) Macroscopic images of extracted rat knees depicting healthy control without any defect, defect filled with MSOD-B and defect filled with MSOD-BΔR1 respectively and microtomography images of same respective constructs. C) Microtomography-based quantification of the sample's bone/mineralized volume over their total volume (ratio). (BV: Bone Volume, TV: Total Volume). No significant difference in BV/TV is observed between MSOD-B & MSOD-BΔR1 samples despite significantly reduced BV/TV ratio in both samples compared to healthy control. One way ANOVA test, n=3 biological replicates, **p < 0.01. D) Image J based quantification of Trabecular thickness (tb.Th) . One way ANOVA test, n=3 biological replicates, n.s = not significant.

Supplementary table 1: Histological grading system The histological graded system is modified and adapted from previously established model as mentioned in materials and methods section. This modified scale incorporates six distinct parameters to comprehensively evaluate the quality of tissue repair

Category	Point
Cell morphology	
Hyaline cartilage	4
Mostly hyaline cartilage or fibrocartilage	3
Mostly fibrocartilage	2
Mostly non-cartilage	1
Non-cartilage	0
Matrix-staining (metachromasia)	
Normal	3
Slightly reduced	2
Markedly reduced	1
No metachromatic staining	0
Surface regularity[a]	
Smooth	3
Moderate	2
Irregular	1
Severely irregular	0
Thickness of cartilage (%)[b]	
121–150	1
81–120	2
51–80	1
0–50	0
Regenerated subchondral bone	
Good	2
Moderate	1
Poor	0
Integration with adjacent cartilage	
Both edges integrated	2
One edge integrated	1
Neither edge integrated	0
Total maximum	16

• [a] Total smooth area of the reparative cartilage compared with the entire area of the cartilage defect.

• [b] Average thickness of the reparative cartilage compared with that of the surrounding cartilage.