Fruitless mating with the exes: the irreversible parthenogenesis in a stick insect

Tomonari NOZAKI[1, 2], Yasuhiko CHIKAMI[2, 3], Koki YANO[3], Ryuta SATO[4], Kenji SUETSUGU[5], and Shingo KANEKO[4]

[1]*Laboratory of Evolutionary Genomics, National Institute for Basic Biology, Okazaki, Aichi 444-8585, Japan*

[2] *Department of Basic Biology, School of Life Science, The Graduate University for Advanced Studies, SOKENDAI, Okazaki, Aichi 444-8585, Japan*

[3] *Division of Evolutionary Developmental Biology, National Institute for Basic Biology, Okazaki, Aichi 444-8585, Japan*

[4]*Graduate School of Symbiotic Systems Science and Technology, Fukushima University, Fukushima, Fukushima 960-1296, Japan*

[5]*Department of Biology, Graduate School of Science, Kobe University, Kobe, Hyogo 657-8501, Japan*

*Author for correspondence:

Tomonari Nozaki; Email: nozaki.t@nibb.ac.jp

Shingo Kaneko; Email: kaneko.shingo@gmail.com

ORCID:

https://orcid.org/0000-0003-4358-8118 (TN)

https://orcid.org/0000-0001-8489-3245 (YC)

https://orcid.org/0000-0001-8960-6727 (KY)

https://orcid.org/0009-0000-5311-9927 (RS)

https://orcid.org/0000-0002-9021-8155 (SK)

https://orcid.org/0000-0002-7943-4164 (KS)

Abstract

Parthenogenetic lineages, common in many animals, have sparked debate about their

evolutionary persistence. While asexuality is expected to ensure reproductive assurance

and provide a demographic advantage, parthenogens should suffer from the lack of gene

shuffling with other individuals. On the other hand, occasional sexual reproduction has

been theoretically predicted to be enough to mitigate the long-term costs of

parthenogenesis. Recent studies have revealed instances of cryptic sex in some old

parthenogenetic lineages, which is most likely mediated by rarely occurring males.

Unlike female traits that rapidly become vestigial under asexuality, traits in males have

been predicted to slowly decay due to the accumulation of neutral mutations over long

evolutionary times. In fact, rare males often retain sexual functions, raising questions

about the asexuality of these long-standing parthenogenetic lineages. Here, we

intensively examined the possibility of sexual reproduction in the Japanese common

stick insect, *Ramulus mikado*, which was also suggested to be an old parthenogen in our

previous work. While asexual female reproduction appears to be quite predominant

throughout Japan, we fortunately obtained the rare males from the field. These males

exhibited typical stick insect male morphology and engaged in mating behaviors with

conspecific females. However, no paternal-specific alleles were detected in the

offspring; all embryos showed genotypes identical to their mothers. Our histological

observations on a few males revealed that they had no sperm in their reproductive

organs, although the degree of decay may be different among the lineages. We also

found that females have sexual organs with signs of degeneration. All these results

demonstrate the irreversible asexual reproduction of *R. mikado* and indicate their long

history as a parthenogenetic species. Our present study provides unique insights into the

maintenance of parthenogenesis and degenerative evolution of sexual traits in ancient

asexual lineages.

Keywords: Obligate parthenogenesis, Trait decay under asexuality, Relaxed selection,

Rare males, Stick insects

Introduction

Sexual reproduction is a widespread strategy among multicellular organisms; on the

other hand, parthenogenesis has repeatedly emerged and is pervasive among almost all

metazoan groups (Suomalainen 1962, Bell 1982, Butlin 2002, Simon et al. 2003, Avise

2008). Furthermore, although it is exceedingly rare, several species have been

recognized as "old" parthenogens, wherein females reproduce solely through parthenogenesis for millions of years (Judson & Normark 1996, Schurko et al. 2009, Schwander et al. 2010). The existence of ancient parthenogens challenges the idea that obligately parthenogenetic species should be short-lived due to the lack of sex and recombination, resulting in a reduced rate of adaptation and increased accumulation of deleterious mutations (Judson & Normark 1996). However, rare sex, which is theoretically predicted to be sufficient to mitigate the long-term costs of parthenogenesis (reviewed in D'Souza & Michiels 2010), has been recently detected in several species with obligate parthenogenesis (Vakhrusheva et al. 2020, Boyer et al. 2021, Freitas et al. 2023, but see Kearney et al. 2022). Therefore, variation in the reproductive modes, including occasional sex rather than the classical bimodal obligate sex and obligate parthenogenesis pattern, should be considered to improve our understanding of the evolutionary persistence of parthenogenesis.

Even after the loss of male individuals, males can often survive in the genomes; in fact, rarely occurring males have been described in many asexual lineages (van der Kooi & Schwander 2014) and are suspected to mediate the rare sex events (Vakhrusheva et al. 2020, Boyer et al. 2021, Freitas et al. 2023). In predominantly parthenogenetic species, females have substantially reduced opportunities for mating with males and have ultimately lost it during evolution (van der Kooi & Schwander 2014, Rayner et al. 2022). Under such conditions, the sexual traits of females would become free from stabilizing selection, which leads to the degeneration of female-specific traits due to the accumulation of neutral mutations or enhancement of the trade-off effect with other traits (Muller 1949, Fong et al. 1995). In contrast to female sexual traits rapidly vestigializing, traits in males that occur rarely within populations have been predicted to decay over much longer evolutionary times simply as a consequence of neutral mutation accumulation (van der Kooi & Schwander 2014, Jalinsky et al. 2020). Consistent with this prediction, many studies have concluded that the "rare males" can be functional ones, i.e., fertile ones (Itow et al. 1984, van der Kooi & Schwander 2014), implying cryptic gene flow in the species (cryptic sex) or between closely related species (e.g., contagious parthenogenesis).

Phasmida, stick and leaf insects, is a well-studied group in terms of the evolution of parthenogenetic lineages (e.g., Pantel 1917, Bedford 1978, Brock 2000, Morgan-Richards et al. 2010, Bast et al. 2018, Nakano et al. 2019). Many species are recognized as obligately parthenogenetic, yet rarely occurring males have often been described (Scali 2009, Brock et al. 2018, reviewed in Table S1). Schwander et al. (2013) reported functional males in ancient asexual species of *Timema*, and that such males can

copulate with the females of sister sexual species and successfully produce offspring. Furthermore, a recent study detected evidence of rare sex in asexual *Timema*, which is most likely mediated by rare males (Freitas et al. 2023). In a parthenogenetic strain of the laboratory stick insect, *Carausius morosus*, with a history of about 100 years, rare males have normal sperm as well as abnormal spermatids in their testis (Pehani 1925, Pijnacker 1964). These studies suggest the potential role of the rare males for the gene flow in the parthenogenetic species, however, the functionality of the males is rarely examined, presumably due to the difficulty in sample collection.

Here, we examined the possibility of sexual reproduction in the Japanese common stick insect, *Ramulus mikado*, which is known as a predominantly parthenogenetic species (Nagashima 2001, Machida et al. 2016, Yano et al. 2021). Males are extremely rare; in fact, there have only ever been about 10 observed cases of *R. mikado* males, despite it being the most common species of stick insects in Japan (Yano et al. 2021, Table S1). We were fortunate to have the opportunity to obtain rare males of this stick insect (Fig. 1A, Table S2) and examined whether the rare males were functional or not. Unexpectedly, we found no genetic contribution of the males to offspring despite their normal behaviors and morphology. Our histological observations revealed that the males did not produce normal sperms and females exhibited signs of degeneration in the sexual organs, highlighting the irreversible parthenogenesis in *R. mikado*.

Methods
Stick insects and their mating behavior
In this study, we collected individuals of *Ramulus mikado* (Rehn 1904), which is a parthenogenetic and wingless stick insect in Japan (Okada 1999, Machida et al. 2016). We used a total of 21 individuals, including six field-collected males and one that occurred in a breeding population (details in Table S2). Of the males, three males (Male#1, 3, and 4) were displayed and mated with females in museums. Male#2 was employed for detailed observation of behavior and sacrificed just after the mating experiment to examine the internal morphology (details below). Male#7 was stored at -20°C from the day after collection and dissected for anatomical observation. All collected individuals were in the late stage of nymph or adult. The legs of all field collected samples were finally kept in 99.5% ethanol for genotypic analysis.

To investigate whether the rare males were able to mate with conspecific females, we observed the sexual behavior of *R. mikado*. The individuals used for the mating

experiment were separately kept in plastic insect cages at room temperature (RT,

approximately 24°C to 28°C) with the food plants until mating. The species is

polyphagous, and the leaves of *Cerasus* sp., *Zelkova* sp., *Quercus* sp., or *Celtis* sp. were

placed in plastic insect cages as their food. Tap water was sprayed once every few days

to provide humidity. We placed Male#2 in the plastic cage with a female. Using a

smartphone camera (iPhone SE, Apple, USA), the mating behaviors of the male

(approaching, mounting, and penis extension and insertion into the female gonopore)

were recorded. This trial was performed three times with different females (Female#3, 4,

and 5) on alternate days. Male#1, 3, and 4, which were displayed in insect museums,

were also placed in insect cages with conspecific females and observed their sexual

behavior. We collected eggs that were oviposited before the mating experiment (virgin

eggs) and those that were oviposited after the experiment (mated eggs) to analyze the

genetic contribution of males to the next generation.

External and internal morphology of males and females

To determine the normality of external morphology, we observed the males after they

were killed (Male#2 and #7) or died (other males). We focused on mating-related

structures, such as claspers and penis, according to previous papers that described *R.*

mikado males (Nagashima 2001, Yano et al. 2021). Then we dissected Male#2 and #7 in

phosphate-buffered saline (PBS: 33 mM KH_2PO_4, 33 mM Na_2HPO_4, pH 6.8) under a

stereomicroscope (SZ61, Olympus, Japan) with fine forceps. The presence or absence of

components of the reproductive system, such as testes and seminal vesicles, was

recorded with reference to Matsuda (1976). For Male#2, one of the testes was then fixed

in 4% paraformaldehyde in PBS for confocal microscopy. Another was fixed with

Bouin's solution (saturated picric acid: formaldehyde: glacial acetic acid=15:5:1) at

25°C overnight and washed with 90% ethanol plus lithium carbonate (Li_2CO_3) for

histological observation. For Male#7, one of the ovary-like structures was fixed in 4%

paraformaldehyde in PBS for confocal microscopy. To gain more detailed insight into

the mating of the species, we also observed female morphology after they oviposited

dozens of eggs. Female#3, 4, and 5 (experienced copulation with Male#2) were

observed externally, and their abdomens were dissected in PBS, to investigate the

presence or absence of spermatheca, i.e., sperm storage organ, bursa copulatrix, i.e.,

copulatory pouch, where the penis reaches, and spermatophore, which is a capsule or

mass containing the spermatozoa and being formed in female bursa copulatrix during

mating (Matsuda 1976, Chapman et al. 2013). Isolated spermatheca, bursa copulatrix

and spermatophores were fixed with FAA fixative (ethanol: formalin: acetic acid =

15:5:1) at 25°C overnight and then preserved in 90% ethanol for histological
examination (details below). As a negative control of mating, we dissected and observed
the bursa copulatrix of two additional females (Female#14 and #15), which were field
collected and assumed to be unmated. External morphology and reproductive systems
of both sexes were recorded using a digital camera (FLOYD-4K, WRAYMER, Japan)
attached to a stereomicroscope (SZ61, Olympus, Japan).

Genotypic analysis on the rare males, females and their offspring
Microsatellite marker analyses were conducted to determine if the rare males originated
from specific lineages. A total of 19 individuals, including the six field-collected males
and one male from a breeding population, were genotyped using 11 polymorphic
microsatellite loci (Table S3, Suetsugu et al. 2023). To assess the genetic involvement of
rare males in the next generation, the genotypes of seven parental pairs and a total of 97
embryos from eggs laid before and after mating were also examined using the
microsatellite markers. Detection of male alleles in eggs laid after mating would
indicate the presence of the rare male's genes in the subsequent generation.

Genomic DNA was extracted from either one leg of the embryo or a portion of the
leg from adult individuals using the DNeasy Blood & Tissue Kit (QIAGEN, Germany),
following the manufacturer's instructions. PCR was conducted in two sets using
multiplex PCR (refer to Table S2). The PCR amplification was carried out in 5 µl
reactions using the QIAGEN Multiplex PCR Kit (QIAGEN, Germany). Each sample
reaction contained 10–20 ng of genomic template DNA, 2.5 µl of Multiplex PCR
Master Mix, 0.2 µM of fluorescently labeled forward primers, and 0.2 µM of reverse
primers (Table S2). The amplification conditions included an initial denaturation at
95 °C for 15 minutes, followed by 33 cycles of denaturation at 94 °C for 30 seconds,
annealing at 57 °C for 1.5 minutes, and extension at 72 °C for 1 minute, with a final
extension at 60 °C for 30 minutes. All thermal cycling conditions were performed using
the T100 thermal cycler (Bio-Rad Laboratories, Inc., USA). Product sizes were
determined using an ABI PRISM 3130 Genetic Analyzer and analyzed with
GeneMapper software (Applied Biosystems, USA).

To assess the kinship relations among genotypes, we constructed a
neighbor-joining tree based on Nei's genetic distance DA (Nei et al. 1983) using
POPULATION 1.2.30 (Langella 2007). The genetic contribution of males to the next
generation was assessed from two perspectives: 1) the inheritance of male-specific
alleles and 2) the presence of genotypic changes in eggs produced after mating that
could be attributed to recombination through Mendelian inheritance.

Microscopic observation of the sexual organs of males and females

To investigate sperm production of rare males, we carefully observed the sexual organs of rare males (Male#2 and #7). For confocal microscopic observation, a fixed testicular lobe or ovary-like structure was washed three times in 0.3% Triton X-100 in PBS (PBS-X) for 15 min for permeabilization. It was then stained with 4,6-diamidino-2-phenylindole (DAPI) (1 μg/mL; Dojindo, Japan) for the nuclei and Alexa Fluor™ 488 Phalloidin (66 nM; Thermo Fisher Scientific, USA) for the cytoskeleton (F-actin). After 1 h at room temperature (from 20 to 25 °C), they were washed three times with PBS-X for 15 min and mounted with VECTASHIELD antifade mounting medium (Vector Laboratories, USA). The morphology of the testis and ovarioles was visualized using fluorescent staining and differential interference contrast microscopy and observed by a confocal laser-scanning microscope (CLSM; FV1000, Olympus, Japan). For histological observation, another testis of Male#2 that was fixed with FAA was dehydrated and cleared using an ethanol-*n*-butanol series. The cleared samples were immersed and embedded in paraffin at 60°C. The paraffin blocks were polymerized at 4°C and cut into 5-μm thick sections using a microtome (RM2155, Leica, Germany). The sections were mounted on microscope slides coated with egg white-glycerin and stained using Delafield's Hematoxylin and Eosin (HE) staining. The stained slides were enclosed with Canada balsam. The slides were observed on an optical microscope (BX50, Olympus, Japan) and photographs were taken using a digital single-lens reflex camera (D7200, Nikon, Japan). To examine the decay of sexual organs in females, such as spermatheca, bursa copulatrix, and spermatophore, samples from three females fixed with FAA were also processed for histological observation.

Results

Investigation on the possibility of sexual reproduction in *R. mikado*

Observation on the mating behavior of rare males in *R. mikado*

To investigate whether the rare male mates with females and whether he displays some mating behavior, we observed the mating of the male collected from Chiba (Male#2) in detail. Both active mating behaviors, i.e., approaching, touching his antennae, and mounting to a female, and the insertion of the penis into the female genital opening were observed (Fig. 1B, C). Other males (Male#1, 3, and 4) also successfully copulated with females (Fig. S1).

<u>Morphology and anatomy of reproductive systems in *R. mikado*</u>

We confirmed that males had an external morphology same as the previous remarks,

such as a slender body, brown color and prominent white lines on both sides of the

thorax and abdomen (Fig. S2, Nagashima 2001, Yano et al. 2021). The males also

possessed an intromittent organ, i.e., penis, along the median of terminal. All males did

not exhibit abnormalities in their external morphology, such as characteristics of

intersex, as shown in *C. morosus* (Pehani 1925, Pijnacker 1964). We found that in a

male (Male#2) there was a typical reproductive system including a pair of testes,

accessory glands and seminal vesicles, as is typical in stick insects (Fig. S1, Fig 2A).

Another dissected male (Male#7) had male-type accessory glands and seminal vesicles

and a penis, but he did not possess testes but ovary-like structures (Fig. 3A, B). We also

dissected three females that mated with Male#2 after a collection of eggs and two virgin

females. Then we described the female characteristics (Fig. S3) and confirmed the

formation of spermatophore (Fig. S4). Micropyles, which are sperm gates; i.e., openings

for sperm entry of insect egg chorion, were observed in *R. mikado* and confirmed to be

present. We found that the eggs had a single micropyle, which is a typical characteristic

of stick insects (see SI Text, Fig. S5, S6) (Bedford 1978).

<u>Genotyping: their origin of males and their no genetic contribution to the next</u>

<u>generation</u>

Our microsatellite analysis indicated the parthenogenetic origin of the rare males, the

lack of kinship among them, and no genetic contribution of males in the next generation.

Six males showed different genotypes to each other, indicating that they were not close

relatives. Male#6 and #7 that were collected at a nearby location showed identical

genotypes. Phylogenetic analysis revealed that the genotypes of three males were

identical to those of females collected from nearby localities, suggesting

parthenogenetic origin of rare males rather than sexual reproduction (Fig. S7). We found

no paternal-specific alleles in eggs oviposited by females mated with males (mated eggs,

Table 1 and Table S5). The genotypic characteristics of the mated eggs were identical to

those of the virgin eggs, indicating the parthenogenetic origin of mated eggs.

The mated eggs were actually virgin. Therefore, we reasonably assumed all

analyzed 97 offspring from seven mothers were of parthenogenetic origin. Embryos

consistently showed almost identical multi-locus genotypes to their mother. For instance,

in loci such as Rmika021 and Rmika091, where the mother exhibits heterozygosity for

two alleles, the mated eggs also showed heterozygosity for the identical two alleles.

However, we found a single case of heterozygous to homozygous transition that should

be attributed to recombination (locus Rmika088, Table S5).

Sexual dysfunction in *R. mikado*

We demonstrated that the normally mating male did not become the father of his

partners' offspring. Moreover, we confirmed that Male#2 could form the

spermatophores in females' bursa copulatrix (Fig. S4) and the presence of sperm entry

on eggs (Fig. S6). To look into the cause of the fruitlessness, we focused on the

gametogenesis of the male. In a bisexual stick insect as a representative of typical

gametes of stick insects, the nuclear and F-actin staining showed that there were many

mature sperms with flagella, forming sperm bundles (see SI Text, Fig. S8A). Also, we

could recognize the sperm cysts containing male germ cells, and some of them

contained spermatids (acrosome was still not formed, Fig. S8B). We observed similar

features in the paraffin sectioning with HE staining as well (Fig. S8C). On the other

hand, we observed no mature sperms and spermatids in the testis of *R. mikado* Male#2.

While there were spermatogonia and some spermatocytes in the spermatogenic cysts,

the nuclei morphology was strongly deformed. We could not observe any sperm flagella

structure. In histological observation, some of the nuclei were deformed/condensed and

strongly stained with hematoxylin (Fig. 2), which are typical characteristics in apoptotic

cells (Nozaki & Matsuura 2021). We also assessed spermatophores isolated from mated

females. Many sperms were observed in the spermatophore produced by males of

bisexual species (Fig. S8D), yet we could not find any sperm-like structure in the *R.*

mikado spermatophore (Fig. S4), suggesting no sperms were transferred during the

ejaculation. Furthermore, we observed a male with ovary-like structures (Male#7) and

confirmed that the male actually had a pair of ovaries, consisting of the ovarioles

containing oocytes (Fig. 3).

We also observed spermatheca and bursa copulatrix of *R. mikado* females at the

histological level (Fig. S9). The spermatheca and bursa copulatrix of *R. mikado*

possessed a thin cuticle wall. Also, the spermatheca of *R. mikado* consisted of an

epithelial layer with many degenerative nuclei. These features were in contrast with

those of females in bisexual species, wherein spermatheca exhibited active secretion,

thickened cuticle and eosinophilic cytoplasm, and bursa copulatrix had thickened cuticle

(Fig. S9), as is typical in insects (Matsuda 1976).

Discussion

The evolutionary persistence of parthenogenesis has long been discussed (Judson &

Normark 1996, Schurko et al. 2009), and recent studies have shown evidence of occasional sexual reproduction in the parthenogenetic species that were thought to have reproduced solely parthenogenetically for millions of years (Vakhrusheva et al. 2020, Boyer et al. 2021, Freitas et al. 2023) (but see [Kearney et al. 2022]). Therefore, obligately parthenogenetic species with a long history may still have the possibility of utilizing or returning to sexual reproduction. However, our study revealed the irreversible parthenogenesis in the Japanese common stick insect *Ramulus mikado*, which had also been an "old" parthenogen (Suetsugu et al. 2023). Both sexes of *R. mikado* had seemingly complete genital organs, including a sperm reservoir in females and a penis in males, and showed usual mating behaviors (Fig. 1, S1-4, Video 1). Nevertheless, we could not detect any paternity of the rare males for offspring laid by their mating partners (Table 1, S1). The females used for histological observation showed that they have sexual organs with signs of degeneration (Fig. S9). Furthermore, we found reproductive dysfunction in dissected males; one did not have any normally matured sperms and spermatids, and only showed abnormal spermatocytes (Male#2, Fig. 2) despite being sacrificed just after mating with females. Another male had ovaries with oocytes instead of testes (Male#7, Fig. 3). All these results strongly suggest that *R. mikado* could no longer return to sexual reproduction. Although not easy to verify, the plasticity of reproductive modes is critical to explain how old or ancient parthenogenetic lineages can avoid the long-term costs of asexuality (Boyer et al. 2021, Freitas et al. 2023). In this mean, the present study provided unique insights into the evolution and the consequence of the loss of sex.

This study supported the idea that the parthenogenesis of *R. mikado* has a long history as an asexual lineage (estimated to be 0.34–0.51 Myr) without cryptic recent gene flow (Suetsugu et al. 2023). Then, how do they overcome the expected long-term cost of asexuality? Our SSR analysis, consistent with the findings of Suetsugu et al. (2023), revealed high heterozygosity in approximately half of the loci and zero or near-zero heterozygosity in the remaining ones (Table S4). We also revealed that mothers and offspring possessed almost identical genotypes, with rare transitions from heterozygous to homozygous due to recombination (Table S5). The result indicates the automixis with central fusion as a mechanism of parthenogenesis in *R. mikado*, avoiding the loss of heterozygosity in individuals and preserving the accumulation of genetic variation across generations, which might explain the long-persistence of the parthenogenesis (Schwander & Crespi 2009). Alternatively, but not exclusively, the historical long-dispersal event, which we have recently shown in *R. mikado*, resulting in a wide distribution range (Suetsugu et al. 2023), might hint at the long history of their

asexuality. Moreover, it is well known that some phasmids including *R. mikado* exhibit
periodical outbreaks in natural forests or plantations (Bedford 1978, Baker 2015, Yano
et al. 2021). The long-term disadvantage of asexual over sexual reproduction such as a
higher extinction rate should be dependent on the effective population size and may
become negligible when population size gets huge (Ross et al. 2013). Therefore, these
factors should be taken into account for future studies on the persistence of
parthenogenesis.

"Rare males" in the parthenogenetic species *R. mikado* were not functional for
reproduction, i.e., sterile, and did not contribute to the next generations. So far,
functionality of "rare males" in parthenogenetic species has been assessed among
several animals including stick insects (Vandel 1931, 1934, Pijnacker 1964, Oliver 1971,
Oliver et al. 1973, Pijnacker et al. 1981, Itow et al. 1984, MacDonald & Browne 1987,
Zchori-Fein et al. 1992, Kajita 1993, Hunter 1999, Giorgini 2001, Adachi-Hagimori et
al. 2011, Schwander et al. 2013, Jalinsky et al. 2020, Kampfraath et al. 2020). However,
such males were potentially functional: they could have some normally mature sperms,
suggesting that the spermatogenesis is *not entirely* degenerated. Therefore, to our
knowledge, this is the first instance of the complete decay of spermatogenesis: the
production of abnormal spermatids only and no genetic contribution to sexual
reproduction. Histological observations of spermatogenesis were conducted on a single
male (Fig. 2), suggesting that the extent of dysfunction may vary among individual
males, each of which is independently produced through parthenogenesis (Fig. S7). All
males investigated did not have paternity, and another dissected male (Male#7) in this
study was intersex in the reproductive system; the male had no testes but ovaries, while
it possessed other male reproductive system (seminal vesicles and male accessory
glands) (Fig. 3). Moreover, a dissected male in a previous report did not possess
testis-like structures (Yano et al. 2021). Therefore, it is likely that functional impairment
in the gonad differentiation/gametogenesis generally occurs in the males of the
predominantly parthenogenetic species, *R. mikado*. In the future studies, it would be
worthwhile to examine whether the genetic basis controlling such sexual development
undergoes degeneration, which should lead to the development instability in males.

van der Kooi and Schwander (2014) proposed that male traits that initially
degenerate in asexual lineages may be related to spermatogenesis. Our findings indicate
that spermatogenesis proceeds to degeneration faster than other sexual traits, such as
male genitalia and mating behavior, in *R. mikado*. These results support the hypothesis
on the decay process of male traits under parthenogenesis. Previous studies also showed
partially abnormal spermatogenesis in rare males with normal external morphology and

behavior in some parthenogenetic animals (MacDonald & Browne 1987, Pijnacker 1964). It has been suggested that spermatogenesis slowly degenerated in parthenogenetic lineages due to the sole effect of genetic drift after the relaxation of stabilizing selection (van der Kooi & Schwander 2014). Based on this assumption, it is plausible that sperm production in the predominantly parthenogenetic lineages degenerates through the following process: 1) the normal spermatogenesis is retained, 2) both the abnormal and normal sperms coexist (the partial decay of spermatogenesis), and 3) the spermatogenesis completely disrupted, and then males eventually become fruitless. Hence, the fruitless males in *R. mikado* might represent the final stage of spermatogenesis decay, supporting the idea that parthenogenesis in the species has a long history as an asexual lineage.

Overall, this study provides a unique and rare example of the irreversible evolution of parthenogenesis. In *R. mikado*, despite the occurrence of males, which display not only characteristic external morphology but also robust mating behavior, the male sex is non-functional and serves as a mere vestige. The finding on the fruitlessness of rare males in *R. mikado* would provide crucial progress in the studies on the decay of male sexual traits after transitioning to asexuality. Although completely decayed spermatogenesis in rare males has been predicted, this is the first report of such instances. Obtaining males in predominantly or obligate parthenogenetic lineages is challenging (Brock et al. 2018); nevertheless, the accumulation of case reports can further improve our understanding of the evolutionary fate of both asexual lineages themselves and sexual traits in both sexes under asexuality.

Acknowledgement

We thank Professors S. Shigenobu and T. Niimi of NIBB for the experimental space and lab equipment. For the sample collection, we thank S. Tanaka, Y. Hayashi, E. Kato, A. Aono, S. Yanagisawa, T. Uchifune, M. Matsumoto, S. Kitamura, M. Yoshida, J. Haga, and T. Hiramatsu, T. Horie and M. Yoshitomi of The Takasegawa Lovers Association, and Professor K. Tojo of Shinshu University. We deeply appreciate T. Takagi and K. Sato for technical advice. We also thank T. Nawa of Nawa Insect Museum, P. Brock and T. Büscher for helpful advice and extensive information for stick insects.

Competing interests

We declare we have no competing interests.

Author contributions

T.N.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, validation, visualization, writing—original draft, writing—review and editing; Y.C.: conceptualization, investigation, methodology, visualization, writing—original draft, writing—review and editing; K.Y.: funding acquisition, investigation, methodology, resources, visualization, writing—original draft, writing—review and editing; R.S.: data curation, formal analysis, investigation, visualization, writing—review and editing; K.S.: conceptualization, funding acquisition, investigation, resources, validation, writing—original draft, writing—review and editing; S.K.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, validation, visualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Data accessibility

All SSR analysis data are available in the supplementary information.

Funding

This work was financially supported by the Japan Society for the Promotion of Science to K. S. (Challenging Exploratory Research No.18K19215), T. N. (Research Fellowship for Young Scientists No. 19J01756) and K. Y. (No. 21J01422). This work was also supported by Competitive Research Funds for Fukushima University Faculty.

Literature cited

Adachi-Hagimori, T., Miura, K. & Abe, Y. (2011). Gene flow between sexual and asexual strains of parasitic wasps: a possible case of sympatric speciation caused by a parthenogenesis-inducing bacterium. Journal of Evolutionary Biology 24, 1254–1262.

Alavi, Y., van Rooyen, A., Elgar, M. A., Jones, T. M., & Weeks, A. R. (2018). Novel microsatellite markers suggest the mechanism of parthenogenesis in *Extatosoma tiaratum* is automixis with terminal fusion. Insect Science. 25, 24–32.

Avise, J. (2008). Clonality: The Genetics, Ecology, and Evolution of Sexual Abstinence in Vertebrate Animals, Oxford Academic, New York.

Baker, E. (2015). The worldwide status of phasmids (Insecta: Phasmida) as pests of agriculture and forestry, with a generalised theory of phasmid outbreaks. Agriculture & Food Security, 4(1), 22.

Bast, J., Parker, D. J., Dumas, Z., Jalvingh, K. M., Tran Van, P., Jaron, K. S., Figuet, E., Brandt, A., Galtier, N., & Schwander, T. (2018). Consequences of asexuality in natural populations: insights from stick insects. Molecular Biology and Evolution 35, 1668–1677.

Bedford, G. O. (1978). Biology and ecology of the Phasmatodea. Annual Review of Enmorology. 23, 125–149.

Bell, G. (1982). The Masterpiece of Nature: The Evolution and Genetics of Sexuality. University of California Press, San Francisco.

Blacket, M. J., Robin, C., Good, R. T., Lee, S. F., & Miller, A. D. (2012). Universal primers for fluorescent labelling of PCR fragments—an efficient and cost-effective approach to genotyping by fluorescence. Molecular Ecology Resources. 12, 456–463.

Boyer, L., Jabbour-Zahab, R., Mosna, M., Haag, C. R., & Lenormand, T. (2021). Not so clonal asexuals: Unraveling the secret sex life of *Artemia parthenogenetica*. Evolution Letters, 5(2), 164–174.

Brock, P. D. (2000). *A complete guide to breeding stick and leaf insects*. TFH Publications.

Brock, P. D., Lee, M., Morgan-Richards, M., & Trewick, S. A. (2018). Missing stickman found: the first male of the parthenogenetic New Zealand phasmid genus *Acanthoxyla* Uvarov, 1944 discovered in the United Kingdom, Atropos Vol 60: 16–23.

Butlin, R. (2002). The costs and benefits of sex: new insights from old asexual lineages. Nature Reviews Genetics, 3(4), 311–317.

Chapman, R. F., Simpson, S. J., Douglas, A. E. (2013). The Insects: Structure and Function, Fifth ed. Cambridge University Press, Cambridge.

Carson, H. L., Chang, L. S., & Lyttle, T. W. (1982). Decay of female sexual behavior under parthenogenesis. Science, 218(4567), 68–70.

D'Souza, T. G., & Michiels, N. K. (2010). The costs and benefits of occasional sex: theoretical predictions and a case study. Journal of Heredity, 101(suppl_1), S34–S41.

Faircloth, B. C. (2008). MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus specific primer design. Molecular ecology resources, 8(1), 92–94.

Freitas, S., Parker, D. J., Labédan, M., Dumas, Z., & Schwander, T. (2023). Evidence for cryptic gene flow in parthenogenetic stick insects of the genus *Timema*, Proceedings of the Royal Society B: Biological Sciences, 290, 20230404

Fong, D. W., Kane, T. C., & Culver, D. C. (1995). Vestigialization and loss of nonfunctional characters. Annual Review of Ecology and Systematics, 26(1), 249–268.

Giorgini, M. (2001). Induction of males in thelytokous populations of *Encarsia meritoria* and *Encarsia protransvena*: a systematic tool. BioControl 46, 427–438.

Hunter, M. S. (1999). The influence of parthenogenesis-inducing *Wolbachia* on the oviposition behaviour and sex-specic developmental requirements of autoparasitoid wasps. Journal of Evolutionary Biology 12, 735–741.

Itow, T., Kobayashi, K., Kubota, M., Ogata, K., Imai, H. T., & Crozier, R. H. (1984). The reproductive cycle of the queenless ant *Pristomyrmex pungens*. Insectes Sociaux, 31(1), 87–102.

Jalinsky, J., Logsdon, J. M., & Neiman, M. (2020). Male phenotypes in a female framework: Evidence for degeneration in sperm produced by male snails from asexual lineages. Journal of Evolutionary Biology, 33, 1050–1059.

Jaron, K. S., Parker, D. J., Anselmetti, Y., Tran Van, P., Bast, J., Dumas, Z., Figuet, E., François, C. M., Hayward, K., Rossier, V., Simon, P., Robinson-Rechavi, M., Galtier, N., & Schwander, T. (2022). Convergent consequences of parthenogenesis on stick insect genomes. Science advances, 8(8), eabg3842.

Judson, O. P., & Normark, B. B. (1996). Ancient asexual scandals. Trends in ecology & evolution, 11(2), 41–46.

Kajita, H. (1993). Induction of males in the thelytokous wasp *Encarsia formosa* Gahan (Hymenoptera: Aphelinidae). Applied Entomology and Zoology, 28(1), 115–117.

Kampfraath, A. A., Dudink, T. P., Kraaijeveld, K., Ellers, J., & Zizzari, Z. V. (2020). Male sexual trait decay in two asexual springtail populations follows neutral mutation accumulation theory. Evolutionary Biology, 47, 285–292.

Kearney, M. R., Jasper, M. E., White, V. L., Aitkenhead, I. J., Blacket, M. J., Kong, J. D., ... & Hoffmann, A. A. (2022). Parthenogenesis without costs in a grasshopper with hybrid origins. Science, 376(6597), 1110–1114.

Langella O., (2007). Populations ver. 1.2. 30: population genetic software (individuals or populations distances, phylogenetic trees). http://bioinformatics.org/~tryphon/populations/

Leach, I. M., Pannebakker, B. A., Schneider, M. V., Driessen, G., Van De Zande, L., & Beukeboom, L. W. (2009). Thelytoky in Hymenoptera with *Venturia canescens*

and *Leptopilina clavipes* as case studies. Lost sex: the evolutionary biology of parthenogenesis, 347–375.

MacDonald, G. H. & Browne, R. A. (1987). Inheritance and reproductive role of rare males in a parthenogenetic population of the brine shrimp, *Artemia parthenogenetica*. Genetica 75, 47–53.

Machida, R., Kano, Y., & Yominaga, O. (2016). *The standard of Polyneoptera in Japan*. Gakken Plus, Tokyo. (In Japanese.)

Matsuda, R. (1976). Morphology and Evolution of the Insect Abdomen. Oxford: Pergamon. 532 pp.

Matsuura, K. (2017). Evolution of the asexual queen succession system and its underlying mechanisms in termites. Journal of Experimental Biology, 220(1), 63–72.

Meirmans, P. G., & Hedrick, P. W. (2011). Assessing population structure: FST and related measures. Molecular ecology resources. 11, 5–18.

Muller, H. J. (1949). The Darwinian and modern conceptions of natural selection. Proceedings of the American Philosophical Society, 93(6), 459–470.

Morgan-Richards, M. A. R. Y., Trewick, S. A., & Stringer, I. A. (2010). Geographic parthenogenesis and the common tea☐tree stick insect of New Zealand. Molecular ecology, 19(6), 1227–1238.

Nagashima, Y. (2001). The Walking Sticks (Phasmida Insect) in Niigata Prefecture. Bulletin of Niigata Seiryo University, 1, 3–10 (in Japanese).

Nakano, M., Morgan-Richards, M., Godfrey, A. J. R., & Clavijo McCormick, A. (2019). Parthenogenetic females of the stick insect *Clitarchus hookeri* maintain sexual traits. Insects, 10(7), 202.

Nakamura, K., & Fukushima, Y. (2022). Seasonal and geographical adaptations in the parthenogenetic stick insect, *Ramulus mikado* (Phasmatodea: Phasmatidae). European Journal of Entomology, 119.

Nei M, Tajima F, & Tateno Y. (1983). Accuracy of estimated phylogenetic trees from molecular data – II. Gene frequency data. Journal of Molecular Evolution. 19, 153–170. Doi:10.1007/BF02300753

Nozaki, T., & Matsuura, K. (2021). Oocyte resorption in termite queens: Seasonal dynamics and controlling factors. Journal of Insect Physiology, 131, 104242.

Okada M. (1999). All About Stick Insects. Tombow Publishing, Osaka, 55 pp. (in Japanese).

Oliver, J. H. (1971). Parthenogenesis in mites and ticks (Arachnida: Acari). American Zoologist 11, 283–299.

Oliver, J. H., Tanaka, K. & Sawada, M. (1973). Cytogenetics of ticks (Acari: Ixodoidea). 12. Chromosome and hybridization studies of bisexual and parthenogenetic *Haemaphysalis longicornis* races from Japan and Korea. Chromosoma 42, 269–288.

Pantel, J. (1917). Description de *Carausius* nouveaux (Orth. Phasmidae), Et notes sur les *Carausius* de l'Inde méridionale, Annales de la Société entomologique de France, 86, 267–306.

Peakall, R., & Smouse, P. E. (2012). GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. Bioinformatics 28, 2537–2539

Pehani, H. (1925). Die Geschlechtszellen der Phasmiden. Zugleich ein Beitrag zur Fortpflanzungsbiologie der Phasmiden. Zeitschrift für wissenschaftliche Zoologie, 125, 167–238.

Pijnacker, L. P. (1964). The cytology, sex determination and parthenogenesis of *Carausius morosus* Br. Dr. thesis, University of Groningen.

Pijnacker, L. P., Ferwerda, M. A. & Helle, W. (1981). Cytological investigations on the female and male reproductive system of the parthenogenetic privet mite *Brevipalpus obovatus* Donnadieu (Phytoptipalpidae, Acari). Acarologia 22, 157–163

Rayner, J. G., Sturiale, S. L. & Bailey, N. W. (2022). The persistence and evolutionary consequences of vestigial behaviours. Biological Reviews, 97(4), 1389–1407.

Ross, L., Hardy, N. B., Okusu, A. & Normark, B. B. (2013). Large population size predicts the distribution of asexuality in scale insects. Evolution, 67(1), 196–206.

Scali, V. (2009). Metasexual stick insects: model pathways to losing sex and bringing it back. Lost sex: the evolutionary biology of parthenogenesis, 317–345.

Scali, V. (2013). A "ghost" phasmid appearance: the male *Bacillus atticus* (Insecta: Phasmatodea). Italian Journal of Zoology, 80(2), 227–232.

Schwander, T. & Crespi, B. J. (2009). Multiple direct transitions from sexual reproduction to apomictic parthenogenesis in *Timema* stick insects. Evolution, 63(1), 84–103.

Schwander, T., Henry, L., & Crespi, B. J. (2011). Molecular evidence for ancient asexuality in *Timema* stick insects. Current Biology, 21(13), 1129–1134.

Schwander, T., Crespi, B. J., Gries, R., & Gries, G. (2013). Neutral and selection-driven decay of sexual traits in asexual stick insects. Proceedings of the Royal Society B: Biological Sciences. 280, 20130823.

Schurko, A. M., Neiman, M., & Logsdon, J. M. (2009). Signs of sex: what we know and

how we know it. Trends in ecology & evolution, 24(4), 208–217.

Simon, J. C., Delmotte, F., Rispe, C., & Crease, T. (2003). Phylogenetic relationships
between parthenogens and their sexual relatives: the possible routes to
parthenogenesis in animals. Biological Journal of the Linnean Society, 79(1),
151–163.

Sinety, R. de (1901). Recherches sur la biologie et l'anatomie de Phasmes. - La Cellule
19: 118–278

Suetsugu, K., Nozaki, T., Hirota, S., Funaki, S., Ito, K., Isagi, Y., Suyama, Y., & Kaneko,
S. (2023). Phylogeographical evidence for historical long-distance dispersal in
the flightless stick insect *Ramulus mikado*, Proceedings of the Royal Society B,
290(2008), 20231708.

Suomalainen, E. (1962). Significance of parthenogenesis in the evolution of insects.
Annual Review of Entomology, 7(1), 349–366.

Vakhrusheva, O. A., Mnatsakanova, E. A., Galimov, Y. R., Neretina, T. V., Gerasimov, E.
S., Naumenko, S. A., ... & Kondrashov, A. S. (2020). Genomic signatures of
recombination in a natural population of the bdelloid rotifer *Adineta vaga*. Nature
Communications, 11(1), 6421.

Vandel, A. (1931). On the existence of males of parthenogenetic origin and its genotypic
constitution of parthenogenetic females of *Trichoniscus* (*Spiloniscus*) *elisabethae*
Herold (Crustaceans, Isopods). Comptes Rendus Hebdomadaires des Seances de
l'Academie des Sciences 193, 887–889.

Vandel, A. (1934). La parthenogenese géographique. II. Lesmâles triploïdes d'origine
parthenogénétique de *Trichoniscus* (*Spiloniscus*) *elisabethae*. Bulletin Biologique
de la France et de la Belgique 68, 419–463.

van der Kooi, C. J., & Schwander, T. (2014). On the fate of sexual traits under
asexuality. Biological Reviews, 89(4), 805–819.

Voelker, R. A., & Kojima, K. I. (1971). Fertility and fitness of XO males in *Drosophila* I.
Qualitative study. Evolution, 119–128.

Yamaguchi, H., & Nakamura, K. (2015). Effects of environmental factors on the
regulation of egg diapause in the walking-stick insect, *Ramulus*
irregulariterdentatus (Phasmatodea: Phasmatidae). European Journal of
Entomology, 112(1), 35–40.

Yano, K., Ozaki, T., Suzuki, T., Yamazaki, H., Nasuno, M., Degawa, Y., & Tojo, K.
(2021). Outbreak of the stick insect, *Ramulus mikado* (Phasmatodea,
Phasmatidae), in the Akashina area of Japan (Azumino City, Nagano Prefecture).
Entomological Science 24, 196–200.

Zchori-Fein, E., Roush, R. T. & Hunter, M. S. (1992). Male production induced by
641   antibiotic treatment in *Encarsia formosa* (Hymenoptera: Aphelinidae), an asexual
642   species. Experientia 48, 102–105.

Table

645 **Table 1.** Selected genotyping results of paternity test

Pair	Category[†]	SSR locus[‡]				
		Rmika056	Rmika021	Rmika033	Rmika077	Rmika050
M#1-F#2	Male	120/120	243/**259**	113/**117**	206/212	223/223
	Female[*]	120/120	243/267	111/113	206/212	223/223
	Mated egg (10)	120/120	243/267	111/113	206/212	223/223
M#2-F#3	Male	**126/126**	243/259	113/**115**	**204/216**	223/223
	Female	120/120	243/259	111/113	214/214	223/223
	Mated egg (10)	120/120	243/259	111/113	214/214	223/223
M#2-F#4	Male	**126/126**	243/259	113/**115**	**204/216**	223/223
	Female	120/120	243/259	111/113	214/214	223/223
	Mated egg (10)	120/120	243/259	111/113	214/214	223/223
M#2-F#5	Male	**130/130**	243/259	113/**115**	**204/216**	223/223
	Female	120/120	243/259	113/117	206/212	223/223
	Mated egg (8)	120/120	243/259	113/117	206/212	223/223
M#3-F#6	Male	120/120	243/243	**111/113**	**212/218**	**223/223**
	Female	120/120	243/289	115/115	208/216	225/225
	Mated egg (15)	120/120	243/289	115/115	208/216	225/225

[†]In parentheses, *n* (number) indicates the *n* of eggs examined (these eggs showed

completely the same genotype as each other)

[‡]Locus that show male-specific alleles (**bold**) are presented (for full data, see Table S5)

[*]Estimated from the genotype of eggs

Figures and captions

Fig. 1. Japanese parthenogenetic stick insect *Ramulus mikado*. **A** photo of a rare male
and a female (both adults). **B** Successful mating. A male (Male#2) mounts on a female
(Female#3), grips the female abdominal tips using "claspers" and inserts their penis into
the female gonopores. **C** A more magnified image of abdominal tips during the mating.
Transparent greenish penis (external genitalia) is inserted into the female gonopore (for
detailed information on morphology and anatomy, see Fig. S2 and S3).

Fig. 2. Detailed observation on the testes of a rare male (Male#2). **A, B** Confocal

images of testes and spermatogenic cysts in a rare male in *R. mikado*. Morphology of

testis, cysts, and spermatids was observed by DAPI-Phalloidin staining or differential

interference contrast (DIC) technique. DNA and F-actin were stained by DAPI (blue or

gray) and Phalloidin (green), respectively. In Male#2 testis, no mature sperm forming

bundles was observed (A). Spermatogenesis seemed to be deformed in the male; in

spermatogenic cysts, there were empty spaces (the red arrow), and the nuclei exhibited

irregular shapes and still large (red arrowheads) (B). **C** Histological images of a testis

and spermatogenic cysts in the same male (Male#2), visualized by EH staining. Almost

all nuclei in the spermatogenic cysts were strongly stained and showed an irregular

shape. Sperm flagella was not observed.

Fig. 3. Detailed observation on the reproductive system of a rare male (Male#7). **A, B** The male possessed ovaries while the structure was irregular (e.g., the reduced number or deformed shape of ovarioles) (A). The ovaries were connected with seminal vesicle and male-type accessory glands in the position near the gonopore (B). Abbreviations used; Sv: seminal vesicle, Ags: accessory glands. **C** Confocal images of ovarioles of a rare male (Male#7), which were visualized by DAPI-Phalloidin staining or differential interference contrast (DIC) technique. DNA and F-actin were stained by DAPI (blue) and Phalloidin (green), respectively. Some of the ovarioles had oocytes containing the yolk, which has strong green-autofluorescence.