Global Pannexin 1-deletion increases tumor-infiltrating lymphocytes in the BRAF/Pten
mouse melanoma model.

Rafael E. Sanchez-Pupo[1], Garth A. Finch[1], Danielle E. Johnston[1], Heather Craig[2], Rober Abdo[1],
Kevin Barr[1], Steven Kerfoot[2], Lina Dagnino[3,4], and Silvia Penuela[1,4]

[1]Department of Anatomy and Cell Biology, Western University, London, Ontario N6A 5C1,
Canada, [2]Department of Microbiology and Immunology, [3]Department of Physiology and
Pharmacology, and [4]Department of Oncology, Division of Experimental Oncology, Schulich
School of Medicine and Dentistry, Western University, London, Ontario N6A 5C1, Canada.

Keywords: Pannexin 1, PANX1, melanoma, immune infiltration, T cell

Running Title: Panx1 influences T cell infiltration in melanoma tumors

Additional Information

Financial Support: This study was supported by a Canadian Institutes of Health Research CIHR
Project grant to Silvia Penuela and Lina Dagnino (FRN 153112).

Corresponding author: Dr. Silvia Penuela. Dept. of Anatomy and Cell Biology, Western
University, London, Ontario N6A 5C1, Canada. Tel.: 519-661-2111, ex. 84735; E-mail:
spenuela@uwo.ca.

Conflict of Interest: The authors declare no potential conflicts of interest of any kind.

ABSTRACT:

Immunotherapies for malignant melanoma seek to boost the anti-tumoral response of CD8$^+$ T cells but have a limited patient response rate, in part due to limited tumoral immune cell infiltration. Genetic or pharmacological inhibition of Pannexin 1 (PANX1) channel-forming protein is known to decrease melanoma cell tumorigenic properties *in vitro* and *ex vivo*. Here, we crossed *Panx1* knockout (*Panx1$^{-/-}$*) mice with the inducible melanoma model: *Braf^{CA}*, *Pten^{loxP}*, *Tyr::CreER^{T2}* (BPC). We found that deleting the *Panx1* gene in mice does not reduce BRAF(V600E)/Pten-driven primary tumor formation or improve survival. Notably, BPC-*Panx1$^{-/-}$* mice tumors exhibited a significant infiltration of CD4$^+$, CD8$^+$ T lymphocytes, and increased Granzyme B$^+$ cells with no changes in the expression of early T cell activation marker CD69, LAG-3 checkpoint receptor or PD-L1 in tumors when compared to BPC-*Panx1$^{+/+}$* genotype. Our results suggest that although *Panx1* deletion does not overturn the aggressive *BRAF*/*Pten*-driven melanoma progression, *in vivo*, it does increase the infiltration of effector immune T cell populations in the tumor microenvironment. We propose that PANX1-targeted therapy could be explored as a strategy to increase tumor-infiltrating lymphocytes to boost anti-tumor immunity.

INTRODUCTION:

Malignant melanoma is a type of skin cancer that develops from melanocytes and becomes a life-threatening disease when it spreads in the body [1]. Depending on the diagnosis and the mutational landscape of the tumor biopsies (e.g., *BRAF* mutation status), targeted therapy with mutant kinase inhibitors or immunotherapies with monoclonal antibodies against immune checkpoint inhibitors are currently being used for melanoma treatment. The use of immune checkpoint inhibitors enhances anti-tumoral T cell-mediated immune responses, but unfortunately, about ~60% of melanoma patients do not respond to this type of therapy [2-4]. Regardless of the treatment options, their effectiveness is usually dampened due to the insufficient effector activity of tumor-infiltrating lymphocytes (TILs) [5], recurrence, and development of multiple resistance mechanisms [6, 7]; therefore, innovative therapy concepts are still warranted.

Pannexin1 (PANX1) is a channel-forming membrane glycoprotein that mediates the passage of nucleotides, ions, and other metabolites involved in intercellular communication [8]. We have previously demonstrated that PANX1 expression correlates with tumor cell aggressiveness in isogenic mouse melanoma cell lines, and its knockdown reduces melanoma tumor size and metastasis to the liver in chick chorioallantoic membrane (CAM) xenografts [9]. More recently, we showed that PANX1 is highly expressed in human melanomas, and genetic or pharmacological targeting of PANX1 decreases the tumorigenic properties of melanoma cells [10]. Mechanistically, PANX1 regulates melanoma cell growth and metabolism through direct interaction with β-catenin and modulation of the Wnt signaling pathway [11]. All these findings place PANX1 as an attractive target for preclinical melanoma research, but it is still unclear whether the silencing of this gene would impact melanoma development *in vivo*.

In addition, PANX1 influences the inflammatory response [12-17] and is expressed in different subtypes of immune cells like macrophages and lymphocytes [12, 18, 19]. Previous evidence shows that PANX1-mediated adenosine-5'-triphosphate (ATP) permeability enables activation and functional enhancement of T cells [16, 17]. On the other hand, the release of ATP through caspase-cleaved PANX1 channels in apoptotic leukemic lymphocytes has been shown to induce the recruitment of inflammatory cells (e.g., monocytes)[12]. Paradoxically, the caspase-dependent PANX1 channel opening in dying macrophages and lymphocytes releases metabolites

that suppress, *in vivo,* proinflammatory signals of myeloid cells in the surrounding tissue [8].

Considering the broad PANX1 expression and different cell types present in the tumor

microenvironment (TME), no studies have specifically addressed whether PANX1 modulates the

recruitment of immune cells in the melanoma TME [13].

Previous histological and pathological examinations of a constitutive global *Panx1* knockout

mouse (*Panx1$^{-/-}$*, *Panx1*KO)(developed by Genentech) [15] showed no overt phenotypes.

However, using the same *Panx1$^{-/-}$* mouse strain, we have found differences in dorsal skin

thickness and delayed wound healing [9, 20]. Moreover, our studies show that in *Panx1* wildtype

mice, normal melanocytes express low levels of PANX1 which are upregulated in melanoma

cells [9]. To gain new insights into the roles of PANX1 in the context of skin cancer we set out

to test the combination of this *Panx1$^{-/-}$* mouse line with a cutaneous melanoma mouse model.

In this work, we crossed the global *Panx1$^{-/-}$* mouse strain with the inducible melanoma model

BRaf$^{CA/+}$, Pten$^{LoxP/Loxp}$, Tyr::CreER^{T2} (abbreviated here as BPC) that harbors two known

oncogenic driver mutations [21-23]. Due to the role of PANX1 in regulating tumor growth and

the immune response [12-17], we sought to evaluate the effects of the *Panx1* germline deletion

on melanoma progression and the tumor immune infiltration of this new hybrid mouse model.

Our results showed that *Panx1* global deletion did not reduce the strong BRAF/Pten-driven

melanoma progression but increased the tumor infiltration of effector immune T cell populations

and Granzyme B^{+} cells. Moreover, gene expression analysis showed that *Panx1* germline

deletion did not impair early T cell immune activation nor affect the expression of checkpoint

receptors in melanoma tumors. Interestingly, female PANX1-deficient BPC mice exhibited

increased spleen size compared to their male counterparts suggesting differences in their immune

phenotype. We anticipate that targeting PANX1 in melanoma may increase the homing of

effector T-lymphocytes to melanoma tumors and improve the effect of immunotherapies.

Materials and Methods

Mouse breeding and genotyping

C57BL/6, Tyr::CreER^{T2}; Braf$^{CA/+}$; Pten$^{LoxP/LoxP}$ (BPC) mice were obtained from Jackson

Laboratory, (Stock # 013590). Mice were kept on a 12:12 h light:dark cycle with normal chow

and water available *ab libitum*. For deletion of *Panx1*, BPC mice were crossed with the global

Panx1$^{-/-}$ mice in a C57BL/6N background strain [20, 15] (Genentech Inc., San Francisco, CA).

Genotyping was performed on genomic DNA derived from the toes using PCR as described in

[22].

Melanoma tumor induction, monitoring and sample collection

All experiments performed on the mice were approved by the Animal Care Committee of the

University Council on Animal Care at the University of Western Ontario, London ON, Canada

(UWO AUP# 2019-070). 3–4-weeks-old BPC mice were used for tumor induction experiments:

Panx1$^{+/+}$ (7 females and 8 males) and *Panx1*$^{-/-}$ (9 females and 4 males). Cutaneous melanomas

were induced with topical application of 4-hydroxytamoxifen (4-HT) (Sigma) on the skin of the

lower back as per protocol in [22]. Mouse body weight and tumor dimensions were monitored

every 3–4 days as soon as the tumors were visible. Measurements were done with a digital

caliper, and tumor volume was calculated using the modified ellipsoid formula $V = (W(\times 2) \times$

$L)/2$, where V is tumor volume, L and W are the longest (length) and shortest (width) tumor

diameters, respectively. The endpoint was considered when the tumor volume (or total combined

tumor volume, in case of multiple tumors per animal) reached 2.0 cm^3, tumors showed signs of

heavy ulceration, or the animals had poor body condition. Euthanasia was carried out using

carbon dioxide asphyxiation followed by cervical dislocation. The spleens were weighed and

measured using a ruler. Draining inguinal lymph nodes compromised with pigmented lesions

were classified as metastatic sites and counted per mouse. Tissue samples were fixed overnight

with Periodate-Lysine-Paraformaldehyde (PLP) fixative or flash-frozen in liquid nitrogen and

stored at –80 ºC until processing for RNA extraction.

Tissue Processing, RNA Extraction, real time-qPCR

Frozen tissues/tumor samples were ground in liquid nitrogen and total RNA was extracted using

TRIzol™ Reagent (Invitrogen) according to the manufacturer's protocol and purified using the

RNAeasy® Plus Mini kit (Qiagen). RNA concentration was determined using an Epoch

Microplate Spectrophotometer (BioTek®). cDNA was generated using the High-Capacity cDNA

Synthesis Kit (Thermo Fisher Scientific; REF 4368814) in a T100™ Thermal Cycler (Bio-Rad).

SYBR®Green Real-Time PCR Master Mix (Bio-Rad) was used along with the primers listed in

Table 1, where mouse *gapdh* and *18S* were the housekeeping gene controls. Real-time qPCR

was performed in a CFX96 Touch TM Real-Time PCR Detection System (Bio-Rad). All the

assays involved at least three technical replicates (n=3), and the relative mRNA expression

analysis was determined using the ΔΔCt method calculated in the Bio-rad CFX Maestro

Software, ver 1.1 (Bio-rad). Between assays, a non-tamoxifen-treated skin BPC-*Panx1*[+/+] sample

was used as the control for inter-plate variation.

Tissue preparation, cryo-sectioning, and immunofluorescence staining

Tumor tissues samples that were fixed in PLP (1% PFA, 0.1 M L-Lysine, 0.01 M sodium

periodate) overnight at 4°C were washed three times for 5 mins in phosphate buffer (0.075 M

sodium phosphate dibasic and 0.025 M sodium phosphate monobasic) and incubated sequentially

in 10%, 20%, and 30% sucrose phosphate solutions, each overnight at 4°C. Tumors were then

embedded and frozen in Fisher Healthcare™ Tissue-Plus™ O.C.T. Compound (Fisher Scientific

Cat.# 23-730-571) and stored at -80 °C until sectioning. Eight-micron-thick serial cryostat

sections were immuno-stained with the fluorescent-conjugated antibodies diluted in 1% BSA,

0.1% Tween-20, and 5% rat-serum blocking buffer. Counterstaining was done with 10 µg/mL of

Hoechst 33342 (Life Technologies Ref.# H3570) for 7 min, at room temperature. Sections were

mounted with ProLong™ Gold Antifade Mountant (ThermoFisher Cat. # P10144).

Antibodies used for immunofluorescence:

Alexa Fluor® 647-conjugated anti-mouse CD4 (clone RM4-5) and anti-mouse CD8a (clone 53-

6.7), (BioLegend, USA, Cat.#: 100530 and 100727, respectively) were used for

immunofluorescence at a final concentration of 2.5 µg/mL and eFluor 660 anti-mouse FOXP3

(clone FJK-16s) eBioscience™ (ThermoFisher Scientific, Cat.# 50-5773-80) was used at 1

µg/mL. eFluor 660 rat IgG2a kappa Isotype Control (eBR2a) (eBioscience™, ThermoFisher

Sci., Cat. # 50-4321-80) and Alexa Fluor® 647 Rat IgG2a, κ Isotype Ctrl (BioLegend, USA, Cat.

#: 400526) were used as controls for staining specificity. Anti-Granzyme B (clone D2H2F)

Rabbit mAb (Cell Signaling Cat. # 17215S) and rabbit anti-Ki67 antibody (Abcam, Cat# ab833)

were used at 1:200 dilution. Goat anti-rabbit AF488 (Life Technologies, Cat. # A-11008) and

Texas Red-X (ThermoFisher Cat. # T6391) secondary antibodies were used at 1:500 dilution in

blocking buffer.

Image acquisition and analysis:

Tile-scan imaging of whole tumor sections (20×) was acquired in an Eclipse Ni-E Fluorescence

Microscope (Nikon) with an ORCA-flash4.0 LT plus (Model C11440) Hamamatsu digital

camera and an S Plan Fluor ELWD Ph1 ADM 20x objective (N/A 0.45 and refractive index of

1). Granzyme B imaging was performed in an LSM 800 Confocal Microscope (Carl Zeiss) using

a Plan-Apochromat LCI Plan-Neofluar 25x (0.8 Imm Korr Water DIC) objective and a 63x/1.40

Oil DIC objective (Zeiss). For the images acquired with the Nikon Fluorescence microscope, the

noise component was removed with Denoise AI utility, and the fluorescence background was

subtracted with the NIS-Elements Advanced Research Analysis software (ver. 5.21.02, Nikon).

Separated tile images per channel were converted to binary images, and automated image

analysis was performed using Cell Profiler Software (ver. 4.0.7, Broad Institute MIT) [24], using

a custom pipeline to determine the percent of nuclei (cells) with positive immunostaining for the

marker over the total number of cells normalized to the area of the field of view. At least two

serial cryosections per tumor were used for quantification.

Statistical Analysis

All the statistical analyses were performed using GraphPad Prism® ver. 8 (San Diego, CA,

USA). Unless otherwise indicated, the results are expressed as the mean ± standard deviation

(SD), and the biological (N) and technical replicates (n) for a given experiment are indicated in

each figure. Animal numbers were estimated by *a priori* power analysis (G*Power ver.3.1 [25])

based on a pilot study of spontaneous tumor incidence between BPC-*Panx1*$^{+/-}$ (heterozygous)

and BPC-*Panx1*$^{-/-}$ mice with an effect size of 1.0 and a power of 0.8. For multiple comparisons

involving the genotype and sex of mice, two-way ANOVA was performed, followed by a Tukey-

Kramer *post hoc* test for unequal sample sizes. For real-time qPCR, $\log_2$ (Relative Normalized

expression) was used as data for statistical comparisons. One-way ANOVA followed by Sidak's

post hoc test was done to analyze relevant mean comparisons in the mRNA expression between

tissues. Outliers were excluded from analysis and were detected using the ROUT method with a

False Discovery Rate of less than 1%. Data normality was verified with the D'Agostino-Pearson

test. Immunofluorescence (IF) quantification data were obtained from individual tumors of four

different mice (at least N=3) per genotype and analyzed with a two-tailed non-parametric Mann-

Whitney test. Statistical significance was considered when $p < 0.05$.

RESULTS

Germline *Panx1*-deletion does not reduce melanoma progression of Braf(V600E)/Pten(del)

mice

To explore the *in vivo* effects of the *Panx1* global deletion on melanoma progression, we crossed

BRaf$^{CA/+}$, *Pten*$^{LoxP/Loxp}$, *Tyr::CreER*^{T2} (BPC) mice with the global *Panx1* knockout (*Panx1*$^{-/-}$,

*Panx1*KO) mouse line, reported on [22, 15]. In the BPC model, a 4-hydroxytamoxifen (4-HT)-

inducible Cre recombinase is used for the melanocyte-specific expression of mutant BRaf^{V600E}—

a constitutively active protein serine kinase—and inactivation of the Pten phosphatase to produce

primary cutaneous malignant melanomas (11). However, as in (23), we observed that

spontaneous melanomas occurred in approximately 70% of the BPC colony in the absence of

tamoxifen exposure. Thus, we started the 4-HT induction experiments in mice between 3–4

weeks of age that looked healthy and had no spontaneous melanoma formation.

Our results showed that regardless of the *Panx1* genotype, melanocytic pigmented lesions arose

from the skin and quickly grew to form cutaneous melanomas as early as three weeks after

topical application of 4-HT. Furthermore, although we applied 4-HT to a small zone in the dorsal

skin of the lower back, we found that, in both animal groups, multiple tumor lesions grew not

only in the application region but also in distant sites (Fig. 1A). As a result, all the visible

melanoma lesions found in the dorsal and ventral/lateral skin of the animals were considered to

determine tumor burden (Fig. 1B) and endpoint conditions. Overall, we observed that BPC-

Panx1$^{-/-}$ mice had similar tumor incidence (Fig. 1B), survival (Fig.1C), and mouse weight among

sexes (Fig.1D) compared to *Panx1*-wildtype counterparts. However, although the total combined

tumor growth over time (Fig.1E) was similar between genotypes, analysis of estimated tumor

growth parameters showed sex differences of the tumor growth in BPC-*Panx1*$^{+/+}$ mice (Fig.1F

and G). BPC-*Panx1*-wildtype male mice had a slightly lower (though not significant) tumor

growth rate (Fig.1G, two-way ANOVA, Sex factor: $F_{1,23} = 4.64$, $p<.05$) with a significant

increase ($p = 0.188$) in tumor doubling time (Fig.1F, two-way ANOVA, Sex factor: $F_{1,23} = 7.86$,

$p<.05$) compared to female counterparts. However, this sex-driven difference was not significant

in BPC-*Panx1*-deficient mice and overall tumor growth did not differ between genotypes. These

results indicated that *Panx1* deletion in this mouse model did not reduce BRAF^{V600E}/Pten-driven

melanoma development.

In general, most melanoma lesions were presented with a combination of melanotic (black) and

amelanotic (white) regions (Fig. 2A). However, a few exceptions (less than 30% of animals per

genotype) also had completely amelanotic tumors primarily distant from the tamoxifen

application site. A classification of the tumors with the presence or complete absence of black

pigmentation rendered no significant differences in their number among genotypes (Fig. 2A). All

mice, regardless of the genotype, developed fast-growing tumors that, in most cases, were

ulcerated by the time of endpoint (Fig. 2B). Since our previous reports indicated that PANX1 is

highly expressed in human melanoma tumors compared to normal skin, we performed real-time-

qPCR in matched samples of non-tamoxifen-treated (adjacent) skin and tamoxifen-induced

tumors. *Panx1* mRNA expression was found to be 3.2-fold increased (Fig. 2C) in tumors of

BPC-*Panx1*$^{+/+}$ mice compared to skin, while undetectable in any tissue sample from BPC-

Panx1$^{-/-}$ mice, as expected. Moreover, mRNA expression of the *S100a1* melanoma marker [26,

27] had a 5.1-fold increase (p<0.05) in tumors of BPC-*Panx1*$^{-/-}$ compared to the skin (Fig. 2D),

indicating that the development of primary tumors occurred independently of *Panx1* expression.

Braf(V600E)/Pten-driven melanoma metastasis to inguinal lymph nodes is not restricted by

***Panx1*-deletion**

Melanoma tumor cells are known for passing through the lymphatic system and entering the

bloodstream to metastasize to other organs [28]. We observed localized and distant clusters of

small black pigmented spots spread in the underside of dorsal skin that was not treated with

tamoxifen, suggesting the presence of melanoma microlesions or that in-transit metastasis may

have occurred (Fig. 3A). Regional black melanoma metastases were found in both right and left

inguinal lymph nodes in both *Panx1* genotypes (Fig. 3B, C, and D) but this was not noticeable in

major organs (e.g., lungs, liver, or brain).

Spleens are significantly enlarged in females of tumor-bearing BPC-*Panx1*$^{-/-}$ mice

Upon measuring the length and weight of spleens (Fig. 4A, B, C), a statistically significant sex-

dependent difference was found in BPC-*Panx1*$^{-/-}$ mice (two-way ANOVA, $F_{1, 20} = 10.91$,

p<0.01). Females had an increased spleen weight index (normalized to body weight) compared

to males of the BPC-*Panx1*$^{-/-}$ cohort but this was not observed in BPC-*Panx1*$^{+/+}$ mice or among

genotypes. A positive correlation between spleen weight and length measurements (Fig. 4D)

(Pearson's coefficients r ≥ 0.7, *p*=0.0247 and *p*=0.0087) in both BPC-*Panx1*$^{+/+}$ and BPC-*Panx1*$^{-/-}$

mice, respectively, confirmed a similar trend in both parameters. However, quantification of the

proliferation marker ki67 by immunostaining (Fig. 4E-F) showed significantly fewer

proliferative splenic cells in tumor-bearing BPC-*Panx1*$^{-/-}$ mice than in the BPC-*Panx1*$^{+/+}$ cohort.

CD8 mRNA is significantly increased in *Panx1*-deficient skin and melanoma tumors

Due to PANX1's role in the inflammatory response [12-17], we sought to examine changes in

the infiltration of immune cells within the TME. We performed a real-time qPCR analysis of

gene expression for markers of immune cell populations known to modulate the immune

response against melanoma. We included primers specific to protein tyrosine phosphatase,

receptor type, C (*Ptprc*) (also known as cluster of differentiation (CD) 45) for leukocytes; CD3

antigen, epsilon polypeptide (*CD3ε*) for T-lymphocytes; CD4 antigen (*CD4*) for T-helper

lymphocytes; CD8 antigen, beta polypeptide (*CD8b*) for cytotoxic T cells; forkhead box P3

(*Foxp3*) for regulatory T cells (T-reg); natural cytotoxicity triggering receptor-1 or NCTR1

(*Ncr1*) of Natural killer (NK) cells and the cell surface glycoprotein F4/80 (*Agre1*) for

monocytes/macrophages. The "immune" markers expression was evaluated in melanoma tumors

and matched samples of non-tamoxifen-treated dorsal skin devoid of any visible melanoma

lesions (referred to as "adjacent skin"). These adjacent skin samples were used to account for

basal immune infiltration levels or skin-resident immune cells [29, 30] at the time of sample

collection.

When analyzing tumors compared with their matched skin samples, we observed a significant

($F_{3, 20}$ = 6.24, $p<0.01$) 4.3 fold-increase expression of CD45 only within the BPC-*Panx1*$^{+/+}$ mice

group (Fig. 5A). The F4/80 mRNA expression ($F_{3, 18}$ = 19.38, $p<0.0001$) was 14.0 ($p<0.0001$)

and 7.7-fold ($p<0.01$) higher in tumors than the skin in BPC-*Panx1*$^{+/+}$ and BPC-*Panx1*$^{-/-}$ mice,

respectively (Fig. 5B). On the other hand, transcripts of NCTR1 ($F_{3, 11}$ = 8.24, $p<0.01$) (Fig. 5C)

were 9.3-fold upregulated ($p<0.01$) in BPC-*Panx1*$^{-/-}$ tumors versus adjacent skin. *CD3ε* (Fig. 5D)

showed a similar expression regardless of the sample type or *Panx1* genotype. While *CD4*

mRNA expression ($F_{3, 11}$ = 27.46, $p<0.0001$) was significantly augmented in the tumors

compared to the skin (Fig. 5E) with a 10.5 ($p<0.01$) and 18.9-fold ($p<0.001$) increase in the

BPC-*Panx1*$^{-/-}$ and BPC-*Panx1*$^{+/+}$, respectively; and 5-fold upregulated ($p<0.05$) in the skin of

BPC-*Panx1*$^{-/-}$ vs. that of BPC-*Panx1*$^{+/+}$. Moreover, *Foxp3* ($F_{3, 16}$ = 9.80, $p<0.001$) was

upregulated (5.9-fold, $p<0.05$, and 15.2-fold, $p<0.01$) in tumors of BPC-*Panx1*$^{+/+}$ and BPC-

Panx1$^{-/-}$ mice, respectively (Fig 5G). Notably, *CD8* ($F_{3, 16}$ = 15.39, $p<0.0001$) was significantly

increased ($p<0.01$) in the skin and tumors (9- and 9.5-fold, respectively) of BPC-*Panx1*$^{-/-}$

compared to BPC-*Panx1*$^{+/+}$ animals (Fig. 5F). Taken together, these results indicated that *Panx1*

deficiency did not impair the overall recruitment of immune cells to the primary tumors in this

melanoma model but had an effect on increasing the transcript expression of CD8 in skin and

tumors of BPC-mice.

Given the similarity in tumor burden among genotypes but increased CD8 mRNA in the BPC-

Panx1[-/-] tumors, we investigated if there were differences in the activation/exhaustion phenotype

of TILs and immunosuppression compared to those of the BPC-*Panx1*[+/+] cohort. We observed a

trend of lower transcript expression (not significant) of *CD69* (early T cell activation marker,

[31]) in BPC-*Panx1*[-/-] tumors (Fig. 6A). Moreover, no differences in expression of the regulatory

immune checkpoint receptor lymphocyte activation gene-3 (*LAG-3*) [32] or programmed cell

death protein-ligand 1(PD-L1) [33] were found between genotypes (Fig.6B, C).

Finally, due to the association of PANX1 channel function and the pro-inflammatory and tumor

promoter cytokine: interleukin-1β (IL-1β), we investigated whether the *IL-1β* expression was

affected within the TME of BPC *Panx1*-deficient mice. IL-1β mRNA levels (Fig. 6D) were

found to be highly variable in the BPC-*Panx1*[-/-] tumors but were not different compared to

tumors of the BPC-*Panx1*[+/+] mice.

Panx1-deletion increases tumor infiltration of CD4[+], CD8[+] T lymphocytes, and Granzyme-

B producing cells

To confirm the differences in tumor T cell infiltration between *Panx1* genotypes we performed

IF staining on PLP-fixed tumor cryosections for CD4, CD8, and Foxp3 lymphocytic markers

(Fig. 7A). Overall, CD4[+] and CD8[+] T cells were seen in small but sparsely distributed clusters

closer to the skin epidermis of the primary melanoma tumors, and Foxp3[+] T cells showed a more

homogeneous distribution within the tumor core irrespective of genotype (Fig. 7A). Overall, the

trends in immune cell infiltration of these markers agreed with our real-time qPCR results, and in

tumors, BPC-*Panx1*[-/-] mice had 4- and 2.4-fold increased staining for positive CD4[+] and CD8[+] T

cells, respectively (Fig.7B). In contrast, the percent of Foxp3[+] T cells in tumors was similar

between genotypes.

We also assessed the IF staining of Granzyme B (GzmB) to determine whether differences in

cytotoxicity of T- or NK cells could be detected within the TME. As shown in Fig.7 E-F, we

observed a significant increase (8.4-fold, $p<0.0001$) in the percentage of GzmB[+] cells in tumors

of BPC-*Panx1*[-/-] compared to BPC-*Panx1*[+/+] mice. Notably, GzmB was mostly localized in

intracellular granules. These results correlate with the higher infiltration of CD8$^+$ T cells and,

likely, NK cells since GzmB can be abundant as cytosolic granules in both cytolytic immune cell

types. Collectively, these results confirmed that the deletion of *Panx1* favored the infiltration of

CD4$^+$ and CD8$^+$ T lymphocytes in the tumors, although the cytotoxic effect seemed to be

insufficient to prevent the tumor progression in BPC-*Panx1*$^{-/-}$ mice.

DISCUSSION

PANX1 channels have attracted considerable attention due to their multiple roles in

inflammation, cell death, and cancer [13, 34]. We have demonstrated that PANX1 influences the

regulation of effector molecules that control melanocyte differentiation (e.g., MITF and β-

catenin) and the metabolism of melanoma cells [10, 9, 11]. Furthermore, we have shown

promising results after silencing *Panx1* expression and reducing the growth and tumorigenicity

of melanoma cell lines *in vitro*. In this work, using the immunocompetent BPC mouse melanoma

model, we did not observe a reduction in Braf(V600E)/Pten(del)-driven melanomas or improve

mice survival upon germline *Panx1* deletion. Of note, our previous *in vitro* and *ex vivo* work

targeting PANX1 in melanoma employed murine B16-F0, -F1, -F10 and human BRAF(V600E)-

mutant melanoma cell lines (A375-P and A375-M2) that do not share the same genomic

alterations of the BPC model [10, 35]. The deletion of *Pten* is known to have a synergistic effect

and dramatically accelerates BRAFV600E-induced melanoma tumors in BPC mice [21, 22, 36].

Therefore, we hypothesize that oncogenic mutations intrinsic to this mouse BPC model may

have overridden our anticipated anti-tumoral effect of *Panx1*-deletion. Since this is a highly

penetrant mouse model, a syngeneic melanoma model may be more suitable to study the effect

of *Panx1*-deletion in melanoma metastasis [37]. Besides, the germline deletion of *Panx1* does

not necessarily imply the same effect previously observed with the use of PANX1 channel

blockers (e.g., with carbenoxolone, probenecid, or spironolactone) due to possible compensation

with other proteins (e.g., connexins or other paralogs) yet to be tested using this or another *in*

vivo melanoma model. We also observed the formation of micro-metastasis to draining inguinal

lymph nodes (a phenotype reported elsewhere [21, 22, 38, 23]) in *Panx1*-deficient mice.

Therefore, we concluded that *Panx1*-deletion does not control the early spread of melanoma.

Furthermore, due to the ubiquitous expression of *Panx1*, its global deletion should have affected

many cell types, including melanoma and immune cells. Therefore, we reason that this scenario

may resemble the effect of an irreversible depletion of PANX1 function in which adaptive or

alternative molecular mechanisms are in play that helped to sustain tumor growth.

Interestingly, we noticed a sex dimorphism in melanoma tumor progression where BPC-$Panx1^{+/+}$

female mice seemed to have a more aggressive disease (Fig.1E, G) than males of the same

cohort. Other researchers have noted this outcome in a C57B/6J background, but the origin of

this difference has not been clarified yet [39]. It has been speculated that this may be due to the

presence of alternative steroid receptors in melanoma cells or technical limitations with

tamoxifen (an estrogen receptor modulator) differentially influencing the mouse sexes in this

melanoma model [39]. This sex-dependency was lost in BPC-$Panx1^{-/-}$ mice, suggesting that

$Panx1$-deletion may influence such a sex-driven phenotype. It is known that sexual dimorphism

exists in the response of $Panx1$-deficient mice in ischemia [40] and epilepsy [41], and research

on this is still lacking in the melanoma and immune cell recruitment contexts. A recent study has

shown that tumor progression and anti-tumoral T cell response are sex-dependent in a syngeneic

B16-F10/BL6 mouse model, however, in contrast to our observation, such study showed that

females had less tumor growth rate and higher $CD4^+$ and $CD8^+$ T cell infiltrates compared to

males [42]. Due to the lack of sample availability at the time of the experiment, we did not

quantify differences in T cell subtypes between samples from different sexes. Therefore, it is

unclear the nature of the loss of such sex disparity and further investigation may be required to

clarify this finding.

To date, there are no reports of major abnormalities or immune-related diseases in the Panx1-

deficient mice (reviewed at [13]). Here, we report for the first time that tumor-bearing BPC-

$Panx1^{-/-}$ female mice have significantly augmented spleen size compared to their male

counterparts. Although the tumor may have caused this condition, we found a lower cell

proliferation (ki67 staining) in BPC-$Panx1^{-/-}$ spleens than in the wildtype cohort with no evident

signs of melanoma metastasis in the spleens of any of the mice. Therefore, this morphological

difference in spleen size does not seem to be caused by an increased $in situ$ proliferation of cells

(resident or in transit) in the spleens of tumor-bearing BPC mice. Due to the short timeframe for

tumor growth (up to 2 months) and the sexual dimorphism found only in BPC-$Panx1^{-/-}$ spleens,

we speculate that developmental differences may account for this phenotypic characteristic of

Panx1⁻ᐟ⁻ mice. Further research is warranted to determine whether global *Panx1*-deletion exerts a

sex-specific systemic immune dysregulation and whether this is reflected in the spleen size.

Tumor-infiltrating lymphocytes (TILs) have already been described in this BPC model [23].

However, others have highlighted that this model does not appear to induce a marked immune

response due to the lack of additional tumoral somatic mutations/neoepitopes [37]. Notably, our

results showed increased TILs in BPC-*Panx1⁻ᐟ⁻* tumors. Interestingly, myeloid-derived

suppressor cells, along with high PD-1 expression on the T cells, have been attributed to causing

a highly immunosuppressive environment in this melanoma model [23]. Although we had a

limited analysis of generic markers of all the sources of possible immunosuppression, we found

no difference in the abundance of tumor-infiltrating T-reg cells between both mice cohorts.

Therefore, it is unclear if subsets of $CD4^+$ T cells (Th1 or Th17), macrophages (M1 or M2), or

myeloid-derived suppressor cells infiltrated the tumors and may have contributed to immune

evasion and melanoma progression. Despite the significant increase of $CD8^+$ T cells in *Panx1*-

deficient melanoma tumors, it is interesting that no changes in markers of TIL activation

(CD69)[43, 44], exhaustion (LAG-3), or tumor immune evasion (PD-L1) were found. These

findings implied that deletion of *Panx1* did not influence major activation/exhaustion phenotypes

of tumor-infiltrating T cells in this BPC model. In addition, our observations of a higher

incidence of $GzmB^+$ cells in the BPC-*Panx1⁻ᐟ⁻* tumors support a notion of an increased immune

cell activation phenotype in the absence of PANX1 that could be exploited to counterbalance the

immunosuppressive or evasive mechanisms of melanoma tumors.

Another important aspect is that PANX1 has been proposed as a potential target to regulate

antitumor immunity in melanoma due to the PANX1-mediated release of proinflammatory

cytokine IL-1β (reviewed in [45]). In this case, it is known that IL-1β in melanoma elicits

inflammation and the expansion of highly immunosuppressive myeloid-derived suppressor cells

[46]. In this study, we found no significant differences in IL-1β mRNA expression among

cohorts suggesting that this immune evasion mechanism may be still in place and contributed to

tumor progression despite the increased TIL in Panx1⁻ᐟ⁻ tumors. Using this same *Panx1⁻ᐟ⁻* mouse

strain, another group showed that marrow-derived macrophages have no difference in the

secretion of IL-1β or inflammasome activation [15]. This underscores the need for research into

the crosstalk between immune and tumor cells under cell-specific PANX1 suppression/inhibition

to better understand the implications of PANX1 targeting in the context of melanoma and immunotherapy.

This transgenic melanoma mouse model was originally developed to harness two significant oncogenic alterations (BRAF^{V600E} and PTEN (loss)), but it is not representative of the larger patient population since only 5.0 % of them had co-occurrence of both mutations (Supplemental Fig. S1A). Furthermore, since *PANX1* expression seems independent of genomic alterations of melanoma (Supplemental Fig. S1B) and is present at all stages of the disease [10], a larger patient population may benefit from PANX1 targeted therapy. Interestingly, *Pten*-null cancers seem resistant to immune checkpoint inhibitory antibodies, where PTEN loss in tumor cells increases the expression of immunosuppressive cytokines inhibiting trafficking and T cell-mediated killing of tumors [47] [48]. Therefore, our results point towards the direction of targeting PANX1 as an alternative to overcome T cell exclusion in immunosuppressive TMEs.

In conclusion, our study demonstrates that a *Panx1* deletion cannot overcome the aggressive tumorigenic effect of Braf(V600E)/Pten(del) driver mutations, but it increases the tumor T cell infiltration with a preference in CD8$^+$ cytotoxic T cells. This could be advantageous since a considerable therapeutic benefit is achieved in patients with a pre-existing CD8$^+$ T cell infiltration within the melanoma microenvironment [49]. Future preclinical studies should include the combination strategies in which PANX1 inhibition could be considered as means to stimulate cytotoxic T-cell tumor infiltration to improve the efficacy of immunotherapies.

Acknowledgments

The authors thank Dr. Saman Maleki-Vareki and Dr. Frank Beier for their input on the article. This study was supported by a Canadian Institutes of Health Research CIHR Project grant to Silvia Penuela and Lina Dagnino (FRN 153112)

Authors' Contributions

S. Penuela: Supervision, conceptualization, data analysis, funding acquisition, and writing; **RE Sanchez-Pupo:** Performed most experimental procedures, data analysis, visualization, methodology, wrote and edited the original draft; **G. Finch:** Data curation, spleen measurements, qPCR experiments, and manuscript editing; **D. Johnston:** Technical assistance, resources management, tumor monitoring, sample collection, colony breeding, and dissections; **R. Abdo:** Dissections and sample collection; **S. Kerfoot, H. Craig:** Collaboration, provided IF protocols,

input for experiments and writing; **L. Dagnino:** Collaboration with grant funding and input on

the writing and data analysis.

The authors declare no conflicts of interest of any kind with the current manuscript.

REFERENCES

1.   Tas F (2012) Metastatic behavior in melanoma: timing, pattern, survival, and influencing factors.
J Oncol 2012, 647684, doi: 10.1155/2012/647684.
2.   Dempke WCM, Fenchel K, Uciechowski P & Dale SP (2017) Second- and third-generation drugs
for immuno-oncology treatment-The more the better? Eur J Cancer 74, 55-72, doi:
10.1016/j.ejca.2017.01.001.
3.   Havel JJ, Chowell D & Chan TA (2019) The evolving landscape of biomarkers for checkpoint
inhibitor immunotherapy. Nat Rev Cancer 19, 133-150, doi: 10.1038/s41568-019-0116-x.
4.   Wolchok JD, Chiarion-Sileni V, Gonzalez R, Rutkowski P, Grob JJ, Cowey CL, Lao CD,
Wagstaff J, Schadendorf D, Ferrucci PF, Smylie M, Dummer R, Hill A, Hogg D, Haanen J, Carlino MS,
Bechter O, Maio M, Marquez-Rodas I, Guidoboni M, McArthur G, Lebbe C, Ascierto PA, Long GV,
Cebon J, Sosman J, Postow MA, Callahan MK, Walker D, Rollin L, Bhore R, Hodi FS & Larkin J (2017)
Overall Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. N Engl J Med 377,
1345-1356, doi: 10.1056/NEJMoa1709684.
5.   Jerby-Arnon L, Shah P, Cuoco MS, Rodman C, Su MJ, Melms JC, Leeson R, Kanodia A, Mei S,
Lin JR, Wang S, Rabasha B, Liu D, Zhang G, Margolais C, Ashenberg O, Ott PA, Buchbinder EI, Haq R,
Hodi FS, Boland GM, Sullivan RJ, Frederick DT, Miao B, Moll T, Flaherty KT, Herlyn M, Jenkins RW,
Thummalapalli R, Kowalczyk MS, Canadas I, Schilling B, Cartwright ANR, Luoma AM, Malu S, Hwu
P, Bernatchez C, Forget MA, Barbie DA, Shalek AK, Tirosh I, Sorger PK, Wucherpfennig K, Van Allen
EM, Schadendorf D, Johnson BE, Rotem A, Rozenblatt-Rosen O, Garraway LA, Yoon CH, Izar B &
Regev A (2018) A Cancer Cell Program Promotes T Cell Exclusion and Resistance to Checkpoint
Blockade. Cell 175, 984-997 e924, doi: 10.1016/j.cell.2018.09.006.
6.   Cabrita R, Mitra S, Sanna A, Ekedahl H, Lovgren K, Olsson H, Ingvar C, Isaksson K, Lauss M,
Carneiro A & Jonsson G (2020) The Role of PTEN Loss in Immune Escape, Melanoma Prognosis and
Therapy Response. Cancers (Basel) 12, doi: 10.3390/cancers12030742.
7.   Trujillo JA, Luke JJ, Zha Y, Segal JP, Ritterhouse LL, Spranger S, Matijevich K & Gajewski TF
(2019) Secondary resistance to immunotherapy associated with beta-catenin pathway activation or PTEN
loss in metastatic melanoma. J Immunother Cancer 7, 295, doi: 10.1186/s40425-019-0780-0.
8.   Medina CB, Mehrotra P, Arandjelovic S, Perry JSA, Guo Y, Morioka S, Barron B, Walk SF,
Ghesquiere B, Krupnick AS, Lorenz U & Ravichandran KS (2020) Metabolites released from apoptotic
cells act as tissue messengers. Nature 580, 130-135, doi: 10.1038/s41586-020-2121-3.
9.   Penuela S, Gyenis L, Ablack A, Churko JM, Berger AC, Litchfield DW, Lewis JD & Laird DW
(2012) Loss of pannexin 1 attenuates melanoma progression by reversion to a melanocytic phenotype. J
Biol Chem 287, 29184-29193, doi: 10.1074/jbc.M112.377176.
10.   Freeman TJ, Sayedyahossein S, Johnston D, Sanchez-Pupo RE, O'Donnell B, Huang K, Lakhani
Z, Nouri-Nejad D, Barr KJ, Harland L, Latosinsky S, Grant A, Dagnino L & Penuela S (2019) Inhibition
of Pannexin 1 Reduces the Tumorigenic Properties of Human Melanoma Cells. Cancers (Basel) 11, doi:
10.3390/cancers11010102.
11.   Sayedyahossein S, Huang K, Li Z, Zhang C, Kozlov AM, Johnston D, Nouri-Nejad D, Dagnino
L, Betts DH, Sacks DB & Penuela S (2021) Pannexin 1 binds beta-catenin to modulate melanoma cell
growth and metabolism. J Biol Chem, 100478, doi: 10.1016/j.jbc.2021.100478.
12.   Chekeni FB, Elliott MR, Sandilos JK, Walk SF, Kinchen JM, Lazarowski ER, Armstrong AJ,
Penuela S, Laird DW, Salvesen GS, Isakson BE, Bayliss DA & Ravichandran KS (2010) Pannexin 1

channels mediate 'find-me' signal release and membrane permeability during apoptosis. Nature 467, 863-867, doi: 10.1038/nature09413.

13.   Crespo Yanguas S, Willebrords J, Johnstone SR, Maes M, Decrock E, De Bock M, Leybaert L, Cogliati B & Vinken M (2017) Pannexin1 as mediator of inflammation and cell death. Biochim Biophys Acta Mol Cell Res 1864, 51-61, doi: 10.1016/j.bbamcr.2016.10.006.

14.   Douanne T, Andre-Gregoire G, Trillet K, Thys A, Papin A, Feyeux M, Hulin P, Chiron D, Gavard J & Bidere N (2019) Pannexin-1 limits the production of proinflammatory cytokines during necroptosis. EMBO Rep 20, e47840, doi: 10.15252/embr.201947840.

15.   Qu Y, Misaghi S, Newton K, Gilmour LL, Louie S, Cupp JE, Dubyak GR, Hackos D & Dixit VM (2011) Pannexin-1 Is Required for ATP Release during Apoptosis but Not for Inflammasome Activation. The Journal of Immunology 186, 6553-6561, doi: 10.4049/jimmunol.1100478.

16.   Woehrle T, Yip L, Elkhal A, Sumi Y, Chen Y, Yao Y, Insel PA & Junger WG (2010) Pannexin-1 hemichannel-mediated ATP release together with P2X1 and P2X4 receptors regulate T-cell activation at the immune synapse. Blood 116, 3475-3484, doi: 10.1182/blood-2010-04-277707.

17.   Woehrle T, Yip L, Manohar M, Sumi Y, Yao Y, Chen Y & Junger WG (2010) Hypertonic stress regulates T cell function via pannexin-1 hemichannels and P2X receptors. J Leukoc Biol 88, 1181-1189, doi: 10.1189/jlb.0410211.

18.   Chen W, Zhu S, Wang Y, Li J, Qiang X, Zhao X, Yang H, D'Angelo J, Becker L, Wang P, Tracey KJ & Wang H (2019) Enhanced Macrophage Pannexin 1 Expression and Hemichannel Activation Exacerbates Lethal Experimental Sepsis. Sci Rep 9, 160, doi: 10.1038/s41598-018-37232-z.

19.   Velasquez S, Malik S, Lutz SE, Scemes E & Eugenin EA (2016) Pannexin1 Channels Are Required for Chemokine-Mediated Migration of CD4+ T Lymphocytes: Role in Inflammation and Experimental Autoimmune Encephalomyelitis. J Immunol 196, 4338-4347, doi: 10.4049/jimmunol.1502440.

20.   Penuela S, Kelly JJ, Churko JM, Barr KJ, Berger AC & Laird DW (2014) Panx1 regulates cellular properties of keratinocytes and dermal fibroblasts in skin development and wound healing. J Invest Dermatol 134, 2026-2035, doi: 10.1038/jid.2014.86.

21.   Damsky WE, Curley DP, Santhanakrishnan M, Rosenbaum LE, Platt JT, Gould Rothberg BE, Taketo MM, Dankort D, Rimm DL, McMahon M & Bosenberg M (2011) beta-catenin signaling controls metastasis in Braf-activated Pten-deficient melanomas. Cancer Cell 20, 741-754, doi: 10.1016/j.ccr.2011.10.030.

22.   Dankort D, Curley DP, Cartlidge RA, Nelson B, Karnezis AN, Damsky WE, Jr., You MJ, DePinho RA, McMahon M & Bosenberg M (2009) Braf(V600E) cooperates with Pten loss to induce metastatic melanoma. Nat Genet 41, 544-552, doi: 10.1038/ng.356.

23.   Hooijkaas AI, Gadiot J, van der Valk M, Mooi WJ & Blank CU (2012) Targeting BRAFV600E in an inducible murine model of melanoma. Am J Pathol 181, 785-794, doi: 10.1016/j.ajpath.2012.06.002.

24.   Lamprecht MR, Sabatini DM & Carpenter AE (2007) CellProfiler: free, versatile software for automated biological image analysis. Biotechniques 42, 71-75, doi: 10.2144/000112257.

25.   Faul F, Erdfelder E, Lang AG & Buchner A (2007) G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods 39, 175-191, doi: 10.3758/bf03193146.

26.   Sviatoha V, Tani E, Kleina R, Sperga M & Skoog L (2010) Immunohistochemical analysis of the S100A1, S100B, CD44 and Bcl-2 antigens and the rate of cell proliferation assessed by Ki-67 antibody in benign and malignant melanocytic tumours. Melanoma Res 20, 118-125, doi: 10.1097/CMR.0b013e3283350554.

27.   Xiong TF, Pan FQ & Li D (2019) Expression and clinical significance of S100 family genes in patients with melanoma. Melanoma Res 29, 23-29, doi: 10.1097/CMR.0000000000000512.

28.   Ubellacker JM, Tasdogan A, Ramesh V, Shen B, Mitchell EC, Martin-Sandoval MS, Gu Z, McCormick ML, Durham AB, Spitz DR, Zhao Z, Mathews TP & Morrison SJ (2020) Lymph protects metastasizing melanoma cells from ferroptosis. Nature 585, 113-118, doi: 10.1038/s41586-020-2623-z.

29.   Nguyen AV & Soulika AM (2019) The Dynamics of the Skin's Immune System. Int J Mol Sci 20,
doi: 10.3390/ijms20081811.
30.   Richmond JM & Harris JE (2014) Immunology and skin in health and disease. Cold Spring Harb
Perspect Med 4, a015339, doi: 10.1101/cshperspect.a015339.
31.   Park SL, Gebhardt T & Mackay LK (2019) Tissue-Resident Memory T Cells in Cancer
Immunosurveillance. Trends Immunol 40, 735-747, doi: 10.1016/j.it.2019.06.002.
32.   Zhang Z, Liu S, Zhang B, Qiao L, Zhang Y & Zhang Y (2020) T Cell Dysfunction and
Exhaustion in Cancer. Front Cell Dev Biol 8, 17, doi: 10.3389/fcell.2020.00017.
33.   Freeman GJ, Long AJ, Iwai Y, Bourque K, Chernova T, Nishimura H, Fitz LJ, Malenkovich N,
Okazaki T, Byrne MC, Horton HF, Fouser L, Carter L, Ling V, Bowman MR, Carreno BM, Collins M,
Wood CR & Honjo T (2000) Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family
member leads to negative regulation of lymphocyte activation. J Exp Med 192, 1027-1034, doi:
10.1084/jem.192.7.1027.
34.   Jiang JX & Penuela S (2016) Connexin and pannexin channels in cancer. BMC Cell Biol 17
Suppl 1, 12, doi: 10.1186/s12860-016-0094-8.
35.   Melnikova VO, Bolshakov SV, Walker C & Ananthaswamy HN (2004) Genomic alterations in
spontaneous and carcinogen-induced murine melanoma cell lines. Oncogene 23, 2347-2356, doi:
10.1038/sj.onc.1207405.
36.   Karreth FA, Tay Y, Perna D, Ala U, Tan SM, Rust AG, DeNicola G, Webster KA, Weiss D,
Perez-Mancera PA, Krauthammer M, Halaban R, Provero P, Adams DJ, Tuveson DA & Pandolfi PP
(2011) In vivo identification of tumor- suppressive PTEN ceRNAs in an oncogenic BRAF-induced mouse
model of melanoma. Cell 147, 382-395, doi: 10.1016/j.cell.2011.09.032.
37.   Meeth K, Wang JX, Micevic G, Damsky W & Bosenberg MW (2016) The YUMM lines: a series
of congenic mouse melanoma cell lines with defined genetic alterations. Pigment Cell Melanoma Res 29,
590-597, doi: 10.1111/pcmr.12498.
38.   Hooijkaas A, Gadiot J, Morrow M, Stewart R, Schumacher T & Blank CU (2012) Selective
BRAF inhibition decreases tumor-resident lymphocyte frequencies in a mouse model of human
melanoma. Oncoimmunology 1, 609-617, doi: 10.4161/onci.20226.
39.   Zhai Y, Haresi AJ, Huang L & Lang D (2020) Differences in tumor initiation and progression of
melanoma in the Braf(CA) ;Tyr-CreERT2;Pten(f/f) model between male and female mice. Pigment Cell
Melanoma Res 33, 119-121, doi: 10.1111/pcmr.12821.
40.   Freitas-Andrade M, Bechberger JF, MacVicar BA, Viau V & Naus CC (2017) Pannexin1
knockout and blockade reduces ischemic stroke injury in female, but not in male mice. Oncotarget 8,
36973-36983, doi: 10.18632/oncotarget.16937.
41.   Aquilino MS, Whyte-Fagundes P, Lukewich MK, Zhang L, Bardakjian BL, Zoidl GR & Carlen
PL (2020) Pannexin-1 Deficiency Decreases Epileptic Activity in Mice. Int J Mol Sci 21, doi:
10.3390/ijms21207510.
42.   Dakup PP, Porter KI, Little AA, Zhang H & Gaddameedhi S (2020) Sex differences in the
association between tumor growth and T cell response in a melanoma mouse model. Cancer Immunol
Immunother 69, 2157-2162, doi: 10.1007/s00262-020-02643-3.
43.   Cibrian D & Sanchez-Madrid F (2017) CD69: from activation marker to metabolic gatekeeper.
Eur J Immunol 47, 946-953, doi: 10.1002/eji.201646837.
44.   Edwards J, Wilmott JS, Madore J, Gide TN, Quek C, Tasker A, Ferguson A, Chen J, Hewavisenti
R, Hersey P, Gebhardt T, Weninger W, Britton WJ, Saw RPM, Thompson JF, Menzies AM, Long GV,
Scolyer RA & Palendira U (2018) CD103(+) Tumor-Resident CD8(+) T Cells Are Associated with
Improved Survival in Immunotherapy-Naive Melanoma Patients and Expand Significantly During Anti-
PD-1 Treatment. Clin Cancer Res 24, 3036-3045, doi: 10.1158/1078-0432.CCR-17-2257.
45.   Varela-Vazquez A, Guitian-Caamano A, Carpintero-Fernandez P, Fonseca E, Sayedyahossein S,
Aasen T, Penuela S & Mayan MD (2020) Emerging functions and clinical prospects of connexins and
pannexins in melanoma. Biochim Biophys Acta Rev Cancer 1874, 188380, doi:
10.1016/j.bbcan.2020.188380.

46.   Tengesdal IW, Dinarello A, Powers NE, Burchill MA, Joosten LAB, Marchetti C & Dinarello
CA (2021) Tumor NLRP3-Derived IL-1β Drives the IL-6/STAT3 Axis Resulting in Sustained MDSC-
Mediated Immunosuppression. Frontiers in Immunology 12, Original Research, doi:
10.3389/fimmu.2021.661323.
47.   Peng W, Chen JQ, Liu C, Malu S, Creasy C, Tetzlaff MT, Xu C, McKenzie JA, Zhang C, Liang
X, Williams LJ, Deng W, Chen G, Mbofung R, Lazar AJ, Torres-Cabala CA, Cooper ZA, Chen PL, Tieu
TN, Spranger S, Yu X, Bernatchez C, Forget MA, Haymaker C, Amaria R, McQuade JL, Glitza IC,
Cascone T, Li HS, Kwong LN, Heffernan TP, Hu J, Bassett RL, Jr., Bosenberg MW, Woodman SE,
Overwijk WW, Lizee G, Roszik J, Gajewski TF, Wargo JA, Gershenwald JE, Radvanyi L, Davies MA &
Hwu P (2016) Loss of PTEN Promotes Resistance to T Cell-Mediated Immunotherapy. Cancer Discov 6,
202-216, doi: 10.1158/2159-8290.CD-15-0283.
48.   Lin Z, Huang L, Li SL, Gu J, Cui X & Zhou Y (2021) PTEN loss correlates with T cell exclusion
across human cancers. BMC Cancer 21, 429, doi: 10.1186/s12885-021-08114-x.
49.   Massi D, Rulli E, Cossa M, Valeri B, Rodolfo M, Merelli B, De Logu F, Nassini R, Del Vecchio
M, Di Guardo L, De Penni R, Guida M, Sileni VC, Di Giacomo AM, Tucci M, Occelli M, Portelli F,
Vallacchi V, Consoli F, Quaglino P, Queirolo P, Baroni G, Carnevale-Schianca F, Cattaneo L, Minisini
A, Palmieri G, Rivoltini L, Mandala M & Italian Melanoma I (2019) The density and spatial tissue
distribution of CD8(+) and CD163(+) immune cells predict response and outcome in melanoma patients
receiving MAPK inhibitors. J Immunother Cancer 7, 308, doi: 10.1186/s40425-019-0797-4.

Tables

Table 1. List of primers used for real-time qPCR analysis

Target gene	Forward primer (5'-3')	Reverse primer (5'-3')
Panx1	ACAGGCTGCCTTTGTGGATTCA	GGGCAGGTACAGGAGTATG
S100a1	CCATGGAGACCCTCATCAAT	TTCTGGACATCCAGGAAGC
CD3ε	ATGCGGTGGAACACTTTCTGG	GCACGTCAACTCTACACTGGT
CD4	GCGAGAGTTCCCAGAAGAAG	AAACGATCAAACTGCGAAGG
CD8β1	AAGAAGCAATGCCCGTTCC	TGAGGGTGGTAAGGCTGCA
Ptprc (CD45)	ATGGTCCTCTGAATAAAGCCCA	TCAGCACTATTGGTAGGCTCC
Foxp3	TTGCCAAGCTGGAAGACTGC	CAGACGGTGCCACCATGACT
ncr1	TGTTGAGAACAGCAGCCTTG	AGGATTATGCATGCCAGACC
Adgre1(F4/80)	CTTTGGCTATGGGCTTCCAGTC	GCAAGGAGGACAGAGTTTATCGTG
*CD69**	GGGCTGTGTTAATAGTGGTCCTC	CTTGCAGGTAGCAACATGGTGG
*LAG-3**	CTCCATCACGTACAACCTCAAGG	GGAGTCCACTTGGCAATGAGCA
*PD-L1**	TGCGGACTACAAGCGAATCACG	CTCAGCTTCTGGATAACCCTCG
*IL-1β**	TGGACCTTCCAGGATGAGGACA	GTTCATCTCGGAGCCTGTAGTG
gapdh	TGTGTCCGTCGTGGATCTGA	TTGCTGTTGAAGTCGCAGGAG
18s	CGGACAGGATTGACAGATTG	CAAATCGCTCCACCAACTAA

596 ***** qSTAR qPCR mouse primer pairs from OriGene Technologies Inc., Rockville, MD, USA, CD69
597 (Cat.# MP201944); *Lag-3* (Cat.# MP207670); *PD-L1* (Cat.# MP201906); *IL-1β* (Cat.# MP206724)

598 **Figure legends**

Figure 1. Global *Panx1* deletion does not improve survival or reduce tumor progression in the Braf(V600E)/Pten-deficient mouse melanoma model. (**A**) Representative picture of *Tyr::CreER; BRaf*$^{CA/+}$*; Pten*$^{-/-}$ (BPC) mice knockout (-/-) for *Panx1* with multiples tumors at endpoint. Region of the tamoxifen application is indicated in the picture. (**B**) Tumor incidence, recorded as the number of visible cutaneous melanoma lesions at endpoint, was similar between both mouse sexes and genotypes. (**C**) Kaplan-Meier survival curves were similar among BPC-*Panx1* wildtype ($^{+/+}$) or BPC-*Panx1*$^{-/-}$ mice (Log-rank test, p = 0.7803). Dotted lines denote 95% confidence intervals. (**D**) Comparison of mice body weight after tumor induction showed no differences among genotypes within the same sex group. Statistical significance was determined by multiple t-tests per timepoint using the Holm-Sidak method without assuming a consistent *SD*. (**E**) Tumor growth curves showed overlap between genotypes, with a group of mice exhibiting slightly delayed tumor growth. Individual lines depict measurements per mouse. (**F, G**) Tumor growth rate and doubling time were similar among genotypes but significantly different between males and females within the BPC-*Panx1*$^{+/+}$ group. Two-way ANOVA showed no interaction between genotype and sex, but significant *p*-values for the sex factor in tumor doubling time ($F_{1,23} = 7.855$, p<.05) and growth rate ($F_{1,23} = 4.642$, p<.05), respectively. Symbols stand for individual data per mouse (BPC-*Panx1*$^{+/+}$, n=15, and BPC-*Panx1*$^{-/-}$, n=13) except for the graph in **D**, where mean ± *SD* are shown. Due to unequal sample sizes per sex, the Tukey-Kramer test was used as *post hoc* test for pairwise comparisons shown in **F** and **G**. Statistical significance was considered when *p*<0.05.

Figure 2. Pigmentation or ulceration of Braf(V600E)/Pten(del) melanoma lesions are not influenced by the global *Panx1* deletion. (A, B) Representative pictures of pigmented and ulcerated tumors. Fractions (out of the total tumors per mice) of pigmented or ulcerated tumors were similar among the genotype and sex groups. Symbols show individual data per mouse and horizontal lines, and error bars represent mean ± *SD*. Data were analyzed with two-way ANOVA. **(C, D)** mRNA expression of *Panx1* and melanoma marker *s100a1* in tumors compared to matched samples of non-tamoxifen treated skin per mouse. *Gapdh* and *18s* were used as reference genes, and the normalized expression is shown relative to one of the BPC-*Panx1*$^{+/+}$ skin samples. Box plots represent the 95% confidence interval (CI) and the median (inner line), with whiskers representing the maximum and minimum values of the gene expression. Symbols display the individual expression data per mouse. Paired t-test was used to compare the log$_2$ (normalized expression) of *Panx1* and one-way ANOVA followed by a Tukey test used to compare the expression of *S100a1*. At least N=5 paired samples were assayed by triplicate (n=3) for real-time-qPCR analysis. Statistical significance is shown as p<0.05(*).

Figure 3. *Panx1* deletion does not prevent the development of Braf/Pten-driven melanoma micro-lesions or metastasis to inguinal lymph nodes in BPC mice. (A) Representative pictures of melanoma micro-lesions found underside the dorsal and ventral/lateral skin and located distant (inset **a**) and nearby (inset **b**) to the application sites of tamoxifen (e.g., lower back dorsal skin). LN: Inguinal draining lymph nodes. **(B)** Representative pictures of right and left inguinal lymph nodes showing black pigmentation indicative of melanoma metastasis. **(C)** Phase-contrast micrograph of an inguinal draining lymph node cryosection exhibiting melanin deposits and pigmented cells (white arrows). **(D)** Quantification of the fraction (out of two inguinal LN per mice) of draining lymph nodes classified with melanoma metastasis based on

the presence or absence of visible pigmented lesions. Symbols correspond to the individual data

per mouse where mean ± *SD* are shown. Data were analyzed with a two-way ANOVA.

Statistical significance was considered when $p < 0.05$.

Figure 4. Sex-dependent differences in spleen size were observed for BPC-*Panx1*^-/- but not

for BPC-*Panx1*^+/+ mice. (A) Representative pictures of BPC mice spleens. **(B)** Morphological

comparison of the spleen measured by length and **(C)** weight index (spleen weight normalized to

body mass) showed a trend of enlarged spleens in female BPC-*Panx1*^-/- mice. Sex-dependent

differences in BPC-*Panx1*^-/- mice were statistically significant (Two-way ANOVA, $F_{1,\,20} = 10.91$,

$p < 0.01$). A Tukey-Kramer test used as *post hoc* test for pairwise comparisons showed a

significantly ($p < 0.01$) higher spleen weight index in BPC-*Panx1*^-/- females than males, but not in

the BPC-*Panx1*^+/+ cohort. Symbols correspond to data from individual mice where mean ± SD

are shown. **(D)** A positive correlation was found between the spleen weight and length,

confirming the trends of increased spleen size in female BPC-*Panx1*^-/- mice observed by both

measurements. **(E)** Representative immunofluorescence micrographs (20x magnification) of

cross-sectional views of the spleen sections stained with the proliferation marker ki67. Yellow

arrows indicate the location of proliferating cells (green). Nuclei (magenta) were counterstained

with Hoechst 33342. Scale bar = 200 µm **(F)** Quantification of ki67 staining in the spleens

showing significant reduction in the % of proliferative cells in spleens of BPC-*Panx1*^-/- mice;

symbols represent individual measurements of at least 12 non-overlapping fields of view from

randomly selected animals: BPC-*Panx1*^+/+ (N=3; n=1 female and n=2 males) and BPC-*Panx1*^-/-

(N=4 females) mice. Box plots represent the 95% confidence interval (CI) and the mean (inner

line), whiskers represent the maximum and minimum counts per image. Data were analyzed with

two-tailed unpaired t-tests. Statistical significance is shown as $p < 0.05$ (*), $p < 0.01$(**) and

$p < 0.0001$(****).

Figure 5. CD8 T-lymphocyte marker is significantly increased in the non-tamoxifen treated

skin and melanoma tumors of BPC-mice lacking PANX1. (A-G) Real-time qPCR analysis of

immune cell markers expression in tamoxifen-induced tumors and matched non-tamoxifen

treated skin. *Gapdh* and *18s* were used as reference genes, and the normalized expression was

calculated relative to one of the BPC-*Panx1*^+/+ skin samples. Box plots represent the 95%

confidence interval (CI) and the median (inner line), with whiskers showing the maximum and

minimum values of the gene expression. Each sample was assayed by triplicate (n=3). The

symbols represent the expression data of each mouse (at least N=3 per genotype). The $\log_2$ of the

expression per group was analyzed by one-way ANOVA followed by a Sidak *post hoc* test. Only

significant *p* values of relevant comparisons between groups are shown as $p < 0.05$ (*), < 0.01(**),

< 0.001(***).

Figure 6. Global *Panx1*-deletion does not affect the activation phenotype of tumor-

infiltrated T-lymphocytes. Real-time qPCR analysis of **(A)** the early T cell immune activation

marker *CD69*, **(B)** checkpoint receptor lymphocyte activation gene-3 (*LAG-3*), **(C)** immune

checkpoint molecule programmed cell death 1-ligand (*PD-L1*), and **(D)** *IL-1β* in tumor samples.

Gapdh was used as a reference gene, and the normalized expression is relative to one of the

BPC-*Panx1*^+/+ skin samples. Box plots represent the 95% confidence interval (CI) and the

median (inner line), with whiskers showing the maximum and minimum values of the gene

expression. Each sample was assayed by triplicate (n=3), and the symbols represent the

expression data of each mouse (at least N=4 per genotype). A t-test compared the mean of the

$\log_2$ of the relative expression per group; statistical significance was considered when $p<0.05$ (*).

Figure 7. Tumor-infiltrating CD4$^+$, CD8$^+$ T-lymphocytes, and Granzyme B producing cells

are significantly increased in BPC-*Panx1*$^{-/-}$ tumors. (A) Representative micrographs (20x,

insets at 60x magnification) showing immunofluorescence staining of primary melanoma tumor

cryosections with immune cell infiltration markers for CD4+, CD8+' and FoxP3+ T-lymphocytes

(shown in cyan). Nuclei (in magenta) was counterstained with Hoechst 33324. **(B-D)**

Immunofluorescence quantification of CD4, CD8, and FoxP3 positive cells for the whole tumor

section. **(E)** Representative granzyme B (GzmB) immunofluorescence staining of a BPC-*Panx1*$^{-/-}$

tumor (fluorescence channels on top and brightfield channel at the bottom) with the intra-tumoral

distribution of GzmB+ cells and black pigmented regions in the tumor. Nuclei are shown in blue,

and GzmB is shown in green. Yellow arrowheads indicate stained immune cells. Scale bars = 40

µm. **(F)** Quantification of GzmB immunostaining. The percent of nuclei with positive immune

marker staining was normalized per area of the field of view (FOV). At least three mice (N=3)

per genotype were examined with at least 12 non-overlapping FOV per tumor. Bar graphs stand

for the median with 95% CI, and individual measurements are depicted with symbols. Data were

compared using a two-tailed Mann-Whitney test. Statistical significance is shown as

$p<0.001$(***) and $p<0.0001$(****).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7