A Non-Coding Variant in *SLC15A4* Drives Lysosomal Deacidification and Lupus Susceptibility

Guru Prashad Maiti[1][†], Manish Kumar Singh[1][†], HariKrishna Reddy-Rallabandi[1], Mehdi Fazel-Najafabadi[1], Loren L. Looger[2], Swapan K. Nath[1][*]

[1]Arthritis and Clinical Immunology Research Program, Oklahoma Medical Research Foundation, Oklahoma City OK, USA

[2]Howard Hughes Medical Institute, Department of Neurosciences, University of California San Diego, La Jolla CA, USA

† Contributed equally

* Correspondence:
Swapan K. Nath, Ph.D.
Swapan-Nath@omrf.org

Keywords: SLC15A4, Luciferase, ChIP-qPCR, CRISPR, Lupus, Lysosomal acidification

Running Title: SLC15A4 Variant in lupus susceptibility

Abstract

Systemic lupus erythematosus (SLE) is a complex autoimmune disease with a strong genetic
basis. Despite the identification of several single nucleotide polymorphisms (SNPs) near the
SLC15A4 gene that are significantly associated with SLE across multiple populations, specific
causal SNP(s) and molecular mechanisms responsible for disease susceptibility are unknown. To
address this gap, we employed bioinformatics, expression quantitative trait loci (eQTLs), and 3D
chromatin interaction analysis to nominate a likely functional variant, rs35907548, in an active
intronic enhancer of *SLC15A4*. Through luciferase reporter assays followed by chromatin
immunoprecipitation-qPCR, we identified significant allele-specific enhancer effects of
rs35907548 in diverse cell lines. The rs35907548 risk allele T is associated with increased
regulatory activity and target gene expression, as shown by eQTLs and chromosome
conformation capture (3C)-qPCR. The latter revealed long-range chromatin interactions between
the rs35907548 enhancer and the promoters of *SLC15A4, GLTLD1*, and an uncharacterized
lncRNA. The enhancer-promoter interactions and expression effects were validated by
CRISPR/Cas9 knock-out of the locus in HL60 promyeloblast cells. Further, KO cells had
dysregulated endolysosomal pH. This study highlights the significance of a non-coding variant,
rs35907548, which may have a direct impact on the underlying regulatory mechanisms
associated with SLE.

1   Introduction

Systemic lupus erythematosus (SLE) is a chronic autoimmune disease characterized by immune attack on the body's tissues and organs, resulting in profound dyshomeostasis and damage to skin, joints, kidneys, cardiovascular system, and nervous system, among others. The disease is very heterogeneous, with each patient presenting uniquely, thus making diagnosis, treatment, and even basic understanding challenging [1]. Gender and ethnicity significantly influence the incidence and severity of SLE, with females and individuals of African, Hispanic, and Asian ancestry being both more prone to SLE and to severe manifestations like kidney disease and frequent hospitalization. Conversely, European-ancestry SLE patients tend to exhibit more skin manifestations but less renal involvement. Therefore, managing SLE and its comorbidities requires a comprehensive internal medicine approach and an understanding of its diverse underlying pathogenic mechanisms [2, 3, 4, 5].

SLE has a significant genetic component, evidenced by familial aggregation, sibling risk ratio, and twin studies [6]. SLE appears to be a highly polygenic disease, with >100 risk loci implicated to date [7, 8]. These SNPs and associated genes are involved in biological pathways related to tolerance, cell signaling, apoptosis, and other critical immune functions [9, 10]. Molecular pathway analysis has revealed different underpinnings of immune system homeostasis and SLE risk in diverse ethnic populations [9, 11, 12], emphasizing the importance of studying the genetics of complex disease in multiple ethnicities – both to elucidate fundamental biochemical pathways and to develop personalized diagnostics and treatments.

SLC15A4 was discovered as an SLE risk locus in 2009 through GWAS on Chinese individuals. Since then, several SNPs have been associated with risk across Asian, European, Hispanic, and African ancestries [9]. However, despite strong genetic association, the actual functional SNP(s) and underlying biological mechanism(s) contributing to SLE pathogenesis are not understood [13, 14].

SLC15A4, also known as peptide/histidine transporter 1 (PHT1), is a member of the solute carrier family 15 of proton-coupled oligopeptide transporters [13, 14, 15, 16]. It is primarily located on the endolysosomal membrane of immune cells, where it transports histidine and bacterially-derived dipeptides such as the NOD2 ligand muramyl dipeptide (MDP). SLC15A4 is crucial for lysosomal acidification, as it generates a proton gradient through transport of the proton acceptor histidine. Moreover, SLC15A4 can recruit the adapter molecule "TLR adaptor interacting with SLC15A4 on the lysosome" (TASL), which regulates Toll-like receptor (TLR) function and promotes downstream signaling through type I interferon and Interferon Response Factor 5 (IRF5) [17]. Therefore, SLC15A4 plays a crucial role in regulating lysosomal function and innate immunity. SLC15A4 deficiency may promote lysosomal dysfunction and impaired autophagy, both associated with various autoimmune diseases, including SLE [18]. Consequently, these findings suggest that pharmacological intervention to restore or supplement the function of SLC15A4 may be a promising therapeutic approach for treating lupus and other endosomal TLR-dependent diseases [18, 19, 20, 21].

Here, we systematically employ bioinformatics to investigate SNPs in the *SLC15A4* locus, identifying rs35907548 as a likely regulatory variant. We show that this variant indeed dictates activity of a potent enhancer, chromatin interactions, expression of *SLC15A4* and nearby genes,

and ultimately plays a decisive role in maintaining endolysosomal acidification, critical for
proper function of immune cells. Our results localize SLE risk of the highly associated *SLC15A4*
locus and give mechanistic insight into its function.

2 Materials and Methods

2.1 Bioinformatics

We utilized RegulomeDB [22] and Ldlink [23] to prioritize SNP regulatory potential. We added
information on *cis*-regulatory elements, including chromatin accessibility (ATAC-seq and
DNase-I hypersensitivity), canonical histone modifications (H3K27ac, H3K4me1, H3K4me3),
and RNA polymerase II (Pol II) binding, from ENCODE [24] data from immune cell lines.
Transcription factor binding sites were taken from ENCODE.

We used ImmuNexUT (Immune Cell Gene Expression Atlas from the University of Tokyo) [25]
data – built on 28 immune cell types from 10 immune diseases and healthy donors – to assess
cell type-specific expression quantitative trait loci (eQTLs) and target gene expression.

2.2 Cell lines

Dual-luciferase reporter assays were performed in HEK293, lymphoblastoid cell lines (LCLs; B-
cell), Jurkat (T-cell), and HL-60 (promyeoloblast). ChIP-qPCR assays were performed in LCLs
(GM18624, GM18603). Chromatin conformation capture (3C) experiments were in LCLs
(GM18603, GM18624). CRISPR/Cas9 gene editing was in HL-60. Endolysosomal pH
measurements were in wild-type and gene-edited HL-60. All cell lines were purchased from
ATCC (American Type Culture Collection). All cells were tested for mycoplasma by PCR and
used between passages 4-7.

2.3 Dual-luciferase assay

We measured enhancer activity with Dual-Luciferase Reporter Assays [26, 27]. Briefly, we
cloned the ~130bp rs35907548 locus into pGL4.26 (Promega) and co-transfected with pGL4.74
(internal control) in HEK293, LCL, Jurkat, and HL-60 cells. After 24 hours, enhancer activity
was measured using a Synergy H1 spectrophotometer (Biotek). Three experimental replicates
were performed per cell type. Statistical significance was assessed by Student's t-test using
GraphPad PRISM; p-value < 0.05 was considered significant.

2.4 Chromatin immunoprecipitation and quantitative PCR (ChIP-qPCR)

ChIP-qPCR assays were performed with Magnify ChIP assay (Cat No. 492024, Thermo-Fisher,
Waltham, MA, USA), following manufacturer's guidelines. 1.5-2 $\times 10^6$ homozygous rs35907548
risk-"TT" and non-risk-"CC" genotype B-lymphoblastoid LCLs (Coriell) were cross-linked with
1% paraformaldehyde, washed, and sonicated. Immunoprecipitation was performed overnight at
4°C with antibodies against individual histone marks or other DNA-binding proteins pre-
incubated with Dynamag magnetic A+G beads. After reverse crosslinking and elution, real-time
qPCR analysis was performed with SYBR Green and primers flanking the rs35907548 region
using an Applied Biosystems 7900HT qPCR machine. Statistical significance was assessed by

Student's t-test using GraphPad PRISM software, and a p-value < 0.05 was considered
significant. Detailed methods are described in Supplementary Materials and Methods.

2.5   Chromatin conformation capture (3C) with quantitative PCR (3C-qPCR)

We used 3C-qPCR [26, 27] in HEK293, LCL, and Jurkat cells. Briefly, cells were suspended in
complete media with 10% FBS (1×10^6/mL media) at 70-80% confluency and cross-linked with
1% paraformaldehyde at room temperature for 10 minutes. Following quenching with 0.2 M
glycine, cross-linked cells were lysed in buffer containing protease inhibitors. Cross-linked
nuclei were purified, suspended in 0.5% SDS, and incubated at 62°C followed by quenching
with Triton X-100. Perforated nuclei were digested with 400 U *HindIII* and *DpnII* at 37°C
overnight and in-nucleus ligated with T4 DNA ligase at 20°C for 4 hours. A small volume of the
digested mixture was reserved to evaluate digestion efficiency. DNA was purified from ligated
chromatin with proteinase K digestion, phenol-chloroform extraction, and alcohol precipitation.
Purified DNA was quantified and diluted for 3C-PCR. Primers were designed to amplify several
promoter regions based on restriction maps. Primers within the rs35907548 enhancer were used
as common primers for other fragments. Primer sequences are in **Supplementary Table 1**.
Cross-linking frequencies were calculated from PCR band intensity. Data were plotted as relative
interaction frequency versus genomic distance from rs35907548.

2.6   CRISPR/Cas9 gene editing

Short-guide RNA (sgRNA)/Cas9 RNP complexes were introduced into HL-60 cells by Neon
Electroporation System [26, 27]. Genomic DNA was extracted three days post-transfection and
Sanger-sequenced to verify deletion. Indel efficiency was determined using TIDE and ICE.
Pooled edited cells were cultured and harvested for gene expression measurements. sgRNA
sequences are in **Supplementary Table 1**. Three experimental replicates were performed per
cell type. Statistical significance was assessed by Student's t-test using GraphPad PRISM; p-
value of < 0.05 was considered significant.

2.7   Quantitative real-time PCR (qRT-PCR)

Pooled CRISPR/Cas9-edited cells were used for RNA purification using the RNA Mini-prep kit
(Zymo Research). Purity and concentration were measured using a NanoDrop
spectrophotometer. Approximately 700 ng total purified RNA was used to generate cDNA with
the iScript cDNA Synthesis kit (Bio-Rad). cDNA was used for PCR to quantify gene expression
for *SLC15A4, GLTD1,* and lncRNA AC069262.1 using qRT-PCR LightCycler 480 Instrument II
(Roche) using specific primers and iTaq Universal SYBR Green Supermix (Bio-Rad). To
normalize gene expression data, 18S rRNA was used as an internal control. Primer sequences are
in **Supplementary Table 1**.

2.8   Endolysosomal pH

We used the pHrodo® Red AM Intracellular pH Indicator (Thermo-Fisher) following
manufacturer's protocol. Briefly, WT and KO cells (0.2×10^6 cells) were stained with 5 μM
pHrodo® Red AM at 37°C for 30 minutes in 96-well plates. Cells were washed with Live Cell
Imaging Media, and standard buffers containing 10 μM nigericin and 10 μM valinomycin were
added to specific wells for 5 minutes to clamp intracellular pH values at 4.5, 5.5, 6.5, and 7.5.

The average cellular fluorescence was measured in triplicate samples using a spectrophotometer
(Synergy H1, Biotek). A standard curve was generated for WT and KO samples, showing a
linear relationship between intracellular pH and relative fluorescence units.

3   Results

3.1   Bioinformatics prioritizes rs35907548 as a likely regulatory SNP

We used diverse bioinformatics tools to assess the regulatory potential of all 77 high-linkage
disequilibrium (LD; r^2>80%) SNPs in and around the *SLC15A4* locus (**Supplementary Table 2**).
These 77 SNPs were generated from 5 significantly associated ($p<5 \times 10^{-8}$) index SNPs
(rs10847697/rs1385374 [28], rs12370194 [29], rs10593112 [30], and rs11059928 [31]).
RegulomeDB2 ranked all SNPs for regulatory potential, prioritizing intronic rs35907548
(**Supplementary Table 2**). We used ENCODE [24]-annotated histone marks, chromatin
accessibility, and RNA Pol II occupancy to identify active chromatin, within which rs35907548
lies (**Figure 1**). ENCODE annotated rs35907548 as a "distal enhancer-like" candidate *cis*-
regulatory element (cCRE). We used PCHi-C chromatin-conformation data to identify several
regions topologically associated with rs35907548 in multiple immune cells (**Figure 1**). To
investigate the impact of rs35907548 on target gene expression, we conducted a thorough search
in publicly available expression quantitative trait locus (eQTL) databases, specifically focusing
on patient-derived primary immune cells [25]. Our findings indicate that rs35907548 acts as an
eQTL for SLC15A4 in various immune cell types (**Supplementary Figure 2**). Notably,
plasmacytoid dendritic cells (pDCs) exhibit the most significant eQTL effect on SLC15A4.
Finally, rs35907548 is in the middle of multiple transcription factor binding sites
(**Supplementary Figure 2**), where the risk allele T is universally conserved among all the
motifs. These transcription factors include Ikaros family zinc finger protein 1 (IKZF1), IKZF3,
E74-like factor 1 (ELF1), Friend leukemia integration 1 transcription factor (FLI1), and ETS1,
all of which are SLE risk genes themselves [32, 33, 34, 35, 36].

3.2   Allele-specific enhancer activity

179 We tested rs35907548 allele-specific enhancer activity in HEK293, Jurkat, HL-60, and LCL B-
180 cells with dual-luciferase reporter enhancer assays. The rs35907548 risk allele (TT) showed
181 higher enhancer activity than the non-risk allele (CC) in all cells (*p*-values: < 0.0001, < 0.001,
182 0.016, 0.0016; **Figure 2a**).

3.3   Long-range chromatin interactions

184 We performed 3C-qPCR between the rs35907548 locus and target gene promoters (**Figure 2b**).
185 The rs35907548 locus interacted with the promoters of *SLC15A4, GLT1D1*, and lncRNA
186 *AC069262*.1 (**Figure 1**).

3.4   Validation with CRISPR-based genome editing

To validate the transcriptional regulatory effects of the rs35907548 locus, we used three short-
guide RNAs (sgRNAs) to delete ~140 bases around rs35907548 in HL-60 cells (**Figure 4a** and
Supplementary Figure 3). The deletion was confirmed with Sanger sequencing, and ICE
analysis demonstrated high indel efficiency (63%) of pooled cells (data not shown).

Subsequently, we determined expression levels of *SLC15A4, GLT1D1*, and lncRNA *AC069262*.1
in KO and WT cells. Expression levels were lower in KO cells than WT (~45%, $p < 0.0001$;
~25%, $p < 0.01$; and ~35%, $p < 0.001$, respectively; **Figure 4b**).

3.5   rs35907548-modulated endolysosomal acidification

196 *SLC15A4* plays a crucial role in maintaining the acidity of the endolysosomal compartment [37].
197 We found that KO cells expressed less *SLC15A4*; consistent with this, KO cells showed a
198 dramatic increase over WT in endolysosomal pH (5.3 *vs*. 4.5, $p < 0.01$; **Figure 5**).

4   Discussion

Identifying true causal variants and their mechanisms of action from broad GWAS peaks is
challenging [38] but an essential prequisite for understanding disease risk and possibilities for
diagnosis and treatment. Several studies have linked *SLC15A4* SNPs to SLE risk in Asians and
Europeans [18, 19, 39], albeit with no experimental validation.

Here, we localize *SLC15A4* SLE risk to the intronic SNP rs35907548, which lies at the center of
an enhancer modulating expression of – and physically interacting with – *SLC15A4* and nearby
genes. Dual-luciferase reporter assays revealed strong enhancer activity, particularly of the risk
"TT" allele, in diverse cell types. Consistently, the risk allele showed substantially higher levels
of binding to active histone marks. We confirmed the activity of the rs35907548 enhancer in
cells by creating CRISPR/Cas9 knock-out cells, which showed significantly lower levels of
SLC15A4, *GLT1D1*, and the uncharacterized lncRNA *AC069262.1*. SLC15A4 is a histidine
transporter located primarily on the endolysosomal membrane, where it plays several pivotal
roles including maintenance of endolysosomal acidification [19, 37], thus regulating protein
degradation, inflammation, endocytosis, and autophagy, among other processes [40, 41, 42].
Accordingly, KO cells failed to properly acidify the endolysosome, demonstrating the profound
effects of the rs35907548 enhancer on cell physiology.

Dysregulation of lysosomal acidification is linked to several autoimmune diseases, including
SLE [21, 43]. In SLE, immune cells including macrophages, dendritic cells, and B-cells
accumulate autophagic vacuoles with impaired acidification, leading to defective autophagy and
abnormal antigen presentation. This dysregulated autophagy, coupled with defective clearance of
apoptotic cells, likely promotes autoimmunity in SLE. Several studies have proposed SLC15A4
as a therapeutic target for SLE and other autoimmune diseases [21, 37]. SLC15A4 agonists or
gene therapy could potentially restore proper functioning of lysosomal activity and a resulting
recovery of immune tolerance. Our study reveals the primary site of SLE risk at *SLC15A4* –
given its critical role in endolysosomal activity regulation, this SNP presumably underlies
association of other autoimmune diseases, as well.

rs35907548 is in the middle of multiple conserved transcription factor binding sites, including
those for the SLE risk gene and white blood cell factors IKZF1, IKZF3, ELF1, FLI1, and ETS1;
the ancestral risk allele T is universally present in the binding motifs. It is likely that proper
binding of these transcription factors is critical to enhancer function, and indeed the risk allele T
shows much higher levels of active chromatin marks and enhancer activity.

The adjacent locus Glycosyltransferase 1 Domain Containing 1 (*GLT1D1*) has also been flagged as an SLE risk locus [44]. GLT1D1 targets programmed cell death-ligand 1 (PD-L1) for glycosylation [45]; glycosylated PD-L1 is strongly immunosuppressive and correlates with B-cell non-Hodgkin's lymphoma progression. In addition to the links to endolysosomal acidification, TLR signaling, and autophagy through SLC15A4, future experiments will investigate the contribution of rs35907548 and other SNPs to SLE risk through *GLT1D1*. The lncRNA *AC069262.1* has not been thoroughly investigated and its role in biological processes and diseases, including SLE, remains largely unexplored. Further investigations are required to uncover its potential significance and contribution to SLE.

5 Data availability statement

All supporting data is available upon request from the corresponding author.

6 Conflict of interest

Authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7 Author Contributions

SKN contributed to conception and design of the study. GPM, MKS, HRR conducted and validated the experiments. MFN, LLL, SKN organized the database and conducted bioinformatics analyses. SKN, GPM, MKS, LLL wrote the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

8 Funding

Research reported in this publication was supported by National Institutes of Health grants R01 AR060366 and R21 AI168943 (SKN).

9 Acknowledgments

The authors wish to sincerely express their appreciation to Dr. Harris for her invaluable advice on setting up the pHrodo experiments. Furthermore, they extend their thanks to Ms. Louise Williamson for her technical assistance during the preparation of the manuscript.

10 Figure Legends

Figure 1. Bioinformatics. Cartoon showing the genomic location of rs35907548 (black vertical line), PCHi-C interactions from rs35907548 to the promoters (rectangle boxes) of the neighboring genes, regulatory histone marks and Pol 2.

Figure 2. Allelic-specific regulatory effects with enhancer containing rs35907548. (A) Luciferase reporter assays on 4 cell types. **(B)** ChIP-qPCR with risk (R) and non-risk (NR) LCLs.

Figure 3. 3C interaction analysis. (A) Schematic representation of rs35907548 region and
neighbor genes region. The SacI restriction enzyme sites were used to design primer. Small
arrows represent the primer location with orientation. The common prime at rs35907548 region
is the common for all other primers at SLC15A4, GLT1D1 and TMEM132C promoter regions.
Big arrow heads represent the transcriptional start site (TSS) of respective gene. **(B-E)** The
Graph represents the relative interaction of rs35907548 regions with different genomic regions in
Cell line HEK, Jurkat **(B, C)** and EBV transformed LCL B-cells **(D, E)**. The relative interaction
frequency for each set of primers represents the intensity of PCR. The 'X' axis is showing the
genomic distance (in kb) of the interacting region in forward and reverse direction from SNP
rs35907548(0 kb).

Figure 4. CRISPR/Cas9-based enhancer deletion. (A) CRISPR-based deletion, **(B)** qPCR-
based target gene expression between WT and KO.

Figure 5. Comparing the endolysosomal acidification between WT and KO cells. (A)
pHrodo^{TM} Red AM stained WT and KO cells under a fluorescent microscope. **(B)** Standard
curve for WT and KO cells using pHrodo^{TM} Red AM with Intracellular pH Calibration Buffer
Kit for the translation of fluorescence ratios into pH. **(C)** pH determined for WT and KO using
standard curve and suggested manufacturer's protocol.

Supplementary Figure 1. Immune cell type-specific eQTLs for rs35907548. The genotypes
and their corresponding sample sizes are shown at the bottom of each box plot. (C/C) is the risk
for rs35907548. USM = unstimulated, SM = stimulated, DN = double negative.

Supplementary Figure 2. Transcription factor binding sites overlapping rs35907548. The
ancestral risk allele T (red oval) produces more favorable motifs for all TFs. Green highlights
bases where the consensus motif is present.

Supplementary Figure 3: CRISPR/Cas9-based enhancer deletion of 139 bp region containing
rs35907548.

References

1.	Crow MK. Pathogenesis of systemic lupus erythematosus: risks, mechanisms and therapeutic targets. Ann Rheum Dis. 2023.

2.	Aguirre A, Izadi Z, Trupin L, Barbour KE, Greenlund KJ, Katz P, et al. Race, Ethnicity, and Disparities in the Risk of End-Organ Lupus Manifestations Following a Systemic Lupus Erythematosus Diagnosis in a Multiethnic Cohort. Arthritis Care Res (Hoboken). 2023;75(1):34-43.

3.	Kabani N, Ginzler EM. Is ethnicity linked to the severity of SLE manifestations? Nat Rev Rheumatol. 2019;15(9):515-6.

4.	Lewis MJ, Jawad AS. The effect of ethnicity and genetic ancestry on the epidemiology, clinical features and outcome of systemic lupus erythematosus. Rheumatology (Oxford). 2017;56(suppl_1):i67-i77.

5.	Maningding E, Dall'Era M, Trupin L, Murphy LB, Yazdany J. Racial and Ethnic Differences in the Prevalence and Time to Onset of Manifestations of Systemic Lupus Erythematosus: The California Lupus Surveillance Project. Arthritis Care Res (Hoboken). 2020;72(5):622-9.

6.	Kuo CF, Grainge MJ, Valdes AM, See LC, Luo SF, Yu KH, et al. Familial Aggregation of Systemic Lupus Erythematosus and Coaggregation of Autoimmune Diseases in Affected Families. JAMA Intern Med. 2015;175(9):1518-26.

7.	Harley ITW, Sawalha AH. Systemic lupus erythematosus as a genetic disease. Clin Immunol. 2022;236:108953.

8.	Molineros JE, Yang W, Zhou XJ, Sun C, Okada Y, Zhang H, et al. Confirmation of five novel susceptibility loci for systemic lupus erythematosus (SLE) and integrated network analysis of 82 SLE susceptibility loci. Hum Mol Genet. 2017;26(6):1205-16.

9.	Owen KA, Bell KA, Price A, Bachali P, Ainsworth H, Marion MC, et al. Molecular pathways identified from single nucleotide polymorphisms demonstrate mechanistic differences in systemic lupus erythematosus patients of Asian and European ancestry. Sci Rep. 2023;13(1):5339.

10.	Theofilopoulos AN, Kono DH, Baccala R. The multiple pathways to autoimmunity. Nat Immunol. 2017;18(7):716-24.

11.	Owen KA, Price A, Ainsworth H, Aidukaitis BN, Bachali P, Catalina MD, et al. Analysis of Trans-Ancestral SLE Risk Loci Identifies Unique Biologic Networks and Drug Targets in African and European Ancestries. Am J Hum Genet. 2020;107(5):864-81.

12.	Wang YF, Wei W, Tangtanatakul P, Zheng L, Lei Y, Lin Z, et al. Identification of Shared and Asian-Specific Loci for Systemic Lupus Erythematosus and Evidence for Roles of Type III Interferon Signaling and Lysosomal Function in the Disease: A Multi-Ancestral Genome-Wide Association Study. Arthritis Rheumatol. 2022;74(5):840-8.

13.	Toyama-Sorimachi N, Kobayashi T. Lysosomal amino acid transporters as key players in inflammatory diseases. Int Immunol. 2021;33(12):853-8.

14.	Yamashita T, Shimada S, Guo W, Sato K, Kohmura E, Hayakawa T, et al. Cloning and functional expression of a brain peptide/histidine transporter. J Biol Chem. 1997;272(15):10205-11.

15.	Daniel H, Kottra G. The proton oligopeptide cotransporter family SLC15 in physiology and pharmacology. Pflugers Arch. 2004;447(5):610-8.

16.	Smith DE, Clémençon B, Hediger MA. Proton-coupled oligopeptide transporter family SLC15: physiological, pharmacological and pathological implications. Mol Aspects Med. 2013;34(2-3):323-36.

17.	Heinz LX, Lee J, Kapoor U, Kartnig F, Sedlyarov V, Papakostas K, et al. TASL is the SLC15A4-associated adaptor for IRF5 activation by TLR7-9. Nature. 2020;581(7808):316-22.

18.	Baccala R, Gonzalez-Quintial R, Blasius AL, Rimann I, Ozato K, Kono DH, et al. Essential requirement for IRF8 and SLC15A4 implicates plasmacytoid dendritic cells in the pathogenesis of lupus. Proc Natl Acad Sci U S A. 2013;110(8):2940-5.

19.	Katewa A, Suto E, Hui J, Heredia J, Liang J, Hackney J, et al. The peptide symporter SLC15a4 is essential for the development of systemic lupus erythematosus in murine models. PLoS One. 2021;16(1):e0244439.

20.	Kobayashi T, Tsutsui H, Shimabukuro-Demoto S, Yoshida-Sugitani R, Karyu H, Furuyama-Tanaka K, et al. Lysosome biogenesis regulated by the amino-acid transporter SLC15A4 is critical for functional integrity of mast cells. Int Immunol. 2017;29(12):551-66.

21.	Rimann I, Gonzalez-Quintial R, Baccala R, Kiosses WB, Teijaro JR, Parker CG, et al. The solute carrier SLC15A4 is required for optimal trafficking of nucleic acid-sensing TLRs and ligands to endolysosomes. Proc Natl Acad Sci U S A. 2022;119(14):e2200544119.

22.	Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, et al. Annotation of functional variation in personal genomes using RegulomeDB. Genome Res. 2012;22(9):1790-7.

23.	Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring population-specific haplotype structure and linking correlated alleles of possible functional variants. Bioinformatics. 2015;31(21):3555-7.

24.	Consortium EP. An integrated encyclopedia of DNA elements in the human genome. Nature. 2012;489(7414):57-74.

25.	Ota M, Nagafuchi Y, Hatano H, Ishigaki K, Terao C, Takeshima Y, et al. Dynamic landscape of immune cell-specific gene regulation in immune-mediated diseases. Cell. 2021;184(11):3006-21.e17.

26.	Fazel-Najafabadi M, Rallabandi HR, Singh MK, Maiti GP, Morris J, Looger LL, et al. Discovery and Functional Characterization of Two Regulatory Variants Underlying Lupus Susceptibility at 2p13.1. Genes (Basel). 2022;13(6).

27.	Singh B MGPZXF-NMBSCSCTCOYCKHKYG. Lupus susceptibility region containing CDKN1B rs34330 mechanistically influences expression and function of multiple target genes, also linked to proliferation and apoptosis. Arthritis Rheumatol. 2021.

28.	Han JW, Zheng HF, Cui Y, Sun LD, Ye DQ, Hu Z, et al. Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus. Nat Genet. 2009;41(11):1234-7.

29.	Sun C, Molineros JE, Looger LL, Zhou XJ, Kim K, Okada Y, et al. High-density genotyping of immune-related loci identifies new SLE risk variants in individuals with Asian ancestry. Nat Genet. 2016.

30.	Langefeld CD, Ainsworth HC, Cunninghame Graham DS, Kelly JA, Comeau ME, Marion MC, et al. Transancestral mapping and genetic load in systemic lupus erythematosus. Nat Commun. 2017;8:16021.

31.	Yin X, Kim K, Suetsugu H, Bang SY, Wen L, Koido M, et al. Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus. Ann Rheum Dis. 2020.

32.   Cai X, Qiao Y, Diao C, Xu X, Chen Y, Du S, et al. Association between polymorphisms
of the IKZF3 gene and systemic lupus erythematosus in a Chinese Han population. PLoS One.
2014;9(10):e108661.
33.   Garrett-Sinha LA, Kearly A, Satterthwaite AB. The Role of the Transcription Factor Ets1
in Lupus and Other Autoimmune Diseases. Crit Rev Immunol. 2016;36(6):485-510.
34.   Hu SJ, Wen LL, Hu X, Yin XY, Cui Y, Yang S, et al. IKZF1: a critical role in the
pathogenesis of systemic lupus erythematosus? Mod Rheumatol. 2013;23(2):205-9.
35.   Sato S, Lennard Richard M, Brandon D, Jones Buie JN, Oates JC, Gilkeson GS, et al. A
critical role of the transcription factor fli-1 in murine lupus development by regulation of
interleukin-6 expression. Arthritis Rheumatol. 2014;66(12):3436-44.
36.   Yang J, Yang W, Hirankarn N, Ye DQ, Zhang Y, Pan HF, et al. ELF1 is associated with
systemic lupus erythematosus in Asian populations. Hum Mol Genet. 2011;20(3):601-7.
37.   Kobayashi T, Shimabukuro-Demoto S, Yoshida-Sugitani R, Furuyama-Tanaka K, Karyu
H, Sugiura Y, et al. The histidine transporter SLC15A4 coordinates mTOR-dependent
inflammatory responses and pathogenic antibody production. Immunity. 2014;41(3):375-88.
38.   Claussnitzer M, Cho JH, Collins R, Cox NJ, Dermitzakis ET, Hurles ME, et al. A brief
history of human disease genetics. Nature. 2020;577(7789):179-89.
39.   He CF, Liu YS, Cheng YL, Gao JP, Pan TM, Han JW, et al. TNIP1, SLC15A4, ETS1,
RasGRP3 and IKZF1 are associated with clinical features of systemic lupus erythematosus in a
Chinese Han population. Lupus. 2010;19(10):1181-6.
40.   Bonam SR, Wang F, Muller S. Lysosomes as a therapeutic target. Nat Rev Drug Discov.
2019;18(12):923-48.
41.   Cao M, Luo X, Wu K, He X. Targeting lysosomes in human disease: from basic research
to clinical applications. Signal Transduct Target Ther. 2021;6(1):379.
42.   Noguchi M, Hirata N, Tanaka T, Suizu F, Nakajima H, Chiorini JA. Autophagy as a
modulator of cell death machinery. Cell Death Dis. 2020;11(7):517.
43.   Harris VM, Harley ITW, Kurien BT, Koelsch KA, Scofield RH. Lysosomal pH Is
Regulated in a Sex Dependent Manner in Immune Cells Expressing CXorf21. Front Immunol.
2019;10:578.
44.   Joo YB, Lim J, Tsao BP, Nath SK, Kim K, Bae SC. Genetic variants in systemic lupus
erythematosus susceptibility loci, XKR6 and GLT1D1 are associated with childhood-onset SLE
in a Korean cohort. Sci Rep. 2018;8(1):9962.
45.   Liu X, Zhang Y, Han Y, Lu W, Yang J, Tian J, et al. Overexpression of GLT1D1 induces
immunosuppression through glycosylation of PD-L1 and predicts poor prognosis in B-cell
lymphoma. Mol Oncol. 2020;14(5):1028-44.
414

Figure 1. **Bioinformatics.** Cartoon showing the genomic location of rs35907548 (black vertical line), PCHi-C interactions from rs35907548 to the promoters (rectangle boxes) of the neighboring genes, regulatory histone marks and Pol 2.
Data: Monocytes, Naive_B, Total_B, Total_CD8, Neutrophils from *Javiere et al. 2015*.

Figure 2. Allelic-specific regulatory effects with enhancer containing rs35907548. (A) Luciferase reporter assays on 4 cell types. **(B)** ChIP-qPCR with risk (R) and non-risk (NR) LCLs

Figure 3: **3C interaction analysis. (A)** Schematic representation of rs35907548 region and neighbor genes region. The SacI restriction enzyme sites were used to design primer. Small arrows represent the primer location with orientation. The common prime at rs35907548 region is the common for all other primers at SLC15A4, GLT1D1 and TMEM132C promoter regions. Big arrow heads represent the transcriptional start site (TSS) of respective gene. (B-E) The Graph represents the relative interaction of rs35907548 regions with different genomic regions in Cell line HEK , Jurkat (B, C) and EBV transformed LCL B-cells (D, E). The relative interaction frequency for each set of primerprimers represents the intensity of PCR. The 'X' axis is showing the genomic distance (in kb) of the interacting region in forward and reverse direction from SNP rs35907548(0 kb).

Figure 4. CRISPR/Cas9-based enhancer deletion. (A) CRISPR-based deletion, (B) qPCR-based target gene expression between WT and KO

Figure 5. Comparing the endolysosomal acidification between WT and KO cells. (A) pHrodo™ Red AM stained WT and KO cells under a fluorescent microscope. (B) Standard curve for WT and KO cells using pHrodo™ Red AM with Intracellular pH Calibration Buffer Kit for the translation of fluorescence ratios into pH. (C) pH determined for WT and KO using standard curve and suggested manufactures protocol.

Supple Fig 1. Immune cell type-specific eQTLs for rs35907548. The genotypes and their corresponding sample sizes are shown at the bottom of each box plot. (C/C) is the risk for rs35907548. USM = unstimulated, SM = stimulated, DN = double negative.

Supp Fig. 2. Transcription factor binding sites overlapping rs35907548. The ancestral risk allele T (red oval) produces more favorable motifs for all TFs. Green highlights bases where the consensus motif is present.

Supple Fig 3: CRISPR/Cas9-based enhancer deletion of 139 bp region containing rs35907548