**Title**

Surface crosslinking of virus-like particles increases resistance to

proteases, low pH, and mechanical stress for mucosal applications

**Authors & Affiliations**

Ahmed Ali [a], Suwannee Ganguillet [a], Yagmur Turgay [a], Tim Keys [a], Erika Causa [b], Ricardo Fradique

[b], Viviane Lutz-Bueno [c,d], Serge Chesnov [e], Chia-wei Lin [e], Verena Lentsch [a], Jurij Kotar [b], Pietro

Cicuta [b], Raffaele Mezzenga [a], Emma Slack [a,*], Milad Radiom [a,c,*]

[a] Institute of Food, Nutrition and Health, ETH Zürich, Zürich, Switzerland
[b] Biological and Soft Systems, Cavendish Laboratory, University of Cambridge, Cambridge CB3
0HE, U.K.
[c] Paul Scherrer Institute PSI, Villigen, Switzerland
[d] Laboratoire Léon Brillouin, CEA-CNRS (UMR-12), CEA Saclay, Université Paris-Saclay, 91191,
Gif-sur-Yvette Cedex, France
[e] University of Zürich/ETH Zürich, Functional Genomics Centre Zürich, Zürich, Switzerland
16

*Milad Radiom (milad.radiom@hest.ethz.ch); Emma Slack (emma.slack@hest.ethz.ch)

**Abstract**

Virus-like particles (VLPs) are emerging as nano-scaffolds in a variety of biomedical

applications including the delivery of vaccine antigens to mucosal surfaces. These soft,

colloidal, and proteinaceous structures (capsids) are nevertheless susceptible to mucosal

environmental factors which limit their usefulness. We addressed this issue by crosslinking

multiple capsid surface reactive residues using polyethylene glycol tethers. Surface

crosslinking enhanced the colloidal stability and mechanical strength of VLPs against low

pH, proteases, and mechanical agitation, while it did not interfere with function as vaccine.

Chemical crosslinking thus offers a viable means to enhance the resilience of VLPs in

mucosal applications.

**Keywords**

Virus-like particle; Polyethylene glycol; Surface-crosslinking; Force spectroscopy; In vitro

human nasal epithelial tissue; Mucus

## Introduction

Virus-like particles (VLPs) have emerged as invaluable nano-carriers in biomedical

applications (*1, 2*). Sensitive cargo can be protected by packaging it inside their capsid

while genetic and/or chemical modifications can be used to deliver vaccine antigens or to

target the cargo to specific cell types. In recent years, various therapeutic molecules have

been integrated into VLPs for immunotherapy (*3*), gene therapy (*4-6*), chemotherapy (*7-*

*9*), as well as for contrast imaging and photothermal therapy (*10, 11*).

As carriers, VLPs must endure environmental challenges and it is likely that their material

properties are an important factor determining their fate (*12*). General key considerations

for "delivery-capable" VLPs include size, colloidal stability, target specificity, and

responsiveness to stimuli for cargo release. For mucosal applications, VLPs are faced

with additional physiological challenges including a thick and adhesive mucus gel, mucus

regeneration and clearance, high concentrations of proteases and secretory

immunoglobulin A (sIgA) antibodies and intermittent mechanical agitation (*13-15*).

Together these effects can significantly reduce the efficacy of VLPs for mucosal

applications (**Figure 1**(a)).

To this end, there is compelling evidence that viruses, viral vectors, and VLPs disintegrate

under mechanical force (*16-25*), and in unfavorable chemical environments (*26, 27*). This

raised the question: can surface crosslinking using macromolecule tethers enhance the

stability of VLP capsids? To address this question, we employed homobifunctional

polyethylene glycol (PEG) to tether surface lysine residues on capsid surface. VLP or viral

vector PEGylation, without crosslinking, has been exploited as a means to enhance

biological function, especially with adenovirus vectors (Adv). For example, PEGylation

protected Adv from neutralizing antibodies, prolonged transgene expression and reduced

immune activation against the vector (*28-30*). Subsequent experiments revealed a

complex dependence on bio-physicochemical properties (*31-33*). Thereby, for example,

conjugation optimization led to reduced anti-vector serum antibody and more persistent

transgene expression (*34*). Other macromolecules were conjugated to bacteriophage Qβ

VLP to shield it from immune recognition (*35, 36*). In other directions, PEG was used to

tether a fluorescent molecule and folic acid to cowpea mosaic virus for intravital vascular

imaging and targeted cargo delivery to cancer cell lines, respectively (*37, 38*).

Our model VLP is assembled from 180 copies of *Acinetobacter* phage coat protein AP205

(*25*). AP205 VLP is a versatile platform for genetic and/or chemical conjugation of various

antigenic molecules thus potentiating a vaccination strategy (*25, 39-41*). We investigated

surface crosslinking of AP205 VLP using PEG tethers. To explore different modes of

PEGylation, the AP205 VLP was reacted with homobifunctional (leading to surface

crosslinking) and monofunctional (leading to simple PEGylation) linkers. We found that

surface crosslinking is a viable toolbox that enhances the various stability profiles of VLPs,

of which resilience to mechanical agitation, pH and protease stress are related to mucosal

applications. Particularly, PEG crosslinking did not shield immune recognition of capsid

coat proteins, enhancing vaccine stability without inhibiting immunogenicity.

# Results

Surface crosslinked and simply PEGylated VLPs were produced using homobifunctional

and monofunctional PEG molecules respectively. The resulting VLPs are named $bPEG_n$-

and $mPEG_n$-AP205 VLP where n is the number of PEG monomers. The functional

group(s) of the PEG molecules was N-hydroxysuccinimide (NHS) ester which reacts with

lysine residues and N-terminus of coat proteins (c.f. **Figure 1**(b)). The sequence of AP205

monomer and the details of the PEG molecules are presented in **S1**. We note that in the

conditions used, the crosslinking was at single VLP level, and we did not observe

aggregation between the particles (**S2**). Transmission electron microscopy (TEM) showed

that the VLPs retained their spherical geometry after the reaction with the linker molecules

and no detectable VLP size change for any lengths of PEG crosslinker tested, n = 9–25,

was observed (**S3**).

**Crosslinking AP205 VLP capsid using polyethylene glycol (PEG) molecules**

Investigations with mass spectrometry and reducing SDS-polyacrylamide gel

electrophoresis (SDS PAGE) revealed that AP205 VLP reaction with homobifunctional

PEG crosslinkers resulted in PEG-crosslinked AP205 dimers, trimers, and possibly higher

oligomers. In the representative SDS PAGE shown in **Figure 1**(c), in the lanes associated

with $bPEG_n$-AP205 VLPs (n = 9 to 25), bands in the MW range 37–50 kDa (marked with

purple arrows/circles) correspond to PEG-crosslinked AP205 dimers, and bands in MW

range 50–75 kDa (marked with orange arrows/pentagons) correspond to PEG-crosslinked

AP205 trimers. Using nano ultra-performance liquid chromatography coupled to mass

spectroscopy, it was found that lysine (K) residues K4 and K61 and K4 and K94 were

involved in the crosslinking reaction (Table S4-7). In **Figure 1**(d), a schematic of PEG

linkages between individual AP205 monomers near the three-fold axis of symmetry is

shown. Using coordinates from the cryo-electron microscopy reconstruction of the AP205

VLP (5LQP) in Visual Molecular Dynamics (VMD) (*41, 42*), the N-N interatomic distances

in K4–K61 linkage were estimated to be 1.8 nm and in K4–K94 linkage to be 1.6 nm and

2.4 nm. Calculation of end-to-end probability distributions of $PEG_9$, $PEG_{17}$ and $PEG_{25}$

linker molecules using worm-like chain model are shown in **Figure 1**(d) (*43*). The

calculated extension lengths are consistent with the interatomic distances. Additional

details regarding the conjugation stoichiometry, mass spectroscopy and SDS PAGE are

found in **S2**, **S4** and **S5**, respectively.

## Enhanced colloidal and enzymatic stability of surface-crosslinked AP205 VLPs

VLPs are potentially interesting scaffolds for the oral or intranasal delivery of biomolecules,

requiring stability in pH-varying environments and in the presence of high concentrations

of proteases (*1, 2, 12*). Therefore, increasing their stability in the complex mucosal

environment is of high translational relevance.

Experiments using dynamic light scattering (DLS) showed that native VLP was stable at

pH 7.4 but aggregated at pH 2.0–6.0 (**Fig. S6-1**). PEG-crosslinking increased the stability

of VLP down to pH ~ 4.0 (**Fig. S6-2**). We then tested the enzymatic stability of PEG-

crosslinked VLPs in pig and mouse gastric fluids. In PBS, an incubation time of 2 h (or

longer) at 37°C did not affect the stability or electrophoretic mobility of the VLPs (*25*)

(**Figure 1**(e)). In pig gastric fluid, which had a pH of 3.0 and high protease concentration,

PEG-crosslinked VLPs, namely $bPEG_9$-AP205 VLP, $bPEG_{17}$-AP205 VLP, and $bPEG_{25}$-

AP205 VLP, showed improved stability after 5 min of incubation, i.e., a band, marked with

arrow, remained at the expected MW, with fewer breakdown products than the unmodified

VLP. At the later time points of 30 min, 1 h and 2 h, the intact portion was disintegrated,

and signs of aggregation and accumulation of material in the wells of the gel appeared.

Interestingly, the PEGylated but not crosslinked VLP, namely $mPEG_{25}$-AP205 VLP,

showed a similar behavior to native VLP and disintegrated in as little as 5 min. This

observation indicates that PEG-crosslinking VLP capsid, but not PEGylation per se, is a

viable approach to overcome the susceptibility of VLPs to colloidal and enzymatic

instability.

Experiments in pig gastric fluid at pH 4.7 corroborated the stabilizing effect of PEG-

crosslinking, but not simple PEGylation, in the presence of proteases (**Fig. S7-1**). At this

pH, the PEG-crosslinked VLPs were stable at all time points. In pig gastric fluid at pH 5.5,

all VLPs were found to be stable over 1 h of incubation (**Fig. S7-2**). Incubation in mouse

gastric fluid at pH 4.6 presented a similar behavior where the PEG-crosslinked VLPs

appeared to be more stable compared to the naked and PEGylated but not crosslinked

VLP (**Fig. S7-3**).

**Figure 1: Susceptibility of virus-like particles (VLPs) to mucosal factors, and their capsid stabilization by means of surface crosslinking using macromolecule tethers.**
(a) VLPs are exposed to physical and chemical challenges on mucosal surfaces. (b) Schematic of AP205 VLP and schematic of one coat protein (CP) dimer (*41*). The N-terminus (blue) of one monomer, the C-terminus (red) of the other monomer, together with several lysine residues are marked. (c) Reducing SDS polyacrylamide gel electrophoresis

of native, PEG-crosslinked (bPEG$_n$-, n = 9–25), and PEGylated but not crosslinked AP205
VLPs (mPEG$_n$-, n = 12 or 25). M is PageRuler Plus Prestained Protein Ladder. The
monomer region is marked with cyan arrows/squares, dimer region with purple
arrows/circles and trimer region with orange arrows/pentagons. (d) Schematic of AP205
CP arrangement near three-fold axis of symmetry and the assignment of lysine residues
K4, K61 and K94 that participate in crosslinking reaction. In the schematic, 3 dimers are
shown with their monomers colored in blue and green. Only one tether (red coil), that is
either of the two of K4–K94 possibilities or the one of K4–K61 possibility, can occur. The
end-to-end probability distributions of bPEG$_n$ tethers (n = 9, 17 and 25) calculated from an
analytical worm-like chain model (*43*). (e) Agarose gel electrophoresis of AP205 VLP,
bPEG$_9$-AP205 VLP, bPEG$_{17}$-AP205 VLP, bPEG$_{25}$-AP205 VLP and mPEG$_{25}$-AP205 VLP
in PBS and in pig gastric fluid (pH 3.0), respectively on the left and right of the gel.
Incubation times varied between 5 min, 30 min, 1 hour and 2 hours at 37°C. Arrows at
time point 5 min indicate persistent PEG-crosslinked AP205 VLPs in gastric fluid. M stands
for marker.
157

**Enhanced stiffness of surface-crosslinked AP205 VLPs**

Susceptibility to mechanical stress sets a limit on potential applications of VLPs in various

indications. This occurs during interactions with cells, e.g. during endocytosis or antigen

presentation to B cells, or when exposed to mucus or fecal flow (*44-50*). Therefore, we

examined the nanomechanical properties of native and PEG-crosslinked VLPs using

atomic force microscopy (AFM) force-indentation spectroscopy.

The schematic of AFM measurements is depicted in **Figure 2**(a) and details are provided

in the methods section. The example shown in **Figure 2**(b) depicts a single representative

AP205 VLP for which the height prior to indentation ( $H_\mathrm{i}$ ) and after indentation ( $H_\mathrm{f}$ ) were

evaluated. **Figure 2**(c) shows the heights distributions of AP205 VLP with the respective

average values. The force versus indentation distance ( $F - \delta$ ) curve was used to

calculate the stiffness, and the force versus indentation time ( $F - t$ ) curve to calculate

the strain rate. An example of these curves for the same VLP (**Figure 2**(b)) is shown in

**Figure 2**(d). The linear part of the response was used to calculate the stiffness, $k = dF/d\delta$

, the total indentation, $\delta = \Delta F/k$ , and the indentation force rate $\dot{F} = dF/dt$ , which was

converted to strain rate via $\gamma = \dot{F}/k\delta$ . After an initial linear response, the VLP yielded at a

threshold force/indentation. The histograms of maximal force and indentation (indent.) of

AP205 VLP together with the average values of these parameters are shown in **Figure**

**2**(e). The physical significance of these parameters is that at a force (or indentation) higher

than the average maximal force (or indentation), yield occurs which significantly reduces

the original stiffness of the VLP (*16-18, 22-25*). The histograms of stiffness and strain rate

of AP205 VLP together with the average values of these parameters are shown in **Figure**

**2**(f) and (g), respectively. For this VLP, we previously showed that the standard linear solid

model (SLSM, **Figure 2**(h)) interprets the stiffness as a function of the strain rate (*25*). We

found that at a strain rate < 100 1/s, the overall mechanical response is governed by $k_0$

in SLSM. We retrieved a good agreement between the current evaluation of AP205

stiffness (47±15 pN/nm) with the previous one (53±23 pN/nm) at low strain rates.

**Figure 2: Mechanical properties of AP205 VLP investigated using AFM force-indentation spectroscopy.** (a) Schematic of AFM calibration on freshly cleaved mica, and force-indentation spectroscopy of HOPG-immobilized VLPs. Single VLPs were localized using PeakForce tapping imaging (base tapping rate of 1 kHz) prior to force

measurements using normal ramp at base approach-retraction rate at 1 Hz. (b) Example
of height evaluation for a single AP205 VLP before and after force application. (c) Height
distributions of AP205 VLP before and after force measurements. (d) Example of force-
indentation distance and force-indentation time curves for AP205 VLP. The linear regime
of force-indentation distance response was used to calculate the stiffness. In the same
range, a linear fit to force-indentation time response gave the indentation force rate. (e, f
and g) Histograms of maximal force and indentation (indent.) (e), stiffness (f) and strain
rate (g) of AP205 VLP. Mean ± standard deviation is reported. The number (n) of AP205
VLPs in repeat measurements is indicated. (h) Standard linear solid model (SLSM). At low
strain rates, there is no contribution from friction (or rate dependent resistance) to
indentation.
202

Similarly, the mechanical properties of PEG-crosslinked VLPs, namely $bPEG_9$-AP205

VLP, $bPEG_{17}$-AP205 VLP, $bPEG_{25}$-AP205 VLP were evaluated, and the results are shown

in **Figure 3**. The initial heights and the heights after a single indentation showed similarity

among the native and PEG-crosslinked VLPs. The distributions of maximal force and

indentation (indent.) however showed variations compared to native VLP. The average

maximal force was found to be higher and accordingly the average maximal indentation

lower compared to native VLP. The increased (decreased) maximal force (indentation) of

PEG-crosslinked VLPs indicated reduced susceptibility to mechanical agitation.

Accordingly, the average stiffness for these VLPs was also higher by a factor of 1.3 to 2.0

than the stiffness of native VLP at similar range of strain rate.

**Figure 3: Enhanced strength and stiffness of PEG-crosslinked AP205 VLPs.** Mechanical properties of PEG-crosslinked AP205 VLPs, namely bPEG$_9$-AP205 VLP, bPEG$_{17}$-AP205 VLP and bPEG$_{25}$-AP205 VLP, and PEGylated but not crosslinked mPEG$_{25}$-AP205 VLP investigated using AFM force-indentation spectroscopy. Distributions of height before and after indentation, maximal force/indentation in linear response regime, and stiffness/strain rate are shown together with mean ± standard deviation. The number (n) of VLPs in repeat measurements is indicated.

We then evaluated capsid size and shell thickness using small angle X-ray scattering

(SAXS). The X-ray scattering spectra of the native and PEGylated VLPs together with fits

to spherical core-shell form factor are shown in **Figure 4**(a). In each case, we found a

good agreement based on the low values for the fitting error ( $\chi^2$ ). In the summary panel

in **Figure 4**(b), it is shown that the diameter ( $D$ ) and shell thickness ( $h$ ) of VLP increased

after PEGylation. In addition, while the VLP diameter remained somewhat constant

between the different lengths of PEG tethers, the shell thickness increased with PEG

length and had its highest value for the PEGylated but not crosslinked VLP. Subsequent

to these evaluations, the elastic modulus was calculated from the relation $E = Dk/2h^2$ ,

**Figure 4**(b) (*51, 52*). The elastic moduli remained somewhat constant ~ 90 MPa among

the native and PEG-crosslinked VLPs. This observation can potentially be explained as

follows. In nanoindentation experiments, the shell compression and bending moduli

contribute to an effective elastic modulus $E$   (*52-54*). Tethered PEG molecules disfavor

bending because it results in chain stretching, therefore increasing the bending modulus.

Reduction in coat protein positive charges after PEG conjugation reduces the inter-atomic

repulsion during compression, therefore decreasing the compression modulus. The higher

stiffness of PEG-crosslinked VLPs compared to native VLP is therefore associated with

an increase in the shell thickness, as well as stretching the PEG molecules (*55*). The effect

of chain stretching became evident where the PEGylated but not crosslinked VLP, namely

mPEG$_{25}$-AP205 VLP, was investigated. This VLP showed a lower elastic modulus and

stiffness compared to its PEG-crosslinked counterpart, namely bPEG$_{25}$-AP205 VLP

(**Figure 3** and **Figure 4**).

**Figure 4: Geometrical and mechanical parameters of native and PEGylated AP205 VLPs.** (a) Small angle X-ray scattering (SAXS) spectra of AP205, bPEG$_9$-AP205, bPEG$_{17}$-AP205, bPEG$_{25}$-AP205 and mPEG$_{25}$-AP205 VLPs, together with fits to spherical core-shell form factor. (b) Summary panel of the geometrical and mechanical properties of the VLPs.

| VLP entry | Stiffness (pN/nm) | Diameter (nm) | Shell thickness (nm) | Elastic modulus (MPa) |
|---|---|---|---|---|
| AP205 | 47±15 | 27.6 | 2.7 | 89±28 |
| bPEG$_9$-AP205 | 59±17 | 28.8 | 3.2 | 83±24 |
| bPEG$_{17}$-AP205 | 89±45 | 28.5 | 3.6 | 98±49 |
| bPEG$_{25}$-AP205 | 94±35 | 28.4 | 4.0 | 83±31 |
| mPEG$_{25}$-AP205 | 76±29 | 28.8 | 4.4 | 57±22 |

**Surface-crosslinking does not restrict mucus translocation of AP205 VLP**

A major feature of intestinal and respiratory sites, where VLP application is relevant, is the production of mucus. Mucus is an adhesive and viscoelastic gel that covers the epithelium and functions to trap and dispose of foreign objects. In the airways, mucus is constantly pushed out via mucociliary clearance (*56, 57*). We used *in vitro* 3D human nasal epithelial tissues to investigate the effect of PEG-crosslinking on translocation across flowing mucus. The tissues were characterized as having fully differentiated mucus producing goblet cells, ciliated cells with motile cilia, as well as other cells of normal human nasal epithelial tissue. The schematic of the experiments is shown in **Figure 5**(a). On delivery of fluorescently labelled VLPs from the apical side, the VLPs translocated through mucus

by diffusion and simultaneously transported in flowing mucus in the direction of mucociliary

clearance to edges of the tissue. The lateral distribution and vertical translocation were

monitored using laser scanning confocal microscopy (LSCM). In **Figure 5**(b),

representative examples of mean fluorescence intensity (MFI) variation with height

collected in the center, [0, 0] mm, prior to VLP delivery (VLP-) and after the delivery of

$mPEG_{25}$-AP205 VLP[GelRed] (VLP+) are shown. In the former case, the location of maximum

MFI was at the height level of the semipermeable membrane which was assigned to 0 µm.

After VLP delivery, the maximum of MFI shifted to a higher height value (about 40 µm)

while the peak associated with semipermeable membrane was observable. The MFI vs

height spectra were normalized (norm. MFI) by the MFI values at height = 0 µm, **Figure**

**5**(b). The normalized MFI spectrum of VLP- was then subtracted from that of VLP+ to give

the net MFI spectrum. The net MFI spectra from the center, [0, 0] mm, together with

spectra at positions [-1, 0] mm, [+1, 0] mm, [0, -1] mm and [0, +1] mm, are shown in **Figure**

**5**(c). These spectra provided information about the height location of VLP deposition and

the net value of MFI at those deposition heights, which corresponded to the translocated

amount of VLPs. In all experiments, it was ensured via bright field microscopy that the cilia

were beating before and after VLP delivery using stablished procedures (*58*).

A typical deposition height and translocated amount of native and PEG-crosslinked VLPs

are shown in **Figure 5**(d) and (e). A similar deposition height between these VLPs was

expected as tissues from the same donor (tissue 3) were used. Furthermore, as shown in

**Figure 5**(e), similar translocated amounts between the native and PEG-crosslinked VLPs

were obtained. This observation indicated that PEG-crosslinking did not restrict nor

increased VLP translocation through mucus. These results were consistent on other

tissues (**S8**). We then investigated the translocation behavior of PEGylated but not

crosslinked VLP, namely $mPEG_{25}$-AP205 VLP[GelRed]. We found that $mPEG_{25}$-AP205

VLP[GelRed] showed a similar deposition height (**Figure 5**(d)), however it had an increased

translocation amount which exceeded native VLP by two folds (**Figure 5**(e)). Particularly

as compared to $bPEG_{25}$-AP205 VLP, this result indicated a higher mucus permeability of

the $mPEG_{25}$-AP205 VLP construct. Consistent with previous reports (*59-61*), a potential

explanation is an effective steric hindrance interactions with mucin glycoproteins from

one-end conjugated PEGs.

**Figure 5: Translocation through mucus of AP205 VLPs in *in vitro* 3D human nasal epithelial tissues.** (a) Schematic of *in vitro* nasal epithelial tissue and delivery of

fluorescently-labelled virus-like particles (VLPs) from the apical side. Using laser scanning
confocal microscopy (objective lens 10x, N.A. 0.3), the lateral distribution and vertical
translocation through mucus of VLPs were monitored at several position 1: [0, 0] mm, 2:
[-1, 0] mm, 3: [+1, 0] mm, 4: [0, -1] mm and 5: [0, +1] mm. (b) Mean fluorescence intensity
(MFI) and normalized MFI (norm. MFI) variation with height collected in the center of tissue
3. MFI and norm. MFI spectra prior to VLP delivery, VLP-, and after the delivery of
mPEG$_{25}$-AP205 VLP$^{GelRed}$, VLP+, are shown. The height of semipermeable membrane is
set to 0 µm. (c) Subtracted (or net) MFI spectra on five positions showing the height
location of VLP deposition and the net MFI. (d, e) The deposition height (d) and the net
MFI corresponding to the translocated amount of VLPs (c) of native and PEGylated VLPs
on tissue 3. (f) Delivery of mPEG$_{25}$-AP205 VLP$^{GelRed}$ to tissue 3. (g) Delivery of mPEG$_{25}$-
AP205 VLP$^{GelRed}$ to trans-cinnamaldehyde-treated tissue 3. The yellow box denotes the
area of mean fluorescence intensity (MFI) calculation and blue circle the height location
of VLP deposition. In these figures, height starts from above the air-liquid interface (ALI)
and increases in the direction of cell layer to below the semipermeable membrane. The
height of the semipermeable membrane is approximatively 250 µm. The individual
confocal images on lateral positions 1 to 6 were taken at a fixed height distance from the
semipermeable membrane.
315

We then investigated the effect of outward-swirling mucociliary clearance on the

deposition of VLPs across the tissue (*62*). mPEG$_{25}$-AP205 VLP$^{GelRed}$ was delivered in

normal tissue condition or when the tissue was treated with trans-cinnamaldehyde.

Analysis of the lateral distribution of deposited VLPs across the tissue (positions 1 to 6)

under normal conditions (**Figure 5**(f)) showed that a noticeable portion of VLP was cleared

to the edge of the plastic insert in flowing mucus while a portion translocated similar to the

observations already discussed. However, the lateral distribution of deposited VLPs was

more uniform across the tissue when ciliary beating was stopped (**Figure 5**(g)). Additional

examples of the deposited VLP distribution when ciliary beating was inhibited are shown

in **SI 9**.

**Compatibility of PEG-crosslinking with vaccine function**

A major application of VLPs is in mucosal delivery of vaccine antigens. We therefore

investigated if PEG-crosslinking may interfere with this function by measuring antibody

induction against coat protein in native and PEG-crosslinked VLPs in a subcutaneous

vaccination model using C57BL/6J specific pathogen free (SPF) mice, i.e. a setting where

vaccine stability is thought not to be a major limitation on performance. Accordingly,

subcutaneous injections of VLPs without additional adjuvant were performed on Day 0

and 10. The mice were sacrificed on Day 20 and anti-AP205 serum IgG titers detected

using enzyme-linked immunosorbent assay (ELISA). The serum antibody titers against

AP205 coat protein are shown in **Figure 6**(a). As expected, native VLP was immunogenic,

and we easily detected specific serum IgG antibodies against AP205 coat protein (*39, 40*).

Surprisingly, PEG-crosslinking, or simple PEGylation of VLP did not shield immune

activation against the coat protein; rather, specific antibody induction was increased.

Potentially several mechanisms could be involved, and one cannot make a direct

correlation with stiffness. Antibody induction against the PEG coating was also

investigated. Consistent with the strong adjuvanticity of VLPs (*63*), the anti-PEG ELISA in

**Figure 6**(b) showed that the injection of PEGylated VLPs, compared to the negative

control, i.e., native VLP, resulted in anti-PEG IgG induction. Furthermore, the anti-PEG

IgG titers seemed to increase with the size of two-end conjugated PEG crosslinker, i.e.,

to be higher in the order $bPEG_9$-AP205 VLP < $bPEG_{17}$-AP205 VLP < $bPEG_{25}$-AP205 VLP.

**Figure 6: Immune activation by PEG-crosslinked AP205 VLPs.** C57BL/6J SPF mice
were subcutaneously injected with native (n = 7), PEG-crosslinked (n = 13), or PEGylated
but not crosslinked (n = 3) AP205 VLPs on Day 0 and Day 10 and sacrificed on Day 20.
Serum antibody (IgG) titers against (a) AP205 coat protein, and (b) PEG was measured
by ELISA. Statistical significance was determined by ordinary one-way ANOVA with
Dunnett's multiple comparisons test, with a single pooled variance on log-normalized data
($*P = 0.0332$; $***P = 0.0021$; $***P = 0.0002$).
356

## Discussion

VLPs have similar physical characteristics to their viruses of origin, including size, proteinaceous core-shell structure, and patchy surface charge distribution, and like viruses are prone to aggregation, disintegration, and sticking to surfaces rendering them non-functional (*64*). Enhancing the stability of VLPs is therefore justified in future developments of VLP biotechnology, particularly for mucosal applications. We investigated surface crosslinking as a means to achieve the latter. We found that it increased the colloidal stability of VLPs at low pH (**S6**) as it protected them against proteases. Specifically, while the native AP205 VLP disintegrated in pig gastric fluid at pH 3.0 and 4.7 in less than 5 min, the surface-crosslinked VLPs presented a longer lifetime in the same fluids (**Figure 1** and **S7**). Susceptibility to mechanical agitation was introduced as another limiting factor to VLP mucosal application. Using different force probing techniques, the endocytosis "pulling" force exerted by a cell onto a virus, B cell antigen extraction, and the force of cilia agitation were measured to vary between a few to hundreds of pN (*44-50*). Although this force range is generally below the yield force values of VLPs, when applied repeatedly as they do *in vivo*, they can disintegrate the capsids due to fatigue (*16-18, 22-25*). Therefore, it is also logical to mechanically reinforce VLPs and surface-crosslinking was found to increase the stiffness and strength of AP205 VLP (**Figure 2** and **Figure 4**).

Mucosal applications of VLPs are faced with other physiological challenges, namely translocation through a thick mucus layer. Particularly in the airways, the mucus is subject to mucociliary clearance which further reduces VLP translocation efficiency by pushing the mucus out of the airways (*56*). Experiments using *in vitro* 3D human nasal epithelial tissues showed an effective translocation of native and surface-crosslinked AP205 VLPs, while mucociliary clearance transported mucus at a speed of about 40–200 µm/s (measured at 1–3 mm from the tissue center, (*62, 65*)) in a swirling rotation to the edges of the tissue (**Figure 5**). Is mucus a poor barrier to VLPs of a size below about 100 nm, including AP205 VLP (27.6 nm), and VLPs of human papilloma virus and Norwalk virus (respectively, 55 nm and 38 nm, (*66*))? Considering viruses that infect mucosal surfaces, including polio (28 nm), hepatitis B (43 nm), adenoviruses (60–90 nm), rotavirus (75 nm), human immunodeficiency virus (120 nm) and herpes simplex virus (180 nm) (*67*), one may conclude that small viruses are capable of deeply penetrating the mucus. This is expected when the pore size of mucus gel (≥ 50 nm) exceeds the size of these viruses. However, penetration is further facilitated by the patchy surface charge distribution on

capsids (*66, 67*). Therefore, structural point mutations as well as macromolecule
entanglement with mucin glycoproteins can alter the diffusion of VLPs in mucus (*68*).

Finally, our results suggest that the material properties of VLPs are relevant variables that
should be further investigated to improve mucosal applications. As schematically shown
in **Figure 1**(a), both chemical, enzymatic, and physical stresses can affect the efficacy of
VLP delivery to the respiratory or intestinal mucosal epithelium. In this work, crosslinking
of multiple surface sites was shown to be a promising approach to improve and extend
the range of VLP mucosal functions by increasing resistance to degradation and improving
physical stability. These modifications actually slightly enhance even parenteral
vaccination indicating that the tethers did not shield the coat proteins from immune
recognition, and that increase biochemical stability may also be beneficial systemically
(**Figure 6**). PEG immunogenicity is currently an unwanted side-effect that may be
overcome with use of alternative crosslinkers in the future. Critically, our work
demonstrates the potential to improve the current VLP system to generate structures
compatible with overall enhanced functionality, strength, stability and immunogenicity at
mucosal surfaces.

# Materials and Methods

### Experimental Design

Surface crosslinking of virus-like particle (VLP) capsids using polyethylene glycol (PEG)
tethers was investigated against physiological challenges to assess its performance on
mucosal surfaces. The model native VLP was derived from the coat protein of
Acinetobacter phage, namely AP205 VLP. PEG tethers included two-end functional PEG
molecules of various molecular weights. The PEG-crosslinked AP205 VLPs were initially
evaluated in terms of the reaction products and the sites of PEG conjugation to reactive
surface sites (using a combination of dynamic light scattering, transmission electron
microscopy, reducing SDS-polyacrylamide gel electrophoresis, electrospray ionization
mass spectroscopy and nano ultra-performance liquid chromatography-MS) and then
investigated for the aggregation behavior against pH (using dynamic light scattering),
disintegration against digestive enzymes from pig and mouse gastric fluid (using agarose
gel electrophoresis), resistance to mechanical force (using atomic force microscopy), and

penetration in mucus in the presence of motile cilia and mucociliary clearance (using *in*

*vitro* 3D human nasal tissues).

**424    Production of AP205 VLP**

The biosynthesis of AP205 VLP is reported in details in our previous study (*25*). Briefly, it

includes the expression of recombinant AP205 coat proteins in bacterial cytoplasm which

then self-assemble into AP205 VLPs. After the expression of VLP, a series of steps

consisting of bacterial lysis, digestion of RNA and DNA contaminants, and extraction of

lipids were performed prior to precipitation and dialysis against PBS to produce purified

AP205 VLPs. Plasmids were generated by custom DNA synthesis (Twist Bioscience, San

Francisco, CA, USA) of the expression cassettes and cloning into the pRSFDuet-1

(Novagen) backbone between the PfoI and BshTI restriction sites. Sequences of the

plasmids are available via the ETH-Zürich research collection at

https://doi.org/10.3929/ethz-b-000556182. The sequence of AP205 monomer and the

description of plasmid are reported in **S1**.

**436    PEGylation of AP205 VLP**

The functional group on the extremities of homobifunctional PEG crosslinker molecules

(bPEG$_n$, n = 5 to 25) and on one extremity of monofunctional PEG molecules (mPEG$_n$, n

= 12 and 25) was NHS-ester which is reactive to primary amine of lysine residues and N-

terminus of AP205 coat proteins. Reactions with native AP205 VLP proceeded at 1–100x

molar ratio of crosslinker molecule to AP205 monomer (

$molar\ ratio = [functional\ moleucle]/[AP205\ CP]$ ). The PEG molecule was dissolved in

anhydrous and molecular sieves dried dimethyl sulfoxide (DMSO, Sigma-Aldrich) to a

concentration of 100 mM. An appropriate amount was then added to AP205 VLP

suspension at 4 mg/ml in PBS. The reaction was left for 24 h at 4°C. Then after, it was

quenched by adding Tris•HCl (Trizma® base, Sigma-Aldrich, and hydrochloric acid 37%,

VWR) at 20–50 mM and incubating for 15 min. Unreacted PEG molecules and Tris were

removed using Zeba Spin Desalting Column (ThermoFisher) with molecular weight cut-off

at 7 kDa. Other reactive molecules including BS[3] and Sulfo-NHS were used similarly;

however, they were initially dissolved to 100 mM in PBS. A summary of the physical

properties of these molecules is provided in **S1**.

**452    Fluorescent staining of VLPs**

We defined a mixing ratio $x = \text{mass of stain}/\text{mass of VLP} = 10^{-2}$, and a final stained VLP

concentration equal to 5.3 mg/ml. (To calculate the mass of stain, we assumed 10,000x

to correspond to 10 mg/ml.) GelRed® (10,000x in water, biotium) or SYBRGold™ (10,000x

in DMSO) were used and staining proceeded for 30 min. To remove unbound dye, the

solution of stained VLP was buffer exchanged with pure PBS using Zeba™ Spin Desalting

Columns (7,000 MWCO, ThermoFisher). We note that since the columns contained

sodium azide, it was important to buffer the column with pure PBS prior to cleaning the

stained VLP solution. The stained VLP solution was finally filtered using 0.2 µm centrifuge

filters (Corning® Costar® Spin-X® centrifuge tube filters) to remove possible aggregates.

The intensity of stained VLPs was monitored over time using laser scanning confocal

microscopy and showed no noticeable reduction over a course of at least two weeks.

**Transmission electron microscopy (TEM)**

4 µl VLP solution (0.1 mg/ml) was deposited on glow-discharged (45 sec, 25 mA, negative

charge) carbon-coated grids (Ted Pella) for 1-5 min, after which the excess liquid was

removed using filter paper. Then the grid was rinsed with 5 µl of PBS to remove loosely

bound VLPs. Subsequently, 3 µl of 2% uranyl acetate was applied and immediately

removed with filter paper prior to a second treatment with 3 µl of 2% uranyl acetate for 20

sec. Excess liquid was again collected using filter paper and the grid was let dry at room

temperature (RT). Data acquisition was performed using a Morgani transmission electron

microscope operating at 100 kV acceleration voltage.

**Electrospray ionization mass spectrometry (ESI-MS)**

ESI-MS was acquired on a Synapt G2-Si quadrupole time-of-flight spectrometer (Waters,

UK) with a capillary voltage of 3 V, a cone voltage of 50 V and a source temperature of

100°C. VLPs at a concentration of 1 mg/ml were reduced by dithiothreitol (DTT) at a final

concentration of 50 mM. The reduction was performed for 1 h at RT and at pH 8. The

samples were acidified with 1% formic acid (Thermo, USA), desalted using C4 Zip Tips

(Millipore, USA) and analyzed in methanol:2-propanol:0.2 % formic acid (30:20:50). The

solutions were infused through a fused silica capillary (ID 75 µm) at a flow rate of 1 µl/min

and sprayed through Pico Tips (ID 30 µm). The last were obtained from New Objective

(Woburn, MA, USA). Recorded m/z data were deconvoluted using the MaxEnt1 software

(Waters, UK) with a resolution of the output mass of 0.5 Da per channel and a Uniform

Gaussian Damage Model at the half-height of 0.5 Da.

**Nano ultra-performance liquid chromatography-MS (nanoUPLC-MS/MS)**

NanoUPLC-MS/MS was acquired on Thermo FUSION mass spectrometer (Thermo,

Germany) coupled to a nano-M-class UPLC system (Waters). BPEG$_9$-AP205 VLP at a

concentration of 1 mg/ml was digested by incubating 3 µl of the sample with 42 µl of 25

mM ammonium bicarbonate (pH 8.5) and 1 µl of TCEP (100 mM) for 1 h at 37°C. Reduced

protein was incubated with 2 µl chloroacetamide (500 mM) for 1 h at 37°C. Then after, 5

µl trypsin (0.1 µg/µl in 10 mM HCl) was added and the solution further incubated overnight

at 37°C. When needed the pH was adjusted to pH 8. At the end of this procedure, the

sample was dried using SpeedVac. Digested and dried sample was dissolved in 20 µl

ddH$_2$O with 0.1% formic acid and diluted 5 times before transferring to autosampler vials.

Peptides were resuspended in 2.5% acetonitrile with 0.1% formic acid and loaded onto a

nanoEase M/Z Symmetry C18 (75 µm × 20 mm, 100 Å, 3 µm particle size) and separated

on a nanoEase M/Z HSS C18 T3 (75 µm × 150 mm, 130 Å, 1.7 µm particle size) at a

constant flow rate of 300 nl/min, with a column temperature of 50°C and a linear gradient

of 2−32% acetonitrile/0.1% formic acid in 79 min, and then 32-45% acetonitrile/0.1%

formic acid in 10 min, followed by a sharp increase to 98% acetonitrile in 2 min and then

held at 98% for another 10 min. Mass spectrometer was operated under data-dependent

acquisition (DDA), one scan cycle comprised of a full scan MS survey spectrum, followed

by up to 12 sequential higher-collisional energy (HCD) MS/MS on the most intense signals

above a threshold of 1e4. Full-scan MS spectra (600–2000 m/z) were acquired in the FT-

Orbitrap at a resolution of 70,000 at 400 m/z, while HCD MS/MS spectra were recorded

in the FT-Orbitrap at a resolution of 35,000 at 400 m/z. HCD was performed with a target

value of 1e5 and normalization collision energy 25 NCE (normalize collisional energy) was

applied. Auto-gain control (AGC) target values were 5e5 for full Fourier transform (FT)

MS. For all experiments, dynamic exclusion was used with a single repeat count, 15 s

repeat duration, and 30 s exclusion duration. The acquired MS data were processed for

identification using the Byonic search engine (PMI). The spectra were searched against

AP205 VLP sequence and AP205 VLP sequence with E. *Coli* database. The following

modifications were included: (a) variable modifications, namely oxidation at N-terminus

methionine and carbamidomethylation at cysteine (C), and (b) user-defined modifications,

namely 478.2565 (C$_{22}$O$_{11}$H$_{38}$) was set for PEG$_9$-crosslinked peptides and 496.2565

(C$_{22}$O$_{12}$H$_{40}$) for one-end PEG$_9$ conjugation at a lysine residue or N-Terminus. In this

analysis, the raw data were analyzed by two software, PMI Byonic search engine and SIM

1.5.5.3 (Spectrum identification machine) (*69*). In addition, two FastA databases with and

without starting methionine were used to confirm the N-terminus.

## SDS-polyacrylamide gel electrophoresis (SDS-PAGE)

521    Denaturing and reducing SDS-PAGE was carried out in 13% Tricine gels. Samples of 3

522    µg of VLP were denatured and reduced in Laemmli sample buffer at 95°C for 5 min prior

523    to electrophoresis. Proteins were visualized by Coomassie staining.

## Dynamic light scattering (DLS)

525    DLS was performed at a fixed angle of 173° by averaging 3 runs of 30-s long each using

526    Zetasizer Nano (Malvern Panalytical). The time correlation function of scattered intensity

527    was analyzed by the cumulant as well as CONTIN methods. The VLP concentration was

528    initially adjusted to 0.1 mg/ml in filtered PBS prior to measurements.

## Gastric fluid collection

Pig gastric content was collected from the stomach of conventionally reared, outbred pigs

with different genetic contributions of French-Swiss Landrace and Swiss Large White.

Total gastric content was collected. Mouse gastric content was obtained by collecting total

stomach content from several specific opportunistic pathogen-free (SPF) C57BL/6J WT

mice. The gastric content from several mice was pooled to have sufficient volume. Pig or

mouse gastric content was centrifuged for 5 min at 16000 rpm to separate the non-

digested food particles. After the collection of supernatant (gastric fluid), it was filtered

using 0.22 µm sterile filter and the pH measured using a laboratory pH-meter.

## Agarose gel electrophoresis

VLPs from 1 mg/ml preparation were diluted 1:10 in pig or mouse gastric fluid and

incubated for 5 min, 30 min, 1 h and 1.5 or 2 h at 37°C. After each incubation, the samples

were immediately placed on ice. Native agarose gel electrophoresis of PBS- or gastric

fluid-treated VLPs was carried out in 0.6% agarose in TAE buffer (40 mM Tris, 0.1% acetic

acid, 1 mM EDTA) with 0.5 µg/ml of ethidium bromide for visualization. Samples of 5 µg

VLP were electrophoresed at 120 Volts for 1 h, then imaged on a UV transilluminator.

## Atomic force microscopy (AFM)

The schematic of AFM measurements is shown in **Figure 2**(a). After an initial calibration,

which also included the calibration of PeakForce tapping mode, on freshly cleaved mica,

the tip was brought in proximity to VLPs immobilized at a concentration of 5 µg/ml on

Highly Oriented Pyrolytic Graphite (HOPG, Ted Pella, Inc.) The calibrated parameters

included stiffness (59.4–82.0 pN/nm) using thermal method and optical lever sensitivity

using cantilever–mica contact region (*70*), and drive3 amplitude sensitivity. During the

transport of cantilever from mica to HOPG, the cantilever remained wet, and the laser spot

was checked to remain on the same spot. The calibration values were re-measured on

mica at the end of the measurements to detect any noticeable variation to the calibrated

parameters. Single VLPs were localized on HOPG by imaging in PeakForce tapping mode

(peak force 50 pN, tapping frequency 1 kHz, amplitude 30 nm). Generally, images of

individual VLPs were acquired at a scan rate of 0.7 Hz with 200×200 or 500×500 nm$^2$ scan

size and a resolution of 64 points per line, leading to a pixel size of 3 nm. After localization,

we used "Point & Shoot" function (NanoScope, Bruker) to place the AFM tip on the apex

of selected VLPs before applying indentation using Ramp mode. The VLPs were indented

up to a threshold force value of 400 pN at a preset frequency of 1 Hz, which resulted in an

indentation force rate of about (4–9)×10$^3$ pN/s. Indented particles were imaged again in

PeakForce tapping to detect signs of degradation, drift, or displacement due to tip lateral

forces. The height of individual VLPs before ($H_i$) and after ($H_f$) indentation were

calculated by placing a trace line over the particles on images previously flattened using

Gwyddion 2.47 (*71*). For stiffness evaluation, we proceeded as follows: deflection versus

piezo displacement was converted to force versus tip separation using protocols written

in Igor Pro (Wavemetrics). Fit to force versus indentation distance ($F-\delta$) curve in the

linear response part of the tip–VLP contact region gave the stiffness ($k = dF/d\delta$) and the

indentation ($\delta = \Delta F/k$). The fit was performed in the force range $50\,\mathrm{pN} \leq F \leq 250\,\mathrm{pN}$, and

only those VLPs with indentation range $2.0 \leq \delta \leq 6.0\,\mathrm{nm}$ were selected for averaging. Only

one indentation per VLP was used for the evaluations. The indentation force rate ($\dot{F}$) was

calculated from a linear fit to the force versus indentation time ($F-t$) curve in the same

region were the stiffness of VLP was calculated. Strain rate was then calculated from the

relation $\dot{\gamma} = \dot{F}/k\delta$. Average stiffness was calculated from at least duplicate independent

VLP preparations. UV-ozone (Novascan) cleaned AC40 BioLever Mini cantilevers

(Bruker) were used. Dimension FastScan (Bruker) was used for imaging and force

spectroscopy.

**Small angle X-ray scattering (SAXS)**

We used a Xeuss 2.0 (Xenocs, France) instrument with a microfocused X-ray source at

the Laboratory Léon Brillouin (LLB NIMBE CEA Saclay). The Cu Kα radiation ($\lambda_{Cu\ K\alpha}$ =

1.5418 Å), and the data were collected by a 2D Pilatus 1 M detector (Dectris, Switzerland).

The scattering vector $q = 4\pi/\lambda \sin(\theta/2)$, with $\theta$ being the scattering angle was calibrated

using silver behenate. Data were collected and azimuthally averaged to produce

unidimensional intensity versus $q$, with $q$ interval of 0.005 to 0.1 Å$^{-1}$. The measurements

were performed at 25°C. The scattering intensity was collected for 2600 s. The intensity

was corrected for sample thickness, transmission, and acquisition time. The modelling of

the integrated data was performed with a core-shell form factor, and a hard sphere

structure factor, when needed, using the SasView software.

***In vitro* 3D human nasal epithelial tissue**

*In vitro* human nasal epithelial tissues from single donors (MucilAir™) were obtained from

Epithelix (Switzerland). The tissues were maintained according to company's protocols

and included tissue 1 from a Caucasian 64-year-old female subject (MD0860), tissue 2

from a Caucasian 46-year-old male subject (MD0871) and tissue 3 from a Caucasian 38-

year-old male subject (MD0774). The subjects were non-smokers and had no history of

respiratory pathology.

**Inhibition of mucociliary clearance**

Ciliary beating was inhibited using trans-cinnamaldehyde (97%, C80687, Sigma-Aldrich).

Trans-cinnamaldehyde was dissolved in MucilAir™ culture medium to a concentration of

30 mM. To improve solubility, freshly prepared solution was sonicated for 15 min and the

solution was kept at 37°C prior to filling the basal channel of MucilAir™ monodonor nasal

epithelial tissue. After 20 min of incubation, the ciliary beating stopped, and no recovery

was observed for up to 2 h (*72*).

**Delivery of VLPs to *in vitro* 3D human nasal epithelial tissue**

Tissues were incubated in a humidified-$CO_2$ chamber at 37°C. After a period of

equilibration 0.5–1 h, 5 µl of VLP solution (5.3 mg/ml) was delivered from the apical side.

The delivery was onto the center of the tissue.

**Laser scanning confocal microscopy (LSCM)**

Lateral distribution and vertical translocation of fluorescently labelled VLPs through mucus

were then monitored using laser scanning confocal microscopy (LSCM, Leica SP5

confocal microscope). We used a 10x objective lens with N.A. 0.3. Excitation wavelength

$\lambda_{ex} = 488\,\text{nm}$ was used for SYBRGold$^{TM}$- stained VLPs and Salmonella vaccine with

detection wavelength $\lambda_{em} = 500 - 590\,\text{nm}$. $\lambda_{ex} = 514\,\text{nm}$ was used for GelRed$^{\circledR}$-stained

VLPs with $\lambda_{em} = 550 - 650\,\text{nm}$. The channel spectral detection was set to HyD (Gain equal

to 100). The mean fluorescence intensity (MFI) variation with height was collected in the

center [0, 0] mm, and on positions [-1, 0] mm, [+1, 0] mm, [0, -1] mm and [0, +1] mm prior

to, during and after VLP delivery. Leica LAS AF software was used for saving the images.

Analysis was performed using ImageJ.

**Mice**

All animal experiments were performed in accordance with Swiss Federal regulations

approved by the Commission for Animal Experimentation of the Kanton Zurich (license

ZH009/2021, Kantonales Veterinäramt Zürich, Switzerland). SPF C57BL/6J WT mice

were used in all experiments. Mice were bred and housed in individually ventilated cages

with a 12 h light/dark cycle in the ETH Phenomics Center (EPIC, RCHCI), ETH Zürich,

and were fed a standard chow diet. All mice included in experiments were 7 weeks or

older and objectively healthy as determined by routine health checks. Wherever possible

an equal number of males and females was used in each experimental group.

**Subcutaneous injection**

629    50 µg of VLP in 100 µl of PBS was subcutaneously injected into the loose skin over the

630    interscapular area of C57BL/6J SPF mice on Day 0 and Day 10. The mice were

631    euthanized on Day 20. Blood was collected by cardiac puncture, and serum separated by

632    spinning 1.1 ml serum gel tubes (Sarstedt) at 10,000 g for 5 min at 20°C in a conventional

633    table-top centrifuge, and heat-inactivated for 30 min at 56°C.

**Enzyme-linked immunosorbent assay (ELISA) for detection of anti-AP205 IgG**

634
**response**

50 µl of AP205 VLP solution (5 µg/ml) was pipetted into an appropriate number of wells of

ELISA plates (4 Nunc MaxiSorp flat bottom 96-well plates) and the VLP allowed to adsorb

overnight at 4°C. The wells were washed three times with wash buffer (0.05% Tween 20

in PBS) and blotted. The wells were blocked using 150 µl of blocking buffer (2% bovine

serum albumin in PBS) for 1.5 h at RT. Subsequently, the wells were washed five times

with wash buffer and blotted. Mouse sera were diluted ten times in blocking buffer, and

sequentially in 1:3 dilution steps added to the wells of ELISA plates. The plates were

incubated for 2 h at RT to allow the binding of serum IgG to AP205 proteins. Goat anti-

mouse IgG-HRP (Sigma) was diluted a thousand-fold in blocking buffer, and 50 µl of

solution added to the wells and incubated for 1 h at RT. The wells were washed five times

using wash buffer and blotted. Finally, 150 µl of freshly prepared substrate solution

(consisting of 0.1 M $NaH_2PO_4$ (substrate buffer), 2,2'-azino-bis(3-ethylbenzothiazoline-6-

sulfonic acid (1–2 mg per 10 ml of substrate buffer), and $H_2O_2$ (10 µl per 10 ml)) was added

to each well. The plate was incubated in the dark for 40 min. On Tecan (Tecan Infinite 200

PRO microplate reader), absorbance was measured at 415 nm. The reference wavelength

for the subtraction of background signal was set to 480 nm.

**Enzyme-linked immunosorbent assay (ELISA) for detection of anti-PEG IgG**

**response**

Mouse sera were diluted ten times in PEG dilution buffer (PEGD50-1, Life Diagnostics,

Inc.), and sequentially in 1:3 dilution steps added to the wells of PEG-BSA coated ELISA

plates (PBSA20PL, Life Diagnostics, Inc.) The plates were incubated for 2 h at RT to allow

antibody binding. The wells were washed five times with 300 µl PEG wash buffer

(PEGW50-20, Life Diagnostics, Inc.) and blotted. Goat anti-mouse IgG-HRP (AP124P,

Sigma-Aldrich) was diluted a thousand-fold in PEG dilution buffer and 100 µl of this

solution was added to each well with an incubation of 1 h at RT. The wells were washed

and blotted again. Finally, 150 µl of freshly prepared substrate solution was added to each

well. The plates were incubated in the dark for 40 min. On Tecan (Tecan Infinite 200 PRO

microplate reader), absorbance was measured at 415 nm. The reference wavelength for

the subtraction of background signal was set to 480 nm.

**Statistical Analysis**

AFM and LSCM results were expressed as mean ± standard deviation analyzed by Igor

Pro from WaveMetrics. *In vivo* mouse vaccination results were assessed using ordinary

one-way ANOVA with Dunnett's multiple comparisons test using GraphPad Prism

(*P = 0.0332; ***P = 0.0021; ***P = 0.0002).

# Acknowledgments

We acknowledge the contributions from Till Germerdonk in DLS measurements, and discussions with Patrick Zueblin. We also acknowledge the time for SAXS measurements provided by the Leon Brillouin Laboratory (LLB), and the support of Fabrice Cousin.

**Funding**

European Research Council (ERC) Grant number 865730 (MR, YT, AA and ES)

Swiss National Science Foundation, Switzerland, rant number IZSEZ0_212991 (MR)

Swiss National Science Foundation, Switzerland, grant numbers 40B2-0_180953 (ES)

Swiss National Science Foundation, Switzerland, grant numbers 310030_185128 (ES)

National Center for Competence in Research in Microbiomes, Switzerland (ES)

Botnar Research Centre for Child Health Multi-investigator project "Microbiota Engineering for Child Health" (ES)

Gebert Rüf Microbials, grant number GR073_17 (ES)

Cystic Fibrosis Trust UK SRC 016  (PC, JK, RF)

EU ITN PhyMot (EC)

**Author contributions**

Conceptualization: MR

Methodology: MR, PC, RM, ES

Investigation: MR, AA, SG, VL, TK, YT, WLB, EC, SC, CWL, JK

Writing—original draft: MR

Writing—review & editing: All authors

Funding acquisition: MR, PC, ES

**Competing interests**

Authors declare that they have no competing interests.

**Data and materials availability**

All data used in the analyses are available without limitation from the first author.

# Supplementary Materials

Further information regarding the Materials and Methods are provided in Supplementary Materials (**S**).

# References

1.  Y. H. Chung, H. Cai, N. F. Steinmetz, Viral nanoparticles for drug delivery, imaging, immunotherapy, and theranostic applications. *Advanced Drug Delivery Reviews* **156**, 214-235 (2020).
2.  H. Tariq, S. Batool, S. Asif, M. Ali, B. H. Abbasi, Virus-Like Particles: Revolutionary Platforms for Developing Vaccines Against Emerging Infectious Diseases. *Frontiers in Microbiology* **12**, (2022).
3.  H. Hu, N. F. Steinmetz, Development of a Virus-Like Particle-Based Anti-HER2 Breast Cancer Vaccine. *Cancers* **13**, (2021).
4.  D. Yin, S. Ling, D. Wang, Y. Dai, H. Jiang, X. Zhou, S. R. Paludan, J. Hong, Y. Cai, Targeting herpes simplex virus with CRISPR-Cas9 cures herpetic stromal keratitis in mice. *Nat Biotechnol* **39**, 567-577 (2021).
5.  S. Banskota, A. Raguram, S. Suh, S. W. Du, J. R. Davis, E. H. Choi, X. Wang, S. C. Nielsen, G. A. Newby, P. B. Randolph, M. J. Osborn, K. Musunuru, K. Palczewski, D. R. Liu, Engineered virus-like particles for efficient in vivo delivery of therapeutic proteins. *Cell* **185**, 250-265.e216 (2022).
6.  J. Mianné, A. Nasri, C. N. Van, C. Bourguignon, M. Fieldès, E. Ahmed, C. Duthoit, N. Martin, H. Parrinello, A. Louis, A. Iché, R. Gayon, F. Samain, L. Lamouroux, P. Bouillé, A. Bourdin, S. Assou, J. De Vos, CRISPR/Cas9-mediated gene knockout and interallelic gene conversion in human induced pluripotent stem cells using non-integrative bacteriophage-chimeric retrovirus-like particles. *BMC Biology* **20**, 8 (2022).
7.  C. Sakai, K. Hosokawa, T. Watanabe, Y. Suzuki, T. Nakano, K. Ueda, M. Fujimuro, Human hepatitis B virus-derived virus-like particle as a drug and DNA delivery carrier. *Biochemical and Biophysical Research Communications* **581**, 103-109 (2021).
8.  E. C. Thuenemann, D. H. T. Le, G. P. Lomonossoff, N. F. Steinmetz, Bluetongue Virus Particles as Nanoreactors for Enzyme Delivery and Cancer Therapy. *Mol Pharm* **18**, 1150-1156 (2021).
9.  S. Panthi, P. T. Schmitt, F. J. Lorenz, B. A. Stanfield, A. P. Schmitt, Paramyxovirus-Like Particles as Protein Delivery Vehicles. *J Virol* **95**, e0103021 (2021).
10. A. Shahrivarkevishahi, M. A. Luzuriaga, F. C. Herbert, A. C. Tumac, O. R. Brohlin, Y. H. Wijesundara, A. V. Adlooru, C. Benjamin, H. Lee, P. Parsamian, J. Gadhvi, N. J. De Nisco, J. J. Gassensmith, PhotothermalPhage: A Virus-Based Photothermal Therapeutic Agent. *J Am Chem Soc* **143**, 16428-16438 (2021).
11. H. Lee, A. Shahrivarkevishahi, J. L. Lumata, M. A. Luzuriaga, L. M. Hagge, C. E. Benjamin, O. R. Brohlin, C. R. Parish, H. R. Firouzi, S. O. Nielsen, L. L. Lumata, J. J. Gassensmith, Supramolecular and biomacromolecular enhancement of metal-free magnetic resonance imaging contrast agents. *Chemical Science* **11**, 2045-2050 (2020).
12. A. Armanious, M. Radiom, R. Mezzenga, Recent Experimental Advances in Probing the Colloidal Properties of Viruses. *Current Opinion in Colloid & Interface Science*, 101703 (2023).
13. V. Sankar, V. Hearnden, K. Hull, D. V. Juras, M. Greenberg, A. Kerr, P. Lockhart, L. Patton, S. Porter, M. Thornhill, Local drug delivery for oral mucosal diseases: challenges and opportunities. *Oral Diseases* **17**, 73-84 (2011).
14. J. O. Morales, K. R. Fathe, A. Brunaugh, S. Ferrati, S. Li, M. Montenegro-Nicolini, Z. Mousavikhamene, J. T. McConville, M. R. Prausnitz, H. D. C. Smyth, Challenges and Future Prospects for the Delivery of Biologics: Oral Mucosal, Pulmonary, and Transdermal Routes. *The AAPS Journal* **19**, 652-668 (2017).
15. S. Hua, Advances in Oral Drug Delivery for Regional Targeting in the Gastrointestinal Tract - Influence of Physiological, Pathophysiological and Pharmaceutical Factors. *Frontiers in Pharmacology* **11**, (2020).
16. I. L. Ivanovska, P. J. d. Pablo, B. Ibarra, G. Sgalari, F. C. MacKintosh, J. L. Carrascosa, C. F. Schmidt, G. J. L. Wuite, Bacteriophage capsids: Tough nanoshells with complex elastic properties. *Proceedings of the National Academy of Sciences* **101**, 7600-7605 (2004).

17. J. P. Michel, I. L. Ivanovska, M. M. Gibbons, W. S. Klug, C. M. Knobler, G. J. L. Wuite, C. F. Schmidt, Nanoindentation studies of full and empty viral capsids and the effects of capsid protein mutations on elasticity and strength. *Proceedings of the National Academy of Sciences* **103**, 6184-6189 (2006).

18. N. Kol, M. Gladnikoff, D. Barlam, R. Z. Shneck, A. Rein, I. Rousso, Mechanical Properties of Murine Leukemia Virus Particles: Effect of Maturation. *Biophysical Journal* **91**, 767-774 (2006).

19. C. Carrasco, A. Carreira, I. A. T. Schaap, P. A. Serena, J. Gómez-Herrero, M. G. Mateu, P. J. d. Pablo, DNA-mediated anisotropic mechanical reinforcement of a virus. *Proceedings of the National Academy of Sciences* **103**, 13706-13711 (2006).

20. N. Kol, Y. Shi, M. Tsvitov, D. Barlam, R. Z. Shneck, M. S. Kay, I. Rousso, A Stiffness Switch in Human Immunodeficiency Virus. *Biophysical Journal* **92**, 1777-1783 (2007).

21. I. Ivanovska, G. Wuite, B. Jönsson, A. Evilevitch, Internal DNA pressure modifies stability of WT phage. *Proceedings of the National Academy of Sciences* **104**, 9603-9608 (2007).

22. A. Llauró, D. Luque, E. Edwards, B. L. Trus, J. Avera, D. Reguera, T. Douglas, P. J. d. Pablo, J. R. Castón, Cargo–shell and cargo–cargo couplings govern the mechanics of artificially loaded virus-derived cages. *Nanoscale* **8**, 9328-9336 (2016).

23. A. Ortega-Esteban, G. N. Condezo, A. J. Pérez-Berná, M. Chillón, S. J. Flint, D. Reguera, C. San Martín, P. J. de Pablo, Mechanics of Viral Chromatin Reveals the Pressurization of Human Adenovirus. *ACS Nano* **9**, 10826-10833 (2015).

24. M. Jiménez-Zaragoza, M. P. L. Yubero, E. Martín-Forero, J. R. Castón, D. Reguera, D. Luque, P. J. de Pablo, J. M. Rodríguez, Biophysical properties of single rotavirus particles account for the functions of protein shells in a multilayered virus. *eLife* **7**, e37295 (2018).

25. M. Radiom, T. Keys, Y. Turgay, A. Ali, S. Peret, S. Chesnov, V. Lutz-Bueno, E. Slack, R. Mezzenga, Mechanical tuning of virus-like particles. *Journal of Colloid and Interface Science* **634**, 963-971 (2023).

26. C. Zeng, L. Scott, A. Malyutin, R. Zandi, P. Van der Schoot, B. Dragnea, Virus Mechanics under Molecular Crowding. *The Journal of Physical Chemistry B* **125**, 1790-1798 (2021).

27. M. Cantero, D. Carlero, F. J. Chichón, J. Martín-Benito, P. J. De Pablo, Monitoring SARS-CoV-2 Surrogate TGEV Individual Virions Structure Survival under Harsh Physicochemical Environments. *Cells* **11**, (2022).

28. C. R. O'Riordan, A. Lachapelle, C. Delgado, V. Parkes, S. C. Wadsworth, A. E. Smith, G. E. Francis, PEGylation of adenovirus with retention of infectivity and protection from neutralizing antibody in vitro and in vivo. *Hum Gene Ther* **10**, 1349-1358 (1999).

29. M. A. Croyle, N. Chirmule, Y. Zhang, J. M. Wilson, PEGylation of E1-deleted adenovirus vectors allows significant gene expression on readministration to liver. *Hum Gene Ther* **13**, 1887-1900 (2002).

30. M. A. Croyle, N. Chirmule, Y. Zhang, J. M. Wilson, "Stealth" adenoviruses blunt cell-mediated and humoral immune responses against the virus and allow for significant gene expression upon readministration in the lung. *J Virol* **75**, 4792-4801 (2001).

31. E. A. Weaver, M. A. Barry, Effects of shielding adenoviral vectors with polyethylene glycol on vector-specific and vaccine-mediated immune responses. *Hum Gene Ther* **19**, 1369-1382 (2008).

32. T. Shimizu, M. Ichihara, Y. Yoshioka, T. Ishida, S. Nakagawa, H. Kiwada, Intravenous administration of polyethylene glycol-coated (PEGylated) proteins and PEGylated adenovirus elicits an anti-PEG immunoglobulin M response. *Biol Pharm Bull* **35**, 1336-1342 (2012).

33. K. L. Lee, S. Shukla, M. Wu, N. R. Ayat, C. E. El Sanadi, A. M. Wen, J. F. Edelbrock, J. K. Pokorski, U. Commandeur, G. R. Dubyak, N. F. Steinmetz, Stealth filaments: Polymer chain length and conformation affect the in vivo fate of PEGylated potato virus X. *Acta Biomater* **19**, 166-179 (2015).

34. Y. Eto, Y. Yoshioka, T. Ishida, X. Yao, T. Morishige, S. Narimatsu, H. Mizuguchi, Y. Mukai, N. Okada, H. Kiwada, S. Nakagawa, Optimized PEGylated adenovirus vector reduces the anti-vector humoral immune response against adenovirus and induces a therapeutic effect against metastatic lung cancer. *Biol Pharm Bull* **33**, 1540-1544 (2010).

35. P. W. Lee, S. A. Isarov, J. D. Wallat, S. K. Molugu, S. Shukla, J. E. Sun, J. Zhang, Y. Zheng, M. Lucius Dougherty, D. Konkolewicz, P. L. Stewart, N. F. Steinmetz, M. J. Hore, J. K. Pokorski, Polymer Structure and Conformation Alter the Antigenicity of Virus-like Particle-Polymer Conjugates. *J Am Chem Soc* **139**, 3312-3315 (2017).

36. S. N. Crooke, J. Zheng, M. S. Ganewatta, S. M. Guldberg, T. M. Reineke, M. G. Finn, Immunological Properties of Protein–Polymer Nanoparticles. *ACS Applied Bio Materials* **2**, 93-103 (2019).

37. J. D. Lewis, G. Destito, A. Zijlstra, M. J. Gonzalez, J. P. Quigley, M. Manchester, H. Stuhlmann, Viral nanoparticles as tools for intravital vascular imaging. *Nature Medicine* **12**, 354-360 (2006).

38. G. Destito, R. Yeh, C. S. Rae, M. G. Finn, M. Manchester, Folic Acid-Mediated Targeting of Cowpea Mosaic Virus Particles to Tumor Cells. *Chemistry & Biology* **14**, 1152-1162 (2007).

39. A. C. Tissot, R. Renhofa, N. Schmitz, I. Cielens, E. Meijerink, V. Ose, G. T. Jennings, P. Saudan, P. Pumpens, M. F. Bachmann, Versatile Virus-Like Particle Carrier for Epitope Based Vaccines. *PLOS ONE* **5**, e9809 (2010).

40. X. Liu, X. Chang, D. Rothen, M. Derveni, P. Krenger, S. Roongta, E. Wright, M. Vogel, K. Tars, M. O. Mohsen, M. F. Bachmann, AP205 VLPs Based on Dimerized Capsid Proteins Accommodate RBM Domain of SARS-CoV-2 and Serve as an Attractive Vaccine Candidate. *Vaccines* **9**, 403 (2021).

41. M. Shishovs, J. Rumnieks, C. Diebolder, K. Jaudzems, L. B. Andreas, J. Stanek, A. Kazaks, S. Kotelovica, I. Akopjana, G. Pintacuda, R. I. Koning, K. Tars, Structure of AP205 Coat Protein Reveals Circular Permutation in ssRNA Bacteriophages. *Journal of Molecular Biology* **428**, 4267-4279 (2016).

42. W. Humphrey, A. Dalke, K. Schulten, VMD: Visual molecular dynamics (http://www.ks.uiuc.edu/Research/vmd/). *Journal of Molecular Graphics* **14**, 33-38 (1996).

43. D. Thirumalai, B. Y. Ha, "Statistical mechanics of semiflexible chains: pA mean field variational approach" in *Theoretical and mathematical models in polymer research,* A. Grosberg, Ed. (Academic Press: San Diego, 1998).

44. Y. Pan, F. Zhang, L. Zhang, S. Liu, M. Cai, Y. Shan, X. Wang, H. Wang, H. Wang, The Process of Wrapping Virus Revealed by a Force Tracing Technique and Simulations. *Advanced Science* **4**, 1600489 (2017).

45. T. Wiegand, M. Fratini, F. Frey, K. Yserentant, Y. Liu, E. Weber, K. Galior, J. Ohmes, F. Braun, D.-P. Herten, S. Boulant, U. S. Schwarz, K. Salaita, E. A. Cavalcanti-Adam, J. P. Spatz, Forces during cellular uptake of viruses and nanoparticles at the ventral side. *Nature Communications* **11**, 32 (2020).

46. Q. Zhang, F. Tian, F. Wang, Z. Guo, M. Cai, H. Xu, H. Wang, G. Yang, X. Shi, Y. Shan, Z. Cui, Entry Dynamics of Single Ebola Virus Revealed by Force Tracing. *ACS Nano* **14**, 7046-7054 (2020).

47. J. Zhang, Y. Huang, M. Sun, T. Song, S. Wan, C. Yang, Y. Song, Mechanosensing view of SARS-CoV-2 infection by a DNA nano-assembly. *Cell Rep Phys Sci* **3**, 101048 (2022).

48. S. Li, Y. Pan, H. Teng, Y. Shan, G. Yang, H. Wang, Revealing the Cell Entry Dynamic Mechanism of Single Rabies Virus Particle. *Chem Res Chin Univ* **38**, 838-842 (2022).

49. K. M. Spillane, P. Tolar, B cell antigen extraction is regulated by physical properties of antigen-presenting cells. *Journal of Cell Biology* **216**, 217-230 (2017).

50. D. B. Hill, V. Swaminathan, A. Estes, J. Cribb, E. T. O'Brien, C. W. Davis, R. Superfine, Force generation and dynamics of individual cilia under external loading. *Biophys J* **98**, 57-66 (2010).

51. M. P. Neubauer, M. Poehlmann, A. Fery, Microcapsule mechanics: From stability to function. *Advances in Colloid and Interface Science* **207**, 65-80 (2014).

52. M. Buenemann, P. Lenz, Elastic properties and mechanical stability of chiral and filled viral capsids. *Physical Review E* **78**, 051924 (2008).

53. L. Menou, Y. C. Salas, L. Lecoq, A. Salvetti, C. F. Moskalenko, M. Castelnovo, Stiffness heterogeneity of small viral capsids. *Physical Review E* **104**, 064408 (2021).

54. M. Buenemann, P. Lenz, Mechanical limits of viral capsids. *Proceedings of the National Academy of Sciences* **104**, 9925-9930 (2007).

55. Z. L. Luo, B. Zhang, H. J. Qian, Z. Y. Lu, S. X. Cui, Effect of the size of solvent molecules on the single-chain mechanics of poly(ethylene glycol): implications on a novel design of a molecular motor. *Nanoscale* **8**, 17820-17827 (2016).

56. R. Bansil, B. S. Turner, The biology of mucus: composition, synthesis and organization. *Advanced Drug Delivery Reviews* **124**, 3-15 (2018).

57. X. M. Bustamante-Marin, L. E. Ostrowski, Cilia and mucociliary clearance. *Cold Spring Harbor perspectives in biology* **9**,  (2017).

58. R. Fradique, E. Causa, C. Delahousse, J. Kotar, L. Pinte, L. Vallier, M. Vila-Gonzalez, P. Cicuta, Assessing motile cilia coverage and beat frequency in mammalian <i>in vitro</i> cell culture tissues. *Royal Society Open Science* **10**, 230185 (2023).

59. Y. Y. Wang, S. K. Lai, J. S. Suk, A. Pace, R. Cone, J. Hanes, Addressing the PEG mucoadhesivity paradox to engineer nanoparticles that "slip" through the human mucus barrier. *Angew Chem Int Ed Engl* **47**, 9726-9729 (2008).

60. C. S. Schneider, Q. Xu, N. J. Boylan, J. Chisholm, B. C. Tang, B. S. Schuster, A. Henning, L. M. Ensign, E. Lee, P. Adstamongkonkul, B. W. Simons, S. S. Wang, X. Gong, T. Yu, M. P. Boyle, J. S. Suk, J. Hanes, Nanoparticles that do not adhere to mucus provide uniform and long-lasting drug delivery to airways following inhalation. *Sci Adv* **3**, e1601556 (2017).

61. J. T. Huckaby, S. K. Lai, PEGylation for enhancing nanoparticle diffusion in mucus. *Advanced Drug Delivery Reviews* **124**, 125-139 (2018).

62. M.-K. Khelloufi, E. Loiseau, M. Jaeger, N. Molinari, P. Chanez, D. Gras, A. Viallat, Spatiotemporal organization of cilia drives multiscale mucus swirls in model human bronchial epithelium. *Scientific Reports* **8**, 2447 (2018).

63. L. Klimek, T. Kündig, M. F. Kramer, S. Guethoff, E. Jensen-Jarolim, C. B. Schmidt-Weber, O. Palomares, M. O. Mohsen, T. Jakob, M. Bachmann, Virus-like particles (VLP) in prophylaxis and immunotherapy of allergic diseases. *Allergo Journal International* **27**, 245-255 (2018).

64. A. Armanious, R. Mezzenga, A Roadmap for Building Waterborne Virus Traps. *JACS Au* **2**, 2205-2221 (2022).

65. H. Matsui, B. R. Grubb, R. Tarran, S. H. Randell, J. T. Gatzy, C. W. Davis, R. C. Boucher, Evidence for periciliary liquid layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways disease. *Cell* **95**, 1005-1015 (1998).

66. S. S. Olmsted, J. L. Padgett, A. I. Yudin, K. J. Whaley, T. R. Moench, R. A. Cone, Diffusion of Macromolecules and Virus-Like Particles in Human Cervical Mucus. *Biophysical Journal* **81**, 1930-1937 (2001).

67. R. A. Cone, Barrier properties of mucus. *Adv Drug Deliv Rev* **61**, 75-85 (2009).

68. G. A. Duncan, N. Kim, Y. Colon-Cortes, J. Rodriguez, M. Mazur, S. E. Birket, S. M. Rowe, N. E. West, A. Livraghi-Butrico, R. C. Boucher, J. Hanes, G. Aslanidi, J. S. Suk, An Adeno-Associated Viral Vector Capable of Penetrating the Mucus Barrier to Inhaled Gene Therapy. *Mol Ther Methods Clin Dev* **9**, 296-304 (2018).

69. D. B. Lima, T. B. de Lima, T. S. Balbuena, A. G. C. Neves-Ferreira, V. C. Barbosa, F. C. Gozzo, P. C. Carvalho, SIM-XL: A powerful and user-friendly tool for peptide cross-linking analysis. *Journal of Proteomics* **129**, 51-55 (2015).

70. H. J. Butt, B. Cappella, M. Kappl, Force measurements with the atomic force microscope: Technique, interpretation and applications. *Surface Science Reports* **59**, 1-152 (2005).

71. D. Nečas, P. Klapetek, Gwyddion: an open-source software for SPM data analysis. *Open Physics* **10**, 181-188 (2012).

72. P. W. Clapp, K. S. Lavrich, C. A. v. Heusden, E. R. Lazarowski, J. L. Carson, I. Jaspers, Cinnamaldehyde in flavored e-cigarette liquids temporarily suppresses bronchial epithelial cell ciliary motility by dysregulation of mitochondrial function. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **316**, L470-L486 (2019).