Network Statistics of the Whole-Brain Connectome of *Drosophila*

Albert Lin[*,1,2], Runzhe Yang[*,1,3], Sven Dorkenwald[1,3], Arie Matsliah[1], Amy R. Sterling[1], Philipp Schlegel[4,5], Szi-chieh Yu[1], Claire E. McKellar[1], Marta Costa[5], Katharina Eichler[5], Alexander Shakeel Bates[4,5,6], Nils Eckstein[7], Jan Funke[7], Gregory S.X.E. Jefferis[4,5], and Mala Murthy[1,✉]

[1]Princeton Neuroscience Institute, Princeton University, Princeton, NJ, USA
[2]Center for the Physics of Biological Function, Princeton University, Princeton, NJ, USA
[3]Computer Science Department, Princeton University, Princeton, NJ, USA
[4]Neurobiology Division, MRC Laboratory of Molecular Biology, Cambridge, UK
[5]Drosophila Connectomics Group, Department of Zoology, University of Cambridge, Cambridge, UK
[6]Centre for Neural Circuits and Behaviour, University of Oxford, Oxford, UK
[7]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, USA
[*] *These authors contributed equally to this work*

Abstract

Animal brains are complex organs composed of thousands of interconnected neurons. Characterizing the network properties of these brains is a requisite step towards understanding mechanisms of computation and information flow. With the completion of the Flywire project, we now have access to the connectome of a complete adult *Drosophila* brain, containing 130,000 neurons and millions of connections. Here, we present a statistical summary and data products of the Flywire connectome, delving into its network properties and topological features. To gain insights into local connectivity, we computed the prevalence of two- and three-node network motifs, examined their strengths and neurotransmitter compositions, and compared these topological metrics with wiring diagrams of other animals. We uncovered a population of highly connected neurons known as the "rich club" and identified subsets of neurons that may serve as integrators or broadcasters of signals. Finally, we examined subnetworks based on 78 anatomically defined brain regions. The freely available data and neuron populations presented here will serve as a foundation for models and experiments exploring the relationship between neural activity and anatomical structure.

Drosophila melanogaster | connectomics | brainwide network analysis
Correspondence: *sseung@princeton.edu*
Correspondence: *mmurthy@princeton.edu*

Introduction

The brain is a complex neural network, and understanding the structure of this network can offer insights into how brains support computations and regulate information flow. In this study, we focus on characterizing the network properties of the recently produced Flywire connectome, the first complete wiring diagram of an adult fly brain (1–3). By exploring the organization of this wiring diagram, we aim to provide derived data, including pre-computed statistics and neuron populations, that will serve as a valuable foundation for future experimental and theoretical work.

Advancements in connectomics have allowed researchers to look beyond specific circuits and characterize neural networks as a whole. Such studies have focused on applying mathematical network theory to biological wiring diagrams, including those of *C. elegans* (302 neurons) (4–7), the larval *Drosophila* brain (3,013 neurons) (8), and several partial connectomes, including from the larval zebrafish brainstem and olfactory bulb (9–11), the mouse olfactory bulb and visual cortex (12–14), and the adult *Drosophila* brain (15–21).

Network statistics computed from wiring diagrams can serve as a basis for comparing different wiring diagrams across brain regions, individuals, developmental stages, or species, enabling researchers to uncover commonalities and differences in neural network organization. Global metrics such as sparsity, connected component size, average path lengths, and small-worldness quantify the interconnectivity and diameter of a network (22–24). Spectral and rich club analyses characterize the robustness of a network's interconnectivity and can identify highly connected neurons that may act as hubs (25). For instance, many of the neurons which form the rich club in *C. elegans* are known to act as command interneurons, suggesting that well-connected nodes in biological networks are likely to have important functions (5, 7, 26). Local metrics, such as the distribution of network motifs, have been studied to uncover organizational principles of neural networks (7, 11, 14, 27–29). Network motifs are recurring patterns of connectivity that are thought of as "building blocks" (30, 31) and reflect the signal propagation and functional properties of complex networks. Several specific motifs—reciprocal connections (5–7, 11, 14), feedforward loops (7, 27, 28), and 3-unicycles (11, 32)—have received significant attention in neuroscience because of their implications for local computation and information flow.

In this paper, we present a large set of network statistics derived from the recently completed Flywire connectome of an adult *Drosophila* brain (**Table 1**). These statistics describe various aspects of the brain's interconnectivity, including path lengths between neurons, frequently traversed neural subpopulations, motif frequencies and strength, and more. We also identify several populations of well-connected neurons, including rich club neurons, highly reciprocal neurons, and neurons which may act as integrators or broadcasters of signals. Additionally, we characterize the connectivity within 78 brain regions and conduct statistical comparisons between the wiring diagrams of different species. With 128,000 neurons, the Flywire connectome is the largest neuron-level reconstruction to date. Thus, the analyses conducted in this study constitute the

	Computed network statistics	
Connected components	strongly connected components	Figure 1d
	weak connected components	Figure 1e
Path length analysis	directed shortest path lengths	Figure 1d
	undirected shortest path lengths	Figure 1e
Percolation analysis	vertex percolation	Figures 1f, g; S1f
	edge percolation	Figure S1a
Rich-club analysis	total-degree rich club	Figure 1h
	in-degree rich club	Figure S1g
	out-degree rich club	Figure S1g
Small-word analysis	clustering coefficient	Table 2
	small-wordness	Equation 1
2-neuron motifs	reciprocity	Table 2; Figures 5c; S5c
	connection strength	Figures 2a, d; 5f, S6a
	neurotransmitter types	Figures 2c, e, f; 5d, e; S5
3-neuron motifs	motif frequencies	Figures 3a; 6a, d; S7a
	motif strength	Figures 3b; 6c; S7b
	neurotransmitter types	Figures 3c, d, e
Large-scale connectivity	degree distribution	Figure 1c
	cell categories	Figure 4
Spectral analysis	forward random walk	Figure S1d
	reversed random walk	Figure S1e
Neuropil subgraphs	internal/external connection weights	Figure S4
	2-neuron motifs	Figures 5, S5, S6
	3-neuron motifs	Figures 6, S7

	Neuron lists available on Codex		# of neurons
2-neuron motifs	reciprocal connection participants	α ⇄ β	77,607
3-neuron motifs	feedforward loop participants	α, β, γ	113,978
	3-unicycle participants	α, β, γ	66,835
N-neuron motifs	highly reciprocal neurons		2,183
	neuropil-specific highly reciprocal neurons (NSRNs)		704
Rich-club analysis	rich-club neurons		40,218
	broadcasters		676
	integrators		638
Spectral analysis	attractors		3,469
	repellers		3,469

Table 1. Data availability. List of data products in this work, including statistics computed in this paper (left) and neuron populations (right). Complete, interactive neuron lists are available online as "Connectivity Tags" on Codex (`codex.flywire.ai`). Definitions for each of these neuron populations can be found in the text, and in **Table S1**.

largest statistical survey yet conducted on a biological network, and the only survey of a complete adult brain. Additionally, for the first time in any connectome, access to synapse-level neurotransmitter predictions (33) enables us to examine a wiring diagram with putative excitatory and inhibitory connections.

The data derived in this work offer a quantitative summary of the network of the adult fly brain, and lay the groundwork for future studies exploring connectivity in the fly. We expect these resources to be utilized both by experimentalists and theorists. For experimentalists, profiling neurons based on their connectivity features and searching populations of well-connected neurons will facilitate the generation of new hypotheses and the identification of key neurons in circuits or regions of interest. Additionally, an understanding of connectivity at the brain scale is crucial for interpreting whole-brain functional activity. From a theoretical perspective, these resources will be useful for modeling circuits at a multitude of scales, and are a vital first step toward developing whole-brain computational models. The neuron populations and motifs we've identified are freely available online at the Connectome Data Explorer (Codex: `codex.flywire.ai`).

Results

Summary of the dataset and definitions

To perform large-scale network analyses, we summarized the synaptic connections between neurons into the following data structure. For each pair of neurons, we sum the total number of synaptic connections to return the weight of their connection. Repeating for all neuron pairs gives us a weighted graph describing the connectome, with 127,978 neurons and over 32 million total thresholded connections, representing the complete *Drosophila* brain (2) (**Figure 1a**). In this paper, we will be using the term "connection" to denote an edge that exists in the network between two neurons, consisting of one or more synapses. The synapses in this dataset were detected automatically (34, 35). To minimize the impact of spurious synapses, we applied a threshold of 5 synapses per connection for all of the analyses conducted in this study except when synapse count distributions are presented (these are shown without threshold) (**Figure 1b, Methods**). We will be using synapse count as a proxy for edge strength in this paper: "stronger" and "weaker" refer to higher or lower synapse counts, respectively.

The Flywire connectome also contains synapse-level neurotransmitter predictions (33). The classifier applied to the dataset discriminates between six neurotransmitters: the fast-acting classical neurotransmitters acetylcholine (ach), GABA (gaba), and glutamate (glut) and the monoamines dopamine (da), octopamine (oct), and seratonin (ser). In the *Drosophila* nervous system, acetylcholine is excitatory and GABA is inhibitory. Glutamate can be either excitatory or inhibitory, but within the brain of the fly it has largely been observed to be inhibitory (36–38).

A key characteristic of the network is the distribution of degrees, which reflects the amount of connectivity between neurons. For any given neuron, the *in-degree* is defined as the number of presynaptic neurons (neurons it receives inputs from), and the *out-degree* is defined as the number of postsynaptic neurons (neurons it sends outputs to). With a threshold of 5 synapses per connection, the average in/out-degree of an intrinsic neuron in

Figure 1. Whole-brain network properties. (a) The FlyWire dataset (1–3) is an EM reconstruction of the complete brain of an adult female *Drosophila melanogaster*, with both hemispheres of the brain and both optic lobes. The volume contains 127,978 neurons and 32 million synapses with a threshold of 5 synapses/connection applied (subsample of synapse locations shown in the inset). **(b)** The distribution of the number of synapses per connected neuron pair. **(c)** The in-degree (number of presynaptic partners) plotted against the out-degree (number of post-synaptic partners, with log-scale x and y-axes. **(d)** Strongly connected components (SCCs) consist of a subset of nodes in a network which are mutually reachable via directed edges. In the fly brain there exists one giant SCC containing 93.3% of all neurons after thresholding at 5 synapses per connection. The distribution of shortest path lengths between neuron pairs within this SCC is plotted. **(e)** Weakly connected components (WCCs) consist of a subset of nodes in a network which are mutually reachable, regardless of edge direction. In the fly brain there exists one giant WCC containing 98.8% of all neurons. The distribution of path lengths between neuron pairs within this WCC is plotted. textbf(f) We examine the role high-degree neurons play in connecting the brain by plotting the sizes of the first two weakly connected components (WCCs) as nodes are removed by total degree. Removal of neurons starting with those with largest degree results in the brain splitting into two WCCs when neurons of approximately degree 50 start to be removed, a deviation from randomly connected networks (dotted lines). The largest surviving total degree as a function of the number of remaining nodes is plotted in gray. textbf(g) Removal of neurons starting with those with smallest degree results in a single giant WCC until all neurons are removed. The smallest surviving total degree as a function of the number of remaining nodes is plotted in gray. **(h)** The rich club coefficient (red) as a function of total degree, compared to the predicted rich-club coefficient of a CFG null model (dotted red). The relative rich club coefficient is plotted in blue. The range over which the relative rich club coefficient is greater than 1.01 is 37 to 93 and is shaded in gray. Because the rich club coefficient of a random network becomes unreliable as it grows increasingly sparse, we take all neurons with total degree > 37 to be part of the rich club.

the brain is 20.5 (2), but the distributions of in-degree and out-degree are not strongly correlated (**Figure 1c**). On average, each connection in the brain consists of approximately 12.6 synapses after the threshold is applied (2). Across the connectome, the probability that any two neurons is connected is 0.000161. This makes the wiring diagram of the fly brain a very sparse matrix when compared to the complete *C. elegans* nervous system, and to the sub-volume wiring diagrams reconstructed in larval zebrafish and mouse (**Table 2**).

Neurons in the brain form a single connected component

To assess the interconnectivity of the neurons in the brain, we searched the connectome for connected components using two sets of criteria. First, we looked for *strongly connected components* (SCCs). All neurons within an SCC are mutually reachable via directed pathways (39). Second, we looked for *weakly connected components* (WCCs), a relaxed criterion in which all neurons within a WCC are mutually reachable, ignoring the directionality of connections.

Despite its sparsity, the brain is highly connected under either criteria – 93.3% of neurons are contained in one giant

SCC, while 98.8% of neurons are contained in one giant WCC (**Figure 1d-e**). These giant connected components persist when either the strongest connections or the weakest connections are pruned (**Figure S1a-b**), indicating that connectivity in the brain is robust: many paths connect neuron pairs. Within the giant SCC, the average shortest directed path length between neuron pairs is 4.42 hops, with every neuron reachable within 13 hops (**Figure 1d**). In the giant WCC, the average shortest undirected path length between neuron pairs is 3.91 hops, with every neuron reachable within 11 hops (**Figure 1e**). These numbers are similar to those found in the hemibrain dataset (19). The short path lengths within both connected components show that despite its size, the fly brain is still relatively shallow when compared to artificial networks (40).

Spectral analysis of the whole-brain network

To better understand the network topology of the brain, we performed a spectral analysis of a random walk in the giant SCC. In this random walk, the transition probability from neuron α to neuron β is $p_{\alpha \to \beta} = \delta_{\alpha \to \beta}/d_\alpha^-$, where d_α^- is the out-degree of neuron α, and $\delta_{\alpha \to \beta} \in \{0, 1\}$ indicates the existence of a connection. Such a random walk converges to a stationary distribution over all neurons in the giant SCC (**Figure S1d**). We found that in this random walk, 3% of neurons were visited 61.2% of the time—the remaining 97% of neurons were visited only 38.8% of the time. These top visited neurons can therefore be classified as *attractor nodes* (41) in the network. These attractor nodes typically make connections in the gnathal ganglia (GNG), a large midline neuropil which both sends and receives information from the periphery and contains a large number of neurons which connect with the ventral nerve cord (VNC).

We also performed a "reverse" walk within the giant SCC, reversing edge directionality so that the transition probability from neuron α to neuron β is $p_{\alpha \to \beta}^{\text{rev}} = \delta_{\beta \to \alpha}/d_\alpha^+$, where d_α^+ is the in-degree of neuron α. The reversed walk also converges to a stationary distribution in which 3% of neurons were visited 42.4% of the time (**Figure S1e**). These highly visited visited neurons in a reverse random walk are *repeller nodes* in the network. Many of these neurons make synapses in the antennal lobes (AL) and medullae (ME) and are close to olfactory and visual sensory neurons, suggesting that neurons in the olfactory and visual systems are more connected than the other sensory modalities.

The brain has a large rich club of neurons

Is the high interconnectivity observed in the fly brain a consequence of a relatively large number of interconnected neurons, or is it dependent on a small number of very highly connected "hub" neurons? To assess this, we constructed survival curves, removing neurons from the network starting with those of highest total degree (in-degree + out-degree). The giant WCC survives until a total degree of 40, at which point the network splits into two WCCs (**Figure 1f**). These two WCCs correspond to a split between the left and right hemispheres, and demonstrate that despite the hemispheric anatomy of the brain, the two hemispheres are highly interconnected: they do not split into separate networks until about 60% of all neurons are removed. Removing neurons from the network by smallest total degree does not result in division of the giant connected component (**Figure 1g**). These results remain qualitatively consistent when neurons are pruned by either in-degree or out-degree alone (**Figure S1f**). This indicates that the interconnectivity of the brain is robust, and not dependent on a small number of highly connected neurons.

Many networks exhibit the "rich club" property (25, 26, 42), in which well connected nodes are preferentially connected to other well connected nodes (see **Methods**). In Flywire, we find that neurons with total degree exceeding 37 are more highly interconnected than one would expect from a randomly connected network, forming a large rich club of 40,218 neurons, approximately 30% of all neurons in the brain (**Figure 1h**). The connection density within this rich club is 0.000870, 5.4 times higher than the overall connection density in the brain. Such a large rich club suggests that the topology of the fly wiring diagram is fairly distributed. This is consistent with the connected component observations, which also suggest a degree of robustness. A rich club analysis considering in-degree alone returns an in-degree threshold of 10, while no rich club is observed when considering out-degree alone (**Figure S1g**).

The fraction of neurons in the rich club is substantially larger in the fly than in *C. elegans*, which has a rich club of 11 neurons (4% of the neurons in the worm) (26). The worm rich club contains known hub neurons, such as the command neurons AVA and AVB. Such highly connected hub neurons do not seem to be present in the fly brain—while there are neurons with very high degrees, there also exist alternate paths between most neuron pairs. The larger rich club we observe in the fly brain relative to that of the worm may be a consequence of scale: a large network can be more robust and flexible by employing a sizeable population of well-connected neurons in a distributed manner rather than relying on a small number of highly connected hubs. We will further examine the properties of this rich club population in the section: ***Large-scale connectivity in the brain***.

Reciprocal and recurrent motifs are overrepresented in the brain

Connection reciprocity is a measure of the amount of direct feedback in the brain: given that neuron α is connected to neuron β, what is the probability that neuron β is connected back to neuron α? Across the whole brain, this connection reciprocity probability is 0.138 (**Table 2**). The connection reciprocity in the is significantly higher than in both the Erdős-Rényi (ER) and configuration (CFG) random null models (**Methods**). The overrepresentation of reciprocal connections in brains is well established, and our results are consistent with previous observations both in *Drosophila* (16, 17, 19) and in other species (5, 7, 14, 27, 28, 43).

We also computed the clustering coefficient, a higher-order connectivity metric which assesses the prevalence of triplet structures in the network irrespective of edge direction: if neuron α and neuron β are connected and neuron α and neuron γ are connected, what is the probability that neuron β and neuron γ are also connected? The clustering coefficient in the brain is

0.0477 (**Table 2**). As was the case with reciprocity, this value of clustering coefficient is higher than in both ER and CFG null models. The high clustering coefficient demonstrates that the network of the fly brain is highly connected and is nonrandom in its structure.

We compared these metrics with two existing whole-animal connectomes, the hermaphrodite and male *C. elegans* (4, 5, 7), and with two sub-volume wiring diagrams, the hindbrain of a larval zebrafish (11) and a region of L2/3 mouse visual cortex (14) (**Table 2**). The whole fly brain is approximately 200 times sparser than the *C. elegans* connectome. Among the neurons entirely contained within the zebrafish hindbrain and mouse visual cortex sub-volumes, connection probability is higher than in the fly. Despite these differences in density, the values of reciprocity and clustering coefficient are comparable across all five datasets. Note that because many neurons in the fish and mouse sub-volumes are truncated, measures of reciprocity and clustering coefficient in these datasets cannot capture long-range reciprocal connections. In addition, the statistics for *C. elegans*, zebrafish hindbrain, and mouse cortex were computed with no synapse threshold, while the fly brain included a threshold of 5 synapses per connection. Varying this threshold in the fly, however, did not significantly alter reciprocity and clustering coefficient values (**Figure S1c, Table S2**).

Small-worldness of the fly brain

A "small-world" network is one in which nodes are highly clustered and path lengths are short (24). High small-worldness coefficients are associated with efficient communication between nodes (44, 45). We quantified the small-worldness of the connectome by comparing it to an Erdős-Rényi graph (ER) (46). The average undirected path length in the ER graph, denoted as ℓ_{rand}, is estimated to be 3.57 hops, similar to the observed average path length in the fly brain's WCC ($\ell_{\text{obs}} = 3.91$). The clustering coefficient (C_{rand}^{Δ}) of the ER graph is only 0.0003, much smaller than the observed clustering coefficient ($C_{\text{obs}}^{\Delta} = 0.0463$) (**Table 2**, **Methods**). The small-worldness coefficient of the fly connectome is:

$$S^{\Delta} = \frac{C_{\text{obs}}^{\Delta}/C_{\text{rand}}^{\Delta}}{\ell_{\text{obs}}/\ell_{\text{rand}}} = 141, \tag{1}$$

significantly higher than that of the *C. elegans* connectome ($S^{\Delta} = 3.21$) and close to that of the internet ($S^{\Delta} = 98.1$) (24), implying highly effective global communication among neurons in the brain.

Strength and neurotransmitter composition of reciprocal connections

The average strength of edges participating in reciprocal connections is higher than the average strength of unidirectional connections (**Figure 2a**). The majority of unidirectional connections are cholinergic (excitatory), while edges participating in reciprocal connections contain fewer cholinergic neurons and more GABAergic neurons (**Figure 2b**). Inhibitory connections in the brain have more synapses on average than excitatory connections (2), which may partially explain the higher average strength of reciprocal connections. The most common reciprocal pairing is between a cholinergic neuron and a

GABAergic neuron and the second most common pairing is acetylcholine-glutamate (**Figure 2c**). Both of these reciprocal motifs are excitatory-inhibitory (E-I), and both are overrepresented when compared to the neurotransmitter frequencies observed for reciprocal connections (**Figure 2b**). Excitatory-excitatory (E-E) acetylcholine-acetylcholine pairs are in contrast underrepresented, as are inhibitory-inhibitory (I-I) GABA-GABA pairs. We observed reciprocal E-I (acetylcholine-GABA and acetylcholine-glutamate) connection strengths to be weakly correlated, while E-E (acetylcholine-acetylcholine) pairs were uncorrelated (**Figure 2d**). Examples of reciprocal neuron pairs are shown in **Figure 2g**.

Reciprocal degree across the neuronal population

Of the 127,978 neurons in the whole brain, 77,607 participate in at least one reciprocal connection: approximately 2 in every 3 neurons. Many neurons in the brain participate in multiple reciprocal connections. To characterize these neurons, we define the *reciprocal degree* as the number of reciprocal connections made by a given neuron (**Figure S2a**). Plotting the distributions of reciprocal degree by neurotransmitter, we observe that the overwhelming majority of neurons with high reciprocal degree ($d^{\text{rec}} > 100$) are GABAergic (**Figures 2e, S2b**), while at lower reciprocal degrees ($d^{\text{rec}} < 100$), all three primary neurotransmitter types are well represented.

What fraction of a neuron's connections are reciprocal? Note that here, we are not considering reciprocity between cell types, but rather between pairs of individual neurons. For most neurons these fractions are low—on average 23% of incoming and 18% of outgoing connections are reciprocal. Plotting the fraction of reciprocal incoming connections against the fraction of reciprocal outgoing connections, we observe only a weak correlation (**Figure S2c**), suggesting that a given neuron's reciprocal degree is not strongly coupled to either its in-degree or its out-degree. Comparing the number of reciprocal connections neurons make to the total number of connections they make by plotting $2\times$ the reciprocal degree against the total degree of neurons (in-degree + out-degree), we again see no relationship (**Figure 2f**). Dividing the neuron population by neurotransmitter, however, we find that neurons of high total degree are mostly GABAergic, and that for many of these neurons, more than half of their total connections are reciprocal (**Figure S2d**). These highly reciprocal neurons may provide large-scale feedback inhibition in the brain (*see below*: ***Identifying neuropil-specific reciprocal neurons***).

Strength and neurotransmitter composition of three-node motifs

The high clustering coefficient of the brain implies an overrepresentation of triplet structures. We determined the frequency at which each of the 12 directed three-node motifs occur in the brain (**Figure 3a**). Feedforward motifs (motifs #1-3) are underrepresented when compared to both ER and CFG null models, while all others, including the highly recurrent motifs (motifs #7-13), are overrepresented. The strengths of edges participating in 3-node motifs are higher than the average edge strength (**Figure 3b**). Complex 3-node motifs which contain reciprocal

Neuronal wiring diagrams	Fruit fly *Drosophila melanogaster* (Dorkenwald et al., 2023)	Nematode *Hermaphrodite C. elegans* (Cook et al., 2019)	Nematode *Male C. elegans* (Cook et al., 2019)	Larval zebrafish (sub-vol.) *Danio rerio* (hindbrain) (Yang et al., 2023)	Mouse (sub-vol.) *Mus musculus* (V1 L2/3) (Turner et al., 2022)
Network size	**127,978 neurons** **2,613,129 connections**	**302 neurons** **3,242 connections**	**364 neurons** **3,467 connections**	**419 neurons** **5,605 connections**	**111 neurons** **659 connections**
Avg. connection strength	**12.61 synapses** 5 ~ 2358	**3.15 synapses** 1 ~ 36	**3.59 synapses** 1 ~ 63	**1.69 synapses** 1 ~ 21	**1.14 synapses** 1 ~ 5
Connection probability $\alpha \dashrightarrow \beta$ Pr	**0.000160** x1	**0.0356** x222 denser than fly	**0.0262** x164 denser than fly	**0.0320** x200 denser than fly	**0.0540** x360 denser than fly
Connection reciprocity $\alpha \rightleftharpoons \beta$ Pr	**0.138** x858 than ER x43.8 than CFG	**0.372** x10.4 than ER x5.03 than CFG	**0.386** x14.7 than ER x6.02 than CFG	**0.113** x3.53 than ER x2.64 than CFG	**0.088** x1.63 than ER x1.33 than CFG
Clustering coefficient α $\beta \quad \gamma$ Pr	**0.0463** x144 than ER x7.57 than CFG	**0.284** x4.06 than ER x1.86 than CFG	**0.331** x6.39 than ER x2.40 than CFG	**0.182** x2.89 than ER x1.90 than CFG	**0.159** x1.51 than ER x1.06 than CFG

Table 2. Connection probabilities, reciprocity, and clustering coefficient in the fly brain and compared across animals. The probability that any two neurons in the fly brain are connected is 0.000160. Connection reciprocity (the probability that two connected neurons are reciprocally connected) in the fly is 0.138, larger than in either an ER or CFG null model. The clustering coefficient (the probability that if neuron α and neuron β are connected and neuron α and neuron γ are connected, then neuron β and neuron γ are also connected, irrespective of directionality) in the fly is 0.0463. Both reciprocity and clustering coefficient are higher than expected with ER and CFG null models. Statistics for the fly were computed with a threshold of 5 synapses per connection. Values for thresholds from 0 to 50 are plotted in **Figure S1c**. Statistics for *C. elegans* were computed for the chemical networks of neurons in hermaphrodite and male worms (5), with no synapse threshold applied. Statistics for larval zebrafish hindbrain (11) and mouse visual cortex (14) were computed with no threshold. Note that any truncated neurons in the zebrafish hindbrain and mouse cortex were excluded from these analyses.

connections tend to be stronger than feedforward motifs.

Examining the neurotransmitter composition of two of these three-node motifs, feedforward loops (motif #4) and 3-unicycles (motif #7) (**Figure 3c**), we found that edges which participate in feedforward loops were predominantly cholinergic, and that the most common neurotransmitter composition for a feedforward loop is three cholinergic neurons, a feedforward excitatory configuration (**Figure 3d**). The next most common compositions contain either one or two inhibitory (GABAergic or glutamatergic) edges. Feedforward loops with one inhibitory edge are likely feedforward inhibition motifs, while loops with two inhibitory edges are likely disinhibition motifs. 3-unicycles in contrast contain a higher proportion of inhibitory GABAergic and glutamatergic neurons, and the three most common 3-unicycle compositions all contain at least one inhibitory neuron (**Figure 3e**). These cycles may act as indirect feedback inhibition circuits. It is interesting to note that the observed neurotransmitter composition frequencies are closer to what may be expected by chance for feedforward loops than they are for 3-unicycles. Examples of neurons which form 3-node motifs are shown in **Figure 3f**.

The fly brain exhibits a high clustering coefficient and an overrepresentation of highly connected 3-neuron motifs. These observations suggest that the local structure of the brain displays a high degree of non-randomness, in line with previous studies in *C. elegans* (5, 7) and in mouse cortex (14, 27, 28). The overrepresentation of feedforward loops (motif #4) has been widely observed in other biological networks (5, 7, 27, 28). This overrepresentation is present in most neuropils in the brain. It is possible that these feedforward loops, which are predominantly excitatory, may form large-scale feedforward structures which span brain regions. 3-unicycles (motif #7) may form recurrent local circuits capable of generating persistent neural activity (11).

Large-scale connectivity in the brain

Within the adult brain, the in-degree and out-degree of neurons are only weakly correlated. Neurons with few inputs and many outputs may serve as broadcasters of signals, while those with many inputs and few outputs may act as integrators. To examine these populations of neurons, we divided the intrinsic rich club neuron population into three categories based on their in-degree and out-degree (**Figure 4a**). While these boundaries are necessarily arbitrary, they nonetheless help us examine the nature of large-scale connectivity in the brain. We divided the rich club neurons by defining *broadcaster neurons* (Region 2) as those for which out-degree $\geq 5\times$ in-degree, and *integrator neurons* (Region 4) as those for which in-degree $\geq 5\times$ out-degree. In the Flywire connectome we find 676 broadcasters and 638 integrators. The remaining intrinsic rich club neurons (37,093) fall into the "large balanced" category (Region 3), including most highly reciprocal neurons. Some examples of broadcasters, integrators, and rich balanced neurons are shown in **Figure 4f**.

When compared to the population of all neurons, rich club neurons are less likely to be cholinergic and more likely to be GABAergic (**Figures 4b, S3a**). Integrator neurons are even less likely to be cholinergic (49%), and include a large fraction of dopaminergic neurons, suggesting that these neurons may be engaged during learning. In contrast, broadcaster neurons are predominantly cholinergic and therefore excitatory (75%). Central brain neurons are dramatically overrepresented in the rich club, while optic lobe intrinsic neurons are underrepresented (**Figures 4c, S3b**). Many integrators are either central brain intrinsic neurons or visual projection neurons. In contrast, few broadcasters are intrinsic to the central brain—many are visual centrifugal neurons or optic lobe intrinsic neurons. These in-

Figure 2. Characterizing reciprocal connections in the brain. (a) Edges that are part of reciprocal connections are stronger on average than unidirectional connections. **(b)** Breakdown of unidirectional and reciprocal edges by neurotransmitter. Unidirectional connections are most likely to be cholinergic. Reciprocal connections are more likely than unidirectional connections to contain a GABAergic neuron. **(c)** The frequency of neurotransmitter pairs forming reciprocal connections, compared to the expected frequency of neurotransmitter pairs under the assumption of independent neurotransmitter choice (red). A majority of reciprocal connections are formed by acetylcholine-GABA pairs. The next most common reciprocal connection type is acetylcholine-glutamate, with acetylcholine-acetylcholine pairs underrepresented. **(d)** Heatmaps of the relative strengths (synapse counts) of the two connections forming acetylcholine-GABA reciprocal pairs (left), acetylcholine-glutamate reciprocal pairs (center), and acetylcholine-acetylcholine reciprocal pairs (right). The strengths of the edges of reciprocal pairs are uncorrelated. Excitatory-inhibitory pairs (acetylcholine-GABA and acetylcholine-glutamate) have higher average strengths than excitatory-excitatory (acetylcholine-acetylcholine) pairs. **(e)** Distributions of reciprocal degree (the number of reciprocal connections a given neuron makes) for cholinergic neurons (left), GABAergic neurons (middle), and glutamatergic neurons (right). GABAergic neurons are more likely to make large numbers of reciprocal connections, while cholingeric neurons are more likely to have smaller numbers of reciprocal connections. **(f)** Scatterplot of 2 times the reciprocal degree of neurons versus their total degree (in-degree + out-degree). Dotted lines indicate a factor of 2 around the $x = y$ line. Large neurons for which reciprocal connections form the majority of their total connections are most likely to be GABAergic. **(g)** Visualizations of exemplar reciprocal neuron pairs. Cell labels are listed where available.

clude a large number of Mi1 and Tm3 neurons, excitatory cells in the medullae (ME) known to play key roles in the motion detection circuit (15, 16, 47). Most neurons are restricted to a single hemisphere—just 11% of neurons have inputs in both hemispheres and 11% have outputs in both hemispheres (**Figure 4d-e**)(2). In comparison, rich club neurons are more likely to have inputs or outputs spanning both hemispheres: 18% and 17%, respectively. This is more common for integrator neurons (23%) than it is for broadcaster neurons (16%). The asymmetry in L/R percentages for broadcaster neurons is due to the large number of medulla-intrinsic broadcasters. Many of these neurons receive inputs from the lamina, and the left lamina of the FAFB sample is still attached to the optic lobe, while the right lamina of the sample is partially detached.

Rich club neurons are closer on average to sensory inputs

To assess the distance of the rich club neurons from sensory inputs, we employed a probabilistic information flow model to determine the relative distance of each neuron (in hops) from a set of seed neurons (**Methods**) (2, 20). The model was run with different sets of seed neurons, each corresponding to a specific sensory modality (olfactory, gustatory, visual, etc.), as well as on the complete set of all sensory inputs, giving us the distance from each neuron in the dataset to each sensory modality. Ranking these distances and normalizing returned the percentile rank of each neuron with respect to each modality. Neurons with percentile rank less than 50% are closer than average to the given sensory input, while neurons with percentile rank greater than

Figure 3. Examining 3-node motifs. (a) The distribution of three-node motifs across the whole brain. Absolute counts of each motif are on the left, and the frequency of each motif relative to that in an ER null model is plotted to the right, together with the average motif frequencies of 100 CFG models (gray violin plots). When we compare the whole-brain network to both ER and CFG null models, we observe an underrepresentation of simple motifs (#1-3) and an overrepresentation of other motifs, particularly highly recurrent motifs (#10, 12, 13). **(b)** The average strength of edges that are part of the 3-node motifs. The dotted line is the average connection strength in the brain. **(c)** Breakdown by neurotransmitter of edges participating in two motifs: feed-forward loops (motif #4) and 3-unicycles (motif #7). Edges in feed-forward loops are more likely to be cholinergic. **(d)** Further examining the neurotransmitter composition of these motifs, we find that feed-forward loops (motif #4) are most likely to be acetylcholine-acetylcholine-acetylcholine, **(e)** while 3-unicycles (motif #7) tend to contain at least one inhibitory edge (glutamate or GABA). **(f)** Visualizations of exemplar 3-node motifs. Cell labels are listed where available.

50% are farther.

The rich club neurons have a mean percentile rank of 44% relative to the set of all sensory inputs (**Figure 4g**). Integrators have a mean percentile rank of 43%, while broadcasters have a mean percentile rank of 53%. The percentile rank of rich club neurons peaks between 10% and 30%, fairly close to the sensory periphery (**Figure 4g**). Integrator neurons are closest, with many having a percentile rank of less than 10%. The distribution of broadcasters is bifurcated, with one peak closer to inputs and another peak far from inputs. The first peak may consist of both neurons propagating sensory information or providing large-scale feedback to sensory systems, while the second peak may consist of downstream feedforward neurons. Examining the ranks with respect to individual sensory modalities, we find that rich club neurons are closer than average to all sensory modalities except the photoreceptors (visual) (**Figures 4g-h, S3c**). The average percentile rank of broadcasters to single sensory inputs tends to be lower than it is for the set of all inputs. This is likely because ranking from a seed population of all inputs will rank integrators before many broadcasters. In contrast, when looking at a single modality, neurons which are primarily connected to a different modality may be ranked later. In most modalities we observe a similar bifurcation in the distribution of broadcaster neuron ranks (**Figures 4g, S3c**).

We can examine the distance of neurons to multiple sensory modalities by plotting the percentile rank of neurons with respect to one modality against the percentile rank of neurons of another modality (**Figure S3d**). Broadcaster and integrator neurons are scattered throughout these distributions, but tend to be closer than average to multiple sensory inputs. These rich

Figure 4. Large-scale neuron connectivity in the brain. (a) Using the in-degree vs. out-degree scatterplot, we can divide the intrinsic rich club neurons into three distinct categories: broadcasters, integrators, and large balanced neurons. Comparing the prevalence of **(b)** neurotransmitters, **(c)** intrinsic superclasses (optic lobe intrinsic, visual projection, visual centrifugal, and central brain intrinsic), and **(d)** input and **(e)** output sides of all intrinsic neurons, rich club neurons, integrators, and broadcasters. **(f)** Examples of rich club neurons in these three categories. **(g)** Applying the information flow model from Schlegel et al. 2021 (2, 20), we determined the percentile rank distributions of rich club, integrator, and broadcaster neuron populations from all inputs to the brain (above), as well as to specific modalities (olfactory below, refer to **Figure S3c** for others). **(h)** Average percentile rank of rich club, integrator, and broadcaster neurons for different modalities. Across all modalities except visual inputs, rich club neurons are closer than average (0.5) to the input layer.

club neurons may be a fruitful starting point when searching for neurons to characterize experimentally. In particular, integrator and broadcaster neurons which are low in rank relative to multiple sensory modalities may be good candidate sites of multi-sensory integration and information propagation.

Differences in connectivity across brain regions

The fly brain consists of a large number of distinct anatomical brain regions, or neuropils (48). The Flywire connectome has been segmented into 78 neuropils (**Figure 5a**), each with different average connection strengths (2). To understand information flow between neuropils, we employed a fractional weighting method accounting for each neuron's projections to and from every neuropil (**Methods**)(2). From these, we computed for each neuropil the relative fraction of internal, external incoming, and external outgoing connection weights (**Figure S4a-b**). These fractions reflect, respectively, the net number of connections within, being received, and being sent from each neuropil.

We find significant differences in these fractions across brain regions: the ellipsoid body (EB) and fan-shaped body (FB) of the central complex have the highest fraction of internal connections, while in other regions, such as the compartments of the mushroom body (MB), the majority of connections are external. Some regions such as the lateral horn (LH) send more external connections than they receive, while others such as the lobula plate (LOP) receive more external connections than they send. The fraction of internal connection weights is not correlated with neuropil size: while large neuropils such as the anterior and posterior ventrolateral protocerebra (AVLP and PVLP) have significant fractions of internal weights, they do not rank the highest. We note that under this classification, internal weights include any neurons with endings outside the brain, such as sensory, ascending, and descending neurons. This likely accounts for the high fraction of internal weights in regions such as the medullae (ME), which receive inputs from R7 and R8 photoreceptors, and the gnathal ganglia (GNG), which connects with large numbers of both ascending and descending neurons. Across the brain, 52% of all connection weights can be classified as internal. Comparing the putative neurotransmitters of the neurons contributing connection weights, we see that internal connections are more likely than external ones to be inhibitory (GABAergic or glutamatergic) (**Figure S4c**). We also see differences in neurotransmitter composition across brain regions (**Figure S4d**).

Prevalence and neurotransmitter composition of reciprocal connections differ across neuropils

To perform motif analyses within each neuropil, we first identified a subnetwork for each neuropil which treats all connections made within that neuropil as edges and includes all neurons connected to these edges (**Figures 5b, S5a**). Different neuropil subnetworks differ notably in both connection strength and density (**Figure S5b**). We computed the reciprocity in each neuropil subnetwork (**Figures 5c, S5c**). Neuropils with particularly high reciprocity probabilities include those in the central complex (FB, EB, and NO) and the two antennal lobes (AL). The relative number of reciprocal connections (reciprocity normalized by neuropil connection density) is high in the mushroom bodies (MB) and medullae (ME) (**Figure S5b**). Note that for these motif analyses, the results for small neuropils such as the cantles (CAN), bulbs (BU), galls (GA), accessory medullae (AME), and ocellar ganglion (OCG) are less interpretable due to the small number of samples.

In most neuropils, as in the whole brain, reciprocal connections are stronger than unidirectional connections, though the ratio of average strengths varies across neuropils (**Figure S5d**). Exceptions include the protocerebral bridge (PB), mushroom body calyces (MB-CA), and bulbs (BU), which have stronger unidirectional connections than reciprocal connections. Comparing the relative prevalence of each neurotransmitter in reciprocal and unidirectional connections, we again see differences between neuropils (**Figures 5d-e, S5d-h**). While reciprocal connections in most neuropils contain fewer cholinergic edges and more GABAergic edges than unidirectional connections, there are notable exceptions, such as in the neuropils of the central complex (FB, EB, PB, and NO). In the compartments of the mushroom body (MB) we find especially large differences in neurotransmitter composition between unidirectional and reciprocal connections. Comparing the strengths of the edges of reciprocal excitatory-inhibitory (acetylcholine-GABA) connections within neuropil subnetworks, we observe that E-I connection strengths are more strongly correlated in some neuropils (such as the FB and NO) than in others (**Figures 5f, S6a-b**). These correlations do not appear to be dependent on neuropil size (**Figure S6c**).

Identifying neuropil-specific reciprocal neurons

We performed a comprehensive search for intrinsic highly reciprocal rich club neurons that make the majority of their connections within a single neuropil, and found 1,863 neurons that meet these criteria (**Figure 5g**). These *neuropil-specific highly reciprocal neurons* (NSRNs) are predominantly inhibitory: 54% are GABAergic and another 10% are glutamatergic (**Figure S6d**). In some neuropils, such as the antennal lobes (AL), medullae (MB), and ellipsoid body (EB), there are many NSRNs, while in other neuropils, such as the superior posterior slopes (SPS) and posteriorlateral protocerebra (PLP), there exist only a handful of such neurons.

Some NSRNs, like the APL neurons in the MB (49, 50), CT1 neurons in the LO (16, 51, 52), or antennal lobe local neurons (ALLNs) (53, 54), have been previously characterized as providing global feedback inhibition in different regions. These neurons tend to be highly branched, with individual processes making reciprocal connections with different feedforward neurons. Some have been shown to have compartmentalized activity, raising the possibility of local computation within these neurons (55–57). Many of the NSRNs identified here have yet to be characterized. They may play similar roles in other circuits—for instance, it is likely that some of the NSRNs found in the AVLP provide feedback to the auditory circuits which span this brain region (58).

Figure 5. Neuropil-specific differences in connectivity. **(a)** An exploded view of the brain showing the brain regions, or neuropils, that the FlyWire dataset is divided into. Each synapse is assigned to a neuropil based on synapse location. **(b)** A schematic showing how neuropil subnetworks are identified for motif analyses. With the standard threshold of 5 synapses per edge applied, all connections composed of synapses within the neuropil of interest (Neuropil A) are treated as edges of the Neuropil A subnetwork. All neurons reached by this set of edges are included in the subnetwork. However connections composed of synapses outside of Neuropil A are not included, even if those connections involve neurons included in the subnetwork. **(c)** The reciprocity within each neuropil subnetwork. Differences in the percentage of **(d)** cholinergic and **(e)** GABAergic neurons between bidirectional (reciprocal) and unidirectional connections across neuropils. Refer to **Figure S5** for the absolute percentages. **(f)** Heatmaps showing the relationship between excitatory and inhibitory connection strengths in reciprocal connections in different brain regions. **(g)** Assessing the number of large (rich club), highly reciprocal neurons which span specific neuropils: making most of their incoming and outgoing connections within a single neuropil and also having a high reciprocal degree. Examples of neurons which meet these criteria are shown.

Figure 6. Differences in three-node motifs across neuropils. (a) Three-node motif distributions for three example neuropils: the EB, AL(R), and MB-ML(R). The frequency of each motif relative to that in an ER null model is plotted to the right, together with the average motif frequencies of 100 CFG models (gray violin plots). Further examples of other neuropils available in **Figure S7a. (b)** Motif frequencies for the 3-node motifs across all 78 neuropil subnetworks, normalized by their respective CFG null models. **(c)** Average strengths of edges participating in 3-node motifs in the different neuropil subnetworks relative to the average 3-node motif strength in each subnetwork. Refer to **Figure S7b** for average strengths relative to average neuropil subnetwork edge strength.

Three-node motifs differ across neuropils in their prevalence and strength

We computed the prevalence of three-node motifs in each neuropil subnetwork, and compared the motif frequencies to ER and CFG random null models constructed for each subnetwork (**Figures 6a, S7a**). Across most neuropils, we observed the same trend as we do across the entire brain: an underrepresentation of feedforward motifs (#1-3) and an overrepresentation of complex motifs (**Figure 6b**). However, there are notable differences between neuropils. In the cantles (CAN), epaulettes (EPA), and gorgets (GOR), for example, the frequency of 3-node motifs was closer to that expected in a CFG null model, while in other neuropils like the ellipsoid body (EB), complex motifs are highly overrepresented (**Figure 6b**).

Feedforward loops (motif #4) are overrepresented in most neuropils, excepting in the fan-shaped body (FB), ellipsoid body (EB), noduli (NO), and mushroom body compartments (MB). This suggests a relative underrepresentation of both feedforward excitatory and feedforward inhibitory circuits in these brain regions. 3-unicycles (motif #7), an indirect feedback inhibition circuit, are overrepresented across the whole brain (**Figure 3c**) but are underrepresented in most neuropils. The notable exceptions, the medullae (ME) and gnathal ganglia (GNG), are very large neuropils and have many sensory inputs. The overrepresentation of 3-unicycles in the meddulae implies the existence of localized cyclic structures within the early visual circuitry. Interestingly, this motif is also overrepresented in the zebrafish oculomotor circuit (11). Motifs #7-10 are underrepresented in the antennal lobes (AL), perhaps a result of the small number of unidirectional edges in these regions. The

most highly connected motifs (#12-13) are particularly overrepresented in the ellipsoid body (EB) and fan-shaped body (FB), consistent with their high reciprocity.

In most neuropils, we find that edges participating in underrepresented motifs are also weaker on average than edges participating in overrepresented motifs (**Figure 6c**). We also observe that in most neuropil subnetworks, edges participating in 3-node motifs are stronger than the average subnetwork edge (**Figure S7b**). This is broadly consistent with the whole-brain 3-node motif strength results. A notable exception is in the laminae (LA), where feedforward connections are strong despite being underrepresented.

Discussion

Here, we have provided a broad overview of the network properties of the *Drosophila* brain, laying the groundwork for identifying neurons and circuit motifs of biological interest and for modeling of particular circuits. In addition to the topology of the neural network, we have taken advantage of spatial information (innervation in different neuropils), neuron class distinctions (sensory versus descending, for example), cell type labels, and neurotransmitter predictions to better contextualize and interpret the network features we uncovered. We compared the statistics of the fly connectome to other wiring diagrams, carried out a comprehensive brain-wide search for 2- and 3-node connectivity motifs, identified highly connected broadcaster and integrator neurons, and identified differences in connectivity in different brain regions. The complete Flywire dataset is freely available online via Codex (Connectome Data Explorer: `codex.flywire.ai`), along with interactive lists of the neurons discussed in this work. These data will allow researchers to profile neurons by their connectivity features and identify key neurons within their circuits or brain regions of interest, a useful resource for hypothesis generation or model development. Experimentally examining highly connected neurons, such as the attractors, repellers, integrators, broadcasters, and NSRNs identified here, may also prove fruitful for linking circuit-level findings with broader activity patterns.

An understanding of how the whole-brain network shapes brain function is particularly important in light of recent experimental findings. A common approach in modern experimental neuroscience is to use anatomical wiring diagrams to generate circuit-level hypotheses, and to test these hypotheses by imaging and perturbing single cells or cell types. However, recent whole-brain imaging experiments, both in the fly (59–61) and in other species (62–69), have revealed brain-wide activity patterns related to both sensory processing (of individual modalities) and simple behaviors (like locomotion). To fully understand distributed computations and information flow in the brain, we must consider interactions not just at the scale of tens of neurons, but at the brain scale. Availability of network statistics at the scale of brain regions, coupled with the broad mesoscale connectivity between brain regions (2), will enable hypothesis generation at the whole-brain scale. Different neuropils serve different functions, and our work now highlights how these different functions are subserved by differences in connection strength, internal connectivity, motif frequency, and

neurotransmitter composition. For example, the central complex (neuropils FB, EB, PB, and NO), which has persistent activity associated with an internal representation of heading (70–75), contains some of the most reciprocal brain regions and has a large number of internal connections. Examination of other neuropil subnetworks may help us generate hypotheses regarding the function of less well-studied neuropils.

In this work, we comprehensively explored 2-node and 3-node motifs, and highlighted several large-scale connectivity patterns by exploring broadcaster (few-to-many), integrator (many-to-few), and highly reciprocal neurons. There remains, of course, a space of larger network motifs to explore. We have integrated the network motif search and visualization tool Vimo (76) into Codex, which allows users to query the Flywire connectome for any network motif of interest.

Limitations

The availability of neurotransmitter predictions greatly enhanced our ability to interpret the circuit motifs we found in the connectome. However, while these predictions are 94% accurate when compared to a set of ground truth neurons, there are several known cases where the predicted neurotransmitter does not align with the known transmitter. When interpreting results on the network scale, we must keep this error rate in mind. Also, monoamines beyond dopamine, octopamine, and seratonin are not accounted for in these predictions. More details on the neurotransmitter predictions are discussed in Eckstein *et al.* (33). It should also be noted that the synaptic connectome does not provide a complete picture of information flow in the brain. We currently do not have a complete map of gap junctions in the fly, and the extent to which extrasynaptic communication (via non-synaptic release of amines or neuropeptides) shapes neural activity in the *Drosophila* brain remains an open question (77–79).

Local circuit motifs are often inferred to be feedforward or feedback connections, with different theorized roles. While we are able to make such inferences on the population level, it can be difficult to place local circuits in the context of global directionality from sensory input to motor output. In this work, we employed an information flow method to rank the neurons by an effective difference from various sensory modalities (2). Ultimately, however, directionality of information flow in particular circuits, especially those in regions of the brain far from sensory inputs or motor outputs, must be determined through functional activity experiments and modeling.

The rich club compensates for anatomical bottlenecks

The anatomy of the fly brain suggests several potential network bottlenecks: one between left hemisphere and right hemisphere and one between the central brain and optic lobes. Only 12% of neurons cross hemispheres and 6% of neurons cross between the central brain and optic lobes (2, 80). Despite these bottlenecks, the brain is robustly interconnected with short path lengths. The large rich club of well-connected neurons may explain these close connections. When compared to the average neuron in the brain, rich club neurons are more likely to contain synapses in both hemispheres, and are also more likely to connect the op-

tic lobes to the central brain. The broad reach of these rich club neurons also keeps path lengths short across these bottlenecks. We may expect the same to be true regarding the ascending and descending neurons which form a bottleneck between the brain and the ventral nerve cord (VNC). Many ascending and descending neurons appear to have high degrees when examined either within the brain or within the VNC. While a wiring diagram of the VNC is now available (81), we await the completion of a complete CNS connectome to determine whether the ascending and descending neurons are members of the rich club.

Comparing connectomes across animals

Comparing network properties across wiring diagrams from different species has the potential to uncover global properties of brain organization. We make several such comparisons in **Table 2**, and have commented on other comparisons throughout the text. The similarities in reciprocity and clustering coefficient across animals, which vary dramatically in both size and connection density, hint at the possibility that some features of circuit architecture may be broadly conserved across biological nervous systems. Comparisons of metrics which are dependent on network topology, however, such as neuron degree or reciprocity and motif frequencies, must be interpreted with care due to differences in proofreading and data resolution. While connectomes in *C. elegans* have been proofread to the level of individual synapses (4–7), in larger connectomics datasets individual synapses are not proofread and instead a threshold on synapses per connection is applied to filter out spurious connections (1, 2, 8, 19). Threshold choice impacts topological metrics, which treat all edges as equivalent. Applying the same threshold across datasets does not resolve this conundrum, as a given number of synapses per connection may have different biological implications across species. It has also been observed, both in this work and in past studies, that different parts of the brain of the fly differ in their connectivity properties (17, 19). It is likely that the same is true in larger, more complex brains as well, meaning that statistics derived from partial wiring diagrams may not be representative.

It been demonstrated in *C. elegans* that there is substantial variability in the connectomes of individuals of the same species (6). Comparisons between the Flywire connectome and hemibrain wiring diagram have already revealed interesting similarities and differences between individual flies, as outlined in our companion paper (80), but more datasets will be needed before we fully understand the amount of variability between individuals in *Drosophila*. The same is expected to be true for zebrafish and mouse connectomes. More whole-brain connectomes are on the horizon, both in *Drosophila* and in other species (82). The network analysis of the fly brain presented here will be a valuable baseline for comparison, both to the connectomes of other *Drosophila* individuals and to the connectomes of other species. As the efficiency of electron microscopy and neural reconstruction continue to increase, it will become possible to better understand which features of these networks are common and which are species- or individual-specific. Such comparative connectomics studies within a single species may shed light on brain development, stereotypy, and learning, while future studies across multiple organisms may elucidate principles of brain evolution, organization, and computation.

Methods

Thresholding connections

All analyses presented in this paper were performed on the $v630$ Snapshot of the Flywire dataset. We chose to use a threshold of 5 synapses per connection between neurons for most of our analyses to reduce the impact of spurious connections. This threshold is also consistent across our companion papers on the Flywire connectome (2, 80). The synapses in the Flywire dataset were detected algorithmically (34, 35). We assessed key statistics as a function the threshold to ensure that our qualitative findings hold over a range of threshold choices (**Figure S1b-c**). We acknowledge that while some of these low synapse number connections may be spurious, it is also likely that a significant number of these low-count connections are real.

Assignment of neurotransmitters to neurons

The neurotransmitter at each synapse was predicted directly from the EM images using a trained convolutional neural network with per-synapse accuracy of 87% (2, 33). The algorithm returns a 1×6 probability vector containing the odds that a given synapse is each of the six primary neurotransmitters in *Drosophila*: ach, gaba, glut, da, oct, or ser. We then averaged these probabilities across all of a neuron's outgoing synapses, under the assumption that each neuron expresses a single outgoing neurotransmitter, to obtain a 1×6 probability vector representing the odds that a given neuron expresses a given neurotransmitter. We then assigned the highest-probability neurotransmitter as the putative neurotransmitter for that neuron. The per-neuron accuracy is 94%.

In cases where the highest probability is $p_1 < 0.2$ and the difference between the top two probabilities $p_1 - p_2 < 0.1$, we classified the neuron as having an uncertain neurotransmitter. In some cases, where the neurotransmitter of a neuron is previously known but differs from the neurotransmitter predicted by the algorithm, the neurotransmitter prediction associated with that neuron was overwritten by the known neurotransmitter.

Cell classifications and labels

84% of neurons are *intrinsic* to the brain, meaning that their projections are fully contained in the brain volume (2). Central brain neurons are fully contained in the central brain, while optic lobe intrinsic neurons are fully contained in the optic lobes. Visual projection neurons have inputs in the optic lobes and outputs in the central brain. Visual centrifugal neurons have inputs in the central brain and outputs in the optic lobe. Sensory neurons are those which are entering the brain from the periphery, and are divided into classes by modality. Refer to our companion paper, Schlegel et al., for more details on classification criteria (80). We also employed annotation labels contributed by the Flywire community (2).

Definitions of in-degree, out-degree, total degree and reciprocal degree

For a given neuron i, the in-degree d_i^+ is the number of incoming synaptic partners the neuron has and the out-degree d_i^- is the number of outgoing synaptic partners the neuron has. The total degree of a neuron i is the sum of in-degree and out-degree:

$$d_i^{\text{tot}} := d_i^+ + d_i^-. \tag{2}$$

The reciprocal degree d_i^{rec} is the number of partners a given neuron form reciprocal connections with. Since each reciprocal connection consists of two edges, we can determine the fraction of reciprocal inputs and outputs as d_i^{rec}/d_i^+ and d_i^{rec}/d_i^-, respectively (**Figure S2c**).

Definitions of connection probability, reciprocity and clustering coefficient

Given the observed wiring diagram as a simple (no self-edges) directed graph $G(V, E)$, the "connection probability" or "density" is the probability that, given an ordered pair of neurons α and β, a directed connection exists from one to the other:

$$p^{\text{conn}} := P[\alpha \to \beta] = \frac{|E|}{|V|(|V|-1)}. \tag{3}$$

The reciprocity is the probability that, given a pair of neurons which are connected α to β, there exists a returning β to α connection:

$$p^{\text{rec}} := P[\beta \to \alpha | \alpha \to \beta]. \tag{4}$$

The (global) clustering coefficient is the probability that for three neurons α, β and γ, given that neurons α and β are connected and neurons α and γ are connected (regardless of directionality), neurons β and γ are connected:

$$C^\Delta := P[\beta \sim \gamma | \alpha \sim \beta \wedge \alpha \sim \gamma]. \tag{5}$$

We computed these metrics both across the whole brain and within brain region (neuropil) subnetworks.

We also systematically quantified the occurrence of distinct directed 3-node motifs within the network, ensuring that duplicates are eliminated: any subgraph involving three unique nodes is counted only once in our analysis. To compute the expected prevalence of specific neurotransmitter motifs (**Figures 2c, 3d-e**) we multiplied the relevant neurotransmitter probabilities for the motif of interest, under the assumption the neurons connect independent of neurotransmitter. We then compared this expectation to the true frequency of motifs with these neurotransmitter combinations.

ER and CFG null models

We probed different statistics of the wiring diagram $G(V, E)$ by comparing them with the statistics of various null models. The simplest null model we employed was a directed version of the Erdős–Rényi model (ER) $\mathcal{G}(V, p)$, where all edges are drawn independently at random, and the connection probability p is set such that the expected number of edges in the ER model equals that observed in the wiring diagram (46). For any nodes $i, j \in V$, the connection probability is constant:

$$P[i \to j] = p = \frac{|E|}{|V|(|V|-1)}. \tag{6}$$

Since reciprocal edges in the wiring diagram are overrepresented when compared to a standard Erdős-Rényi (ER) model, we adopted a generalized Erdős–Rényi model (gER), which preserves the expected number of reciprocal edges. The gER model $\mathcal{G}(V, p^{\text{uni}}, p^{\text{bi}})$ has two parameters, uni-directional connection probability p^{uni} and bi-directional connection probability p^{bi}, both of which are set to match the wiring diagram. To do this, we defined the sets of unidirectional and bidirectional edges as:

$$E^{\text{uni}} := \{(i,j) | (i,j) \in E \wedge (j,i) \notin E\}, \tag{7}$$

$$E^{\text{bi}} := \{(i,j) | (i,j) \in E \wedge (j,i) \in E\}. \tag{8}$$

For any nodes i and j:

$$P[i \not\overset{\to}{\leftarrow} j] = P[i \overset{\leftarrow}{\not\to} j] = p^{\text{uni}} = \frac{|E^{\text{uni}}|}{|V|(|V|-1)}, \tag{9}$$

$$P[i \overset{\leftarrow}{\to} j] = p^{\text{bi}} = \frac{|E^{\text{bi}}|}{|V|(|V|-1)}, \tag{10}$$

$$P[i \overset{\not\leftarrow}{\not\to} j] = 1 - 2p^{\text{uni}} - p^{\text{bi}}. \tag{11}$$

All edges between unordered node pairs were drawn independently and at random.

In line with previous work (11, 60), we also employed a directed configuration model (CFG), $\mathcal{G}(V, \{d_i^+\}, \{d_i^-\})$, which preserves degree sequences during random rewiring. We sampled 1,000 random graphs uniformly from a configuration space of graphs with the same degree sequences as the observed graph by applying the switch-and-hold algorithm (83), where we randomly select two edges in each iteration and swap their target endpoints under the condition that doing so does not introduce self-loops or multiple edges (switch), or else keep them unchanged (hold).

Spectral analysis

Given a strongly connected graph $G(V, E)$ and its 0-1 adjacency matrix $A \in \mathbb{R}_{\geq 0}^{n \times n}$, where A_{ij} indicates the existence of a connection from neuron j to neuron i, one can construct an irreducible Markov chain on the strongly connected graph with a transition matrix $P_{ij} := A_{ij}/\sum_k A_{kj}$ giving the transition probability from j to i. The Perron-Frobenius theorem guarantees that P has a unique positive right eigenvector π with eigenvalue 1, and therefore that π is the stationary distribution of the Markov chain. We constructed such a transition matrix for the connectome and determined the eigenvector π.

We also defined a "reverse" Markov chain with a transition matrix $P_{ij}^{\text{rev}} := A_{ji}/\sum_k A_{jk}$ giving the transition probability from j to i. P^{rev} also has a unique positive right eigenvector π^{rev} with eigenvalue 1. **Figures S1d and S1e** show the stationary distribution of forward and reversed Markov chains, respectively.

The normalized symmetric Laplacian of the Markov chain P is

$$\mathcal{L} = I - \frac{1}{2}\left(\Pi^{1/2}P_\alpha\Pi^{-1/2} + \Pi^{-1/2}P^\intercal\Pi^{1/2}\right), \quad \textbf{(12)}$$

where $\Pi := \texttt{Diag}(\pi)$ and I is the identity matrix. Similarly, we defined $\mathcal{L}^{\text{rev}}$ for the reverse Markov chain. The eigen-spectra of $\mathcal{L}$ and $\mathcal{L}^{\text{rev}}$ are shown in **Figures S1d** and **S1e**, respectively. The gaps between eigenvalues indicate the conductance properties of the graph.

Finding rich club neurons

We employed the standard rich club formulation to quantify the rich club effect (25). The rich club coefficient $\Phi(k)$ at a given degree value (k), with all nodes with degree $< k$ pruned, is the number of existing connections in the surviving subnetwork divided by the total possible connections in the surviving subnetwork:

$$\Phi(k) = \frac{M_k}{N_k(N_k-1)}, \quad \textbf{(13)}$$

where N_k neurons in the network with degree $\geq k$ and M_k is the number of connections between such neurons.

To control for the fact that high-degree nodes have a higher probability of connecting to each other by chance, we normalized the rich club coefficient to the average rich club value of 5 samples from a CFG null model. (**Figure 1h**):

$$\Phi_{norm}(k) = \frac{\Phi(k)}{\langle\Phi_{CFG}(k)\rangle}. \quad \textbf{(14)}$$

We conducted only five samples because the rich club computation on such a large network is costly (the time complexity is $O(|V|\log|V|)$). The standard method of determining the rich club threshold is to look for values of k for which $\Phi_{norm}(k) > 1 + n\sigma$, where σ is the standard deviation of $\Phi_{CFG}(k)$ and n is chosen arbitrarily. However, since the standard deviation from our five samples is extremely small, we chose instead to define the onset and the offset thresholds of the rich club as $\Phi_{norm}(k) > 1.01$ (1% denser than the CFG random networks).

We computed the rich club coefficient in three different ways, by sweeping by total degree (Figure 1h), in-degree, and out-degree (Figure S1g), progressively moving from small to large values. As we observed, when the total degrees of the remaining nodes surpass 37, the network becomes denser compared to randomized networks. Once the minimal total degree reaches 93, the network becomes as sparse as the randomized counterpart. Therefore, we classified neurons with total degrees above 37 as "rich club" neurons because they exhibit denser interconnections when considered as a subnetwork. In terms of in-degree, the range for denser-than-random connectivity is between 10 and 54. Considering out-degree alone did not reveal any specific onset or offset threshold for rich club behavior, as the subnetwork always remains sparser than random.

Definitions of broadcaster neurons, integrator neurons, and neuropil-specific recurrent neurons

To identify broadcaster neurons, we filtered the intrinsic rich club neurons ($d^{\text{tot}} > 37$) for those which had an out-degree was at least 5 times higher than their in-degree:

$$d^- \geq 5 \times d^+. \quad \textbf{(15)}$$

Similarly, we identified integrator neurons by filtering the intrinsic rich club neurons for those which had an in-degree was at least 5 times higher than their out-degree:

$$d^+ \geq 5 \times d^-. \quad \textbf{(16)}$$

Rich club neurons which did not fall into either category were defined as "large balanced" neurons. This analysis was limited to intrinsic neurons–those which have all of their inputs and outputs within the brain–to avoid spurious identification of afferent or efferent neurons as broadcasters or integrators.

When identifying large recurrent neuropil-specific neurons (**Figure 5g**) we applied the following criteria. First, the neurons were intrinsic and met the rich club criteria. Second, at least 50% of the neuron's incoming connections were contained within the subnetwork of a single neuropil. Third, at least 50% of the neuron's outgoing connections were contained within the same neuropil.

Neuron ranking

We employed a probabilistic connectome flow model previously published in Schlegel et al. 2021 to determine the ranking of neurons relative to various sensory neuron populations (2, 20). This method ignores the sign of connections. Starting from a set of user-defined seed neurons, the model traverses the wiring diagram probabilistically: in each iteration the chance that a neuron is added to the traversed set increases linearly with the fractions of synapses it is receiving from neurons already in the traversed set. When this likelihood reaches 30%, the neuron is guaranteed to be added to the traversed set. The process is then repeated until the entire network graph has been traversed. The iteration in which a neuron was added corresponds to the distance in hops it was from the seed neurons. For each set of seed neurons, the model was run 10,000 times. The distance used to determine the rank of any given neuron was the average iteration in which it was added to the traversed set.

We ran this model using the following subsets of sensory neurons as seeds: olfactory receptor neurons, gustatory receptor neurons, mechanosensory Johnston's Organ neurons, head and neck bristle mechanosensory neurons, thermosensory neurons, hygrosensory neurons, visual projection neurons, visual photoreceptors, ocellar photoreceptors and ascending neurons. We also ran the model using the set of all of the input neurons as seed neurons. All neurons in the brain were then ranked by their traversal distance from each set of starting neurons, and this ranking was normalized to return a percentile rank.

Identifying neuropil subnetworks

Most of the neurons in the *Drosophila* brain have soma at the surface of the brain. Therefore, they cannot be associated to neuropils (brain regions) based on their soma locations. Synapses, however, can be associated with neuropils. To perform motif analyses at the level of individual neuropils, we identified neuropil subnetworks based on the the connections made

by the synapses contained within each neuropil volume. All connections within the neuropil of interest are taken as edges of this subnetwork, and all neurons connected to these edges are included (**Figure 5b**). The number of neurons associated with each neuropil subnetwork is plotted in **Figure S5d**. Note that if two neurons both in a given neuropil subnetwork share a connection which occurs in a different neuropil, that connection is not included as an edge in the given subnetwork.

Determining information flow between neuropils

To determine the contributions a single neuron makes to information flow between neuropils, we first applied two simplifying assumptions: (1) that information flow through the neuron can be approximated by the fraction of synapses in a given region and (2) that inputs and outputs can be treated independently. Employing these two assumptions we constructed a matrix representing the projections of a single neuron between neuropils. The fractional inputs of a given neuron are a $1 \times N$ vector containing the fraction of incoming synapses the neuron has in each of the N neuropils, and the fractional outputs are a similar vector containing the fraction of outgoing synapses in each of the N neuropils. We multiplied these vectors against each other to generate the $N \times N$ matrix of the neuron's fractional weights, with a total weight of one. Summing these matrices across all neurons produced a matrix of neuropil-to-neuropil connectivity, or projectome (see Figure 4 of Dorkenwald et al., 2023) (2).

From the neuropil-to-neuropil connectivity matrix we determined the total weight of internal connections—those within a given neuropil—by identifying the neurons which contribute to the diagonal of the matrix. We likewise determined the weight external connections—either incoming to the neuropil or outgoing from the neuropil—by looking at the off-diagonals. These data were used to construct the analyses in **Figure S5a-c**.

Code and software

The analyses presented in this paper were performed in Python with the numpy and graph-tool (84) packages, and in MATLAB (standard toolboxes). Some 3D renders were generated in Cinema4D. The Flywire data is available online via Codex (Connectome Data Explorer): codex.flywire.ai. Neuron annotations, neurotransmitter information, and compact data downloads are available via Codex, along with neuron lists generated in this work, including neurons participating in 2-node and selected 3-node motifs, rich club neurons, broadcaster and integrator neurons, and neuropil-specific reciprocal neurons.

AUTHOR CONTRIBUTIONS

AL, RY, and SD analyzed the data. SD, PS, and AM curated the data and made it available for download. AM developed software and data analysis tools. AM and ARS built the Codex online platform. SCY, CEM, MC, KE, and PS trained and managed Flywire proofreaders. ASB, NE, GSXEJ and JF provided neurotransmitter information. CEM and ARS provided Flywire community support and training. AL, RY, SD, ARS, and PS generated figure panels. AL, RY, and MM wrote the manuscript with feedback from the other authors. MM supervised the project.

ACKNOWLEDGEMENTS

We thank Sebastian Seung, Davi Bock and the members of the Murthy, Seung, Jefferis, and Clandinin labs for their advice on the project and comments on the manuscript. AL was supported by the NSF through the Center for the Physics of Biological Function (PHY-1734030). GSXEJ was supported by Wellcome Trust Collaborative Awards 203261/Z/16/Z and 220343/Z/20/Z, the Neuronex2 award (MRC_MC_EX_MR/T046279/1), and the MRC (MC-U105188491). MM was supported by NIH BRAIN Initiative grants RF1 MH117815, RF1 MH129268 and U24 NS126935. We also acknowledge support from the Princeton Neuroscience Institute and assistance from Google.

References

1. Sven Dorkenwald, Claire Mckellar, Thomas Macrina, Nico Kemnitz, Kisuk Lee, Jingpeng Wu, Sergiy Popovych, Alexander Bae, Eric Mitchell, Barak Nehoran, Zhen Jia, Jonathan Zung, Derrick Brittain, Forrest Collman, Chris Jordan, William Silversmith, Christa Baker, David Deutsch, Sandeep Kumar, Austin Burke, Jay Gager, James Hebditch, Merlin Moore, Sarah Morejohn, Ben Silverman, Kyle Willie, Ryan Willie, Mala Murthy, and H Sebastian Seung. Flywire: Online community for whole-brain connectomics. *Nature Methods*, 19:119–128, 2022.

2. Sven Dorkenwald, Arie Matsliah, Amy R Sterling, Philipp Schlegel, Szi chieh Yu, Claire E. McKellar, Albert Lin, Marta Costa, Katharina Eichler, Yijie Yin, Will Silversmith, Casey Schneider-Mizell, Chris S Jordan, Derrick Brittain, Akhilesh Halageri, Kai Kuehner, Oluwaseun Ogedengbe, Ryan Morey, Jay Gager, Krzysztof Kruk, Eric Perlman, Runzhe Yang, David Deutsch, Doug Bland, Marissa Sorek, Ran Lu, Thomas Macrina, Kisuk Lee, J. Alexander Bae, Shang Mu, Barak Nehoran, Eric Mitchell, Sergiy Popovych, Jingpeng Wu, Zhen Jia, Manuel Castro, Nico Kemnitz, Dodam Ih, Alexander Shakeel Bates, Nils Eckstein, Jan Funke, Forrest Collman, Davi D Bock, Gregory S X E Jefferis, H. Sebastian Seung, Mala Murthy, and the FlyWire Consortium. Neuronal wiring diagram of an adult brain. *bioRxiv*, 2023. doi: 10.1101/2023.06.27.546656.

3. Zhihao Zheng, J. Scott Lauritzen, Eric Perlman, Camenzind G. Robinson, Matthew Nichols, Daniel Milkie, Omar Torrens, John Price, Corey B. Fisher, Nadiya Sharifi, Steven A. Calle-Schuler, Lucia Kmecova, Iqbal J. Ali, Bill Karsh, Eric T. Trautman, John A. Bogovic, Philipp Hanslovsky, Gregory S.X.E. Jefferis, Michael Kazhdan, Khaled Khairy, Stephan Saalfeld, Richard D. Fetter, and Davi D. Bock. A complete electron microscopy volume of the brain of adult drosophila melanogaster. *Cell*, 174(3):730–743.e22, 2018. ISSN 0092-8674. doi: https://doi.org/10.1016/j.cell.2018.06.019.

4. John G White, E Southgate, J N Thomson, and Sydney Brenner. The structure of the nervous system of the nematode Caenorhabditis elegans. *Philosophical Transactions of the Royal Society of London B*, 314:1–340, 1986.

5. Steven J Cook, Travis A Jarrell, Christopher A Brittin, Yi Wang, Adam E Bloniarz, Maksim A Yakovlev, Ken CQ Nguyen, Leo T-H Tang, Emily A Bayer, Janet S Duerr, et al. Whole-animal connectomes of both caenorhabditis elegans sexes. *Nature*, 571(7763):63–71, 2019.

6. Daniel Witvliet, Ben Mulcahy, James K. Mitchell, Yaron Meirovitch, Daniel R. Berger, Yuelong Wu, Yufang Liu, Wan Xian Koh, Rajeev Parvathala, Douglas Holmyard, Richard L. Schalek, Nir Shavit, Andrew D. Chisholm, Jeff W. Lichtman, Aravinthan D.T. Samuel, and Mei Zhen. Connectomes across development reveal principles of brain maturation. *Nature*, 596(7871): 257–261, 2021. ISSN 14764687. doi: 10.1038/s41586-021-03778-8.

7. Lav R. Varshney, Beth L. Chen, Eric Paniagua, David H. Hall, and Dmitri B. Chklovskii. Structural properties of the Caenorhabditis elegans neuronal network. *PLoS Comput. Biol.*, 7(2), 2011. ISSN 1553734X. doi: 10.1371/journal.pcbi.1001066.

8. Michael Winding, Benjamin D. Pedigo, Christopher L. Barnes, Heather G. Patsolic, Youngser Park, Tom Kazimiers, Akira Fushiki, Ingrid V. Andrade, Avinash Khandelwal, Javier Valdes-Aleman, Feng Li, Nadine Randel, Elizabeth Barsotti, Ana Correia, Richard D. Fetter, Volker Hartenstein, Carey E. Priebe, Joshua T. Vogelstein, Albert Cardona, and Marta Zlatic. The connectome of an insect brain. *Science*, 379(6636):eadd9330, 2023. doi: 10.1126/science.add9330.

9. Ashwin Vishwanathan, Alexandro D. Ramirez, Jingpeng Wu, Alex Sood, Runzhe Yang, Nico Kemnitz, Dodam Ih, Nicholas Turner, Kisuk Lee, Ignacio Tartavull, William M. Silversmith, Chris S. Jordan, Celia David, Doug Bland, Mark S. Goldman, Emre R. F. Aksay, H. Sebastian Seung, and the Eyewirers. Predicting modular functions and neural coding of behavior from a synaptic wiring diagram. *bioRxiv*, 2022. doi: 10.1101/2020.10.28.359620.

10. Adrian A. Wanner and Rainer W. Friedrich. Whitening of odor representations by the wiring diagram of the olfactory bulb. *Nature Neuroscience 2020 23:3*, 23:433–442, 1 2020. ISSN 1546-1726. doi: 10.1038/s41593-019-0576-z.

11. Runzhe Yang, Ashwin Vishwanathan, Jingpeng Wu, Nico Kemnitz, Dodam Ih, Nicholas Turner, Kisuk Lee, Ignacio Tartavull, William M. Silversmith, Chris S. Jordan, Celia David, Doug Bland, Amy Sterling, Mark S. Goldman, Emre R. F. Aksay, H. Sebastian Seung, and the EyeWirers. Cyclic structure with cellular precision in a vertebrate sensorimotor neural circuit. *Current Biology*, 2023.

12. Davi D Bock, Wei-Chung Allen Lee, Aaron M Kerlin, Mark L Andermann, Greg Hood, Arthur W Wetzel, Sergey Yurgenson, Edward R Soucy, Hyon Suk Kim, and R Clay Reid. Network anatomy and in vivo physiology of visual cortical neurons. *Nature*, 471(7337):177–182, 2011.

13. Wei Chung Allen Lee, Vincent Bonin, Michael Reed, Brett J. Graham, Greg Hood, Katie Glattfelder, and R. Clay Reid. Anatomy and function of an excitatory network in the visual cortex. *Nature 2016 532:7599*, 532:370–374, 3 2016. ISSN 1476-4687. doi: 10.1038/nature17192.

14. Nicholas L Turner, Thomas Macrina, J Alexander Bae, Runzhe Yang, Alyssa M Wilson, Casey Schneider-Mizell, Kisuk Lee, Ran Lu, Jingpeng Wu, Agnes L Bodor, et al. Reconstruction of neocortex: Organelles, compartments, cells, circuits, and activity. *Cell*, 185(6):1082–1100, 2022.

15. Shin Ya Takemura, Arjun Bharioke, Zhiyuan Lu, Aljoscha Nern, Shiv Vitaladevuni, Patricia K. Rivlin, William T. Katz, Donald J. Olbris, Stephen M. Plaza, Philip Winston, Ting Zhao, Jane Anne Horne, Richard D. Fetter, Satoko Takemura, Katerina Blazek, Lei Ann Chang, Omotara Ogundeyi, Mathew A. Saunders, Victor Shapiro, Christopher Sigmund, Gerald M. Rubin, Louis K. Scheffer, Ian A. Meinertzhagen, and Dmitri B. Chklovskii. A visual motion detection circuit suggested by drosophila connectomics. *Nature 2013 500:7461*, 500:175–181, 8 2013. ISSN 1476-4687. doi: 10.1038/nature12450.

16. Shin-ya Takemura, Yoshinori Aso, Toshihide Hige, Allan Wong, Zhiyuan Lu, C Shan Xu, Patricia K Rivlin, Harald Hess, Ting Zhao, Toufiq Parag, Stuart Berg, Gary Huang, William Katz, Donald J Olbris, Stephen Plaza, Lowell Umayam, Roxanne Aniceto, Lei-Ann Chang, Shirley Lauchie, Omotara Ogundeyi, Christopher Ordish, Aya Shinomiya, Christopher Sigmund, Satoko Takemura, Julie Tran, Glenn C Turner, Gerald M Rubin, and Louis K Scheffer. A connectome of

a learning and memory center in the adult *Drosophila* brain. *eLife*, 6:e26975, jul 2017. ISSN 2050-084X. doi: 10.7554/eLife.26975.

17. Jane Anne Horne, Carlie Langille, Sari McLin, Meagan Wiederman, Zhiyuan Lu, C Shan Xu, Stephen M Plaza, Louis K Scheffer, Harald F Hess, and Ian A Meinertzhagen. A resource for the *Drosophila* antennal lobe provided by the connectome of glomerulus va1v. *eLife*, 7:e37550, nov 2018. ISSN 2050-084X. doi: 10.7554/eLife.37550.

18. Feng Li, Jack Lindsey, Elizabeth C Marin, Nils Otto, Marisa Dreher, Georgia Dempsey, Ildiko Stark, Alexander Shakeel Bates, Markus William Pleijzier, Philipp Schlegel, Aljoscha Nern, Shinya Takemura, Nils Eckstein, Tansy Yang, Audrey Francis, Amalia Braun, Ruchi Parekh, Marta Costa, Louis Scheffer, Yoshinori Aso, Gregory S.X.E. Jefferis, L. F. Abbott, Ashok Litwin-Kumar, Scott Waddell, and Gerald M Rubin. The connectome of the adult drosophila mushroom body provides insights into function. *eLife*, 9:1–217, 2020. ISSN 2050084X. doi: 10.7554/eLife.62576.

19. Louis K. Scheffer, C. Shan Xu, Michal Januszewski, Zhiyuan Lu, Shin Ya Takemura, Kenneth J. Hayworth, Gary B. Huang, Kazunori Shinomiya, Jeremy Maitin-Shepard, Stuart Berg, Jody Clements, Philip M. Hubbard, William T. Katz, Lowell Umayam, Ting Zhao, David Ackerman, Tim Blakely, John Bogovic, Tom Dolafi, Dagmar Kainmueller, Takashi Kawase, Khaled A. Khairy, Laramie Leavitt, Peter H. Li, Larry Lindsey, Nicole Neubarth, Donald J. Olbris, Hideo Otsuna, Eric T. Trautman, Masayoshi Ito, Alexander S. Bates, Jens Goldammer, Tanya Wolff, Robert Svirskas, Philipp Schlegel, Erika R. Neace, Christopher J. Knecht, Chelsea X. Alvarado, Dennis A. Bailey, Samantha Ballinger, Jolanta A. Borycz, Brandon S. Canino, Natasha Cheatham, Michael Cook, Marisa Dreher, Octave Duclos, Bryon Eubanks, Kelli Fairbanks, Samantha Finley, Nora Forknall, Audrey Francis, Gary Patrick Hopkins, Emily M. Joyce, Sungjin Kim, Nicole A. Kirk, Julie Kovalyak, Shirley A. Lauchie, Alanna Lohff, Charli Maldonado, Emily A. Manley, Sari McLin, Caroline Mooney, Miatta Ndama, Omotara Ogundeyi, Nneoma Okeoma, Christopher Ordish, Nicholas Padilla, Christopher Patrick, Tyler Paterson, Elliott E. Phillips, Emily M. Phillips, Neha Rampally, Caitlin Ribeiro, Madelaine K. Robertson, Jon Thomson Rymer, Sean M. Ryan, Megan Sammons, Anne K. Scott, Ashley L. Scott, Aya Shinomiya, Claire Smith, Kelsey Smith, Natalie L. Smith, Margaret A. Sobeski, Alia Suleiman, Jackie Swift, Satoko Takemura, Iris Talebi, Dorota Tarnogorska, Emily Tenshaw, Temour Tokhi, John J. Walsh, Tansy Yang, Jane Anne Horne, Feng Li, Ruchi Parekh, Patricia K. Rivlin, Vivek Jayaraman, Marta Costa, Gregory S.X.E. Jefferis, Kei Ito, Stephan Saalfeld, Reed George, Ian A. Meinertzhagen, Gerald M. Rubin, Harald F. Hess, Viren Jain, and Stephen M. Plaza. A connectome and analysis of the adult drosophila central brain. *eLife*, 9:1–74, 2020. ISSN 2050084X. doi: 10.7554/ELIFE.57443.

20. Philipp Schlegel, Alexander Shakeel Bates, Tomke Stürner, Sridhar R Jagannathan, Nikolas Drummond, Joseph Hsu, Laia Serratosa Capdevila, Alexandre Javier, Elizabeth C Marin, Asa Barth-Maron, Imaan FM Tamimi, Feng Li, Gerald M Rubin, Stephen M Plaza, Marta Costa, and Gregory S X E Jefferis. Information flow, cell types and stereotypy in a full olfactory connectome. *eLife*, 10:e66018, may 2021. ISSN 2050-084X. doi: 10.7554/eLife.66018.

21. Brad K Hulse, Hannah Haberkern, Romain Franconville, Daniel Turner-Evans, Shin-ya Takemura, Tanya Wolff, Marcella Noorman, Marisa Dreher, Chuntao Dan, Ruchi Parekh, Ann M Hermundstad, Gerald M Rubin, and Vivek Jayaraman. A connectome of the *Drosophila* central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. *eLife*, 10:e66039, oct 2021. ISSN 2050-084X. doi: 10.7554/eLife.66039.

22. Daniel M Busiello, Samir Suweis, Jorge Hidalgo, and Amos Maritan. Explorability and the origin of network sparsity in living systems. *Scientific reports*, 7(1):12323, 2017.

23. Vito Latora and Massimo Marchiori. Efficient behavior of small-world networks. *Physical review letters*, 87(19):198701, 2001.

24. Mark D Humphries and Kevin Gurney. Network 'small-world-ness': a quantitative method for determining canonical network equivalence. *PloS one*, 3(4):e0002051, 2008.

25. Vittoria Colizza, Alessandro Flammini, M Angeles Serrano, and Alessandro Vespignani. Detecting rich-club ordering in complex networks. *Nature physics*, 2(2):110–115, 2006.

26. Emma K. Towlson, Petra E. Vértes, Sebastian E. Ahnert, William R. Schafer, and Edward T. Bullmore. The rich club of the c. elegans neuronal connectome. *Journal of Neuroscience*, 33 (15):6380–6387, 2013. ISSN 0270-6474. doi: 10.1523/JNEUROSCI.3784-12.2013.

27. Sen Song, Per Jesper Sjöström, Markus Reigl, Sacha Nelson, and Dmitri B Chklovskii. Highly nonrandom features of synaptic connectivity in local cortical circuits. *PLoS biology*, 3(3):e68, 2005.

28. Rodrigo Perin, Thomas K Berger, and Henry Markram. A synaptic organizing principle for cortical neuronal groups. *Proceedings of the National Academy of Sciences*, 108(13):5419–5424, 2011.

29. Daniel Udvary, Philipp Harth, Jakob H. Macke, Hans Christian Hege, Christiaan P.J. de Kock, Bert Sakmann, and Marcel Oberlaender. The impact of neuron morphology on cortical network architecture. *Cell Reports*, 39, 4 2022. ISSN 22111247. doi: 10.1016/j.celrep.2022.110677.

30. R. Milo, S. Shen-Orr, S. Itzkovitz, N. Kashtan, D. Chklovskii, and U. Alon. Network motifs: Simple building blocks of complex networks. *Science*, 298(5594):824–827, 2002. doi: 10.1126/science.298.5594.824.

31. Olaf Sporns and Rolf Kötter. Motifs in brain networks. *PLOS Biology*, 2(11):null, 10 2004. doi: 10.1371/journal.pbio.0020369.

32. James E Ferrell Jr. Feedback loops and reciprocal regulation: recurring motifs in the systems biology of the cell cycle. *Current opinion in cell biology*, 25(6):676–686, 2013.

33. Nils Eckstein, Alexander Shakeel Bates, Andrew Champion, Michelle Du, Yijie Yin, Philipp Schlegel, Alicia Kun-Yang Lu, Thomson Rymer, Samantha Finley-May, Tyler Paterson, Ruchi Parekh, Sven Dorkenwald, Arie Matsliah, Szi-Chieh Yu, Claire McKellar, Amy Sterling, Katharina Eichler, Marta Costa, Sebastian Seung, Mala Murthy, Volker Hartenstein, Gregory S.X.E. Jefferis, and Jan Funke. Neurotransmitter classification from electron microscopy images at synaptic sites in drosophila melanogaster. *bioRxiv*, 2023. doi: 10.1101/2020.06.12.148775.

34. Larissa Heinrich, Jan Funke, Constantin Pape, Juan Nunez-Iglesias, and Stephan Saalfeld. Synaptic cleft segmentation in non-isotropic volume electron microscopy of the complete drosophila brain. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 11071 LNCS:317–325, 2018. ISSN 16113349. doi: 10.1007/978-3-030-00934-2_36/COVER.

35. Julia Buhmann, Arlo Sheridan, Caroline Malin-Mayor, Philipp Schlegel, Stephan Gerhard, Tom Kazimiers, Renate Krause, Tri M. Nguyen, Larissa Heinrich, Wei Chung Allen Lee, Rachel Wilson, Stephan Saalfeld, Gregory S.X.E. Jefferis, Davi D. Bock, Srinivas C. Turaga, Matthew Cook, and Jan Funke. Automatic detection of synaptic partners in a whole-brain drosophila electron

microscopy data set. *Nature Methods 2021 18:7*, 18:771–774, 6 2021. ISSN 1548-7105. doi: 10.1038/s41592-021-01183-7.

36. Wendy W. Liu and Rachel I. Wilson. Glutamate is an inhibitory neurotransmitter in the drosophila olfactory system. *Proceedings of the National Academy of Sciences of the United States of America*, 110:10294–10299, 6 2013. ISSN 00278424. doi: 10.1073/PNAS.1220560110/SUPPL_FILE/PNAS.201220560SI.PDF.

37. Ellena v. McCarthy, Ying Wu, Tagide deCarvalho, Christian Brandt, Guan Cao, and Michael N. Nitabach. Synchronized bilateral synaptic inputs to drosophila melanogaster neuropeptidergic rest/arousal neurons. *Journal of Neuroscience*, 31(22):8181–8193, 2011. ISSN 0270-6474. doi: 10.1523/JNEUROSCI.2017-10.2011.

38. Sebastian Molina-Obando, Juan Felipe Vargas-Fique, Miriam Henning, Burak Gür, T Moritz Schladt, Junaid Akhtar, Thomas K Berger, and Marion Silies. On selectivity in the *Drosophila* visual system is a multisynaptic process involving both glutamatergic and gabaergic inhibition. *eLife*, 8:e49373, sep 2019. ISSN 2050-084X. doi: 10.7554/eLife.49373.

39. Robert Tarjan. Depth-first search and linear graph algorithms. *SIAM journal on computing*, 1 (2):146–160, 1972.

40. Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 770–778, 2016.

41. David F Gleich. Pagerank beyond the web. *siam REVIEW*, 57(3):321–363, 2015.

42. Martijn P Van Den Heuvel and Olaf Sporns. Rich-club organization of the human connectome. *Journal of Neuroscience*, 31(44):15775–15786, 2011.

43. Wilfrid Rall, G.M. Shepherd, T.S. Reese, and M.W. Brightman. Dendrodendritic synaptic pathway for inhibition in the olfactory bulb. *Experimental Neurology*, 14(1):44–56, 1966. ISSN 0014-4886. doi: https://doi.org/10.1016/0014-4886(66)90023-9.

44. Duncan J Watts and Steven H Strogatz. Collective dynamics of 'small-world'networks. *nature*, 393(6684):440–442, 1998.

45. John Guare. *Six degrees of separation: A play*. Vintage, 1990.

46. Edgar N Gilbert. Random graphs. *The Annals of Mathematical Statistics*, 30(4):1141–1144, 1959.

47. Alexander Arenz, Michael S. Drews, Florian G. Richter, Georg Ammer, and Alexander Borst. The temporal tuning of the drosophila motion detectors is determined by the dynamics of their input elements. *Current Biology*, 27:929–944, 4 2017. ISSN 09609822. doi: 10.1016/j.cub.2017.01.051.

48. Kei Ito, Kazunori Shinomiya, Masayoshi Ito, J Douglas Armstrong, George Boyan, Volker Hartenstein, Steffen Harzsch, Martin Heisenberg, Uwe Homberg, Arnim Jenett, Haig Keshishian, Linda L Restifo, Wolfgang Rössler, Julie H Simpson, Nicholas J Strausfeld, Roland Strauss, and Leslie B Vosshall. A systematic nomenclature for the insect brain. *Neuron*, 81: 755–765, 2014. ISSN 08966273. doi: 10.1016/j.neuron.2013.12.017.

49. Xu Liu and Ronald L. Davis. The gabaergic anterior paired lateral neuron suppresses and is suppressed by olfactory learning. *Nature Neuroscience 2008 12:1*, 12:53–59, 11 2008. ISSN 1546-1726. doi: 10.1038/nn.2235.

50. Yanying Wu, Qingzhong Ren, Hao Li, and Aike Guo. The gabaergic anterior paired lateral neurons facilitate olfactory reversal learning in drosophila. *Learning & Memory*, 19:478–486, 10 2012. ISSN 1072-0502. doi: 10.1101/LM.025726.112.

51. Kazunori Shinomiya, Shin Ya Takemura, Patricia K. Rivlin, Stephen M. Plaza, Louis K. Scheffer, and Ian A. Meinertzhagen. A common evolutionary origin for the on- and off-edge motion detection pathways of the drosophila visual system. *Frontiers in Neural Circuits*, 9:152337, 7 2015. ISSN 16625110. doi: 10.3389/FNCIR.2015.00033/BIBTEX.

52. Kazunori Shinomiya, Gary Huang, Zhiyuan Lu, Toufiq Parag, C. Shan Xu, Roxanne Aniceto, Namra Ansari, Natasha Cheatham, Shirley Lauchie, Erika Neace, Omotara Ogundeyi, Christopher Ordish, David Peel, Aya Shinomiya, Claire Smith, Satoko Takemura, Iris Talebi, Patricia K. Rivlin, Aljoscha Nern, Louis K. Scheffer, Stephen M. Plaza, and Ian A. Meinertzhagen. Comparisons between the on-and off-edge motion pathways in the drosophila brain. *eLife*, 8, 2019. ISSN 2050084X. doi: 10.7554/ELIFE.40025.

53. Shawn R. Olsen and Rachel I. Wilson. Lateral presynaptic inhibition mediates gain control in an olfactory circuit. *Nature 2008 452:7190*, 452:956–960, 3 2008. ISSN 1476-4687. doi: 10.1038/nature06864.

54. Cory M. Root, Kaoru Masuyama, David S. Green, Lina E. Enell, Dick R. Nässel, Chi Hon Lee, and Jing W. Wang. A presynaptic gain control mechanism fine-tunes olfactory behavior. *Neuron*, 59:311–321, 7 2008. ISSN 08966273. doi: 10.1016/j.neuron.2008.07.003.

55. Matthias Meier and Alexander Borst. Extreme compartmentalization in a drosophila amacrine cell. *Current Biology*, 29:1545–1550.e2, 5 2019. ISSN 09609822. doi: 10.1016/j.cub.2019.03.070.

56. Hoger Amin, Anthi A Apostolopoulou, Raquel Suárez-Grimalt, Eleftheria Vrontou, and Andrew C Lin. Localized inhibition in the *Drosophila* mushroom body. *eLife*, 9:e56954, sep 2020. ISSN 2050-084X. doi: 10.7554/eLife.56954.

57. Jonathan E. Schenk and Quentin Gaudry. Nonspiking interneurons in the drosophila antennal lobe exhibit spatially restricted activity. *eNeuro*, 10, 1 2023. ISSN 2373-2822. doi: 10.1523/ENEURO.0109-22.2022.

58. Christa A. Baker, Claire McKellar, Rich Pang, Aljoscha Nern, Sven Dorkenwald, Diego A. Pacheco, Nils Eckstein, Jan Funke, Barry J. Dickson, and Mala Murthy. Neural network organization for courtship-song feature detection in drosophila. *Current Biology*, 32: 3317–3333.e7, 8 2022. ISSN 18790445. doi: 10.1016/J.CUB.2022.06.019/ATTACHMENT/16F13A9C-98D3-4D28-AC95-31F85990C9AE/MMC3.XLSX.

59. Diego A Pacheco, Stephan Y Thiberge, Eftychios Pnevmatikakis, and Mala Murthy. Auditory activity is diverse and widespread throughout the central brain of drosophila. *Nat Neurosci*, 24: 93–104, 2021. doi: https://doi.org/10.1038/s41593-020-00743-y.

60. Maxwell H Turner, Kevin Mann, and Thomas R Clandinin. The connectome predicts resting-state functional connectivity across the drosophila brain. *Current Biology*, 31(11):2386–2394, 2021.

61. Luke E. Brezovec, Andrew B. Berger, Shaul Druckmann, and Thomas R. Clandinin. Mapping the neural dynamics of locomotion across the drosophila brain. *bioRxiv*, 2022. doi: 10.1101/2022.03.20.485047.

62. Kelsey M Hallinen, Ross Dempsey, Monika Scholz, Xinwei Yu, Ashley Linder, Francesco Randi, Anuj K Sharma, Joshua W Shaevitz, and Andrew M Leifer. Decoding locomotion from population

neural activity in moving *C. elegans*. *eLife*, 10:e66135, jul 2021. ISSN 2050-084X. doi: 10.7554/eLife.66135.

63. Eviatar Yemini, Albert Lin, Amin Nejatbakhsh, Erdem Varol, Ruoxi Sun, Gonzalo E. Mena, Aravinthan D.T. Samuel, Liam Paninski, Vivek Venkatachalam, and Oliver Hobert. NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal Identification in C. elegans. *Cell*, 184:272–288, 2021. doi: 10.1016/j.cell.2020.12.012.

64. Timothy W Dunn, Yu Mu, Sujatha Narayan, Owen Randlett, Eva A Naumann, Chao Tsung Yang, Alexander F Schier, Jeremy Freeman, Florian Engert, and Misha B Ahrens. Brain-wide mapping of neural activity controlling zebrafish exploratory locomotion. *eLife*, 5(MARCH2016): 1–29, 2016. ISSN 2050084X. doi: 10.7554/eLife.12741.

65. Martin Haesemeyer, Drew N Robson, Jennifer M Li, Alexander F Schier, and Florian Engert. A Brain-wide Circuit Model of Heat-Evoked Swimming Behavior in Larval Zebrafish. *Neuron*, 98 (4):817–831, 2018. ISSN 10974199. doi: 10.1016/j.neuron.2018.04.013.

66. Yu Mu, Davis V. Bennett, Mikail Rubinov, Sujatha Narayan, Chao Tsung Yang, Masashi Tanimoto, Brett D. Mensh, Loren L. Looger, and Misha B. Ahrens. Glia Accumulate Evidence that Actions Are Futile and Suppress Unsuccessful Behavior. *Cell*, 178(1):27–43, 2019. ISSN 10974172. doi: 10.1016/j.cell.2019.05.050.

67. Simon Musall, Matthew T. Kaufman, Ashley L. Juavinett, Steven Gluf, and Anne K. Churchland. Single-trial neural dynamics are dominated by richly varied movements. *Nature Neuroscience 2019 22:10*, 22:1677–1686, 9 2019. ISSN 1546-1726. doi: 10.1038/s41593-019-0502-4.

68. Sarah L. West, Justin D. Aronson, Laurentiu S. Popa, Kathryn D. Feller, Russell E. Carter, William M. Chiesl, Morgan L. Gerhart, Aditya C. Shekhar, Leila Ghanbari, Suhasa B. Kodandaramaiah, and Timothy J. Ebner. Wide-field calcium imaging of dynamic cortical networks during locomotion. *Cerebral Cortex*, 32:2668–2687, 6 2022. ISSN 1047-3211. doi: 10.1093/CERCOR/BHAB373.

69. Albert Lin, Daniel Witvliet, Luis Hernandez-Nunez, Scott W. Linderman, Aravinthan D. T. Samuel, and Vivek Venkatachalam. Imaging whole-brain activity to understand behaviour. *Nature Reviews Physics*, 0123456789, 2022. doi: 10.1038/s42254-022-00430-w.

70. Johannes D. Seelig and Vivek Jayaraman. Neural dynamics for landmark orientation and angular path integration. *Nature 2015 521:7551*, 521:186–191, 5 2015. ISSN 1476-4687. doi: 10.1038/nature14446.

71. Daniel Turner-Evans, Stephanie Wegener, Hervé Rouault, Romain Franconville, Tanya Wolff, Johannes D. Seelig, Shaul Druckmann, and Vivek Jayaraman. Angular velocity integration in a fly heading circuit. *eLife*, 6, 5 2017. ISSN 2050084X. doi: 10.7554/ELIFE.23496.

72. Jonathan Green, Atsuko Adachi, Kunal K. Shah, Jonathan D. Hirokawa, Pablo S. Magani, and Gaby Maimon. A neural circuit architecture for angular integration in drosophila. *Nature 2017 546:7656*, 546:101–106, 5 2017. ISSN 1476-4687. doi: 10.1038/nature22343.

73. Sung Soo Kim, Ann M. Hermundstad, Sandro Romani, L. F. Abbott, and Vivek Jayaraman. Generation of stable heading representations in diverse visual scenes. *Nature 2019 576:7785*, 576:126–131, 11 2019. ISSN 1476-4687. doi: 10.1038/s41586-019-1767-1.

74. Yvette E. Fisher, Jenny Lu, Isabel D'Alessandro, and Rachel I. Wilson. Sensorimotor experience remaps visual input to a heading-direction network. *Nature 2019 576:7785*, 576:121–125, 11 2019. ISSN 1476-4687. doi: 10.1038/s41586-019-1772-4.

75. Tatsuo S. Okubo, Paola Patella, Isabel D'Alessandro, and Rachel I. Wilson. A neural network for wind-guided compass navigation. *Neuron*, 107:924–940.e18, 9 2020. ISSN 10974199. doi: 10.1016/J.NEURON.2020.06.022/ATTACHMENT/C924EF477-E422-4151-BD5C-593A3D2B7916/MMC1.PDF.

76. Jakob Troidl, Simon Warchol, Jinhan Choi, Jordan Matelsky, Nagaraju Dhanyasi, Xueying Wang, Brock Wester, Donglai Wei, Jeff W. Lichtman, Hanspeter Pfister, and Johanna Beyer. Vimo visual analysis of neuronal connectivity motifs. *bioRxiv*, 2022. doi: 10.1101/2022.12.09.519772.

77. Paul H. Taghert and Michael N. Nitabach. Peptide neuromodulation in invertebrate model systems. *Neuron*, 76:82–97, 10 2012. ISSN 0896-6273. doi: 10.1016/J.NEURON.2012.08.035.

78. Dick R. Nässel and Meet Zandawala. Recent advances in neuropeptide signaling in drosophila, from genes to physiology and behavior. *Progress in Neurobiology*, 179:101607, 8 2019. ISSN 0301-0082. doi: 10.1016/J.PNEUROBIO.2019.02.003.

79. Dick R. Nässel and Meet Zandawala. Endocrine cybernetics: neuropeptides as molecular switches in behavioural decisions. *Open Biology*, 12, 7 2022. ISSN 20462441. doi: 10.1098/RSOB.220174.

80. Philipp Schlegel, Yijie Yin, Alexander Shakeel Bates, Sven Dorkenwald, Katharina Eichler, Paul Brooks, Daniel S Han, Marina Gkantia, Marcia dos Santos, Eva J Munnelly, Griffin Badalamente, Laia Serratosa Capdevila, Varun Aniruddha Sane, Markus William Pleijzier, Imaan F M Tamimi, Christopher R Dunne, Irene Salgarella, Alexandre Javier, Siqi Fang, Eric Perlman, Tom Kazimiers, Sridhar R Jagannathan, Arie Matsliah, Amy R Sterling, Szi chieh Yu, Claire E McKellar, FlyWire Consortium, Marta Costa, H. Sebastian Seung, Mala Murthy, Volker Hartenstein, Davi D Bock, and Gregory S X E Jefferis. A consensus cell type atlas from multiple connectomes reveals principles of circuit stereotypy and variation. *bioRxiv*, 2023. doi: 10.1101/2023.06.27.546055.

81. Shin ya Takemura, Kenneth J Hayworth, Gary B Huang, Michal Januszewski, Zhiyuan Lu, Elizabeth C Marin, Stephan Preibisch, C Shan Xu, John Bogovic, Andrew S Champion, Han SJ Cheong, Marta Costa, Katharina Eichler, William Katz, Christopher Knecht, Feng Li, Billy J Morris, Christopher Ordish, Patricia K Rivlin, Philipp Schlegel, Kazunori Shinomiya, Tomke Stürner, Ting Zhao, Griffin Badalamente, Dennis Bailey, Paul Brooks, Brandon S Canino, Jody Clements, Michael Cook, Octave Duclos, Christopher R Dunne, Kelli Fairbanks, Siqi Fang, Samantha Finley-May, Audrey Francis, Reed George, Marina Gkantia, Kyle Harrington, Gary Patrick Hopkins, Joseph Hsu, Philip M Hubbard, Alexandre Javier, Dagmar Kainmueller, Wyatt Korff, Julie Kovalyak, Dominik Krzemiński, Shirley A Lauchie, Alanna Lohff, Charli Maldonado, Emily A Manley, Caroline Mooney, Erika Neace, Matthew Nichols, Omotara Ogundeyi, Nneoma Okeoma, Tyler Paterson, Elliott Phillips, Emily M Phillips, Caitlin Ribeiro, Sean M Ryan, Jon Thomson Rymer, Anne K Scott, Ashley L Scott, David Shepherd, Aya Shinomiya, Claire Smith, Natalie Smith, Alia Suleiman, Satoko Takemura, Iris Talebi, Imaan FM Tamimi, Eric T Trautman, Lowell Umayam, John J Walsh, Tansy Yang, Gerald M Rubin, Louis K Scheffer, Jan Funke, Stephan Saalfeld, Harald F Hess, Stephen M Plaza, Gwyneth M Card, Gregory SXE Jefferis, and Stuart Berg. A connectome of the male drosophila ventral nerve cord. *bioRxiv*, 2023. doi: 10.1101/2023.06.05.543757.

82. Larry F. Abbott, Davi D. Bock, Edward M. Callaway, Winfried Denk, Catherine Dulac, Adrienne L. Fairhall, Ila Fiete, Kristen M. Harris, Moritz Helmstaedter, Viren Jain, Narayanan Kasthuri, Yann LeCun, Jeff W. Lichtman, Peter B. Littlewood, Liqun Luo, John H.R. Maunsell, R. Clay Reid, Bruce R. Rosen, Gerald M. Rubin, Terrence J. Sejnowski, H. Sebastian Seung, Karel Svoboda, David W. Tank, Doris Tsao, and David C. Van Essen. The mind of a mouse. *Cell*, 182(6): 1372–1376, 2020. ISSN 0092-8674. doi: https://doi.org/10.1016/j.cell.2020.08.010.

83. Yael Artzy-Randrup and Lewi Stone. Generating uniformly distributed random networks. *Physical Review E*, 72(5):056708, 2005.

84. Tiago P. Peixoto. The graph-tool python library. *figshare*, 2014. doi: 10.6084/m9.figshare.1164194.

Supplement

Supplemental figures

Figure S1. Supplement for Figure 1. The effects of edge percolation on the size of the largest WCC when **(a)** large connections are removed first and when **(b)** small connections are removed first. **(c)** The sizes of the first two SCCs as a function of the synapse threshold. Spectral analysis of the whole-brain network with **(d)** forward and **(e)** reverse walks. In each case, the stationary probability distributions are shown, as well as the distribution of neuropils in which the inputs and outputs of the top 3000 most visited neurons are located. Renders of the top 3% attractor (red) and repeller (green) neurons are also shown. The top 0.3% are rendered in darker colors. textbf(f) The sizes of the first two weakly connected components (WCCs) as nodes are removed by in-degree or out-degree. Removal of neurons starting with those with largest in-degree (top left) or largest out-degree (top right) result in the brain splitting into two WCCs when neurons of approximately degree 100 start to be removed, a deviation from randomly connected networks (dotted lines). Removal of neurons starting with those with smallest in-degree (bottom left) or smallest out-degree (bottom right) results in a single giant WCC until all neurons are removed. **(g)** The rich club coefficient (red) as a function of in-degree (top) and out-degree (bottom), compared to the predicted rich-club coefficient of a CFG null model (dotted red). The relative rich club coefficient is plotted in blue.

	Neuron lists available on Codex		Definitions
2-neuron motifs	reciprocal connection participants	α ⇄ β	all neurons that participate in reciprocal connections.
3-neuron motifs	feedforward loop participants		neurons that participate in feedforward loop motifs consisting of unidirectional connections: β→α, α→γ and β→γ, precisely.
	3-unicycle participants		neurons that participate in 3-unicycles consisting of unidirectional connections: β→α, α→γ and γ→β, precisely.
N-neuron motifs	highly reciprocal neurons		neurons with the numbers of reciprocal edges ≥ 0.5 x total-degrees.
	neuropil-specific highly reciprocal neurons (NSRNs)		intrinsic rich-club and highly reciprocal neurons with ≥ 50% of incoming connections, and ≥ 50% outgoing connections are contained in the same neuropils, respectively.
Rich-club analysis	rich-club neurons		high-degree neurons that are densely connected with other high-degree neurons (total-degree is higher than 37).
	broadcasters		intrinsic rich-club neurons with out-degrees ≥ 5 x in-degree.
	integrators		intrinsic rich-club neurons with in-degrees ≥ 5 x out-degrees.
Spectral analysis	attractors		top 3% most visited neurons in a forward random walk over the largest strongly connected component.
	repellers		top 3% most visited neurons in a reversed random walk over the largest strongly connected component.

Table S1. Supplement for Table 1. Definitions for all neuron populations identified in this paper.

Neuronal wiring diagrams	Fruit fly (no threshold) *Drosophila melanogaster* (Dorkenwald et al., 2023)	Fruit fly (≥ 5 synapses) *Drosophila melanogaster* (Dorkenwald et al., 2023)
Network size	127,978 neurons 14,680,950 connections	127,978 neurons 2,613,129 connections
Avg. connection strength	3.59 synapses 1 ~ 2358	12.61 synapses 5 ~ 2358
Connection probability α ⇢ β Pr	0.000896 x5.62	0.000160 x1
Connection reciprocity α ⇄ β Pr	0.265 x293 than ER x46.1 than CFG	0.138 x858 than ER x43.8 than CFG
Clustering coefficient	0.108 x59.7 than ER x9.17 than CFG	0.0463 x144 than ER x7.57 than CFG

Table S2. Supplement for Table 2. Network statistics of the fly connectome with no threshold on the number of synapses per connection (left) and a threshold of 5 synapses per connection.

Figure S2. Supplement for Figure 2. **(a)** Distribution of reciprocal degree (gray) alongside distributions of in-degree (red) and out-degree (blue). **(b)** Distributions of reciprocal degree for glut, da, oct, and ser neurons. **(c)** Heatmap showing the fraction of reciprocal incoming connections versus the fraction of reciprocal outgoing connections. Dotted lines indicate a factor of 2 around the $x = y$ line.

Figure S3. Supplement for Figure 4. In-degree vs. out-degree scatterplots showing broadcaster, rich balanced, and integrator neurons plotted by **(a)** the putative neurotransmitter of each neuron and **(b)** the superclass of each neuron. **(c)** Percentile rank distributions of rich club, integrator, and broadcaster neuron populations from various input modalities. **(d)** Scatterplots of percentile rank from one sensory modality on each axis. Broadcaster neurons are highlighted in teal and integrator neurons are highlighted in purple.

Figure S4. Internal and external connections across neuropils. (a) The number and **(b)** relative fraction of neuron weights in each neuropil making connections internal to that neuropil, external incoming connections, and external outgoing connections. **(c)** Comparing the neurotransmitter composition of all internal and all external neuron weights across the whole brain and **(d)** by neuropil.

Figure S5. Supplement for Figure 5. (a) The number of neurons included in each neuropil subnetwork. **(b)** The average connection strength (no synapse threshold applied) of connections made in each neuropil (above), and the connection density of each neuropil (below). **(c)** Reciprocity normalized by connection density for all 78 neuropils. **(d)** Average reciprocal (bidirectional) connection strength normalized by average unidirectional connection strength in all neuropils. **(e)** The relative fraction of glutamatergic neurons participating in bidirectional and unidirectional connections. Absolute percentages of **(f)** acetylcholine, **(g)** GABA, and **(h)** glutamate occurrence in unidirectional and bidirectional connections within each neuropil subnetwork.

Figure S6. Additional supplement for Figure 5. (a) Heatmaps showing the relationship between excitatory and inhibitory (GABA) connection strengths in reciprocal connections in different brain regions. **(b)** Excitatory-inhibitory correlations (Pearson r-score) for all neuropils. **(c)** These correlations do not appear to be correlated with neuropil subnetwork size. **(d)** The neurotransmitter composition of the population of neuropil-specific highly reciprocal neurons (NSRNs).

Figure S7. Supplement for Figure 6. **(a)** Three-node motif distributions for additional neuropils. The frequency of each motif relative to that in an ER null model is plotted to the right, together with the average motif frequencies of 100 CFG models (gray violin plots). **(b)** Average strengths of edges participating in 3-node motifs in the different neuropil subnetworks relative to the average edge strength in each subnetwork.