SNP-Slice: A Bayesian nonparametric framework to resolve SNP haplotypes in mixed infections

Nianqiao P. Ju[1*], Jiawei Liu[2] and Qixin He[2*]

[1*]Department of Statistics, Purdue University, 150 N. University Street, West Lafayette, 47907, Indiana, USA.
[2]Department of Biological Sciences, Purdue University, 915 W. State Street, West Lafayette, 47907, Indiana, USA.

*Corresponding author(s). E-mail(s): nianqiao@purdue.edu;
heqixin@purdue.edu;
Contributing authors: liu3594@purdue.edu;

Abstract

Multi-strain infection is a common yet under-investigated phenomenon of many pathogens. Currently, biologists analyzing SNP information have to discard mixed infection samples, because existing downstream analyses require monogenomic infection inputs. Such a protocol impedes our understanding of the real genetic diversity, co-infection patterns, and genomic relatedness of pathogens. A reliable tool to learn and resolve the SNP haplotypes from polygenomic data is an urgent need in molecular epidemiology. In this work, we develop a slice sampling Markov Chain Monte Carlo algorithm, named SNP-Slice, to learn not only the SNP haplotypes of all strains in the populations but also which strains infect each host. Our method reconstructs SNP haplotypes and allele frequencies accurately without reference panels and outperforms the state of art methods at estimating the multiplicity of infections, allele frequencies, and heterozygosity. We illustrate the performance of SNP-Slice on empirical malaria and HIV datasets.

Keywords: multiplicity of infection, multi-strain infections, Bayesian nonparametrics, Markov Chain Monte Carlo, slice sampling, malaria, HIV, SNP

1 Introduction

Multi-strain infection is a common yet under-investigated phenomenon of many viral, bacterial, fungal, and protozoan diseases [1]. For example, 30-80% of individuals in endemic malaria transmission regions often carry multiple, genetically-distinct strains of the same pathogen species [2, 3]. The presence of multiple strains in individuals often results in poor treatment outcomes [4] or increased virulence from within-host competition [5]. Detailed analyses of population-level parasite diversity, relatedness, and regions subject to selection require monogenomic infection inputs. Therefore, isolates with a high complexity of infection are often excluded from these analyses, which amounts to the majority of samples from high-endemic regions [6, 7]. In addition to the unfortunately wasted resources, it remains difficult to understand the parasite diversity patterns where the knowledge is most needed. Therefore, a reliable tool to learn and resolve the SNP haplotypes from polygenomic data is an urgent need in molecular epidemiology.

Earlier algorithmic efforts on resolving strains focused on metagenomic studies, in which they often rely on deep sequencing coverage per strain ($> 50\times$) or existing strain reference libraries [8]. However, field samples with mixed infections usually contain low concentrations of parasite DNA [9], and correspond poorly to reference libraries. In recent years, several reference-free methods have been developed for bacterial and malarial infections. However, most of the tools only estimate the number of strains infecting each host, known as the multiplicity of infection (MOI), without resolving the constituent strains [10–12]. While SplitStrains [13] and DEploid [14] can infer haplotypes, they are limited to 2-4 strains per infection. The majority of these algorithms focus on resolving mixed infections within one host, without utilizing information from population-level haplotypes and allele frequencies. Thus, it is imperative to develop methods with a focus on population-level inference, without heavy parametrization, to ensure a wide application to different diseases.

In this work, we develop SNP-Slice, a family of slice sampling Markov Chain Monte Carlo algorithms, to infer SNP haplotypes infecting the population as well as whether they appear in each host. In other words, we aim to fully resolve all multi-strain infection information in terms of the haplotype dictionary (feature extraction) and host-strain assignment (feature allocation). In addition, many epidemiological characteristics, such as MOI, allele frequency, heterozygosity, and Fws can be derived from our inference results (Section 4.6). In the SNP-Slice (SS) framework, we use the Indian Buffet Process (IBP) prior [15, 16] to describe strain assignments (Section 4.3), which imposes a clustering structure among the hosts and strains, and allows each host to be infected with any combination of the (potentially infinitely many) strains. In IBP, a single concentration parameter α controls the feature allocation structure as well as the size of the haplotype dictionary, which makes the SS method simpler to tune than competing methods that require more model parameters. Moreover, we propose and examine three novel observational models (Bin, Pois, and NegBin) that tailor to SNP read counts data and accommodate unbalanced within-host strain frequencies (Section 4.2). We also introduce an SS-Cat method to study SNP categorical data that indicates the heterozygosity status at each site. Overall, the SNP-Slice framework

includes four algorithms: SS-Bin, SS-Pois, SS-NegBin, and SS-Cat. In terms of computational contributions, SNP-Slice efficiently solves the challenge of high-dimensional Bayesian inference and yields accurate reconstruction of the haplotype dictionary as well as their presence in hosts. Extensive experiments demonstrate the effectiveness of the proposed methods on simulated malaria mixed infection datasets. Based on our results, we recommend that analysts use SNP read count data when available and that SS-NegBin or SS-Cat be used to analyze field data (SS-NegBin for high transmission regions and SS-Cat for low transmission regions). From an evolutionary biology perspective, the SNP-Slice framework will help recover unexplored information from molecular surveillance of disease transmission, evaluation of transmission control, and detailed monitoring of disease progression within hosts. We demonstrate its performance on HIV-1 time series data (Section 2.8) and malaria field datasets (Section 2.7) and its ability to discover new biological insights from resolved haplotype information.

2 Results

2.1 SNP-Slice outperforms existing methods at MOI and allele frequencies estimations across simulation scenarios and data types.

The SNP-Slice framework simultaneously learns the haplotype dictionary (D) and its host-strain assignment (A). We performed comprehensive benchmarks on realistic synthetic datasets across scenarios of varying transmission intensity (Section A.1).

First, we compared SNP-Slice's performance to the state-of-the-art MOI estimation method The Real McCOIL (hereafter RM) [11]. Examining the error of the estimated MOI against the true MOI ($\widehat{M_i} - M_i$), we find SNP-Slice to be more accurate than RM, both on the population level (Table 1) and stratified by true MOI values (Figure 1). In particular, SNP-Slice achieves exact MOI recovery ($M_i = \widehat{M_i}$) for at least 60% of the individuals, across simulated infection scenarios; and our estimations are within ± 1 from the truth for more than 75% of the individuals (Table 1). When the transmission rate is low (Scenario 1), inference algorithms using heterozygosity status input (SS-Cat and RM-Cat) yield better MOI estimation than algorithms using SNP count input, and 99.8% of MOI estimates are within ± 1 of the truth. As the transmission rate increases (Scenario 2&3), RM-Cat and SS-Cat begin to underestimate MOI. Under the highest transmission intensity, while RM-Prop makes an average error of -1.2, SS-Pois, SS-NegBin can control estimation error below 0.4 in absolute value.

SNP-Slice also outperformed RM at estimating allele frequencies (AF;Section 4.6). We reported the bias and root-mean-squared error (RMSE) of their allele frequency estimations in Table 2. While all methods appear to underestimate AF, the SS-based methods have smaller AF estimation errors than RM-based methods across all transmission intensity levels. Count-based methods are more accurate than RM-Cat and SS-Cat, which uses categorical input. However, SS-Cat achieves higher estimation accuracy than RM-Prop, whose input, SNP read counts, is actually more informative than that of SS-Cat. This demonstrates SS's ability to effectively extract information and interpret mixed infection data.

Table 1 Summary of allele frequency estimation bias and RMSE under different malaria transmission intensities.

Method	% of $M_i = \hat{M}_i$	% of $|M_i - \hat{M}_i| \leq 1$	Bias	RMSE
		Scenario 1		
RM-Prop	91.4	97.2	0.10	0.43
SS-Bin	87.2	91.5	0.27	0.84
SS-Pois	89.8	93.4	0.20	0.69
SS-NegBin	92.2	96.0	0.13	0.53
RM-Cat	92.1	98.6	0.08	0.36
SS-Cat	**97.3**	**99.8**	**-0.004**	**0.20**
		Scenario 2		
RM-Prop	70.0	80.0	-0.64	1.73
SS-Bin	66.2	73.9	0.88	1.88
SS-Pois	70.3	82.3	0.32	1.26
SS-NegBin	74.7	**86.2**	**0.10**	**1.10**
RM-Cat	68.4	80.9	-0.31	1.58
SS-Cat	**76.2**	83.7	-0.58	1.48
		Scenario 3		
RM-Prop	58.0	72.0	-1.20	2.90
SS-Bin	59.1	72.6	0.94	2.08
SS-Pois	60.3	**75.8**	**-0.20**	**1.89**
SS-NegBin	61.5	**75.8**	-0.41	2.04
RM-Cat	57.9	75.1	-0.48	2.31
SS-Cat	**64.3**	73.3	-1.32	2.83

Table 2 Bias and RMSE of AF estimation.

Method	Scenario 1		Scenario 2		Scenario 3	
	Bias	RMSE	Bias	RMSE	Bias	RMSE
RM-Prop	-0.012	0.019	-0.047	0.092	-0.044	0.090
SS-Bin	-0.008	0.020	**-0.009**	**0.021**	**-0.019**	**0.035**
SS-Pois	-0.005	0.016	-0.014	0.031	-0.022	0.040
SS-NegBin	**-0.004**	**0.013**	-0.015	0.029	-0.021	0.038
RM-Cat	-0.012	0.019	-0.061	0.107	-0.063	0.110
SS-Cat	-0.006	0.017	-0.041	0.092	-0.036	0.088

2.2 SNP-Slice accurately recovers heterzygosity of individuals.

We investigated SNP-Slice's ability to learn the individual heterozygosity, which reflects the genetic diversity of strains within a host (Section 4.6) We highlight that only inference methods that recover both strain dictionary and strain allocations are able to provide heterozygosity estimations. As a result, competing methods such as RM cannot perform such an important task. As shown in Table 3, SNP-Slice recovers individual heterozygosities with high precision. Although all methods slightly over-estimate heterozygosity, the highest estimation bias across all malaria transmission

Fig. 1 Comparison of SNP-Slice's MOI estimation performance to The Real McCOIL, stratified by true MOI. Boxplots of error in MOI estimation $(M_i - \hat{M}_i)$ in Scenario 1. The shaded region indicates the estimation error within ± 1. Results are the combination of 128 independent SNP-Slice algorithm repetitions after removing identical duplicate strains for each individual, and each SNP-Slice chain length is set to 10,000 maximum iterations if the input data is read counts (SS-NegBin and SS-Pois), and 5,000 maximum iterations if the input data type is categorical (SS-Cat). The RM results are also based on 128 repeats with 10,000 iterations each. More details about the parameters for running the algorithm are listed in the supplementary material. For visual clarity, a random subset of 5% points in each case is shown.

intensities is only 0.01 with the count-based methods. We also found that SS is more accurate when the transmission intensity is lower.

Table 3 Bias and RMSE of individual heterzygosity.

Method	Scenario 1		Scenario 2		Scenario 3	
	Bias	RMSE	Bias	RMSE	Bias	RMSE
SS-Bin	**0.0003**	**0.003**	**0.003**	**0.009**	0.008	0.025
SS-Pois	0.0008	0.005	0.003	0.010	**0.005**	**0.023**
SS-NegBin	0.0007	0.005	0.003	0.012	0.005	0.025
SS-Cat	0.001	0.012	0.008	0.031	0.021	0.052

2.3 SNP-Slice accurately recovers the strain dictionary.

While accurate heterozygosity estimations already imply an accurate recovery of the haplotype dictionary, we further studied the quality of the inferred haplotypes learned by comparing them with the underlying true haplotypes. First, we define the dissimilarity between two haplotypes to be the proportion of sites with different alleles: Two

identical haplotypes have a dissimilarity of 0 and two opposite haplotypes have a dissimiliarty of 1. Naturally, the similarity between two haplotypes is one minus their dissimilarity. Second, we investigated how many of the true haplotypes are recovered well by SNP-Slice. We define the recovery score of each true strain to be the largest similarity between this true strain D_k and any inferred strain $\widehat{D_r}$ in the dictionary learned by SNP-Slice:

$$\mathrm{rec}_k = \max_r \mathrm{sim}(D_k, \widehat{D_r}) = 1 - \min_r \mathrm{dis}(D_k, \widehat{D_r}). \tag{1}$$

In Table 4, we report the percentage of true strains with a recovery score higher than various cutoff values. We see that count-based methods have similar performance and it is more challenging to recover the haplotypes as the transmission intensity increases. Under low transmission rate, more than $\approx 78\%$ of the strains can be exactly recovered by SS, and for $\approx 99\%$ of the true haplotypes we can find an inferred haplotype that is more than 80% similar to it. Even in the highest transmission intensity situation, more than $\approx 72\%$ of the true haplotypes are recovered at the 80% cutoff value.

Table 4 The percentage of true strains obtaining a recovery score higher than various cutoff values. True and inferred dictionaries have been simplified by merging strains that share identical SNP type at $\geq \frac{92}{96}$ of total sites with others.

Scenario	Cutoff	SS-Cat	SS-Pois	SS-Bin	SS-NegBin
1	100%	77.3%	76.8%	74.5%	**78.2%**
	$\geq 95\%$	81.4%	83.1%	81.7%	**83.4%**
	$\geq 90\%$	85.1%	88.4%	**88.9%**	87.9%
	$\geq 80\%$	95.3%	98.5%	98.5%	**98.8%**
2	100%	41.8%	45.6%	44.5%	**45.8%**
	$\geq 95\%$	46.6%	**52.2%**	52.1%	51.7%
	$\geq 90\%$	70.9%	77.7%	**80.5%**	76.8%
	$\geq 80\%$	86.0%	91.4%	**92.3%**	90.5%
3	100%	15.5%	16.1%	**16.8%**	16.1%
	$\geq 95\%$	18.6%	22.1%	**23.2%**	21.8%
	$\geq 90\%$	33.3%	43.6%	**44.7%**	42.4%
	$\geq 80\%$	61.3%	73.9%	**76.3%**	71.7%

We found a negative association between the true strains' relative strain abundance within the population and their recovery rates (Figure 2). This suggests that the poorly recovered strains are less abundant than those recovered well. In conclusion, SNP-Slice constructs a strain dictionary that resembles the dominating strains in the local population.

2.4 Potential cause of bias in MOI due to minor allele sites.

Although major alleles should appear more frequently than minor alleles in the population, there can be some hosts that have more minor allele sites than major allele sites. In Figure 3, we investigated MOI estimations on these atypical hosts. For hosts

Fig. 2 The recovery scores of true haplotypes and their relative abundance in the population. For visual clarity, a random subset of 5% points in each case is shown.

with $< 20\%$ minor-dominated sites, all methods control estimation error below ± 1. When the fraction of minor-dominated sites increases above 0.4, the bias of RM is around -3, but, in contrast, the average error from count-based SS methods is only around ± 1.

Fig. 3 Error of MOI estimation in Scenario 2, where hosts are organized based on the fraction of sites with more minor allele reads than major allele reads. For visual clarity, a random subset of 5% points is displayed for each method.

2.5 SNP-Slice is robust to unbalanced within host strain frequency

When modeling SNP read count data, SS-Bin, SS-Pois, and SS-NegBin make the assumption that all strains within the same host appear with equal frequency. This modeling assumption facilities efficient computations. However, within-host competition of strains due to the difference in growth rates or drug sensitivity often leads to

unbalanced composition of strain frequencies. We tested the robustness of the current model to the violation of this balancing assumption, by simulating an SNP read count dataset where strains within the same host have unbalanced strain frequencies. SS-NegBin still outperformed RM-Prop when estimating MOI, AF, and heterozygosity. Also, comparing Table 5 with the bias and RMSE reported for Scenario 2 in Tables 1,2, and 3, the performance of SS-NegBin does not deteriorate when the balanced-frequency assumption is violated. This is because SS-NegBin is designed to be more robust to violations of model assumptions, as we will explain in Section 4.2.

Table 5 Comparison of SS-NegBin and RM methods, under unbalanced within host strain frequency.

	SS-NegBin	RM-Prop
MOI		
Bias	**0.36**	-0.74
RMSE	**1.54**	1.82
AF		
Bias	**-0.010**	-0.050
RMSE	**0.028**	0.093
Heterzygosity		
Bias	**-0.002**	×
RMSE	**0.026**	×

2.6 SNP-Slice is robust to heterogeneous SNP sampling efforts

Average read counts from genomic sequencing might differ substantially between isolate samples, especially from host tissue or blood samples, because the amount of parasite DNA and quality contained in these samples vary significantly. Signal intensities from microarray may also vary significantly, in which certain samples have weak signals. Here we test whether SNP-Slice is robust to situations when certain samples have very low read count information or weak SNP signal intensity. We simulated observations where a certain percentage of the hosts have only 1/10 of the total reads compared to the rest of the hosts, and tested whether our recommended method of SS-NegBin can still maintain its previously reported high accuracy. We report the bias and RMSE of MOI, heterozygosity, and allele frequency for SS-NegBin and RM in Table 6. Since RM only uses SNP read ratio data, it is not sensitive to heterogeneity in sampling efforts. While heterogeneous sampling efforts can affect the performance of SS-NegBin, it still outperforms RM on all estimands. Moreover, we find that increasing the percentage of hosts with small total reads introduces a negative bias to MOI estimations and increases the RMSE of all estimations.

Table 6 Summary of SS and RM accuracy, when some hosts have small total read counts.

	10% low reads		25% low reads		50% low reads	
	SS-NegBin	RM-Prop	SS-NegBin	RM-Prop	SS-NegBin	RM-Prop
MOI						
Bias	**0.07**	-0.65	**-0.09**	-0.66	**-0.25**	-0.66
RMSE	**1.11**	1.74	**1.21**	1.73	**1.23**	1.76
AF						
Bias	**-0.016**	-0.047	**-0.023**	-0.048	**-0.024**	-0.048
RMSE	**0.030**	0.092	**0.041**	0.092	**0.043**	0.092
Heterozygosity						
Bias	**0.002**	$\times$	**0.002**	$\times$	**0.002**	$\times$
RMSE	**0.013**	$\times$	**0.015**	$\times$	**0.018**	$\times$

2.7 Empirical data evaluation on malaria in Uganda

We applied SNP-Slice to the cross-sectional survey data in Uganda to illustrate the performance of SNP-Slice on malaria field data. This survey dataset, previously investigated in [11, 17] with the RM method, contains the Sequenom assay of 105 SNPs from dried blood samples collected from random households of three sub-counties in Uganda. Due to prevalent missing data, the final included loci (L) and individuals (N) per sub-county are: Nagongera, 63L/462N; Walukuba, 49L/48N; Kihihi, 52L/74N. We re-analyzed the categorical data indicating heterozygosity status (from [11]) using the RM algorithm and compared its performance with SS-Cat. Using the same setup described in [11], we obtained the same empirical distribution of inferred MOI from RM-Cat as in Fig. 2 of [11]. RM-Cat suggests that Nagongera and Walukuba have a long and rugged tail of high MOIs with a median MOI of 5, whereas Kihihi has the highest proportion of hosts with MOI = 1, and a median MOI of 1, despite a small proportion of MOI > 6 (Fig. 4). SS-Cat also suggested larger proportions of mixed infections from Nagongera and Walukuba (median MOI: 3) than Kihihi, where single-infection dominates (Fig. 4). However, the distributions of MOI inferred from SS-Cat are much more concentrated, right-skewed, smooth, and do not have long and uneven tails. The maximum MOI inferred for Nagongera, Walukuba, and Kihihi are 11, 7, and 6 from SS-Cat, respectively, compared with 21, 16, and 9 from RM-Cat. The dramatic inference differences between the two methods may be attributed to the highly varying estimates from RM-Cat when the true MOI is larger than 3 (see Fig. 1). It is worth noting that RM-Cat is very sensitive to the user-specified value of `maxMOI`, while inferences from SS-Cat do not require knowing the maximum MOI value a priori and are not sensitive to the value of concentration parameter α. When we reduced the `maxMOI` value from 25 (used in [11]) to 15, the distributions of RM-inferred MOIs shifted towards the left(Figure B3). The long tail and rugged shape of the distribution, however, did not change. Given that the data includes all age classes, immunity gradually increases with age, and transmission is uniform in these localities, MOIs are expected to follow a distribution similar to either Poisson or negative binomial distributions [18]. Therefore, SNP-Slice yields a more realistic estimation for the distribution of individual MOIs in all three populations.

Fig. 4 Comparison of individual MOI in three regions of Uganda estimated by RM and SS. The dotted vertical line indicates the median of the estimated MOI.

2.8 Empirical data evaluation on HIV

We applied SNP-Slice to HIV-1 infection data, obtained from [19]. Unlike malaria, strain diversity in HIV does not arise from multiple infections or co-transmitted multiple strains. Instead, HIV has an extremely high mutation rate (0.34 mutations per replication cycle [20]) and forms new strains as viral load increases during the entire infection period. The constant appearance of new strains initiates an arms race with the immune system and significantly prolongs the duration of infection. SNP-Slice is still applicable for such type of disease to infer the within-host strain compositions at different time points. Here we inferred the strain dynamics within individual patients using SS-NegBin. SS-NegBin learned not only the changing pattern of MOI, the strain haplotype dictionary, but also when each strain appeared and disappeared in the evolutionary history of each patient. The number of strain haplotypes is usually low in the early stage of infection, but increases as the disease progresses (Figure 5 Top). As indicated by the lengths of horizontal line segments, some ancestral haplotypes persist throughout the tracked infection years, while most newly evolved haplotypes persist for shorter periods. This agrees with the observation that reversion to ancestral HIV-1 is common because they usually represent the optimal form for the virus [19]. Changes in inferred heterozygosity follow a similar trend to that of viral load, indicating that our inference result is reasonable as mutants increase when the viral load increases.

3 Discussion

Our method is a parsimonious, effective, and robust tool to analyze SNP read count data in the presence of mixed infections, because 1) uses only one parameter α, which controls individual-MOI as well as the size of the haplotype dictionary, 2) achieves high estimation accuracy on important epidemiological characteristics such as multiplicity of infection, allele frequency, and heterozygosity across infection scenarios and 3) does not require reference panels. SNP-Slice extends beyond the capacity of existing methods because of its ability to learn the SNP haplotype dictionary and the

Fig. 5 Top: appearance and disappearance of latent strains for each HIV patient. Bottom: level of heterozygosity (left axis) and viral load (red line, right axis).

strain assignments simultaneously with high accuracy without using reference panels. When estimating regions with prevalent mixed infections, one should use read counts data instead of heterozygosity status data if possible, as the former contains more detailed mixed infection information for resolving higher MOIs. Our experiments have confirmed that count-based methods are more accurate than SS-Cat, with the only exception of estimating MOI in the low transmission intensity setting. Our analysis suggests that SS-NegBin is preferred for studying field samples, where the observation noise can be high. Even in the most challenging scenario, $> 75\%$ of the MOI estimations from SS-NegBin are within ± 1 of the true value. SS-Pois performs as well as SS-NegBin in our simulated data when transmission intensity is high (Scenario 3), but might also be attributed to that we use Poisson distribution to simulate read counts ((Section A.1)). Since SS-NegBin is designed to be more robust to violations of model assumptions (Section 4.2), it is still the top recommended method.

Our re-analysis of the Uganda malaria SNP array data differs significantly from the original analysis in [11], but roughly agrees with another study that sequenced the major antigen gene *var* DBLα types [21]. Unique DBLα types sampled from Nagongera, Walukuba, and Kihihi were, on average, 118, 98, and 68 in [21] respectively. Each parasite genome usually carries around 50-60 unique *var* genes. Because more than 75% of types only appeared once in these populations, we can safely assume each strain carries around 40 unique types, translating to a mean MOI of 3.0, 2.5, and 1.7 in Nagongera, Walukuba, and Kihihi. The maximum MOI in Nagongera and

Walukuba does not exceed 5.3 from the *var* type data, or 6-7 from SS-Cat, while using RM-Cat would lead to the conclusion that many individuals have MOI larger than 10, which is unrealistic given the transmission intensity in the region.

Although SNP-Slice is designed to resolve mixed infections in a local population, it can also be applied directly to a longitudinal sample of strain evolution within a host [19, 22]. When applied to the time series of HIV-1 within patients, SS recovered haplotype dynamics in HIV and achieved reasonable performance. Zanini et al. [19] showed that the phylogenetic tree of V3 region built from patient 1 displays one low haplotype diversity period between two episodes of bursts of new haplotypes, while Patient 3 has a more steady rate of strain replacement. We recovered a similar pattern in Figure 5 regarding the dynamics of strains.

We discuss some limitations of SNP-Slice in its current form and point out several ways to extend this framework. First, SS characterizes representative hosts carrying common haplotypes in the population better than atypical hosts with rare haplotypes, for example, those hosts investigated in Section 2.4 and rare haplotypes in Figure 2. This is unsurprising because SS pools information across hosts to improve haplotype inferences, hence favoring common haplotypes and typical hosts. Following this logic, SS is intended to study mixed infections in a local population. We do not recommend using SS on artificially pooled samples from different localities unless there is enough overlap in the haplotype dictionaries across these populations. Finally, because the SNP-Slice framework reconstructs the haplotype dictionary from scratch, it requires that the number of hosts in the dataset be large enough to contain the most abundant strains in the local population. While SS is not intended to analyze small datasets or single host information, if necessary, the framework can be extended to incorporate existing haplotype reference panels so that one can apply the method separately to each host.

In summary, we developed a general methodology for resolving mixed infections on hosts infected by a population of parasites. SNP-Slice can be widely applied to any pathogen SNP panels as long as they contain biallelic markers and have information on read count, SNP intensity signal, or heterozygosity status. The resolved haplotype information recovered from polygenomic data by SNP-Slice enables many standard analyzing tools for understanding genomic relatedness, diversity patterns, and evolutionary trends. Since SNP-Slice can accurately estimate host-level heterozygosity and construct population-level haplotype dictionaries, it is expected that SNP-Slice will become a standard tool for disease surveillance in molecular epidemiology.

4 Methods

4.1 Problem formulation

Our data consists of SNP read counts (or heterozygosity status) for N people at L sites. For each host i and each site j, we observe Y_{ij}^0 reads from major alleles and Y_{ij}^1 reads from minor alleles, summing to $R_{ij} = Y_{ij}^0 + Y_{ij}^1$ total reads. When read counts data is not available, one observes heterozygosity status (0: only major alleles, 1: only minor alleles, 0.5: heterozygous). Given the data, SNP-Slice aims to resolve mixed infections by estimating two matrices: the feature dictionary D and a feature allocation A. The

feature dictionary $D \in \{0,1\}^{K \times P}$ represents the underlying strains at the population level. Each row of D represents the haplotype of one strain: $D_{kj} = 0$ indicates the kth strain carries a major allele at site j and $D_{kj} = 1$ indicates the site has a minor allele. The allocation matrix $A \in \{0,1\}^{N \times K}$ describes the composition of strains. Each row of A describes which strains are present in the corresponding host: $A_{ik} = 1$ indicates the ith host is infected with the kth strain. The size of the haplotype dictionary, K, is usually unknown. All length L binary sequences are candidates for the haplotype dictionary. We can imagine that there are $K = 2^L \approx \infty$ potential haplotypes and only a finite number of them showed up in the local population.

4.2 Three novel noise models tailored to SNP counts.

If we only observe heterozygosity status (0,1, or 0.5), we can use the categorical model proposed in [11]. Here we propose three novel noise models tailored to SNP read counts data. We describe observational models linking the latent variables (A, D) and the observed quantities Y, R. Given A and D, a host i has $M_i = \sum_{r=1}^{K} A_{ir}$ strains, and here M_i is the MOI host i. At each site j, this host should have $T_{ij} = \sum_{r=1}^{K} A_{ir}(1 - D_{rj})$ major allies.

To begin with, we introduce a Binomial model for Y_{ij}, where

$$Y_{ij}^0 \sim \text{Binomial}\left(R_{ij}, \frac{T_{ij}}{M_i}\right) \tag{2}$$

independently for each host and each site. We also propose two more flexible models: the Poisson model with

$$Y_{ij}^0 \sim \text{Poisson}\left(\frac{R_{ij} \cdot T_{ij}}{M_i}\right), \tag{3}$$

and the Negative Binomial model

$$Y_{ij}^0 \sim \text{NegBin}\left(R_{ij}, \frac{1}{1 + T_{ij}/M_i}\right). \tag{4}$$

We analytically compare the models in Table 7. They have the same expected values for Y_{ij} and the same maximum likelihood estimator (MLE) for T_{ij} when the individual MOI M_i is known. The Poisson and NegBin models progressively introduce higher variance to Y_{ij}, and therefore should be more robust to model miss-specifications such as balanced strain weights during observation. This is consistent with our empirical evaluations in Section 2.5. In Section 2.6, we also compared their robustness to heterogeneous sampling efforts across hosts, which is frequent in field samples.

4.3 Bayesian nonparametric inference with the IBP prior.

We aim to uncover latent strain structures behind the observed SNP data. A key assumption is that the observed data manifests a finite subset of infinitely many latent structures. Instead of specifying a fixed number strains in the population or an upper bound in it, our method infers the size of the haplotype dictionary from the empirical

Table 7 Comparison of three noise models.

Model	$\mathbb{E}(Y_{ij})$	$\text{Var}(Y_{ij})$	MLE of T_{ij} given M_i
Bin	$R_{ij} \cdot T_{ij}/M_i$	$R_{ij} \cdot (T_{ij}/M_i)(1 - T_{ij}/M_i)$	$M_i \cdot Y_{ij}/R_{ij}$
Pois	$R_{ij} \cdot T_{ij}/M_i$	$R_{ij} \cdot T_{ij}/M_i$	$M_i \cdot Y_{ij}/R_{ij}$
NegBin	$R_{ij} \cdot T_{ij}/M_i$	$R_{ij} \cdot (T_{ij}/M_i)(1 + T_{ij}/M_i)$	$M_i \cdot Y_{ij}/R_{ij}$

data. Additionally, mixed infections imply that each host can be infected with any combination of the strains. These modeling assumptions lead us to choose the India Buffet Process, an infinite dimensional probability distribution over sparse binary matrices with a finite number of rows (representing hosts) and an unbounded number of columns (representing features/strains). IBP generates sparse matrices with clustering structure. SNP-Slice only uses one parameter α, which controls the number of unique strains in a finite sample. IBP has been used successfully in imaging and natural language processing. In this work, we use it as a prior to describe the feature allocation matrix A.

4.4 Stick-breaking Construction for the IBP

The IBP is inspired by a culinary metaphor: Consider a restaurant that serves infinitely many dishes, each of them representing one feature/strain, where customers (hosts) eat existing dishes based on their popularity and try $\text{Pois}(\alpha)$ numbers of new dishes. This representation is conducive to understanding the clustering structure in feature assignments, where popular features tend to become more popular as the number of hosts increases. However, here we present a stick-breaking representation of the IBP [15] that facilitates our MCMC computations. Let $\mu_{(1)} > \mu_{(2)} > \ldots > \mu_{(K)}$ be a decreasing ordering of $(\mu_1, \mu_2, \ldots, \mu_K)$ where each $\mu_k \sim \text{Beta}(\frac{\alpha}{K}, 1)$ independently. Let A be a binary matrix of size $N \times K$, where $A_{ik} \sim \text{Bernoulli}(\mu_{(k)})$. Here α is a concentration parameter and K is a truncation level that corresponds to the unknown dictionary size. In the limit of $K \to \infty$, we have $A \sim \text{IBP}(\alpha)$. The stick-breaking metaphor begins with a stick of unit length. One recursively breaks a length $\mu_{(k)}$ stick into two smaller pieces at the $\nu_k \sim \text{Beta}(\alpha, 1)$ relative point, and the next headpiece shall have length $\mu_{(k+1)} = \mu_{(k)} \nu_k$. The length of $\mu_{(k)}$ shall correspond with the frequency of the kth most popular feature in A. As the stick-breaking process continues, the remaining stick will be so small that all subsequent features are inactive. While we can continue the process until $k \to \infty$, there should only be a finite number of active features, denoted as K^+.

4.5 The SNP-Slice algorithm

We develop an algorithm, named SNP-Slice, to approximate the posterior distribution of A, D given SNP counts. The method is based on slice sampling [23], which can handle complex distributions, including the posterior distributions induced by the three non-conjugate observation models tailored to SNP read counts data. SNP-Slice utilizes the stick-breaking construction for the IBP, which includes stick lengths μ as an auxiliary variable. The sampling problem is challenging, because all three latent

variables are infinite-dimensional: μ has infinitely many entries, A has infinitely many columns, and D has infinitely many rows.

4.5.1 Slice variable s

Following [15], we facilitate computations by introducing an auxiliary slice variable

$$s \mid A, \mu \sim \text{Uniform}(0, \mu^\star = \min_{k:\exists i, A_{ik}=1} \mu_{(k)}) \tag{5}$$

where $\mu^\star = \min_{k:\exists i, A_{ik}=1} \mu_{(k)})$ is the shortest stick length of all active features. Importantly, this construction guarantees that s is less than the stick length corresponding to any active feature, so those features associated with lengths of $\mu_{(l)} < s$ must be inactive. As a result, knowing $A_{il} = 0$ for all $\mu_{(l)} < s$, we can fix the parts of (A, D) corresponding to inactive features $\mu_{(l)} < s$. After adding s, the state space becomes $\{\mu, A, D, s\}$. SNP-Slice proceeds by updating these variables iteratively while keeping the joint posterior distribution invariant. We describe the Monte Carlo updating rules for the other three variables in the following.

4.5.2 Data-adaptive truncation level

Our implementation maintains a finite number of sticks to circumvent the infinite dimensionality of the state space and increase the dimensions of μ, A, D when needed. The slice variable s determines this finite truncation level, which adapts to the complexity of the data. When $s \leq \mu_{(k)}$ for all features, we will break more sticks to create an inactive feature, sampling according to a distribution with (unnormalized) probability density function

$$p(\mu_{(k)} \mid \mu_{(k-1)}, A_{\cdot l} = 0 \; \forall l \geq k) \propto \exp\left[\alpha \sum_{i=1}^{N} \frac{1}{i}(1 - \mu_{(k)})^i\right] \mu_{(k)}^{\alpha-1}(1 - \mu_{(k)})^N. \tag{6}$$

We only need one feature with $\mu_{(k)} \leq s$, since the construction forces all subsequent features to also be inactive.

4.5.3 Update μ

Potentially active features ($\mu_{(k)} \geq s$) each have infected $c_k = \sum_{i=1}^{N}$ hosts. The conditional distribution of $\mu_{(k)}$ follows the $\text{Beta}(c_k, N + 1 - c_k)$ distribution restricted to the interval $(\mu_{(k+1)}, \mu_{(k-1)})$. Features with $\mu_{(k)} < s$ are definitely inactive, and we can update their stick lengths using Eq.(6)

4.5.4 Update D

The conditional posterior distribution for the dictionary is $p(D \mid \mu, A, s, Y) \propto p(D)p(Y \mid A, D)$, which is independent of μ and s. We can iteratively update each atom D_{kj} in the dictionary with samples from the Bernoulli distribution. The two conditional probabilities are $p(D_{kj} = 1 \mid A, Y, D_{-k,j}) \propto \rho \prod_{i=1}^{N} p(Y_{ij} \mid A_i, D_{kj} = 1, D_{-(kj)})$

and $p(D_{kj} = 0 \mid A, Y, D_{-k,j}) \propto (1 - \rho) \prod_{i=1}^{N} p(Y_{ij} \mid A_i, D_{kj} = 0, D_{-(kj)})$. We only need to update the active features/strains, because the inactive features do not enter the likelihood $p(Y \mid A, D)$ specified by the three observation models.

4.5.5 Update A

Similar to updating D, we can iteratively update each atom A_{ik} with a sample from the Bernoulli distribution, following the conditional distribution $p(A \mid D, \mu, s, Y) \propto p(A \mid \mu)p(s \mid A, \mu)p(Y \mid A, D)$. The two conditional probabilities are: $p(A_{ik} = 1 \mid \text{rest}) \propto \frac{\mu_{(k)}}{\mu^{(\star,1)}} \prod_{j=1}^{P} p(Y_{ij} \mid A_{ik} = 1, A_{i,-k}, D)$ and $p(A_{ik} = 0 \mid \text{rest}) \propto \frac{1-\mu_{(k)}}{\mu^{(\star,0)}} \prod_{j=1}^{P} p(Y_{ij} \mid A_{ik} = 0, A_{i,-k}, D)$. The denominators, namely $\mu^{(\star,1)}$ and $\mu^{(\star,0)}$, are needed when a change in allocations A_{ik} affects which feature is the last active feature.

4.6 Epidemiological Quantities Derived from SNP-Slice Results

Many genetic and epidemiological quantities can be derived from SNP-Slice results to benefit molecular biology research and public health decisions. In Section 2, we used MOI, AF, and host-level heterozygosity to assess the accuracy of our methods.

At the population level, one can calculate minor allele frequencies (MiAF) per locus with $f_j = \left(\sum_{i=1}^{N} \sum_{k=1}^{K^+} A_{ik} D_{kj} \right) / \left(\sum_{i=1}^{N} \sum_{k=1}^{K^+} A_{ik} \right)$. Major allele frequency is $F_j = 1 - f_j$. Heterozygosity in the local parasite population H_S can be calculated with $H_S = \frac{2}{L} \sum_{j=1}^{L} f_j \cdot F_j$.

For each host, one can calculate MOI ($M_i = \sum_{k=1}^{K^+} A_{ik}$), heterozygosity ($H_i$), and $F_{\text{ws},i}$. Within host heterozygosity [24], $H_i \in [0, 0.5]$, evaluates the average level of allelic diversity over all the sites. It can be computed from SNP-Slice outputs with

$$H_i = 1 - \frac{1}{L} \sum_{j=1}^{L} \left[\left(\frac{T_{ij}}{M_i} \right)^2 + \left(M_i - \frac{T_{ij}}{M_i} \right)^2 \right] = \frac{2}{L} \sum_{j=1}^{L} \frac{T_{ij}(M_i - T_{ij})}{M_i^2}. \tag{7}$$

A single-infection host has $H_i = 0$. Multi-strain infections have $H_i > 0$, and heterozygosity is the highest (=0.5) when a host has two strains that are exactly opposite of each other. Fws [25] is another metric that reflects the complexity of infection in each host. It can be calculated with $F_{ws,i} = 1 - H_i/H_S$.

Acknowledgments. We would like to thank Dr. Frédéric Labbé for guiding the implementation of malaria-SNP simulator. Q. He and N.P.Ju acknowledges support from Purdue University's Showalter Trust Research Award. Also, this project was funded, in part, with support from the Indiana Clinical and Translational Sciences Institute funded, in part by Grant Number UL1TR002529 from the National Institutes of Health, National Center for Advancing Translational Sciences, Clinical and Translational Sciences Award. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Authors' contributions

N. Ju implemented the SNP-Slicer algorithm. J. Liu analyzed empirical and simulated data. N. Ju and Q. He jointly conceived and supervised the work. N. Ju wrote the first draft. All authors revised the manuscript.

Appendix A   Supplementary Methods

A.1   Simulated Malaria Infection Data

Our simulation experiments in Sections 2.1-2.6 uses simulated SNP data generated from an agent-based stochastic simulator of malaria transmission. The simulator tracks parasite strain composition and host immunity [6, 26–28]. At the beginning of the simulation, each parasite carries a randomly sampled SNP set defined by the MAF per site.

In Section 2, we tested three different levels of transmission intensity that encompass endemic transmissions in Africa to seasonal transmissions in Asia or South America. The scenarios 1-3 represent simulations with transmission rates from low to high. In the simulation process, we mainly kept using the default parameter values from [28], assuming identical initial SNP allele frequencies and no linkage disequilibrium among loci. We simulated 200 SNP loci in each scenario, and later sampled 96 of them with minor allele frequency > 0.1. We recorded all 430 individuals with active infections at the ending time point of simulation in Scenario 1. For Scenario 2 and 3, we randomly sampled 200 infected individuals since the total numbers of infected were too large. The few parameters varying among scenarios are listed in Table A1.

Table A1 Parameters for simulated malaria infection data using the stochastic simulator from [28]. The parameters listed here are named according to the literature, where n_genes_initial means the number of genes in the initial gene pool, and n_alleles_per_locus_initial is the initial number of alleles for each epitope locus. Biting rate refers to per mosquito biting rate, and the actual daily biting rate equals this rate multiplied by daily mosquito population size (we used default values from [28]).

Scenario	n_genes_initial	n_alleles_per_locus_initial	biting_rate
1	2000	200	5×10^{-5}
2	10000	1000	1×10^{-4}
3	20000	2000	5×10^{-4}

After simulating the infection process, we simulated minor and major allele read counts per host per site to mimic a short-read next-generation sequencing process. On average, the number of total read counts R_{ij} is around 115, which is typical in high-quality SNP reads datasets. Under the balanced frequency scenario, the number of major allele reads was simulated according to the Poisson model Eq. (3). For the unbalanced strain simulations in Section 2.5, we first sampled the weight of each strain

according to a Dirichlet distribution and used the weighted major allele frequency to simulate major read counts, also according to the Poisson model.

In Section 2, when analyzing MOI, we did an extra process to remove the duplicated strains with identical SNP sequences. If a host contained multiple strains sharing identical SNP sequences, they were counted as one strain when calculating MOI. Such removals were not performed in calculating allele frequency and heterozygosity in order not to lose genetic information.

A.2 Preparing Empirical Datasets for SNP-Slice

In Section 2.7, we applied SNP-Slice to SNP categorical data (homozygous or heterozygous) in three regions of Uganda [11, 17]. The filtering of sampled hosts and sites were processed according to the criteria described in [11].

We obtained HIV with-in host dynamics data from [19]. Since HIV virus mutations often involve more than two nucleotides, we first filter the bi-allelic SNP sites with the cutoff value of 1%. When the proportion of either major or minor allele reads at a site is below 1% of the total reads, we convert that read count to zero. Only 3 patients were included in our analysis since the others did not have enough time points in data.

A.3 Parameter Values for SS and RM in Section 2

SNP-Slice: In each scenario, we ran 128 MCMC chains in parallel and all tables and figures include results from these 128 chains. Each MCMC chain continues for a maximum of 10,000 iterations if read count data is used as input (SS-NegBin, SS-Pois, and SS-Bin) and 5,000 if categorical data is used as input (SS-Cat). At least half of the maximum iteration is run as the burn-in process. MCMC is ended when the local maximum likelihood estimator is not changed for 10% of the maximum iteration. The only exception is 'Nagongera' of Uganda, where the number of hosts is large in this population (462 individuals). Due to time constraints, we set up the maximum MCMC iteration to be 500. As seen in Figure 4, using only 500 MCMC iterations of SS-Cat already yields more realistic results than using RM-Cat.

For simulated malaria infection data, since the average MOI values in scenarios 1-3 are respectively 1.27, 2.58, and 3.90, the concentration parameter α is set as 1.5, 3, and 4. However, we have tested that SNP-Slice is not sensitive to this parameter. For the same scenario, the average inferred MOI would still be very similar even if different values of α are used as input, indicating the robustness of SNP-Slice inference. The parameters for the probability of erroneously calling in the categorical method are set to be as low as $e_1 = e_2 = 0.001$.

For both real infection datasets (HIV and malaria), we use $\alpha = 3$ since SNP-Slice is not sensitive to this parameter. For empirical malaria in Uganda, we follow [11] and use $e_1 = e_2 = 0.05$.

The Real McCOIL: In [11], the total number of MCMC iterations is 100,000 for simulation data. Our trials have found out that setting the total number of MCMC iterations to be 10,000 and 100,000 could generate almost the same summary statistics across 128 chains, so we run 128 chains with 10,000 iterations for all RM cases. The initial MOI is always set to 1.

For empirical malaria infection data in Uganda, we keep using the parameter values used in [11], including $e_1 = e_2 = 0.05$ and 25 as the upper bound of MOI (`maxMOI`).

For simulated malaria infection data, we use $e_1 = e_2 = 0$ when running the categorical method as indicated by [11]. We use `maxMOI` 8, 12, and 16 as `maxMOI` in scenarios 1-3. As we will discuss later, RM-Cat is quite sensitive to this parameter. The default value of this parameter is 25, but here we choose these values to be around or a little less than the maximum inferred MOI by all methods of SNP-Slice to avoid getting extremely high values of MOI from RM.

Appendix B   Supplementary Results

B.1   Accuracy of AF and heterozygosity estimations

In Section 2.1, we reported the bias and RMSE of AF estimations numerically in Table 2. Here we visualize the accuracy of AF estimations in Scenario 2 in Figure B1. The average error of SS-Bin, SS-Pois, and SS-NegBin are around -0.01, which is much smaller than typical values of true AFs in the population.

Fig. B1 True allele frequency and their estimations from SS-NegBin, SS-Pois, SS-Bin, and SS-Cat. A randomly selected subset of 5% of the points are displayed for visual clarity.

Similarly, we visualize the accuracy of heterozygosity estimations in Figure B2, as a supplement to the numerical results reported in Table 3.

Fig. B2 True heterozygosity and their estimations from SS-NegBin, SS-Pois, SS-Bin, and SS-Cat. A randomly selected subset of 5% of the points are displayed for visual clarity.

B.2 Sensitivity of RM to `maxMOI`

As described in Section 2.7, the RM-Cat method generates MOI distributions with long and rugged tails. Here we change the user-specified upper bound of MOI (`maxMOI`) to 15 instead of 25 and rerun RM analysis (Fig. 4). Although the MOI distributions in the new setting still have long tails, we observe notable changes in the inferred MOI distribution, particularly for Nagongera and Walukuba. With `maxMOI`=25, RM estimated the maximum MOI to be 21 and 16 for these two populations. However, when setting `maxMOI`=15, the largest inferred MOI was reduced to 14 and 11 respectively. Higher transmission regions, like Noagonera and Walukuba, are more sensitive to the value of `maxMOI`, which impacts our estimations on high MOI individuals. Notably, SNP-Slice is not sensitive to the value of the concentration parameter α, which enables it to outperform The Real McCOIL in generating more realistic MOI distributions with minimal prior information.

Fig. B3 Comparison of individual MOI in three regions of Uganda estimated by RM, when `maxMOI` is set to 15 (blue) instead of 25 (green, used in Fig. 4). The dotted vertical lines indicate medians of the estimated MOI.

References

[1] Balmer, O. & Tanner, M. Prevalence and implications of multiple-strain infections. *The Lancet Infectious Diseases* **11**, 868–878 (2011). URL https://www.sciencedirect.com/science/article/pii/S1473309911702419.

[2] Tiedje, K. E. *et al.* Seasonal Variation in the Epidemiology of Asymptomatic Plasmodium falciparum Infections across Two Catchment Areas in Bongo District, Ghana. *The American Journal of Tropical Medicine and Hygiene* **97**, 199–212 (2017). URL http://www.ajtmh.org/content/journals/10.4269/ajtmh.16-0959.

[3] Ntoumi, F. *et al.* Age-dependent carriage of multiple plasmodium falciparum merozoite surface antigen-2 alleles in asymptomatic malaria infections. *The American journal of tropical medicine and hygiene* **52**, 81–88 (1995).

[4] de Roode, J. C., Culleton, R., Bell, A. S. & Read, A. F. Competitive release of drug resistance following drug treatment of mixed Plasmodium chabaudi

infections. *Malaria Journal* **3**, 33 (2004). URL https://doi.org/10.1186/1475-2875-3-33.

[5] Frank, S. A. Models of parasite virulence. *The Quarterly review of biology* **71**, 37–78 (1996).

[6] He, Q. *et al.* Networks of genetic similarity reveal non-neutral processes shape strain structure in Plasmodium falciparum. *Nature Communications* **9**, 1817 (2018). URL https://www.nature.com/articles/s41467-018-04219-3.

[7] Otto, T. D. *et al.* Evolutionary analysis of the most polymorphic gene family in falciparum malaria. *Wellcome Open Research* **4**, 193 (2019). URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7001760/.

[8] Anyansi, C., Straub, T. J., Manson, A. L., Earl, A. M. & Abeel, T. Computational methods for strain-level microbial detection in colony and metagenome sequencing data. *Frontiers in Microbiology* **11** (2020). URL https://www.frontiersin.org/articles/10.3389/fmicb.2020.01925.

[9] Carpi, G. *et al.* Whole genome capture of vector-borne pathogens from mixed dna samples: a case study of borrelia burgdorferi. *BMC genomics* **16**, 1–11 (2015).

[10] Assefa, S. A. *et al.* estMOI: estimating multiplicity of infection using parasite deep sequencing data. *Bioinformatics* **30**, 1292–1294 (2014). URL https://doi.org/10.1093/bioinformatics/btu005.

[11] Chang, H.-H. *et al.* THE REAL McCOIL: A method for the concurrent estimation of the complexity of infection and snp allele frequency for malaria parasites. *PLoS computational biology* **13**, e1005348 (2017).

[12] Sobkowiak, B. *et al.* Identifying mixed mycobacterium tuberculosis infections from whole genome sequence data. *BMC genomics* **19**, 1–15 (2018).

[13] Gabbassov, E., Moreno-Molina, M., Comas, I., Libbrecht, M. & Chindelevitch, L. Splitstrains, a tool to identify and separate mixed mycobacterium tuberculosis infections from wgs data. *Microbial Genomics* **7** (2021).

[14] Zhu, S. J., Almagro-Garcia, J. & McVean, G. Deconvolution of multiple infections in Plasmodium falciparum from high throughput sequencing data. *Bioinformatics* **34**, 9–15 (2018). URL https://doi.org/10.1093/bioinformatics/btx530.

[15] Teh, Y. W., Grür, D. & Ghahramani, Z. Meila, M. & Shen, X. (eds) *Stick-breaking construction for the indian buffet process.* (eds Meila, M. & Shen, X.) *Proceedings of the Eleventh International Conference on Artificial Intelligence and Statistics*, Vol. 2 of *Proceedings of Machine Learning Research*, 556–563 (PMLR, San Juan, Puerto Rico, 2007). URL https://proceedings.mlr.press/v2/teh07a.html.

[16] Griffiths, T. L. & Ghahramani, Z. The indian buffet process: An introduction and review. *Journal of Machine Learning Research* **12** (2011).

[17] Nankabirwa, J. I. *et al.* Estimating malaria parasite prevalence from community surveys in uganda: a comparison of microscopy, rapid diagnostic tests and polymerase chain reaction. *Malaria journal* **14**, 1–11 (2015).

[18] Hill, W. G. & Babiker, H. A. Estimation of numbers of malaria clones in blood samples. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **262**, 249–257 (1995).

[19] Zanini, F. *et al.* Population genomics of intrapatient HIV-1 evolution. *Elife* **4**, e11282 (2015).

[20] Mansky, L. M. & Temin, H. M. Lower in vivo mutation rate of human immunodeficiency virus type 1 than that predicted from the fidelity of purified reverse transcriptase. *Journal of virology* **69**, 5087–5094 (1995).

[21] Ruybal-Pesántez, S. *et al.* Population genomics of virulence genes of plasmodium falciparum in clinical isolates from uganda. *Scientific Reports* **7**, 11810 (2017).

[22] Sharon, I. *et al.* Time series community genomics analysis reveals rapid shifts in bacterial species, strains, and phage during infant gut colonization. *Genome research* **23**, 111–120 (2013).

[23] Neal, R. M. Slice sampling. *The Annals of Statistics* **31**, 705–767 (2003).

[24] Nei, M. *Molecular evolutionary genetics* (Columbia University Press, New York, 1987).

[25] Manske, M. *et al.* Analysis of plasmodium falciparum diversity in natural infections by deep sequencing. *Nature* **487**, 375–379 (2012).

[26] Pilosof, S. *et al.* Competition for hosts modulates vast antigenic diversity to generate persistent strain structure in plasmodium falciparum. *PLoS biology* **17**, e3000336 (2019).

[27] He, Q., Pilosof, S., Tiedje, K. E., Day, K. P. & Pascual, M. Frequency-dependent competition between strains imparts persistence to perturbations in a model of plasmodium falciparum malaria transmission. *Frontiers in ecology and evolution* **9**, 633263 (2021).

[28] Labbé, F. *et al.* Neutral vs. non-neutral genetic footprints of plasmodium falciparum multiclonal infections. *PLOS Computational Biology* **19**, e1010816 (2023).