Chitosan stimulates root hair callose deposition and inhibits root hair growth

Matěj Drs[1,2], Samuel Haluška[1,2], Eliška Škrabálková[1,2], Pavel Krupař[2], Lucie Brejšková[1], Karel Muller[1], Natalia Serrano[1], Andrea Potocká[1], Aline Voxeur[3], Samantha Vernhettes[3], Jitka Ortmannová[1], George Caldarescu[2], Matyáš Fendrych[2], Martin Potocký[1], Viktor Žárský[1,2], Tamara Pečenková[1,2*]

[1] Institute of Experimental Botany of the Czech Academy of Sciences, Rozvojová 263, 165 02, Prague 6, Czech Republic

[2] Department of Experimental Plant Biology, Faculty of Science, Charles University, Viničná 5, 128 44, Prague 2, Czech Republic

[3] Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000, Versailles, France

*corresponding author pecenkova@ueb.cas.cz

Abstract

Although angiosperm plants have a general capacity to react after the immunity elicitor chitin or chitosan treatment by the cell wall callose deposition, this response in particular cell types and its evolutionary conservation is not understood. Here we show that also the growing root hairs (RHs) of Arabidopsis can respond to a mild (0.001%) chitosan treatment by the callose deposition and by a deceleration of the RH growth. We demonstrate that the glucan synthase-like 5 (GSL5)/PMR4 is vital for chitosan-induced callose deposition but not for RH growth inhibition. Upon the higher chitosan concentration (0.01%) treatment, RHs do not deposit callose, while growth inhibition is prominent. To understand the specificities of the low and high concentration chitosan treatments, we analysed the PTI signalling components, gene expression, and RH cellular endomembrane and cytoskeleton modifications upon these treatments. Importantly, chitosan-induced callose deposition is also present in the functionally analogous and evolutionarily only distantly related RH-like structures rhizophores (lycopod) and rhizoids (bryophytes). Our results point to the RH callose deposition as a conserved strategy of soil-anchoring plant cells (rhizoids/rhizophores/RHs) to deal with mild biotic stress. At the same time, high chitosan concentration prominently disturbs intracellular dynamics, tip-localised endomembrane compartments and RH growth, precluding callose deposition.

Introduction

Root hairs (RHs) are unicellular plant structures crucial for several aspects of root function: mechanical anchorage of a plant in a substrate, the absorption of water and essential nutrients due to increased root surface, and in root-microbe interactions. In the model plant *Arabidopsis thaliana*, RH cell fate is determined by a position-dependent signalling cascade governed by specific transcription factors, leading to the RH initiation, bulging and cylindrical/polarised tip growth (Carol and Dolan, 2002; Dolan, 1996; Parker et al., 2000; Vissenberg et al., 2020). Several root hair-defective (RHD) genes, F-actin cytoskeletal meshwork and secretory pathway play a prominent role in this process (Baluška et al., 2000; Ichikawa et al., 2014). The tip growth is supported by constant modification of the preexisting cell wall (CW), as documented by the vital importance of several cellulose and hemicellulose biosynthesis participants (Cavalier et al., 2008; Park et al., 2011).

In aerial plant parts, when a pathogen attacks the plant cell, pathogen-associated molecular patterns (PAMP) triggered immunity (PTI) and potentially effector-triggered immunity (ETI) are activated (Jones and Dangl, 2006; Ngou et al., 2021). One of the earliest defence responses to a pathogen attack is the deposition of β-1,3 glucan polysaccharide callose by which the plant reinforces cell walls and blocks the pathogen entry (Jacobs et al., 2003). In Arabidopsis, among 12 callose synthase genes, only one of them, *GSL5* or *PMR4* (*POWDERY MILDEW RESISTANT 4*), is expressed upon pathogenic infection or elicitor treatment, such as fungal cell wall derived chitin and chitosan, but also endogenous damage-associated patterns (DAMP)(Denoux et al., 2008; Gómez-Gómez et al., 1999; Luna et al., 2011; Verma and Hong, 2001; Vogel and Somerville, 2000). It has been recently shown that this callose synthase is responsible also for the root hair callose deposition upon phosphate starvation (Okada et al., 2023).

In contrast to shoots, the roots are constantly exposed to a mixture of microorganisms, and plants need to prevent the over-activation of immunity in roots, mainly by restricting defence responses to specific root zones and cell layers (Chuberre et al., 2018). The PTI in roots also comprises the ROS production, sporadic and patchy callose deposition and growth rate modifications upon PAMP stimulation (Badri et al., 2008; Chuberre et al., 2018; Millet et al., 2010; Rich-Griffin et al., 2020; Zhang et al., 2022). Upon injury of RH, callose can be deposited locally in the RH cell wall as well, however, the identity of the callose synthase involved remains unknown (Galway et

al., 2011). The RH can react to pathogenic bacteria (e. g. *Pseudomonas syringae*), similarly

to the beneficial ones, by the enhancement of the RH growth (Pečenková et al., 2017;

Zamioudis et al., 2013). Unlike living bacteria, elicitor treatments in Arabidopsis did

not provoke prominent RH growth changes, except for the DAMP PEP1 (Okada et al.,

2021).

Both RH tip growth and pathogen sensing are under the control of receptor

kinases, which further modulate the homeostasis of ROS and Ca2+. In both

development and defence, the ligands binding to the cognate receptors, leads to the

activation of Ca2+ channels CYCLIC NUCLEOTIDE GATED CHANNELS (CNGCs)

(Ladwig et al., 2015)) which produce transients with specific patterns, further

transmitted to transcriptional reprogramming of defence (or symbiosis) (Yuan et al.,

2017). Recent studies have reported that CNGC6, CNGC9 and CNGC14 are crucial for

RH tip polar growth (Brost et al., 2019; Ladwig et al., 2015). A phosphorylation

activation of mitogen-activated protein kinases (MAPK) cascade is another

convergence point of signalling pathways. Root hair MAPK3/6 signalling cascade

plays a role in the RH growth (Rentel et al., 2004), as well as in plant immunity

(including chitin-activated defence), together with the second MAPK4-comprising

cascade (rev. in (Zhang and Zhang, 2022)).

The simplest way to study defence responses is by applying a single PAMP or

immunity elicitor such as chitin or its derivate chitosan. Chitin is an insoluble polymer

of β-1,4-linked N-acetylglucosamine, which is in Arabidopsis recognized by a complex

of receptor and co-receptors - Chitin Elicitor Receptor Kinase 1 (CERK1) and lysin

motif receptor kinases 4 and 5 (LYK4 and 5) (Cao et al., 2014; Miya et al., 2007; Wan et

al., 2012). The chitin recognition activates the production of chitinases by plants; to

avoid this recognition and hydrolysis by chitinases, fungi secrete chitin de-N-

acetylases to convert chitin to chitosan. Fully deacetylated polymer units can still

activate Ca^{2+} signalling-related subset of cellular responses, independent of the

CERK1/ROS/MAPK pathway, which in case of overactivation provokes cell death (Ye

et al., 2020).

Chitosan is typically obtained from marine crustaceans′ chitin by chemical

deacetylation (up to 90%) with molecular weights (Mw) ranging from oligo- (<5 kDa)

to polysaccharides (70–100 kDa). Due to the protonation of glucosamine, chitosan

molecules carry an overall positive charge. Chitosan functions as a PAMP (Iriti et al.,

2006), but also as an antimicrobial factor causing intracellular ROS burst with

subsequent oxidation of pathogen cell membrane fatty acids and plasma membrane

permeabilization (Lopez-Moya et al., 2017). Interestingly, high doses of chitosan in the rhizosphere of Arabidopsis, tomato or barley significantly arrest root development, probably via interference with auxin and gibberellic acid regulation or via activation of jasmonic (JA) and salicylic acid (SA) related genes in roots (Lopez-Moya et al., 2017; Suwanchaikasem et al., 2022).

In this report, we compare the elicitation of plant immunity with two different chitosan concentrations, low 0.001%, further referred to as LCC, and high (0.01% HCC), focusing on RH reactions. We bring evidence of the plant capability to adjust the RH rate growth, as well as to reinforce the RH cell wall by the callose deposition. The observed RH-specific effects are dependent on the chitosan concentration, but not its molecular size, and are present also in the functionally analogous but evolutionary non-related structures, namely rhizoids of the liverwort *Marchantia polymorpha* and rhizophores of the lycophyte *Selaginella uncinata*, as well as in the regenerating protoplasts of *Physcomitrium patens*. Further, on the subcellular level, we tested the responsiveness of RH endomembranes and cytoskeleton to chitosan treatments. To understand the impact of the chitosan treatments on overall plant growth and defence, we performed Ca^{2+} and MAPK signalling assays and RNAseq analysis, pointing to limited but significant changes caused by LCC, compared to HCC treatment. Out of these experiments, we could conclude that the RH callose deposition is the consequence of the mild biotic stress affecting the cells with the intensive growth rate, while the stronger biotic stress provokes almost immediate and more prominent tip-localised endomembrane changes and RH growth cessation. Our results thus reveal the unexpected aspects of the RH growth-defence trade-off, and set the basis for possible future modifications of the general plant resistance by mild and environmentally low-impact treatments.

Results

Chitosan treatment induces concentration-dependent root growth changes and callose deposition also in root hairs

When the 5-7 day-old *A. thaliana* seedlings in liquid medium were treated with several different concentrations of chitosan 0.0001% - 0.01%, 24 h post treatment we could observe the ectopic callose deposition in cotyledon leaves, but also in roots and RHs (Figure 1A). The patterns were dependent on used concentration - while LCC (0.001%) induced small callose deposits in leaves, no primary root growth inhibition

and prominent callose depositions in regions close to the root tips, the HCC (0.01%)
induced larger callose deposits in leaves, prominent primary root growth inhibition
and no callose in root hairs. Conversely, plants incubated with very low chitosan
concentration (0.0001%) were indistinguishable from the mock treatment by both the
lack of prominent callose deposition and no inhibition of the primary root growth
(Figure 1A, B). Given the striking response of RH to specific chitosan concentration
(LCC, 0.001%), we wondered if it is caused by the particular chitosan composition. We
therefore tested the effect of the same low concentration for several other chitosan
variants with different Mw, viscosity, and solubility. The observed effect on the callose
deposition along RH was consistent for all analysed chitosan variants but not for the
non-deacetylated chitin, confirming that the RH callose deposition is chitosan-specific
(Figure 2). When the seedlings were treated by LCC in distilled water, which does not
support the RH growth such as liquid 1/2MS, there was no callose deposition in RH,
pointing to the fact that the callose is deposited only in newly growing RH exposed to
chitosan (Supplementary figure 1A). Similarly, the application of HCC provokes
strong   RH   growth   inhibition   without   observable   RH   callose   deposition
(Supplementary figure 1B).

We could conclude that chitosan provokes specific responses in both above-
ground organs and root hairs, depending on the applied concentrations.

Callose deposition is PMR4-dependent and uncoupled from the RH growth inhibition

In order to uncover the key players in the RH callose deposition, we decided to
analyse the RH modifications in the *gsl5/pmr4* mutant lacking the callose synthase
isoform responsible for the pathogen-induced callose deposition. The LCC-induced
callose deposition was completely abolished in the *pmr4* mutant (Figure 3A). Further
on, when using the GFP-tagged PMR4 construct (Sabol et al., in preparation), we could
observe a higher accumulation of PMR4 on the tip of the newly growing RH 4h after
the chitosan addition in comparison to the mock treatment (Figure 3B). Along with the
callose *pmr4* mutant, we tested *cals8-1,* which is involved in the biotic stress-related
regulation of plasmodesmatal permeability, as well as a range of secretory, defence-
related (including signalling co-receptor *bak1,* chitin receptor *cerk1*) and phytohormone
mutants, however, for all of them we found the RH callose pattern similar to the one
in WT/Col-0 (Supplementary figure 1C).

While the 24 h treatment by HCC completely arrested the growth of new RHs,
the LCC caused a gradual decrease in RH growth (Figure 3C), and we hypothesised
that it may be coupled with callose deposition and rigidification of RH cell wall.
However, the LCC-induced RH growth decline in the *pmr4* mutant that does not
deposit callose was similar to that of WT/Col-0 (Figure 3A). We also monitored both
callose deposition and root hair growth in WT after the LCC addition, showing that
the callose starts to accumulate already at 4h time point, while the average inhibition
of the root hair growth starts to be evident after 8h, reaching significant difference from
mock treatment approximately 12 h after addition of chitosan (Figure 3D). This
difference in the time course pattern further confirms the independence of the RH
growth decline and callose deposition.

The LCC-induced callose deposition in RHs is thus PMR4-dependent and
uncoupled from the mechanisms underlying slower RH growth.

Low concentration chitosan treatment triggers weak PTI-related signalling

We further wanted to uncover differences in general activation of PTI responses
between the LCC and HCC treatments. For that purpose, we first followed calcium
spiking using sensor R-GECO construct (Keinath et al., 2015) in a 20 min treatment of
seedlings by the two concentrations of chitosan. There was only a mild Ca^{2+} increase
in spiking in LCC treatment compared to HCC treatment, which triggered more
prominent peaks of intracellular Ca^{2+} increase (Figure 4A). Besides their intensity, the
two treatments also differ in their patterns and peaking time, evidencing early
specificity of the LCC versus HCC. Interestingly, despite the obvious employment of
the Ca^{2+} channels in chitosan sensing, *cngc* mutants have unaffected deposition of the
callose to the RH cell walls, indicative of higher importance of the CNGC-mediated
signalling for the RH growth than for the callose deposition (Supplementary figure
1C).

Additionally, we assayed another parallel early signalling PTI pathway, MAPK
activation, after the two chitosan concentrations 30 min long treatments. Similarly, a
western blot analysis of MAPK activation by phosphorylation (quantified as a ratio of
band intensities obtained for the phosphorylated part and whole pool MAPKs)
demonstrates significant MAPK activation in both cases. However, HCC causes a more
prominent effect on the MAPK signalling pathway than the LCC (Figure 4B). The

distribution of activation along the three tested MAPKs - MAPK3,6 and 4 was similar for both treatments.

These results confirm that the LCC treatment, similarly to the HCC one, activates PTI-related signalisation, nevertheless, the pattern and intensities of these activations significantly differ.

Chitosan modulates rapid calcium and endomembrane dynamics but not actin cytoskeleton in growing root hairs

Having shown distinct fast calcium responses after LCC and HCC treatment, we next analysed the spatiotemporal details of the Ca^{2+} reporter R-GECO within the RH cells. In mock treatment, the very tips of RH undergo constitutive pulsatile changes of R-GECO signal intensity, while the atrichoblasts´ R-GECO pattern transiently changes upon the buffer exchange during the mock treatment, and much more prominently upon chitosan treatments (Figure 5A, Supplementary video). Interestingly, upon both the LCC and HCC, a wave of R-GECO signal often starting within 5-7 min at the RH tips and moving to the RH cell interior and further on through trichoblast toward other root cells was observed (Figure 5B, Supplementary video).

We then tested if and how the chitosan-triggered early signalling event affects the intracellular dynamics. We focused on well-growing RH close to the root tip, and after applying mock, LCC and HCC treatments, we followed intracellular changes in localisation and dynamics of vacuolar and cytoskeletal genetic markers, as well as endocytic dye FM4-64 in a real-time approach (Figure 6). We monitored treated RH for 0-30 minutes. First, we could observe an immediate and significant root hair growth cessation in the case of HCC-treated seedlings, often accompanied by RH damaging and bursting. In addition, for LCC we could also observe significantly lower average growth rates. Nevertheless, out of the obtained data distribution, we could conclude that in the case of this treatment, RH mostly either stop or prominently slow down their growth, or they grow at rates similar to mock-treated root hairs (Figure 6A). Interestingly, the vacuolar dynamics does not follow the trend found for growth and the RHs react to the two treatments contradictory – while LCC causes small and insignificant enhancement in vacuolar dynamics, HCC impairs the vacuolar dynamics in the RH tip area (Figure 6B). The cessation of vacuolar movement in the RH tip upon HCC treatment may be related to observation obtained for the same treatment on the

enhanced FM4-64 dye accumulation in RH endocytotic/vesicular compartment,
probably as a consequence of collapsing endomembrane trafficking (Figure 6C).

Since vacuolar dynamics is often governed by the actin cytoskeleton, we tested
the effect of various chitosan concentrations on the organization and dynamics of
filamentous actin in RHs. Somewhat unexpectedly, there was no difference in the actin
organisation during the early response to chitosan, as dynamics of actin-decorating
fimbrin-GFP were indistinguishable among the mock, LCC, and HCC treated plants
(Supplementary figure 2).

The microscopical observations thus confirmed that the LCC, despite
provoking prominent changes in CW composition and RH growth inhibition in later
time points, does not impose a prominent stress that would disturb endomembrane
compartments in early time points. On the other hand, HCC treatment significantly
inhibits RH growth, rapidly affects calcium gradient and oscillations in RH tip, blocks
vacuolar dynamics and provokes accumulation of endocytotic/FM4-64 marked
compartment.

Chitosan induces changes in gene expression and cell wall components

To obtain a more general view on chitosan responses and to gain another insight
into distinct plant reactions to low and high chitosan treatment, we performed the
RNAseq analysis of liquid medium-grown Arabidopsis seedlings treated with mock,
LCC or HCC for 24h. The general responsiveness of plants to LCC is far less dramatic
than what is observed on the level of RHs, as only 58 differentially expressed genes
(DEGs) were identified for LCC, and 8507 for HCC, comprising participants of plant
PTI responses; most of the DEGs found for LCC are also present in the HCC DEG gene
set, while only 11 genes are specifically regulated by LCC (Table I; Supplementary
table I). Out of these 11 DEGs, 10 genes are downregulated, and only one, AtLRX6
(Leucine-rich repeat-extensin 6; AT3G22800), upregulated and found to belong to
leucine-rich repeat/extension-type proteins, reported previously to play a role in RH
growth. We have also identified genes which were differentially expressed in LCC and
HCC treatments - in contrast to HCC which upregulates their expression, LCC
downregulates (bottom 5 genes in the Table I). Interestingly, when the same LCC was
applied on mature rosette leaves, both RNAseq and cell wall composition analysis
revealed differences that are more prominent and more similar to expression patterns

provoked by other elicitor treatments (similarly to the HCC-treated seedlings; Supplemental figure 3A).

The restricted number of DEGs provoked by LCC is in concordance with milder Ca^{2+} and MAPK responses activation and with limited morphological changes observed mainly on the level of newly growing RHs, implying that the RH callose encasement is an effective but limited and localised strategy for the plant to overcome mild biotic threats, and to proceed with undisturbed development.

Even though our results are centred around seedlings' roots reactivity to chitosan, in order to obtain a more general view of the LCC impact on plants, we analysed LCC-caused gene expression of adult rosettes, and we encountered the results with those from the cell wall profiling of the LCC-treated rosette leaves (Supplementary figure 4; (Voxeur et al., 2019)). Both analyses accordingly revealed the affection of pectin synthesis and modification/methylation states upon LCC. We next realised an enzymatic fingerprinting of the homogalacturonans and xyloglucans using a commercial polygalacturonase and an endo-cellulase, respectively, from 5-weeks-old Arabidopsis leaves treated in presence of mock and LCC. We analysed the oligosaccharide produced by LC-MS (Paterlini et al., 2022). It revealed that LCC treatment likely triggered the demethylesterification and acetylesterification since the relative amounts of GalA4Me decreased and GalA3 Ac increased, respectively. Chitosan treatment also modified the structure and composition of hemicellulose building blocks. It remains to be further assessed whether, and how the CW components modifications take part in the RH responsiveness to chitosan.

Callose deposition is conserved in evolutionarily unrelated structures functionally analogous to root hairs

Because the callose deposition was preserved in all tested signalling and/or secretory pathway mutants, we speculated that it is a robust reaction which might be evolutional conserved and present also in non-angiosperm plant lineages and their stress responses. In a similar experimental setup with plants cultivated in the liquid medium, we were able to induce the growth of the morphological structures functionally analogous to root hairs, namely rhizoids in the liverwort *Marchantia polymorpha* and the rhizophores of spikemoss *Selaginella uncinata*. When the plants were treated for 24 h with LCC, *M. polymorpha* rhizoids growing out of an asexual reproductive body gemma were enclosed by callose, similarly to rhizophore surface

cells of *S. uncinata* (Figure 7). Also, no callose depositions were detected in mock, while HCC treatment prevented their appearance, mirroring the situation with RH of *A. thaliana*.

We further verified if the relation of chitosan treatment and callose response can also be established for different unicellular model – regenerating protoplasts derived from the moss *P. patens* tip growing protonemal cells (Brejšková et al., 2021; Cove et al., 2009). We could indeed observe a prominent callose deposition caused by LCC, while HCC treatment had a deleterious effect on protoplast viability and regeneration (Supplementary figure 4A). Interestingly, in contrast to *P. patens* regenerating protoplasts, above-ground tissue-derived protoplasts of *A. thaliana* did not deposit callose while regenerating in the presence of chitosan (Supplementary figure 4B).

These results confirmed that the potential of RH and RH-like structure to deposit callose is an evolutionary old intrinsic feature. This capacity, however, may be exclusive for the tissue derived from the non-aerial plant parts, exhibiting fast cellular growth and newly generating the cell wall.

Discussion

In this work, we uncovered the RHs capability to perceive the chitosan as an elicitor, and depending on its concentration, to deposit callose to the cell wall and/or to inhibit RH growth. The two processes are not mutually dependent, and the two studied chitosan concentrations, the low (LCC) and the high ones (HCC), may induce plant responses that overlap in their quality and quantity, but also those which are specific. We also demonstrate that the shift in treatment stringency toward the milder concentrations of the stressor can reveal new defence- and development-related phenotypes.

Phytopathological studies often deal with approaches involving high elicitor concentrations necessary to provoke responses that could be easily observed and quantified (Aslam et al., 2009; Luna et al., 2011). We speculate that the low chitosan concentration (LCC, 0.001%), used in our study, imposes a mild biotic stress, which young plants can sustain, and to which they react by the adjustments on the level of RHs (as the most exposed cell type) by slowing down their growth rate and by fortifying the cell wall with callose. Besides RHs, the LCC stimuli are also perceived by other root parts and also above-ground organs, as our results from Ca^{2+} and MAPK assays showed. Interestingly, prolonged LCC treatment had no impact on the growth of young plants, and they are not affected in their further development, implying an

accommodation to these low chitosan concentrations, also observable in quenching of early phases of Ca^{2+} spiking patterns. Our results thus point mainly to the quantitative difference between the LCC and HCC treatments. Nevertheless, in agreement with RNAseq and CW analysis, the *Pseudomonas syringae* susceptibility assays showed an absolute enhancement of resistance of plants to bacteria when applied with LCC, thus strengthening our hypothesis that the mild biotic stress provides an optimal score for defence and growth trade-off costs (Supplementary figure 5A).

Similarly, the RNAseq data from LCC- and HCC-treated seedlings corroborate the existence of primarily quantitative differences between the two treatments since only a minority of DEGs was found to be specific for LCC. Nevertheless, the presence of minor qualitative differences is legible as well. For instance, it has been reported that the HCC treatment, which impairs plant growth, also affects the auxin pathway. In contrast, LCC has only a minor effect on auxin signalling enhancement, as demonstrated using the DR5::GFP auxin reporter line (Supplemental figure 5B). Despite the scarcity of the LCC-specific candidates, we consider that the role of one of the identified DEGs, LRX10, is worthy of further verification, since this protein, together with related pollen tube-expressed isoforms, has been found to be involved in the pollen tube cell wall pectins and callose deposition, important for the process of fertilisation (Wang et al., 2018).

Callose deposition is typically described as an early defence response in leaf tissue, where it serves mainly to seal the plasmodesmatal connection between cells to prevent pathogen spread and nutrient withdrawal (Epel, 2009) or to prevent pathogens' entrance (Meyer et al., 2009; Ortmannová et al., 2022)), but also in primary root epidermis after the defence-eliciting treatments, displaying a similar spotty pattern as found in leaves (Millet et al., 2010). In agreement with this defence involvement, callose deposition is repressed in the establishment of mutualistic interactions (rev. (German et al., 2023)). Besides the defence-related functions, callose also serves as a mechanical support in the new cell wall formation, in the pollen exine formation and germinating-pollen plug, but also in a functional specialisation of trichome basal parts (Kubátová et al., 2019; Kulich et al., 2015; Ma et al., 2021; Motomura et al., 2022). The RH callose deposition observed by us is tightly coupled with a state of the elongating root cells, and in the newly growing root hairs in the zones closest to the root tip. Therefore, we hypothesise that the RH callose coating is the consequence of two processes going along – of the elicitor caused defence signalling, as evidenced by the dependance on the typically pathogen-induced isoform of callose synthase PMR4, but also of the fragile and immature state of the cell wall

undergoing synthesis, which was additionally confirmed on our experiments with regenerating protoplasts that readily deposit callose upon the LCC treatment (Dehors et al., 2019), as well on recently demonstrated abiotic stress and starvation related RH callose deposition (Okada et al., 2023).

Both LCC / HCC treated rhizoids/rhizophores of *Marchantia polymorpha* and *Selaginella uncinata*, despite their evolutionary unrelated origins, provide the same morphological and subcellular responses, including the employment of the evolutionarily conserved callose synthases in terrestrial plants. This functional conservation might have an origin in the callose contribution to the newly formed post-meiotic cell wall of spores or pollen. Secondarily, these independently-evolved mechanisms may have also been involved in the onset of responses of plant cells to the hostile environmental conditions of land plants, also including pathogen threat (Herburger and Holzinger, 2015; Ks et al., 2020).

Unlike LCC, the HCC has a prominent effect on the appearance and subsequent loss of complexity of the RH tip-localised portion of the vacuolar compartment. The HCC causes the vacuolar withdrawal and dismission of the complex tubular tip-localised vacuolar growing parts. Given the massive uptake of FM4-64 dye in HCC-treated RHs, this may be the consequence of the enlargement of the endocytic zone in the RH tip, or an unrelated endomembrane response. The prompt response of vacuolar dynamics to chitosan may suggest that, besides a proposed role for vacuolar calcium channels in mechanosensing of the tip-growing cells, a part of the pathogen sensing might reside on the tonoplast as well (Radin et al., 2021).

Despite the critical importance of the actin cytoskeleton in both RH growth and responsiveness to pathogens, in our experiment, at an early time points, there was no significant modification of fimbrin-GFP-labelled actin filaments upon chitosan treatments (Supplementary figure 3, see also (Porter and Day, 2016; Ringli et al., 2002; Sun et al., 2021; Vaškebová et al., 2018)). It would be interesting to verify if this observation might be related to the different requirements for the callose- and cellulose synthases localization along the plasma membrane by the microtubules and the actin, respectively, similarly to what has been found for the tobacco pollen tube (Cai et al., 2011).

In addition to demonstrating the usefulness of mild chemical/eliciting treatments of plants for opening new perspectives in plant studies, we are also suggesting the simple experimental setup based on a single, exposed, and sensitive cells. This approach could be further employed for gaining the knowledge on the root

and RH-related immunity, as well as in the studies of the overall plant fitness in relation to the rhizosphere conditions.

Conclusions

The effect of the low concentration chitosan treatment on the callose accumulation in Arabidopsis root hairs and similar effect in Marchantia and Selaginella rhizoids or rhizophores, expose their unexpected deeply rooted capacity to react by the callose accumulation in response to this fungal elicitor. The root hair cell wall callose deposition is PMR4/GSL5-dependent and occurs in parallel, but independently of the root hair growth inhibition. The higher chitosan concentration treatments cause prominent root hair growth inhibition that precludes the callose deposition, leading to the phenotypic difference between the two treatments. The intensive root hair growth inhibition is probably the consequence of the emphatic defence alerting, as well as interference with the root tip-localised vacuolar and endocytic compartments.

Material and methods

Plant cultivation

For seedlings and plants cultivation, seeds were surface sterilised (5 min in 70% ethanol, 2 × 5 min in 10% commercial bleach, rinsed three times in sterile distilled water) and stratified for 2–3 days at 4°C. Seeds were then germinated and grown on vertical ½MS agar plates (half-strength Murashige and Skoog salts, Duchefa Biochemie, supplemented with 1% sucrose, vitamin mixture, and 1.6% plant agar, Duchefa Biochemie) at 21 °C and 16 h of light per day for five-seven days. For cultivation of plants, seedlings that were 7 d-old were transferred into Jiffy Products International pellets and grown at 22 °C and 16 h of light per day in growth rooms.

The following previously published Arabidopsis mutant and transgenic lines were used in this study: *aux1-21* (CS9584, (Bennett et al., 1996)), *bak1-4* (CS71777; (Kemmerling et al., 2007)), *cals8-1* (SALK_037603; (Cui and Lee, 2016)), *cerk1-1* (pst14772; (Miya et al., 2007)), *cngc14-1* (SALK206460; (Shih et al., 2015)), *cngc2-3* (SALK_066908; Chin et al., 2013), *exo70H4-1* (SALK_023593;(Kulich et al., 2015)), *npr1* (CS3726; (Cao et al., 1994)), *pmr4-1* (CS67159; (Vogel and Somerville, 2000)), *rbohD-3* (N9555, Torres et al., 2002 11756663), UQ10::GFP-PMR4 (Kulich et al., 2018),

promPMR4::GFP-PMR4 (Sabol et al, in preparation), DR5-GFP (Heisler et al., 2005), fimbrin-GFP (Wang et al., 2004), and vacuolar ATPase Vti-GFP (gifted by Karen Schumacher, Heidelberg). The wild type (WT) Col-0 was used as control. When needed, primers used for the genotyping were the same as used in the above-cited original studies.

Chitosan treatment

Plants were propagated in vitro for 5-7 d on vertical plates, and then transferred by forceps into liquid ½ MS (control) or liquid ½ MS with chitosan added (100x and 1000x diluted stock solution of the 1% chitosan in 1% acetic acid), usually 4-8 plants per 3 ml, in 6-multiwell plates (Nunclon, Thermo Fisher Scientific). Plants were cultivated for 24 h 16/8 h light/dark at 21–22°C without shaking. After 24 h of incubation, plants were transferred to 3 ml of ethanol for 6-24 h at room temperature for destaining, with mild shaking, and subsequently transferred into 3ml of 150 mM K2HPO4. After 1h, aniline blue (final concentration 0,005% solution in 150 mM K2HPO4) was added and seedlings stained for additional 24 h with mild shaking and observed under the microscope. Additional verifications confirmed that there were no differences between the mock treatment with and without corresponding dilution of acetic acid. Several chitosans used for analyses were purchased from Merck, Germany (Cat. Nos C3646, 523682, 448877 and 419419), as well as chitin (C7170).

Root and root hair phenotype examination

The plant responses to pathogens were monitored by microscopy and documented by photography. The seedlings were placed on microscopic glass, their primary root stretched and lengths measured. Microscopic analysis of root tips and root hairs was performed using an Olympus BX51 microscope with attached DP50 camera (Olympus; Psm) or Zeiss AxioImager ApoTome2 microscope (Pst). For each treatment/genotype 5-10 plants were photographed and analysed. Root hair lengths were determined for the first 20-40 root hairs starting from the root tip for each plant; for statistical analysis. Callose deposition was analysed using the pre-set conditions for bright-field and DAPI and objective 5x or 10x.

Protoplasts preparation

The moss *Physcomitrium patens* Gransden strain was routinely propagated in vitro on BCD medium supplemented with ammonium tartrate dibasic (BCDAT), according to (Cove et al., 2009), in a climate chamber (16 h of light/8 h of dark, 25 °C, illuminated by fluorescent tubes at 50-70 μmol m^{-2} sec^{-1}). Protonemal tissue (6-7 days after propagation) from cellophane-grown culture was transferred into the

1%Driselase/8%mannitol solution. When filamentous structure became invisible (approx. 1 h at RT), protoplast suspension was filtered via/through 100 µm steel mesh. Protoplasts were diluted in 8% mannitol up to 10 ml and spun down (700 rpm for 4 min, without brake) and this step was repeated three times. Protoplast sediment was carefully resuspended between centrifugations to avoid aggregation of protoplasts. Finally protoplasts were incubated in the liquid regeneration medium (BCDAT+ 5mM NH_4+ 6% w/v mannitol + 10 mM $CaCl_2$) overnight at RT. Arabidopsis protoplasts were prepared according to (Yoo et al., 2007). Briefly, Arabidopsis leaves from 3-4 weeks old plants grown under short day conditions (8 h og light per day) were cut onto 2 mm stripes. The 2g of prepared plant tissue were submerged in 37 °C warm protoplast solution (0.4M mannitol, 1.5% cellulase, 0.4% macerozyme, 20mM MES pH 5.7, 20mM KCl, 0.1% BSA, 10mM CaCl2) and vacuum has been applied for 30 min. The submerged leaves were kept in the dark in RT. After 3 h, the flasks with protoplasts were gently swirled and the quality of protoplasts was checked with a microscope. The protoplast solution was washed with a WI buffer (4mM MES pH 5.7, 0.5M mannitol, 20mM KCl), filtered through miracloth and clarified with multiple centrifugations (3x100g, 2 min). For the recovery of protoplasts, 0.7 ml of WI solution in a 12-well plate was applied. Both types of protoplasts were left to recover for 3h, and were subsequently treated with chitosan or mock for 24 h. Protoplasts were stained with aniline blue and the final condition of protoplasts monitored with the microscope Olympus BX51 microscope with attached DP50 camera.

Selaginella and Marchantia propagation and callose deposition assay

For sterile *Selaginella uncinata* ('Comenius' strain) culture establishment, excised fragments containing meristems were washed with distilled H_2O and surface sterilized with 70% EtOH (1 minute) and subsequently with 40% commercial bleach (SAVO)/2% sodium hypochlorite for 15 minutes and rinsed with a sterile water twice (Park et al., 2020) . Axenic regenerants were subsequently vegetatively propagated on ½ MS media with 0.6% agar under standard conditions (22°C, 16/8 light/dark, ca. 180 µmol/m²/s). Chitosan treatments were performed on whole regenerating plantlets interchangeably with the *Arabidopsis thaliana* approach.

Marchantia polymorpha Tak-1 has been maintained on ½ Gamborg media ((Gamborg et al., 1968) ; G0210 Duchefa) supplemented with 1,2% agar under standard conditions (22°C, 16/8 light/dark, ca. 180 µmol/m²/s). Chitosan treatments were done with gemmae analogously to an *Arabidopsis thaliana* procedure, with a slight change, where the liquid ½ MS has been replaced with ½ Gamborg media supplemented with 1% sucrose to enhance the rhizoids initiation**.**

Confocal and Real Time microscopy

Small agar block with *in vitro* cultivated 6-day-old seedlings was gently cutted from the cultivation plate and transferred into sterile microscopic chambers containing 100 ul of liquid ½ MS. Chambers were covered with lids and left overnight in a cultivation chamber. For real time treatments we used custom designed 3D printed perfusion add-on which fits into the microscopic chamber and allows exchange of media inside the microscopic chamber during microscopy session. To mediate the exchange we used medical grade silicone tubes mounted onto a peristaltic pump and perfusion add-on with speed approximately 100 ul/min. All microscopic samples were 10 min pretreated with the mock solution right before capturing, chitosan treatment was applied directly during the microscopic session. For mock treatments the liquid ½MS was used. For chitosan treatments, 0.001% chitosan (LCC) or 0.01% chitosan for HCC were added into the ½ MS. For the FM4-64 (Thermofisher scientific) staining we used the same perfusion settings and solutions just with the addition of 0.001% FM4-64 dye (1 mM stock solution). Samples were treated for 15 minutes before observation.

Real Time imaging was done on the Spinning disc (SD) microscope Nikon (Eclipse Ti-E, inverted) with Yokogawa CSU-W1 SD unit (50mm) equipped with Omicron LightHUB ULTRA light source, Plan-Apochromat L 20x/0.75 or Plan-Apochromat LS 40x/1.25 WI objectives were used for capturing. Microscope is operated by NIS Elements 5.30 software. GFP was observed using a 488 nm excitation laser with Semrock brightline Em 525/30 filter. FM4-64 was observed using a 488 nm excitation laser with Semrock brightline Em 641/75 filter.

Analysis of vacuolar dynamics

Dynamics of vacuoles were analysed using *stack difference* task inside *mutli kymograph* plugin ImageJ (FIJI) followed by analysis of *moments* thresholded area of the result image. See supplementary material *VD_macro.ijm*.

FM4-64 uptake analysis

Average Z projections of single RH were taken with subsequent analysis of signal intensity within inner cell space (ImageJ - FIJI).

Ca2+ signalling reporter imaging and evaluation

A. thaliana seedlings expressing R-GECO1 Ca2+ signalling reporter were vertically grown in a growth chamber at 23 °C by day (16 h), 18 °C by night (8 h), 60% humidity and light intensity of 100 μmol photons m^{-2} s^{-1} (Keinath et al., 2015). Microfluidics experiments were performed using four-day-old seedlings placed into a

microfluidic chip and enclosed by a cover glass (Serre et al., 2021). Channel one contained a control solution (½MS, fluorescein dextran 6 μg/ml), while channel two contained a treatment solution (½MS, 0.01% or 0.001% chitosan). The media flow rate was 3 μl/min (OBI1, MFS2 Elveflow, and Elveflow software ESI (v.3.04.1). The system was mounted to a vertical microscope stage and kept for 15 min plant recovery. The media exchange was indicated by fluorescein-dextran signal intensity.

Plants were imaged using a vertical stage Zeiss Axio Observer 7 with a Yokogawa CSU-W1-T2 spinning disk unit with 50-μm pinholes equipped with a VS-HOM1000 excitation light homogenizer (Visitron Systems), objective Zeiss Plan-Apochromat ×10/0.45. Images were acquired using VisiView software (Visitron Systems, v.4.4.0.14). R-GECO1 was excited by a 561 nm laser, and fluorescein dextran by a 488 nm laser. The signal was detected using a PRIME-95B Back-Illuminated sCMOS camera (1,200 × 1,200 pixels; Photometrics). The seedlings were imaged every 13 s for 60 min. The signal intensity of R-GECO1 was measured inside the root at the root late elongation zone using ImageJ Fiji software (Schindelin et al., 2012). The intensity of the background was measured in areas not affected by the root response and subtracted from the R-GECO1 values. The intensity of fluorescein-dextran was measured outside of the root. The data were normalised using division by initial fluorescence intensity values (F/F0). As the initial value F0, we used the average fluorescence intensity at five-time points before treatment.

RNA isolation and RNAseq analysis

RNA was isolated from young, 6-7-day-old whole seedlings (transferred from vertical 1/2MS plates to liquid ½ MS for treatments), and adult (5-6 week old) rosette leaves of *A. thaliana* plants, using the RNeasy Plant kit (Qiagen) according to the manufacturer's instructions. Isolated RNAs were stabilized by GenTegra technology microtubes (GenTegra, Pleasanton, California, USA). Strand-specific cDNA libraries were constructed from polyA enriched RNA and sequenced on the Illumina NovaSeq6000 platform with subsequent analysis performed by Eurofins. The RNA-seq data used in this study are deposited in the National Center for Biotechnology Information Gene Expression Omnibus database.

Protein extraction, western blot and MAPK assay

Total protein extracts were isolated from 5–7-day-old *A. thaliana* seedlings according to procedure described in Fernandez et al., 2020. Briefly, untreated and chitosan treated tissue samples were ground in liquid nitrogen and dissolved in an extraction buffer containing Phosphostop, and the concentration of proteins

determined by the Bio-Rad protein assay kit with BSA as the standard. The extracts were denatured by boiling in 6× SDS loading buffer. The protein samples were separated by 10% SDS-PAGE and analysed by Western blot using the α-p44/42-ERK antibody (SAB4301578), anti-MAPK3 (M8318, Sigma), anti-MAPK4 (A6979, Sigma), and anti-MAPK6 (A7104, Sigma). The primary antibodies were incubated with the membranes for 3 h at room temperature in the blocking solution. Horseradish peroxidase-conjugated antibodies (anti-rabbit and anti-mouse; Promega) were applied followed by chemiluminescent ECL detection (Amersham) by the Bio-Rad documentation system. Using the Gel Analysis function of ImageJ, signal intensities for protein bands were determined for each treatment from 3 different samples. Loading consistency was examined by staining the membrane with Ponceau S.

Cell wall preparation and enzymatic fingerprinting

For cell wall analysis plants have been propagated in Jiffy pellets for 5-6 weeks, 10/14 h of light/dark regime. Plants were treated by dipping into the mock (distilled water with 0.005% Silwett) or LCC chitosan (distilled water with 0.005% Silwett and 0.001% chitosan), and left for 24 h in the growth chamber. Treated and non-treated samples were submerged in 96% (v/v) ethanol and boiled at 70°C for 10 min. The pellets were collected by centrifugation (13000 g for 10 min) and dried in a speed vacuum concentrator at 30°C overnight. Samples were digested with 1 U/mg DW sample of *Aspergillus aculeatus* endo-polygalacturonase M2 (Megazyme, Bray, Ireland) (Paterlini et al., 2022) in 50 mM ammonium acetate buffer (pH 5) at 37 °C for 18 h. Samples were centrifuged at 13000 rpm for 10 min and 100 µL of the supernatants were transferred into vials. The oligosaccharides released from digestion were analysed according to (Paterlini et al., 2022).

Bacterial flooding assay

Flooding assay was performed with modifications according to Ishiga et al. 2012 (Pečenková et al., 2020). Briefly, plants grown in vitro (½ MS, at 21°C, 12 / 12 h of light / dark per day, for 10 -12 days) were pretreated with mock (water with 0.0025% Silwet), or with chitosan (LCC, diluted in water with 0.0025% Silwet), both by flooding, and decanting after 2 min. After 24 h, the *P. syringae pv. tomato* DC3000 (Pst) OD = 0.05, was used for flooding for 2 min: the mock pretreated group was inoculated with either chitosan and Pst (chit + Pst), or with Pst only, together with the chitosan pretreated group (24 h chit + Pst) After the incubation, plants (three to four seedlings in one sample, four samples for each line, in three replications) were homogenized, homogenates subsequently diluted in series, plated put, and after approximately 36 h of plates incubation on 28 °C colonies counted.

Image analysis and statistics

Image processing software was employed for image data quantification. Roots and root hair sizes were analysed using AnalySIS (Soft Imaging System GmbH, Germany) or ImageJ (Schneider et al., 2012) software. The numerical data obtained were processed using Microsoft Excel.

To determine statistical significance, Student's t-test and ANOVA tests were conducted either using Excel or on-line calculators (http://in-silico.net, and http://statpages.info/anova1sm.html). P-values of less than 0·05 indicated a significant difference among the various groups. For multiple-comparison experiments, the Tukey Honestly Significant Difference (HSD) post hoc test was used.

Contribution

M.D. performed root hair confocal microscopy, designed the experimental setup for the real time observation of root hairs, performed microscopy and image analysis, prepared RNA for RNAseq analysis, prepared figures and wrote the manuscript. S. H. performed Selaginella and Marchantia experiments, and wrote the corresponding parts in the manuscript. E. Š. performed MAP kinases assays. P. K. and M. F. performed Ca2+-spiking recording and analysis. L. B. and J. O. prepared protoplasts and analysed their cell wall. K. M. and N. S. analysed RNAseq data. A. P. performed Pst susceptibility assays. A. V. and S. V. did the cell wall fingerprinting analysis. G. C. verified callose deposition in mutant lines. M. P. and V. Ž. planned the experiments and wrote the manuscript. T. P. performed root hair growth and callose deposition analysis, prepared figures and wrote the manuscript.

Acknowledgment

We would like to thank to Norbert Zlámal from the Bratislava`s Comenius University Botanical Garden for sharing *Selaginella uncinata*, Adriana Jelínková and Edita Janková Drdová (IEB CAS) for sharing Arabidopsis mutants and markers seeds, and Jana Šťovíčková for technical assistance. This work was supported by the Ministry of Education, Youth and Sports project CZ.02.1.01/0.0/0.0/16_019/0000738, 8J19FR001 MEYS Czech-French mobility program and Charles University Grant Agency (GAUK) project 415322.

 Legend to figures

Figure 1. Concentration dependent modes of chitosan-triggered callose deposition.
Different concentrations of chitosan cause different patterns of callose depositions 24
h post treatment (p. t.) in both cotyledons and roots (root hairs); mock treatment as a
control below (bar = 100 μm). B) Concentration dependent primary root growth
inhibition 24 h p. t. Small letters indicate significance of differences based on the
ANOVA test.

Figure 2. Root hair growth during the chitosan treatments. A) The low chitosan
concentration (0.001%, LCC) of several other chitosan variants has the same effect on
RH callose deposition (bar = 100 μm).

Figure 3. Callose deposition in LCC is PMR4-dependent. A) A *pmr4* mutant root hair
(RH) growth does not deposit callose upon the LCC treatment; nevertheless, the RH
growth is suppressed to the similar extent as in the Col-0 control. Small letters indicate
significance of differences based on the ANOVA test. B) Upon the LCC treatment, at 4
h p. t. GFP-PMR4 accumulates on the tips of growing RH, further confirming a role for
PMR4 in RHs callose deposition (marked by white asterisks). C) The time course of
decline of the RH growth during the LCC treatment; the significant growth inhibition
is observable at 12 h p.t. D) Timecourse of callose deposition reveals significant
changes between the mock and LCC already at 4 h p. t. Bars = 100 μm in A), C) and
D) and 10 μm in B).

Figure 4. Chitosan-triggered signalling in *A. thaliana* seedlings. A) Ca^{2+} spiking
analysed as R-GECO sensor fluorescence in root cells upon chitosan treatment shows
slightly later onset with mild oscillations and long duration within the ca 45 min frame
of LCC treatment, while HCC induces prominent Ca^{2+} spiking with subsequent drop
in signal intensity. B) Western blot analysis of MAPK activation by phosphorylation
upon 30 min of HCC and LCC treatments quantified as the ratio of band intensities
obtained for phosphorylated portion versus the whole pool MAPKs. Both treatments
cause significant increase in MPK3/6/4 phosphorylation in comparison to the mock
treatment, also the difference between the LCC- and HCC-caused phosphorylations is
significant, with LCC causing almost two-time less MAPKs phosphorylation.

Figure 5. R-GECO signal patterns upon mock, LCC and HCC treatment; below are
kymographs showing R-GECO dynamics within the selected RH (white rectangles in
upper row) in a time range of approximately 7 min after treatments; bars = 100 μm.

Figure 6. Subcellular dynamics during the early time points of LCC and HCC
treatments. A) The real-time analysis of the root hairs growth during the mock, LCC,

and HCC treatments based on the bright field images. The plot shows the RH growth rates for the mock, LCC and HCC treatments; small letters indicate significant differences based on Kruskal - Wallis test. B) The real-time changes in the vacuolar dynamics quantified as a vacuole-occupied area in the RH tip, imaged using the tonoplast marker Vti-GFP. The plot displays the differences between two consecutive frames; the examples of the differential signal are shown on the right; examples of vacuolar dynamics are shown below. Small letters indicate significant differences based on the ANOVA test. Bar = 20 µm. C) The analyses of the endocytosed FM4-64 dye after 15 min of treatments for mock, LCC, and HCC. The endocytic compartment stained by FM4-64 in the RH tip area is prominently enhanced in case of the HCC treatments. Small letters indicate significant differences based on Kruskal -Wallis test. On the right are examples of maximal intensity projections of FM4 -64 stained root hairs. Bar = 10 µm.

Figure 7. Callose deposition is conserved in structures analogous to root hairs. *M. polymorpha (liverwort)* rhizoids growing out of gemma are enclosed by callose when incubated for 24 h with LCC. The same effect is observable also on spikemoss rhizophores (*S. uncinata*). Bar = 100 µm.

Table I. Eleven genes (out of 58 DEGS; most of them with high RH expression) are specifically affected by LCC, when compared to HCC (out of 8508 DEGs), as shown by the RNAseq analysis performed on 6 d.o. plants cultivated in liquid medium (mock, LCC, HCC; 24h); 5 genes below the line in the Table are differentially expressed in LCC in comparison to HCC (fold change < 1 = downregulation, fold change > 1 = upregulation).

Supplementary figure 1. A) Cultivation in water does not favour the RH growth and callose deposition; the RH callose reaction requires a rich cultivation medium in order for RHs to grow, and callose deposits to be formed. B) The growth of new RH is inhibited in early time points (4 and 8 h p. t.) in high chitosan concentration (HCC; bars = 100 µm). C) Arabidopsis mutants in secretion, and defence and phytohormone signalling have ability to deposit root hair callose similarly to Col-0. Bars = 100 µm.

Supplementary figure 2. Actin dynamics is not significantly altered during the early time points of the HCC treatment, as observed using fimbrin-GFP marker and quantified using the variance between the two time points during the observation.

Supplementary figure 3. A) Numbers of specific and common DEGs among three different types of experiments with chitosan treatments. B) Cell wall changes triggerd by LCC treatment. Enzymatic fingerprinting of homogalacturonans (upper graph) and xyloglucans (lower graph) from cell walls of WT mock and LCC treated leaves. Data represents mean ±SD, n=4, *P<0.05, **P<0.01, student's t test. OGs are named GalAxMeyAcz. x, y and z indicate the DP and the number of methyl- and acetylester groups, respectively. GalA: Galacturonic acid, Me: methylester group, Ac: Acetylester group. Xyloglucan fragment nomenclature follows that described in Fry et al. (1993): G stands for glucose; X for xylose-glucose; L for galactose-xylose-glucose; F for fucose-galactose-xylose-glucose.

Supplementary figure 4. Chitosan LCC treatment induces deposition of callose in moss *P. patens* regenerating protoplasts (A) but not in leaf mesophyll-derived Arabidopsis regenerating protoplasts (B), as assayed by aniline blue staining. Bars = 100 μm.

Supplemental figure 5 A) LCC has prominent effect on amplification of bacteria *P. syringae pv. tomato* (Pst) in flooding experiment *in planta*, as chitosan/LCC added with Pst (chit + Pst) abolishes bacterial amplification in leaf tissue (red arrow points to "null" colony counts), while chitosan pretreatment 24 h prior to Pst inoculation (24 h chit + Pst) causes only small, but significant decrease in colony counts (upper graph); the effect of adding chitosan together with bacteria may not be a direct interference with bacteria amplification, as shown by the experiment of LCC treated Pst *in vitro* (lower graph). B) LCC has a minor effect on auxin signalling enhancement as demonstrated using the DR5::GFP reporter line (bars = 100 μm).

Bibliography

Aslam SN, Erbs G, Morrissey KL, Newman M-A, Chinchilla D, Boller T, Molinaro A, Jackson RW, Cooper RM. 2009. Microbe-associated molecular pattern (MAMP) signatures, synergy, size and charge: influences on perception or mobility and host defence responses. *Mol Plant Pathol* **10**:375–387. doi:10.1111/j.1364-3703.2009.00537.x

Badri DV, Loyola-Vargas VM, Du J, Stermitz FR, Broeckling CD, Iglesias-Andreu L, Vivanco JM. 2008. Transcriptome analysis of Arabidopsis roots treated with signaling compounds: A focus on signal transduction, metabolic regulation and secretion. *New Phytol* **179**:209–223. doi:10.1111/j.1469-8137.2008.02458.x

Baluška F, Salaj J, Mathur J, Braun M, Jasper F, Šamaj J, Chua NH, Barlow PW, Volkmann D. 2000. Root hair formation: F-actin-dependent tip growth is initiated by local assembly of profilin-supported F-actin meshworks accumulated within expansin-enriched bulges. *Dev Biol* **227**:618–632. doi:10.1006/dbio.2000.9908

Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA. 1996. Arabidopsis AUX1 gene: a permease-like regulator of root gravitropism. *Science* **273**:948–950. doi:10.1126/science.273.5277.948

Brejšková L, Hála M, Rawat A, Soukupová H, Cvrčková F, Charlot F, Nogué F, Haluška S, Žárský V. 2021. SEC6 exocyst subunit contributes to multiple steps of growth and development of Physcomitrella (Physcomitrium patens). *Plant J* **106**:831–843. doi:10.1111/tpj.15205

Brost C, Studtrucker T, Reimann R, Denninger P, Czekalla J, Krebs M, Fabry B, Schumacher K, Grossmann G, Dietrich P. 2019. Multiple cyclic nucleotide-gated channels coordinate calcium oscillations and polar growth of root hairs. *Plant J Cell Mol Biol* **99**:910–923. doi:10.1111/tpj.14371

Cai G, Faleri C, Del Casino C, Emons AMC, Cresti M. 2011. Distribution of callose synthase, cellulose synthase, and sucrose synthase in tobacco pollen tube is controlled in dissimilar ways by actin filaments and microtubules. *Plant Physiol* **155**:1169–1190. doi:10.1104/pp.110.171371

Cao H, Bowling SA, Gordon AS, Dong X. 1994. Characterization of an Arabidopsis Mutant That Is Nonresponsive to Inducers of Systemic Acquired Resistance. *Plant Cell* **6**:1583–1592. doi:10.1105/tpc.6.11.1583

Cao Y, Liang Y, Tanaka K, Nguyen CT, Jedrzejczak RP, Joachimiak A, Stacey G. 2014. The kinase LYK5 is a major chitin receptor in Arabidopsis and forms a chitin-induced complex with related kinase CERK1. *eLife* **3**:e03766. doi:10.7554/eLife.03766

Carol RJ, Dolan L. 2002. Building a hair: tip growth in Arabidopsis thaliana root hairs. *Philos Trans R Soc Lond B Biol Sci* **357**:815–821. doi:10.1098/rstb.2002.1092

Cavalier DM, Lerouxel O, Neumetzler L, Yamauchi K, Reinecke A, Freshour G, Zabotina OA, Hahn MG, Burgert I, Pauly M, Raikhel NV, Keegstra K. 2008. Disrupting two Arabidopsis thaliana xylosyltransferase genes results in plants deficient in xyloglucan, a major primary cell wall component. *Plant Cell* **20**:1519–1537. doi:10.1105/tpc.108.059873

Chin K, DeFalco TA, Moeder W, Yoshioka K. 2013. The Arabidopsis cyclic nucleotide-gated ion channels AtCNGC2 and AtCNGC4 work in the same signaling pathway to regulate pathogen defense and floral transition. *Plant Physiol* **163**:611–624. doi:10.1104/pp.113.225680

Chuberre C, Plancot B, Driouich A, Moore JP, Bardor M, Gügi B, Vicré M. 2018. Plant immunity is compartmentalized and specialized in roots. *Front Plant Sci* **871**:1692–1692. doi:10.3389/fpls.2018.01692

Cove DJ, Perroud P-F, Charron AJ, McDaniel SF, Khandelwal A, Quatrano RS. 2009. Culturing the moss Physcomitrella patens. *Cold Spring Harb Protoc* **2009**:pdb.prot5136. doi:10.1101/pdb.prot5136

Cui W, Lee J-Y. 2016. Arabidopsis callose synthases CalS1/8 regulate plasmodesmal permeability during stress. *Nat Plants* **2**:16034. doi:10.1038/nplants.2016.34

Dehors J, Mareck A, Kiefer-Meyer M-C, Menu-Bouaouiche L, Lehner A, Mollet J-C. 2019. Evolution of Cell Wall Polymers in Tip-Growing Land Plant Gametophytes: Composition, Distribution,

Functional Aspects and Their Remodeling. *Front Plant Sci* **10**:441.
doi:10.3389/fpls.2019.00441

Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Ferrari S, Ausubel FM,
Dewdney J. 2008. Activation of defense response pathways by OGs and Flg22 elicitors in
Arabidopsis seedlings. *Mol Plant* **1**:423–445. doi:10.1093/mp/ssn019

Dolan L. 1996. Pattern in the root epidermis: An interplay of diffusible signals and cellular geometry.
Ann Bot **77**:547–553. doi:10.1006/anbo.1996.0069

Epel BL. 2009. Plant viruses spread by diffusion on ER-associated movement-protein-rafts through
plasmodesmata gated by viral induced host beta-1,3-glucanases. *Semin Cell Dev Biol*
20:1074–1081. doi:10.1016/j.semcdb.2009.05.010

Galway ME, Eng RC, Schiefelbein JW, Wasteneys GO. 2011. Root hair-specific disruption of cellulose
and xyloglucan in AtCSLD3 mutants, and factors affecting the post-rupture resumption of
mutant root hair growth. *Planta* **233**:985–999. doi:10.1007/s00425-011-1355-6

Gamborg OL, Miller RA, Ojima K. 1968. Nutrient requirements of suspension cultures of soybean root
cells. *Exp Cell Res* **50**:151–158. doi:10.1016/0014-4827(68)90403-5

German L, Yeshvekar R, Benitez-Alfonso Y. 2023. Callose metabolism and the regulation of cell walls
and plasmodesmata during plant mutualistic and pathogenic interactions. *Plant Cell Environ*
46:391–404. doi:10.1111/pce.14510

Gómez-Gómez L, Felix G, Boller T. 1999. A single locus determines sensitivity to bacterial flagellin in
Arabidopsis thaliana. *Plant J Cell Mol Biol* **18**:277–284. doi:10.1046/j.1365-313x.1999.00451.x

Heisler MG, Ohno C, Das P, Sieber P, Reddy GV, Long JA, Meyerowitz EM. 2005. Patterns of auxin
transport and gene expression during primordium development revealed by live imaging of
the Arabidopsis inflorescence meristem. *Curr Biol CB* **15**:1899–1911.
doi:10.1016/j.cub.2005.09.052

Herburger K, Holzinger A. 2015. Localization and Quantification of Callose in the Streptophyte Green
Algae Zygnema and Klebsormidium: Correlation with Desiccation Tolerance. *Plant Cell Physiol*
56:2259–2270. doi:10.1093/pcp/pcv139

Ichikawa M, Hirano T, Enami K, Fuselier T, Kato N, Kwon C, Voigt B, Schulze-Lefert P, Baluška F, Sato
MH. 2014. Syntaxin of plant proteins SYP123 and SYP132 mediate root hair tip growth in
Arabidopsis thaliana. *Plant Cell Physiol* **55**:790–800. doi:10.1093/pcp/pcu048

Jacobs AK, Lipka V, Burton RA, Panstruga R, Strizhov N, Schulze-Lefert P, Fincher GB. 2003. An
Arabidopsis Callose Synthase, GSL5, Is Required for Wound and Papillary Callose Formation.
Plant Cell **15**:2503–2513. doi:10.1105/tpc.016097

Jones JDG, Dangl JL. 2006. The plant immune system. *Nature* **444**:323–329. doi:10.1038/nature05286

Keinath NF, Waadt R, Brugman R, Schroeder JI, Grossmann G, Schumacher K, Krebs M. 2015. Live Cell
Imaging with R-GECO1 Sheds Light on flg22- and Chitin-Induced Transient [Ca2+]cyt Patterns
in Arabidopsis. *Mol Plant* **8**:1188–1200. doi:10.1016/j.molp.2015.05.006

Kemmerling B, Schwedt A, Rodriguez P, Mazzotta S, Frank M, Qamar SA, Mengiste T, Betsuyaku S,
Parker JE, Müssig C, Thomma BPHJ, Albrecht C, de Vries SC, Hirt H, Nürnberger T. 2007. The
BRI1-associated kinase 1, BAK1, has a brassinolide-independent role in plant cell-death
control. *Curr Biol CB* **17**:1116–1122. doi:10.1016/j.cub.2007.05.046

Kentaro Okada, Koei Yachi, Tan Anh Nhi Nguyen, Satomi Kanno, Chika Tateda, Tae-Hong Lee, Uyen
Nguyen, Shunsuke Miyashima, Kei Hiruma, Kyoko Miwa, Takaki Maekawa, Michitaka
Notaguchi, Yusuke Saijo. 2023. Defense-related callose synthase *PMR4* promotes
root hair callose deposition and adaptation to phosphate deficiency in *Arabidopsis
thaliana*. *bioRxiv* 2023.07.05.547890. doi:10.1101/2023.07.05.547890

Ks R, Ra L, Rd W, Ha O, A M. 2020. Callose in sporogenesis: Novel composition of the inner spore wall
in hornworts. *Plant Syst Evol Entwicklungsgeschichte Syst Pflanz* **306**:16. doi:10.1007/s00606-
020-01631-5

Kubátová Z, Pejchar P, Potocký M, Sekereš J, Žárský V, Kulich I. 2019. Arabidopsis Trichome Contains
Two Plasma Membrane Domains with Different Lipid Compositions Which Attract Distinct
EXO70 Subunits. *Int J Mol Sci* **20**:3803. doi:10.3390/ijms20153803

Kulich I, Vojtíková Z, Glanc M, Ortmannová J, Rasmann S, Žárský V. 2015. Cell wall maturation of
Arabidopsis trichomes is dependent on exocyst subunit EXO70H4 and involves callose
deposition. *Plant Physiol* **168**:120–131. doi:10.1104/pp.15.00112

Kulich I, Vojtíková Z, Sabol P, Ortmannová J, Neděla V, Tihlaříková E, Žárský V. 2018. Exocyst Subunit
EXO70H4 Has a Specific Role in Callose Synthase Secretion and Silica Accumulation. *Plant
Physiol* **176**:2040–2051. doi:10.1104/pp.17.01693

Ladwig F, Dahlke RI, Stührwohldt N, Hartmann J, Harter K, Sauter M. 2015. Phytosulfokine regulates
growth in arabidopsis through a response module at the plasma membrane that includes
cyclic nucleotide-gated channel17, H+-ATPase, and BAK1. *Plant Cell* **27**:1718–1729.
doi:10.1105/tpc.15.00306

Lopez-Moya F, Escudero N, Zavala-Gonzalez EA, Esteve-Bruna D, Blázquez MA, Alabadí D, Lopez-
Llorca LV. 2017. Induction of auxin biosynthesis and WOX5 repression mediate changes in
root development in Arabidopsis exposed to chitosan. *Sci Rep* **7**:16813. doi:10.1038/s41598-
017-16874-5

Luna E, Pastor V, Robert J, Flors V, Mauch-Mani B, Ton J. 2011. Callose deposition: a multifaceted
plant defense response. *Mol Plant-Microbe Interact MPMI* **24**:183–193. doi:10.1094/MPMI-
07-10-0149

Ma X, Wu Y, Zhang G. 2021. Formation pattern and regulatory mechanisms of pollen wall in
Arabidopsis. *J Plant Physiol* **260**:153388. doi:10.1016/j.jplph.2021.153388

Meyer D, Pajonk S, Micali C, O'Connell R, Schulze-Lefert P. 2009. Extracellular transport and
integration of plant secretory proteins into pathogen-induced cell wall compartments. *Plant J
Cell Mol Biol* **57**:986–999. doi:10.1111/j.1365-313X.2008.03743.x

Millet YA, Danna CH, Clay NK, Songnuan W, Simon MD, Werck-Reichhart D, Ausubel FM. 2010. Innate
immune responses activated in Arabidopsis roots by microbe-associated molecular patterns.
Plant Cell **22**:973–990. doi:10.1105/tpc.109.069658

Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, Shirasu K, Narusaka Y, Kawakami N, Kaku H, Shibuya
N. 2007. CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in Arabidopsis.
Proc Natl Acad Sci U S A **104**:19613–19618. doi:10.1073/pnas.0705147104

Motomura K, Sugi N, Takeda A, Yamaoka S, Maruyama D. 2022. Possible molecular mechanisms of
persistent pollen tube growth without de novo transcription. *Front Plant Sci* **13**:1020306.
doi:10.3389/fpls.2022.1020306

Ngou BPM, Ahn H-K, Ding P, Jones JDG. 2021. Mutual potentiation of plant immunity by cell-surface
and intracellular receptors. *Nature* **592**:110–115. doi:10.1038/s41586-021-03315-7

Okada K, Kubota Y, Hirase T, Otani K, Goh T, Hiruma K, Saijo Y. 2021. Uncoupling root hair formation
and defence activation from growth inhibition in response to damage-associated Pep
peptides in Arabidopsis thaliana. *New Phytol* **229**:2844–2858. doi:10.1111/nph.17064

Ortmannová J, Sekereš J, Kulich I, Šantrůček J, Dobrev P, Žárský V, Pečenková T. 2022. Arabidopsis
EXO70B2 exocyst subunit contributes to papillae and encasement formation in antifungal
defence. *J Exp Bot* **73**:742–755. doi:10.1093/jxb/erab457

Park K, Jang BK, Lee HM, Cho JS, Lee CH. 2020. An Efficient Method for In Vitro Shoot-Tip Culture and
Sporophyte Production Using Selaginella martensii Spring Sporophyte. *Plants Basel Switz
***9**:235. doi:10.3390/plants9020235

Park S, Szumlanski AL, Gu F, Guo F, Nielsen E. 2011. A role for CSLD3 during cell-wall synthesis in
apical plasma membranes of tip-growing root-hair cells. *Nat Cell Biol* **13**:973–980.
doi:10.1038/ncb2294

Parker JS, Cavell AC, Dolan L, Roberts K, Grierson CS. 2000. Genetic interactions during root hair
morphogenesis in Arabidopsis. *Plant Cell* **12**:1961–1974. doi:10.1105/tpc.12.10.1961

Paterlini A, Sechet J, Immel F, Grison MS, Pilard S, Pelloux J, Mouille G, Bayer EM, Voxeur A. 2022.
Enzymatic fingerprinting reveals specific xyloglucan and pectin signatures in the cell wall
purified with primary plasmodesmata. *Front Plant Sci* **13**.

Pečenková T, Janda M, Ortmannová J, Hajná V, Stehlíková Z, Žárský V. 2017. Early Arabidopsis root
hair growth stimulation by pathogenic strains of Pseudomonas syringae. *Ann Bot* **120**:437–

446. doi:10.1093/aob/mcx073

Porter K, Day B. 2016. From filaments to function: The role of the plant actin cytoskeleton in
pathogen perception, signaling and immunity. *J Integr Plant Biol* **58**:299–311.
doi:10.1111/jipb.12445

Radin I, Richardson RA, Coomey JH, Weiner ER, Bascom CS, Li T, Bezanilla M, Haswell ES. 2021. Plant
PIEZO homologs modulate vacuole morphology during tip growth. *Science* **373**:586–590.
doi:10.1126/science.abe6310

Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS,
Hirt H, Knight MR. 2004. OXI1 kinase is necessary for oxidative burst-mediated signalling in
Arabidopsis. *Nature* **427**:858–861. doi:10.1038/nature02353

Rich-Griffin C, Eichmann R, Reitz MU, Hermann S, Woolley-Allen K, Brown PE, Wiwatdirekkul K,
Esteban E, Pasha A, Kogel KH, Provart NJ, Ott S, Schäfer P. 2020. Regulation of cell type-
specific immunity networks in arabidopsis roots. *Plant Cell* **32**:2742–2762.
doi:10.1105/TPC.20.00154

Ringli C, Baumberger N, Diet A, Frey B, Keller B. 2002. ACTIN2 is essential for bulge site selection and
tip growth during root hair development of Arabidopsis. *Plant Physiol* **129**:1464–1472.
doi:10.1104/pp.005777

Serre NBC, Kralík D, Yun P, Slouka Z, Shabala S, Fendrych M. 2021. AFB1 controls rapid auxin
signalling through membrane depolarization in Arabidopsis thaliana root. *Nat Plants* **7**:1229–
1238. doi:10.1038/s41477-021-00969-z

Shih HW, Depew CL, Miller ND, Monshausen GB. 2015. The cyclic nucleotide-gated channel CNGC14
regulates root gravitropism in arabidopsis thaliana. *Curr Biol* **25**:3119–3125.
doi:10.1016/j.cub.2015.10.025

Sun H, Zhu X, Li C, Ma Z, Han X, Luo Y, Yang L, Yu J, Miao Y. 2021. Xanthomonas effector XopR hijacks
host actin cytoskeleton via complex coacervation. *Nat Commun* **12**:4064.
doi:10.1038/s41467-021-24375-3

Suwanchaikasem P, Idnurm A, Selby-Pham J, Walker R, Boughton BA. 2022. Root-TRAPR: a modular
plant growth device to visualize root development and monitor growth parameters, as
applied to an elicitor response of Cannabis sativa. *Plant Methods* **18**:46. doi:10.1186/s13007-
022-00875-1

Vaškebová L, Šamaj J, Ovecka M. 2018. Single-point ACT2 gene mutation in the Arabidopsis root hair
mutant der1-3 affects overall actin organization, root growth and plant development. *Ann
Bot* **122**:889–901. doi:10.1093/aob/mcx180

Verma DP, Hong Z. 2001. Plant callose synthase complexes. *Plant Mol Biol* **47**:693–701.
doi:10.1023/a:1013679111111

Vissenberg K, Claeijs N, Balcerowicz D, Schoenaers S. 2020. Hormonal regulation of root hair growth
and responses to the environment in Arabidopsis. *J Exp Bot* **71**:2412–2427.
doi:10.1093/JXB/ERAA048

Vogel J, Somerville S. 2000. Isolation and characterization of powdery mildew-resistant Arabidopsis
mutants. *Proc Natl Acad Sci U S A* **97**:1897–1902. doi:10.1073/pnas.030531997

Voxeur A, Habrylo O, Guénin S, Miart F, Soulié M-C, Rihouey C, Pau-Roblot C, Domon J-M, Gutierrez
L, Pelloux J, Mouille G, Fagard M, Höfte H, Vernhettes S. 2019. Oligogalacturonide production
upon Arabidopsis thaliana-Botrytis cinerea interaction. *Proc Natl Acad Sci U S A* **116**:19743–
19752. doi:10.1073/pnas.1900317116

Wan J, Tanaka K, Zhang X-C, Son GH, Brechenmacher L, Nguyen THN, Stacey G. 2012. LYK4, a lysin
motif receptor-like kinase, is important for chitin signaling and plant innate immunity in
Arabidopsis. *Plant Physiol* **160**:396–406. doi:10.1104/pp.112.201699

Wang X, Wang K, Yin G, Liu X, Liu M, Cao N, Duan Y, Gao H, Wang W, Ge W, Wang J, Li R, Guo Y. 2018.
Pollen-Expressed Leucine-Rich Repeat Extensins Are Essential for Pollen Germination and
Growth. *Plant Physiol* **176**:1993–2006. doi:10.1104/pp.17.01241

Wang Y-S, Motes CM, Mohamalawari DR, Blancaflor EB. 2004. Green fluorescent protein fusions to
Arabidopsis fimbrin 1 for spatio-temporal imaging of F-actin dynamics in roots. *Cell Motil*

Cytoskeleton **59**:79–93. doi:10.1002/cm.20024

Ye W, Munemasa S, Shinya T, Wu W, Ma T, Lu J, Kinoshita T, Kaku H, Shibuya N, Murata Y. 2020.
Stomatal immunity against fungal invasion comprises not only chitin-induced stomatal
closure but also chitosan-induced guard cell death. *Proc Natl Acad Sci U S A* **117**:20932–
20942. doi:10.1073/pnas.1922319117

Yoo S-D, Cho Y-H, Sheen J. 2007. Arabidopsis mesophyll protoplasts: a versatile cell system for
transient gene expression analysis. *Nat Protoc* **2**:1565–1572. doi:10.1038/nprot.2007.199

Yuan P, Jauregui E, Du L, Tanaka K, Poovaiah BW. 2017. Calcium signatures and signaling events
orchestrate plant–microbe interactions. *Curr Opin Plant Biol* **38**:173–183.
doi:10.1016/j.pbi.2017.06.003

Zamioudis C, Mastranesti P, Dhonukshe P, Blilou I, Pieterse CMJ. 2013. Unraveling root
developmental programs initiated by beneficial Pseudomonas spp. Bacteria. *Plant Physiol*
162:304–318. doi:10.1104/pp.112.212597

Zhang J, Li Y, Bao Q, Wang H, Hou S. 2022. Plant elicitor peptide 1 fortifies root cell walls and triggers
a systemic root-to-shoot immune signaling in Arabidopsis. *Plant Signal Behav* **17**:2034270.
doi:10.1080/15592324.2022.2034270

Zhang M, Zhang S. 2022. Mitogen-activated protein kinase cascades in plant signaling. *J Integr Plant*
Biol **64**:301–341. doi:10.1111/jipb.13215

Figure 1. Concentration dependent modes of chitosan-triggered callose deposition. Different concentrations of chitosan cause different patterns of callose depositions 24 h post treatment (p. t.) in both cotyledons and roots (root hairs); mock treatment as a control below (bar = 100 μm). B) Concentration dependent primary root growth inhibition 24 h p. t. Small letters indicate significance of differences based on the ANOVA test.

Figure 2. Root hair growth during the chitosan treatments. A) The low chitosan concentration (0.001%, LCC) of several other chitosan variants has the same effect on RH callose deposition (bar = 100 μm).

Figure 3. Callose deposition in LCC is PMR4-dependent. A) A *pmr4* mutant root hair (RH) growth does not deposit callose upon the LCC treatment; nevertheless, the RH growth is suppressed to the similar extent as in the Col-0 control. Small letters indicate significance of differences based on the ANOVA test. B) Upon the LCC treatment, at 4 h p. t. GFP-PMR4 accumulates on the tips of growing RH, further confirming a role for PMR4 in RHs callose deposition (marked by white asterisks). C) The time course of decline of the RH growth during the LCC treatment; the significant growth inhibition is observable at 12 h p.t. D) Timecourse of callose deposition reveals significant changes between the mock and LCC already at 4 h p. t. Bars = 100 μm in A), C) and D) and 10 μm in B).

Figure 4. Chitosan-triggered signalling in *A. thaliana* seedlings. A) Ca²⁺ spiking analysed as R-GECO sensor fluorescence in root cells upon chitosan treatment shows slightly later onset with mild oscillations and long duration within the ca 45 min frame of LCC treatment, while HCC induces prominent Ca²⁺ spiking with subsequent drop in signal intensity. B) Western blot analysis of MAPK activation by phosphorylation upon 30 min of HCC and LCC treatments quantified as the ratio of band intensities obtained for phosphorylated portion versus the whole pool MAPKs. Both treatments cause significant increase in MPK3/6/4 phosphorylation in comparison to the mock treatment, also the difference between the LCC- and HCC-caused phosphorylations is significant, with LCC causing almost two-time less MAPKs phosphorylation.

Figure 5. R-GECO signal patterns upon mock, LCC and HCC treatment; below are kymographs showing R-GECO dynamics within the selected RH (white rectangles in upper row) in a time range of approximately 7 min after treatments; bars = 100 μm.

Figure 6. Subcellular dynamics during the early time points of LCC and HCC treatments. A) The real-time analysis of the root hairs growth during the mock, LCC, and HCC treatments based on the bright field images. The plot shows the RH growth rates for the mock, LCC and HCC treatments; small letters indicate significant differences based on Kruskal - Wallis test. B) The real-time changes in the vacuolar dynamics quantified as a vacuole-occupied area in the RH tip, imaged using the tonoplast marker Vti-GFP. The plot displays the differences between two consecutive frames; the examples of the differential signal are shown on the right; examples of vacuolar dynamics are shown below. Small letters indicate significant differences based on the ANOVA test. Bar = 20 μm. C) The analyses of the endocytosed FM4-64 dye after 15 min of treatments for mock, LCC, and HCC. The endocytic compartment stained by FM4-64 in the RH tip area is prominently enhanced in case of the HCC treatments. Small letters indicate significant differences based on Kruskal -Wallis test. On the right are examples of maximal intensity projections of FM4 -64 stained root hairs. Bar = 10 μm.

mock	LCC	HCC	
			M. polymorpha
			S. uncinata

Figure 7. Callose deposition is conserved in structures analogous to root hairs. *M. polymorpha (liverwort)* rhizoids growing out of gemma are enclosed by callose when incubated for 24 h with LCC. The same effect is observable also on spikemoss rhizophores (*S. uncinata*). Bar = 100 μm.

Table I. Eleven genes (out of 59 DEGS; most of them with good RH expression) are specifically affected by LCC, when compared to HCC chitosan treatment (out of 8508 DEGs); 5 genes below the line in the Table are differentially expressed in LCC in comparison to HCC (fold change < 1 = downregulation, fold change > 1 = upregulation).

gene identifier	gene name	protein function	fold change LCC	fold change HCC
AT1G72900	ERF11	ERF domain protein 11	0.49	insig.
AT2G22500	DIC1	uncoupling protein 5	0.59	insig.
AT2G46400	WRKY46	DNA-binding protein 46	0.36	insig.
AT3G22800		leucine-rich repeat (LRR)/extensin	1.85	insig.
AT3G56400	WRKY70	WRKY DNA-binding protein 70	0.48	insig.
AT4G08950	EXO	phosphate-responsive 1 family protein	0.70	insig.
AT4G11280	ACS6	1-aminocyclopropane-1-carboxylic acid (ACC) synthase 6	0.47	insig.
AT4G24380		involved in 10-formyltetrahydrofolate biosynthesis	0.42	insig.
AT5G41750		disease resistance protein (TIR-NBS-LRR class) family	0.42	insig.
AT5G59550		zinc finger (C3HC4-type RING finger) family protein	0.63	insig.
AT5G64870		SPFH/Band 7/PHB domain-containing, flotillin	0.49	insig.
AT1G72900		Toll-Interleukin-Resistance (TIR) domain-containing protein	0.57	2.95
AT2G16660		major facilitator superfamily protein	0.52	1.30
AT2G34930		disease resistance family protein LRR	0.62	1.79
AT3G50930	BCS1	cytochrome BC1 synthesis	0.44	2.12
AT5G42380	CML37	calmodulin like 37	0.32	2.50