Smith-Magenis syndrome protein Rai1 regulates body weight homeostasis through

hypothalamic Bdnf-producing neurons and TrkB signalling

Sehrish Javed[1,2], Ya-Ting Chang[1,2], Yoobin Cho[1,2], Yu-Ju Lee[1,2], Hao-Cheng Chang[1,2], Minza

Haque[1,2], Yu Cheng Lin[1,2], Wei-Hsiang Huang[1,2],*

Affiliations:

[1]Department of Neurology and Neurosurgery, Centre for Research in Neuroscience, McGill

University, Montréal, QC H3G 1A3, Canada

[2]Brain Repair and Integrative Neuroscience Program, The Research Institute of the McGill

University Health Centre, Montréal, QC H3G 1A3, Canada

*Corresponding author. Email: wei-hsiang.huang@mcgill.ca

Abstract

Retinoic acid-induced 1 (*RAI1*) haploinsufficiency causes Smith-Magenis syndrome (SMS), a genetic disorder with symptoms including hyperphagia, hyperlipidemia, severe obesity, and autism phenotypes. Rai1 is a transcriptional regulator with a pan-neural expression pattern and hundreds of downstream targets. The mechanisms linking neural *Rai1* to body weight regulation remain unclear. Here we find that hypothalamic brain-derived neurotrophic factor (Bdnf)-TrkB signalling is disrupted in SMS (*Rai1*$^{+/-}$) mice. Selective Rai1 loss from all Bdnf-producing cells or from Bdnf-producing neurons in the paraventricular nucleus of the hypothalamus (PVH) induced obesity in mice. Electrophysiological recordings revealed that *Rai1* ablation increased inhibitory synaptic transmission to PVH^{Bdnf} neurons and decreased intrinsic neuronal excitability. Chronic treatment of SMS mice with a partial agonist of tropomyosin receptor kinase B (TrkB), the cognate Bdnf receptor, delayed obesity onset. This treatment also partially rescued disrupted lipid profiles, insulin intolerance, and stereotypical repetitive behaviour in SMS mice. These data argue that Rai1 regulates body weight and metabolic function through hypothalamic Bdnf-producing neurons and that targeting TrkB signalling might improve associated SMS phenotypes.

Introduction

Obesity is an increasing global concern: 20% of the world's adult population is expected to become obese by 2025 if current trends persist (Collaboration, 2017). However, the complex environmental, behavioural, and genetic contributions to obesity make this a difficult condition to model and address. Monogenic obesity disorders provide a consistent system where the pathophysiology, genetic predisposition, and neurobiology underlying phenotypes can potentially provide insights that translate to polygenic obesity (Loos & Yeo, 2022).

One such monogenic disorder is Smith-Magenis syndrome (SMS; OMIM #182290) (Smith *et al*, 1986), a neurogenetic condition frequently associated with obesity. 80% of SMS cases are caused by interstitial deletions of the 17p11.2 chromosomal region containing a dosage-sensitive gene, *retinoic acid-induced 1* (*RAI1*) (Slager *et al*, 2003). More than 90% of SMS patients are overweight (40% > 95th percentile weight, 40% > 85th percentile weight) and exhibit over-eating tendencies by adolescence (Alaimo *et al*, 2015; Burns *et al*, 2010; Gandhi *et al*, 2022). 20% of SMS patients do not have 17p11.2 deletions but instead carry *RAI1* point mutations that are still frequently associated with obesity (Boot *et al*, 2021; Edelman *et al*, 2007). These clinical data indicate that RAI is critical for energy homeostasis underlying body weight regulation.

The human RAI1 protein shares >80% sequence identity with the mouse Rai1 protein, allowing us to study Rai1 function *in vivo* (Huang *et al*, 2016; Javed *et al*, 2020). We previously found that *Rai1* encodes a transcriptional regulator that binds to the promoter or enhancer regions of its target genes, most of which are involved in neurodevelopment and synaptic transmission (Huang *et al.*, 2016). SMS (*Rai1*[+/-]) mice exhibit obesity associated with hyperphagia and increased adiposity (Burns *et al.*, 2010; Huang *et al.*, 2016). Interestingly, Rai1 may have a sexually

dimorphic function, as female SMS mice exhibit more pronounced obesity than males (Burns *et*

al., 2010; Huang *et al.*, 2016).

Using a homozygous *Rai1* conditional knockout (*Rai1*-cKO) mice, we found that *Rai1* deletion

in either subcortical excitatory neurons (vGlut2$^+$) or single-minded homolog 1 (Sim1$^+$) neurons

induced obesity in mice (Huang *et al.*, 2016). Sim1$^+$ neurons reside in several brain regions

including the nucleus of the lateral olfactory tract, the supraoptic nucleus, the medial amygdala,

and the paraventricular nucleus of the hypothalamus (PVH), a region that is critical for body

weight regulation (Balthasar *et al*, 2005). Sim1$^+$ neurons in the PVH are composed of several

molecularly defined cell types including those that express oxytocin, vasopressin, corticotropin-

releasing hormone, somatostatin, melanocortin-4-receptor (MC4R), growth hormone-releasing

hormone (GHRH), or brain-derived neurotrophic factor (Bdnf) (An *et al*, 2015). Deleting *Bdnf*

from the PVH results in hyperphagic obesity and reduced locomotion (An *et al.*, 2015). While

PVH neuronal activity is reduced during feeding (Xu *et al*, 2019), activation of PVH^{Bdnf} neurons

results in weight loss (Wu & Xu, 2022). We recently found that Rai1 promotes *Bdnf* expression

in hypothalamus throughout life (Javed *et al*, 2021). We hypothesized that Rai1 may be

connected to Bdnf-producing neurons in PVH, explaining how a neural transcription factor

regulates body weight homeostasis.

Bdnf regulates energy metabolism by binding to its cognate receptor tropomyosin receptor

kinase B (TrkB) (An *et al.*, 2015) and may thus provide a potential therapeutic target for SMS.

However, the link between *Rai1,* Bdnf, and SMS symptoms is currently unclear. Patients

carrying monogenic mutations in either *BDNF* (Gray *et al*, 2006) or *NTRK2* (encoding TrkB)

(Yeo *et al*, 2004) experience severe hyperphagic obesity. Moreover, obesity and associated

metabolic dysfunction such as impaired glucose and insulin sensitivity are correlated with

impaired PI3K-Akt pathway, a TrkB downstream target (Su *et al*, 2021). Given that (1) *Bdnf*

overexpression during early embryonic development or (2) targeting *Bdnf* overexpression to the

PVH of adolescent SMS ($Rai1^{+/-}$) mice protected them from weight gain and metabolic defects

(Javed *et al.*, 2021), and that (3) a point mutation in *RAI1* disrupts its ability to transcriptionally

enhance *Bdnf* expression *in vitro* (Abad *et al*, 2018), pharmacological targeting of Bdnf signaling

is an attractive and yet unexplored therapeutic avenue for SMS.

Building on our previous discoveries, here we examine the neurobiological functions of Rai1 in

Bdnf-producing neurons, specifically those that reside in the PVH. Importantly, we evaluated the

involvement of the Bdnf-TrkB pathways in SMS pathogenesis using a small molecule drug

(LM22A-4) to enhance TrkB signalling. Our work indicates that PVH^{Bdnf} neurons depend on

Rai1 to maintain normal neuronal activity and regulate body weight homeostasis, and TrkB

signalling dysfunction contributes to obesity associated with SMS mice.

Results

Rai1 deficiency dysregulates specific downstream targets of neurotrophin in the mouse hypothalamus

Multiple lines of evidence indicated that SMS mice show decreased *Bdnf* mRNA expression in the hypothalamus (Burns *et al.*, 2010; Huang *et al.*, 2016; Javed *et al.*, 2021). To determine if Bdnf protein levels were consequently altered in the hypothalamus, we performed enzyme-linked immunosorbent assay (ELISA) and found that Bdnf protein levels were significantly reduced in 7-week-old pre-symptomatic SMS mice relative to controls (**Fig 1A**). To quantify if molecular pathways associated with Bdnf-TrkB signalling were altered, we subjected hypothalamic tissue lysates from 7-week-old pre-obese SMS and control (Ctrl) mice to reverse phase protein array (RPPA) (**Fig 1B**). We found that control and SMS samples clustered separately in principle component analysis (**Fig 1C**).

Hierarchical clustering analysis indicated that the majority of differentially expressed phosphorylated proteins (**Fig 1D**) and total proteins (**Figure 1—figure supplement 1**) were downregulated in SMS mice. We mapped all proteins, as well as phospho-proteins, that were differentially expressed in Ctrl and SMS samples onto the interconnected biological network and identified a subset of connected nodes downregulated in SMS (FDR < 10e-5) belong to the neurotrophin, PI3K-Akt, and mTOR signalling pathways (**Fig 1E**). These pathways are known to play a critical role in regulating body weight homeostasis and can be activated by Bdnf either directly (neurotrophin pathway) or indirectly (PI3K-Akt and mTOR pathways) (Cota *et al*, 2006; Schultze *et al*, 2012). The expression levels of PIK3CA, a common node in these pathways, were significantly reduced in the SMS mice compared to controls (**Fig 1F**). Moreover, the expression levels of phosphorylated NF-κB (nuclear factor-kappa B), a downstream target of the

neurotrophin pathway, were significantly reduced in the SMS group (**Fig 1G**). Together, these

results suggest that *Rai1* loss impairs specific pathways, including neurotrophin signalling, and

indicate that dysfunction of Bdnf-producing cells could contribute to SMS pathogenesis.

Bdnf-producing cells depend on Rai1 to regulate body weight homeostasis

To target Bdnf-producing cells, we used a $Bdnf^{Cre/+}$ mouse line that carries a viral 2A

oligopeptide and Cre recombinase immediately upstream of Bdnf's endogenous STOP codon

(Tan *et al*, 2016). We crossed the $Bdnf^{Cre/+}$ mice with tdTomato-Ai9 reporter mice and found that

Ai9 signals faithfully reflected endogenous Bdnf protein expression in the PVH (**Figure 2—**

figure supplement 1A-E). More than 50% of Bdnf-producing cells within the PVH were

positive for phosphorylated TrkB (p-TrkB) (**Figure 2—figure supplement 1F**). Moreover, 20%

of p-TrkB^{+} PVH cells express Bdnf, suggesting that Bdnf could have an autocrine mode of action

in this cell type (**Figure 2—figure supplement 1F-J**). We also found that Rai1 is expressed in

PVH^{Bdnf} cells that are distinct from larger magnocellular cells such as the PVH oxytocin neurons

(**Figure 2—figure supplement 2A-I**). Specifically, 55% of PVH^{Bdnf} neurons express Rai1,

whereas only 2% of PVH^{Oxytocin} neurons express Rai1 (**Figure 2—figure supplement 2I**). These

data suggest that Rai1 is enriched in Bdnf-producing neurons in the mouse PVH.

To determine if Bdnf-producing cells depend on Rai1 to regulate body weight, we deleted one or

both copies of *Rai1* using the $Bdnf^{Cre/+}$ and the conditional $Rai1^{fl/fl}$ mice (Huang *et al.*, 2016) (**Fig**

2A, Figure 2—figure supplement 3A). Conditional knockout mice (cKO: $Bdnf^{Cre/+}$; $Rai1^{fl/fl}$)

showed a significant reduction in Rai1 expression in PVH^{Bdnf} neurons compared to controls (**Fig**

2B-C, Figure 2—figure supplement 3B). The total number of Bdnf-producing cells in the PVH

remained similar in Ctrl and cKO mice (**Figure 2—figure supplement 3C**), indicating that Rai1

loss did not affect PVH neurogenesis or survival.

We monitored body weight weekly and found that male and female cKO mice showed a

significant increase in body weight beginning at 15 weeks of age when compared to controls

(Ctrl: $Bdnf^{Cre/+}$) (**Fig 2D**, **Figure 2—figure supplement 4A**). However, conditional *Rai1*

heterozygous mice ($Bdnf^{Cre/+}$; $Rai1^{fl/+}$) did not show a pathological increase in body weight

(**Figure 2—figure supplement 3D**, **Figure 2—figure supplement 4B**), suggesting that non-

Bdnf-producing cells also contribute to SMS pathogenesis. EcoMRI found that 26-week-old

cKO male and female mice showed an increased proportion of fat mass but not lean mass (**Fig**

2E, **Figure 2—figure supplement 4C**). Further dissection of fat tissues from different organs

showed an increased mass of subcutaneous inguinal (S.C. Ing.) fat in cKO mice of both sexes

and increased mass of epididymal adipose tissues (eWAT) in female cKOs (**Fig 2F**, **Figure 2—**

figure supplement 4D).

In contrast, the total mass of brown adipose tissue (BAT) remained unchanged between groups

(**Fig 2F**, **Figure 2—figure supplement 4D**). In both male and female cKO mice, we found

hypertrophic eWAT adipocytes (**Fig 2G**, **Figure 2—figure supplement 4E**) and elevated blood

leptin levels (**Fig 2H**, **Figure 2—figure supplement 4F**). However, there was no significant

alteration in blood lipids such as triglycerides (TG), high-density lipoprotein (HDL), low-density

lipoproteins (LDL), and very low-density lipoprotein (VLDL) in either male or female cKO mice

(**Figure 2—figure supplement 3E-G**, **Figure 2—figure supplement 4G-I**). Altogether, these

data suggest that obesity caused by ablating *Rai1* from Bdnf-producing cells is associated with

increased fat mass, fat deposition, and leptin levels.

Obesity could result from increased food intake, reduced energy expenditure, or both. Similar to

SMS mice, male and female cKO mice did not have defects in respiratory exchange rates when

measured by indirect calorimetry (**Figure 2—figure supplement 3H**, **Figure 2—figure**

supplement 4J). Interestingly, cKO mice showed sex-specific differences in energy expenditure.

During the dark phase, female cKO mice showed a reduction in energy expenditure (**Fig 2I**) and

decrease in total locomotor activity (**Fig 2J**). In contrast, total food intake remained unchanged

(**Fig 2K**). Male cKO mice became hyperphagic (**Figure 2—figure supplement 4K**), showed

increased energy expenditure during the dark phase (**Figure 2—figure supplement 4L**), and

exhibited normal locomotor activity when compared to controls (**Figure 2—figure supplement**

4M). These results suggest that obesity in male cKO mice is attributable to overeating despite an

increase in energy expenditure.

To determine if Rai1 loss from Bdnf-producing cells impacts metabolic function, a glucose

tolerance test (GTT) was performed. Interestingly, cKO mice of both sexes showed normal blood

glucose levels right before the test but became hyperglycemic 20 minutes (female cKO) and 45

minutes (male cKO) after glucose administration (**Fig 2L**, **Figure 2—figure supplement 4N**).

This change was accompanied by a significantly higher area under the curve (AUC) for both

groups (**Figure 2—figure supplement 3I**, **Figure 2—figure supplement 4O**). While the male

cKO mice remained hyperglycemic throughout the GTT (**Figure 2—figure supplement 4P**), the

glucose levels in female cKO mice returned to normal towards the end (**Fig 2L**). We also found

increased blood insulin levels during the GTT in both sexes (**Figure 2—figure supplement 3J**,

Figure 2—figure supplement 4P). In the insulin tolerance test (ITT), female cKO mice showed

moderately elevated glucose levels 20 minutes into the test (**Figure 2—figure supplement 3K**).

By contrast, male cKO mice showed considerably higher glucose levels than controls in the ITT

(~6.5 ng/ml for cKO and ~2.0ng/ml for Ctrl) during the 60- to 120-minute intervals (**Figure 2—**

figure supplement 4Q). Altogether, these experiments demonstrate that Bdnf-producing cells

depend on Rai1 to regulate body weight, adiposity, glucose and insulin tolerance in mice.

Rai1 **deletion reduces PVH**^{Bdnf} **neuronal excitability**

Bdnf-producing neurons are widely distributed in the hypothalamus, being found in the

dorsomedial hypothalamus (Unger *et al*, 2007), ventromedial hypothalamus (Xu *et al*, 2003), and

PVH (An *et al.*, 2015). To understand how *Rai1* loss affects intrinsic neuronal excitability and

synaptic transmission of PVH^{Bdnf} neurons that underlie energy homeostasis (An *et al.*, 2015), we

fluorescently labelled them with Cre-dependent tdTomato (**Fig. 3A**). We found that a

hyperpolarized step followed by increasing the holding voltage at resting membrane potentials

(Rm) was able to initiate rebound firing or rebound repetitive firing in most PVH^{Bdnf} neurons

(**Figure 3—figure supplement 1A**). Hyperpolarization-induced I_H currents and low

depolarization events were observed in some PVH^{Bdnf} neurons (**Figure 3—figure supplement**

1A). cKO PVH^{Bdnf} neurons showed reduced action potential (AP) firing frequency at a holding

voltage of -60mV compared to Ctrl PVH^{Bdnf} neurons (**Fig 3B-C**). The resting membrane

potential (Rm) of cKO (-52mV) neurons showed a 10mV positive shift compared to Ctrl neurons

(-62mV) (**Fig 3D**). We also found that fewer cKO PVH^{Bdnf} neurons showed spontaneous firing at

Rm (**Fig 3E**, Ctrl:10/13; cKO:5/12) and at holding voltage of -70mV (**Fig 3F**, Ctrl:6/12;

cKO:1/12) compared to Ctrl PVH^{Bdnf} neurons. The threshold of the first AP initiated by a ramp

current injection (1 nA/sec) from Rm was significantly increased in cKO neurons (**Fig 3G-H**).

Because the higher the holding voltage is, the narrower the shoulder and the lower the amplitude

of APs are (**Figure 3—figure supplement 1B**), we compared the AP waveforms at a holding

voltage of -60mV, close to the Rm of both genotypes. The AP of the cKO group exhibited a

lower peak amplitude (measuring from the threshold to peak) (**Fig 3I**), a normal threshold (**Fig**

3J), and half-width (**Figure 3—figure supplement 1C**). We found that both passive and AP

properties of cKO PVH^{Bdnf} neurons were altered. The Rm of cKO neurons shifted to a more

positive voltage than the gating threshold of voltage-gated sodium channels (6/12 were more

positive than -55mV), which was unfavourable for their transition from the inactivation state to

the closed state. Combined with the higher AP threshold of cKO neurons, these changes

contribute to a decreased excitability in cKO PVH^{Bdnf} neurons.

PVN neurons receive glutamatergic and GABAergic inputs and are tonically inhibited by

GABAergic neurons (Colmers & Bains, 2018; Gao *et al*, 2017). The rightward shift of the

probability of miniature inhibitory postsynaptic current (mIPSC) frequency indicated that cKO

PVH^{Bdnf} neurons received a higher frequency of inhibitory inputs, although the average

frequency (**Fig 3K**) was not significantly different between groups. The probability of mIPSC

amplitude also showed a right shift, suggesting an increased mIPSC postsynaptic response (**Fig**

3L, Figure 3—figure supplement 1D). Furthermore, we observe an increased area under the

curve (**Fig 3M**), indicating that *Rai1* loss induced a larger total outward current upon the

inhibitory inputs. Interestingly, we found that cKO PVH^{Bdnf} neurons had a smaller cellular

diameter (**Figure 3—figure supplement 1E**), indicating that the increased inhibitory currents

were unlikely due to changes in passive membrane properties associated with increased cell size.

Overall, our data argue that *Rai1* deletion decreases the excitability of PVH^{Bdnf} neurons and

likely increases pre-synaptic inhibitory transmission and post-synaptic inhibitory responses,

resulting in decreased AP firing.

Rai1 expression is necessary for PVH^Bdnf neurons to regulate energy homeostasis

Reduced PVH^Bdnf neuronal activity in cKO mice suggested that Rai1 loss in Bdnf-producing PVH neurons might be sufficient to induce defective energy homeostasis *in vivo*. To test this hypothesis by deleting Rai1 only in PVH^Bdnf neurons but not other Bdnf-producing cells, we performed PVH^Bdnf-specific *Rai1* deletion using a CRISPR/Cas9 system packaged into AAV9. Two viruses were delivered; one encoding a reporter molecule (GFP) and three single guide RNAs targeting the *Rai1* gene (sgRai1), and the other carrying Cre-dependent *Staphylococcus aureus* CRISPR-associated protein 9 (saCas9) (**Fig 4A**). Hereafter, our experiments focus on female SMS mice since they showed more consistent defects in fat distribution (Burns *et al.*, 2010) and body weight regulation (Huang *et al.*, 2016) than males. Bilateral administration of two AAVs into the PVH of female *Bdnf^Cre/+* mice (Exp group) resulted in a significant loss of Rai1 protein expression in PVH^Bdnf neurons when compared to *Bdnf^Cre/+* mice injected with an AAV encoding the sgRai1 and GFP (Ctrl group) (**Fig 4B-C**, **Figure 4—figure supplement 1A**). Quantification of AAV-infected (GFP⁺) Bdnf neurons showed that while the total number of Bdnf⁺ (**Figure 4—figure supplement 1B**) and Bdnf⁺ GFP⁺ (**Figure 4—figure supplement 1C**) cells was not different in the two groups, the percentage of AAV-infected Bdnf⁺ cells that express Rai1 was significantly reduced in the Exp group (**Fig 4C**, **Figure 4—figure supplement 1D-E**). Deleting *Rai1* from the PVH^Bdnf neurons resulted in a significant increase in body weight beginning at 17 weeks of age (**Fig 4D**). Similar to the cKO (*Bdnf^Cre/+; Rai1^fl/fl*) mice, obesity in 26-week-old Exp mice was due to increased fat mass (**Fig 4E**) and more specifically, increased subcutaneous inguinal (S.C. ing) and epididymal white adipose tissues (eWAT) but not BAT (**Fig 4F**). We also found that eWAT adipocytes became hypertrophic (**Fig 4G**). Similar to the cKO mice, food intake did not differ between Exp and Ctrl groups (**Fig 4H**), likely due to the

high level of satiety signals in the brain caused by increased blood leptin levels (**Fig 4I**). We also

found that energy expenditure (**Fig 4J**), respiratory exchange rate (**Figure 4—figure**

supplement 1F), and blood lipid parameters, including TG (**Figure 4—figure supplement 1G**),

HDL (**Figure 4—figure supplement 1H**), LDL and VLDL levels (**Figure 4—figure**

supplement 1I), were similar in Exp and Ctrl groups, consistent with the cKO data (**Figure 4—**

figure supplement 14). Interestingly, Exp mice showed significantly less locomotor activity

during the dark phase than the Ctrl mice (**Fig 4K**). A GTT indicated that Exp mice have a

hyperglycemic profile (**Fig 4L**), accompanied by a slightly higher area under the curve for blood

glucose levels (p=0.05) (**Fig 4M**). Insulin levels in Exp mice were also significantly elevated 15

minutes after glucose administration (**Fig 4N**).

While Exp mice showed a trend towards increased glucose levels in insulin tolerance tests, the

change was not statistically different (**Figure 4—figure supplement 1J**). Altogether, these

experiments indicate that conditional deletion of Rai1 from PVH^{Bdnf} neurons during adolescence

impairs the regulation of body weight, adiposity, glucose tolerance, and locomotion. These

changes recapitulate most phenotypes observed in cKO mice with *Rai1* deleted in Bdnf-

producing cells from early development.

Pharmacologically enhancing TrkB signalling partially alleviates obesity in SMS mice

Our findings so far indicate that multiple pathways, including neurotrophin signalling, are

dysregulated in SMS mice, and that Rai1 loss from Bdnf-producing cells, including PVH^{Bdnf}

neurons, impairs energy homeostasis *in vivo*. Next we sought to examine the contribution of

Bdnf-TrkB dysfunction in SMS pathogenesis by pharmacologically activating downstream

targets of the Bdnf-TrkB pathway.

We systematically treated mice with LM22A-4, a small molecule Bdnf loop domain mimetic

(Massa *et al*, 2010). To measure the bioavailability of LM22A-4 in the SMS brain, we

simultaneously administered LM22A-4 intranasally (5 mg/kg body weight) and intraperitoneally

(50 mg/kg body weight), using the highest doses used in previous studies (Canals *et al*, 2004;

Chang *et al*, 2006; Gu *et al*, 2022; Kron *et al*, 2012; Ogier *et al*, 2007). We then harvested

forebrain and hypothalamic tissues one- or three-hours after drug administration. Spectrometric

analysis found a significantly higher LM22A-4 concentration in both brain regions one-hour

post-treatment, while its concentration decreased three hours after treatment (**Figure 5—figure**

supplement 1A). Consistent with the reduced neurotrophin signalling observed in our proteomic

analysis, saline-treated SMS mice showed reduced phosphorylated-Akt (p-Akt; **Fig 4A**, **Figure**

5—figure supplement 1B) levels in the hypothalamus. LM22A-4 treatment significantly

increased the levels of p-Akt in the SMS mice 1-hr post-treatment (**Fig 4A**, **Figure 5—figure**

supplement 1B), indicating increased signalling downstream of the Bdnf-TrkB pathway.

We next examined if chronic LM22A-4 treatment was sufficient to reverse or modify the

progression of the obesity phenotype in adult SMS mice. Specifically, we treated SMS mice

daily from 8 to 14 weeks of age, during which SMS mice began to gain significantly more

weight. Saline-treated SMS mice (**Fig 5C**) were significantly heavier than saline-treated control

mice from 9 weeks of age onward. Interestingly, LM22A-4-treated SMS mice showed a similar

body weight to saline-treated control littermates until 13 weeks of age but became obese by 14

weeks of age (**Fig 5D**). Consistent with a partial rescue, LM22A-4-treated SMS mice were

significantly less obese than saline-treated SMS mice throughout the treatment period (**Fig 5E**).

Control mice treated with LM22A-4 did not show a further reduction in body weight (**Figure**

5—figure supplement 1C), indicating that the therapeutic effects were specific to SMS

pathogenesis.

We measured food intake and found that LM22A-4 did not significantly rescue increased food

intake in SMS mice. (**Fig 5F**). SMS mice showed normal energy expenditure (**Figure 5—figure**

supplement 1D) and respiratory exchange rate (**Figure 5—figure supplement 1E**), neither of

which was altered by LM22A-4 treatment. Interestingly, we did observe a partial rescue of

locomotor activity in LM22A-4-treated SMS mice that was not significantly different from

saline-treated Ctrl or SMS mice (**Fig 5G**). We also found that LM22A-4 treatment partially

rescued fat deposition (**Fig 5H**) and completely rescued HDL levels (**Fig 5I**) in SMS mice.

During an insulin tolerance test, we found that saline-treated SMS mice showed increased

baseline insulin levels (**Fig 5J**) and blood glucose levels during the test (**Fig 5K**), both of which

were partially rescued by LM22A-4 treatment.

In addition to a deficit in energy homeostasis, SMS mice showed two highly reproducible

behavioral deficits: increased stereotypic rearing in the open field and decreased social

dominance in the tube test (Huang *et al*, 2018; Rao *et al*, 2017). Saline-treated SMS mice

showed significantly more rearing behavior when compared to the saline-treated controls (**Fig**

5L). Interestingly, LM22A-4 treatment reduced the repetitive rearing behaviour in SMS mice

(**Fig 5L**). LM22A-4 treatment did not change repetitive behaviors in control mice (**Figure 5—**

figure supplement 1F), suggesting the effect is specific to *Rai1* haploinsufficiency. However,

we did not observe a rescue of social dominance deficits in SMS mice (**Figure 5—figure**

supplement 2A-E). Together, these data indicate that LM22A-4 treatment delayed the onset of

obesity and fully rescued repetitive behaviour in SMS mice, suggesting that Bdnf-TrkB

signalling dysfunction contributes to changes in the mouse SMS brain linked to obesity and

behavioral phenotypes.

Discussion

The hypothalamus, and Bdnf-TrkB signalling in this region, control body weight by regulating

the intricate balance between energy intake and expenditure (Xu & Xie, 2016). The role of Bdnf-

producing neurons in human obesity disorders and how the activity of these cells is regulated

remained poorly studied. We found that Rai1 deficiency, linked to Smith-Magenis syndrome

(SMS), regulates Bdnf expression and maintains intrinsic excitability and inhibitory synaptic

transmissions in PVH^{Bdnf} neurons.

Consistent with this mechanism, we found that removing Rai1 from Bdnf-producing neurons or

PVH^{Bdnf} neurons results in imbalanced energy homeostasis and results in obesity. A small

molecule drug LM22A-4 led to increased Akt phosphorylation and partially rescued locomotor

activity, insulin intolerance, and HDL levels, while delaying the onset of obesity in SMS mice.

These findings highlight a potential role for hypothalamic Bdnf-producing neurons, specifically

PVH^{Bdnf} neurons, in SMS pathogenesis and highlight Bdnf signalling as a potentially druggable

pathway to improve energy homeostasis defects and repetitive behaviours commonly observed in

SMS patients (Burns *et al.*, 2010; Moss *et al*, 2009).

The involvement of different PVH subtypes in human obesity is only beginning to be elucidated.

Dysfunction of oxytocin PVH neurons is linked to the Prader-Willi syndrome (Lee *et al*, 2012),

another genetic disorder associated with obesity. Since Rai1 expression is enriched in PVH^{Bdnf}

neurons but not PVH^{Oxytocin} neurons, our data suggest that distinct pathophysiological

mechanisms underlie SMS and Prader-Willi syndrome.

Conditional knockout analysis allowed us to specifically map the cell-specific origin of

neuropathological phenotypes, such as SMS-like symptoms in mice. Our previous work found

that deleting *Rai1* from cortical excitatory neurons using an *Emx1^{Cre}* allele (Gorski *et al*, 2002),

from the GABAergic neurons using a *Gad2^{Cre}* allele (Taniguchi *et al*, 2011), or from astrocytes

using a *GFAP^{Cre}* allele (Garcia *et al*, 2004) did not induce SMS-like obesity (Huang *et al.*, 2016).

In contrast, obesity could be induced by ablating *Rai1* from Sim1-lineage neurons and, to a lesser

extent, SF1-lineage neurons (Huang *et al.*, 2016). This analysis suggests that Rai1 expression in

PVH is more important than in ventromedial nucleus of the hypothalamus (VMH) for body

weight regulation. Interestingly, embryonic deletion of *Bdnf* from SF1-lineage neurons including

the VMH only resulted in modest weight gain (Unger *et al.*, 2007). By contrast, deleting *Bdnf*

from Sim1-lineage neurons including the PVH resulted in a more severe obesity phenotype (An

et al., 2015). This is consistent with the idea that Bdnf expression in PVH is critical for energy

homeostasis. We found that Rai1 expression in Bdnf-producing cells regulates energy

homeostasis in a sexually dimorphic manner. Specifically, female cKO mice showed normal

food intake, decreased energy expenditure, and reduced locomotor activity. In contrast, male

cKO mice showed increased food intake with no changes in locomotor activity. Intriguingly,

energy expenditure was increased in male cKO mice, likely due to the higher demand to

maintain normal activity levels in obese mice. In both sexes, cKO mice showed an increased fat

mass and adiposity with no change in lean mass, highlighting an important role for Rai1 in

regulating fat deposition. Our data also indicate that Bdnf-producing cells are only partially

responsible for obesity in SMS: male and female cKO mice become overweight by 15 weeks, in

contrast to SMS mice which become obese by 9 weeks of age, suggesting that non-Bdnf neurons

contribute to SMS phenotypes.

We found the metabolic phenotypes induced by deleting *Rai1* from Bdnf-producing cells were

specifically due to *Rai1* deletion in PVH[Bdnf] neurons. Therefore, our study focused on PVH[Bdnf]

neurons. Electrophysiological experiments showed that most wild-type PVH[Bdnf] neurons display

high frequency spontaneous AP firing (1-10 Hz) at the resting membrane potential. The majority

of PVH[Bdnf] neurons fire sharp APs held at a hyperpolarized potential and show repetitive

overshoot during the current ramp test. Loss of Rai1 significantly reduced intrinsic neuronal

excitability and increased inhibitory synaptic transmissions onto PVH[Bdnf] neurons. Because

GABAergic Rai1 loss did not induce obesity (Huang *et al.*, 2016) and most Bdnf-producing

neurons are glutamatergic (Canals *et al*, 2001), the increased mIPSC amplitude was likely due to

postsynaptic changes in GABA receptor functions caused by Rai1 loss. The ionic changes in

PVH[Bdnf] neurons that precisely explain these excitability changes await future investigations.

Different groups of PVH neurons show phasic bursting or slowly inactivating delayed rectifier

potassium conductance, which are properties mediated by different potassium channels (Lee *et*

al., 2012). Given that our previous RNA-seq experiment found that hypothalamic Rai1 loss

induces misexpression of potassium channel subunits *Kcnc2* and *Kcnj11* (Huang *et al.*, 2016),

Rai1-dependent potassium channels are likely involved in altered neuronal activity. Furthermore,

we found fewer low threshold depolarization events in cKO PVH[Bdnf] neurons during the

hyperpolarized voltage step, suggesting a potential involvement of HCN channels.

Our previous work found that Rai1 loss in the dentate gyrus resulted in neuronal

hyperexcitability (Chang *et al*, 2022b). Here we demonstrated that Rai1 loss in the hypothalamus

resulted in neuronal hypoactivity, suggesting that Rai1 regulates neuronal excitability in a cell

type-specific manner. Reduced PVH^{Bdnf} neuronal activity upon Rai1 loss could potentially

explain obesity, consistent with a previous report that found chemogenetic activation of PVH^{Bdnf}

neurons promotes negative energy balance by decreasing feeding and increasing energy

expenditure (Wu & Xu, 2022).

In future work, it would be interesting to dissect the sex-specific role of Rai1 in regulating the

functional development of PVH^{Bdnf} neurons. In addition to a role for Rai1 in regulating neuronal

activity, it is possible that Rai1 regulates other aspects of PVH^{Bdnf} neuronal function, including

projections to other brain regions to regulate energy homeostasis. In this regard, it would be

interesting to dissect how Rai1 controls circuit connectivity of PVH^{Bdnf} neurons and how Rai1

loss in PVH^{Bdnf} neurons affects nearby PVH^{TrkB} neurons that receive Bdnf signaling and are

linked to body weight regulation (An *et al*, 2020).

At the molecular level, our RPPA data show that *Rai1* depletion disrupted multiple intracellular

signalling pathways and resulted in the hypoactivation of specific phospho-signalling proteins.

Multiple early studies support the notion that Rai1 directly promotes Bdnf expression. First,

hypothalamic *Bdnf* mRNA levels were reduced in SMS mice (Burns *et al.*, 2010; Javed *et al.*,

2021) or mice with *Rai1* conditional deletions (Huang *et al.*, 2016). Second, Rai1 protein directly

binds to a promoter region of *Bdnf* (Huang *et al.*, 2016). Finally, when Rai1 expression was

induced using CRISPR activation, *Bdnf* mRNA levels were significantly elevated (Chang *et al*,

2022a).

To lend further support, here we show that SMS mice have reduced hypothalamic Bdnf protein

levels, as well as reduced activation of a TrkB downstream target Akt. Reduced Akt activity is

frequently associated with insulin resistance and obesity in mice (Shao *et al*, 2000) and children

(Su *et al.*, 2021). We also found that obesity, increased HDL levels, and insulin insensitivity in

SMS mice could be partially ameliorated using a small molecule drug (LM22A-4) that promotes

Akt phosphorylation. LM22A-4 penetrates the blood-brain barrier reasonably well and has been

shown to ameliorate symptoms of other neurological disorders associated with Bdnf dysfunction

including Rett syndrome, Huntington's disease, Dravet syndrome, and refractory neonatal

seizures (Canals *et al.*, 2004; Chang *et al.*, 2006; Gu *et al.*, 2022; Kron *et al.*, 2012; Ogier *et al.*,

2007). The partial rescue of SMS phenotypes that we observed contrasts with our previous

findings that found (1) AAV-mediated Bdnf overexpression in PVH at 3 weeks of age and (2)

genetically overexpressing Bdnf from early development could fully prevent the development of

obesity in SMS mice (Javed *et al.*, 2021). It is therefore possible that further increasing TrkB

signalling activation and/or earlier intervention might improve therapeutic outcomes.

Alternatively, our RPPA data showed that in addition to disruption of neurotrophin signalling,

multiple pathways including mTOR were downregulated in SMS. Increasing hypothalamic

mTOR activity has been shown to decrease food intake and body weight (Cota *et al.*, 2006).

Therefore, it is possible that enhancing both mTOR and TrkB signalling could achieve stronger

rescue of obesity in SMS mice. Nevertheless, these data support the concept that drugging TrkB

signalling may improve stereotypical repetitive behavior and partially alleviate metabolic

functions in a mouse model of SMS. Delayed symptom onset could provide a window of

opportunity for other interventions to further modify disease progression, an area requiring

further investigation.

This study has several limitations. First, whether the PVH^{Bdnf} neurons depend on Rai1 to regulate

adaptive thermogenesis awaits future analysis. It is known that a subset of posterior PVH^{Bdnf}

neurons project to the spinal cord and regulate thermogenesis (An *et al.*, 2015). However, we

found the weights of brown adipocytes were similar in control and cKO mice. Moreover, SMS

mice have normal respiratory exchange rate and there are currently no reports of defective

adaptive thermogenesis in SMS patients. This suggests that adaptive thermogenesis is not

impacted in SMS. Another limitation is that we did not test different doses, durations, and time

points for LM22A-4 treatment. The precise mechanisms by which LM22A-4 improves body

weight, metabolic function, and repetitive behaviours also remain unknown. Nevertheless, our

work identifies Rai1 as a novel regulator of the neuronal excitability of PVH^{Bdnf} neurons. This

regulation is critical for regulating food intake and energy expenditure. Our evidence suggests

that Bdnf-producing neurons are involved in SMS pathogenesis and that enhancing Bdnf-TrkB

signalling should be explored as a future therapeutic avenue for obesity and metabolic conditions

associated with reduced neurotrophin signalling (Xu & Xie, 2016).

Materials and Methods

Animal Management

The animal protocol was approved by the animal care committee of the Montreal General

Hospital and was performed according to the guidelines of the Canadian Council on Animal

Care. Mice were group-housed in 12-hour light/12-hour dark cycles with an *ad libitum* supply of

regular chow and water. The exception was in the fasting studies when the animals were food

deprived for 5 hours and in CALMS and food intake studies when animals were single-housed in

a 12-hour light/12-hour dark cycle with an *ad libitum* supply of regular chow and water. All mice

were maintained in a C57Bl/6J background. $Bdnf^{Cre/Cre}$, *Rosa-tomato*$^{td/td}$ (Ai9), and $Act\beta^{Cre/Cre}$

mice were purchased from the Jackson laboratory. The $Rai1^{fl/fl}$ strain was previously generated

and characterized by us (Huang *et al.*, 2016). SMS mice ($Rai1^{+/-}$: whole-body germline *Rai1*

heterozygous knockout that genetically mimics SMS) were generated by crossing germline

$Act\beta^{Cre/Cre}$ mice with $Rai1^{fl/+}$. To generate conditional knockout mice, $Bdnf^{Cre/+}$; $Rai1^{fl/+}$ mice

were crossed with $Rai1^{fl/+}$ to create control and experimental groups from the same littermate.

Metabolic profiling

Metabolic profiling was conducted in three different experimental conditions: male and female

conditional knockout mice, female virus-injected $Bdnf^{Cre/+}$ mice, and female drug- or saline-

injected SMS and control mice. Metabolic profiling included body weight and body composition

analysis, food intake measurement, indirect calorimetry for locomotion activity, energy

expenditure (EE), respiratory exchange rate (RER), measurement of serum lipids and leptin

levels, adipose tissue histology analysis, and glucose and insulin tolerance tests. All experiments

were performed as previously conducted experiments (Javed *et al.*, 2021). A brief description is

detailed below.

Body weight and body composition analysis

The body weight of animals was measured weekly from week 3 (weaning age) onward. Body

composition, including total fat and lean mass, were assessed using a nuclear EchoMRI whole-

body analyzer. Toward the end of experiments, animals were euthanized to collect brown

adipose, visceral, subcutaneous (inguinal) and ependymal white adipose tissues and weighed

using an analytical scale (Sartorius).

Food Intake Measurement

Mice were single-housed for food intake measurement. Food intake was measured over a one-

week period during which food was weighed prior to placement in the cage with minimal

bedding. Each morning remaining food inside the cage was weighed. The amount of total food

consumed was averaged over 7 days.

Indirect Calorimetry

A Comprehensive Lab Animal Monitoring System (CLAM) (Columbus instruments) was used to

measure the RER and EE (normalized over lean mass). Animals were singled-housed in the

CLAMS apparatus at 21°C (70°F) in a light–dark cycle matching their housing conditions for 24

hours (acclimation), followed by 48 hours of measurement.

Serum lipid level determination

A fluorescence-based assay kit (Abcam ab65390) was used to measure serum lipid levels. The

manufacturer's instructions were followed except that we used half-area black 96-well

microplates. Fluorescence measurements at 535/587 nm (Ex/Em) were recorded with the Ensight

instrument from PerkinElmer. A new standard curve ranging from 0 to 10 μg/ml was produced

for each microplate to accurately determine lipid levels in serum samples. Standards and samples

were loaded and analyzed twice.

Histology of adipose tissue

Ependymal white adipose tissues (WAT) were excised immediately after the animals were

euthanized and fixed with 10% formalin in PBS. The fixed tissues were dehydrated, embedded in

paraffin, and sectioned. Two 5-micrometre thick sections were taken from each animal and

stained with hematoxylin and eosin. Images of sections were then captured using an Aperio

ScanScope CS slide scanner at 20X optical magnification, and the resulting images were saved

in jpeg format using the ImageScope software. The images were manually cleaned to remove

non-adipose tissues such as lymph nodes. An automated method using the Adiposoft plugin

version 1.16 was utilized to determine adipocyte size and number from the jpeg images in Fiji

version 2.1.0/1.53c. The Adiposoft settings used excluded cells on edges and set a minimum and

maximum cell diameter of 20 and 300 μm, respectively. An experimental value of 2.49 μm per

pixel was determined using the Aperio scale bar and used for Adiposoft. To ensure accuracy, two

images from each animal were analyzed blindly, examining more than 1600 cells per mouse and

more than 9100 cells per condition. Adipocyte size frequencies were calculated using the

frequency function in Excel.

Glucose and insulin tolerance tests

For the glucose tolerance test, experimental mice were food deprived for 5 hours with *ad libitum*

water supply. A bolus of glucose (1.5g/kg of lean body weight) was then administered

intraperitoneally and glycemia was measured from tail vein blood samples using the Accu-chek

Performa glucometer at T0 (before the injection) as well as at 15, 30, 45, 60, 90 and 120 min

(after the injection). Tail vein blood samples were collected via a capillary for insulin assays at

T0, 15 and 30 min. For insulin tolerance tests, experimental mice were food-deprived for 5 hours

with *ad libitum* access to water. A bolus of insulin (1 U/kg of lean body weight) was

administered via an IP injection and glycemia was measured from blood sampled at the tail vein

using a Accu-chek Performa glucometer at T0 (before the injection), as well as at 15, 30, 45, 60,

90 and 120 min (after the injection). Tail vein blood samples were collected via a capillary for

insulin assays at T0.

Stereotaxic Injections

$Bdnf^{Cre/+}$ mice were anesthetized using isoflurane, and the ears were barred onto a stereotaxic

device (David Kopf Instruments). Adeno-associated viruses (AAVs, volume = 250 nl) were

bilaterally injected into the PVH using the following coordinates relative to the Bregma: -0.6 mm

anteroposterior (AP), ±0.35 mm mediolateral (ML) and -5.52 mm dorsoventral (DV). A

description of the AAV constructs and titre used is detailed below:

AAV9-CMV7-DIO-saCas9 (Viral title: >1 × 10^13), Vector Biolab

AAV9-CMV7-sgRai1-GFP (Viral title: >1 ×10^9), Applied Biological Materials Inc. Three

sgRNAs were pooled to target Rai1 (sgRNA1: TTCCTCGCCAGAGTAGCGCC, sgRNA2:

CCCAGCCTCATGATAGGCCG, sgRNA3: CCCAGCCTCATGATAGGCCG).

Immunostaining of mouse brain tissues

Immunostaining experiments were performed as previously described (Huang *et al*, 2012).

Briefly, mice were anesthetized and perfused with 1X PBS and 4% formaldehyde, and the brains

were dissected and fixed in 4% formaldehyde overnight. Fixed brains were then transferred to

30% sucrose solution and subsequently embedded in the OCT solution. The brains were then

sectioned (40μm), washed in PBS, permeabilized with 0.5% Triton X-100, and incubated with

10% normal donkey serum in PBS for 2 hours. The sections were then incubated overnight with

primary antibodies (Rai1: 1:500 dilution, RRID:AB_2921229 (Huang *et al.*, 2016); Bdnf: 1:500,

RRID:AB_10862052, Abcam ab108319; p-TrkB: 1:500 dilution, RRID:AB_2721199, Millipore

ABN1381; oxytocin: 1:1000 dilution, RRID:AB_11212999, Millipore MAB5296) at 4°C on a

shaker. The following day, sections were washed with 1X PBS twice for 10 minutes and

incubated with a secondary antibody (1:1000 dilution) for 3 hours at room temperature. Sections

were washed with 1X PBS once before mounting on glass slides with the DAPI-mounting

(DAPI-Fluoromount-G, SouthernBiotech) and then imaged using a confocal microscope

(Olympus FV-1000 confocal laser scanning microscope) with 20x or 40x objectives (NA 0.85

and 1.3, respectively). Z-stacks were taken with a step size of 1.0 μm with a 1024 x 1024

resolution. Images were acquired as a stack with at least 13 optical sections. Image analysis was

performed using the Fiji software.

Real-Time Quantitative Reverse Transcription PCR (qRT-PCR)

qRT-PCR was performed using mRNAs isolated from hypothalamic tissue samples. After

isolation of total RNA, mRNAs were reverse-transcribed with the SuperScript III First-Strand

Synthesis System (Thermo-Fisher). Quantitative PCR reactions were conducted using SsoFast

EvaGreen Supermix on a Bio-Rad qPCR system.

Enzyme-linked immunosorbent assay (ELISA)

ELISA was performed using hypothalamic tissues from 7-week-old SMS mice to assess the

mature Bdnf protein levels using a commercially available Bdnf ELISA kit (Mature BDNF

Rapid ELISA Kit: Human, Mouse, Rat).

Western blotting analysis

Mice were quickly decapitated, brains were rapidly removed, and hypothalamic tissues were

collected, weighed, and snap-frozen with liquid nitrogen. Hypothalamic tissues were lysed using

a tissue lysis buffer containing RIPA buffer (5M NaCl, 1M PH 8.0 Tris HCl, 1% NP.40, 10%

$C_{24}H_{39}NaO_4$, and 20% SDS) and the Halt™ Protease and Phosphatase Inhibitor Cocktail

(Thermo-Fisher). Lysates were first sonicated on ice for 15 seconds followed by centrifuging at

10000 RPM at 4°C for 10 minutes. The supernatant was collected in a separate Eppendorf tube,

and a BCA assay was performed to quantify protein levels. Proteins were then separated using

SDS-PAGE gel and transferred onto nitrocellulose membranes. To determine the

phosphorylation levels of Akt, membranes were cut and blocked in 5% milk in tris-buffered

saline (TBS) for 1 hour and then incubated overnight at 4°C with anti-p-Akt (Ser473) antibody

(1:1000 dilution, RRID:AB_2315049, Cell Signaling Technology 4060T) and anti-H3 antibody

(internal control: 1:1000 dilution, RRID:AB_302613, Abcam ab1791). The next day, membranes

were washed with TBST three times for five minutes before incubating with a HRP-conjugated

secondary antibody (1:10000 dilution, Thermo Fisher 200-403-095S). Membranes were later

stripped by incubating in 0.5M NaOH for 10 min and then re-probed for total Akt levels by

incubating with anti-Akt (C67E7) antibody (1:1000 dilution, Cell Signaling Technology 4691T).

The density of each immunoreactive phospho-protein band was expressed as a fraction of the

band for the total Akt protein in the same lane.

Reverse-phase protein array (RPPA)

Reverse-phase protein array (RPPA) was used for precise high-throughput quantification of total

and phospho-protein levels in different signalling pathways. Protein lysates from hypothalamic

tissues were extracted and transferred to 384-well microarray plates to print individual slides.

The slides were then labelled with validated antibodies that recognize total protein and specific

phosphorylation sites. SYPRO Ruby staining was used to measure total protein levels. The data

obtained from RPPA were normalized for individual samples. Each antibody's raw signal

intensity (SI) was determined by subtracting the background from the antibody-specific signals

during image analysis. The normalized antibody SI was expressed using the formula $N = A - C/T$

x M, where N is the normalized SI, A is the raw antibody SI, C is the negative control SI, T is the

SI of total protein, and M is the median SI of total protein from a spot within the same sample

group. Antibodies with SI less than 200 were filtered out, and quality control (QC) scores were

established to discard poor-quality reactions. The coefficient of variation (CV) score was

determined for each antibody and defined as the percentage of samples on a slide with a CV <

20%. Antibodies with a median technical triplicate CV >20% were also excluded from further

analysis. T-tests were performed for each antibody to identify differentially expressed proteins in

the SMS compared to control groups. PCA plots were generated for all samples, annotated with

their experimental group, and hierarchical clustering was performed using a Python analysis tool.

Enrichment analysis on the differentially expressed proteins was performed using an online data

resource string: version 11.5.

Electrophysiology of acute brain slices

Mouse brains were removed and placed in ice-cold carbonated slicing artificial cerebrospinal

fluid (aCSF), which contained (in mM) 124 NaCl, 3 KCl, 1.4 NaH_2PO_4, 26 $NaHCO_3$, 1.3 Mg

SO4, 10 D-glucose, 0.5 $CaCl_2$, and 3.5 $MgCl_2$. Coronal sections (300 μm) were cut on a

vibratome. Slices were allowed to recover at 32 °C for 40 minutes and then at 30 °C for

30 minutes to 6 hours in the recording medium, carbonated aCSF (124 NaCl, 3 KCl, 1.4

NaH_2PO_4, 26 $NaHCO_3$, 1.3 Mg SO4, 10 D-glucose, 2.4 $CaCl_2$, in mM, ~300 mOsm). Signals

were recorded with a 5× gain, low-pass filtered at 2 kHz, digitized at 10 kHz (Molecular

Devices Multiclamp 700B) and analyzed with pClamp 11 (Molecular Devices). Whole-cell

recordings were made using 3–5 MΩ pipettes filled with an internal solution that contained

(in mM) 131 potassium gluconate, 8 NaCl, 20 KCl, 2 EGTA, 10 HEPES, 2 MgATP and 0.3

Na3GTP (pH 7.2, with KOH, 300–310 mOsm with sucrose). For miniature current recording,

the internal solution contained (in mM) 130 gluconate acid, 8 CsCl, 1 NaCl, 2 EGTA, 0.8

CsOH, 10 HEPES, 2 MgATP and 0.3 Na3GTP (pH 7.2, with CsOH, 300–310 mOsm adjusted

with sucrose). mEPSCs were recorded in the presence of 1 µM TTX at a holding potential of -

70 mV. mIPSCs were recorded at a holding potential of +10 mV. The mIPSCs were analyzed

from events during a 2-minute continuous recording that contained > 15 events/neuron.

Additional calculations, statistical analysis, and plotting was performed in Microsoft Excel or

Prism (GraphPad).

Spectrometric analysis of LM22A-4 levels in the brain

We assessed the efficacy of LM22A-4, a small molecule pharmacological agent, in 8-week-old

SMS and control mice. LM22A-4, chemical name N1, N3, N5tris (2hydroxyethyl) 1, 3, 5

benzenetricarboxamide (Molecular formula: $C_{15}H_{21}N_3O_6$, Formula weight: 339.3), was

manufactured by Cayman Chemical with purity specification >98%. A stock solution was made

by first dissolving LM22A-4 in DMSO (solubility =30mg/ml) and then diluting it with 0.9%

isotonic sterilized saline solution (Sigma). To maximize brain concentrations, LM22A-4 was

administered both intranasally (5mg/kg) and intraperitoneally (50mg/kg), as previously reported

(Simmons *et al*, 2013). The bioavailability of this drug in the hypothalamus using these doses

and routes of administration was assessed with the reverse phase LC triple-quadrupole tandem

mass spectroscopic detection (LC-MS/MS) by Absorption Systems. Mice were euthanized, and

hypothalamic tissue was collected at either 1 h or 3 h post-injection (n= 4/timepoint). Known

amounts of LM22A-4 were added to the blank sample to generate a standard curve to verify test

accuracy.

LM22A-4 treatment

Mice were divided into four groups: Saline-injected control, Saline-injected SMS, LM22A-4-injected SMS, and LM22A-4-injected control mice. Mice were injected daily, intra-nasally (5mg/kg) and intraperitoneally (50mg/kg), for 7 weeks. Body weight was measured between 8 and 14 weeks of age. Food intake was measured from 14 to 15 weeks of age. After 15 weeks of age, mice were euthanized to collect brown adipose, visceral, subcutaneous (inguinal) and ependymal white adipose tissues and weighed using an analytical scale (Sartorius). Hypothalamic brain tissues were collected for western blot analysis. Metabolic profiling was performed on a separate cohort of mice injected for two weeks prior to the start of experiments as well as at specific weeks during the experiment.

Vertical rearing

To measure vertical repetitive rearing, mice were taken into the testing room and allowed to habituate for 1 h prior to testing. During the test, mice were placed in the center of a 45 cm (W) × 45 cm (D) × 40 cm (H) square arena, and their movement was recorded for 10 minutes.

Statistical analysis

Sample sizes used were based on our previously studies (Huang *et al.*, 2016; Javed *et al.*, 2021). The data were subjected to statistical analysis for significance by using GraphPad Prism 0.9. Significance was defined as having a p-value < 0.05. The significance levels were expressed as follows: * p< 0.05, **p<0.01, ***p<0.0001, **** p<0.0001. The statistical tests used for each analysis (e.g., Student's t-test, Fisher's exact test, analysis of variance [ANOVA]) are indicated in the text and legends.

Data Availability

This study includes no data deposited in external repositories.

Acknowledgements

Sehrish Javed was supported by a doctoral award from the Fonds de Recherche du Québec -

Santé (FRQS). This work was supported by the Rare Diseases: Models & Mechanisms Network

and Smith-Magenis syndrome Research Foundation.

Author contributions

Sehrish Javed: Conceptualization; formal analysis; investigation; data curation; visualization;
writing. **Ya-Ting Chang**: Formal analysis; investigation; data curation; visualization. **Yoobin**
Cho: Formal analysis; investigation. **Yu-Ju Lee**: Formal analysis; investigation. **Hao-Cheng**
Chang: Formal analysis; investigation. **Minza Haque**: Formal analysis; investigation. **Yu**
Cheng Lin: Formal analysis; investigation. **Wei-Hsiang Huang**: Conceptualization; supervision;
funding acquisition; visualization; writing; project administration.

Disclosure and competing interests statement

The authors do not have any conflicts of interest to disclose relating to this work.

References

Abad C, Cook MM, Cao L, Jones JR, Rao NR, Dukes-Rimsky L, Pauly R, Skinner C, Wang Y, Luo F *et al* (2018) A Rare De Novo RAI1 Gene Mutation Affecting BDNF-Enhancer-Driven Transcription Activity Associated with Autism and Atypical Smith-Magenis Syndrome Presentation. *Biology (Basel)* 7

Alaimo JT, Barton LV, Mullegama SV, Wills RD, Foster RH, Elsea SH (2015) Individuals with Smith-Magenis syndrome display profound neurodevelopmental behavioral deficiencies and exhibit food-related behaviors equivalent to Prader-Willi syndrome. *Res Dev Disabil* 47: 27-38

An JJ, Kinney CE, Tan JW, Liao GY, Kremer EJ, Xu B (2020) TrkB-expressing paraventricular hypothalamic neurons suppress appetite through multiple neurocircuits. *Nat Commun* 11: 1729

An JJ, Liao GY, Kinney CE, Sahibzada N, Xu B (2015) Discrete BDNF Neurons in the Paraventricular Hypothalamus Control Feeding and Energy Expenditure. *Cell Metab* 22: 175-188

Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, Ferreira M, Tang V, McGovern RA, Kenny CD *et al* (2005) Divergence of melanocortin pathways in the control of food intake and energy expenditure. *Cell* 123: 493-505

Boot E, Linders CC, Tromp SH, van den Boogaard MJ, van Eeghen AM (2021) Possible underreporting of pathogenic variants in RAI1 causing Smith-Magenis syndrome. *Am J Med Genet A* 185: 3167-3169

Burns B, Schmidt K, Williams SR, Kim S, Girirajan S, Elsea SH (2010) Rai1 haploinsufficiency causes reduced Bdnf expression resulting in hyperphagia, obesity and altered fat distribution in mice and humans with no evidence of metabolic syndrome. *Hum Mol Genet* 19: 4026-4042

Canals JM, Checa N, Marco S, Akerud P, Michels A, Perez-Navarro E, Tolosa E, Arenas E, Alberch J (2001) Expression of brain-derived neurotrophic factor in cortical neurons is regulated by striatal target area. *J Neurosci* 21: 117-124

Canals JM, Pineda JR, Torres-Peraza JF, Bosch M, Martin-Ibanez R, Munoz MT, Mengod G, Ernfors P, Alberch J (2004) Brain-derived neurotrophic factor regulates the onset and severity of motor dysfunction associated with enkephalinergic neuronal degeneration in Huntington's disease. *J Neurosci* 24: 7727-7739

Chang HC, Lee YJ, Javed S, Haque M, Chang YT, Lin YC, Oram C, Huang WH (2022a) rAAV-CRISPRa therapy corrects Rai1 haploinsufficiency and rescues selective disease features in Smith-Magenis syndrome mice. *J Biol Chem*: 102728

Chang Q, Khare G, Dani V, Nelson S, Jaenisch R (2006) The disease progression of Mecp2 mutant mice is affected by the level of BDNF expression. *Neuron* 49: 341-348

Chang YT, Kowalczyk M, Fogerson PM, Lee YJ, Haque M, Adams EL, Wang DC, DeNardo LA, Tessier-Lavigne M, Huguenard JR *et al* (2022b) Loss of Rai1 enhances hippocampal excitability and epileptogenesis in mouse models of Smith-Magenis syndrome. *Proc Natl Acad Sci U S A* 119: e2210122119

Collaboration NCDRF (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 390: 2627-2642

Colmers PLW, Bains JS (2018) Balancing tonic and phasic inhibition in hypothalamic corticotropin-releasing hormone neurons. *J Physiol* 596: 1919-1929

Cota D, Proulx K, Smith KA, Kozma SC, Thomas G, Woods SC, Seeley RJ (2006) Hypothalamic mTOR signaling regulates food intake. *Science* 312: 927-930

Edelman EA, Girirajan S, Finucane B, Patel PI, Lupski JR, Smith AC, Elsea SH (2007) Gender,
genotype, and phenotype differences in Smith-Magenis syndrome: a meta-analysis of 105 cases.
Clin Genet 71: 540-550
Gandhi AA, Wilson TA, Sisley S, Elsea SH, Foster RH (2022) Relationships between food-
related behaviors, obesity, and medication use in individuals with Smith-Magenis syndrome. *Res*
Dev Disabil 127: 104257
Gao H, Molinas AJR, Miyata K, Qiao X, Zsombok A (2017) Overactivity of Liver-Related
Neurons in the Paraventricular Nucleus of the Hypothalamus: Electrophysiological Findings in
db/db Mice. *J Neurosci* 37: 11140-11150
Garcia AD, Doan NB, Imura T, Bush TG, Sofroniew MV (2004) GFAP-expressing progenitors
are the principal source of constitutive neurogenesis in adult mouse forebrain. *Nat Neurosci* 7:
1233-1241
Gorski JA, Talley T, Qiu M, Puelles L, Rubenstein JL, Jones KR (2002) Cortical excitatory
neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage. *J*
Neurosci 22: 6309-6314
Gray J, Yeo GS, Cox JJ, Morton J, Adlam AL, Keogh JM, Yanovski JA, El Gharbawy A, Han JC,
Tung YC *et al* (2006) Hyperphagia, severe obesity, impaired cognitive function, and
hyperactivity associated with functional loss of one copy of the brain-derived neurotrophic factor
(BDNF) gene. *Diabetes* 55: 3366-3371
Gu F, Parada I, Yang T, Longo FM, Prince DA (2022) Chronic partial TrkB activation reduces
seizures and mortality in a mouse model of Dravet syndrome. *Proc Natl Acad Sci U S A* 119
Huang WH, Guenthner CJ, Xu J, Nguyen T, Schwarz LA, Wilkinson AW, Gozani O, Chang HY,
Shamloo M, Luo L (2016) Molecular and Neural Functions of Rai1, the Causal Gene for Smith-
Magenis Syndrome. *Neuron* 92: 392-406
Huang WH, Tupal S, Huang TW, Ward CS, Neul JL, Klisch TJ, Gray PA, Zoghbi HY (2012)
Atoh1 governs the migration of postmitotic neurons that shape respiratory effectiveness at birth
and chemoresponsiveness in adulthood. *Neuron* 75: 799-809
Huang WH, Wang DC, Allen WE, Klope M, Hu H, Shamloo M, Luo L (2018) Early adolescent
Rai1 reactivation reverses transcriptional and social interaction deficits in a mouse model of
Smith-Magenis syndrome. *Proc Natl Acad Sci U S A*
Javed S, Lee YJ, Xu J, Huang WH (2021) Temporal dissection of Rai1 function reveals brain-
derived neurotrophic factor as a potential therapeutic target for Smith-Magenis syndrome. *Hum*
Mol Genet 31: 275-288
Javed S, Selliah T, Lee YJ, Huang WH (2020) Dosage-sensitive genes in autism spectrum
disorders: From neurobiology to therapy. *Neurosci Biobehav Rev* 118: 538-567
Kron M, Howell CJ, Adams IT, Ransbottom M, Christian D, Ogier M, Katz DM (2012) Brain
activity mapping in Mecp2 mutant mice reveals functional deficits in forebrain circuits,
including key nodes in the default mode network, that are reversed with ketamine treatment. *J*
Neurosci 32: 13860-13872
Lee SK, Lee S, Shin SY, Ryu PD, Lee SY (2012) Single cell analysis of voltage-gated potassium
channels that determines neuronal types of rat hypothalamic paraventricular nucleus neurons.
Neuroscience 205: 49-62
Loos RJF, Yeo GSH (2022) The genetics of obesity: from discovery to biology. *Nat Rev Genet*
23: 120-133

Massa SM, Yang T, Xie Y, Shi J, Bilgen M, Joyce JN, Nehama D, Rajadas J, Longo FM (2010)
Small molecule BDNF mimetics activate TrkB signaling and prevent neuronal degeneration in
rodents. *J Clin Invest* 120: 1774-1785
Moss J, Oliver C, Arron K, Burbidge C, Berg K (2009) The prevalence and phenomenology of
repetitive behavior in genetic syndromes. *J Autism Dev Disord* 39: 572-588
Ogier M, Wang H, Hong E, Wang Q, Greenberg ME, Katz DM (2007) Brain-derived
neurotrophic factor expression and respiratory function improve after ampakine treatment in a
mouse model of Rett syndrome. *J Neurosci* 27: 10912-10917
Rao NR, Abad C, Perez IC, Srivastava AK, Young JI, Walz K (2017) Rai1 Haploinsufficiency Is
Associated with Social Abnormalities in Mice. *Biology (Basel)* 6
Schultze SM, Hemmings BA, Niessen M, Tschopp O (2012) PI3K/AKT, MAPK and AMPK
signalling: protein kinases in glucose homeostasis. *Expert Rev Mol Med* 14: e1
Shao J, Yamashita H, Qiao L, Friedman JE (2000) Decreased Akt kinase activity and insulin
resistance in C57BL/KsJ-Leprdb/db mice. *J Endocrinol* 167: 107-115
Simmons DA, Belichenko NP, Yang T, Condon C, Monbureau M, Shamloo M, Jing D, Massa
SM, Longo FM (2013) A small molecule TrkB ligand reduces motor impairment and
neuropathology in R6/2 and BACHD mouse models of Huntington's disease. *J Neurosci* 33:
18712-18727
Slager RE, Newton TL, Vlangos CN, Finucane B, Elsea SH (2003) Mutations in RAI1
associated with Smith-Magenis syndrome. *Nat Genet* 33: 466-468
Smith AC, McGavran L, Robinson J, Waldstein G, Macfarlane J, Zonona J, Reiss J, Lahr M,
Allen L, Magenis E (1986) Interstitial deletion of (17)(p11.2p11.2) in nine patients. *Am J Med*
Genet 24: 393-414
Su X, Gu D, Xu L, Liang Z, Luo X, Yang P, Yang J (2021) PI3K/Akt pathway expression in
children with different obesity degrees and its relationship with glucolipid metabolism and
insulin resistance. *Am J Transl Res* 13: 6592-6598
Tan CL, Cooke EK, Leib DE, Lin YC, Daly GE, Zimmerman CA, Knight ZA (2016) Warm-
Sensitive Neurons that Control Body Temperature. *Cell* 167: 47-59 e15
Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsiani D, Kvitsani D, Fu Y, Lu J *et al*
(2011) A resource of Cre driver lines for genetic targeting of GABAergic neurons in cerebral
cortex. *Neuron* 71: 995-1013
Unger TJ, Calderon GA, Bradley LC, Sena-Esteves M, Rios M (2007) Selective deletion of Bdnf
in the ventromedial and dorsomedial hypothalamus of adult mice results in hyperphagic behavior
and obesity. *J Neurosci* 27: 14265-14274
Wu SW, Xu B (2022) Rapid and Lasting Effects of Activating BDNF-Expressing PVH Neurons
on Energy Balance. *eNeuro* 9
Xu B, Goulding EH, Zang K, Cepoi D, Cone RD, Jones KR, Tecott LH, Reichardt LF (2003)
Brain-derived neurotrophic factor regulates energy balance downstream of melanocortin-4
receptor. *Nat Neurosci* 6: 736-742
Xu B, Xie X (2016) Neurotrophic factor control of satiety and body weight. *Nat Rev Neurosci*
17: 282-292
Xu Y, Lu Y, Cassidy RM, Mangieri LR, Zhu C, Huang X, Jiang Z, Justice NJ, Xu Y, Arenkiel BR
et al (2019) Identification of a neurocircuit underlying regulation of feeding by stress-related
emotional responses. *Nat Commun* 10: 3446

Yeo GS, Connie Hung CC, Rochford J, Keogh J, Gray J, Sivaramakrishnan S, O'Rahilly S,
Farooqi IS (2004) A de novo mutation affecting human TrkB associated with severe obesity and
developmental delay. *Nat Neurosci* 7: 1187-1189

Figure 1: *Rai1* haploinsufficiency disrupts hypothalamic proteomic profile in mice

(A) ELISA showing that Bdnf protein levels in hypothalamic tissues were significantly reduced

in SMS mice compared to controls (n=4/genotype). *p<0.05, student's t-test.

(B) A schematic diagram showing the experimental strategy. Hypothalamic tissue lysates from 7-

week-old control (n = 5, 3 technical triplicates/sample) and SMS (n=8, 3 technical

triplicates/sample) mice were subjected to RPPA that utilizes a cocktail of 240 validated

antibodies to probe several key intracellular signalling pathways by quantifying the expression

levels of total and phosphorylated proteins

(C) Principal component analysis segregates the SMS and Ctrl groups where PC1 explains 47%

of the variance, and PC2 explains 25% of the variance.

(D) Heat map showing hierarchical clustering of differentially expressed phospho-proteins: most

showed reduced levels in SMS samples.

(E) Protein-protein interaction (PPI) analysis showing differentially expressed proteins and

phospho-proteins in a crosstalk network. PPI enrichment p-value: < 1.0e-16. The nodes represent

the proteins/phospho-proteins and the lines indicate physical interactions. Associations with

specific molecular pathways are indicated as color coded outer rings.

(F-G) Protein levels of two neurotrophin pathway-related nodes, PIK3CA **(F)** and phospho-NF-

kB **(G)**, which were significantly reduced in SMS samples. ****p<0.0001, student's t-test.

Figure 2: Rai1 is required in Bdnf-producing cells to regulate energy metabolism

(A) A schematic diagram showing selective deletion of *Rai1* from Bdnf-producing cells in mice.

(B) Representative images showing that in Ctrl mice (*Bdnf*$^{Cre/+}$; *Ai9*), many PVH^{Bdnf} neurons (magenta) express Rai1 (cyan, double-positive cells were indicated with yellow arrowheads). By contrast, PVH^{Bdnf} neurons lack Rai1 expression in the cKO group (yellow arrowheads) (*Bdnf*$^{Cre/+}$; *Rai1*$^{fl/fl}$; *Ai9*). Scale bars = 20μm.

(C) Percentage of PVH^{Bdnf} neurons co-expressing Rai1 in Ctrl (n=3) and cKO (n=3) mice. ****$p < 0.0001$, student's t-test.

(D) Female cKO mice (n = 12) showed a significant weight gain when compared to female Ctrl mice (n = 10).

(E) Body composition was measured with Echo-MRI, showing an increased fat mass in 26-week-old mice.

(F) Fat mass of brown adipocytes (BAT), subcutaneous inguinal (S.C.ing) and epididymal white adipose tissue (eWAT) in Ctrl and cKO mice.

(G) Representative images showing eWAT adipocyte hypertrophy of the cKO mice (left). Scale bar = 500μm. Frequency distribution of adipocytes at each cellular size (right).

(H) Female cKO mice showed significantly increased blood leptin levels.

(I) Female cKO mice showed reduced energy expenditure during the dark phase. ns indicates not significantly different.

(J) Female cKO mice showed reduced locomotor activity during the dark phase. ns indicates difference is not significant.

(K) Female cKO mice showed similar food intake as control mice. ns indicates not significant.

(L) Glucose tolerance test showing that cKO mice became hyperglycemic 20 minutes after intraperitoneal glucose administration, suggesting glucose intolerance.

Data are shown as mean±SEM. Statistics for **D-G**, **I-J**, **L**: ns indicates not significantly different, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, one or two-way ANOVA with Šidák's multiple comparisons test.

Figure 3: Rai1 loss reduces intrinsic neuronal excitability and enhances inhibitory synaptic transmission of PVH^Bdnf neurons

(A) A representative image showing a patched PVH^{Bdnf} neuron labelled by *Bdnf^{Cre}*-dependent tdTomato fluorescence signals.

(B-C) Representative traces of spontaneous firing of control (black) and cKO (red) neurons at a holding voltage of -60mV **(B)**. The average spontaneous AP firing frequencies are shown in **(C)**.

(D) The resting membrane potentials of control (black) and cKO (red) PVH^{Bdnf} neurons.

(E-F) The number of control (black) and cKO (red) PVH^{Bdnf} neurons that showed spontaneous firing (solid) or are silent (open) at resting membrane potentials **(e)** and at a holding voltage of -70mV **(f)**.

(G) Representative traces of elicited APs in control (black) and cKO (red) PVH^{Bdnf} neurons responding to an inject current (bottom, blue), which ramps up from the resting membrane potentials at 1 nA per second for 500ms.

(H) The threshold of the first AP initiated by a current ramp in control (black) and cKO (red) PVH^{Bdnf} neurons.

(I-J) The amplitude **(I)** and threshold **(J)** for the first AP firing at the holding voltage of -60mV in control (black) and cKO (red) PVH^{Bdnf} neurons.

(K) The cumulative fractions of mIPSC frequency measured in control (black) and cKO(red) PVH^{Bdnf} neurons. The average frequency of events from a 2-min stable recording of each neuron is shown in the inset.

(L) The cumulative fractions of mIPSC amplitude of control (black) and cKO (red) PVH^{Bdnf} neurons. The average amplitude of events from a 2-min stable recording of each neuron is normalized to its capacitance and shown in the inset.

(M) The area under the curve of each mIPSC event from a 2-min recording is averaged. Controls are shown in black, and cKOs are shown in red.

Data are shown as mean±SEM. Statistics: Neuron numbers are in brackets (more than 3 mice/genotype); *$p<0.05$, unpaired student's t-tests.

Figure 4: Selective *Rai1* ablation in PVH^{Bdnf} neurons induces obesity

(A) A schematic showing stereotaxic injection of adeno-associated viruses (AAVs) expressing Cas9 protein and single-guide RNAs targeting Rai1 (sgRai1) to delete *Rai1* from the PVH^{Bdnf} neurons of *Bdnf^{Cre/+}* female mice. Scale bar = 50µm.

(B) Representative images showing that Rai1 immunoreactivity (cyan) was dramatically reduced in *Bdnf*$^{Cre/+}$*; Ai9* mice injected with both sgRai1 and DIO-saCas9 viruses (Exp group, bottom row) into the PVH. In contrast, many PVH^{Bdnf} neurons in *Bdnf*$^{Cre/+}$*; Ai9* mice injected only with the sgRai1 virus showed Rai1 expression (Ctrl group, top row).

(C) The percentage of PVH^{Bdnf} neurons co-expressing Rai1 and GFP (virus) was significantly reduced in the Exp group, indicating a successful *Rai1* deletion.

(D) PVH^{Bdnf} neuron-specific *Rai1* deletion induced body weight gain in Exp mice.

(E) Body composition was measured with Echo-MRI, showing an increased fat mass deposition in 26-week-old Exp mice.

(F) Fat deposition analysis shows that the Exp group has significantly more subcutaneous inguinal (S.C.ing) and epididymal white adipose tissue (eWAT) mass (grams). ns indicates not significantly different.

(G) Representative images showing eWAT adipocyte hypertrophy of the Exp group (left). Scale bar = 500μm. Frequency distribution of adipocytes at each cellular size (right). ns indicates not significantly different.

(H) Exp and control mice showed similar food intake. ns indicates not significant, student's t-test.

(I) Blood leptin levels were significantly increased in the Exp mice. *$p < 0.05$, student's t-test.

(J) Exp and control mice showed similar energy expenditure.

(K) Exp mice showed reduced locomotor activity in the dark phase.

(L) Exp mice became hyperglycemic during the glucose tolerance test.

(M) Measurement of the area under the curve (AUC) in Exp and Ctrl mice during the glucose tolerance test. $p = 0.0522$, student's t-test.

(N) Exp mice showed increased plasma insulin levels during the glucose tolerance test.

932   Data are shown as mean±SEM. Statistics for **C-G**, **J-L**, **N**: ns indicates not significantly

933   different, *$p<0.05$, **$p < 0.01$, ***$p<0.001$, ****$p < 0.0001$, one or two-way ANOVA with

934   Šidák's multiple comparisons test.

936
Figure 5: LM22A-4 treatment partially alleviates obesity and stereotypic behaviour in
adult SMS mice

(A) Representative western blot showing p-Akt, total Akt, Histone 3 (H3, a loading control)
levels in the hypothalamus of saline-treated Ctrl, saline-treated SMS and LM22A-4-treated SMS
mice.

(B) Quantification showing phosphorylation deficits in Akt protein levels of the hypothalamus
were reversed by LM22A-4 treatment in SMS mice.

(C-E) Body weight of saline-injected control, saline-injected SMS, and LM22A-4 injected SMS
mice. Saline or LM22A-4 treatment periods were highlighted in orange. Ns indicates not
significantly different.

(F) Food intake (in grams) of saline-injected control, saline-injected SMS, and LM22A-4
injected SMS mice.

(G) Locomotor activity (number of beam breaks) for saline-injected control, saline-injected
SMS, and LM22A-4 injected SMS mice.

(H) Fat deposition analysis showing that the saline-injected SMS group has significantly more
subcutaneous inguinal (S.C.ing) and epididymal white adipose tissue (eWAT) mass (grams) than
saline-injected Ctrl mice. S.C.ing and eWAT mass in LM22A-4 injected mice is comparable to
the saline-injected mice.

(I) Insulin tolerance test performed on saline-injected control, saline-injected SMS, and LM22A-
4 injected SMS mice, two-week post daily administration of LM22A-4 or saline, shows
reductions in the blood glucose levels of LM22A-4 treated SMS mice. * Shows significance
between saline injected Ctrl and saline injected SMS group.

(J) Insulin levels in saline-injected control, saline-injected SMS, and LM22A-4 injected SMS
mice right before the start of insulin tolerance test (time point 0).

(K) High density lipoprotein (HDL) levels were restored to normal in LM22A-4 injected SMS
mice.

(L) Stereotypic rearing behaviour was fully rescued in LM22A-4 treated SMS mice.

Data are shown as mean±SEM. Statistics for **B-L**: ns indicates not significantly different,
*$p<0.05$, **$p < 0.01$, ***$p<0.001$, ****$p < 0.0001$, one or two-way ANOVA with Šidák's
multiple comparisons test.

Figure 1—figure supplement 1

Fig 1-figure supplement 1: Differentially expressed proteins in the hypothalamus of SMS mice

A heat map showing hierarchical clustering of differentially expressed proteins in the hypothalamus of SMS mice.

Figure 2—figure supplement 1

979 **Fig 2-figure supplement 1: The *Bdnf^Cre* allele labels Bdnf-producing neurons in the**

980 **paraventricular nucleus of hypothalamus**

981 (A) Schematic showing a transgenic mouse expressing *Bdnf^Cre*-dependent td-Tomato (Ai9)

982 signals in Bdnf-expressing cells.

983 **(B-E)** Colocalization of *Bdnf^Cre*-dependent td-Tomato signals with endogenous Bdnf protein.

984 Black: DAPI, Magenta: Bdnf-producing cells, Green: endogenously Bdnf. Scale bar: 50 µm.

985 **(F)** Quantification showing the percentage of Bdnf-producing cells that express p-TrkB protein

986 (left bar, magenta) and the percentage of p-TrkB-expressing cells that express Bdnf (right bar,

987 green) in the PVH. Data are shown as mean±SEM.

988 **(H-J)** Representative images showing that Bdnf-producing cells and p-TrkB signals are

989 colocalized in the PVH. Black: DAPI, Magenta: Bdnf-producing cells, Green: phospho-TrkB[816]

990 (Scale bar: 50µm).

991

992

Fig 2-figure supplement 2: Rai1 expression in Bdnf-expressing but not oxytocin-expressing magnocellular PVH neurons

(A-H) Representative images DAPI expression (blue, **A**), oxytocin (green, **B**), Bdnf-producing cells (Ai9 reporter, magenta, **C**), and Rai1 protein (cyan, **D**) in the PVH. Note that Bdnf-producing neurons are distinct from the oxytocin-expressing neurons (**E**) and that Rai1 is more selectively expressed in the Bdnf-expressing neurons (**F-H**).

(I) Quantification showing the percentage of Rai1$^+$ cells that co-express either Bdnf (left bar, magenta) or oxytocin (right bar, green) in the PVH. n=3 mice/group. Data are shown as mean±SEM.

Fig 2-figure supplement 3: Ablating *Rai1* from the Bdnf-producing cells induces body weight gain and defective energy homeostasis

(A) Schematic representation of the conditional knockout mice carrying either normal Rai1 alleles and the Ai9 reporter in Bdnf-producing cells (top) or homozygous *Rai1* deletion and the Ai9 reporter in the Bdnf-producing cells.

(B) Representative images showing the expression of DAPI (white), Rai1-expressing cells (green), and Bdnf-expressing cells (Ai9 reporter, magenta) in the PVH. The top panel shows the Ctrl group with Rai1-Ai9 co-expression (yellow arrowheads) and the bottom panel shows

reduced Rai1 and Ai9 colocalization (yellow arrowheads). Note that Rai1 expression was still

detected in non-Bdnf neurons in the PVH.

(C) Quantification showing the percentage of Bdnf-producing cells was not altered by *Rai1*

deletion, indicating that Rai1 loss did not impair the generation or survival of PVH^{Bdnf} neurons.

ns indicates not significantly different, student's t-test.

(D) The *Bdnf*$^{Cre/+}$ and *Bdnf*$^{Cre/+}$; *Rai1*$^{fl/+}$ female mice had similar body weights.

(E-G) Blood parameter analysis of triglycerides (TG), high-density lipoprotein (HDL), low-

density lipoprotein and very low-density lipoprotein (LDL+VLDL) did not differ between Ctrl

and cKO mice. ns indicates not significantly different, student's t-test.

(H) Respiratory exchange rate did not differ between Ctrl and cKO mice.

(I) In the glucose tolerance test, the cKO mice showed an increased area under curve (AUC).

***$p<0.001$, student's t-test.

(J) In the glucose tolerance test, cKO mice showed increased insulin levels at the beginning of

the test and at 30 minutes after glucose administration.

(K) In the insulin tolerance test, blood glucose level after insulin injection.

Data are shown as mean±SEM. Statistics for **D**, **H, J-K**: ns indicates not significantly different,

*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, two-way ANOVA with Šidák's multiple

comparisons test.

1032

1033

1034

Fig 2-figure supplement 4

Fig 2-figure supplement 4: Metabolic profiles of male cKO mice lacking Rai1 expression in the Bdnf-producing cells

(A) Male cKO mice gained significantly more weight than Ctrl mice beginning at 15 weeks of age.

(B) Male $Bdnf^{Cre/+}$ and $Bdnf^{Cre/+}$; $Rai1^{fl/+}$ mice had similar body weights.

(C) Echo-MRI analysis showed that 26-week-old male cKO mice had a significantly increased body weight due to increased fat but not lean mass. **p<0.01, ****p<0.0001, student's t-test.

(D) Fat disposition analysis showing the weight of brown adipocytes (BAT), subcutaneous inguinal (S.C.ing), and epididymal white adipose tissues (eWAT). ns indicates not significantly different, *p<0.05, **p<0.01, student's t-test.

(E) Male cKO mice showed eWAT cell hypertrophy. Top: Representative images of the eWAT tissues in Ctrl and cKO mice. Scale bar = 500μm. Bottom: Frequency distribution of cellular sizes.

(F-I) Blood parameter analysis showing that male cKO mice showed significantly increased leptin levels **(F)** without alterations in TG **(G)**, HDL **(H)**, LDL+VLDL **(I)**. *p< 0.05, student's t-test.

(J) No differences were found in respiratory exchange rate in male Ctrl and cKO mice.

(K) Male cKO mice showed a significantly increased food intake compared to Ctrl littermates. *p< 0.05, student's t-test.

(L) Male cKO mice showed increased energy expenditure at the dark phase.

(M) Male Ctrl and cKO mice showed similar locomotor activities.

(N-P) Glucose tolerance test shows that cKO male mice are glycemic 45 minutes post intraperitoneal glucose administration **(N)**, increased AUC **(O)**, despite significantly higher insulin levels before and 30 minutes after glucose administration **(P)**, suggesting potential insulin insensitivity.

(O) Insulin tolerance test showing that male cKO mice showed significantly higher blood glucose levels 60 minutes after intraperitoneal insulin administration. **p<0.01, student's t-test.

1063   Data are shown as mean±SEM. Statistics for **A-B, E, J, L-N**, **P-Q**: ns indicates not significantly

1064   different, *$p<0.05$, **$p < 0.01$, ***$p<0.001$, ****$p < 0.0001$, one- or two-way ANOVA with

1065   Šidák's multiple comparisons test.

Figure 3—figure supplement 1

1066

Fig 3-figure supplement 1: The cellular and synaptic properties of control and Rai1-

deficient PVH^{Bdnf} neurons

(A) Representative traces of rebound and rebound repetitive firing of PVH^{Bdnf} neurons in control

(black) and cKO (red).

(B) Representative AP waveforms at holding voltages of -70mV, -60mV, and -50mV in control

(left) and -50mV and -45mV in cKO (right).

(C) The half-width of AP initiated at a holding voltage of -60mV (neuronal numbers in brackets,

p=0.374, student's t-test).

(D) Representative traces of miniature recordings in control (black) and cKO (red) PVH^{Bdnf}

neurons. The traces were recorded with a Cs-based internal at a holding voltage of -70mV in

aCSF (top), spontaneous mEPSCs were recorded in aCSF containing 1uM TTX at a holding

voltage of -70mV (middle), and mIPSCs were recorded at holding voltage of +10mV (bottom).

(E) Quantification showing that the cellular size of cKO PVH^{Bdnf} neurons was significantly

smaller than Ctrl neurons. ****p<0.0001, student's t-test.

1081

1082

Figure 4—figure supplement 1

Fig 4-figure supplement 1: Metabolic profile of mice lacking *Rai1* in PVH^{Bdnf} neurons

(A) Representative images showing co-expression of virus-infected cells (GFP/green), *Bdnf^{Cre}*-dependent Ai9 signals (Bdnf cells, magenta), and endogenous Rai1 (cyan). Yellow arrowheads indicate Bdnf neurons infected with AAV and in these cells, Rai1 signals were diminished in Exp but not Ctrl mice. Scale bar = 50μm.

(B-C) The total number of PVH^{Bdnf} neurons **(B)** and total number of PVH^{Bdnf} neurons infected with AAVs **(C)** per slice showing that Rai1 deletion or virus infection did not alter the total number of PVH^{Bdnf} neurons.

(D) Rai1 expression was significantly reduced in PVH^{Bdnf} neurons infected with both AAVs

when compared to PVH^{Bdnf} neurons infected with sgRai1-GFP AAV alone.

(E) The number of PVH^{Bdnf} neurons that co-express Rai1 was significantly reduced in the Exp

group. **(F)** Respiratory exchange rate remained similar in Ctrl and Exp mice.

(G-I) Blood parameter analysis showing the levels of TG, HDL, and LDL+VLDL were similar

in Ctrl and Exp mice.

(J) In the insulin tolerance test, Exp group showed a trend towards increased blood glucose level

after insulin injection.

Data are shown as mean±SEM. Statistics for **F**, **J**: ns indicates not significantly different, two-

way ANOVA. Statistics for **B-E**, **G-I**: **$p < 0.01$, student's t-test.

1102

1103

1104

Figure 5—figure supplement 1

Figure 5—figure supplement 1: LM22A-4 treatment does not disrupt energy homeostasis and repetitive rearing in the Ctrl mice

(A) LM22A-4 concentration in the hypothalamus and forebrain of SMS mice 1 hour (black dots) and 3 hours (red dots) after simultaneous intranasal and IP injections.

(B) Four independent sets of western blotting analyses showing reduced p-Akt levels in the hypothalamus of saline-treated SMS mice, which were increased by LM22A-4 treatment. Total Akt protein was used as a control.

(C) Similar body weight of Ctrl mice treated with either saline or LM22A-4.

(D) Energy expenditure was not different between groups in the light vs. dark phase.

1114 **(E)** Respiratory exchange rates were not different between groups in the light vs. dark phase.

1115 **(F)** Repetitive rearing behaviour in Ctrl mice was not altered by LM22A-4 treatment. ns
1116 indicates not significantly different, student's t-test.

1117 Data are shown as mean±SEM. Statistics for **C-E**: ns indicates not significantly different, two-
1118 way ANOVA. Statistics for **F**: student's t-test.

1119

Figure 5—figure supplement 2

1120

Figure 5—figure supplement 2: LM22A-4 treatment in adult SMS mice is insufficient to improve social interaction deficit

(A) Schematic showing the tube test that assesses social interaction between stranger mice of different genotypes.

(B) 7-week-old Ctrl mice won significantly more than SMS mice before the LM22A-4 treatment.

(C) 10-week-old saline-injected Ctrl mice won significantly more than the saline-injected SMS mice (two weeks post-LM22A-4 treatment).

(D) 10-week-old saline-injected Ctrl mice won significantly more than LM22A-4 injected SMS mice.

(E) 10-week-old saline-injected Ctrl mice showed a similar winning rate to LM22A-4-injected SMS mice.

Data are shown as mean±SEM. Statistics for **B-E**, ns indicates not significantly different, *p<0.05, ***p<0.001, ****p<0.0001, student's t-test.