Smith-Magenis syndrome protein RAI1 regulates body weight homeostasis through

hypothalamic BDNF-producing neurons and neurotrophin downstream signalling

Sehrish Javed[1,2], Ya-Ting Chang[1,2], Yoobin Cho[1,2], Yu-Ju Lee[1,2], Hao-Cheng Chang[1,2], Minza

Haque[1,2], Yu Cheng Lin[1,2], Wei-Hsiang Huang[1,2],*

Affiliations:

[1]Department of Neurology and Neurosurgery, Centre for Research in Neuroscience, McGill

University, Montréal, QC H3G 1A3, Canada

[2]Brain Repair and Integrative Neuroscience Program, The Research Institute of the McGill

University Health Centre, Montréal, QC H3G 1A3, Canada

*Corresponding author. Email: wei-hsiang.huang@mcgill.ca

Abstract

Retinoic acid-induced 1 (*RAI1*) haploinsufficiency causes Smith-Magenis syndrome (SMS), a genetic disorder with symptoms including hyperphagia, hyperlipidemia, severe obesity, and autism phenotypes. RAI1 is a transcriptional regulator with a pan-neural expression pattern and hundreds of downstream targets. The mechanisms linking neural *Rai1* to body weight regulation remain unclear. Here we find that hypothalamic brain-derived neurotrophic factor (BDNF) and its downstream signalling are disrupted in SMS (*Rai1*$^{+/-}$) mice. Selective *Rai1* loss from all BDNF-producing cells or from BDNF-producing neurons in the paraventricular nucleus of the hypothalamus (PVH) induced obesity in mice. Electrophysiological recordings revealed that *Rai1* ablation decreased the intrinsic excitability of PVH^{BDNF} neurons. Chronic treatment of SMS mice with LM22A-4 engages neurotrophin downstream signalling and delayed obesity onset. This treatment also partially rescued disrupted lipid profiles, insulin intolerance, and stereotypical repetitive behaviour in SMS mice. These data argue that RAI1 regulates body weight and metabolic function through hypothalamic BDNF-producing neurons and that targeting neurotrophin downstream signalling might improve associated SMS phenotypes.

Introduction

Obesity is an increasing global concern: 20% of the world's adult population is expected to become obese by 2025 if current trends persist (Collaboration, 2017). However, the complex environmental, behavioural, and genetic contributions to obesity make this a complex condition to model and address. Monogenic obesity disorders provide a consistent system where the pathophysiology, genetic predisposition, and neurobiology underlying phenotypes can potentially provide insights that translate to polygenic obesity (Loos & Yeo, 2022).

One such monogenic disorder is Smith-Magenis syndrome (SMS; OMIM #182290) (Smith *et al*, 1986), a neurogenetic condition frequently associated with obesity. 80% of SMS cases are caused by interstitial deletions of the 17p11.2 chromosomal region containing a dosage-sensitive gene, *retinoic acid-induced 1* (*RAI1*) (Slager *et al*, 2003). Most SMS patients are overweight (40% > 95th percentile weight, 80% > 85th percentile weight) and exhibit over-eating tendencies by adolescence (Alaimo *et al*, 2015; Burns *et al*, 2010; Gandhi *et al*, 2022). 20% of SMS patients do not have 17p11.2 deletions but instead carry *RAI1* point mutations that are still frequently associated with obesity (Boot *et al*, 2021; Edelman *et al*, 2007). These clinical data indicate that RAI is critical for body weight regulation.

The human RAI1 protein shares >80% sequence identity with the mouse RAI1 protein, allowing us to study RAI1 function *in vivo* (Huang *et al*, 2016; Javed *et al*, 2020). We previously found that *Rai1* encodes a transcriptional regulator that binds to the promoter or enhancer regions of its target genes, most of which are involved in neurodevelopment and synaptic transmission (Huang *et al.*, 2016). SMS (*Rai1*$^{+/-}$) mice exhibit obesity associated with hyperphagia and increased adiposity (Burns *et al.*, 2010; Huang *et al.*, 2016). Interestingly, RAI1 may have a sexually

dimorphic function, as female SMS mice exhibit more pronounced obesity than males (Burns *et*

al., 2010; Huang *et al.*, 2016).

Using a homozygous *Rai1* conditional knockout (*Rai1*-cKO) mice, we found that *Rai1* deletion

in either subcortical excitatory neurons (vGlut2[+]) or single-minded homolog 1 (Sim1[+]) neurons

induced obesity in mice (Huang *et al.*, 2016). Sim1[+] neurons reside in several brain regions

including the nucleus of the lateral olfactory tract, the supraoptic nucleus, the medial amygdala,

and the paraventricular nucleus of the hypothalamus (PVH), a region that is critical for body

weight regulation (Balthasar *et al*, 2005). Sim1[+] neurons in the PVH are composed of several

molecularly defined cell types, including those that express oxytocin, vasopressin, corticotropin-

releasing hormone, somatostatin, melanocortin-4-receptor (MC4R), growth hormone-releasing

hormone (GHRH), or brain-derived neurotrophic factor (BDNF) (An *et al*, 2015). Deleting *Bdnf*

from the PVH results in obesity due to increased food intake, reduced energy expenditure, and

reduced locomotion (An *et al.*, 2015). While PVH neuronal activity is reduced during feeding

(Xu *et al*, 2019), activation of PVH[BDNF] neurons results in weight loss (Wu & Xu, 2022). We

recently found that RAI1 promotes *Bdnf* expression in the hypothalamus throughout life (Javed

et al, 2021). We hypothesized that RAI11 may be connected to BDNF-producing neurons in

PVH, explaining how a neural transcription factor regulates body weight homeostasis.

BDNF regulates energy metabolism by binding to its cognate receptor tropomyosin receptor

kinase B (TRKB) (An *et al.*, 2015) and may thus provide a potential therapeutic target for SMS.

However, the link between *RAI1, BDNF*, and SMS symptoms is unclear. Patients carrying

monogenic mutations in either *BDNF* (Gray *et al*, 2006) or *NTRK2* (encoding TRKB) (Yeo *et al*,

2004) experience severe hyperphagic obesity. Moreover, obesity and associated metabolic

dysfunctions such as impaired glucose and insulin sensitivity are correlated with impaired PI3K-

AKT pathway, a TRKB downstream target (Su *et al*, 2021). Given that (1) *Bdnf* overexpression

during early embryonic development or (2) targeting *Bdnf* overexpression to the PVH of

adolescent SMS ($Rai1^{+/-}$) mice protected them from weight gain and metabolic defects (Javed *et*

al., 2021), and that (3) a point mutation in *RAI1* disrupts its ability to transcriptionally enhance

BDNF expression *in vitro* (Abad *et al*, 2018), pharmacological targeting of neurotrophin

downstream signalling is an attractive and yet unexplored therapeutic avenue for SMS.

Building on our previous discoveries, here we examine the neurobiological functions of RAI1 in

BDNF-producing neurons, specifically those that reside in the PVH. Importantly, we evaluated

the involvement of the BDNF-TRKB pathways in SMS pathogenesis using a small molecule

drug (LM22A-4) to enhance neurotrophin downstream signalling. Our work indicates that

PVH^{BDNF} neurons depend on RAI1 to maintain normal neuronal activity and regulate body

weight homeostasis, and dysfunction of BDNF downstream signalling contributes to obesity

associated with SMS mice.

 Results

 Rai1 **deficiency dysregulates specific downstream targets of neurotrophin in the mouse**

 hypothalamus

Multiple lines of evidence indicated that SMS mice show decreased *Bdnf* mRNA expression in

the hypothalamus (Burns *et al.*, 2010; Huang *et al.*, 2016; Javed *et al.*, 2021). To determine if

BDNF protein levels were consequently altered in the hypothalamus, we performed an enzyme-

linked immunosorbent assay (ELISA) and found that BDNF protein levels were significantly

reduced in 7-week-old pre-symptomatic SMS mice relative to controls (**Fig 1A**). To quantify if

molecular pathways associated with BDNF downstream signalling were altered, we subjected

hypothalamic tissue lysates from 7-week-old pre-obese SMS and control (Ctrl) mice to reverse

phase protein array (RPPA) (**Fig 1B**). Control and SMS samples are clustered separately in the

principal component analysis (**Fig 1C**).

Hierarchical clustering analysis indicated that most differentially expressed phosphorylated

proteins (**Fig 1D**) and total proteins (**Figure 1—figure supplement 1**) were downregulated in

SMS mice. We mapped total proteins, as well as phospho-proteins, that were differentially

expressed in Ctrl and SMS samples onto the interconnected biological network and identified a

subset of connected nodes downregulated in SMS (FDR < 10e-5) belong to the neurotrophin,

PI3K-AKT, and mTOR signalling cascades (**Fig 1E**). These molecules are known to play a

critical role in regulating body weight homeostasis and can be activated by BDNF either directly

(neurotrophin pathway) or indirectly (PI3K-AKT and mTOR pathways) (Cota *et al*, 2006;

Schultze *et al*, 2012). The expression levels of PIK3CA, a common node in these pathways, were

significantly reduced in the SMS mice compared to controls (**Fig 1F**). Moreover, the expression

levels of phosphorylated NF-κB (nuclear factor-kappa B), a downstream target of the

neurotrophin pathway, were significantly reduced in the SMS group (**Fig 1G**). Together, these

results suggest that *Rai1* loss impairs specific pathways, including neurotrophin signalling, and

indicate that dysfunction of BDNF-producing cells could contribute to SMS pathogenesis.

BDNF-producing cells depend on RAI1 to regulate body weight homeostasis

To target BDNF-producing cells, we used a $Bdnf^{Cre/+}$ mouse line that carries a viral 2A

oligopeptide and Cre recombinase immediately upstream of *Bdnf*'s endogenous STOP codon

(Tan *et al*, 2016). We crossed the $Bdnf^{Cre/+}$ mice with tdTomato-Ai9 reporter mice and found that

Ai9 signals faithfully reflected endogenous BDNF protein expression in the PVH (**Figure 2—**

figure supplement 1A-E). We also found that RAI1 is expressed in PVH^{BDNF} cells that are

distinct from larger magnocellular cells, such as the PVH oxytocin neurons (**Figure 2—figure**

supplement 2A-I). Specifically, 55% of PVH^{BDNF} neurons express RAI1, whereas only 2% of

$PVH^{Oxytocin}$ neurons express RAI1 (**Figure 2—figure supplement 2I**). These data suggest that

RAI1 is enriched in BDNF-producing neurons in the mouse PVH.

To determine if BDNF-producing cells depend on RAI1 to regulate body weight, we deleted one

or both copies of *Rai1* using the $Bdnf^{Cre/+}$ and the conditional $Rai1^{fl/fl}$ mice (Huang *et al.*, 2016)

(**Fig 2A, Figure 2—figure supplement 3A**). Conditional knockout mice (cKO: $Bdnf^{Cre/+}$;

$Rai1^{fl/fl}$) showed a significant reduction in Rai1 expression in PVH^{BDNF} neurons compared to

controls (**Fig 2B-C**, **Figure 2—figure supplement 3B**). The total number of BDNF-producing

cells in the PVH remained similar in Ctrl and cKO mice (**Figure 2—figure supplement 3C**),

indicating that RAI1 loss did not affect PVH neurogenesis or survival.

We monitored body weight weekly and found that male and female cKO mice showed a

significant increase in body weight beginning at 15 weeks of age compared to controls (Ctrl:

$Bdnf^{Cre/+}$) (**Fig 2D**, **Figure 2—figure supplement 4A**). However, conditional *Rai1* heterozygous

mice ($Bdnf^{Cre/+}$*; Rai1$^{fl/+}$*) did not show a pathological increase in body weight (**Figure 2—figure**

supplement 3D, **Figure 2—figure supplement 4B**), suggesting that non-BDNF-producing cells

also contribute to SMS pathogenesis. EcoMRI found that 26-week-old cKO male and female

mice showed increased fat mass but not lean mass (**Fig 2E**, **Figure 2—figure supplement 4C**).

Further dissection of fat tissues from different organs showed an increased mass of subcutaneous

inguinal (S.C. Ing.) fat in cKO mice of both sexes and an increased mass of epididymal adipose

tissues (eWAT) in female cKOs (**Fig 2F**, **Figure 2—figure supplement 4D**).

In contrast, the total mass of brown adipose tissue (BAT) remained unchanged between groups

(**Fig 2F**, **Figure 2—figure supplement 4D**). In both male and female cKO mice, we found

hypertrophic eWAT adipocytes (**Fig 2G**, **Figure 2—figure supplement 4E**) and elevated blood

leptin levels (**Fig 2H**, **Figure 2—figure supplement 4F**). However, there was no significant

alteration in blood lipids such as triglycerides (TG), high-density lipoprotein (HDL), low-density

lipoproteins (LDL), and very low-density lipoprotein (VLDL) in either male or female cKO mice

(**Figure 2—figure supplement 3E-G**, **Figure 2—figure supplement 4G-I**). These data suggest

that obesity caused by ablating *Rai1* from BDNF-producing cells is associated with increased fat

mass, fat deposition, and leptin levels.

Obesity could result from increased food intake, reduced energy expenditure, or both. Similar to

SMS mice, male and female cKO mice did not have defects in respiratory exchange rates when

measured by indirect calorimetry (**Figure 2—figure supplement 3H**, **Figure 2—figure**

supplement 4J). Interestingly, cKO mice showed sex-specific differences in energy expenditure.

During the dark phase, female cKO mice showed reduced energy expenditure (Fig 2I) and

decreased total locomotor activity (**Fig 2J**). In contrast, total food intake remained unchanged

(**Fig 2K**). Male cKO mice became hyperphagic (**Figure 2—figure supplement 4K**), showed

increased energy expenditure during the dark phase (**Figure 2—figure supplement 4L**), and

exhibited normal locomotor activity when compared to controls (**Figure 2—figure supplement**

4M). These results suggest that obesity in male cKO mice is attributable to overeating despite

increased energy expenditure.

To determine if *Rai1* loss from BDNF-producing cells impacts metabolic function, a glucose

tolerance test (GTT) was performed. Interestingly, cKO mice of both sexes showed normal blood

glucose levels right before the test but became hyperglycemic 20 minutes (female cKO) and 45

minutes (male cKO) after glucose administration (**Fig 2L**, **Figure 2—figure supplement 4N**).

This change was accompanied by a significantly higher area under the curve (AUC) for both

groups (**Figure 2—figure supplement 3I**, **Figure 2—figure supplement 4O**). While the male

cKO mice remained hyperglycemic throughout the GTT (**Figure 2—figure supplement 4N**), the

glucose levels in female cKO mice returned to normal towards the end (**Fig 2L**). We also found

increased blood insulin levels during the GTT in both sexes (**Figure 2—figure supplement 3J**,

Figure 2—figure supplement 4P). In the insulin tolerance test (ITT), female cKO mice showed

moderately elevated glucose levels 20 minutes into the test (**Figure 2—figure supplement 3K**).

By contrast, male cKO mice showed considerably higher glucose levels than controls in the ITT

(~6.5 ng/ml for cKO and ~2.0ng/ml for Ctrl) during the 60- to 120-minute intervals (**Figure 2—**

figure supplement 4Q). These experiments demonstrate that BDNF-producing cells depend on

RAI1 to regulate body weight, adiposity, glucose and insulin tolerance in mice.

***Rai1* deletion reduces PVH[BDNF] neuronal excitability**

BDNF-producing neurons are widely distributed in the hypothalamus, being found in the

dorsomedial hypothalamus (Unger *et al*, 2007), ventromedial hypothalamus (Xu *et al*, 2003), and

PVH (An *et al.*, 2015). To understand how *Rai1* loss affects intrinsic neuronal excitability and

synaptic transmission of PVH[BDNF] neurons that underlie energy homeostasis (An *et al.*, 2015),

we fluorescently labelled them with Cre-dependent tdTomato (**Fig. 3A**). We found that a

hyperpolarized step followed by increasing the holding voltage at resting membrane potentials

(Rm) was able to initiate rebound firing or rebound repetitive firing in most PVH[BDNF] neurons

(**Figure 3—figure supplement 1A**). Hyperpolarization-induced I_H currents and low

depolarization events were observed in some PVH[BDNF] neurons (**Figure 3—figure supplement**

1A). cKO PVH[BDNF] neurons showed reduced action potential (AP) firing frequency at a holding

voltage of -60mV compared to Ctrl PVH[BDNF] neurons (**Fig 3B-C**). The resting membrane

potential (Rm) of cKO (-52mV) neurons showed a 10mV positive shift compared to Ctrl neurons

(-62mV) (**Fig 3D**). We also found that fewer cKO PVH[BDNF] neurons showed spontaneous firing

at Rm (**Fig 3E**, Ctrl:10/13; cKO:5/12) and at a holding voltage of -70mV (**Fig 3F**, Ctrl:6/12;

cKO:1/12) compared to Ctrl PVH[BDNF] neurons. The threshold of the first AP initiated by a ramp

current injection (1 nA/sec) from Rm was significantly increased in cKO neurons (**Fig 3G-H**).

Because a wider shoulder and lower AP amplitude were observed when held at more depolarized

potentials (**Figure 3—figure supplement 1B**), we compared the AP waveforms at a holding

voltage of -60mV, close to the Rm of both genotypes. The AP of the cKO group exhibited a

lower peak amplitude (measuring from the threshold to peak) (**Fig 3I**), a normal threshold (**Fig**

3J), and a normal half-width (**Figure 3—figure supplement 1C**). In summary, we found that

both passive and AP properties of cKO PVH[BDNF] neurons were altered due to RAI1 loss. The

Rm of cKO neurons shifted to a more positive voltage than the gating threshold of voltage-gated

sodium channels (6/12 were more positive than -55mV), which was unfavourable for their

transition from the inactivation state to the closed state. Combined with the higher AP threshold

of cKO neurons, these changes contribute to a decreased excitability in cKO PVH[BDNF] neurons.

PVN neurons receive glutamatergic and GABAergic inputs and are tonically inhibited by

GABAergic neurons (Colmers & Bains, 2018; Gao *et al*, 2017). We observed a slight rightward

shift of the probability of miniature inhibitory postsynaptic current (mIPSC) frequency in cKO

PVH[BDNF] neurons, although the average frequency (**Fig 3K**) was not significantly different

between groups. The probability of mIPSC amplitude also showed a right shift without a

significant change (**Fig 3L, Figure 3—figure supplement 1D**). However, we observed a

significantly increased area under the curve (**Fig 3M**). Interestingly, we found that cKO

PVH[BDNF] neurons had a smaller cellular diameter (**Figure 3—figure supplement 1E**). Overall,

our data argue that *Rai1* deletion decreases the excitability and AP firing of PVH[BDNF] neurons

without significant changes in synaptic inhibitory transmission.

RAI1 is necessary for PVH[BDNF] neurons to regulate energy homeostasis

Reduced PVH[BDNF] neuronal activity in cKO mice suggested that RAI1 loss in BDNF-producing

PVH neurons might be sufficient to induce defective energy homeostasis *in vivo*. To test this

hypothesis by deleting *Rai1* only in PVH[BDNF] neurons but not other BDNF-producing cells, we

performed PVH[BDNF]-specific *Rai1* deletion using a CRISPR/Cas9 system packaged into AAV9.

Two viruses were delivered: one encoding a reporter molecule (GFP) and three single guide

RNAs targeting the *Rai1* gene (sgRai1), and the other carrying Cre-dependent *Staphylococcus*

aureus CRISPR-associated protein 9 (saCas9) (**Fig 4A**). Hereafter, our experiments focus on

female SMS mice since they showed more consistent defects in fat distribution (Burns *et al.*,

2010) and body weight regulation (Huang *et al.*, 2016) than males. Bilateral administration of

two AAVs into the PVH of female $Bdnf^{Cre/+}$ mice (Exp group) at 3 weeks of age resulted in a

significant loss of Rai1 protein expression in PVH^{BDNF} neurons when compared to $Bdnf^{Cre/+}$ mice

injected with an AAV encoding the *sgRai1* and GFP (Ctrl group) (**Fig 4B-C**, **Figure 4—figure**

supplement 1A). Quantification of AAV-infected (GFP^+) BDNF neurons showed that while the

total number of $BDNF^+$ (**Figure 4—figure supplement 1B**) and $BDNF^+$ GFP^+ (**Figure 4—**

figure supplement 1C) cells were not different in the two groups, the percentage of AAV-

infected $BDNF^+$ cells that express RAI1 was significantly reduced in the Exp group (**Fig 4C**,

Figure 4—figure supplement 1D-E). Moreover, the virus expression did not spread to the non-

PVH neighboring nuclei (**Figure 4—figure supplement 1F**). Deleting *Rai1* from the PVH^{BDNF}

neurons resulted in a significant increase in body weight beginning at 17 weeks of age (**Fig 4D**).

Similar to the cKO ($Bdnf^{Cre/+}; Rai1^{fl/fl}$) mice, obesity in 26-week-old Exp mice was due to

increased fat mass (**Fig 4E**) and more specifically, increased subcutaneous inguinal (S.C. ing)

and epididymal white adipose tissues (eWAT) but not BAT (**Fig 4F**). We also found that eWAT

adipocytes became hypertrophic (**Fig 4G**). Similar to the cKO mice, food intake did not differ

between Exp and Ctrl groups (**Fig 4H**), likely due to the high level of satiety signals in the brain

caused by increased blood leptin levels (**Fig 4I**). We also found that energy expenditure (**Fig 4J**),

respiratory exchange rate (**Figure 4—figure supplement 1G**), and blood lipid parameters,

including TG (**Figure 4—figure supplement 1H**), HDL (**Figure 4—figure supplement 1I**),

LDL and VLDL levels (**Figure 4—figure supplement 1J**), were similar in Exp and Ctrl groups,

consistent with the cKO data (**Figure 2**). Interestingly, Exp mice showed significantly less

locomotor activity during the dark phase than the Ctrl mice (**Fig 4K**). A GTT indicated that Exp

mice have a hyperglycemic profile (**Fig 4L**), accompanied by a slightly higher area under the

curve for blood glucose levels (p=0.05) (**Fig 4M**). Insulin levels in Exp mice were also

significantly elevated 15 minutes after glucose administration (**Fig 4N**).

While Exp mice showed a trend towards increased glucose levels in insulin tolerance tests, the

change was not statistically different (**Figure 4—figure supplement 1K**). Altogether, these

experiments indicate that conditional deletion of *Rai1* from PVH[BDNF] neurons during

adolescence impairs the regulation of body weight, adiposity, glucose tolerance, and locomotion.

These changes recapitulate most phenotypes observed in cKO mice with *Rai1* deleted in BDNF-

producing cells from early development.

Pharmacologically enhancing neurotrophin downstream signalling partially alleviates

obesity in SMS mice

Our findings so far indicate that multiple pathways, including neurotrophin signalling, are

dysregulated in SMS mice, and that Rai1 loss from BDNF-producing cells, including PVH[BDNF]

neurons, impairs energy homeostasis *in vivo*. Next we sought to examine the contribution of

BDNF dysfunction in SMS pathogenesis by pharmacologically activating downstream targets of

the neurotrophin pathway.

We systematically treated mice with LM22A-4, a potential small molecule BDNF loop domain

mimetic (Massa *et al*, 2010). To measure the bioavailability of LM22A-4 in the SMS brain, we

simultaneously administered LM22A-4 intranasally (5 mg/kg body weight) and intraperitoneally

(50 mg/kg body weight), using the highest doses used in previous studies (Canals *et al*, 2004;

Chang *et al*, 2006; Gu *et al*, 2022; Kron *et al*, 2012; Ogier *et al*, 2007). We then harvested

forebrain and hypothalamic tissues one or three hours after drug administration. The

spectrometric analysis found a significantly higher LM22A-4 concentration in both brain regions

one-hour post-treatment, while its concentration decreased three hours after treatment (**Figure**

5—figure supplement 1A). Consistent with the reduced neurotrophin signalling observed in our

proteomic analysis, saline-treated SMS mice showed reduced phosphorylated-AKT (p-AKT; **Fig**

5A, **Figure 5—figure supplement 1B**, **Figure 5—source data 1, Figure 5—figure**

supplement 1 source data 1) levels in the hypothalamus. LM22A-4 treatment significantly

increased the levels of p-AKT in the SMS mice 1-hr post-treatment (**Fig 5A**, **Figure 5—figure**

supplement 1B, **Figure 5—source data 1**, **Figure 5—figure supplement 1 source data 1**),

indicating increased signalling downstream of the BDNF pathway.

We next examined if chronic LM22A-4 treatment was sufficient to reverse or modify the

progression of the obesity phenotype in adult SMS mice. Specifically, we treated SMS mice

daily from 8 to 14 weeks of age, during which SMS mice began to gain significantly more

weight. Saline-treated SMS mice (**Fig 5C**) were significantly heavier than saline-treated control

mice from 9 weeks of age onward. Interestingly, LM22A-4-treated SMS mice showed a similar

body weight to saline-treated control littermates until 13 weeks of age but became obese by 14

weeks of age (**Fig 5D**). Consistent with a partial rescue, LM22A-4-treated SMS mice were

significantly less obese than saline-treated SMS mice throughout the treatment period (**Fig 5E**).

Control mice treated with LM22A-4 did not show a further reduction in body weight (**Figure**

5—figure supplement 1C), indicating that the therapeutic effects were specific to SMS

pathogenesis.

We measured food intake and found that LM22A-4 did not significantly rescue increased food

intake in SMS mice (**Fig 5F**). SMS mice showed normal energy expenditure (**Figure 5—figure**

supplement 1D) and respiratory exchange rate (**Figure 5—figure supplement 1E**), neither of

which was altered by LM22A-4 treatment. Interestingly, we did observe a partial rescue of

locomotor activity in LM22A-4-treated SMS mice that was not significantly different from

saline-treated Ctrl or SMS mice (**Fig 5G**). We also found that the LM22A-4 treatment partially

rescued fat deposition (**Fig 5H**) and completely rescued HDL levels (**Fig 5I**) in SMS mice.

During an insulin tolerance test, we found that saline-treated SMS mice showed increased

baseline insulin levels (**Fig 5J**) and blood glucose levels during the test (**Fig 5K**), both of which

were partially rescued by LM22A-4 treatment.

In addition to a deficit in energy homeostasis, SMS mice showed two highly reproducible

behavioural deficits: increased stereotypic rearing in the open field and decreased social

dominance in the tube test (Huang *et al*, 2018; Rao *et al*, 2017). Saline-treated SMS mice

showed significantly more rearing behaviour when compared to the saline-treated controls (**Fig**

5L). Interestingly, the LM22A-4 treatment reduced the repetitive rearing behaviour in SMS mice

(**Fig 5L**). LM22A-4 treatment did not change repetitive behaviours in control mice (**Figure 5—**

figure supplement 1F), suggesting the effect is specific to *Rai1* haploinsufficiency. However,

we did not observe a rescue of social dominance deficits in SMS mice (**Figure 5—figure**

supplement 2A-E). Together, these data indicate that LM22A-4 treatment delayed the onset of

obesity and fully rescued repetitive behaviour in SMS mice, suggesting that neurotrophin

downstream signalling dysfunction contributes to changes in the mouse SMS brain linked to

obesity and behavioural phenotypes.

Discussion

The hypothalamus and neurotrophin signalling in this region control body weight by regulating

the intricate balance between energy intake and expenditure (Xu & Xie, 2016). The role of

BDNF-producing neurons in human obesity disorders and how the activity of these cells is

regulated remained poorly studied. We found that *Rai1* deficiency, linked to Smith-Magenis

syndrome (SMS), regulates BDNF expression and maintains intrinsic excitability in PVH^{BDNF}

neurons.

Consistent with this mechanism, we found that removing RAI1 from BDNF-producing neurons

or PVH^{BDNF} neurons results in imbalanced energy homeostasis and obesity. A small molecule

drug LM22A-4 led to increased AKT phosphorylation and partially rescued locomotor activity,

insulin intolerance, and HDL levels, while delaying the onset of obesity in SMS mice. These

findings highlight a potential role for hypothalamic BDNF-producing neurons, specifically

PVH^{BDNF} neurons, in SMS pathogenesis and highlight BDNF signalling as a potentially

druggable pathway to improve energy homeostasis defects and repetitive behaviours commonly

observed in SMS patients (Burns *et al.*, 2010; Moss *et al*, 2009).

The involvement of different PVH subtypes in human obesity is only beginning to be elucidated.

Dysfunction of oxytocin PVH neurons is linked to the Prader-Willi syndrome (Lee *et al*, 2012),

another genetic disorder associated with obesity. Since *Rai1* expression is enriched in PVH^{BDNF}

neurons but not PVH^{Oxytocin} neurons, our data suggest that distinct pathophysiological

mechanisms underlie SMS and Prader-Willi syndrome.

The conditional knockout analysis allowed us to precisely map the cell-specific origin of

neuropathological phenotypes, such as SMS-like symptoms in mice. Our previous work found

that deleting *Rai1* from cortical excitatory neurons using an *Emx1^{Cre}* allele (Gorski *et al*, 2002),

from the GABAergic neurons using a *Gad2^{Cre}* allele (Taniguchi *et al*, 2011), or from astrocytes

using a $Gfap^{Cre}$ allele (Garcia *et al*, 2004) did not induce SMS-like obesity (Huang *et al.*, 2016).

In contrast, obesity could be induced by ablating *Rai1* from Sim1-lineage neurons and, to a lesser

extent, SF1-lineage neurons (Huang *et al.*, 2016). This analysis suggests that Rai1 expression in

PVH is more important than in the ventromedial nucleus of the hypothalamus (VMH) for body

weight regulation. Interestingly, Although BDNF is required in the VMH and DMH to regulate

body weight (Unger *et al.*, 2007), embryonic deletion of *Bdnf* from the SF1-lineage populations

including the VMH did not result in obesity (Kamitakahara *et al*, 2016; Yang *et al*, 2016). By

contrast, deleting *Bdnf* from Sim1-lineage neurons including the PVH resulted in a more severe

obesity phenotype (An *et al.*, 2015). This is consistent with the idea that BDNF expression in

PVH is critical for energy homeostasis. We found that *Rai1* expression in BDNF-producing cells

regulates energy homeostasis in a sexually dimorphic manner. Specifically, female cKO mice

showed normal food intake, decreased energy expenditure, and reduced locomotor activity. In

contrast, male cKO mice showed increased food intake with no changes in locomotor activity.

Intriguingly, energy expenditure was increased in male cKO mice, likely due to the higher

demand to maintain normal activity levels in obese mice. In both sexes, cKO mice showed an

increased fat mass and adiposity with no change in lean mass, highlighting an essential role for

Rai1 in regulating fat deposition. Our data also indicate that BDNF-producing cells are only

partially responsible for obesity in SMS: male and female cKO mice become overweight by 15

weeks, in contrast to SMS mice which become obese by 9 weeks of age, suggesting that non-

BDNF neurons contribute to SMS phenotypes.

Notably, mice lacking one copy of *Rai1* in the BDNF-producing cells do not exhibit

obesity, whereas SMS patients and SMS mice show pronounced obesity (Burns *et al.*, 2010;

Huang *et al.*, 2016; Smith *et al*, 2005). This indicates that although reduced *Bdnf* expression and

BDNF-producing neurons contribute to regulating body weight, additional molecular changes

and other hypothalamic populations also play important roles in regulating body weight

homeostasis in SMS. Our RPPA data suggest that mTOR signalling is also misregulated in

addition to the reduced activation of the neurotrophin downstream cascades. Hypothalamic

mTORC1 is crucial to regulate glucose release from the liver, peripheral lipid metabolism, and

insulin sensitivity (Burke *et al*, 2017; Caron *et al*, 2016; Smith *et al*, 2015), while mTORC2

regulates glucose tolerance and fat mass (Kocalis *et al*, 2014). How the impaired mTOR

signalling contributes to energy homeostasis defects in SMS and the therapeutic potential of

targeting this pathway to treat SMS-related obesity remains unclear and warrants future

investigation.

What additional RAI1-dependent hypothalamic cell types residing in brain regions other than

PVH regulate obesity in SMS? Other important cell types such as TRKB neurons within the

PVH (An *et al*, 2020) and several RAI1-expressing hypothalamic nuclei including the arcuate

nucleus, ventromedial nucleus of the hypothalamus (VMH), and lateral hypothalamus all play

important roles in regulating energy homeostasis. POMC- and AGRP-expressing neurons within

the arcuate nucleus are known to regulate food intake and glucose and insulin homeostasis

(Quarta *et al*, 2021; Vohra *et al*, 2022). Therefore, *Rai1* function in these neurons could

contribute to obesity in SMS, a topic that awaits future investigation.

We found the metabolic phenotypes induced by deleting *Rai1* from BDNF-producing cells were

specifically due to *Rai1* deletion in PVH^{BDNF} neurons. Therefore, our study focused on PVH^{BDNF}

neurons. Electrophysiological experiments showed that most wild-type PVH^{BDNF} neurons display

high-frequency spontaneous AP firing (1-10 Hz) at the resting membrane potential. Most

PVH^{BDNF} neurons fire sharp APs held at a hyperpolarized potential and show repetitive

392 overshoot during the current ramp test. Loss of *Rai1* significantly reduced the intrinsic neuronal

393 excitability of PVH^{BDNF} neurons. Because GABAergic *Rai1* loss did not induce obesity (Huang

394 *et al.*, 2016) and most BDNF-producing neurons are glutamatergic (Canals *et al*, 2001). The

395 ionic changes in PVH^{BDNF} neurons that precisely explain these excitability changes await future

396 investigations. Different groups of PVH neurons show phasic bursting or slowly inactivating

397 delayed rectifier potassium conductance, which are properties mediated by different potassium

398 channels (Lee *et al.*, 2012). Given that our previous RNA-seq experiment found that

399 hypothalamic Rai1 loss induces misexpression of potassium channel subunits *Kcnc2* and *Kcnj11*

400 (Huang *et al.*, 2016), RAI1-dependent potassium channels are likely involved in altered neuronal

401 activity. Furthermore, we found fewer low threshold depolarization events in cKO PVH^{BDNF}

402 neurons during the hyperpolarized voltage step, suggesting a potential involvement of HCN

403 channels.

404 Our previous work found that *Rai1* loss in the dentate gyrus resulted in neuronal

405 hyperexcitability (Chang *et al*, 2022b). Here we demonstrated that *Rai1* loss in the hypothalamus

406 resulted in neuronal hypoactivity, suggesting that RAI1 regulates neuronal excitability in a cell

407 type-specific manner. Reduced PVH^{BDNF} neuronal activity upon *Rai1* loss could potentially

408 explain obesity, consistent with a previous report that found chemogenetic activation of

409 PVH^{BDNF} neurons promotes negative energy balance by decreasing feeding and increasing

410 energy expenditure (Wu & Xu, 2022).

411 It is plausible that RAI1 regulates the expression of genes encoding inward rectifier K$^+$ channels,

412 which regulate neuronal activity and potentially energy homeostasis. For instance, KIR6 (a

413 family of ATP-sensitive potassium channels, K$_{ATP}$) is widely expressed in the hypothalamus.

414 Deleting the hypothalamic KIR6.2 subunit impairs K$_{ATP}$ channel function and glucose tolerance

(Miki *et al*, 2001; Parton *et al*, 2007). Moreover, reduced expression of hypothalamic GIRK4

(encoding an inwardly rectifying potassium channel) causes obesity (Perry *et al*, 2008).

GABAergic neurotransmission from arcuate AGRP-expressing neurons to the PVH neurons is

important to increase appetite by favouring hyperphagia (Atasoy *et al*, 2012). Disrupting the

composition of these ion channels could contribute to reduced PVH^{BDNF} neuronal firing, which

awaits further investigations.

Female mice with a conditional knockout of *Rai1* from BDNF-producing neurons do not

display a noteworthy difference in food intake. Conversely, their male counterparts exhibit a

significant increase in food intake. Although SMS individuals of both genders tend to overeat,

male patients who are obese show significantly higher food consumption than their female

counterparts (Gandhi *et al.*, 2022). This observation raises the possibility that *Rai1* regulates

eating behaviours through multiple cell types in the hypothalamus and that a male-specific

involvement of BDNF-producing neurons in regulating food intake, potentially provides a

neurobiological basis for the observed pattern in SMS patients (Gandhi et al., 2022). In future

work, it would be interesting to dissect the sex-specific role of RAI1 in regulating the functional

development of PVH^{BDNF} neurons. In addition to a role for RAI1 in regulating neuronal activity,

RAI1 may regulate other aspects of PVH^{BDNF} neuronal function, including projections to other

brain regions to regulate energy homeostasis. In this regard, it would be interesting to dissect

how RAI1 controls the circuit connectivity of PVH^{BDNF} neurons and how *Rai1* loss in PVH^{BDNF}

neurons affects nearby PVH^{TRKB} neurons that receive BDNF signalling and are linked to body

weight regulation (An *et al.*, 2020).

At the molecular level, our RPPA data show that *Rai1* depletion disrupted multiple intracellular

signalling pathways and resulted in the hypoactivation of specific phospho-signalling proteins.

Multiple early studies support the notion that Rai1 directly promotes *Bdnf* expression. First,

hypothalamic *Bdnf* mRNA levels were reduced in SMS mice (Burns *et al.*, 2010; Javed *et al.*,

2021) or mice with *Rai1* conditional deletions (Huang *et al.*, 2016). Second, the RAI1 protein

directly binds to a promoter region of *Bdnf* (Huang *et al.*, 2016). Finally, when *Rai1* expression

was induced using CRISPR activation, *Bdnf* mRNA levels were significantly elevated (Chang *et*

al, 2022a).

To lend further support, here we show that SMS mice have reduced hypothalamic BDNF

protein levels, as well as reduced activation of a TRKB downstream target AKT. Reduced AKT

activity is frequently associated with insulin resistance and obesity in mice (Shao *et al*, 2000) and

children (Su *et al.*, 2021). We also found that obesity, increased HDL levels, and insulin

insensitivity in SMS mice could be partially alleviated using a small molecule drug (LM22A-4)

that promotes AKT phosphorylation. LM22A-4 penetrates the blood-brain barrier reasonably

well and has been shown to ameliorate symptoms of other neurological disorders associated with

BDNF dysfunction including Rett syndrome, Huntington's disease, Dravet syndrome, and

refractory neonatal seizures (Canals *et al.*, 2004; Chang *et al.*, 2006; Gu *et al.*, 2022; Kron *et al.*,

2012; Ogier *et al.*, 2007). We recognize that while several in vivo studies have demonstrated the

potential of LM22A-4 in targeting neurotrophin downstream signalling (Kron *et al*, 2014; Li *et*

al, 2017), an in vitro analysis failed to demonstrate the ability of LM22A-4 to activate TRKB

directly (Boltaev *et al*, 2017). Therefore, the precise mechanism by which LM22A-4 enhances

AKT cascades in the mammalian brain remains unclear and awaits further investigations. In the

hypothalamus of SMS mice, LM22A-4 could indirectly engage neurotrophin downstream PI3K-

AKT pathway through the G protein-coupled receptor-dependent transactivation of the TRKB

receptor (Domeniconi & Chao, 2010) or other unknown mechanisms. Moreover, while LM22A-

4 may have potential side effects, we found that wild-type mice treated with LM22A-4 did not

show a further decrease in body weight, suggesting limited side effects regarding body weight

regulation.

The partial rescue of SMS phenotypes that we observed contrasts with our previous findings that

found (1) AAV-mediated BDNF overexpression in PVH at 3 weeks of age and (2) genetically

overexpressing BDNF from early development could fully prevent the development of obesity in

SMS mice (Javed *et al.*, 2021). Therefore, further increasing TRKB signalling activation and

earlier intervention might improve therapeutic outcomes. Alternatively, our RPPA data showed

that in addition to disruption of neurotrophin signalling, multiple pathways including mTOR

were downregulated in SMS. Increasing hypothalamic mTOR activity has been shown to

decrease food intake and body weight (Cota *et al.*, 2006). Therefore, it is possible that enhancing

both mTOR and TRKB signalling could achieve a more robust rescue of obesity in SMS mice.

Nevertheless, these data support the concept that drugging TRKB signalling may improve

stereotypical repetitive behaviour and partially alleviate metabolic functions in a mouse model of

SMS. Delayed symptom onset could provide a window of opportunity for other interventions to

further modify disease progression, an area requiring further investigation.

This study has several limitations. First, whether the PVH[BDNF] neurons depend on Rai1 to

regulate adaptive thermogenesis awaits future analysis. It is known that a subset of posterior

PVH[BDNF] neurons project to the spinal cord and regulate thermogenesis (An *et al.*, 2015).

However, we found the weights of brown adipocytes were similar in control and cKO mice.

Moreover, SMS mice have normal respiratory exchange rates and there are currently no reports

of defective adaptive thermogenesis in SMS patients. This suggests that adaptive thermogenesis

is not impacted in SMS. Another limitation is that we did not test different doses, durations, and

time points for LM22A-4 treatment. The precise mechanisms by which LM22A-4 improves

body weight, metabolic function, and repetitive behaviours remain unknown. Nevertheless, our

work identifies RAI1 as a novel regulator of the neuronal excitability of PVH[BDNF] neurons. This

regulation is critical for regulating food intake and energy expenditure. Our evidence suggests

that BDNF-producing neurons are involved in SMS pathogenesis and that enhancing BDNF-

TRKB signalling should be explored as a future therapeutic avenue for obesity and metabolic

conditions associated with reduced neurotrophin signalling (Xu & Xie, 2016).

Materials and Methods

Animal Management

The animal protocol (MUHC-8127) was approved by the animal care committee of the Montreal

General Hospital and was performed according to the guidelines of the Canadian Council on

Animal Care. Mice were group-housed in 12-hour light/12-hour dark cycles with an *ad libitum*

supply of regular chow and water. The exception was in the fasting studies when the animals

were food deprived for 5 hours and in CALMS and food intake studies when animals were

single-housed in a 12-hour light/12-hour dark cycle with an *ad libitum* supply of regular chow

and water. All mice were maintained in a C57Bl/6J background. $Bdnf^{Cre/Cre}$ (RRID:

IMSR_JAX:030189), $Rosa26^{Ai9/Ai9}$ (RRID: IMSR_JAX:007909), and $Actb^{Cre/Cre}$ (RRID:

IMSR_JAX:019099) mice were purchased from the Jackson laboratory. The $Rai1^{fl/fl}$ (RRID:

IMSR_JAX:029103) strain was previously generated and characterized by us (Huang *et al.*,

2016). SMS mice ($Rai1^{+/-}$: whole-body germline $Rai1$ heterozygous knockout that genetically

mimics SMS) were generated by crossing germline $Actb^{Cre/Cre}$ mice with $Rai1^{fl/+}$. To generate

conditional knockout mice, $Bdnf^{Cre/+}$; $Rai1^{fl/+}$ mice were crossed with $Rai1^{fl/+}$ to create control

and experimental groups from the same littermate. 7 mice with malocclusion or lesions were

excluded from the study: 4 with malocclusion (2 females and 2 males) and 3 with lesions (3

females).

Metabolic profiling

Metabolic profiling was conducted in three experimental conditions: male and female

conditional knockout mice, female virus-injected $Bdnf^{Cre/+}$ mice, and female drug- or saline-

injected SMS and control mice. Metabolic profiling included body weight and body composition

analysis, food intake measurement, indirect calorimetry for locomotion activity, energy

expenditure (EE), respiratory exchange rate (RER), measurement of serum lipids and leptin

levels, adipose tissue histology analysis, and glucose and insulin tolerance tests. All experiments

were performed as previously described (Javed *et al.*, 2021). A brief description is detailed

below.

Body weight and body composition analysis

The body weight of animals was measured weekly from week 3 (weaning age) onward. After the

25^{th} week, mice were subjected to body composition analysis. Body composition, including total

fat and lean mass, was assessed using a nuclear EchoMRI whole-body analyzer. Toward the end

of the experiments, animals were euthanized to collect brown adipose, visceral, subcutaneous

(inguinal) and ependymal white adipose tissues and weighed using an analytical scale

(Sartorius).

Food Intake Measurement

Mice were single-housed for food intake measurement. Food intake was measured over a one-

week period during which food was weighed prior to placement in the cage with minimal

bedding. Each morning, the remaining food inside the cage was weighed. The amount of total

food consumed was averaged over 7 days.

Indirect Calorimetry

A Comprehensive Lab Animal Monitoring System (CLAM) (Columbus instruments) was used to

measure the RER and EE (normalized over lean mass). Animals were singled-housed in the

CLAMS apparatus at 21°C (70°F) in a light–dark cycle matching their housing conditions for 24

hours (acclimation), followed by 48 hours of measurement.

Serum lipid level determination

A fluorescence-based assay kit (Abcam ab65390) was used to measure serum lipid levels. The

manufacturer's instructions were followed except that we used half-area black 96-well

microplates. Fluorescence measurements at 535/587 nm (Ex/Em) were recorded with the Ensight

instrument from PerkinElmer. A new standard curve ranging from 0 to 10 μg/ml was produced

for each microplate to accurately determine lipid levels in serum samples. Standards and samples

were loaded and analyzed twice.

Histology of adipose tissue

Ependymal white adipose tissues (WAT) were excised immediately after the animals were

euthanized and fixed with 10% formalin in PBS. The fixed tissues were dehydrated, embedded in

paraffin, and sectioned. Two 5-micrometre thick sections were taken from each animal and

stained with hematoxylin and eosin. Images of sections were then captured using an Aperio

ScanScope CS slide scanner at 20X optical magnification, and the resulting images were saved

in jpeg format using the ImageScope software. The images were manually cleaned to remove

non-adipose tissues such as lymph nodes. An automated method using the Adiposoft plugin

version 1.16 was utilized to determine adipocyte size and number from the jpeg images in Fiji

version 2.1.0/1.53c. The Adiposoft settings used excluded cells on edges and set a minimum and

maximum cell diameter of 20 and 300 μm, respectively. An experimental value of 2.49 μm per

pixel was determined using the Aperio scale bar and used for Adiposoft. To ensure accuracy, two

images from each animal were analyzed blindly, examining more than 1600 cells per mouse and

more than 9100 cells per condition. Adipocyte size frequencies were calculated using the

frequency function in Excel.

Glucose and insulin tolerance tests

For the glucose tolerance test, experimental mice were food deprived for 5 hours with *ad libitum*

water supply. A bolus of glucose (1.5g/kg of lean body weight) was then administered

intraperitoneally and glycemia was measured from tail vein blood samples using the Accu-chek

Performa glucometer at T0 (before the injection) as well as at 15, 30, 45, 60, 90 and 120 min

(after the injection). Tail vein blood samples were collected via a capillary for insulin assays at

T0, 15 and 30 min. For insulin tolerance tests, experimental mice were food-deprived for 5 hours

with *ad libitum* access to water. A bolus of insulin (1 U/kg of lean body weight) was

administered via an IP injection and glycemia was measured from blood sampled at the tail vein

using an Accu-chek Performa glucometer at T0 (before the injection), as well as at 15, 30, 45, 60,

90 and 120 min (after the injection). Tail vein blood samples were collected via a capillary for

insulin assays at T0.

Stereotaxic Injections

Bdnf^{Cre/+} mice were anesthetized using isoflurane, and the ears were barred onto a stereotaxic

device (David Kopf Instruments). Adeno-associated viruses (AAVs, volume = 250 nl) were

bilaterally injected into the PVH using the following coordinates relative to the Bregma: -0.6 mm

anteroposterior (AP), ±0.35 mm mediolateral (ML) and -5.52 mm dorsoventral (DV). A

description of the AAV constructs and titre used is detailed below:

AAV9-CMV7-DIO-saCas9 (Viral title: >1 × 10^13), Vector Biolab

AAV9-CMV7-sgRai1-GFP (Viral title: >1 ×10^9), Applied Biological Materials Inc. Three

sgRNAs were pooled to target Rai1 (sgRNA1: TTCCTCGCCAGAGTAGCGCC, sgRNA2:

CCCAGCCTCATGATAGGCCG, sgRNA3: CCCAGCCTCATGATAGGCCG).

Immunostaining of mouse brain tissues

Immunostaining experiments were performed as previously described (Huang *et al*, 2012).

Briefly, mice were anesthetized and perfused with 1X PBS and 4% formaldehyde, and the brains

were dissected and fixed in 4% formaldehyde overnight. Fixed brains were transferred to 30%

sucrose solution and subsequently embedded in the OCT solution. The brains were then

sectioned (40µm), washed in PBS, permeabilized with 0.5% Triton X-100, and incubated with

10% normal donkey serum in PBS for 2 hours. The sections were then incubated overnight with

primary antibodies (Rai1: 1:500 dilution, RRID:AB_2921229 (Huang *et al.*, 2016); BDNF:

1:500, RRID:AB_10862052, Abcam ab108319; p-TRKB: 1:500 dilution, RRID:AB_2721199,

Millipore ABN1381; oxytocin: 1:1000 dilution, RRID:AB_11212999, Millipore MAB5296) at

4°C on a shaker. The following day, sections were washed with 1X PBS twice for 10 minutes

and incubated with a secondary antibody (1:1000 dilution) for 3 hours at room temperature.

Sections were washed with 1X PBS once before mounting on glass slides with the DAPI-

mounting (DAPI-Fluoromount-G, SouthernBiotech) and then imaged using a confocal

microscope (Olympus FV-1000 confocal laser scanning microscope) with 20x or 40x objectives

(NA 0.85 and 1.3, respectively). Z-stacks were taken with a step size of 1.0 μm with a 1024 x

1024 resolution. Images were acquired as a stack with at least 13 optical sections. Image analysis

was performed using the Fiji software.

Real-Time Quantitative Reverse Transcription PCR (qRT-PCR)

qRT-PCR was performed using mRNAs isolated from hypothalamic tissue samples. After the

isolation of total RNA, mRNAs were reverse-transcribed with the SuperScript III First-Strand

Synthesis System (Thermo-Fisher). Quantitative PCR reactions were conducted using SsoFast

EvaGreen Supermix on a Bio-Rad qPCR system.

Enzyme-linked immunosorbent assay (ELISA)

ELISA was performed using hypothalamic tissues from 7-week-old SMS mice to assess the

mature BDNF protein levels using a commercially available BDNF ELISA kit (Mature BDNF

Rapid ELISA Kit: Human, Mouse, Rat).

Western blotting analysis

Mice were quickly decapitated, brains were rapidly removed, and hypothalamic tissues were

collected, weighed, and snap-frozen with liquid nitrogen. Hypothalamic tissues were lysed using

a tissue lysis buffer containing RIPA buffer (5M NaCl, 1M PH 8.0 Tris HCl, 1% NP.40, 10%

$C_{24}H_{39}NaO_4$, and 20% SDS) and the Halt™ Protease and Phosphatase Inhibitor Cocktail

(Thermo-Fisher). Lysates were first sonicated on ice for 15 seconds and centrifuged at 10000

RPM at 4°C for 10 minutes. The supernatant was collected in a separate Eppendorf tube, and a

BCA assay was performed to quantify protein levels. Proteins were then separated using SDS-

PAGE gel and transferred onto nitrocellulose membranes. To determine the phosphorylation

levels of AKT, membranes were cut and blocked in 5% milk in tris-buffered saline (TBS) for 1

hour and then incubated overnight at 4°C with anti-p-AKT (Ser473) antibody (1:1000 dilution,

RRID: AB_2315049, Cell Signaling Technology 4060T) and anti-H3 antibody (internal control:

1:1000 dilution, RRID: AB_302613, Abcam ab1791). The next day, membranes were washed

with TBST three times for five minutes before incubating with an HRP-conjugated secondary

antibody (1:10000 dilution, Thermo Fisher 200-403-095S). Membranes were later stripped by

incubating in 0.5M NaOH for 10 min and then re-probed for total Akt levels by incubating with

anti-Akt (C67E7) antibody (1:1000 dilution, Cell Signaling Technology 4691T). The density of

each immunoreactive phospho-protein band was expressed as a fraction of the band for the total

AKT protein in the same lane.

Reverse-phase protein array (RPPA)

A reverse-phase protein array (RPPA) was used for precise high-throughput quantification of

total and phospho-protein levels in different signalling pathways. Protein lysates from

hypothalamic tissues were extracted and transferred to 384-well microarray plates to print

individual slides. The slides were then labelled with validated antibodies recognizing total

protein and specific phosphorylation sites. SYPRO Ruby staining was used to measure total

protein levels. The data obtained from RPPA were normalized for individual samples. Each

antibody's raw signal intensity (SI) was determined by subtracting the background from the

antibody-specific signals during image analysis. The normalized antibody SI was expressed

using the formula $N = A - C/T \times M$, where N is the normalized SI, A is the raw antibody SI, C is

the negative control SI, T is the SI of total protein, and M is the median SI of total protein from a

spot within the same sample group. Antibodies with SI less than 200 were filtered out, and

quality control (QC) scores were established to discard poor-quality reactions. The coefficient of

variation (CV) score was determined for each antibody and defined as the percentage of samples

on a slide with a $CV < 20\%$. Antibodies with a median technical triplicate $CV > 20\%$ were also

excluded from further analysis. T-tests were performed for each antibody to identify

differentially expressed proteins in the SMS compared to control groups. PCA plots were

generated for all samples, annotated with their experimental group, and hierarchical clustering

was performed using a Python analysis tool. The vertical axis of the dendrogram represents the

dissimilarity (measured as distance) between protein expressions, and the horizontal axis

represents the individual test samples. The colour code (ranging from red to yellow to green)

specifies the expression levels of different proteins, where red indicates low expression, yellow

indicates intermediate expression, and green indicates high expression. Enrichment analysis on

the differentially expressed proteins was performed using an online data resource string: version

11.5.

Electrophysiology of acute brain slices

Mouse brains were removed and placed in ice-cold carbonated slicing artificial cerebrospinal

fluid (aCSF), which contained (in mM) 124 NaCl, 3 KCl, 1.4 NaH_2PO_4, 26 $NaHCO_3$, 1.3 Mg

SO4, 10 D-glucose, 0.5 CaCl$_2$, and 3.5 MgCl$_2$. Coronal sections (300 µm) were cut on a

vibratome. Slices were allowed to recover at 32 °C for 40 minutes and then at 30 °C for

30 minutes to 6 hours in the recording medium, carbonated aCSF (124 NaCl, 3 KCl, 1.4

NaH$_2$PO$_4$, 26 NaHCO$_3$,1.3 Mg SO4, 10 D-glucose, 2.4 CaCl$_2$, in mM, ~300 mOsm). Signals

were recorded with a 5× gain, low-pass filtered at 2 kHz, digitized at 10 kHz (Molecular

Devices Multiclamp 700B) and analyzed with pClamp 11 (Molecular Devices). Whole-cell

recordings were made using 3–5 MΩ pipettes filled with an internal solution that contained

(in mM) 131 potassium gluconate, 8 NaCl, 20 KCl, 2 EGTA, 10 HEPES, 2 MgATP and 0.3

Na3GTP (pH 7.2, with KOH, 300–310 mOsm with sucrose). For miniature current recording,

the internal solution contained (in mM) 130 gluconate acid, 8 CsCl, 1 NaCl, 2 EGTA, 0.8

CsOH, 10 HEPES, 2 MgATP and 0.3 Na3GTP (pH 7.2, with CsOH, 300–310 mOsm adjusted

with sucrose). mEPSCs were recorded in the presence of 1 µM TTX at a holding potential of -

70 mV. mIPSCs were recorded at a holding potential of +10 mV. The mIPSCs were analyzed

from events during a 2-minute continuous recording that contained > 15 events/neuron.

Additional calculations, statistical analysis, and plotting were performed in Microsoft Excel or

Prism (GraphPad).

Spectrometric analysis of LM22A-4 levels in the brain

We assessed the efficacy of LM22A-4, a small molecule pharmacological agent, in 8-week-old

SMS and control mice. LM22A-4, chemical name N1, N3, N5tris (2hydroxyethyl) 1, 3, 5

benzenetricarboxamide (Molecular formula: C$_{15}$H$_{21}$N$_3$O$_6$, Formula weight: 339.3), was

manufactured by Cayman Chemical with purity specification >98%. A stock solution was made

by first dissolving LM22A-4 in DMSO (solubility =30mg/ml) and then diluting it with 0.9%

isotonic sterilized saline solution (Sigma). To maximize brain concentrations, LM22A-4 was

administered both intranasally (5mg/kg) and intraperitoneally (50mg/kg), as previously reported

(Simmons *et al*, 2013). The bioavailability of this drug in the hypothalamus using these doses

and routes of administration was assessed with the reverse phase LC triple-quadrupole tandem

mass spectroscopic detection (LC-MS/MS) by Absorption Systems. Mice were euthanized, and

hypothalamic tissue was collected at either 1 h or 3 h post-injection (n= 4/timepoint). Known

amounts of LM22A-4 were added to the blank sample to generate a standard curve to verify test

accuracy.

LM22A-4 treatment

Mice were divided into four groups: Saline-injected control, Saline-injected SMS, LM22A-4-

injected SMS, and LM22A-4-injected control mice. Mice were injected daily, intra-nasally

(5mg/kg) and intraperitoneally (50mg/kg), for 7 weeks. Body weight was measured between 8

and 14 weeks of age. Food intake was measured from 14 to 15 weeks of age. After 15 weeks of

age, mice were euthanized to collect brown adipose, visceral, subcutaneous (inguinal) and

ependymal white adipose tissues and weighed using an analytical scale (Sartorius).

Hypothalamic brain tissues were collected for western blot analysis. Metabolic profiling was

performed on a separate cohort of mice injected for two weeks prior to the start of experiments

as well as at specific weeks during the experiment.

Vertical rearing

To measure vertical repetitive rearing, mice were taken into the testing room and allowed to

habituate for 1 h prior to testing. During the test, mice were placed in the center of a 45 cm (W)

× 45 cm (D) × 40 cm (H) square arena, and their movement was recorded for 10 minutes.

Statistical analysis

The sample sizes used were based on our previous studies (Huang *et al.*, 2016; Javed *et al.*,

2021). The data were subjected to statistical analysis for significance by using GraphPad Prism

0.9. Significance was defined as having a p-value < 0.05. The significance levels were as

follows: * p< 0.05, **p<0.01, ***p<0.0001, **** p<0.0001. The statistical tests used for each

analysis (e.g., Student's t-test, analysis of variance [ANOVA]) are indicated in the text and

legends.

Data Availability

726   This study includes no data deposited in external repositories.

Acknowledgements

729   Sehrish Javed was supported by a Fonds de Recherche du Québec - Santé (FRQS) doctoral

730   award. This work was supported by the Rare Diseases: Models & Mechanisms Network and the

731   Smith-Magenis syndrome Research Foundation.

References

Abad C, Cook MM, Cao L, Jones JR, Rao NR, Dukes-Rimsky L, Pauly R, Skinner C, Wang Y, Luo F *et al* (2018) A Rare De Novo RAI1 Gene Mutation Affecting BDNF-Enhancer-Driven Transcription Activity Associated with Autism and Atypical Smith-Magenis Syndrome Presentation. *Biology (Basel)* 7

Alaimo JT, Barton LV, Mullegama SV, Wills RD, Foster RH, Elsea SH (2015) Individuals with Smith-Magenis syndrome display profound neurodevelopmental behavioral deficiencies and exhibit food-related behaviors equivalent to Prader-Willi syndrome. *Res Dev Disabil* 47: 27-38

An JJ, Kinney CE, Tan JW, Liao GY, Kremer EJ, Xu B (2020) TrkB-expressing paraventricular hypothalamic neurons suppress appetite through multiple neurocircuits. *Nat Commun* 11: 1729

An JJ, Liao GY, Kinney CE, Sahibzada N, Xu B (2015) Discrete BDNF Neurons in the Paraventricular Hypothalamus Control Feeding and Energy Expenditure. *Cell Metab* 22: 175-188

Atasoy D, Betley JN, Su HH, Sternson SM (2012) Deconstruction of a neural circuit for hunger. *Nature* 488: 172-177

Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, Ferreira M, Tang V, McGovern RA, Kenny CD *et al* (2005) Divergence of melanocortin pathways in the control of food intake and energy expenditure. *Cell* 123: 493-505

Boltaev U, Meyer Y, Tolibzoda F, Jacques T, Gassaway M, Xu Q, Wagner F, Zhang YL, Palmer M, Holson E *et al* (2017) Multiplex quantitative assays indicate a need for reevaluating reported small-molecule TrkB agonists. *Sci Signal* 10

Boot E, Linders CC, Tromp SH, van den Boogaard MJ, van Eeghen AM (2021) Possible underreporting of pathogenic variants in RAI1 causing Smith-Magenis syndrome. *Am J Med Genet A* 185: 3167-3169

Burke LK, Darwish T, Cavanaugh AR, Virtue S, Roth E, Morro J, Liu SM, Xia J, Dalley JW, Burling K *et al* (2017) mTORC1 in AGRP neurons integrates exteroceptive and interoceptive food-related cues in the modulation of adaptive energy expenditure in mice. *Elife* 6

Burns B, Schmidt K, Williams SR, Kim S, Girirajan S, Elsea SH (2010) Rai1 haploinsufficiency causes reduced Bdnf expression resulting in hyperphagia, obesity and altered fat distribution in mice and humans with no evidence of metabolic syndrome. *Hum Mol Genet* 19: 4026-4042

Canals JM, Checa N, Marco S, Akerud P, Michels A, Perez-Navarro E, Tolosa E, Arenas E, Alberch J (2001) Expression of brain-derived neurotrophic factor in cortical neurons is regulated by striatal target area. *J Neurosci* 21: 117-124

Canals JM, Pineda JR, Torres-Peraza JF, Bosch M, Martin-Ibanez R, Munoz MT, Mengod G, Ernfors P, Alberch J (2004) Brain-derived neurotrophic factor regulates the onset and severity of motor dysfunction associated with enkephalinergic neuronal degeneration in Huntington's disease. *J Neurosci* 24: 7727-7739

Caron A, Labbe SM, Lanfray D, Blanchard PG, Villot R, Roy C, Sabatini DM, Richard D, Laplante M (2016) Mediobasal hypothalamic overexpression of DEPTOR protects against high-fat diet-induced obesity. *Mol Metab* 5: 102-112

Chang HC, Lee YJ, Javed S, Haque M, Chang YT, Lin YC, Oram C, Huang WH (2022a) rAAV-CRISPRa therapy corrects Rai1 haploinsufficiency and rescues selective disease features in Smith-Magenis syndrome mice. *J Biol Chem*: 102728

Chang Q, Khare G, Dani V, Nelson S, Jaenisch R (2006) The disease progression of Mecp2 mutant mice is affected by the level of BDNF expression. *Neuron* 49: 341-348

Chang YT, Kowalczyk M, Fogerson PM, Lee YJ, Haque M, Adams EL, Wang DC, DeNardo LA, Tessier-Lavigne M, Huguenard JR *et al* (2022b) Loss of Rai1 enhances hippocampal excitability and epileptogenesis in mouse models of Smith-Magenis syndrome. *Proc Natl Acad Sci U S A* 119: e2210122119

Collaboration NCDRF (2017) Worldwide trends in body-mass index, underweight, overweight, and
obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9
million children, adolescents, and adults. *Lancet* 390: 2627-2642

Colmers PLW, Bains JS (2018) Balancing tonic and phasic inhibition in hypothalamic corticotropin-
releasing hormone neurons. *J Physiol* 596: 1919-1929

Cota D, Proulx K, Smith KA, Kozma SC, Thomas G, Woods SC, Seeley RJ (2006) Hypothalamic mTOR
signaling regulates food intake. *Science* 312: 927-930

Domeniconi M, Chao MV (2010) Transactivation of Trk receptors in spinal motor neurons. *Histol
Histopathol* 25: 1207-1213

Edelman EA, Girirajan S, Finucane B, Patel PI, Lupski JR, Smith AC, Elsea SH (2007) Gender, genotype, and
phenotype differences in Smith-Magenis syndrome: a meta-analysis of 105 cases. *Clin Genet* 71: 540-550

Gandhi AA, Wilson TA, Sisley S, Elsea SH, Foster RH (2022) Relationships between food-related behaviors,
obesity, and medication use in individuals with Smith-Magenis syndrome. *Res Dev Disabil* 127: 104257

Gao H, Molinas AJR, Miyata K, Qiao X, Zsombok A (2017) Overactivity of Liver-Related Neurons in the
Paraventricular Nucleus of the Hypothalamus: Electrophysiological Findings in db/db Mice. *J Neurosci* 37:
11140-11150

Garcia AD, Doan NB, Imura T, Bush TG, Sofroniew MV (2004) GFAP-expressing progenitors are the
principal source of constitutive neurogenesis in adult mouse forebrain. *Nat Neurosci* 7: 1233-1241

Gorski JA, Talley T, Qiu M, Puelles L, Rubenstein JL, Jones KR (2002) Cortical excitatory neurons and glia,
but not GABAergic neurons, are produced in the Emx1-expressing lineage. *J Neurosci* 22: 6309-6314

Gray J, Yeo GS, Cox JJ, Morton J, Adlam AL, Keogh JM, Yanovski JA, El Gharbawy A, Han JC, Tung YC *et al*
(2006) Hyperphagia, severe obesity, impaired cognitive function, and hyperactivity associated with
functional loss of one copy of the brain-derived neurotrophic factor (BDNF) gene. *Diabetes* 55: 3366-
3371

Gu F, Parada I, Yang T, Longo FM, Prince DA (2022) Chronic partial TrkB activation reduces seizures and
mortality in a mouse model of Dravet syndrome. *Proc Natl Acad Sci U S A* 119

Huang WH, Guenthner CJ, Xu J, Nguyen T, Schwarz LA, Wilkinson AW, Gozani O, Chang HY, Shamloo M,
Luo L (2016) Molecular and Neural Functions of Rai1, the Causal Gene for Smith-Magenis Syndrome.
Neuron 92: 392-406

Huang WH, Tupal S, Huang TW, Ward CS, Neul JL, Klisch TJ, Gray PA, Zoghbi HY (2012) Atoh1 governs the
migration of postmitotic neurons that shape respiratory effectiveness at birth and chemoresponsiveness
in adulthood. *Neuron* 75: 799-809

Huang WH, Wang DC, Allen WE, Klope M, Hu H, Shamloo M, Luo L (2018) Early adolescent Rai1
reactivation reverses transcriptional and social interaction deficits in a mouse model of Smith-Magenis
syndrome. *Proc Natl Acad Sci U S A*

Javed S, Lee YJ, Xu J, Huang WH (2021) Temporal dissection of Rai1 function reveals brain-derived
neurotrophic factor as a potential therapeutic target for Smith-Magenis syndrome. *Hum Mol Genet* 31:
275-288

Javed S, Selliah T, Lee YJ, Huang WH (2020) Dosage-sensitive genes in autism spectrum disorders: From
neurobiology to therapy. *Neurosci Biobehav Rev* 118: 538-567

Kamitakahara A, Xu B, Simerly R (2016) Ventromedial hypothalamic expression of Bdnf is required to
establish normal patterns of afferent GABAergic connectivity and responses to hypoglycemia. *Mol Metab*
5: 91-101

Kocalis HE, Hagan SL, George L, Turney MK, Siuta MA, Laryea GN, Morris LC, Muglia LJ, Printz RL,
Stanwood GD *et al* (2014) Rictor/mTORC2 facilitates central regulation of energy and glucose
homeostasis. *Mol Metab* 3: 394-407

Kron M, Howell CJ, Adams IT, Ransbottom M, Christian D, Ogier M, Katz DM (2012) Brain activity
mapping in Mecp2 mutant mice reveals functional deficits in forebrain circuits, including key nodes in
the default mode network, that are reversed with ketamine treatment. *J Neurosci* 32: 13860-13872
Kron M, Lang M, Adams IT, Sceniak M, Longo F, Katz DM (2014) A BDNF loop-domain mimetic acutely
reverses spontaneous apneas and respiratory abnormalities during behavioral arousal in a mouse model
of Rett syndrome. *Dis Model Mech* 7: 1047-1055
Lee SK, Lee S, Shin SY, Ryu PD, Lee SY (2012) Single cell analysis of voltage-gated potassium channels that
determines neuronal types of rat hypothalamic paraventricular nucleus neurons. *Neuroscience* 205: 49-
62
Li W, Bellot-Saez A, Phillips ML, Yang T, Longo FM, Pozzo-Miller L (2017) A small-molecule TrkB ligand
restores hippocampal synaptic plasticity and object location memory in Rett syndrome mice. *Dis Model*
Mech 10: 837-845
Loos RJF, Yeo GSH (2022) The genetics of obesity: from discovery to biology. *Nat Rev Genet* 23: 120-133
Massa SM, Yang T, Xie Y, Shi J, Bilgen M, Joyce JN, Nehama D, Rajadas J, Longo FM (2010) Small molecule
BDNF mimetics activate TrkB signaling and prevent neuronal degeneration in rodents. *J Clin Invest* 120:
1774-1785
Miki T, Liss B, Minami K, Shiuchi T, Saraya A, Kashima Y, Horiuchi M, Ashcroft F, Minokoshi Y, Roeper J *et al*
(2001) ATP-sensitive K+ channels in the hypothalamus are essential for the maintenance of glucose
homeostasis. *Nat Neurosci* 4: 507-512
Moss J, Oliver C, Arron K, Burbidge C, Berg K (2009) The prevalence and phenomenology of repetitive
behavior in genetic syndromes. *J Autism Dev Disord* 39: 572-588
Ogier M, Wang H, Hong E, Wang Q, Greenberg ME, Katz DM (2007) Brain-derived neurotrophic factor
expression and respiratory function improve after ampakine treatment in a mouse model of Rett
syndrome. *J Neurosci* 27: 10912-10917
Parton LE, Ye CP, Coppari R, Enriori PJ, Choi B, Zhang CY, Xu C, Vianna CR, Balthasar N, Lee CE *et al* (2007)
Glucose sensing by POMC neurons regulates glucose homeostasis and is impaired in obesity. *Nature* 449:
228-232
Perry CA, Pravetoni M, Teske JA, Aguado C, Erickson DJ, Medrano JF, Lujan R, Kotz CM, Wickman K (2008)
Predisposition to late-onset obesity in GIRK4 knockout mice. *Proc Natl Acad Sci U S A* 105: 8148-8153
Quarta C, Claret M, Zeltser LM, Williams KW, Yeo GSH, Tschop MH, Diano S, Bruning JC, Cota D (2021)
POMC neuronal heterogeneity in energy balance and beyond: an integrated view. *Nat Metab* 3: 299-308
Rao NR, Abad C, Perez IC, Srivastava AK, Young JI, Walz K (2017) Rai1 Haploinsufficiency Is Associated
with Social Abnormalities in Mice. *Biology (Basel)* 6
Schultze SM, Hemmings BA, Niessen M, Tschopp O (2012) PI3K/AKT, MAPK and AMPK signalling: protein
kinases in glucose homeostasis. *Expert Rev Mol Med* 14: e1
Shao J, Yamashita H, Qiao L, Friedman JE (2000) Decreased Akt kinase activity and insulin resistance in
C57BL/KsJ-Leprdb/db mice. *J Endocrinol* 167: 107-115
Simmons DA, Belichenko NP, Yang T, Condon C, Monbureau M, Shamloo M, Jing D, Massa SM, Longo FM
(2013) A small molecule TrkB ligand reduces motor impairment and neuropathology in R6/2 and BACHD
mouse models of Huntington's disease. *J Neurosci* 33: 18712-18727
Slager RE, Newton TL, Vlangos CN, Finucane B, Elsea SH (2003) Mutations in RAI1 associated with Smith-
Magenis syndrome. *Nat Genet* 33: 466-468
Smith AC, Magenis RE, Elsea SH (2005) Overview of Smith-Magenis syndrome. *J Assoc Genet Technol* 31:
163-167
Smith AC, McGavran L, Robinson J, Waldstein G, Macfarlane J, Zonona J, Reiss J, Lahr M, Allen L, Magenis
E (1986) Interstitial deletion of (17)(p11.2p11.2) in nine patients. *Am J Med Genet* 24: 393-414

Smith MA, Katsouri L, Irvine EE, Hankir MK, Pedroni SM, Voshol PJ, Gordon MW, Choudhury AI, Woods A,
Vidal-Puig A *et al* (2015) Ribosomal S6K1 in POMC and AgRP Neurons Regulates Glucose Homeostasis
but Not Feeding Behavior in Mice. *Cell Rep* 11: 335-343
Su X, Gu D, Xu L, Liang Z, Luo X, Yang P, Yang J (2021) PI3K/Akt pathway expression in children with
different obesity degrees and its relationship with glucolipid metabolism and insulin resistance. *Am J*
Transl Res 13: 6592-6598
Tan CL, Cooke EK, Leib DE, Lin YC, Daly GE, Zimmerman CA, Knight ZA (2016) Warm-Sensitive Neurons
that Control Body Temperature. *Cell* 167: 47-59 e15
Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsiani D, Kvitsani D, Fu Y, Lu J *et al* (2011) A resource
of Cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex. *Neuron* 71: 995-1013
Unger TJ, Calderon GA, Bradley LC, Sena-Esteves M, Rios M (2007) Selective deletion of Bdnf in the
ventromedial and dorsomedial hypothalamus of adult mice results in hyperphagic behavior and obesity.
J Neurosci 27: 14265-14274
Vohra MS, Benchoula K, Serpell CJ, Hwa WE (2022) AgRP/NPY and POMC neurons in the arcuate nucleus
and their potential role in treatment of obesity. *Eur J Pharmacol* 915: 174611
Wu SW, Xu B (2022) Rapid and Lasting Effects of Activating BDNF-Expressing PVH Neurons on Energy
Balance. *eNeuro* 9
Xu B, Goulding EH, Zang K, Cepoi D, Cone RD, Jones KR, Tecott LH, Reichardt LF (2003) Brain-derived
neurotrophic factor regulates energy balance downstream of melanocortin-4 receptor. *Nat Neurosci* 6:
736-742
Xu B, Xie X (2016) Neurotrophic factor control of satiety and body weight. *Nat Rev Neurosci* 17: 282-292
Xu Y, Lu Y, Cassidy RM, Mangieri LR, Zhu C, Huang X, Jiang Z, Justice NJ, Xu Y, Arenkiel BR *et al* (2019)
Identification of a neurocircuit underlying regulation of feeding by stress-related emotional responses.
Nat Commun 10: 3446
Yang H, An JJ, Sun C, Xu B (2016) Regulation of Energy Balance via BDNF Expressed in Nonparaventricular
Hypothalamic Neurons. *Mol Endocrinol* 30: 494-503
Yeo GS, Connie Hung CC, Rochford J, Keogh J, Gray J, Sivaramakrishnan S, O'Rahilly S, Farooqi IS (2004) A
de novo mutation affecting human TrkB associated with severe obesity and developmental delay. *Nat*
Neurosci 7: 1187-1189

Figures and Figure legends

Figure 1: *Rai1* haploinsufficiency disrupts hypothalamic proteomic profile in mice

(A) ELISA showing that BDNF protein levels in hypothalamic tissues were significantly reduced in SMS mice compared to controls (n=4/genotype). *p<0.05, unpaired t-test, two-tailed.

(B) A schematic diagram showing the experimental strategy. Hypothalamic tissue lysates from 7-week-old control (n = 5, technical triplicates/sample) and SMS (n=8, technical triplicates/sample) mice were subjected to RPPA that utilizes a cocktail of 240 validated antibodies to probe several key intracellular signalling pathways by quantifying the expression levels of total and phosphorylated proteins

(C) Principal component analysis segregates the SMS and Ctrl groups where PC1 explains 47% of the variance, and PC2 explains 25% of the variance.

(D) Heat map showing hierarchical clustering of differentially expressed phospho-proteins: most showed reduced levels in SMS samples.

(E) Protein-protein interaction (PPI) analysis showing differentially expressed proteins and phospho-proteins in a crosstalk network. PPI enrichment p-value: < 1.0e-16. The nodes represent the proteins/phospho-proteins and the lines indicate physical interactions. Associations with specific molecular pathways are shown as color-coded outer rings.

(F-G) The protein levels of two neurotrophin pathway-related nodes, PIK3CA (**F**) and phospho-NF-kB (**G**), were significantly reduced in SMS samples. ****p<0.0001, unpaired t-test, two-tailed

Figure 2: RAI1 is required in BDNF-producing cells to regulate energy metabolism

(A) A schematic diagram showing selective deletion of *Rai1* from BDNF-producing cells in mice.

(B) Representative images showing that in Ctrl mice (*Bdnf*$^{Cre/+}$; *Ai9*), many PVH^{BDNF} neurons (magenta) express RAI1 (cyan, double-positive cells were indicated with yellow arrows). By

contrast, PVH^{BDNF} neurons lack RAI1 expression in the cKO group (*Bdnf$^{Cre/+}$; Rai1$^{fl/fl}$; Ai9*).

Scale bars = 20µm.

(C) Percentage of PVH^{BDNF} neurons co-expressing RAI1 in Ctrl (n=3) and cKO (n=3) mice.

unpaired t-test, two-tailed, ****$p < 0.0001$.

(D) Female cKO mice (n = 12) showed a significant weight gain when compared to female Ctrl

mice (n = 10). Two-way ANOVA with Šidák's multiple comparisons test. *$p < 0.05$, ***$p < 0.001$.

(E) Body composition was measured with Echo-MRI, showing an increased fat mass in 26-

week-old mice. ns indicates the difference is not significant. Two-way ANOVA with Šidák's

multiple comparisons test. ****$p < 0.0001$.

(F) Fat mass of brown adipocytes (BAT), subcutaneous inguinal (S.C.ing) and epididymal white

adipose tissue (eWAT) in Ctrl and cKO mice. ns indicates the difference is not significant. Two-

way ANOVA with Šidák's multiple comparisons test. **$p < 0.01$, ****$p < 0.0001$.

(G) Representative images showing eWAT adipocyte hypertrophy of the cKO mice (left). Scale

bar = 500µm. Frequency distribution of adipocytes at each cellular size (right). Two-way

ANOVA with Šidák's multiple comparisons test. ***$p < 0.001$, ****$p < 0.0001$.

(H) Female cKO mice showed significantly increased blood leptin levels. unpaired t-test, two-

tailed, **$p < 0.01$.

(I) Female cKO mice showed reduced energy expenditure during the dark phase. ns indicates not

significantly different. Two-way ANOVA with Šidák's multiple comparisons test. *$p < 0.05$.

(J) Female cKO mice showed reduced locomotor activity during the dark phase. ns indicates the

difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. *$p < 0.05$.

(K) Female cKO mice showed similar food intake as control mice. ns indicates not significant.

unpaired t-test, two-tailed.

(L) Glucose tolerance test showing that female cKO mice became hyperglycemic 20 minutes

after intraperitoneal glucose administration, suggesting glucose intolerance. Two-way ANOVA

with Šidák's multiple comparisons test. *$p < 0.05$, ***$p < 0.001$.

Data are shown as mean±SEM.

Figure 3: *Rai1* loss reduces intrinsic neuronal excitability and enhances inhibitory synaptic transmission of PVH^BDNF neurons

(A) A representative image showing a patched PVH^BDNF neuron labelled by *Bdnf^Cre*-dependent tdTomato fluorescence signals.

(B-C) Representative traces of spontaneous firing of control (black) and cKO (red) neurons at a holding voltage of -60mV **(B)**. The average spontaneous AP firing frequencies are shown in **(C)**. *$p<0.05$, unpaired student's t-tests.

(D) The resting membrane potentials of control (black) and cKO (red) PVH^BDNF neurons. **$p<0.01$, unpaired student's t-tests.

(E-F) The number of control (black) and cKO (red) PVH^BDNF neurons that showed spontaneous firing (open) or are silent (solid) at resting membrane potentials **(E)** and a holding voltage of -70mV **(F)**.

(G) Representative traces of elicited APs in control (black) and cKO (red) PVH^BDNF neurons responding to an inject current (bottom, blue), which ramps up from the resting membrane potentials at 1 nA per second for 500ms.

(H) The threshold of the first AP initiated by a current ramp in control (black) and cKO (red) PVH^BDNF neurons. *$p<0.05$, unpaired student's t-tests.

(I-J) The amplitude **(I)** and threshold **(J)** for the first AP firing at the holding voltage of -60mV in control (black) and cKO (red) PVH^BDNF neurons. *$p<0.05$, unpaired student's t-tests.

(K) The cumulative fractions of mIPSC frequency were measured in control (black) and cKO(red) PVH^BDNF neurons. The average frequency of events from a 2-minute stable recording of each neuron is shown in the inset, unpaired student's t-tests.

(L) The cumulative fractions of mIPSC amplitude of control (black) and cKO (red) PVH^BDNF neurons. The average amplitude of events from a 2-min stable recording of each neuron is normalized to its capacitance and shown in the inset, unpaired student's t-tests.

(M) The area under the curve of each mIPSC event from a 2-minute recording is averaged. Controls are shown in black, and cKOs are shown in red. unpaired t-tests, two-tailed. *p<0.05.

Data are shown as mean±SEM. Statistics: neuron numbers are in brackets (more than 3 mice/genotype)

Figure 4: Selective *Rai1* ablation in PVH^BDNF neurons induces obesity

(A) A schematic showing stereotaxic injection of adeno-associated viruses (AAVs) expressing Cas9 protein and single-guide RNAs targeting *Rai1* (sgRai1) to delete *Rai1* from the PVH^BDNF neurons of *Bdnf*^Cre/+ female mice. Scale bar = 50μm.

(B) Representative images showing that RAI1 immunoreactivity (cyan) was dramatically reduced in *Bdnf*^Cre/+*; Ai9* mice injected with both sgRai1 and DIO-saCas9 viruses (Exp group n =3, bottom row) into the PVH. In contrast, many PVH^BDNF neurons in *Bdnf*^Cre/+*; Ai9* mice injected only with the sgRai1 virus showed RAI1 expression (Ctrl group n =3, top row).

(C) The percentage of PVH^BDNF neurons co-expressing RAI1 and GFP (virus) was significantly reduced in the Exp group, indicating a successful *Rai1* deletion. unpaired t-test, two-tailed, ****p<0.0001

(D) PVH^BDNF neuron-specific *Rai1* deletion induced body weight gain in Exp mice (Ctrl group: n=5, Exp group: n =5). Two-way ANOVA with Šidák's multiple comparisons test. *p<0.05, **p < 0.01, ****p < 0.0001.

(E) Body composition was measured with Echo-MRI, showing an increased fat mass deposition in 26-week-old Exp mice. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. ***p<0.001, ****p < 0.0001.

(F) Fat deposition analysis shows that the Exp group has significantly more subcutaneous inguinal (S.C.ing) and epididymal white adipose tissue (eWAT) mass (grams). ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. *p<0.05, ****p < 0.0001.

(G) Representative images showing eWAT adipocyte hypertrophy of the Exp group (left). Scale bar = 500µm. Frequency distribution of adipocytes at each cellular size (right). ns indicates not significantly different. Two-way ANOVA with Šidák's multiple comparisons test. $*p<0.05$, $***p<0.001$.

(H) Exp and control mice showed similar food intake. ns indicates not significantly different, unpaired t-test, two-tailed.

(I) Blood leptin levels were significantly increased in the Exp mice. unpaired t-test, two-tailed, $*p<0.05$.

(J) Exp and control mice showed similar energy expenditure. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test.

(K) Exp mice showed reduced locomotor activity in the dark phase. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. $*p<0.05$.

(L) Exp mice became hyperglycemic during the glucose tolerance test. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. $*p<0.05$.

(M) Measurement of the area under the curve (AUC) in Exp and Ctrl mice during the glucose tolerance test. unpaired t-test, two-tailed, $p=0.05$.

(N) Exp mice showed increased plasma insulin levels during the glucose tolerance test. Two-way ANOVA with Šidák's multiple comparisons test, $**p<0.01$.

Data are shown as mean±SEM.

Figure 5: LM22A-4 treatment partially alleviates obesity and stereotypic behaviour in adult SMS mice

(A) Representative western blot showing p-AKT, total AKT, Histone 3 (H3, a loading control) levels in the hypothalamus of saline-treated Ctrl, saline-treated SMS and LM22A-4-treated SMS mice.

(B) Quantification showing deficits in p-AKT levels of the hypothalamus were reversed by LM22A-4 treatment in SMS mice. Unpaired t-test, two-tailed, $*p<0.05$.

(C-E) Body weight of saline-injected control, saline-injected SMS, and LM22A-4 injected SMS mice. Saline or LM22A-4 treatment periods were highlighted in orange. ns indicates not significantly different. Two-way ANOVA with Šidák's multiple comparisons test. $*p<0.05$, $**p<0.01$, $****p<0.0001$.

(F) Food intake (in grams) of saline-injected control, saline-injected SMS, and LM22A-4 injected SMS mice. ns indicates the difference is not significant. unpaired t-test, two-tailed, $*p<0.05$, $**p<0.01$.

(G) Locomotor activity (number of beam breaks) for saline-injected control, saline-injected SMS, and LM22A-4 injected SMS mice. ns indicates the difference is not significant. unpaired t-test, two-tailed, $*p<0.05$.

(H) Fat deposition analysis shows that the saline-injected SMS group has significantly more subcutaneous inguinal (S.C.ing) and epididymal white adipose tissue (eWAT) mass (grams) than saline-injected Ctrl mice. S.C.ing and eWAT mass in LM22A-4 injected mice is comparable to the saline-injected mice. Two-way ANOVA with Šidák's multiple comparisons test. $**p<0.01$, $***p<0.001$, $****p<0.0001$.

(I) High-density lipoprotein (HDL) levels were restored to normal in LM22A-4 injected SMS mice. ns indicates the difference is not significant. unpaired t-test, two-tailed, $**p<0.01$, $***p<0.001$.

(J) Insulin levels in saline-injected control, saline-injected SMS, and LM22A-4 injected SMS mice right before the start of the insulin tolerance test (time point 0). ns indicates the difference is not significant. unpaired t-test, two-tailed, $*p<0.05$.

(K) Insulin tolerance test performed on saline-injected control, saline-injected SMS, and LM22A-4 injected SMS mice, two weeks post daily administration of LM22A-4 or saline, shows reductions in the blood glucose levels of LM22A-4 treated SMS mice. * Shows significance between saline-injected Ctrl and saline-injected SMS group. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test. $**p < 0.01$.

(L) Stereotypic rearing behaviour was fully rescued in LM22A-4 treated SMS mice. ns indicates the difference is not significant. unpaired t-test, two-tailed. $*p < 0.05$, $**p < 0.01$.

Data are shown as mean±SEM.

1069

Fig 1-figure supplement 1: Differentially expressed proteins in the hypothalamus of SMS mice

A heat map showing hierarchical clustering of differentially expressed proteins in the hypothalamus of SMS mice.

1074

1075

Fig 2-figure supplement 1: The *Bdnf^Cre* allele labels BDNF-producing neurons in the paraventricular nucleus of the hypothalamus

(A) Schematic showing a transgenic mouse expressing *Bdnf^Cre*-dependent td-Tomato (Ai9) signals in BDNF-expressing cells.

(B-E) Colocalization of *Bdnf^Cre*-dependent td-Tomato signals with endogenous BDNF protein. Black: DAPI, Magenta: BDNF-producing cells; Green: endogenous BDNF. Scale bar: 50 μm, yellow arrowheads indicate Ai9 marked BDNF cells coexpressing endogenous BDNF.

1083

1084

Fig 2-figure supplement 2: RAI1 expression in BDNF-expressing but not oxytocin-expressing magnocellular PVH neurons

(A-H) Representative images of DAPI expression (blue, **A**), oxytocin (green, **B**), BDNF-producing cells (Ai9 reporter, magenta, **C**), and RAI1 protein (cyan, **D**) in the PVH. Note that BDNF-producing neurons are distinct from the oxytocin-expressing neurons (**E**) and that RAI1 is more selectively expressed in the BDNF-expressing neurons (**F-H**).

(I) Quantification showing the percentage of RAI1$^+$ cells that co-express either BDNF (left bar, magenta) or oxytocin (right bar, green) in the PVH. n=3 mice/group. Data are shown as mean±SEM.

1094

1095

Fig 2-figure supplement 3: Ablating *Rai1* from the BDNF-producing cells induces body weight gain and defective energy homeostasis

(A) Schematic representation of the conditional knockout mice carrying either normal *Rai1* alleles and the Ai9 reporter in BDNF-producing cells (top) or homozygous *Rai1* deletion and the Ai9 reporter in the BDNF-producing cells.

(B) Representative images showing the expression of DAPI (white), RAI1-expressing cells (green), and BDNF-expressing cells (Ai9 reporter, magenta) in the PVH. The top panel shows the Ctrl group with RAI1-Ai9 co-expression (yellow arrowheads) and the bottom panel shows reduced RAI1 and Ai9 colocalization (yellow arrowheads). Note that RAI1 expression was still detected in non-BDNF neurons in the PVH.

(C) Quantification showing the percentage of BDNF-producing cells was not altered by *Rai1* deletion, indicating that RAI1 loss did not impair the generation or survival of PVH^{BDNF} neurons. ns indicates not significantly different, unpaired t-test, two-tailed.

(D) The *Bdnf*$^{Cre/+}$ and *Bdnf*$^{Cre/+}$; *Rai1*$^{fl/+}$ female mice had similar body weights. ns indicates not significantly different. Two-way ANOVA with Šidák's multiple comparisons test.

(E-G) Blood parameter analysis of triglycerides (TG), high-density lipoprotein (HDL), low-density lipoprotein and very low-density lipoprotein (LDL+VLDL) did not differ between Ctrl and cKO mice. ns indicates not significantly different, unpaired t-test, two-tailed.

(H) The respiratory exchange rate did not differ between Ctrl and cKO mice. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test.

(I) In the glucose tolerance test, the cKO mice showed an increased area under the curve (AUC). ***$p < 0.001$, unpaired t-test, two-tailed.

(J) In the glucose tolerance test, cKO mice showed increased insulin levels at the beginning of the test and 30 minutes after glucose administration. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons tests, *$p < 0.05$, **$p < 0.01$.

(K) In the insulin tolerance test, blood glucose level is measured after insulin injection. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test, $*p < 0.05$.

Data are shown as mean±SEM.

Fig 2-figure supplement 4: Metabolic profiles of male cKO mice lacking RAI1 expression in the BDNF-producing cells

(A) Male cKO mice gained significantly more weight than Ctrl mice beginning at 15 weeks of age. Two-way ANOVA with Šidák's multiple comparisons test, $*p < 0.05$, $**p<0.01$, $****p < 0.0001$.

(B) Male $Bdnf^{Cre/+}$ and $Bdnf^{Cre/+}$; $Rai1^{fl/+}$ mice had similar body weights. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test.

(C) Echo-MRI analysis showed that 26-week-old male cKO mice had a significantly increased body weight due to increased fat but not lean mass. ns indicates the difference is not significant. unpaired t-test, two-tailed, $**p<0.01$, $****p<0.0001$.

(D) Fat disposition analysis showing the weight of brown adipocytes (BAT), subcutaneous inguinal (S.C.ing), and epididymal white adipose tissues (eWAT). ns indicates the difference is not significant. unpaired t-test, two-tailed, $*p<0.05$, $**p<0.01$.

(E) Male cKO mice showed eWAT cell hypertrophy. Top: Representative images of the eWAT tissues in Ctrl and cKO mice. Scale bar = 500μm. Bottom: Frequency distribution of cellular sizes. Two-way ANOVA with Šidák's multiple comparisons test, $*p < 0.05$, $**p<0.01$, $***p<0.001$, $****p < 0.0001$.

(F-I) Blood parameter analysis showing that male cKO mice showed significantly increased leptin levels **(F)** without alterations in TG **(G)**, HDL **(H)**, and LDL+VLDL **(I)**. ns indicates the difference is not significant. unpaired t-test, two-tailed. $*p< 0.05$.

(J) No differences were found in the respiratory exchange rate in male Ctrl and cKO mice. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test.

(K) Male cKO mice showed a significantly increased food intake compared to Ctrl littermates., unpaired t-test, two-tailed, $*p < 0.05$.

(L) Male cKO mice showed increased energy expenditure at the dark phase. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test, $*p<0.05$.

(M) Male Ctrl and cKO mice showed similar locomotor activities. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons test.

(N-P) Glucose tolerance test shows that cKO male mice are glycemic 45 minutes post intraperitoneal glucose administration **(N)** and had increased AUC (unpaired t-test, two-tailed) **(O)**, despite significantly higher insulin levels before and 30 minutes after glucose administration **(P)**, suggesting potential insulin insensitivity. ns indicates the difference is not significant. Two-way ANOVA with Šidák's multiple comparisons tests, $*p<0.05$, $**p < 0.01$, $***p<0.001$.

(Q) Insulin tolerance test showing that male cKO mice showed significantly higher blood glucose levels 60 minutes after intraperitoneal insulin administration. Two-way ANOVA with Šidák's multiple comparisons test, $*p<0.05$, $**p < 0.01$.

Data are shown as mean±SEM.

Fig 3-figure supplement 1: The cellular and synaptic properties of control and *Rai1*-deficient PVH^{BDNF} neurons

(A) Representative traces of rebound and rebound repetitive firing of PVH^{BDNF} neurons in control (black) and cKO (red).

(B) Representative AP waveforms at holding voltages of -80mV, -60mV, and -50mV in control (left) and -50mV and -45mV in cKO (right).

(C) The half-width of AP initiated at a holding voltage of -60mV (neuronal numbers in brackets, ns indicates the difference is not significant, unpaired t-test, two-tailed).

(D) Representative traces of miniature recordings in control (black) and cKO (red) PVH^{BDNF} neurons. The traces were recorded with a Cs-based internal at a holding voltage of -70mV in aCSF (top), spontaneous mEPSCs were recorded in aCSF containing 1uM TTX at a holding voltage of -70mV (middle), and mIPSCs were recorded at holding voltage of +10mV (bottom).

(E) Quantification showing that the cellular size of cKO PVH^{BDNF} neurons was significantly smaller than Ctrl neurons. unpaired t-test, two-tailed. ****$p<0.0001$.

Fig 4-figure supplement 1: Metabolic profile of mice lacking *Rai1* in PVH^{BDNF} neurons

(A) Representative images showing co-expression of virus-infected cells (GFP/green), *Bdnf^{Cre}*-dependent Ai9 signals (BDNF cells, magenta), and endogenous RAI1 (cyan). Yellow arrowheads indicate BDNF neurons infected with AAV and in these cells, RAI1 signals were diminished in Exp but not Ctrl mice. Scale bar = 50μm.

(B-C) The total number of PVH^{BDNF} neurons **(B)** and total number of PVH^{BDNF} neurons infected with AAVs **(C)** per slice show that *Rai1* deletion or virus infection did not alter the total number of PVH^{BDNF} neurons. ns indicates the difference is not significant. unpaired t-test, two-tailed.

(D) *Rai1* expression was significantly reduced in PVH^{BDNF} neurons infected with both AAVs compared to PVH^{BDNF} neurons infected with sgRai1-GFP AAV alone. unpaired t-test, two-tailed, **$p<0.01$.

(E) The number of PVH^{BDNF} neurons that co-express RAI1 was significantly reduced in the Exp group. unpaired t-test, unpaired, **$p<0.01$.

(F) Representative image showing lack of viral expression in the PVH neighbouring nuclei. ARC: arcuate nucleus. VMH: ventromedial nucleus of the hypothalamus. 3V: third ventricle. Scale bar = 50μm.

(G) The respiratory exchange rate remained similar in Ctrl and Exp mice. ns indicates the difference is not significant. Two-way ANOVA, Šídák's multiple comparisons test.

(H-J) Blood parameter analysis showing TG, HDL, and LDL+VLDL levels were similar in Ctrl and Exp mice. ns indicates the difference is not significant. unpaired t-test, two-tailed.

(K) In the insulin tolerance test, the Exp group showed a trend towards increased blood glucose levels after insulin injection. Two-way ANOVA, Šídák's multiple comparisons test.

Data are shown as mean±SEM.

Figure 5—figure supplement 1: LM22A-4 treatment does not disrupt energy homeostasis and repetitive rearing in the Ctrl mice

(A) LM22A-4 concentration in the hypothalamus and forebrain of SMS mice 1 hour (black dots) and 3 hours (red dots) after simultaneous intranasal and IP injections.

(B) Four independent sets of western blotting analyses showing reduced p-AKT levels in the hypothalamus of saline-treated SMS mice, which were increased by LM22A-4 treatment. Total AKT protein was used as a control.

(C) Similar body weight of Ctrl mice treated with either saline or LM22A-4. ns indicates the difference is not significant. Two-way ANOVA, Šídák's multiple comparisons test.

(D) Energy expenditure was not different between groups in the light vs. dark phase. ns indicates the difference is not significant. Two-way ANOVA, Šídák's multiple comparisons test.

(E) Respiratory exchange rates were not different between groups in the light vs. dark phase. ns indicates the difference is not significant. Two-way ANOVA, Šídák's multiple comparisons test.

(F) Repetitive rearing behaviour in Ctrl mice was not altered by LM22A-4 treatment. ns indicates the difference is not significant. unpaired t-test, two-tailed.

Data are shown as mean±SEM.

Figure 5—figure supplement 2: LM22A-4 treatment in adult SMS mice is insufficient to

improve social interaction deficit

(A) Schematic showing the tube test that assesses social interaction between stranger mice of
different genotypes.

(B) 7-week-old Ctrl mice won significantly more than SMS mice before the LM22A-4 treatment.
unpaired t-test, two-tailed. ****$p < 0.0001$.

(C) 10-week-old saline-injected Ctrl mice won significantly more than the saline-injected SMS
mice (two weeks post-LM22A-4 treatment). unpaired t-test, two-tailed, ***$p < 0.001$.

(D) 10-week-old saline-injected Ctrl mice won significantly more than LM22A-4 injected SMS
mice. unpaired t-test, two-tailed, *$p < 0.05$.

(E) 10-week-old saline-injected SMS mice showed a similar winning rate to LM22A-4-injected
SMS mice. ns indicates the difference is not significant. unpaired t-test, two-tailed.

Data are shown as mean±SEM.

Figure 1—figure supplement 1

A

BDNF-producing cells

td-Tomato (Ai9)

B — DAPI
C — AI9 (BDNF)
D — anti-BDNF
E — AI9 (BDNF)/anti-BDNF

Figure 2—figure supplement 1

Fig 2-figure supplement 3

Fig 2-figure supplement 4

Figure 3—figure supplement 1

Figure 4—figure supplement 1

Figure 5—figure supplement 1

Figure 5—figure supplement 2

Uncropped western blots for staining against antibodies for P-AKT & AKT.
The membrane was stripped to stain for AKT after staining for p-AKT.
Before staining, the membrane was cut at band 72KDa and below 53 KDa.
The yellow box represents the cropped region for images shown in this
manuscript.

Set 2: Figure 5—figure supplement 1

Set 4: Figure 5—figure supplement 1

~60kD →

~60kD →   p-AKT

~60kD →

~60kD →   AKT

Figure 5A, same as Set 3 in Figure 5—figure supplement 1

p-AKT

AKT

H3

Set 1: Figure 5—figure supplement 1

~60kD →

~60kD →

AKT