ASB3 expression aggravates inflammatory bowel disease by targeting TRAF6 protein

stability and affecting the intestinal microbiota

Short Title: The role of ASB3 in IBD

Mingyang Cheng [1,2,3,4#], Bin Xu[5#], Yu Sun [1,2,3,4], Junhong Wang [1,2,3,4], Yiyuan Lu [1,2,3,4], Chunwei

Shi [1,2,3,4], Tianxu Pan [1,2,3,4], Wenhui Zhao [1,2,3,4], Xiaoxu Li [1,2,3,4], Xiaomei Song [6], Jianzhong Wang

[1,2,3,4], Nan Wang [1,2,3,4], Wentao Yang [1,2,3,4], Yanlong Jiang [1,2,3,4], Haibin Huang [1,2,3,4], Guilian Yang

[1,2,3,4], Yan Zeng [1,2,3,4*], Dongqin Yang[7*], Chunfeng Wang [1,2,3,4*], Xin Cao [1,2,3,4*]

1 College of Veterinary Medicine, Jilin Agricultural University, Changchun, China.

2 Jilin Provincial Key Laboratory of Animal Microecology and Healthy Breeding, Jilin

Agricultural University, Changchun, China.

3 Jilin Provincial Engineering Research Center of Animal Probiotics, Jilin Agricultural University,

Changchun, China.

4 Key Laboratory of Animal Production and Product Quality Safety of Ministry of Education, Jilin

Agricultural University, Changchun, China.

5 Department of General Surgery, Shanghai 10th People's Hospital, Tongji University School

of Medicine, Shanghai 200072, China

6 Department of Gastroenterology, Chongqing General Hospital, Chongqing

7 Central laboratory, Huashan Hospital, Fudan University

# These authors contributed equally

*Correspondence: Yan Zeng, zengyan@jlau.edu.cn, Dongqin Yang, kobesakura@fudan.edu.cn,

Chunfeng Wang, wangchunfeng@jlau.edu.cn, Xin Cao, xinc@jlau.edu.cn.

Abstract

E3 ubiquitin ligase (E3) plays a vital role in regulating inflammatory responses by mediating

ubiquitination. Previous studies have shown that ankyrin repeat and SOCS box-containing protein

3 (ASB3) is involved in immunomodulatory functions associated with cancer. However, the

impact of ASB3 on the dynamic interplay of microbiota and inflammatory responses in

inflammatory bowel disease (IBD) is unclear. Here we systematically identify the E3 ligase ASB3

as a facilitative regulator in the development and progression of IBD. ASB3$^{-/-}$ mice are resistant to

DSS-induced colitis. IκBα phosphorylation levels and production of proinflammatory factors

IL-1β, IL-6, and TNF-α were reduced in colonic tissues of ASB3$^{-/-}$ mice compared to WT mice.

This colitis-resistant phenotype was suppressed after coprophagic microbial transfer and reversed

after combined antibiotic clearance of the microbiota. In addition, ASB3 specifically interacts with

TRAF6 and degrades it via the proteasomal pathway. Further analysis revealed that ASB3

enhances K48-linked polyubiquitination of TRAF6 through the interaction of its SOCS box

(SOCX) domain with the TRAF6 C-terminal domain. Taken together, Our findings demonstrates

that ASB3 is associated with dysregulation of the colitis microbiota and promotes

proinflammatory factors production by disrupting the TRAF6 stability. Strategies to limit the

protein level of ASB3 in intestinal epithelial cells may help in the treatment of colitis.

Keywords: Inflammatory bowel disease; ASB3; Intestinal epithelial cells; Intestinal microbiota;

TRAF6

1. Introduction

Inflammatory bowel disease (IBD) is a heterogeneous group of chronic inflammatory diseases, including Crohn's disease (CD) and ulcerative colitis (UC), which is usually triggered by multiple complex factors.[1] During IBD, various immune cell populations in the intestinal mucosa promote an inflammatory response by secreting proinflammatory factors. Meanwhile, intestinal epithelial cells (IECs) play an essential role in regulating intestinal immune homeostasis by maintaining the integrity of the intestinal mucosal barrier and regulating the functions of intestinal immune cells.[2] IBD-associated genetic defects, epithelial barrier defects, dysregulated immune response, diet, and antibiotic use, among other factors, lead to early dysbiosis of the gut lamina propria (LP), which may precede the development of clinically overt disease.[3] Continued inflammation may lead to late dysbiosis, manifested by an overall reduction in microbial diversity and loss of beneficial symbionts, leading to further exacerbation of the inflammatory response.[4] Currently, the treatment of IBD is limited to immunotherapy. Therefore, understanding the host gene-microbiota causality and interactions in colitis provides new insights for developing therapeutic regimens with synergistic microbiota.[5]

Previous studies have shown that the E3 ubiquitin ligase, one of the critical regulators of autophagy and apoptosis, plays an essential role in normal development, tumor progression, and tissue homeostasis.[6] E3 ubiquitin ligases promote protein degradation through the ubiquitin-proteasome system (UPS). One class of E3 ubiquitin ligases is called SOCS family proteins because of the SOCS box domain.[7] Ankyrin (ANK) repeat and SOCS box containing protein 3 (ASB3) is a member of the ASB family. This family has 18 members, and the protein structure includes the N-terminal ankyrin repeat and the C-terminal SOCS cassette.[8] The

N-terminal domain of the ASB3 protein is responsible for substrate recognition, and the

C-terminal SOCS box performs E3 enzyme function.

Ubiquitination modification is a crucial posttranslational modification (PTM) in the inflammatory

response, affecting the function of essential proteins of the signaling pathway.[9] Dysregulation of

E3 ubiquitin ligase-related response processes has been reported in IBD and numerous

immune-related diseases.[10] For instance, ASB1 promotes TAB2 stability by inhibiting

K48-linked polyubiquitination, which leads to enhanced NF-κB and MAPK activation.[11]

MARCH3 negatively regulates IL-6/STAT3 signaling to limit colitis by targeting IL-6Rα

ubiquitination of IL-6Rα at K401 and subsequent degradation.[12] In addition, TRIM34 regulates

intestinal inflammation by controlling Muc2 cytokinesis in colonic goblet cells (GCs).[13] Thus,

more elucidation of the functions and targets of E3 ubiquitin ligases will provide additional

avenues for the treatment of infectious diseases. To date, despite new evidence of the importance

of ASB3 regulation in cancer development, its role in colonic inflammation has not been

evaluated.

Here, we investigated the physiological role of ASB3 in the intestine and found that

ASB3-deficient mice exhibited reduced susceptibility to dextran sodium sulfate (DSS)-induced

colitis. Further experiments showed that ASB3 deficiency directly protects against Bacteroidetes

and inhibits *Lactobacillus* overgrowth, thus favoring the growth of *Akkermansia*. Moreover, we

demonstrate that IECs ASB3 modulates proinflammatory factors by specifically enhancing

K48-linked polyubiquitination of TRAF6. This novel mechanism precedes the dysregulation of

gut ecology and reinforces the role of ASB3 in maintaining inflammatory signaling and a healthy

gut microbial ecology.

2. Results

2.1. ASB3 expression is upregulated in IBD patients and in mice with DSS-induced colitis

To explore the potential role of ASB3 in IBD, we examined the expression levels of ASB3 in the colonic tissues of UC and CD patients. The results showed that the mRNA expression of ASB3 was upregulated in both UC (n=5) and CD (n=5) compared to controls (Figure 1a, c). Consistent with the transcriptional level, the expression of ASB3 protein was enhanced at the site of IBD inflammation and was accompanied by damage to the intestinal barrier (Figure 1b, d). We then verified the level of ASB3 in the colons of mice. There was an increase in IκBα phosphorylation and in ASB3 expression in the colonic tissue from DSS-treated FVB mice (Figure 1e, f). Furthermore, similar results were observed in colon organoids and HT-29 cells stimulated with TNF-α (Figure 1g-i), suggesting that intestinal inflammation is accompanied by abnormally high ASB3 expression. We also observed that ASB3 expression levels were higher in the colonic mucosa of IBD patients than in the colonic mucosa of uninflamed donors (Figure 1j). In conclusion, abnormally high expression of ASB3 was positively correlated with inflammation in IBD tissues.

2.2. ASB3$^{-/-}$ mice are resistant to DSS-induced colitis

To further investigate the function of ASB3 in the development of colitis, ASB3$^{-/-}$ and WT mice were stimulated with 3% DSS for 6 days and then allowed to recover with normal drinking water for 2 days. The lack of ASB3 expression in the healthy state did not lead to anatomical abnormalities or spontaneous inflammation of the intestine (Figure S1b-d, Supporting

Information). Nevertheless, our results showed that ASB3$^{-/-}$ mice had a significantly lower rate of

weight loss than WT mice after DSS challenge (Figure 2a). We observed that WT mice died

during the acute phase of DSS challenge, while all ASB3$^{-/-}$ mice survived (Figure 2b). In addition,

the reduced severity of colitis observed in ASB3$^{-/-}$ mice was manifested by less diarrhea and fewer

bloody stools, longer colonic length, and reduced clinical scores (Figure 2c-f). To assess this

hypoinflammatory phenotype, we analyzed histopathological sections of the colon. Consistent

with the clinical features described above, ASB3$^{-/-}$ mice exhibited less severe submucosal

inflammatory cell infiltration and less severe epithelial or crypt erosion than WT mice (Figure 2g).

Epithelial cells include enterocytes, goblet cells, enteroendocrine cells, Paneth cells, and M cells,

which are tightly connected and covered with mucus. Intestinal barrier damage and epithelial cell

depletion are regarded as important characteristics associated with mucosal inflammation.[14] We

further confirmed by qPCR and Alcian blue staining that ASB3 expression facilitated a reduction

in the number of cupped cells (Figure 2h, i). We also found that the absence of ASB3 expression

during inflammation attenuated the damage to intestinal epithelial cells (Figure 2j). Our findings

suggest that ASB3 deficiency protects against DSS-induced colonic inflammation.

2.3. ASB3 exacerbates colitis by promoting the release of proinflammatory factors

The development of IBD is closely related to intestinal cytokine storms.[15] To determine whether

ASB3 regulates the production of proinflammatory cytokines, we examined the production of

IL-1β, IL-6, and TNF-α in the mouse colon.[16] As expected, the transcript levels and secretion

levels of IL-1β, IL-6, and TNF-α in the colon of ASB3$^{-/-}$ mice after DSS challenge were much

lower than those in WT mice (Figure 3a, b). Moreover, ASB3 deficiency resulted in diminished

phosphorylation of NF-κB-dependent proteins, exemplified by IκBα (Figure 3c). These findings

establish a link between ASB3 expression and increased IBD exacerbation and solidify that ASB3

expression may exacerbate colonic inflammation through excessive activation of immune

signaling. To a great extent, mucosal immune cells directly alter the function of epithelial cells,

thus adapting organ function to changing demands.[17] The ILC population, which plays a critical

regulatory role in intestinal inflammation, attracted our attention. We observed a significant

decrease in the percentage of ILC3s in Lin- cells in colonic lamina propria lymphocytes (cLPLs)

of WT mice but not in ASB3$^{-/-}$ mice after administration of the drug compared to controls (Figure

3d). WT mice exhibited loss of the IL-22^{+} ILC3 subpopulation in colonic LPLs and reduced

mRNA expression of IL-22 (Figure 3e-g). ILC3-derived IL-22 is essential for tissue repair.[18]

Hence, we are also more convinced by this evidence of the importance of ASB3 expression in

maintaining homeostasis in the intestine.

2.4. Attenuated colitis in ASB3$^{-/-}$ mice is partially dependent on altered microbiota

Currently, the etiology of IBD is thought to be caused by inappropriate interactions between host

cells and the gut microbiota secondary to increased inflammation involving the overgrowth of

harmful opportunistic bacteria or pathogens and the loss of beneficial bacteria.[19] To determine

whether regulation of microbiota by ASB3 resulted in changes in specific bacteria, we performed

a high-throughput sequencing analysis of 16S rDNA isolated from WT and ASB3$^{-/-}$ mice before

and after DSS challenge. These bacteria were originally derived from the same ASB3$^{+/-}$ parents in

our animal facility. The petal and box plots show that the ASB3$^{-/-}$ mice had a lower relative

abundance of fecal bacteria than the WT mice (Figure S2a, b, Supporting Information). Notably,

principal coordinate analysis revealed no segregation between genotypes at steady state. However,

distinct colonies were formed in ASB3$^{-/-}$ mice after DSS challenge (Figure S2c, Supporting

Information). At the phylum level, the abundance of Bacteroidetes was elevated in ASB3$^{-/-}$ mice

before and after DSS treatment, and the abundance of Muribaculaceae was reduced (Figure S2d,

Supporting Information). Furthermore, at the genus level, ASB3$^{-/-}$ mice exhibited a reduced

abundance of *Lactobacillus* and an increased abundance of *Akkermansia* (Figure S2e, Supporting

Information). To determine whether the slowing of colitis observed in ASB3$^{-/-}$ mice was associated

with the composition of the intestinal microbiota, we performed microbiota-transfer studies by

cohousing mice, which leads to exchange of the microbiota through coprophagia. Six- to

eight-week-old WT and ASB3$^{-/-}$ mice were either housed individually (SiHo mice) or cohoused

(CoHo mice) for 6 weeks and then treated with 3% DSS (Figure 4a). We found that SiHo WT

mice and SiHo ASB3$^{-/-}$ mice continued to have significantly different rates of weight loss after

receiving DSS (Figure 4b). However, ASB3$^{-/-}$ mice cohoused with WT mice (CoHo ASB3$^{-/-}$ mice)

exhibited similar rates of weight loss as WT mice (CoHo WT mice) in the same littermates (Figure

4c). Notably, CoHo ASB3$^{-/-}$ mice showed acute phase death on day 9, a trend similar to that of WT

mice after either littermates or non-littermates DSS administration (Figure 4d). Furthermore,

CoHo ASB3$^{-/-}$ and their WT cage companions were similar across almost all measures, including

colon length, disease activity score, and histopathological changes (Figure 4e-g). We also

demonstrated that microbial exchange somewhat alleviated IκBα phosphorylation and

inflammatory factor (IL-1β, IL-6, and TNF-α) expression in the colons of CoHo WT mice (Figure

4h, i). We note that while the transferred microorganisms reduced the enteritis phenotype within

the same litter of WT mice, the difference was narrowed but still existed. We speculate that the

role in IBD pathogenesis of inappropriate host gene-microbe interactions may be causal. To test

this, we first cleared the intestinal flora using a combination of antibiotics and then induced colitis

using DSS (Figure 4j). Interestingly, despite the reversal of the colitis phenotype in ASB3$^{-/-}$ mice,

the phosphorylation level of IκBα in the colon was lower than that in WT mice (Figure 4k). These

data suggest that ASB3 expression may preferentially determine the substrate of intestinal

inflammation, amplifying microbial signals through many inflammatory factors, thereby

exacerbating intestinal barrier damage and the development of colonic inflammation.

2.5. ASB3 promotes NF-κB activation by interacting specifically with TRAF6

A recent study showed that overexpression of ASB17 increased the level of LPS-mediated NF-κB

activation in THP-1 cells.[20] However, the mechanism of ASB3-induced NF-κB activation is

unclear. To determine the role of ASB3 in NF-κB signaling pathway activation, we stimulated WT

and ASB3-overexpressing 293T cells with TNF-α and found that overexpression of ASB3

significantly upregulated the phosphorylation level of IκBα (Figure 5a). In Toll-like receptor or

IL-1 signaling, Myd88-dependent NF-κB activation is linked by multiple signaling molecules,

including TNF receptor associated factor 6 (TRAF6), inhibitor kappa B kinase α (IKKα), inhibitor

kappa B kinase β (IKKβ), TGF-β-activated kinase 1 (TAK1), and TAK1-binding protein 2

(TAB2).[21] We next used an NF-κB reporter gene to screen for key activation sites in

inflammatory pathways. The results showed that ASB3 activated NF-κB reporter gene activity

mediated only by TRAF6 but not by Myd88 and IKKβ (Figure 5b). We also found that

overexpression of ASB3 upregulated IκBα phosphorylation mediated by TRAF6 (Figure 5c).

TRAF6 is most widely described as a positive regulator of NF-κB signaling, and we speculate that

ASB3 involvement in NF-κB activation may involve TRAF6 targeting. We then transfected

labeled ASB3 and other signaling molecules into 293T cells and identified the proteins interacting

with ASB3 by immunoprecipitation. TRAF6 protein, involved in the NF-κB signaling pathway,

was identified as an interaction partner of ASB3 (Figure 5d). Furthermore, confocal imaging

confirmed the specific colocalization of ASB3 and TRAF6 (Figure 5e). We also used *in vitro*

transcription to further demonstrate that ASB3 interacts explicitly with TRAF6 but not TRAF3

(Figure 5f). The same phenomenon was also confirmed in colonic tissue (Figure S3a, b,

Supporting Information). Previous studies have shown that TRAF6 contains an RF domain, a

COIL domain, a ZnF domain, and a TRAF-C domain.[22] ASB3 is composed of two main

functional structure domains, including the ANK domain and the SOCX domain. We found that

ASB3 interacts only with the C-terminal structural domain of TRAF6 (Figure 5g). Surprisingly,

both the ANK domain and the SOCX domain were able to interact with TRAF6 (Figure 5h).

Taken together, these results demonstrate that TRAF6 is essential for ASB3-mediated NF-κB

activation.

2.6. ASB3 destabilizes TRAF6 by enhancing K48-linked polyubiquitination

We previously observed that overexpression of ASB3 specifically upregulated TRAF6-mediated

NF-κB reporter gene activity. We hypothesized that ASB3 might regulate the stability of TRAF6.

To test this assumption, we cotransfected HA-tagged ASB3 with flag-tagged Myd88, TRAF6, or

IKKβ and performed immunoblotting analysis. Overexpression of ASB3 specifically

downregulated TRAF6 expression but not TAB1 and IKKβ expression (Figure 6a). The

ubiquitin-proteasome and autophagy-lysosome pathways are the major systems controlling protein

degradation in eukaryotic cells.[23] We found that ASB3-mediated TRAF6 degradation was

restored only by the proteasome inhibitor MG132 but not by the autophagy inhibitor

3-methylamide (3-MA), the lysosomal inhibitor ammonium chloride (NH$_4$Cl), or the pan-Caspase

inhibitor ZVAD (Figure 6b).

To determine which ubiquitin chain of TRAF6 is modulated by ASB3, we analyzed

ASB3-mediated ubiquitin mutants and the degradation of TRAF6. We found that ASB3 strongly

mediates K48-linked polyubiquitination of TRAF6 in a mammalian overexpression system and in

an IBD sample, but did not potentiate K63-linked ubiquitination of TRAF6 (Figure 6c-e) (Figure

S4, Supporting Information). Adding MG132 led to the consistent conclusion that although

TRAF6 protein expression was restored, the K48-linked polyubiquitin chains still accumulated

(Figure 6f, g). To further explore which structural domain of ASB3 is required for enhanced

K48-linked polyubiquitination of TRAF6, we continued to use ASB3 truncates and examined their

effect on TRAF6 ubiquitination. The results showed that removing the SOCX domain eliminated

ASB3-mediated endogenous K48-linked polyubiquitination of TRAF6 (Figure 6h). In contrast,

truncation of the ANK domain does not eliminate TRAF6 exogenous K48-linked

polyubiquitination (Figure 6i). In conclusion, ASB3 enhances K48-linked polyubiquitination of

TRAF6 through the interaction of its SOCX domain with the TRAF6 C-terminal domain.

2.7. ASB3 disrupts TRAF6 stability in IECs exacerbates lethal inflammation

Since we observed that ASB3 deficiency eliminates K48-linked polyubiquitination of TRAF6 in

mouse colon tissue (Figure 7a), we hypothesize that the mechanism of destabilizing regulation of

TRAF6 by ASB3 may occur in IECs. Next, we constructed a model of colonic organoid

inflammation and found that after TNF-α treatment, organoids from WT mice exhibited abnormal

cell morphology and attenuated TRAF6 protein compared to those from ASB3$^{-/-}$ mice (Figure S5a,

Supporting Information) (Figure 7b). qPCR analysis showed that deletion of ASB3 in organoids

was effective in alleviating the overexpression of proinflammatory factors (Figure 7c). To further

confirm the effect of ASB3 on TRAF6 in IECs, we examined the types of ubiquitinated

modifications. The results showed that ASB3 expression enhanced K48-linked polyubiquitination

of TRAF6 in the organoid (Figure 7d). In addition, similar results were obtained with IECs

dissociated from mouse colon tissue (Figure 7e). Collectively, these results reveal that ASB3

triggers massive release of proinflammatory factors from intestinal epithelial cells by targeting

TRAF6 destabilization.

3. Discussion

Ubiquitination plays an essential role in the pathogenesis and development of IBD.

Ubiquitin-modifying enzymes (UMEs) coordinate the optimal ubiquitination of target proteins

through synergistic effects, thereby maintaining gut homeostasis. However, in some reactions,

these UMEs are aberrantly expressed and thus play a negative role. In the present study, we

demonstrated that excessive activation of the E3 ligase ASB3 promotes the secretion of

proinflammatory factors. In addition, we identified another key role of ASB3 in regulating gut

microbiota imbalance by amplifying inflammatory signals. Mechanistically, ASB3 specifically

enhanced K48-linked polyubiquitination, leading to TRAF6 destabilization. In this manner,

proinflammatory factors are released in large quantities, the intestinal microecology and immune

system are modulated, and damage due to inflammation is exacerbated. Thus, this study reveals a

novel strategy for maintaining intestinal homeostasis by ASB3 protein-regulated inflammatory

responses (Figure 8).

A significant enhanced in ASB3 expression was reported in foci sites from UC and CD patients,

compared to healthy sites (Figure 1a-d, j). This phenomenon of exacerbated mucosal inflammation

was confirmed in both our animal and organoid models (Figure 1e-h). The role of ASB3 in

intestinal diseases is poorly studied and is currently only found to be upregulated in COAD

(Figure S1a, Supporting Information).[24] However, one study reported that ASB3 could inhibit

CRC metastasis by delaying the epithelial-mesenchymal transition.[25] Our current studies showed

that ASB3$^{-/-}$ mice had downregulated susceptibility to experimental colitis and preserved intestinal

barrier function (Figure 2). Interestingly, enhanced ASB3 expression is causally related to

impaired intestinal homeostasis by the massive secretion of colonic proinflammatory factors. Thus,

aberrant expression of ASB3 may have significant pathophysiological consequences.

ASB3 overexpression inhibited TRAF6 protein stability (Figure 6a) but not transcription

(Figure S4a, Supporting Information). We found that ASB3 specifically interacted with TRAF6

and degraded TRAF6 through K48-linked polyubiquitination (Figure 6c-g). A previous study has

reported that increased expression of zinc finger protein A20 inhibited TRAF6/NF-κB

activation.[26] TRAF6 autophagic degradation has also been shown to block NF-κB signaling.[27]

This may seem paradoxical at first; TRAF6 destabilization in our study still resulted in NF-κB

expression and inflammatory factor secretion. Surprisingly, however, we noted in one experiment

that mice lacking TRAF6 in IECs exhibited exacerbated DSS-induced inflammatory responses

that TLR-related signaling molecules (Myd88 and TRIF) did not trigger.[28] TRAF6 signaling

independent of TLRs likely limits DSS-induced colitis not by mediating intrinsic functions

involved in IECs homeostasis, but instead by inducing the release of epithelial cytokines such as

TGF-β to exert direct or indirect anti-inflammatory functions.[29] MALT1 has been reported to

regulate STAT3 during IBD recovery to positively promote mucosal healing.[30] This explains the

upregulation of IL-22 expression in the colon, which may be achieved through interactions

between IECs and ILCs. Numerous studies have shown that the TLR-induced NF-κB signaling

pathway depends on regulation of E3 ligases.[31] A20 was shown to be an important regulator of

NF-κB and to have a bidirectional function in inflammatory stabilization.[32] In fact, we found that

ASB3-mediated destabilization of TRAF6 activated NF-κB *in vivo* and *in vitro*, which may be a

novel mechanism for disrupting the homeostasis of inflammation maintained by TRAF6.

In the present study, we found a negative correlation between ASB3 expression and Lgr5 (Figure

S5b, Supporting Information). Recent reports have confirmed that the E3 ubiquitin ligase TRIM27

maintains intestinal homeostasis by activating Wnt signaling to promote the self-renewal of $Lgr5^+$

ISCs.[33] Intestinal homeostasis requires the integrity of the intestinal epithelial barrier to be

maintained, and epithelial cells maintain dynamic homeostasis by coordinating with other

intestinal cell populations and with cytokines.[34] Whether ASB3 affects ISCs to drive epithelial

cell proliferation and repair needs to be investigated in depth in the future with the help of

organoid models. Immune cells in ASB3-deficient mice behaved normally at steady state, and

intestinal ILC3 numbers were upregulated in inflammatory states (Figure S6, Supporting

Information). Our study also showed more severe intestinal inflammation accompanied by a

reduced amount of ILC3s, possibly limited by the degree of epithelial cell inflammation (Figure

3d). The immune-related roles played by intestinal epithelial cells are primarily driven by

gut-resident natural immune cells. These cell populations depend on the regulation of several

molecules during IBD. DR3-driven loss of ILC3s in the large intestine is an essential factor

contributing to the exacerbation of colitis,[35] and mice lacking CD93 expression in ILC3s also

exhibit impaired IL-22 production and increased colonic inflammation.[36] In another study,

metabolic reprogramming during intestinal LTi cell or ILC3 activation was dependent on the

regulation of the immune checkpoint PD-1.[37] ASB3 has been reported to degrade TNF-R2 via the

proteasome pathway,[8] and whether there is a correlation between this and DR3 remains to be

explored. Our data confirm that targeting ASB3/TRAF6 interactions largely affects the tissue

repair function of ILC3s by disrupting the intestinal barrier. However, we currently have no direct

evidence to suggest that ASB3 is involved in ILC3 regulation.

We observed that ASB3-deficient mice acquired colitis resistance by protecting more

Bacteroidetes species while reducing the sudden increase in *Lactobacillus* (Figure S2, Supporting

Information). Intestinal immunity and homeostasis are regulated by epithelial cell/immune cell

interactions and require fine-tuned communication between the host and microbes. The growth of

certain pathogenic bacteria usually manifests as IBD-associated microbial dysbiosis at the cost of

loss of protective commensal bacteria. Host genetic mutations often amplify this disorder. ANG1

deficiency antagonizes the growth of Lachnospiraceae by increasing the level of *α-Proteobacteria*,

leading to dysbiosis of the intestinal ecology.[38] Similarly, Nlrp12 deficiency led to a reduction in

microbiota diversity, loss of protective intestinal commensal strains of Lachnospiraceae, and an

increase in the highly pathogenic bacterium Erysipelotrichaceae.[5] CARD9 deletion reduces

members of *L. reuteri*, *Allobaculum*, and the phylum Actinobacteria and weakens intestinal

tryptophan catabolic function.[39] Although the gut microbiota changes underlying IBD are not

unique, most of the Bacteroidetes species identified in one study were depleted in IBD, and the

abundance of *Streptococcus* and *Lactobacillus* was increased.[40] Members of Muribaculaceae and

Lachnospiraceae have been identified as significant utilizers of mucin monosaccharides, and

members of Rikenomycetaceae and Bacteroidetaceae are also thought to have functions in the

utilization of mucin monosaccharides.[41] Interconversion of mucin monosaccharide-utilizing

intestinal commensal bacteria caused by ASB3 deficiency before and after DSS challenge also

implies that ASB3 may have potential effects on the intestinal mucus layer.

In this study we report a previously unidentified function of ASB3 deficiency in ameliorating

DSS-induced colitis. Our data suggest that high expression of ASB3 in intestinal epithelial cells

under pathological conditions mediates TRAF6 destabilization, thereby inducing proinflammatory

factor release and impaired intestinal barrier. ASB3 exacerbates the interaction between abnormal

immune signaling and dysbiotic microbiota, creating a vicious cycle that leads to persistent

inflammation and ecological dysregulation in the gut. This evidence increases our understanding

of the pathogenesis of colitis. Interventions of ASB3-mediated proinflammatory pathway and

ASB3-synergistic microbiota may have potential relevance for clinical IBD applications.

4. Experimental Section

Human Samples

Human colon samples from IBD patients were provided by the tissue bank in accordance with its

regulations and approved by the Ethics Committee of Tongji University and Shanghai Tenth

People's Hospital.

Mice

ASB3[-/-] mice were generated using PiggyBac transposon-based targeting technology and were

kindly provided by Professor Dongqin Yang (Shanghai Fudan University). Mice used in this study

were housed under specific-pathogen-free (SPF) conditions and had an FVB genetic background.

To avoid the effect of familial heritability on the colony, ASB3[-/-] mice and the wild-type (WT)

controls were all generated from the same heterozygous (HE) ASB3[+/-] parents. WT, HE, and

ASB3[-/-] mice were identified by PCR amplification and DNA sequencing with the primers

5′-CTGAGATGTCCTAAATGCACAGCG-3′, 5′-CCATGACCAAACCCAATTTACACAC-3′, and

5′-TTGTTTTTTTTCCCCCCTAGACAGG-3′, respectively. For the cohousing experiment,

4-week-old mice from the same mother were divided into either individually reared (SiHo) or

cohoused with age- and sex-matched mice (CoHo) for 6 weeks. For antibiotic treatment, mice

were given an ABX cocktail (33.2 mg ampicillin, 33.2 mg neomycin, 33.2 mg metronidazole, and

16.7 mg vancomycin) orally for 5 consecutive days.[42]

Induction of DSS-mediated acute colitis.

Six- to eight-week-old sex-matched mice were induced to develop acute colitis with 3.0% (w/v)

DSS (MW 36-50 kDa; MP Biomedical) dissolved in sterile, distilled water ad libitum for

experimental days 1 to 6 and were then provided regular water for another 2 days.[43] The DSS

solution was made freshly every day.

Cells, organoids and plasmids

Human embryonic kidney cells (HEK293T) and human colorectal adenocarcinoma cells (HT-29)

were cultured in Dulbecco's modified Eagle's medium (DMEM) and RPMI-1640 supplemented

with 10% (v/v) fetal bovine serum (FBS) and 2 mM glutamine with penicillin (100

U/ml)/streptomycin (100 mg/ml). Intestinal epithelial cells (IECs) or crypts were dissociated from

colonic segments of WT or ASB3$^{-/-}$ mice. Isolated colonic crypts were resuspended in DMEM/F12

medium, counted and resuspended in colonic organoid growth medium and matrix gel (Corning)

at a 1:1 ratio.[44] The medium was changed every other day, and after 7 days, the organoid was

stimulated with 150 ng/ml mouse recombinant TNF-α (PeproTech) for 24 h.[45] All cells were

cultured and maintained at 37 °C with 5% CO_2.

Plasmids for Flag-tagged Myd88, TRIF, TRAF3, TRAF6, TAB1, IKKβ, Myc-tagged ubiquitin

(Myc-K48), NF-κB-Luc and pRL-TK (internal control luciferase reporter plasmid) used in the

study were described previously.[23] Plasmids for HA-tagged ASB3, ASB3 (ΔANK), and ASB3

(ΔSOCX) were kindly provided by Prof. Dongqin Yang (Fudan University, Shanghai, China).

Plasmids for Flag-tagged TRAF6 (aa 260-522), TRAF6 (aa 110-522), and TRAF6 (aa 1-349) were

kindly provided by Prof. Qiyun Zhu (Lanzhou Veterinary Research Institute, Chinese Academy of

Agricultural Sciences, China).

Histology

For histopathological analysis, colon tissue from WT and ASB3$^{-/-}$ mice was fixed in 4%

paraformaldehyde solution overnight for processing and embedded in paraffin wax according to

standard procedures. Then, 3 μm sections were taken and stained with hematoxylin and eosin or

Alcian blue. The stained sections were scanned with a microscope (Leica).

Confocal microscopy

Confocal microscopy was performed as previously described.[46] The indicated plasmids were cotransfected into 293T cells and collected after 24 h. Cells were fixed with 4% paraformaldehyde for 15 min at room temperature and subsequently blocked and permeabilized with 0.5% Triton X-100 5% skim milk 1 h at 4 °C. After blocking, slides for immunofluorescence were incubated with fluorescein-conjugated secondary antibodies (1:500) overnight at 4°C in the dark and mounted with DAPI (Beyotime) (1:1000). Images were acquired with a Zeiss microscope (LSM 710) to visualize stained cells.

Statistical analysis

Data are presented as the mean ± SD unless otherwise indicated. All samples were analyzed using GraphPad Prism 8 software. Data were analyzed by unpaired two-tailed Student's t test or two-way ANOVA with Sidak's test to correct for multiple comparisons. Probability (p) values of < 0.05 were considered significant: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ and $****P < 0.0001$; n.s., not significant.

Figure legends

Figure. 1 Abnormal ASB3 expression levels in IBD patients and in mice with DSS-induced colitis.

a The mRNA expression of ASB3, IL-1β, IL-6, and TNF-α was measured in both UC groups (n=5) relative to controls. **b** Representative immunoblotting of p-IκBα, IκBα, ASB3, villin, EpCAM and

β-actin protein expression in UC. **c** The mRNA expression of ASB3, IL-1β, IL-6, and TNF-α was

measured in both CD groups (n=5) relative to controls. **d** Representative immunoblotting of

p-IκBα, IκBα, ASB3, villin, EpCAM and β-actin protein expression in CD. **e** The expression of

p-IκBα, IκBα, ASB3, and β-actin proteins in the colons of DSS-treated or untreated mice was

detected by Western blotting. **f** The mRNA expression levels of ASB3 in the colons of DSS-treated

or untreated mice were determined by qPCR assay. **g** The expression of p-IκBα, IκBα, ASB3, and

β-actin proteins in TNF-α-treated or untreated organoids was detected by Western blotting. **h** The

mRNA expression levels of ASB3 in the organoids of TNF-α-treated or untreated mice were

determined by qPCR assay. **i** HT-29 cells were treated with TNF-α (150 ng/ml) for 12 h, then

stained with indicated antibody and secondary antibody. The nuclei were stained by DAPI. Scale

bars, 50 μm. **j** Representative IHC staining (left) and quantification (right) of ASB3 in colon

tissues collected from IBD patients. Scale bars, 50 μm. P values less than 0.05 were considered

statistically significant (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$). Differences

between groups in **a**, **c**, **f**, **h** and **j** by unpaired Student's t test.

Figure. 2 ASB3 deficiency provides protection against DSS-induced colitis.

a, b Body weight (**a**) and survival (**b**) of conventionally raised wild-type (WT) and ASB3$^{-/-}$ mice

treated with 3% DSS (above horizontal axes) (WT, n=7; ASB3$^{-/-}$, n=7); the results are presented

relative to initial values, set as 100% (throughout). **c** Representative images of diarrhea/bloody

stools observed on day 6. **d, e** Colon length on day 8. **f** DAI scores were determined on day 6. **g**

Representative images of pathological H&E stained colon sections collected on day 8. Scale bar,

100 μm. **h** The mRNA expression levels of Muc-2 and Reg3γ were determined by qPCR assay. **i**

Representative images of Alcian Blue staining of colon tissues of WT and ASB3[-/-] mice. Scale bar,

250 or 100 μm. **j** IF for EpCAM in the colon; representative image of 3 mice/genotype. Scale bar,

100 μm. Differences between groups in **e**, **f**, **h** were determined by unpaired Student's t test; in **a**

by One-way ANOVA; in **b** by log-rank test.

Figure. 3 ASB3 deficiency inhibits the synthesis and release of proinflammatory cytokines

a qPCR analysis of IL-1β, IL-6, and TNF-α mRNA expression in colon tissue sections from WT

(n=5) and ASB3[-/-] mice (n=5) treated with 3% DSS. **b** Colonic secretion of cytokines. **c** The

expression of p-IκBα, IκBα, ASB3, and β-actin proteins in colon tissues from WT (n=6) and

ASB3[-/-] mice (n=5) treated with or without 3% DSS was detected by Western blotting. **d**

Frequencies of ILC3s in cLPLs from WT and ASB3[-/-] mice treated with or without 3% DSS. **e**

Proportions of IL-22[+] ILC3 cells in the colon of WT (ASB3[+/+]) and ASB3[-/-] mice treated with 3%

DSS. **f** Frequencies and absolute number of IL-22[+] ILC3s in cLPLs from WT (n=9) and ASB3[-/-]

(n=9) mice treated with 3% DSS. **g** The mRNA expression levels of IL-22 were determined by

qPCR assay. Differences were determined in **b**, **c**, **d**, **f** by unpaired Student's t test; in **a**, **g** by

One-way ANOVA.

Figure. 4 Commensal microbiota from ASB3[-/-] mice enhances protection against DSS-induced

experimental colitis in WT mice.

a Schematic representation of the SiHo-CoHo strategy and DSS treatment. **b-d** Body weight (**b, c**)

and survival (**d**) of SiHo-WT, SiHo-ASB3[-/-], Coho-WT or Coho-ASB3[-/-] mice during

DSS-induced colitis. **e-g** Colon length (**e**) representative H&E staining (**f**) and DAI (**g**) of WT,

ASB3$^{-/-}$, Coho-WT or Coho-ASB3$^{-/-}$ mice on day 8 after DSS induction. Scale bar, 50 µm. **h** The

protein expression of p-IκBα, IκBα, and β-actin was detected by Western blotting. **i** The mRNA

expression levels of IL-1β, IL-6, and TNF-α were determined by qPCR assay. **j** Schematic

representation of the DSS challenge after ABX treatment. **k** The protein expression of p-IκBα,

IκBα and β-actin was detected by Western blotting. Difference were determined in **e**, **g**, **h** by

unpaired Student's t test; in **d** by log-rank test.

Figure. 5 ASB3 specifically interacts with TRAF6.

a HEK293T cells were transfected with HA-ASB3 plasmid or a vector control. 24 h after

transfection, the cells were treated with TNF-α (150 ng/ml) for the indicated times. Cell lysates

were separated by SDS-PAGE and analyzed by immunoblotting with the indicated antibodies. **b**

HEK293T cells were transfected with the indicated plasmids along with control vector or

increased amounts of ASB3 expression plasmids. Reporter assays were performed 24 h after

transfection. **c** HEK293T cells were cotransfected with HA-ASB3 and Flag-TRAF6 plasmids or a

vector control. **d, f-h** The indicated plasmids were transfected into HEK293T cells or transcribed

in vitro. Then, coimmunoprecipitation and immunoblotting analyses were performed with the

indicated antibodies. **e** HEK293T cells were transfected with HA-ASB3, Flag-TRAF6 or

Flag-TRAF3 plasmids for 24 h and then stained with Flag antibody or HA primary antibody and

secondary antibody. The nuclei were stained with DAPI. The fluorescence intensity profile of

DAPI (blue), HA-ASB3 (green) and Flag-TRAF6 or Flag-TRAF3 (red) was measured along the

line drawn by ZEN Blue. Scale bars, 50 µm. Difference were determined in **b** by unpaired

Student's t test.

Figure. 6 ASB3 promotes the K48-linked polyubiquitination of TRAF6.

a HEK293T cells transfected with Flag-TRAF6, Flag-TAB1, or Flag-IKKβ with increasing

amounts of HA-ASB3 for 24 h before immunoblotting analysis. **b** HEK293T cells were

transfected with the indicated plasmids for 20 h and then treated with 3-methyladenine (3-MA)

(10 mM), NH_4Cl (20 mM), MG132 (10 μM), or ZVAD (20 μM) for 6 h. The cell lysates were then

analyzed by immunoblotting with the indicated antibodies. **c** HEK293T cells transfected with

Flag-TRAF6, Myc-ubiquitin, or its mutants [KO, which all but one lysine residues were

simultaneously mutated to arginines (K-only)] together with control and ASB3 plasmids were

pre-treated with MG132 (20 μM) for 6h. The cells were then subjected to coimmunoprecipitation

and immunoblotting analysis with the indicated antibodies. **d** HEK293T cells transfected with

Flag-TRAF6 and Myc-K48 with the increased amount of HA-ASB3 for 24 h. **e** HEK293T cells

transfected with Flag-TRAF6 with the increased amount of HA-ASB3 for 24 h. **f** HEK293T cells

transfected with Flag-TRAF6, Myc-K48 together with control and ASB3 plasmids were pretreated

with MG132 (20 μM) for 6 h, and the cells were then subjected to coimmunoprecipitation and

immunoblotting analysis with the indicated antibodies. **g** HEK293T cells transfected with

Flag-TRAF6 with increasing amounts of HA-ASB3 for 20 h and then treated with MG132 (20 μM)

for 6 h. **h, i** HEK293T cells were transfected with the indicated truncated plasmids for 20 h and

then treated with or without MG132 (20 μM) for 6 h, and samples were collected at the indicated

times. The cells were use in ubiquitination assays with the indicated antibodies.

Figure. 7 ASB3 promotes NF-κB signaling by limiting TRAF6 stability in IECs.

a Total colonic tissue proteins from DSS-treated WT and ASB3$^{-/-}$ mice. The samples were use in

ubiquitination assays with the indicated antibodies. **b** Representative H&E staining and IF in the

organoid. Scale bar, 50 μm or 100 μm. **c** The mRNA expression of IL-1β, IL-6, TNF-α, and Muc-2

was measured in both TNF-α-treated organoids relative to controls. **d** Total organoid proteins from

TNF-α-treated WT and ASB3$^{-/-}$ mice. **e** Total protein of IECs from dissociated colonic tissue from

DSS-treated WT and ASB3$^{-/-}$ mice. The samples were used in ubiquitination assays with the

indicated antibodies. Difference were determined in **c** by One-way ANOVA.

Figure. 8 ASB3 aggravates intestinal epithelial inflammation and perturbs gut microbiota

homeostasis by targeting TRAF6 destabilization.

Under pathological conditions, ASB3 is upregulated in intestinal epithelial cells, leading to

TRAF6 degradation via augmented K48-linked polyubiquitination. This results in the activation of

TLR-independent NF-κB signaling, compromising the integrity of the intestinal epithelium and

exacerbating the imbalance between the protective probiotic *Akkermansia* and *Lactobacillus*. Our

study underscores the significance of ASB3 as a crucial mediator for preserving intestinal mucosal

homeostasis by interacting with TRAF6.

Data availability

The 16S rRNA sequencing of the fecal microbiota has been deposited on the Sequence Read

Archive website with the BioProject accession number PRJNA1000707 (Temporary Submission

ID: SUB13720433).

Acknowledgments

We thank Dr. Bin Xu (Department of General Surgery, Shanghai 10th People's Hospital, Tongji

University) for donating the CD patients sample. We thank Prof. Dongqin Yang (Central

laboratory, Huashan Hospital, Fudan University) for donating the UC patients samples, ASB3$^{+/-}$

mice and HA-tagged ASB3 plasmids. We thank Prof. Qiyun Zhu (Lanzhou Veterinary Research

Institute, Chinese Academy of Agricultural Sciences) for donating the Flag-tagged TRAF6

truncated plasmids.

Author contributions

MC performed the majority of the experiments. BX, YS, JW, YL, CS, WZ, XL and XS contributed

and/or supported us for some experiments. BX, and DY provided/analyzed human

material/samples. XC, YZ and MC contributed toward the animal studies and wrote the

manuscript. JW, NW, WY, YJ and HH participated in the inflammation evaluation experiments.

MC and BX analysed and interpreted data. MC, YZ, GY, CW and XC conceived and designed the

experiments, interpreted the data and wrote the manuscript. XC acts as guarantor for the present

study. All authors discussed the results and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (32273043,

32202890, U21A20261, 81970506), the Science and Technology Development Program of

Changchun City (21ZY42) and China Agriculture Research System of MOF and MARA

(CARS-35).

Conflicts of interest The authors disclose no conflicts of interest.

References

[1] Y. Shan, M. Lee, E. Chang, *Annual review of medicine* **2022**, *73*, 455, https://doi.org/10.1146/annurev-med-042320-021020.

[2] S. Mehandru, J. Colombel, *Nature reviews. Gastroenterology & hepatology* **2021**, *18* (2), 83, https://doi.org/10.1038/s41575-020-00399-w.

[3] R. Caruso, B. Lo, G. Núñez, *Nature reviews. Immunology* **2020**, *20* (7), 411, https://doi.org/10.1038/s41577-019-0268-7.

[4] K. Glassner, B. Abraham, E. Quigley, *The Journal of allergy and clinical immunology* **2020**, *145* (1), 16, https://doi.org/10.1016/j.jaci.2019.11.003.

[5] L. Chen, J. Wilson, M. Koenigsknecht, W. Chou, S. Montgomery, A. Truax, W. Brickey, C. Packey, N. Maharshak, G. Matsushima, S. Plevy, V. Young, R. Sartor, J. Ting, *Nature immunology* **2017**, *18* (5), 541, https://doi.org/10.1038/ni.3690.

[6] W. Zhang, F. Liu, Z. Che, M. Wu, Z. Tang, J. Liu, D. Yang, *Science China. Life sciences* **2019**, *62* (12), 1692, https://doi.org/10.1007/s11427-018-9505-0.

[7] R. Li, C. Chen, J. He, L. Zhang, L. Zhang, Y. Guo, W. Zhang, K. Tan, J. Huang, *Virology* **2019**, *532*, 55, https://doi.org/10.1016/j.virol.2019.04.004.

[8] A. Chung, Y. Guan, Z. Yuan, J. Albina, Y. Chin, *Molecular and cellular biology* **2005**, *25* (11), 4716, https://doi.org/10.1128/mcb.25.11.4716-4726.2005.

[9] J. Ruan, D. Schlüter, M. Naumann, A. Waisman, X. Wang, *Trends in molecular medicine* **2022**, *28* (4), 304, https://doi.org/10.1016/j.molmed.2022.01.006.

[10] Y. Xiao, Q. Huang, Z. Wu, W. Chen, *Immunobiology* **2020**, *225* (6), 152026, https://doi.org/10.1016/j.imbio.2020.152026.

[11] P. Hou, P. Jia, K. Yang, Z. Li, T. Tian, Y. Lin, W. Zeng, F. Xing, Y. Chen, C. Li, Y. Liu, D. Guo, *Proceedings of the National Academy of Sciences of the United States of America* **2021**, *118* (3), https://doi.org/10.1073/pnas.2015416118.

[12] H. Lin, L. Feng, K. Cui, L. Zeng, D. Gao, L. Zhang, W. Xu, Y. Sun, H. Shu, S. Li, *Cellular & molecular immunology* **2021**, *18* (12), 2648, https://doi.org/10.1038/s41423-021-00799-1.

[13] Q. Lian, S. Yan, Q. Yin, C. Yan, W. Zheng, W. Gu, X. Zhao, W. Fan, X. Li, L. Ma, Z. Ling, Y. Zhang, J. Liu, J. Li, B. Sun, *Cellular & molecular immunology* **2021**, *18* (2), 350, https://doi.org/10.1038/s41423-020-0366-2.

[14] Y. Yao, G. Kim, S. Shafer, Z. Chen, S. Kubo, Y. Ji, J. Luo, W. Yang, S. Perner, C. Kanellopoulou, A. Park, P. Jiang, J. Li, S. Baris, E. Aydiner, D. Ertem, D. Mulder, N. Warner, A. Griffiths, C. Topf-Olivestone, M. Kori, L. Werner, J. Ouahed, M. Field, C. Liu, B. Schwarz, C. Bosio, S. Ganesan, J. Song, H. Urlaub, T. Oellerich, S. Malaker, L. Zheng, C. Bertozzi, Y. Zhang, H. Matthews, W. Montgomery, H. Shih, J. Jiang, M. Jones, A. Baras, A. Shuldiner, C. Gonzaga-Jauregui, S. Snapper, A. Muise, D. Shouval, A. Ozen, K. Pan, C. Wu, M. Lenardo, *Cell* **2022**, *185* (7), 1172, https://doi.org/10.1016/j.cell.2022.02.013.

[15] I. Sayed, K. Suarez, E. Lim, S. Singh, M. Pereira, S. Ibeawuchi, G. Katkar, Y. Dunkel, Y. Mittal, R. Chattopadhyay, M. Guma, B. Boland, P. Dulai, W. Sandborn, P. Ghosh, S. Das, *The FEBS journal* **2020**, *287* (18), 3967, https://doi.org/10.1111/febs.15236.

[16] W. Fan, C. Ding, S. Liu, X. Gao, X. Shen, M. De Boevre, Z. Gao, M. Li, S. Zhang, Y. Miao, W. Guan, G.
Liu, L. Yan, S. De Saeger, S. Song, *Cell reports* **2022**, *41* (2), 111454,
https://doi.org/10.1016/j.celrep.2022.111454.

[17] A. Diefenbach, S. Gnafakis, O. Shomrat, *Immunity* **2020**, *52* (3), 452,
https://doi.org/10.1016/j.immuni.2020.02.016.

[18] B. Wang, J. Lim, T. Kajikawa, X. Li, B. Vallance, N. Moutsopoulos, T. Chavakis, G. Hajishengallis, *Cell*
reports **2019**, *26* (6), 1614, https://doi.org/10.1016/j.celrep.2019.01.054.

[19] Q. Zhao, C. Maynard, *Gut microbes* **2022**, *14* (1), 2041342,
https://doi.org/10.1080/19490976.2022.2041342.

[20] P. Wan, G. Yang, S. Zhang, Y. Zhang, Y. Jia, X. Che, Z. Luo, P. Pan, G. Li, X. Chen, Q. Zhang, W. Zhang,
Q. Tan, Y. Li, J. Wu, *Frontiers in cellular and infection microbiology* **2022**, *12*, 759077,
https://doi.org/10.3389/fcimb.2022.759077.

[21] M. Hayden, S. Ghosh, *Cell* **2008**, *132* (3), 344, https://doi.org/10.1016/j.cell.2008.01.020.

[22] Z. Chen, Y. Zeng, Y. Wei, Q. Wang, M. Liu, B. Zhang, J. Liu, Q. Zhu, S. Xu, *Virology* **2022**, *568*, 1,
https://doi.org/10.1016/j.virol.2022.01.001.

[23] Y. Zeng, S. Xu, Y. Wei, X. Zhang, Q. Wang, Y. Jia, W. Wang, L. Han, Z. Chen, Z. Wang, B. Zhang, H.
Chen, C. Lei, Q. Zhu, *PLoS pathogens* **2021**, *17* (2), e1009300,
https://doi.org/10.1371/journal.ppat.1009300.

[24] L. Mu, Z. Han, S. Yu, A. Wang, D. Chen, S. Kong, Y. Gu, L. Xu, A. Liu, R. Sun, Y. Long, *Frontiers in*
immunology **2022**, *13*, 842524, https://doi.org/10.3389/fimmu.2022.842524.

[25] W. Du, Z. Lu, W. Ye, X. Fu, Y. Zhou, C. Kuang, J. Wu, Z. Pan, S. Chen, R. Liu, W. Huang, *Chinese*
journal of cancer **2017**, *36* (1), 11, https://doi.org/10.1186/s40880-017-0180-0.

[26] H. Deng, Q. Deji, W. Zhaba, J. Liu, S. Gao, Y. Han, M. Zhou, C. Wang, *Frontiers in immunology* **2021**,
12, 623256, https://doi.org/10.3389/fimmu.2021.623256.

[27] T. Deng, B. Hu, X. Wang, S. Ding, L. Lin, Y. Yan, X. Peng, X. Zheng, M. Liao, Y. Jin, W. Dong, J. Gu, J.
Zhou, *Autophagy* **2022**, *18* (12), 2781, https://doi.org/10.1080/15548627.2022.2047384.

[28] K. Vlantis, A. Polykratis, P. Welz, G. van Loo, M. Pasparakis, A. Wullaert, *Gut* **2016**, *65* (6), 935,
https://doi.org/10.1136/gutjnl-2014-308323.

[29] P. Beck, I. Rosenberg, R. Xavier, T. Koh, J. Wong, D. Podolsky, *The American journal of pathology*
2003, *162* (2), 597, https://doi.org/10.1016/s0002-9440(10)63853-9.

[30] L. Wittner, L. Wagener, J. J. Wiese, I. Stolzer, S. M. Krug, E. Naschberger, R. Jackstadt, R. Beyaert, R.
Atreya, A. A. Kühl, G. Sturm, M. Gonzalez-Acera, J. V. Patankar, C. Becker, B. Siegmund, Z. Trajanoski, B.
Winner, M. F. Neurath, M. Schumann, C. Günther, *International Journal of Molecular Sciences* **2023**, *24*
(8), 7402.

[31] Y. Lu, R. Huang, J. Ying, X. Li, T. Jiao, L. Guo, H. Zhou, H. Wang, A. Tuersuntuoheti, J. Liu, Q. Chen, Y.
Wang, L. Su, C. Guo, F. Xu, Z. Wang, Y. Lu, K. Li, J. Liang, Z. Huang, X. Chen, J. Yao, H. Hu, X. Cheng, Y.
Wan, X. Chen, N. Zhang, S. Miao, J. Cai, L. Wang, C. Liu, W. Song, H. Zhao, *Signal transduction and*
targeted therapy **2022**, *7* (1), 185, https://doi.org/10.1038/s41392-022-00985-1.

[32] X. Yu, W. Li, Q. Deng, H. Liu, X. Wang, H. Hu, Y. Cao, Z. Xu-Monette, L. Li, M. Zhang, Z. Lu, K. Young,
Y. Li, *Blood* **2021**, *137* (12), 1615, https://doi.org/10.1182/blood.2020004918.

[33] J. Wang, D. Zhao, Z. Lei, P. Ge, Z. Lu, Q. Chai, Y. Zhang, L. Qiang, Y. Yu, X. Zhang, B. Li, S. Zhu, L.
Zhang, C. Liu, *Cellular & molecular immunology* **2023**, *20* (2), 158,
https://doi.org/10.1038/s41423-022-00963-1.

[34] Y. Xie, Y. Zhao, L. Shi, W. Li, K. Chen, M. Li, X. Chen, H. Zhang, T. Li, Y. Matsuzawa-Ishimoto, X. Yao,
D. Shao, Z. Ke, J. Li, Y. Chen, X. Zhang, J. Cui, S. Cui, Q. Leng, K. Cadwell, X. Li, H. Wei, H. Zhang, H. Li, H.
Xiao, *The Journal of clinical investigation* **2020**, *130* (4), 2111, https://doi.org/10.1172/jci133264.
[35] J. Li, W. Shi, H. Sun, Y. Ji, Y. Chen, X. Guo, H. Sheng, J. Shu, L. Zhou, T. Cai, J. Qiu, *Nature*
communications **2019**, *10* (1), 3371, https://doi.org/10.1038/s41467-019-11304-8.
[36] J. Huang, H. Lee, X. Zhao, J. Han, Y. Su, Q. Sun, J. Shao, J. Ge, Y. Zhao, X. Bai, Y. He, X. Wang, X.
Wang, C. Dong, *Immunity* **2021**, *54* (4), 673, https://doi.org/10.1016/j.immuni.2021.03.018.
[37] D. Wu, L. Hu, M. Han, Y. Deng, Y. Zhang, G. Ren, X. Zhao, Z. Li, P. Li, Y. Zhang, S. Chen, J. Li, Y. Shi, J.
Xue, P. Wang, C. Zhong, *Nature metabolism* **2022**, *4* (7), 867,
https://doi.org/10.1038/s42255-022-00595-9.
[38] D. Sun, R. Bai, W. Zhou, Z. Yao, Y. Liu, S. Tang, X. Ge, L. Luo, C. Luo, G. Hu, J. Sheng, Z. Xu, *Gut* **2021**,
70 (4), 666, https://doi.org/10.1136/gutjnl-2019-320135.
[39] B. Lamas, M. Richard, V. Leducq, H. Pham, M. Michel, G. Da Costa, C. Bridonneau, S. Jegou, T.
Hoffmann, J. Natividad, L. Brot, S. Taleb, A. Couturier-Maillard, I. Nion-Larmurier, F. Merabtene, P.
Seksik, A. Bourrier, J. Cosnes, B. Ryffel, L. Beaugerie, J. Launay, P. Langella, R. Xavier, H. Sokol, *Nature*
medicine **2016**, *22* (6), 598, https://doi.org/10.1038/nm.4102.
[40] M. Schirmer, A. Garner, H. Vlamakis, R. Xavier, *Nature reviews. Microbiology* **2019**, *17* (8), 497,
https://doi.org/10.1038/s41579-019-0213-6.
[41] F. Pereira, K. Wasmund, I. Cobankovic, N. Jehmlich, C. Herbold, K. Lee, B. Sziranyi, C. Vesely, T.
Decker, R. Stocker, B. Warth, M. von Bergen, M. Wagner, D. Berry, *Nature communications* **2020**, *11* (1),
5104, https://doi.org/10.1038/s41467-020-18928-1.
[42] H. Li, J. Xie, X. Guo, G. Yang, B. Cai, J. Liu, M. Yue, Y. Tang, G. Wang, S. Chen, J. Guo, X. Qi, D. Wang,
H. Zheng, W. Liu, H. Yu, C. Wang, S. Zhu, F. Guo, *Gut microbes* **2022**, *14* (1), 2127456,
https://doi.org/10.1080/19490976.2022.2127456.
[43] S. Cui, C. Wang, W. Bai, J. Li, Y. Pan, X. Huang, H. Yang, Z. Feng, Q. Xiang, L. Fei, L. Zheng, J. Huang,
Q. Zhang, Y. Wu, Y. Chen, *Science advances* **2020**, *6* (43), https://doi.org/10.1126/sciadv.aaz7290.
[44] H. Miyoshi, T. Stappenbeck, *Nature protocols* **2013**, *8* (12), 2471,
https://doi.org/10.1038/nprot.2013.153.
[45] J. Zhang, L. Cen, X. Zhang, C. Tang, Y. Chen, Y. Zhang, M. Yu, C. Lu, M. Li, S. Li, B. Lin, T. Zhang, X.
Song, C. Yu, H. Wu, Z. Shen, *Redox biology* **2022**, *56*, 102469,
https://doi.org/10.1016/j.redox.2022.102469.
[46] M. Cheng, M. Kanyema, Y. Sun, W. Zhao, Y. Lu, J. Wang, X. Li, C. Shi, J. Wang, N. Wang, W. Yang, Y.
Jiang, H. Huang, G. Yang, Y. Zeng, C. Wang, X. Cao, *Journal of virology* **2023**, *97* (2), e0192322,
https://doi.org/10.1128/jvi.01923-22.

a

ASB3 mRNA expression (Folds)

p=0.0049

Ctrl   UC

b

| | #1 | #2 | #3 | #4 | Patients |
| | C UC | C UC | C UC | C UC | |

α-ASB3 —55 kD
α-pIκBα —40 kD
α-villin —100 kD
α-Epcam —35 kD
α-β-actin —40 kD

c

ASB3 mRNA expression (Folds)

p=0.0337

Ctrl   CD

d

| | #1 | #2 | #3 | #4 | Patients |
| | C CD | C CD | C CD | C CD | |

α-ASB3 —55 kD
α-pIκBα —40 kD
α-villin —100 kD
α-Epcam —35 kD
α-β-actin —40 kD

e

| | WT | WT+DSS |
| | #1 #2 #3 #4 | #1 #2 #3 |

α-p-IκBα —40 kD
α-IκBα —40 kD
α-ASB3 —55 kD
α-β-actin —40 kD

f

ASB3

Rel. mRNA Level

*

WT   WT+DSS

g

Organoid

| | WT | WT+TNF-α |
| | #1 #2 | #1 #2 |

α-p-IκBα —40 KD
α-IκBα —40 KD
α-ASB3 —55 KD
α-β-actin —40 KD

h

ASB3

Rel. mRNA Level

****

WT   WT+TNF-α

i

AF488-ASB3

Ctrl   TNF-α

10 μm

j

α-ASB3

healthy   IBD

ASB3 IHC Score

*

Healthy   IBD

a

mosue colon

IP:TRAF6
IB:ub-(WT)
IB:ub-(K48)
IB:ASB3
IB:TRAF6

Lysate
IB:TRAF6
IB:β-actin

b

Organoid

WT ASB3⁻/⁻ WT ASB3⁻/⁻

Cy5-TRAF6

c

IL-1β *IL-6* *TNF-α* *Muc-2*

Rel. mRNA Level

Ctrl TNF-α

■ WT ■ ASB3⁻/⁻

****

d Organoid

WT ASB3⁻/⁻
#1 #2 #1 #2

IP:TRAF6
IB:ub-(K48)
IB:TRAF6
IB:ASB3

Lysate
IB:ub-(K48)
IB:TRAF6
IB:ASB3
IB:β-actin

e IECs

WT ASB3⁻/⁻
#1 #2 #3 #1 #2 #3

IP:TRAF6
IB:ub-(K48)
IB:TRAF6
IB:ASB3

Lysate
IB:ub-(K48)
IB:TRAF6
IB:ASB3
IB:β-actin