Nanoscale analysis of human G1 and metaphase chromatin *in situ*

Jon Ken Chen[1], Tingsheng Liu (刘廷生)[1], Shujun Cai (蔡舒君)[1], Weimei Ruan[2], Cai Tong Ng[1], Jian Shi[1], Uttam Surana[2,3], and Lu Gan (甘露)[1,*]

[1] Department of Biological Sciences and Centre for BioImaging Sciences, National University of Singapore, Singapore 117543

[2] Institute of Molecular and Cell Biology and Agency for Science Technology and Research, 61 Biopolis Drive, Singapore 138673

[3] Department of Pharmacology, National University of Singapore, Singapore 117543

* Correspondence: lu@anaphase.org

Keywords:
Chromatin, mitosis, mammalian cells, nucleosome, compaction, cryo-ET

Abbreviations used:
cryo-ET: electron cryotomography / cryo-electron cryotomography
Megacomplex: multi-megadalton globular complexes

New terms:
Dense irregular bodies

ABSTRACT
The structure of chromatin at the nucleosome level inside cells is mysterious. Here we
present *in situ* electron cryotomography analyses of chromatin in both G1 and
metaphase RPE-1 cells. G1 nucleosomes are concentrated in globular chromatin
domains and metaphase nucleosomes are concentrated in the chromatids.
Classification analysis reveals that canonical mononucleosomes, ordered stacked
dinucleosomes, and mononucleosomes with a disordered gyre-proximal density are
abundant in both cell-cycle states. Class averages that have more than two stacked
nucleosomes or that have side-by-side dinucleosomes are not detected, suggesting that
groups of more than two nucleosomes are heterogeneous. Large multi-megadalton
structures are abundant in G1 nucleoplasm, but not found in G1 chromatin domains and
metaphase chromatin. The macromolecular phenotypes studied here represent a
starting point for the comparative analysis of compaction in normal and unhealthy
human cells, other cell-cycle states, other organisms, and *in vitro* chromatin assemblies.

INTRODUCTION

Eukaryotic chromosomes are long nucleoprotein polymers (chromatin) whose basic units are called nucleosomes. The canonical nucleosome is a 10-nm wide, 6-nm thick cylindrical structure that has an eight-histone protein core surrounded by 1.65 left-handed turns (145 – 147 base pairs) of ordered DNA (Hewish & Burgoyne, 1973; Kornberg, 1974; Luger *et al*, 1997; Olins & Olins, 1974). Sequential nucleosomes are connected by short stretches of linker DNA. Groups of nucleosomes (oligonucleosomes) are believed to make higher-order interactions. Electron microscopy studies of purified and cellular samples have led to diverse higher-order chromatin models, including the 30-nm fiber (Finch & Klug, 1976), arc-shaped nucleosome stacks (Dubochet & Noll, 1978), liquid-like chromatin (Eltsov *et al*, 2008; McDowall *et al*, 1986), zig-zags (Bednar *et al*, 1998), 5- to 24-nm-thick fibers (Ou *et al*, 2017), plates (Chicano *et al*, 2019), 100- to 200-nm periodic structures (Hayashida *et al*, 2021), and 200-nm-wide slinkies of sequentially stacked nucleosomes (Sedat *et al*, 2022a; Sedat *et al*, 2022b). These models have not been adequately tested *in situ* in the absence of traditional electron-microscopy sample-preparation perturbations.

In metazoans, chromosomes are decompacted during interphase (Huxley & Zubay, 1961) and compacted to individualized chromatids in mitosis (Flemming, 1880). Super-resolution microscopy studies of interphase mammalian cells have detected small irregular clusters of nucleosomes called clutches (Ricci *et al*, 2015) and 160- to 500-nm-wide chromatin-rich regions called "chromatin domains" (Barth *et al*, 2020; Fang *et al*, 2018; Li *et al*, 2021; Miron *et al*, 2020; Nagashima *et al*, 2019; Nozaki *et al*, 2017; Otterstrom *et al*, 2019; Xiang *et al*, 2018; Xu *et al*, 2018). Chromatin domains have also been indirectly detected by super-resolution microscopy of mitotic cells (Nozaki *et al.*, 2017). Correct chromatin organization is required for important mitotic phenotypes like transcriptional repression and accurate segregation. Recent cryo-EM advancements make it possible to study these rearranged states at the nucleosome level in 3-D.

Electron cryotomography (cryo-ET) is a popular form of cryo-EM that is used to study the *in situ* 3-D structure of cells in a frozen-hydrated life-like state, without the artifacts associated with the chemical fixation, dehydration, and staining (Ng & Gan, 2020). *In situ* cryo-ET studies of eukaryotic chromatin have been done in microbial (Cai *et al*, 2018b; Chen *et al*, 2016; Gan *et al*, 2013; Tan *et al*, 2023; Tan *et al*, 2024), human (Cai *et al*, 2018a; Hou *et al*, 2023), and insect cells (Eltsov *et al*, 2018; Fatmaoui *et al*, 2022). These studies confirmed that the chromatin of multiple species is irregular *in situ* (Itoh *et al*, 2021). Importantly, cryo-ET can reveal nucleosome-level chromatin structure (Cai *et al.*, 2018a; Eltsov *et al.*, 2018), non-chromatin globular multi-megadalton complexes (megacomplexes), and "pockets" – regions that are devoid of megacomplexes and nucleosome-like particles (Cai *et al.*, 2018b). When groups of macromolecular

80 complexes that have very similar conformations are abundant, i.e., they are ordered,
81 they manifest as unambiguous class averages in 3-D classification and subtomogram
82 averaging analysis. *In situ* cryo-ET is therefore an essential tool to test different
83 chromatin compaction models and to discover previously undescribed molecular
84 phenotypes.
85
Here we present cryo-ET analysis of chromatin in cryo-FIB-milled RPE-1 cells, arrested
in G1 phase (herein abbreviated to G1) and in metaphase. These cell-cycle states
represent the extremes of both chromosome compaction and of transcriptional activity.
Our study reveals differences in chromatin structure spanning length scales from ~2 nm
to more than 100 nm, from the mononucleosome to the chromatin domain level.
Subtomogram classification and averaging analyses reveal canonical nucleosomes and
other nucleosome-containing assemblies not seen in previous studies. Remapping
shows that G1 nucleosomes pack into domains and that metaphase chromatin does not
have internal domain-like structure. These findings reveal structural features that are
either shared or that deviate in two vastly different cell-cycle states.

RESULTS

Cell cycle synchronization

We used a pharmacological block-and-release protocol to enrich hTERT-immortalized human retinal pigment epithelial (RPE-1) cells in both G1 and metaphase (Scott *et al*, 2020). First, G1 cells are enriched by arrest with palbociclib. The cells are then released from G1, allowed to grow, and treated with nocodazole to arrest at prometaphase. Finally, the cells are washed free of nocodazole and immediately incubated in MG132 to allow progression to and then arrest at metaphase (Figure S1A). This mitotic arrest is reversible (Scott *et al.*, 2020), which we confirmed by following the loss of mitotic markers histone H3 phosphorylated at serine 10 and Cyclin B1 (Figure S1B). The cytology of both G1 and metaphase cells and chromosomes were verified by immunofluorescence microscopy (Figure S1, C and D). Herein, we refer to palbociclib- and MG132-arrested cells as G1 and metaphase cells, respectively.

In addition to morphological differences, an important phenotypic difference between interphase and mitotic chromatin is the repression of transcription in mitosis. To confirm these phenotypes in synchronized RPE-1 cells, we used antibodies specific for the RNA polymerase II (RNAPII) large subunit Rpb1's C-terminal heptad repeats YSPTSPS phosphorylated at S5 (CTD-S5P) and at S2 (CTD-S2P). These epitopes indicate initiating RNAPII and elongating RNAPII, respectively (Harlen & Churchman, 2017; Komarnitsky *et al*, 2000). As expected, both forms of RNAPII were detected in G1 cell nuclei and were largely excluded from the chromatin of metaphase cells (Figure 1A). These immunofluorescence results are consistent with the near-global repression of chromatin transcription in metaphase cells.

Optimization of cryoprotectants for cellular cryo-ET

In cellular cryo-EM and cryo-ET experiments, cells are frozen and imaged in conditions that immobilize the water molecules in a glassy "vitreous ice" state that resembles the life-like disordered form (Dubochet *et al*, 1988). Cryo-EM images of intact eukaryotic cells have poor contrast because cells are generally too thick. Therefore, frozen-hydrated cellular samples are made much thinner by either cryosectioning (Ladinsky *et al*, 2006; Studer *et al*, 2014) or cryo-FIB milling (Hayles *et al*, 2007; Marko *et al*, 2006; Medeiros *et al*, 2018; Rigort *et al*, 2010; Schaffer *et al*, 2015; Villa *et al*, 2013). We did pilot cryo-ET experiments on metaphase RPE-1 cells prepared by sequential thymidine block and Taxol-induced arrest (before the Scott protocol was published). These cells were then self-pressurized frozen, which is a method we have previously used to prepare frozen-hydrated yeast for cryosections (Cai *et al.*, 2018b; Ma *et al*, 2022; Ng *et al*, 2019; Ng *et al*, 2020), with dextran as an extracellular cryoprotectant. Compared to standard high-pressure freezing, self-pressurized freezing is faster and cheaper (Han *et*

al, 2012; Yakovlev & Downing, 2011). In this method, a cellular sample is sealed in a
copper tube, then submerged in liquid-ethane cryogen. The cellular material that is
closest to the tube's inner-wall surface forms crystalline ice. This ice expands and
pressurizes the material that is cooling deeper inside the tube, suppressing ice-crystal
formation there. Unfortunately, our pilot samples showed diffraction-contrast features
(Dubochet *et al.*, 1988) in the tilt-series projection images (Figure S2, A and B),
indicating that the ice within the frozen cells was crystalline instead of vitreous.
Furthermore, the crevasse artifacts were more severe and the cryosections were more
often detached from the carbon support in our RPE-1 samples than in our yeast ones.
In the cryotomograms, compacted chromosomes have the expected dimensions
(hundreds of nanometers) and are surrounded by ribosomes (Figure S2, C and D), as
previously observed *in situ* (Eltsov *et al.*, 2008; McDowall *et al.*, 1986). Chromatin
densities in our pilot samples appeared string-like, rather than granular, which may be
an artifact from the crystalline ice. While these experiments show that our RPE-1
cryosections are inadequate as cryo-ET samples, they suggest that mitotic
chromosomes can potentially be located due to different image features relative to the
surrounding cytoplasm.
153
An alternative and more-popular way to prepare frozen-hydrated cells is plunge
freezing. While the region of the cell at the mammalian nuclear periphery is just thin
enough (~5 to 10 μm) to be plunge-frozen in unmodified growth medium (medium) in a
vitreous state (Cai *et al.*, 2018a; Mahamid *et al*, 2016), both metaphase cells and the
regions in the nuclei interior are thicker than this limit. Ice-crystal formation in thick
plunge-frozen biological cryo-ET samples can be suppressed by prior treatment with
cryoprotectants (Bäuerlein *et al*, 2022; Cai *et al*, 2022; Creekmore *et al*, 2024; Harapin
et al, 2015; Jentoft *et al*, 2023). Glycerol and DMSO are popular cryoprotectants used to
prepare cell-line cryostocks, so we treated cells with each of these cryoprotectants at
different concentrations and then characterized their morphology by timelapse light
microscopy (Figure S3A). RPE-1 cells that were incubated with 6% and higher
concentrations of DMSO, but not glycerol, showed detachment from the tissue culture
plate and shrinkage of the nucleus after a 6-minute treatment (Figure S3, B and C). At
the highest concentration (9%) of DMSO and glycerol tested, we found that DMSO-
treated cells start detaching in 2 minutes. Therefore, at the highest – and most
cryoprotective – concentrations of DMSO, incubations need to be shorter than 2
minutes, which was done for all following experiments. While this cryoprotectant screen
suggests that glycerol is better than DMSO, the experiments below show that DMSO
treatment allows for higher image contrast and that the treatment time can be shortened
to a few seconds.
174

To determine the effects of cryoprotectants on ice-crystal formation and image contrast,
we performed cryo-ET of cryo-FIB-milled cryolamellae of G1 cells that had been treated
with different concentrations of DMSO and glycerol. Due to the time-consuming nature
of these tests, we used G1 cells, which are easier to prepare, and we used DMSO to
screen the cryoprotectant concentration. Grids bearing the G1 RPE-1 cells were briefly
immersed in PBS+Ca+Mg plus cryoprotectant, then immediately mounted in the
Vitrobot, blotted manually and then plunged into propane-ethane mix (details in
Methods). Because cryoprotection is expected to change with sample thickness, we
kept our analysis to regions of approximate equal thickness, namely the nuclear
periphery, which could be located by the distinctive double membranes of the nuclear
envelope. Tilt series of these samples show that ice-crystal diffraction contrast occupied
large regions in samples treated with ≤ 3% cryoprotectant, smaller regions in 6%
cryoprotectant (Figure S4), and were undetectable in most samples in 9%
cryoprotectant.
189
Next, we collected cryo-ET data of cells in 9% glycerol for comparison with 9% DMSO,
in both defocus phase contrast and with Volta phase plate (VPP). VPP data have more
low-resolution contrast than defocus phase-contrast data (Danev *et al*, 2014; Fukuda *et*
al, 2015). This extra contrast facilitates the detection of macromolecular complexes in
challenging samples like cryotomograms of cells, especially if contrast is further lowered
by cryoprotectant. We found that cells treated with 9% glycerol produce cryotomograms
with unacceptably low contrast, even when the VPP is used (Figure S5). Therefore, a
brief 9% DMSO treatment offers the best combination of cryoprotection and contrast. In
all subsequently collected tilt series, only those without diffraction-contrast artifacts were
used for further analysis.
200
DMSO has been shown to affect the stability of some protein complexes *in vitro* (Chan
et al, 2017). To control for the potential disruption of nucleosomes in 9% DMSO, we
plunge-froze HeLa oligonucleosomes in either pure storage buffer or storage buffer plus
9% DMSO, followed by cryo-ET with a VPP. Cryotomogram densities showed the
characteristic beads-on-string motifs are preserved in 9% DMSO (Figure S6).
Nucleosomes have a conspicuous double-gyre motif (two 10-nm-long parallel linear
densities) in cryotomographic slices (Figure S6, insets). This double-gyre motif is
evident when the nucleosome is oriented with its superhelical axis parallel to the ice
surface, and corresponds to side and gyre views (Zhou *et al*, 2019). These observations
suggest that the 9% DMSO treatment does not disrupt nucleosomes.
211
To further test if DMSO treatment alters nucleosome structure, we performed
subtomogram analysis of the oligonucleosome samples (details in Methods). This
workflow was originally established using nuclear lysates (Cai *et al*, 2018c) and has

three major steps. First, nucleosome-like particles are template-matched using a
rounded-cylinder reference. Template matching done this way is analogous to single-
particle cryo-EM automated particle picking using a Gaussian (2-D) blob that has
dimensions similar to the complex under study. Given the simplistic nature of the
reference, template-matching hits are not considered to be "identified" nucleosomes, so
we refer to them as candidate nucleosomes. Second, canonical nucleosomes are
identified based on their structure by sequential 2-D and then 3-D subtomogram
classification. Alternatively, the subtomograms are 3-D classified directly (2-D
classification skipped), resulting in more nucleosomes detected, though at a cost of
computation time (Tan *et al.*, 2023). Third, the subtomogram class averages are
inspected for known structures, which are subsequently refined in 3-D. Canonical
nucleosome class averages were detected in both the absence of DMSO (Figure S7A)
and presence of 9% DMSO (Figure S7B). Refinement of the canonical nucleosome
classes that have the highest-resolution features resulted in a 19 – 19.5 Å resolution
subtomogram averages; both classes resemble the canonical nucleosome with some
ordered linker DNA as expected of oligonucleosome samples.
231
DMSO – and also glycerol – may also have unexpected effects on cellular health.
Immunofluorescence of G1 and metaphase cells treated with 9% DMSO or 9% glycerol
show a similar CTD-S5P (Figure 1, B and C) and CTD-S2P (Figure S8) distributions
compared to untreated cells. These confocal microscopy data suggest that large-scale
chromatin structure is not disrupted by these cryoprotectant treatments. To control for
the induction of apoptosis, we performed immunoblot analysis for the apoptotic marker,
cleaved poly (ADP-ribose) polymerase-1 (PARP-1) (Kaufmann, 1989; Kaufmann *et al*,
1993). Untreated negative control G1 and metaphase cells did not show cleaved PARP-
1 while positive-control cells treated with the apoptosis inducer staurosporine (Bertrand
et al, 1994) showed cleaved PARP-1 (Figure S9). Neither DMSO nor glycerol treatment
induced cleaved PARP-1 (Figure S9), indicating that brief cryoprotectant treatments
with these cryoprotectants did not induce apoptosis. In summary, brief immersion in
medium containing 9% DMSO both cryoprotects plunge-frozen cells without killing them
and preserves nucleosome structure. In the cryo-ET experiments described below, most
samples were cryoprotected in medium with 9% DMSO, with additional control samples
prepared in medium with either no cryoprotectant or 9% glycerol.
248
Overview of the G1 nucleus interior
G1-arrested cells were immersed in medium containing 9% DMSO, plunge-frozen, cryo-
FIB milled, and then imaged by VPP cryo-ET. In the G1 cell nucleus cryotomographic
slices, we observed masses of nucleosome-like densities, both near the nuclear
envelope (Figure 2A) and further in the nuclear interior (Figure 2B). An additional
example cryotomographic slice from another G1 cell is shown in Figure S10. To control

for the cryolamellae quality, we template matched cytoplasmic ribosomes in G1
cryolamellae (Figure S11) and then directly 3-D classified the candidate ribosomes to
yield 684 ribosome subtomograms. Note that the low number of ribosomes results from
the targeting of nuclei, meaning that cytoplasm occupies small portions of the fields of
view. The refined average is at ~33 Å resolution (Figure S11B) and has the same
handedness as and comparable features to density maps of human ribosome single-
particle cryo-EM reconstructions (Khatter *et al*, 2015) simulated at similar resolutions
(Figure S11C).
263
Densities that resemble the nucleosome gyre view are abundant in the locally
concentrated groups (chromatin domains; see below) in both perinuclear and
intranuclear regions (Figure 2, C and E). In addition to nucleosome-like densities, the
G1 nucleus contains conspicuous non-nucleosome features, of which the most
abundant are megacomplexes (Figure 2D). Because the best-characterized nuclear
megacomplexes are preribosomes, we template matched them using a spherical
reference. We performed both 2-D and 3-D classification, yielding one class average of
candidate preribosomes (Figure S12A). The preribosome average resembles the 60S
complex, which is apparent when the refined structure is oriented to the large subunit of
the 80S ribosome (Figure S12B). In addition to preribosomes, G1 nuclei have dense
bodies with irregular shapes and dimensions more than 40 nm (Figure 2F); because
they are much larger than megacomplexes, herein we refer to them as "dense irregular
bodies". The dense irregular bodies do not have any double-gyre motifs, suggesting
that they do not contain nucleosomes. They are also not a result of DMSO treatment
because we have, in retrospect, observed them in both glycerol-cryoprotected RPE-1
cells and in other cells that were not treated with DMSO (see below).
280
G1 chromatin domains have multiple ordered nucleosome species
The mammalian interphase nucleus contains both chromatin-rich and chromatin-poor
regions. This uneven chromatin distribution is visible in confocal images of RPE-1 cells
(Figure 1), consistent with recent confocal and super-resolution studies of other cell
types (Barth *et al.*, 2020; Cho *et al*, 2022; Fang *et al.*, 2018; Li *et al.*, 2021; Miron *et al.*,
2020; Nagashima *et al.*, 2019; Nozaki *et al.*, 2017; Otterstrom *et al.*, 2019; Xiang *et al.*,
2018; Xu *et al.*, 2018). The super-resolution studies have named the 100- to 500-nm-
wide compact chromatin-rich regions "chromatin domains". Because the nucleosome-
rich regions that we see in G1 cells have dimensions compatible with the chromatin
domains seen by super-resolution microscopy, herein we call them "chromatin
domains", and for brevity, we refer to the chromatin-poor regions as nucleoplasm.
292
To get insight into the nucleosome organization of G1 cells, we performed
subtomogram analysis using a cylindrical reference, which minimizes model bias. We

sped up this initial analysis by doing template matching with a larger grid spacing of 21
nm. The expectation is that a subset of nucleosomes will be missed, which is
acceptable as long as the analysis provides a representative sample of structures within
this large dataset. We analyzed the chromatin domains by direct 3-D classification of
the template-matched candidate nucleosomes. To limit the effects of neighboring
particles in the alignment and classification, we used a tight 110 Å diameter spherical
mask. As a positive control, we restricted subtomogram analysis to the chromatin
domains. We found canonical nucleosome class averages (Figure S13A) like we saw in
HeLa (Cai *et al.*, 2018a). These class averages have the unmistakable features of
canonical nucleosomes: a ~10-nm-long double-gyre density that follows a left-handed
~1.65 turn helical path. Next, we performed a negative-control subtomogram analysis of
the cytoplasm, which does not have chromatin. As expected, the cytoplasm does not
have any nucleosome class averages (Figure S13B).
308
To increase the number of detectable nucleosomes, we repeated the template matching
with a smaller 7 nm grid spacing. Three-dimensional classification of the resultant hits
revealed 4 canonical-nucleosome class averages (Figure S14A). Two of these class
averages had an extra density near one face, suggesting the presence of a nearby
complex that was potentially excluded by the tight spherical mask. Because
cryotomographic slices had densities that resemble stacked nucleosomes (Figure 2, B
and D), we repeated 3-D classification, except using a stacked-double-cylinder
reference and a mask large enough (240 Å sphere) to accommodate two nucleosomes
(Figure S14B). Classification with these new parameters revealed two classes that each
had a canonical nucleosome and an extra density near the face. The subtomograms
belonging to these two classes were pooled and subjected to a second round of 3-D
classification, using the larger mask and one of the nucleosome class averages as the
reference (Figure S14C). This second round of classification revealed a stacked
dinucleosome class average and six class averages that have an ordered
mononucleosome plus an extra, disordered density. In the stacked dinucleosome, both
nucleosome densities have unambiguous features of mononucleosomes with DNA
gyres well resolved; herein we refer to this class as ordered stacked dinucleosomes.
Here, the term "ordered" implies that there are sufficient numbers of canonical
nucleosome pairs that are oriented and positioned with very low heterogeneity, resulting
in a class average with two unambiguous canonical nucleosome densities. Stacked
dinucleosomes have a broader range of inter-nucleosome orientations and positions
would be too heterogeneous to resolve as a separate, unambiguous class average.
331
To potentially detect more subtomograms that contain a canonical nucleosome, we
repeated the first round of 3-D classification with the larger mask and using the stacked
dinucleosome class average as the reference (Figure S14D). This modified

classification strategy revealed three types of class averages: (1) canonical
nucleosomes, (2) a stacked dinucleosome, and (3) a canonical nucleosome with a large
noisy density near the DNA gyres (Figure S14D). The subtomograms of each class
average type were subjected to a third round of classification, resulting in seven classes
from group 1, three classes from group 2, and five classes from group 3 (Figure S14E).
Group 1's class averages primarily differ in the length of the linker DNA and the noisy
densities near the nucleosome face, so we refined each class average separately.
Group 2's classes differ primarily in their noise densities, so we pooled them for
refinement. Group 3's class averages have gyre-proximal densities at variable positions,
so we refined each class separately. The 13 refined class averages have resolutions
ranging from 24 to 34 Å (Figure S15, Movie S1).
346
Group 1's seven refined mononucleosome class averages differ by the length of
ordered linker DNA (Figure 3A). Furthermore, three of these class averages have
asymmetry in the lengths of the two linker DNAs. Note that the difference in the
subtomogram averages' linker DNA lengths may result either from the absolute length
of the individual nucleosomes' linkers or from the heterogeneity of the linker DNA
structure (see Discussion). When rendered at a lower contour level, the classes have a
heterogeneous extra density near one of the faces (Figure S15A), indicating that
another complex is present, though at a lower occupancy (more details below).
Remapping analysis in the later section suggests that some of gyre-proximal densities
have contributions from other nucleosomes.
357
The two nucleosomes of the ordered stacked dinucleosome class average (Group 2)
differ in the lengths of ordered linker DNA (Figure 3B and S15B). As with the Group 1
mononucleosomes, the linker DNA density could appear shorter in one of the
dinucleosome average's members due to either short linker DNA or variable linker DNA
conformation in the contributing nucleosomes (see Discussion). Earlier single-particle
cryo-EM analysis of recombinant oligonucleosome arrays revealed dinucleosome
stacking (Dombrowski *et al*, 2022; Garcia-Saez *et al*, 2018; Song *et al*, 2014). Both the
tetranucleosome and twisted-fiber arrays show two types of dinucleosome stacking:
closely packed dinucleosomes (Type I interface) and less-closely packed nucleosomes
(Type II interface). Following the nomenclature in these studies, the G1 stacked
dinucleosome structures we observe have the Type II interface (Figure 3B).
369
Nucleosomes with a gyre-proximal density (Group 3) are conformationally variable
(Figure 3C and S15C). Each class average has a canonical nucleosome – albeit with
lower-resolution features than in the mononucleosome and stacked dinucleosome
classes. These class averages refined lower resolutions than the other groups (30 – 34
Å vs. low ~20's Å resolution; Figures S15, D – F). Compared to the other groups, the

disc views of this group are undersampled (Figure S15G), meaning that the members of
this class are more abundant than the numbers detected here. When rendered at a
lower contour level, an additional noisy density becomes visible proximal to the DNA
gyres. Even though the additional densities appear noisy, they are not uniformly
distributed around the gyre. Instead, their centers of mass are at four different
superhelical locations (seven when the nucleosome's pseudo-twofold symmetry is
considered); see the analysis with docked nucleosome crystal structure below.
382
Other ordered multiple-nucleosome motifs are not detected in G1
We also attempted to detect ordered sets of nucleosomes packed in other
arrangements. One such arrangement is the side-by-side dinucleosome. The original
mask (240 Å sphere) is big enough to accommodate a side-by-side dinucleosome.
However, the isotropic shape of the mask can enclose other nearby particles, which, if
heterogeneous, would interfere with both alignment and classification. We therefore
designed a mask that better conforms to the longer aspect ratio needed to
accommodate two nucleosomes packed side-by-side. We then performed another
round of 3-D classification on the subtomograms from Group 3 using the new mask and
reference. However, none of the resultant class averages revealed two canonical
nucleosome densities (Figure S16A).
394
The presence of ordered stack dinucleosomes raised the possibility of stacks with more
than two ordered nucleosomes. To test if ordered stacking extends beyond two
nucleosomes in G1 chromatin domains, we modified our classification workflow to
accommodate either three or four stacked nucleosomes (Figure S16B). We used
stacked tubular masks instead of a spherical one and we used the stacked
dinucleosome class average as the reference for an additional round of 3-D
classification on the subtomograms from Group 2. No ordered tri- or tetranucleosome
stacks emerged from 3-D classification. A few classes have noisy densities
above/below the upper-most/bottom-most ordered nucleosome, which may correspond
to an additional heterogeneously oriented/positioned nucleosome or other
macromolecular complex. Because the ordered stacked trinucleosome and
tetranucleosome would be part of stacks with even more ordered nucleosomes
(pentanucleosomes, hexanucleosomes, etc.), class averages with larger ordered
stacked nucleosome assemblies would also be absent.
409
Ordered stacked dinucleosomes not detected in glycerol-treated cells and have a
different structure in non-cryoprotected cells
To control for the effects of DMSO treatment *in situ*, we also performed cryo-ET on G1
cells that were cryoprotected in 9% glycerol (Figure S17, A and B). The nucleosomes in
these cells were also enriched in the domains (Figure S17, C – E), but the DNA gyres

were less-defined than in the DMSO-treated cells. Stacked nucleosomes (Figure S17, C
– E) and dense irregular bodies were also found in the nucleoplasm (Figure S17, F). By
visual inspection, we could not find any notable differences between the glycerol- and
DMSO-cryoprotectant nuclei cryotomogram densities.
419
To determine if glycerol and DMSO treatment lead to differences in chromatin structure
that is too fine to directly visualize, we performed 3-D classification and averaging
analysis. Template-matching and 3-D classification analysis using the same strategy as
for DMSO-treated cells produced class averages of mononucleosomes and
nucleosomes with gyre-proximal densities (Figure S18). No ordered stacked
dinucleosome classes were found. Group 1's class averages refined to 29–34 Å
resolution while group 3's class averages refined to 31 Å resolution (Figure S19).
427
We also performed Volta cryo-ET analysis of G1 cells that were not treated with any
cryoprotectants. Note that these cells were damaged by crystalline ice, whose presence
could be detected in the tilt-series diffraction-contrast features exemplified in our DMSO
concentration screen (Figure S4). Volta cryotomographic slices of perinuclear regions
had greater contrast (Figure S20, A and B) than similar regions imaged in either DMSO-
or glycerol-cryoprotected cells. Enlargements show that the density difference between
nucleosomes and nucleoplasm is more pronounced (Figure S20, C – E) than in cells
cryoprotected with DMSO or glycerol. The nucleoplasm between the macromolecular
complexes appears emptier than in both the DMSO- and glycerol-cryoprotected cells.
This empty-nucleoplasm phenotype, like the string-like chromatin phenotype of the
cryosections, may have resulted from the aggregation artifact that occurs in the
absence of vitreous freezing (Dubochet & Sartori Blanc, 2001; Kellenberger, 1987).
440
To characterize the changes at higher resolution, we did classification analysis using
the workflow described above (Figure S21, A – D), but only detected convincing Group
1 (mononucleosome) class averages. We did not detect any convincing group 2
(ordered stacked dinucleosome) and group 3 (nucleosomes with gyre-proximal
densities) class averages. Group 1's class averages refined to 29 – 56 Å resolution
(Figure S22); note that two of the class averages (2 and 9) have only a few hundred
subtomograms and are therefore at much-lower resolution. Given the lower contrast
cryotomograms of the glycerol-cryoprotected cells and the ice-crystal artifacts in the
non-cryoprotected cells, the analysis below reports only on the DMSO-treated cells, with
caveats where appropriate.
451
G1 chromatin domains are irregular and have few megacomplexes and pockets
To visualize the overall shape of these chromatin-rich regions, we used the EMAN2
convolutional neural network tool (Chen *et al*, 2017; Tang *et al*, 2007) to train and then

automatically segment the chromatin domains (Figure S23). Segmentations were done
at both the nuclear periphery and the nuclear interior (more examples in Figure S24).
Because the cryolamellae are thinner than the chromatin domains, the segmentation
samples a random slice through each domain. Nevertheless, all chromatin domain
segmentations have irregular shapes, suggesting that the domains are themselves
irregular because random sections through platonic bodies like spheres and ovoids
would have circular and oval profiles, respectively.
462
To visualize the nucleosomes' *in situ* 3-D distribution, we remapped them into the
volume of the original cryotomogram along with the preribosomes and (cytoplasmic)
ribosomes (Figure 3D, Movie S2). Consistent with the appearance of cryotomographic
slices, the nucleosomes were not mixed with the preribosomes (megacomplexes).
Instead, they are concentrated in regions that span several hundred nm as expected of
chromatin domains. To determine if nucleosomes belonging to each group had a biased
distribution, we remapped the mononucleosomes, ordered stacked dinucleosomes, or
nucleosomes with gyre-proximal densities into three separate maps (Figure S25), but
did not find any bias or clustering of any of these classes. Furthermore, separate
remaps of each class (7 for mononucleosomes, 1 for stacked dinucleosomes, 5 for
nucleosomes with gyre-proximal densities) did not reveal any biased distributions
(Figure S26). The remapped model shows that the chromatin domains have irregular
shapes, like we expected based on the segmentations. Stacks with four nucleosomes
are also present in the remapped model (Figure 3D, inset II), but are not ordered
because their numbers were too low to detect as a class average with multiple
unambiguous nucleosome densities; see similar analysis of metaphase chromatin
below.
480
Our previous cryo-ET study of fission yeast mitotic chromatin brought our attention to
two non-chromatin features – megacomplexes and regions devoid of macromolecular
complexes, which we termed pockets (Cai *et al.*, 2018b). After inspecting 21
cryotomograms, we found only one candidate intra-domain megacomplex. In these
cryotomograms, we did not find any unambiguous pockets such as the ones seen in
metaphase chromatin (see below). In summary, G1 chromatin domains have few non-
chromatin features that we have previously seen in compacted fission yeast chromatin.
488
Canonical nucleosome classes were not obtained from the nucleoplasm
Mammalian chromatin domains are thought to be connected by short stretches of
chromatin "linker filaments" that extend through the nucleoplasm (Li *et al.*, 2021; Miron
et al., 2020). The composition of these linker filaments is unknown: they could be mostly
naked DNA, oligonucleosome chains, or compacted chromatin fibers. In our cellular
cryotomograms, we are unable to see naked DNA, but we can locate canonical

nucleosomes by class averaging – if they are abundant. We therefore performed
template matching followed by 3-D classification on subtomograms that were outside of
the chromatin domains, i.e., within the nucleoplasm and some cytoplasm. The
nucleoplasm is more constitutionally heterogeneous than the chromatin domains, so we
used 200 classes, of which 157 remained non-empty after the run completed. Only one
canonical-nucleosome class average was found (Figure S27), but remapping showed
that they came from the periphery of the chromatin domains due to imperfect seeding of
the initial search points (not shown). None of the other class averages bear
resemblance to a nucleosome. Therefore, either the linker filament nucleosomes are too
rare to manifest as a nucleoplasmic class average in our dataset or the linker-filament
nucleosomes have structures that we do not yet recognize.
506
Overview of RPE-1 metaphase chromatin
We next prepared DMSO-cryoprotected cryo-FIB-milled metaphase-arrested cells and
then recorded VPP cryo-ET data. Unlike G1 cells, in which the chromatin is separated
from cytoplasm by a high-contrast easy-to-recognize nuclear envelope, metaphase
chromosomes are in contact with the cytoplasm due to mitotic nuclear-envelope
breakdown. This difference complicates metaphase-chromosome targeting for cryo-ET.
Based on clues from our pilot cryotomograms of cryosections, we found that in the
lower-magnification survey images of cryolamellae, cytoplasm regions of cryolamellae
have granular and linear densities from ribosomes, membranes, and cytoskeleton; in
contrast metaphase chromatin have more uniform densities (Figure S28). These visual
cues greatly facilitated target selection. In cryotomographic slices, metaphase chromatin
appear as large regions enriched with nucleosome-like particles and without any
cytoplasmic constituents such as ribosomes, bundles of ~10-nm-diameter filaments
(Figure 4A), and microtubule bundles (Figure 4B). Another metaphase cell example is
shown in Figure S29.
522
Ribosomes are located near the surface of the compacted chromosomes (Figure 4A),
consistent with previous analysis of both HeLa cell cryosections (Eltsov *et al*., 2008),
U2OS cryolamellae (Zhao & Jensen, 2022), our cryosections (Figure S2), and isolated
metaphase chromosomes (Nishino *et al*, 2012). Spindle microtubules that are inserted
into metaphase chromosomes were previously observed in U2OS cells (Kamasaki *et al*,
2013) and are also seen in the RPE-1 chromosomes (Figure 4B). Nucleosomes are
also irregularly packed in the metaphase chromosome surface and interior, without any
large higher-order chromatin structures like 30-nm fibers (Scheffer *et al*, 2011). In RPE-
1 metaphase chromosomes, pockets are present (Figure S29), albeit rarer than in G1.
However, neither megacomplexes nor dense irregular bodies were seen inside any of
the metaphase chromatin regions in our dataset.
534

Mononucleosomes and stacked dinucleosomes in metaphase chromatin

The most conspicuous macromolecular complexes in metaphase chromatin are nucleosome-like particles (Figure 4, C – F). We performed template matching, subtomogram classification, and averaging of metaphase nucleosomes using the same image-processing parameters and workflow as for G1 chromatin (Figure S30, A – E). The individually refined mononucleosome class averages of group 1 have linker DNA of different lengths (Figure 5A). Unlike in G1 chromatin, we did not find a class average that resembles the nucleosome core particle, which has barely detectable linker DNA. Some of the metaphase class averages with shorter linker length appear to be less well-defined, suggesting that they are more heterogeneous. We also observed a mononucleosome class with asymmetric linker DNA lengths. As with the G1 mononucleosome averages, the differences in metaphase mononucleosome linker DNA lengths could arise from multiple factors (see Discussion).

Metaphase chromatin also has numerous densities that resemble stacked nucleosomes (Figure 4, C – F). Accordingly, subtomogram analysis revealed ordered stacked metaphase dinucleosome class averages (Figures S30 C – E and 5B). We also found class averages of nucleosomes with a noisy density proximal to the gyres (Figure S31C, Figure 5C). We refined these class averages to 26 Å (mononucleosomes and stacked dinucleosomes) and 32 – 35 Å (nucleosomes with gyre-proximal densities) (Figure S31, Movie S3). When rendered at a lower contour level, the metaphase stacked dinucleosome and some of the mononucleosome class averages show an extra density near the nucleosome face, like what is seen in G1 (more details below).

We performed the modified search for ordered side-by-side dinucleosomes like we did for G1 chromatin and did not detect any candidates in the metaphase chromatin either (Figure S32A). Likewise, using the same modified strategy as for G1 chromatin, we attempted – and failed – to detect ordered stacked trinucleosomes in metaphase chromatin (Figure S32B). In summary, ordered stacked dinucleosomes, and nucleosomes with gyre-proximal densities are abundant in both G1 and metaphase cells, but ordered structures with two side-by-side or more than two stacked nucleosomes are not.

Next, we remapped the refined class averages of all three nucleosome groups and the ribosomes (Figure 5D). The remapped ribosomes localized exclusively outside the chromosome, consistent with the absence of megacomplexes inside. Remapping revealed a few sets of stacks with more than two nucleosomes (Figure 5D, insets I – IV), suggesting that stacked tri- and tetranucleosomes are present (Figures 4D and E, Movie S4). These tri- and tetranucleosome stacks are too few to manifest as class

574 averages with unambiguous nucleosomes. Therefore, nucleosome stacking can extend
575 beyond two nucleosomes *in situ* in both G1 and metaphase, albeit with less order.
576
As with the G1 cryotomograms, we also remapped the mononucleosomes, ordered
stacked dinucleosomes, and nucleosomes with gyre-proximal densities as three
separate groups (Figure S33), but did not find any distribution bias. Likewise, we
remapped each class (8 for mononucleosomes, 1 for ordered stacked dinucleosomes,
and 4 for nucleosomes with gyre-proximal densities) and did not find biased
distributions of these class averages either (Figure S34).
583
Detailed remapping analysis of nucleosome packing
At least one class average from each group has an extra density, which is visible at a
sufficiently low contour level. To test if these densities are from other nucleosomes, we
remapped all the class averages at low contour levels and then shaded them in different
colors (Figure 6). This visualization scheme allowed us to see if any of the 'extra'
densities overlapped with the nucleosome densities of neighboring class averages. In
both G1 (Figure 6A) and metaphase (Figure 6B), we found (1) examples where the
extra density partially overlapped a nearby remapped nucleosome (black arrowheads),
(2) the extra density did not overlap with a remapped nucleosome but did partially
overlap with a nucleosome-like density in the tomographic slice (gray arrowheads), and
(3) examples of no overlap with either a remapped nucleosome or a nucleosome-like
density in the cryotomographic slice (white arrowhead). In the latter case, there was
some overlap with unknown structures, suggesting that non-nucleosome complexes
may also contribute to the extra densities. In summary, both nucleosomes and non-
nucleosomes contribute to the heterogeneous extra densities in our class averages.
599
Comparative analysis of G1 and metaphase nucleosomes *in situ*
The organization of RPE-1 nucleosomes can be explored in greater detail by
comparison with crystal structures. To simplify the analysis, we assigned the group 1
mononucleosome class averages into only two classes, based on either short or long
linker DNA lengths. These two classes were then refined (Figure S35), separately for
G1 and metaphase nucleosomes. Next, we docked nucleosome and chromatosome
crystal structures as rigid bodies in the refined class averages from both G1 (Figure 7A)
and metaphase (Figure 7B). The main source of variation is the length of ordered linker
DNA in the mononucleosome and the stacked dinucleosome structures. Both G1 and
metaphase stacked dinucleosomes have a center-to-center spacings of ~6.8 nm.
Comparison of the nucleosomes with gyre-proximal densities shows that the centers of
mass of these densities populate the following approximate superhelical locations: 3.8,
4.8, 5.5 and 6 in G1 chromatin and 3.4, 3.9, 4.2 and 5 in metaphase chromatin. For
nucleosomes with gyre-proximal densities, metaphase nucleosomes do not show

interactions near the DNA entry/exit site whereas G1 nucleosomes do (superhelical
location 6).
616
To assess how the populations of the ordered nucleosome-containing subtomograms
differ in G1 and metaphase, we calculated the ratios of class averages of stacked
dinucleosomes plus nucleosomes with gyre-proximal densities relative to the canonical
mononucleosome class averages (all linker DNA lengths) in each cell-cycle state (Table
1). Note that this analysis is limited because (1) the total number of canonical vs. non-
canonical nucleosomes is unknown and (2) only ordered stacked dinucleosomes are
considered here. We found modest differences in the class-average populations:
compared to G1 cells, metaphase cells have ~23% more stacked dinucleosomes and
~24% fewer nucleosomes with gyre-proximal densities relative to mononucleosomes
(Table 1; raw values are in Tables S6 and S7).

DISCUSSION

At the micron length scale, interphase and mitotic chromatin have large morphological differences. Our study reveals both similarities and differences between G1 and metaphase chromatin from the nanometer to the hundred-nanometer length scales *in situ*. We are also able to compare the 3-D organization of intracellular nucleosomes to those studied in recent *in vitro* chromatin samples. The following discussion starts with the smallest length scale and goes to the largest.

***In situ* nucleosome structure and its role in higher-order structure**
RPE-1 cells have canonical nucleosome class averages that differ in the length of ordered linker-DNA densities. The class averages that have long ordered linker DNA in a crossed configuration resemble the chromatosome, which is a nucleosome with an additional 10 to 25 bp of ordered linker DNA (Simpson, 1978; Zhou *et al*, 2015). A crystal structure of the chromatosome with 25 extra base pairs of ordered DNA per arm (Garcia-Saez *et al.*, 2018) could be rigid-body docked into the subtomogram average of the nucleosome classes that have long linker DNA, suggesting that their *in situ* structure resembles the *in vitro* one. The averages that have shorter linker densities can arise from nucleosomes that indeed have shorter linker DNA, nucleosomes that have longer but structurally variable linker DNA, or a combination of both. A mononucleosome class average with ultra-short linker DNA was detected in G1 cells but not in metaphase cells, suggesting that linker DNA ordering may be a consequence of chromatin compaction. The class averages of nucleosomes with gyre-proximal densities all have ultra-short linker DNA, in both G1 and metaphase; these averages could result from nucleosomes that have ultra-short linker DNA or those that have more DNA structural heterogeneity. Nucleosome class averages that have short and long ordered linker DNA were also seen in HeLa, T-lymphoblast, and budding yeast cells (Cai *et al.*, 2018a; Hou *et al.*, 2023; Tan *et al.*, 2023), though canonical nucleosome class averages (regardless of linker DNA length) are much rarer in yeast.

Nucleosome linker DNA orientation and length variability may both contribute to the nucleosome class averages that have short linker-DNA densities. Previous work on isolated chromosomes and oligonucleosomes *in vitro* produced tomograms of unambiguous nucleosomes (Beel *et al*, 2021; Zhang *et al*, 2023; Zhang *et al*, 2022). While it is possible to directly visualize DNA in purified chromatin, which contains nucleosome and DNA densities, such a task is not feasible *in situ*, which contains many non-chromatin densities. Denoising may facilitate the visualization of chromatin densities, but may introduce errors if the training data is biased, for example with simulated densities. Further improvements in image quality, sample preparation, and

666 denoising algorithms may facilitate the annotation and measurement of DNA in
667 unaveraged subtomograms *in situ*.
668
Because we did not enforce C2 symmetry in our subtomogram analysis, the
mononucleosome class averages show linker DNA length asymmetry. This asymmetry
is also evident in the *in situ* class averages of HeLa and budding yeast cells (Cai *et al.*,
2018a; Tan *et al.*, 2023). Nucleosomes with asymmetric linker DNA structure have been
seen in cryo-EM studies of nucleosomes with histones and DNA from cellular lysates
(Arimura *et al*, 2021); reconstituted oligonucleosomes within small condensates (Zhang
et al., 2022); digested native oligonucleosomes (Jentink *et al*, 2023); decompacted
isolated mitotic chromosomes (Beel *et al.*, 2021); and reconstituted chromatosomes
(Bednar *et al*, 2017; Wang *et al*, 2021b; Zhou *et al*, 2021). In contrast, nucleosomes
with symmetric linker DNA have been found *in vitro* in oligonucleosome arrays
assembled from the Widom "601" sequence and that have more than two ordered
nucleosomes (Dombrowski *et al.*, 2022; Ekundayo *et al*, 2017; Garcia-Saez *et al.*, 2018;
Jentink *et al.*, 2023; Lewis *et al*, 2021; Schalch *et al*, 2005; Song *et al.*, 2014; Zhou *et al*,
2022). Taking into account the previous cryo-EM studies, our *in situ* data, and
mesoscale simulations (Collepardo-Guevara & Schlick, 2014), we suggest that *in situ*
linker DNA conformational variability may suppress the formation of ordered
oligonucleosome structures that have more than two nucleosomes.
686
Type II stacked dinucleosomes are abundant in DMSO-cryoprotected cells
DMSO-cryoprotected G1 and metaphase cells have ordered stacked dinucleosomes, in
which DNA gyres and some linker DNA are resolved in both nucleosomes within the
same class average. The *in situ* ordered stacked nucleosomes resemble those studied
in vitro that interact via a Type II interface, (Dombrowski *et al.*, 2022; Garcia-Saez *et al.*,
2018; Song *et al.*, 2014), in which the H4 N-terminal residues from one nucleosome
contact the H2A-H2B acidic patch of its neighbor. The absence of symmetry
enforcement allowed us to see that in the ordered stacked dinucleosome, the two
nucleosomes have a ~10 bp difference in ordered linker DNA. We did not detect
ordered trinucleosome stacks, meaning that the orientation and/or position of the third
nucleosome is variable. Furthermore, while stacks with three nucleosomes are seen in
cryotomographic slices (Figure 4) and stacks with up to four non-ordered nucleosomes
were seen in remapped models (Figures 3D, 5D, 6), such arrangements are rare
compared to ordered stacked dinucleosomes. The rarity of ordering beyond two
nucleosomes is consistent with the absence of the larger ordered structures (fibers and
plates; see below).
703
Our observation that ordered stacked dinucleosomes were not detected in glycerol-
treated cells could result either from insufficient signal-to-noise ratio due to the glycerol

or buffer-induced differences in abundance; further investigation will be needed to
distinguish these possibilities. The non-cryoprotected G1 cells did not have an ordered
stacked dinucleosome class, but did have a class average that contained two
nucleosome-like densities that vaguely resembles a stacking interaction. It is unclear to
what extent ice-crystal artifacts contribute to the absence of ordered stacked
dinucleosomes in these samples. Even though unambiguous ordered stacked
nucleosomes are not present in all three cryoprotection conditions tested, it is clear that
stacking is abundant, as can be commonly seen in tomographic slices. Furthermore, the
other nucleosome-nucleosome interaction (gyre-proximal) can be seen in a subset of
remapped models. These experiments suggest that the detection of some ordered
nucleosome-nucleosome assemblies may depend on the biochemical environment,
quality of freezing, and image contrast, and will require further investigation.
718
Nucleosome stacking has been observed in multiple contexts, with varying degrees of
order. Stacking is found in preparations of mononucleosomes (Bilokapic *et al*, 2018;
Dubochet & Noll, 1978; Eltsov *et al.*, 2018; Zhou *et al*, 2018); reconstituted nucleosome
arrays (Dombrowski *et al.*, 2022; Ekundayo *et al.*, 2017; Garcia-Saez *et al.*, 2018; Geiss
et al, 2014; Jentink *et al.*, 2023; Lewis *et al.*, 2021; Schalch *et al.*, 2005; Song *et al.*,
2014; Takizawa *et al*, 2020; Zhou *et al.*, 2022); isolated natural chromatin (*ex vivo*) from
chicken erythrocytes, starfish spermatozoids, picoplankton, and budding yeast (Cai *et
al.*, 2018c; Scheffer *et al*, 2012); isolated chicken erythrocyte nuclei (Scheffer *et al.*,
2011); *in situ* in interphase HeLa, T-lymphoblast, KE37, non-mitotic fly brain, and fly
embryo cells (Cai *et al.*, 2018a; Eltsov *et al.*, 2018; Fatmaoui *et al.*, 2022; Hou *et al.*,
2023). In contrast, nucleosome stacking was not observed in either budding or fission
yeast *in situ* (Cai *et al.*, 2018b; Tan *et al.*, 2023). In summary, these observations show
that extensive nucleosome stacking is favorable both *in vitro* and *ex vivo* under some
buffer conditions; limited nucleosome stacking is favorable *in situ* in RPE-1 cells; and
that nucleosome stacking is unfavorable *in situ* in yeasts.
734
Other structures that have an ordered nucleosome
The detection of only one group of ordered dinucleosome class averages (stacked,
Type II) indicates that other kinds of inter-nucleosome interactions – for example those
required for side-by-side ordered interactions – are too few to manifest as an
unambiguous class average. We did, however, detect multiple mononucleosome
classes that have a gyre-proximal density of unknown composition. These densities are
not distributed uniformly around the nucleosome disc; instead, their centers of masses
occupy ~7 SHL positions. If these densities did occupy all possible positions around the
gyres, then we probably would not be able to detect them because of their increased
positional heterogeneity. Many nuclear complexes were shown to interact with the
nucleosome gyres, including transcription factors (Michael *et al*, 2020; Nishimura *et al*,

2022; Tanaka *et al*, 2020; Wang *et al*, 2021a), remodelers (Armache *et al*, 2019; Bacic
et al, 2021; Baker *et al*, 2021; Chittori *et al*, 2019; Farnung *et al*, 2020; Farnung *et al*,
2017; Liu *et al*, 2017; Nodelman *et al*, 2022; Yan *et al*, 2019), histone-mark interactors
(Finogenova *et al*, 2020; Grau *et al*, 2021; Kasinath *et al*, 2021; Liu *et al*, 2021; Poepsel
et al, 2018) and RNA polymerase (Ehara *et al*, 2019; Ehara *et al*, 2022; Farnung *et al*,
2022; Farnung *et al*, 2018; Kujirai *et al*, 2018). Other gyre-interacting complexes include
those that either ensconce much of the nucleosome or that bind to a distorted
nucleosome (Chen *et al*, 2022b; Dodonova *et al*, 2020; Eustermann *et al*, 2018;
Farnung *et al*, 2021; Han *et al*, 2020; He *et al*, 2020; Liu *et al*, 2020; Mashtalir *et al*,
2020; Patel *et al*, 2019; Wang *et al*, 2023; Wang *et al*, 2022b, Yuan, 2022; Willhoft *et al*,
2018). Our inspection of remapped subtomogram averages suggest that both
nucleosome and non-nucleosome complexes may contribute to these gyre-proximal
densities. Because many of these complexes are related to transcription, which is
repressed in metaphase and active in G1, we speculate that nucleosomes may
contribute more to the gyre-proximal densities in metaphase than in G1 cells.
761
Canonical nucleosomes are located in irregular domains in G1 but not metaphase
In RPE-1 G1 cell nuclei, most canonical nucleosomes are concentrated within the
conspicuous chromatin domains. These domains have been seen in textbook traditional
electron-microscopy images and have been studied with renewed interest using
advanced light microscopies (Barth *et al.*, 2020; Cho *et al.*, 2022; Fang *et al.*, 2018; Li *et*
al., 2021; Miron *et al.*, 2020; Nagashima *et al.*, 2019; Nozaki *et al.*, 2017; Otterstrom *et*
al., 2019; Xiang *et al.*, 2018; Xu *et al.*, 2018). Consistent with these super-resolution
studies, chromatin domains have irregular shapes throughout the G1 RPE-1 nucleus, in
the perinuclear region in a cryotomogram of an interphase HeLa cell cryolamella (Cai *et*
al., 2018a), and in electron spectroscopic images of mouse fibroblast cells (Strickfaden
et al, 2020). In contrast, domain-like features are not seen in the popular model yeasts
S. cerevisiae and *S. pombe*, which have uniform chromatin distributions throughout the
interphase nucleoplasm (Cai *et al.*, 2018b; Chen *et al.*, 2016; Tan *et al.*, 2023).
775
Single-molecule localization studies have revealed the presence of domain-like features
in mitotic cells. PALM imaging of mitotic mammalian cells revealed domains ranging
from ~140 to 200 nm. (Nozaki *et al.*, 2017). STORM imaging of histones in fixed mitotic
cells (Xu *et al.*, 2018) and of DNA in isolated metaphase chromosomes (Wang *et al*,
2022a) have revealed "nanodomains" – small ~50- to 90-nm granular features
corresponding to locally concentrated chromatin. In our cryotomograms of metaphase
cells, the canonical nucleosomes are uniformly distributed within the compacted
chromosomes (with the exception of pockets; see below), i.e., we did not observe
separate domain-like features within the metaphase chromatin. Human metaphase
chromatin has also been studied *in situ* in 3-D by Chrom-EMT (Ou *et al.*, 2017) and

cryo-ET of cryo-FIB-milled cells (Zhao & Jensen, 2022). Neither of these EM-based
studies reported any nanodomain-like features in mitotic chromatin. It is unclear what
factors explain the visibility of mitotic chromosome domains and nanodomains in light-
microscopy but not in electron microscopy data.
790
The size and shapes of oligonucleosome condensates have been characterized by both
light and electron microscopy (Gibson *et al*, 2019; Maeshima *et al*, 2016; Sanulli *et al*,
2019; Strickfaden *et al.*, 2020; Zhang *et al.*, 2022). Condensates of dodecameric arrays
can grow to more than 4 μm (Gibson *et al.*, 2019; Sanulli *et al.*, 2019), which are an
order of magnitude larger than G1 chromatin domains but the same order as mitotic
chromosomes. Furthermore, condensates of reconstituted arrays have circular profiles,
presumably because they are 2-D projections of spherical bodies (Strickfaden *et al.*,
2020). In contrast, condensates of fractionated (natural) chromatin are irregular
(Maeshima *et al.*, 2016). Subsequent studies using defined '601' chromatin arrays
confirmed that longer arrays (with 62 nucleosome repeats) generate irregular
condensates (Chen *et al*, 2022a) whereas arrays with 17 or fewer nucleosomes form
rounded condensates (Chen *et al.*, 2022a; Gibson *et al*, 2023). Condensates of
reconstituted tetranucleosomes appear irregular during the early, spinodal phase of *in*
vitro condensation and then fuse into larger round ones (Zhang *et al.*, 2022). In
summary, *in vitro* condensates of either natural chromatin, long artificial arrays, or
tetranucleosomes at the early stages of compaction more closely resemble *in situ* G1
chromatin domains than condensates of (short) dodecameric arrays. It remains
unknown how other nuclear factors, like histone post-translational modifications,
remodelers, and chaperones would affect *in vitro* condensates.
810
Large-scale nucleosome packing is irregular in G1 and metaphase
The earliest *in situ* cryo-EM study of mammalian chromosomes was done on Chinese
hamster ovary and HeLa cells. This study proposed that the mitotic chromatin inside
these cells is irregular and "liquid-like" (McDowall *et al.*, 1986), not highly ordered like
30-nm fibers (Finch & Klug, 1976). Our cryo-ET results here show that RPE-1 chromatin
in both G1 and metaphase is also irregular. We did not observe 30-nm fibers, which
have unmistakable dimensions and compact shapes in cryotomograms (Scheffer *et al.*,
2011). The RPE-1 data reported here therefore adds to the growing list of *in situ* cryo-
EM and cryo-ET studies in picoalgae, budding yeast, fission yeast, HeLa, T-
lymphoblast, colorectal adenocarcinoma, and fly neuron cells (Cai *et al.*, 2018a; Cai *et*
al., 2018b; Chen *et al.*, 2016; Eltsov *et al.*, 2018; Eltsov *et al.*, 2008; Fatmaoui *et al.*,
2022; Gan *et al.*, 2013; Hou *et al.*, 2023), which all report that frozen-hydrated
eukaryotic chromatin packing is irregular *in situ*.
824

Recent studies of purified metaphase chromosomes proposed additional models of
higher-order chromatin structure: metaphase chromatin plates in frozen-hydrated
samples, in which nucleosomes are packed edge-to-edge in monolayers (plates), with
interdigitation between adjacent plates (Chicano *et al.*, 2019), and 100- to 200-nm
periodic structures in fixed and heavy-metal-stained samples (Hayashida *et al.*, 2021),
in which the repeat direction is parallel to the chromatid long axis. Our metaphase-cell
data did not show evidence of edge-to-edge packing or 100- to 200-nm periodic
structures. The simplest explanation for these differences is from structural changes
induced by sample preparation (isolated vs. *in situ* chromosomes). Our data does not
rule out the alternative possibility in which the plates are irregular, with nucleosomes
having large deviations relative to the ordered edge-to-edge arrangement depicted in
the earlier study. For the 100- to 200-nm periodic structures, we do not know alternative
hypotheses to explain their absence in our – and others' – cryo-EM and cryo-ET data.
838
A cryo scanning transmission electron microscopy cryotomography study proposed
another higher-order chromatin model that involves long-range ordered packing:
sequential nucleosomes make a continuous stack in both interphase and mitosis (Sedat
et al., 2022a; Sedat *et al.*, 2022b). In this model, termed "slinky", the linker DNA follows
a coiled path along stacks of consecutive nucleosomes akin to the hypernucleosomes
of archaeal chromatin (Bowerman *et al*, 2021; Henneman *et al*, 2018), which was also
seen in reconstituted telomere-like chromatin arrays (Soman *et al*, 2022). This slinky is
proposed to coil into 100- to 300-nm gyruses, leaving a nucleosome-free tunnel in the
middle. None of our class averages have the key feature needed for slinkies: linker DNA
that is curved enough to connect to sequential nucleosomes in the stack. Furthermore,
our cryotomograms do not show any large nucleosome-free regions in either the
chromatin domains or metaphase chromosomes that would account for the hole in the
middle of a slinky gyrus.
852
Non-nucleosome features of G1 nuclei and metaphase chromatin
We previously observed irregular intranuclear regions that we termed "pockets" in
cryotomograms of *S. pombe* cells. Pockets are conspicuous because they lack
macromolecular complexes and megacomplexes. In G1 chromatin domains, we did not
find any instances of pockets, but we did find a few unambiguous ones in metaphase
chromatin. With the exception of pockets, the RPE-1 metaphase chromatin distribution
is uniform and resembles that seen in other cryo-EM and cryo-ET studies of mitotic
human cells (Eltsov *et al.*, 2008; Zhao & Jensen, 2022). We did not detect
mesh/network-like nucleosome-free regions analogous to the "reticular pattern" of
negatively stained cells in a previous study (Ou *et al.*, 2017). Nevertheless, the
presence of pockets in metaphase chromosomes suggests that the compaction is not
maximized.

865
No megacomplexes were seen in the interior of the chromatin domains or mitotic
chromatin, indicating that the crowding disfavors access by very large complexes. In
contrast, mitotic budding and fission yeasts both have megacomplexes mixed with the
chromatin (Cai *et al.*, 2018b; Chen *et al.*, 2016; Ng *et al.*, 2019). This molecular-packing
difference may be related to differences in mitotic transcription: both yeasts have high
levels of mitotic transcription (Cho *et al*, 1998; Oliva *et al*, 2005; Peng *et al*, 2005;
Rustici *et al*, 2004; Spellman *et al*, 1998) whereas mammalian chromatin undergoes
genome-wide transcriptional repression (Flemming, 1880; Prescott & Bender, 1962;
Taylor, 1960). Note that mammalian mitotic chromatin exhibits transient access to small
complexes such as transcription factors (Teves *et al*, 2016) and permits very low levels
of mitotic transcription (Palozola *et al*, 2017). High-resolution correlative light
microscopy and cryo-ET analysis will be needed to relate the 3-D structure of the
nucleosomes to the transcriptional state.
879
The G1 nucleoplasm has many heterogeneous structures
Unlike the canonical-nucleosome-rich chromatin domains, the chromatin-poor G1
nucleoplasm is constitutionally heterogeneous. Our subtomogram analysis did not
produce canonical nucleosome class averages in the nucleoplasm. There are at least
two distinct possible explanations. If the nucleoplasmic nucleosomes have the canonical
structure but too few, then the resultant class averages would be either too noisy to be
recognizable as canonical nucleosomes or may be mis-classified. Alternatively, if the
nucleoplasmic nucleosomes are abundant but have non-canonical structures, then we
would miss them because we do not yet know their structures. The linker-filament
structures remain to be determined (Li *et al.*, 2021; Miron *et al.*, 2020).
890
Nuclear bodies that are larger than nucleosomes are abundant in the nucleoplasm. The
most conspicuous nucleoplasmic features are the megacomplexes, which have been
seen in HeLa and yeast nuclei (Cai *et al.*, 2018a; Cai *et al.*, 2018b; Chen *et al.*, 2016;
Ng *et al.*, 2019). Some of the megacomplexes are preribosomes because they have the
expected size (~20 to 30 nm) (Erdmann *et al*, 2021; Lau *et al*, 2021). Indeed, we have
detected one class average that resembles the large ribosomal subunit. The identities of
the vast majority of megacomplexes remain to be determined.
898
The G1 nucleoplasm also contains dense irregular bodies, which have dimensions that
exceed 40 nm. Dense irregular bodies are also present in our cryotomogram of an
interphase HeLa cell (Cai *et al.*, 2018a) and in *S. cerevisiae* cells (Ng *et al.*, 2019),
though we previously did not appreciate their differences from megacomplexes. No
such bodies have been observed in *S. pombe* yet. Dense irregular bodies are too big to
be preribosomes and are unlikely to harbor canonical nucleosomes because they lack

the characteristic double-gyre motifs. Further work is needed to determine the identity of
these dense irregular bodies and their relationship to megacomplexes (if any), diverse
nuclear condensates that have been studied by light microscopy (Stanek & Fox, 2017;
Zhu & Brangwynne, 2015), or ribonucleoprotein complexes (Pacheco-Fiallos *et al*,
2023).
910
Metaphase chromosome periphery
The metaphase chromosome's surface, termed the chromosome periphery, is
associated with non-nucleosomal complexes such as nucleolar and ribosomal proteins
and RNAs (Booth *et al*, 2016; Nishino *et al.*, 2012; Ohta *et al*, 2010; Stenstrom *et al*,
2020). Ribosomes are present on the surface of metaphase chromosomes, reminiscent
of those seen adjacent to HeLa and U2OS metaphase chromosomes *in situ* (Eltsov *et
al.*, 2008; Zhao & Jensen, 2022) and the ones that co-purify with isolated chromosomes
(Nishino *et al.*, 2012). The co-purified ribosomes generate a ~30-nm peak in small-angle
X-ray scattering experiments that was previously (and erroneously) attributed to 30-nm
chromatin fibers (Langmore & Paulson, 1983; Nishino *et al.*, 2012). The chromosome
periphery is organized by the proliferation marker Ki-67, which extends ~100 nm from
the chromatin surface and keeps the chromosomes individualized via a surfactant
function (Booth *et al*, 2014; Cuylen *et al*, 2016). Even though the Ki-67 protein (> 300
kDa) is more massive than a nucleosome (200 kDa) – much of its sequence is predicted
to be intrinsically disordered (Remnant *et al*, 2021), meaning that it has an extended
structure. Long peptide-wide proteins are too thin to see in cryotomograms of cells,
which would explain the absence of filamentous densities in chromosome-periphery
regions of our cryotomograms. Instead, our cryotomograms show that the
macromolecular contents of the nuclear periphery are both heterogeneous and lack
order, consistent with the disordered nature of Ki-67.
931
Conclusions
Our study has revealed *in situ* similarities and differences of G1 and metaphase
chromatin across multiple length scales (Table 2). Both G1 chromatin domains and
metaphase chromosomes are densely packed with canonical nucleosomes and are
largely devoid of megacomplexes. Canonical nucleosome class averages have variable
ordered-linker-DNA lengths, though only G1 cells have a nucleosome class with ultra-
short linker DNA. In both G1 and metaphase, a subset of canonical nucleosome classes
has asymmetric linker DNA structure. Both G1 and metaphase chromatin have ordered
stacked dinucleosomes classes, but no oligomers with more than two ordered
nucleosomes. In detecting these ordered dinucleosomes, we show how to bootstrap a
classification workflow to detect oligomers with ordered subunits. Remapping analysis
has shown that the additional nucleosomes interact with both other nucleosomes and
non-nucleosome complexes at their faces and gyres, and that these interactions are

variable as evidenced in the absence of unambiguous class average densities.
Metaphase canonical nucleosomes are uniformly distributed: there are no chromatin
domains within, which raises the question of how compaction and decompaction
proceed in terms of interphase chromatin domains. The structural cell-biology
phenotypes presented here give multiple starting points for further nanoscale analysis
and additional hypotheses about interphase and mitotic human nuclei and chromatin *in*
situ.

MATERIALS AND METHODS

Cell culture

hTERT-RPE-1 cells (human retina epithelium cells immortalized with hTERT; CRL-4000™) were obtained from ATCC (Manassas, VA). The cells were grown in an incubator set to 37°C and 5% CO_2. The cells were cultured in complete medium consisting of DMEM/F-12 with GlutaMAX™ medium (Gibco, 10565-018; Thermo Fisher Scientific (TFS), Waltham, MA) supplemented with 10% fetal bovine serum (Sigma, F9665) and 1% penicillin-streptomycin (Pen-strep) (Gibco, 15140-122). Cell washing was done with PBS+Ca+Mg, consisting of Phosphate-buffered saline (PBS, Vivantis, PB0344-1L; Vivantis Technologies Sdn. Bhd., Selangor Darul Ehsan, Malaysia), 0.1 mg/mL $CaCl_2$ (Sigma, 12095), and 0.1 mg/mL $MgCl_2$ (Sigma, M8266-100G). Details of all the reagents and equipment used are summarized in Table S1.

Taxol-arrested cells for pilot experiments

Taxol-arrested cells were prepared following our previous work (Ruan *et al*, 2019). The cells were first synchronized in G1 as follows. Cells were seeded in culture flasks or dishes and incubated for 24 hours. They were then subject to depleted serum culturing for 24 hours. The cells were then further treated with 3mM thymidine for another 24 hours. To arrest the cells in mitosis, they were released from the thymidine block for 4 hours by replacement of fresh culture medium. Next, the cells were incubated with 800 μM Taxol for 12 hours. The mitotic cells were collected by shake-off.

G1 and metaphase cell-cycle arrest for both LM and cryo-ET of lamellae

The cell-cycle synchronization was done following a published protocol (Scott *et al*., 2020), with modifications. Adherent cells (destined for G1 arrest) were prepared differently for confocal microscopy versus cryo-ET. For confocal experiments, RPE-1 cells were seeded in a 35 mm μ-dish at a density of 4.8×10^5 cells/mL. After a 24-hour incubation, 1000× palbociclib (Selleckchem, S1116; Selleck Chemicals, Houston, TX), dissolved in 0.22 μm filtered Milli Q water, was added to the medium to a final concentration of 1 μM and then incubated for a further 18 hours. For cryo-ET experiments, gold grids were placed on the cover of a 35 mm Petri dish and exposed to UV light for 15 minutes in a biosafety cabinet. Then the gold grids were washed by incubating 10 minutes in complete medium + Pen-strep in a 35 mm dish, in a shaker at 20 RPM. The complete medium was then discarded and replaced with fresh medium. To arrest cells in G1 phase, RPE-1 cells were seeded in a 35 mm Petri dish where two gold grids were positioned in advance at a density of 2.0×10^5 cells/mL. After a 24-hour incubation, 1000× palbociclib was added to the medium to a final concentration of 1 μM and then followed by an 18-hour incubation. Then, the grids were picked up and

991   immersed in DMEM/F-12 with GlutaMAX™ medium containing 9% DMSO (Sigma,
992   472301) for 5 seconds as cryoprotectant before plunge-freezing (see below).
993

Metaphase cells were synchronized the same way for all experiments (confocal
microscopy, immunoblots, cryo-ET). RPE-1 cells were plated into a 25 cm^2 flask at a
density of 8.0×10^4 cells/mL. After a 24-hour incubation, palbociclib was added to a final
concentration of 1 µM and the cells were then incubated for 18 hours (as above). The
medium was then pipetted out and the cells were rinsed three times with 1 mL pre-
warmed PBS+Ca+Mg. The cells were then cultured in fresh complete medium for 8
hours. Then nocodazole (Sigma, SML1665) (5 mg/mL stock in DMSO) was added into
the medium to a final concentration of 50 ng/mL. After a 12-hour incubation, the cells
were harvested by both shaking the flask and tapping it horizontally against the bench.
Next, the cell suspension medium was washed four times by centrifugation at 1000 $\times$ g
for 3 minutes, decanting the supernatant, and cell-pellet resuspension with 3 mL
PBS+Ca+Mg. The last resuspension was done in complete culture medium with 10 µM
MG132 (Sigma, M7449). After a 70-minute incubation, the cells were detached by
shake off and flask tapping (see above), centrifuged at 1000 $\times$ g for 3 minutes. The cell
pellet is used for the subsequent experiments.
1009

Immunoblots
Both nocodazole- and MG132-arrested cells were prepared as above. Nocodazole-
arrested cells were equally divided into four tubes, corresponding to: no washout and
30, 60, and 120 minutes after nocodazole washout. In a separate experiment, MG132-
arrested cells were also equally divided into four tubes, corresponding to: no washout
and, 30, 60, and 120 minutes after MG132 washout. The cell washes were done with
PBST that was pre-warmed to 37°C in a water bath. Time zero of the nocodazole-
washout experiment corresponds to when the first wash started. All cells were washed
with ice-cold PBST before the addition of lysis buffer. Ice-cold protein extraction lysis
buffer was composed of RIPA (Sigma, R0278), Protease inhibitor cocktail (Sigma,
P8340) and Phosphatase Inhibitor Cocktail 3 (Sigma, P0044) with a volume ratio of
100:1:1. One hundred µL lysis buffer was added per 1.0×10^6 cells and mixed by
vortexing for 6 seconds. In total, 5 vortex cycles were done with a 6 minute wait
between each vortex. The lysate mixture was centrifuged at 16,000 $\times$ g for 20 minutes.
The supernatant was aspirated into a clean tube and stored in −80°C until all groups
were ready for total protein concentration determination via BCA assay (TFS, 23227).
An equal volume of 2✕ Laemmli buffer was added into the cell lysates and mixed by
vortexing briefly for less than 5 seconds. The mixture was boiled at 95°C for 5 minutes,
then centrifuged at 16,000 $\times$ g for 5 minutes. The denatured protein mixture was divided
into aliquots of about 20 µg for either immediate electrophoresis or storage in −80°C.
1030

Twenty µg of protein was loaded in each well of a Mini-protean TGX gel (Bio-Rad
4561083; Hercules, CA) with Precision Plus Protein Dual Color Standards (Bio-Rad
1610374) or MagicMark™ XP Western Protein Standard (TFS, LC5602) as the protein
ladder. Electrophoresis was done using constant voltage at 90 V for 30 minutes,
followed by 120 V for another 30 minutes until the blue dye almost reached the bottom
of the gel. The ladder and proteins in the gel were transferred to the PVDF membrane
(Bio-Rad 1704156) by the semi-dry method, with constant current 2.5 A for 3 minutes.
The membrane was blocked in 1× TBST (Tris-buffered saline with 0.1% Tween 20)
containing 5% nonfat milk at 23°C for 1 hour. Next, the membrane was rinsed with
TBST for 5 seconds, then incubated with primary rabbit polyclonal antibody to beta
tubulin (Abcam ab6046; Cambridge, United Kingdom), rabbit polyclonal anti-phospho-
histone H3 (Ser10) antibody (Abcam, ab5176), or cyclin B1 (GNS1) mouse monoclonal
IgG1 (Santa Cruz, sc-245; Dallas, TX) overnight at 4°C. All antibodies were diluted
1:500 in blocking buffer. After three sequential 5-minute washes with TBST, the
membrane previously incubated with anti-Cyclin B1 antibody was incubated with m-
IgGκ BP-HRP secondary antibody (Santa Cruz, sc-516102) for 1 hour at 23°C, and
membranes previously incubated with anti-beta tubulin antibody or anti-phospho-histone
H3 (Ser10) antibody were incubated with mouse anti-rabbit IgG-HRP secondary
antibody (Santa Cruz, sc-2357) for 1 hour at 23°C. After 3 sequential 5-minute washes
in TBST, the membranes were incubated with Clarity Western ECL substrate solution
(Bio-Rad 1705061) at 23°C for 5 minutes, then imaged with ImageQuant LAS4000
system (Cytiva, Marlborough, MA). Full, uncropped blots are shown in Figure S36. A
summary of all the antibodies used is in Table S2.
1054
Apoptosis control experiments
RPE1 cells were incubated in complete medium containing 1 µM staurosporine for 6
hours to induce apoptosis as a positive control. To control for the carrier, another group
of RPE1 cells was incubated in complete medium containing 0.1% DMSO for 6 hours.
Five more flasks of RPE1 cells were arrested in G1 phase. One flask did not have any
cryoprotectant added and was used as the negative control while the other four flasks
were treated with cryoprotectants. To test for apoptosis induction by cryoprotectants,
flasks of G1 cells was incubated with either 4 mL 9% DMSO or 4 mL 9% glycerol in
DMEM/F12 medium. These treatments lasted either 1 minute or 10 minutes and were
stopped by discarding the cryoprotectant-containing medium and rinsing with 2 mL cold
PBS 3 times. For apoptosis analysis in metaphase cells, five flasks of metaphase-
arrested RPE1 cells were aliquoted into 15 mL tubes. The cells destined for
cryoprotectant treatment were centrifuged at 200 $\times$ g for 3 minutes, then the
supernatant was removed via pipetting. Next the cell pellets were resuspended in 0.5
mL DMEM/F12 medium containing either 9% DMSO or 9% glycerol. These cells were
incubated for either 1 minute or 10 minutes. The cryoprotectant was achieved by

dilution with 5 mL PBS followed by three washes with 2 mL cold PBS. After the washes,
the proteins were extracted and analyzed by immunoblots with rabbit monoclonal
antibodies against PARP1 (Abcam, ab191217) and with immunoblots against beta
tubulin as the loading control.
1075

Conventional fluorescence microscopy
For morphological analysis, RPE-1 cells were imaged using a Nikon Eclipse Ti
microscope (Nikon Instruments Inc., Tokyo, Japan) equipped with a Plan Fluor 10× Ph1
objective lens. To obtain fields of view of multiple cells, Z-stack images were acquired
on an LSM 900 AiryScan microscope controlled by ZEN 3.1 software, using a 20× 0.8
NA Plan-Apochromat under confocal mode (Zeiss, Jena, Germany). Z-projection at
maximum intensity was done in FIJI (Schindelin *et al*, 2012).
1083

Immunofluorescence
All washes were done as follows. Attached cells were washed 3 times with PBST
(PBS+Ca+Mg with 0.1% Tween 20 (Sinopharm, T20087687, Beijing, China)) for 5
minutes on a shaker at 20 RPM. Suspended cells were washed 3 times with PBST for 5
minutes, except the first wash after fixation (see fixation protocol below), in which PBST
was replaced with PBST containing 0.5% BSA (Sigma, A9647). For each wash, the
suspended cells were centrifuged at 600 × g for 3 minutes, then supernatant was
decanted, the cell pellet was resuspended with 1 mL PBST using a P1000 micropipette,
and then incubated 2 minutes on a rotator at 20 RPM.
1093

Palbociclib-synchronized G1 (attached) cells were washed three times with
PBS+Ca+Mg after the completion of the drug treatment. The cells were then fixed with
4% paraformaldehyde (Electron Microscopy Sciences, 15714, Hatfield, PA) in PBS for
10 minutes at 23°C, then washed 3 times for 5 minutes each in PBST, then
permeabilized with 0.25% Triton X-100 (TFS, Alfa Aesar A16046) in PBS for 10 minutes
at 23°C. To block non-specific binding, the cells were then washed 3 × 5 minutes with
PBST, then incubated with PBST containing 1% BSA and 22.5 mg/mL glycine (Bio-Rad
1610718) for 30 minutes at 23°C. Next, the cells were immunostained with primary
antibodies: rabbit polyclonal to RNAPII CTD repeat YSPTSPS (phospho S2) (1:300,
Abcam ab5095), rabbit polyclonal to RNAPII CTD repeat YSPTSPS (phospho S5)
(1:300, Abcam ab5131)), or Alexa Fluor-conjugated antibody (Alexa Fluor® 647) mouse
monoclonal anti-histone H3 (phospho S10) antibody (1:250, Abcam ab196698)), in
PBST with 1% BSA, on a rotator with slow rotation, overnight at 4°C. The cells were
then incubated with Alexa Fluor 488 goat anti-rabbit IgG (H+L) secondary antibody
(1:500, TFS A-11008) in Milli Q purified water containing 1% BSA for 1 hour at 23°C.
Finally, the cells were incubated with 300 nM 4′,6-diamidino-2-phenylindole (DAPI, TFS

D1306) in PBS for 5 minutes. After three sequential 5-minute washes with PBST, the
cells were ready for imaging.
1112
To harvest nocodazole- and MG132-arrested cells after the completion of the drug
treatment, the flasks were shaken and tapped against the bench. The detached cells
were collected in a 15 mL tube and centrifuged at 600 $\times$ g for 3 minutes. Cell pellets
were resuspended with 1 mL PBST and transferred to a 1.5 mL microfuge tube for all
following incubations and washes, which are different from the G1 cells (attached to the
dish). After the last wash before imaging, most of the supernatant was removed, leaving
about 20 µL. The cell pellets were resuspended in this remaining supernatant with a
micropipette using a 200 µL tip. Finally, 5 µL of this cell suspension was mounted on a
20 mm Ø coverslip (Paul Marienfeld GmbH & Co. KG, 0112600, Lauda-Königshofen,
Germany) with 15 µL Prolong Gold antifade reagent and then cured 20 to 24 hours prior
to imaging. All treatments between the harvesting and blocking steps were done in 15
mL conical tubes. After blocking, the washing was done in 1.5 mL microfuge tubes. All
reagents including their concentrations and incubation durations were the same as
above G1 cells.
1127
To obtain a large field of view containing multiple cells, Z-stacks were acquired with a
Zeiss LSM 900 confocal microscope (see above). To observe details of single cells, Z-
stacks were acquired with the same microscope, but using a 63$\times$ 1.4 NA Plan-
Apochromat oil-immersion objective and operated in Airyscan super-resolution mode. Z-
stacks (3-D) were projected using Fiji's "3D Project" brightest point method. All the
confocal microscopy details are summarized in Table S3.
1134
Self-pressurized freezing
Taxol-treated RPE-1 cells were scraped or shaken off in 1$\times$ PBS. The cells were further
pelleted by centrifugation at 600 $\times$ g for 10 minutes. For 100 µL cell pellets, 25 µL 50%
w/v dextran 6000 (Sigma 31388) was added and mixed gently as the extracellular
cryoprotectant, yielding a final concentration of 12.5% dextran. The cells were loaded
into copper tubes, which were sealed and then dropped into liquid ethane for self-
pressurized freezing.
1142
Vitreous sectioning
Cryomicrotomy was done in an UC7/FC7 cryomicrotome (Leica Microsystems, Vienna,
Austria). RPE-1 cells were cut into 100-nm-thick (nominal) cryosections using a 35°
diamond knife (Diatome Cryo35, Nidau, Switzerland). We used the dual-
micromanipulator method (Ladinsky *et al.*, 2006; Ng *et al.*, 2020; Studer *et al.*, 2014).
The cryosections were attached to a continuous carbon grid (EMS CF200-Cu-UL,
Hatfield, PA) using a Crion device (Leica) in "charge" mode.

1150

Plunge-freezing of oligonucleosomes

1151
HeLa oligonucleosomes (EpiCypher 16-0003, Durham, NC; sold as "polynucleosomes")
in storage buffer (20 mM HEPES, pH 7.5, 1 mM EDTA) was diluted 1:1 in a buffer
containing 20 mM HEPES-KOH, pH 7.8, 2 mM $MgCl_2$. The additional $MgCl_2$ was added
to increase the local oligonucleosome concentration to facilitate targeting but was of
insufficient concentration; since the same $MgCl_2$-containing buffer was used in all the
cryoprotectant test samples, the conclusions were independent of $MgCl_2$. For
preparations containing cryoprotectant, the sample was also supplemented with DMSO
to a final concentration of 9% (v/v). Each glow-discharged Quantifoil R2/4 200 mesh
copper grid (Quantifoil Micro Tools GmbH, Jena, Germany) was then deposited with 4
µL sample and plunge-frozen in liquid ethane using a Vitrobot Mark IV (TFS) operated
at 22°C and 100% humidity.

1163

Plunge-freezing of RPE-1 cells

1164
RPE-1 cells were plunge-frozen in 63/37 propane-ethane mixture (Tivol *et al*, 2008)
using a Vitrobot Mark IV (TFS) operated at 100% humidity and 37°C. For G1-arrested
RPE-1 cells that were grown on C-flat 2/4 200 mesh gold grids (Protochips, Morrisville,
NC), the grid was held with a Vitrobot tweezer and then immersed for 5 seconds in
DMEM/F12 with GlutaMAX™ medium that was supplemented with either 9% DMSO or
9% glycerol as cryoprotectant. For the no-cryoprotection control, the grid was immersed
in fresh DMEM/F12 with GlutaMAX™ only. The grid was then loaded on the Vitrobot
and manually blotted from the back using Whatman® Grade 1 filter paper before being
plunge-frozen. Note that in the earlier screening experiments to find the optimal
cryoprotectant concentration (Figure S4), the grids were immersed in PBS+Ca+Mg that
were supplemented with 0 – 9% DMSO. Subsequent sample preparation was
performed using DMEM/F12 with GlutaMAX™ instead (see below).

1177

Metaphase-arrested RPE-1 cells were first harvested as a suspension and concentrated
to 1.5×10^6 cells/mL. Immediately before plunge freezing, the cell suspension was mixed
with an equal volume of fresh DMEM/F12 with GlutaMAX™ medium containing either
18% DMSO or 18% glycerol, bringing the final DMSO (or glycerol) concentration to 9%.
For the no-cryoprotection control, the cells were resuspended in fresh DMEM/F12 with
GlutaMAX™ only. 4 µL of the cell suspension was then deposited onto C-flat 2/4 200
mesh gold grids (without any coatings) that were glow discharged at 15 mA for 15
seconds, before being manually back-blotted and plunge-frozen as described above.

1186

Cryo-FIB milling

1187
Cryolamellae of plunge-frozen RPE-1 cells were prepared using a Helios NanoLab 600
DualBeam (TFS) equipped with a PolarPrep 2000 Cryo Transfer System (Quorum

Technologies, Laughton, UK) (Medeiros *et al.*, 2018). The samples were first coated
with a layer of organometallic platinum using an in-chamber gas injection system that
was positioned 9 mm from the grid center and operated for approximately 10 to 18
seconds flow time at 28°C (Hayles *et al.*, 2007). Cryo-FIB milling was then performed
with the Helios stage tilted to 12°. The bulk of the cellular material was first removed
using the FIB at 30 kV 2.8 nA, followed by further thinning of the cryolamellae at a lower
current of 0.28 nA. The cryolamellae were then polished to a final nominal thickness of
~150 nm at 48 pA.
1198
Electron cryotomography
A summary of the cryo-ET sample preparation, imaging, and processing parameters are
reported in Table S4. Additional details of each tilt series are reported in Table S5.
Cryo-ET data of both RPE-1 cryolamellae and purified HeLa oligonucleosome samples
were collected using SerialEM (Mastronarde, 2005) on a Titan Krios (TFS) equipped
with a K3 camera (AMETEK, Berwyn, PA) and a BioContinuum energy filter (AMETEK)
operating in zero-loss mode with a 20 eV slit width. Tilt series were acquired in super-
resolution mode with a bin ×1 pixel size of 3.4 Å at the specimen level. These data were
collected in the dose-symmetric tilt scheme and using either defocus phase contrast at
−5 µm defocus or Volta phase contrast at −0.5 µm defocus (both nominal values). The
K3 data was collected after TFS changed their Volta phase plate design, requiring that
the phase-plate heater be operated at ~370 mW power.
1211
To speed up the setup of the tilt series collection on SerialEM, we used the Python
scripts described by Weis *et. al* (Weis *et al*, 2021) to generate virtual maps of regions
interest from low-magnification montage images of cryolamellae or grid squares. HeLa
oligonucleosome samples were imaged using a tilt range of ±60° and a tilt increment of
2°, for a nominal total dose of 120 e$^-$/Å^2 per tilt series. For cryolamellae samples, a
stage pre-tilt of −14° was applied so that the cryolamellae are approximately
perpendicular to the electron-optical axis when the first image is recorded. This pre-tilt
angle was determined by trial-and-error. Cryolamellae tilt series were collected using a
tilt range of −70° to +42° and a 2° tilt increment. The nominal total dose per tilt series for
cryolamellae samples was 110 e$^-$/Å^2.
1222
To more accurately target regions of interest for cryolamellae imaging, we first
generated longer exposure (~0.1 e$^-$/Å^2 per tile) low-magnification montage images of
our samples. Montage tiles were acquired at −90 µm defocus using an unbinned pixel
size of 19.2 Å (4,800× magnification) and a 2-second exposure time. In this imaging
condition, chromosome regions and cytoplasm of metaphase-arrested RPE-1 cells
show different density features (Figure S28). Microtubules are also frequently seen
surrounding the chromosome regions. For G1-arrested cells, nuclear regions were

identified based on cytological cues. The nucleus periphery regions were found based
on the presence of the nuclear envelope. If the nuclear envelope was not visible,
imaging positions were selected based on the absence of the cytoplasmic components
listed above.
1234
Super-resolution movie frames were aligned and Fourier cropped to 3.4 Å pixel size
using MotionCor2 (Zheng *et al*, 2017). Tilt series stacks were aligned by patch tracking
and reconstructed into cryotomograms at 6.8 Å voxel size using the IMOD program
Etomo (Kremer *et al*, 1996; Mastronarde, 1997; Mastronarde & Held, 2017). All
subsequent template matching and subtomogram analyses were performed using
cryotomograms at 6.8 Å voxel size.
1241
Automated segmentation of chromatin regions in the cryotomograms was performed
using EMAN2 (Chen *et al.*, 2017). For the convolutional neural network training, 64 x 64
tiles containing either G1 chromatin domains or metaphase chromosome regions were
extracted from the cryotomograms (binned to a voxel size of 13.5 Å) and used as
positive training examples. Tiles from the cytoplasm, nucleoplasm and lamella surface
contaminants were used as negative examples.
1248
For the cryosection pilot experiments, both defocus phase-contrast and Volta phase
contrast tilt series were collected using the same Titan Krios as above, but with a
Falcon II camera (TFS) in integration mode at a 12.4 or 7.3 Å pixel size as the specimen
level, all controlled with Tomo4 (TFS). A nominal defocus of −10 μm (defocus phase
contrast) or −0.5 μm (VPP) was applied. Bi-directional tilt series, starting at 0°, were
collected with a ±60° tilt range and 2° tilt increment. These data were collected before
TFS implemented a change in Volta phase plate design, so the phase-plate heater was
operated at lower than 150 mW power.
1257
Denoising for visualization
A subset of cryotomograms were denoised for visualization purposes; the subtomogram
analysis steps in the following sections were done on non-denoised data. MotionCor2
was used to generate tilt series images where either the odd- or even-numbered raw
movie frames were excluded. The odd- and even-frames only tilt series stacks were
then reconstructed at 6.8 Å voxel size in IMOD and used as the input for denoising in
cryo-CARE (Buchholz *et al*, 2019). The default parameters were used for cryo-CARE,
with the exception of the "n_tiles" value in the prediction settings which was changed to
5. All cryotomogram slices shown in Figures 2 – 5 were created from denoised data. A
copy of these figures was also created using non-denoised data and can be found as
Figures S37 – S40. Figures S17 and S20 were also created from denoised data.
1269

Template matching

Candidate nucleosome subtomograms were picked from bin ×2 cryotomograms (6.8 Å pixel size; low-pass filtered using *mtffilter* with mean and sigma values of 0.17 and 0.05, respectively, with the roll-off starting at 40 Å resolution) by template matching using the PEET package (Heumann, 2023; Heumann *et al*, 2011; Nicastro *et al*, 2006). A featureless soft-edged cylinder ~10 nm diameter and ~5.5 nm tall was generated as the template-matching reference using the Bsoft program *beditimg* (Heymann, 2022). Cryotomograms of interest were seeded with a grid of points (from which translational searches initialize) that were spaced ~7 nm apart using the PEET program *gridInit*. For efficiency, we excluded seeded points that were located outside chromatin domains or metaphase chromosome regions by using the segmented version of the cryotomograms as masks. At the end of the PEET template matching run, duplicate removal was performed on the candidate points, with a distance threshold of ~5.5 nm.

Globular complexes such as ribosomes and preribosomes were picked using PEET, as described above with three differences: 1) A featureless sphere of ~24 nm diameter was used as the reference. 2) The initial grid of points seeded were ~21 nm apart. 3) Duplicate removal was performed with a 13.5 nm distance threshold.

The coordinates of the template matching subtomograms (candidate nucleosomes/ribosomes/preribosomes) were then exported to RELION for subtomogram averaging, classification, and refinement. The remaining "junk" subtomograms were excluded at the end of each classification run.

Subtomogram analysis of mononucleosomes

Classification and subtomogram averaging were performed using RELION 3 (Kimanius *et al*, 2016; Zivanov *et al*, 2018). A summary of the workflow is shown in Figures S14 and S30. 64×64×64 voxel subtomograms corresponding to the template-matching coordinates in PEET were extracted in RELION and subjected to 3-D classification. To analyze for mononucleosomes, the subtomograms were classified using a featureless cylinder as the reference and a 110 Å-diameter spherical mask (Figures S14A & S30A).

To analyze for ordered stacked dinucleosomes, the same set of extracted subtomograms from the mononucleosome PEET run were subjected to 3-D classification, using a double-stacked featureless cylinder as the reference and a larger 240 Å-diameter spherical mask. The resulting nucleosome classes from the double-cylinder 3-D classification run were selected and subjected to a second round of 3-D classification, where an ordered stacked dinucleosome class was observed (Figures S14 & S30, B & C). In an attempt to pick up more nucleosome particles, the original set of extracted subtomograms were again subjected to 3-D classification, this time using

the ordered stacked dinucleosome average as the reference, low-pass filtered to 60 Å in
RELION (Figures S14D & S30D). Nucleosome averages from this classification run
were manually placed into three groups: 1) mononucleosomes; 2) ordered stacked
dinucleosomes; and 3) mononucleosomes with an additional proximal density.
Subtomograms from each of these three groups were subjected to a second round of 3-
D classification separately (Figures S14E & S30E). For Group 1, the resulting
mononucleosome classes were manually separated into two further groups
corresponding to the presence of either short or long linker DNA densities and then
subjected to 3-D refinement separately. 3-D classification for Group 2 resulted in
several ordered stacked dinucleosome classes, which were refined as a single group.
For Group 3, we detected several mononucleosome classes with an additional density
at different positions around the DNA gyres. Each of these Group 3 classes were
subjected to separate 3-D refinements (Figures S15 and S31). The details are
summarized in Table S6 and S7.
1324

Subtomogram analysis of other nucleosome and multi-nucleosome structures
To detect the presence of ordered side-by-side dinucleosomes and stacked tri-/tetra-
nucleosomes, further 3-D classification was performed on the nucleosome particles in
Groups 2 and 3 (Figures S15 and S31, panels B and C), respectively. Custom masks
were generated to include the nucleosome densities seen in the subtomogram
averages; and to enclose additional volume where we anticipate more nucleosomes
may be found. Figures S16 and S32 show a rendering of the reference and mask
combination used for each of the 3-D classification runs. These custom masks were
created in UCSF Chimera (Pettersen *et al*, 2004) by manually positioning multiple
featureless cylinders of 15 nm-diameter and 10 nm-height, and then combining them
into a single volume using the following Chimera commands:

```
1336 vop new empty size 64 gridSpacing 6.766
1337 vop maximum [cyl #1] [cyl #2] [cyl #3] onGrid [empty #]
```

1338

1339   The multi-cylinder masks' edges were then smoothed by Gaussian filtering with the
1340   command:

```
1341 vop gaussian [combined cylinder #] sDev 67.66
```

1342

1343   References used during these 3-D classification runs were subtomogram averages
1344   generated during the refinement step from either Group 2 or 3, all low-pass filtered to 60
1345   Å in RELION. The angular search range was restricted to ±15° for these 3-D
1346   classification runs.
1347

Subtomogram analysis of ribosomes and preribosomes
Template matching hits for candidate ribosomes in the cytoplasm were subjected to
direct 3-D classification using a 24-nm-diameter sphere as a reference. The

unambiguous ribosome class averages were pooled and then subjected to refinement.
For preribosome subtomogram averaging, template matching hits from both the nuclear
region and the cytoplasm was included during RELION classification. Candidate
preribosomes were first 2-D classified, whereupon classes that did not have large
complexes were removed. Next, 3-D classification was performed on the remaining
particles, using the refined ribosome class average generated in the previous step as a
reference. An initial low-pass filter of 60 Å was applied to the ribosome reference in
RELION, during the 3-D classification run.
1359
Docking of nucleosome crystal structures
The docking of nucleosome crystal structures into our refined subtomogram averages
was performed using UCSF Chimera. The nucleosome crystal structure (PDB 1AOI)
(Luger *et al.*, 1997) was docked into nucleosome subtomogram averages that have
short linker DNAs. The chromatosome crystal structure (PDB 5NL0) (Garcia-Saez *et al.*,
2018) was docked into subtomogram averages with longer linker DNAs. The histone
tails were removed from the atomic models prior to docking into our subtomogram
averages. Briefly, the corresponding atomic model was manually aligned into the
subtomogram average, and then the fit-in-map function was executed. For the fit-in-map
function, the option to simulate a density map from the atomic model at 20 Å was used.
1370
For the stacked dinucleosome averages, the Chimera "hide dust" function was used to
hide the nucleosome density with the shorter linker DNA. PDB 5NL0 was then fitted to
the remaining visible nucleosome. The hidden nucleosome density was next rendered
visible using the command "`sop invertShown`", and then docked by PDB 1AOI. For
the mononucleosome subtomogram averages that have an additional proximal density,
PDB 1AOI was fitted to the nucleosome region, with the proximal density still rendered.
1377
The fitting correlation coefficients for the G1 nucleosomes subtomogram averages were
as follows: mononucleosome, short linker = 0.78; mononucleosome, long linker = 0.78;
stacked dinucleosome, nucleosome 1 = 0.71; stacked dinucleosome, nucleosome 2 =
0.67; nucleosome with proximal densities = 0.65 – 0.73. For the metaphase nucleosome
subtomogram averages: mononucleosome, short linker = 0.77; mononucleosome, long
linker = 0.77; stacked dinucleosome, nucleosome 1 = 0.68; stacked dinucleosome,
nucleosome 2 = 0.66; nucleosome with proximal densities = 0.70 – 0.72.
1385
1386 **Simulation of ribosome densities**
1387 Coordinates for the single-particle reconstruction of human ribosomes (PDB 4UG0)
1388 (Khatter *et al.*, 2015) were used as input for density map simulations. The EMAN2
1389 program *e2pdb2mrc.py* was used to create an .mrc file using the following command:

```
e2pdb2mrc.py --apix 6.766 --box 64 --res [res] --center 4UG0.pdb
4UG0.mrc
```
The option [res] corresponds to the resolution of the output map. Three maps were created, covering resolutions (25 – 35Å) typical of averages of the order of ~1,000 subtomograms, such as the one obtained from the G1 cell cryotomograms.

Remapping

Remapping of nucleosome, ribosome, and preribosome particles was performed using a custom python script *ot_remap_v2.py* (available at https://github.com/anaphaze/ot-tools). The coordinate and orientation information for each particle from the RELION refined .star files was imposed on the corresponding subtomogram average, which was then written into a new .mrc file. Remapped models were visualized using UCSF Chimera.

Data sharing

A copy of all the tilt series raw data and cryotomograms presented in this paper (Table S5) was deposited in EMPIAR (Iudin *et al*, 2016) as EMPIAR-#####. Subtomogram averages of the RPE-1 nucleosomes, dinucleosomes, nucleosomes with gyre-proximal densities (Figures 3A – C and 5A – C), cytoplasmic ribosomes (Figure S11B), and preribosomes (Figure S12B) have been deposited at EMDB. The EMD serial numbers are shown in Tables S6 – S9.

Contributions

JC, TL, SC, WR, CTN - experiments, project design, writing; US - project design, writing; JS - training; LG - project design, writing, training.

ACKNOWLEDGEMENTS

We thank the CBIS microscopy staff for support and training. We thank Paulo D'Avino for advice on RPE-1 synchronization. JC, TL, and LG were supported by a Singapore Ministry of Education grant MOE2019-T2-1-140. WR and US were funded by the Biomedical Research Council of A*STAR (Agency for Science, Technology and Research), Singapore.

REFERENCES

Arimura Y, Shih RM, Froom R, Funabiki H (2021) Structural features of nucleosomes in interphase and metaphase chromosomes. *Mol Cell* 81: 4377-4397 e4312

Armache JP, Gamarra N, Johnson SL, Leonard JD, Wu S, Narlikar GJ, Cheng Y (2019) Cryo-EM structures of remodeler-nucleosome intermediates suggest allosteric control through the nucleosome. *Elife* 8

Bacic L, Gaullier G, Sabantsev A, Lehmann LC, Brackmann K, Dimakou D, Halic M, Hewitt G, Boulton SJ, Deindl S (2021) Structure and dynamics of the chromatin remodeler ALC1 bound to a PARylated nucleosome. *Elife* 10

Baker RW, Reimer JM, Carman PJ, Turegun B, Arakawa T, Dominguez R, Leschziner AE (2021) Structural insights into assembly and function of the RSC chromatin remodeling complex. *Nat Struct Mol Biol* 28: 71-80

Barth R, Bystricky K, Shaban HA (2020) Coupling chromatin structure and dynamics by live super-resolution imaging. *Sci Adv* 6

Bäuerlein FJB, Pastor-Pareja JC, Fernández-Busnadiego R (2022) Cryo-electron tomography of native Drosophila tissues vitrified by plunge freezing. *bioRxiv*

Bednar J, Garcia-Saez I, Boopathi R, Cutter AR, Papai G, Reymer A, Syed SH, Lone IN, Tonchev O, Crucifix C *et al* (2017) Structure and Dynamics of a 197 bp Nucleosome in Complex with Linker Histone H1. *Mol Cell* 66: 384-397 e388

Bednar J, Horowitz RA, Grigoryev SA, Carruthers LM, Hansen JC, Koster AJ, Woodcock CL (1998) Nucleosomes, linker DNA, and linker histone form a unique structural motif that directs the higher-order folding and compaction of chromatin. *Proc Natl Acad Sci U S A* 95: 14173-14178

Beel AJ, Azubel M, Mattei PJ, Kornberg RD (2021) Structure of mitotic chromosomes. *Mol Cell* 81: 4369-4376 e4363

Bertrand R, Solary E, O'Connor P, Kohn KW, Pommier Y (1994) Induction of a common pathway of apoptosis by staurosporine. *Exp Cell Res* 211: 314-321

Bilokapic S, Strauss M, Halic M (2018) Cryo-EM of nucleosome core particle interactions in trans. *Sci Rep* 8: 7046

Booth DG, Beckett AJ, Molina O, Samejima I, Masumoto H, Kouprina N, Larionov V, Prior IA, Earnshaw WC (2016) 3D-CLEM Reveals that a Major Portion of Mitotic Chromosomes Is Not Chromatin. *Mol Cell* 64: 790-802

Booth DG, Takagi M, Sanchez-Pulido L, Petfalski E, Vargiu G, Samejima K, Imamoto N, Ponting CP, Tollervey D, Earnshaw WC *et al* (2014) Ki-67 is a PP1-interacting protein that organises the mitotic chromosome periphery. *Elife* 3: e01641

Bowerman S, Wereszczynski J, Luger K (2021) Archaeal chromatin 'slinkies' are inherently dynamic complexes with deflected DNA wrapping pathways. *Elife* 10

Buchholz TO, Krull A, Shahidi R, Pigino G, Jekely G, Jug F (2019) Content-aware image restoration for electron microscopy. *Methods Cell Biol* 152: 277-289

Cai S, Böck D, Pilhofer M, Gan L (2018a) The in situ structures of mono-, di-, and trinucleosomes in human heterochromatin. *Mol Biol Cell* 29: 2450-2457

Cai S, Chen C, Tan ZY, Huang Y, Shi J, Gan L (2018b) Cryo-ET reveals the macromolecular reorganization of S. pombe mitotic chromosomes in vivo. *Proc Natl Acad Sci U S A* 115: 10977-10982

Cai S, Song Y, Chen C, Shi J, Gan L (2018c) Natural chromatin is heterogeneous and self-associates in vitro. *Mol Biol Cell* 29: 1652-1663

Cai S, Wu Y, Guillen-Samander A, Hancock-Cerutti W, Liu J, De Camilli P (2022) In situ architecture of the lipid transport protein VPS13C at ER-lysosome membrane contacts. *Proc Natl Acad Sci U S A* 119: e2203769119

Chan DS, Kavanagh ME, McLean KJ, Munro AW, Matak-Vinkovic D, Coyne AG, Abell C (2017) Effect of DMSO on Protein Structure and Interactions Assessed by Collision-Induced Dissociation and Unfolding. *Anal Chem* 89: 9976-9983

Chen C, Lim HH, Shi J, Tamura S, Maeshima K, Surana U, Gan L (2016) Budding yeast chromatin is dispersed in a crowded nucleoplasm in vivo. *Mol Biol Cell* 27: 3357-3368

Chen M, Dai W, Sun SY, Jonasch D, He CY, Schmid MF, Chiu W, Ludtke SJ (2017) Convolutional neural networks for automated annotation of cellular cryo-electron tomograms. *Nature methods* 14: 983-985

Chen Q, Zhao L, Soman A, Arkhipova AY, Li J, Li H, Chen Y, Shi X, Nordenskiold L (2022a) Chromatin Liquid-Liquid Phase Separation (LLPS) Is Regulated by Ionic Conditions and Fiber Length. *Cells* 11

Chen X, Wang X, Liu W, Ren Y, Qu X, Li J, Yin X, Xu Y (2022b) Structures of +1 nucleosome-bound PIC-Mediator complex. *Science* 378: 62-68

Chicano A, Crosas E, Oton J, Melero R, Engel BD, Daban JR (2019) Frozen-hydrated chromatin from metaphase chromosomes has an interdigitated multilayer structure. *EMBO J* 38

Chittori S, Hong J, Bai Y, Subramaniam S (2019) Structure of the primed state of the ATPase domain of chromatin remodeling factor ISWI bound to the nucleosome. *Nucleic Acids Res* 47: 9400-9409

Cho NH, Cheveralls KC, Brunner AD, Kim K, Michaelis AC, Raghavan P, Kobayashi H, Savy L, Li JY, Canaj H *et al* (2022) OpenCell: Endogenous tagging for the cartography of human cellular organization. *Science* 375: eabi6983

Cho RJ, Campbell MJ, Winzeler EA, Steinmetz L, Conway A, Wodicka L, Wolfsberg TG, Gabrielian AE, Landsman D, Lockhart DJ *et al* (1998) A genome-wide transcriptional analysis of the mitotic cell cycle. *Mol Cell* 2: 65-73

Collepardo-Guevara R, Schlick T (2014) Chromatin fiber polymorphism triggered by variations of DNA linker lengths. *Proc Natl Acad Sci U S A* 111: 8061-8066

Creekmore BC, Kixmoeller K, Black BE, Lee EB, Chang YW (2024) Ultrastructure of human brain tissue vitrified from autopsy revealed by cryo-ET with cryo-plasma FIB milling. *Nat Commun* 15: 2660

Cuylen S, Blaukopf C, Politi AZ, Muller-Reichert T, Neumann B, Poser I, Ellenberg J, Hyman AA, Gerlich DW (2016) Ki-67 acts as a biological surfactant to disperse mitotic chromosomes. *Nature* 535: 308-312

Danev R, Buijsse B, Khoshouei M, Plitzko JM, Baumeister W (2014) Volta potential phase plate for in-focus phase contrast transmission electron microscopy. *Proc Natl Acad Sci U S A* 111: 15635-15640

Dodonova SO, Zhu F, Dienemann C, Taipale J, Cramer P (2020) Nucleosome-bound SOX2 and SOX11 structures elucidate pioneer factor function. *Nature* 580: 669-672

Dombrowski M, Engeholm M, Dienemann C, Dodonova S, Cramer P (2022) Histone H1
binding to nucleosome arrays depends on linker DNA length and trajectory. *Nat*
Struct Mol Biol 29: 493-501
Dubochet J, Adrian M, Chang JJ, Homo JC, Lepault J, McDowall AW, Schultz P (1988)
Cryo-electron microscopy of vitrified specimens. *Quarterly reviews of biophysics*
21: 129-228
Dubochet J, Noll M (1978) Nucleosome arcs and helices. *Science* 202: 280-286
Dubochet J, Sartori Blanc N (2001) The cell in absence of aggregation artifacts. *Micron*
32: 91-99
Ehara H, Kujirai T, Fujino Y, Shirouzu M, Kurumizaka H, Sekine SI (2019) Structural
insight into nucleosome transcription by RNA polymerase II with elongation
factors. *Science* 363: 744-747
Ehara H, Kujirai T, Shirouzu M, Kurumizaka H, Sekine SI (2022) Structural basis of
nucleosome disassembly and reassembly by RNAPII elongation complex with
FACT. *Science* 377: eabp9466
Ekundayo B, Richmond TJ, Schalch T (2017) Capturing Structural Heterogeneity in
Chromatin Fibers. *J Mol Biol* 429: 3031-3042
Eltsov M, Grewe D, Lemercier N, Frangakis A, Livolant F, Leforestier A (2018)
Nucleosome conformational variability in solution and in interphase nuclei
evidenced by cryo-electron microscopy of vitreous sections. *Nucleic Acids Res*
Eltsov M, Maclellan KM, Maeshima K, Frangakis AS, Dubochet J (2008) Analysis of
cryo-electron microscopy images does not support the existence of 30-nm
chromatin fibers in mitotic chromosomes in situ. *Proc Natl Acad Sci U S A* 105:
19732-19737
Erdmann PS, Hou Z, Klumpe S, Khavnekar S, Beck F, Wilfling F, Plitzko JM,
Baumeister W (2021) In situ cryo-electron tomography reveals gradient
organization of ribosome biogenesis in intact nucleoli. *Nat Commun* 12: 5364
Eustermann S, Schall K, Kostrewa D, Lakomek K, Strauss M, Moldt M, Hopfner KP
(2018) Structural basis for ATP-dependent chromatin remodelling by the INO80
complex. *Nature* 556: 386-390
Fang K, Chen X, Li X, Shen Y, Sun J, Czajkowsky DM, Shao Z (2018) Super-Resolution
Imaging of Individual Human Sub-Chromosomal Regions In Situ Reveals
Nanoscopic Building Blocks of Higher-Order Structure. *ACS Nano*
Farnung L, Ochmann M, Cramer P (2020) Nucleosome-CHD4 chromatin remodeler
structure maps human disease mutations. *Elife* 9
Farnung L, Ochmann M, Engeholm M, Cramer P (2021) Structural basis of nucleosome
transcription mediated by Chd1 and FACT. *Nat Struct Mol Biol* 28: 382-387
Farnung L, Ochmann M, Garg G, Vos SM, Cramer P (2022) Structure of a backtracked
hexasomal intermediate of nucleosome transcription. *Mol Cell* 82: 3126-3134
e3127
Farnung L, Vos SM, Cramer P (2018) Structure of transcribing RNA polymerase II-
nucleosome complex. *Nat Commun* 9: 5432
Farnung L, Vos SM, Wigge C, Cramer P (2017) Nucleosome-Chd1 structure and
implications for chromatin remodelling. *Nature* 550: 539-542

Fatmaoui F, Carrivain P, Grewe D, Jakob B, Victor J-M, Leforestier A, Eltsov M (2022)
Cryo-electron tomography and deep learning denoising reveal native chromatin
landscapes of interphase nuclei. *bioRxiv*
Finch JT, Klug A (1976) Solenoidal model for superstructure in chromatin. *Proc Natl*
Acad Sci U S A 73: 1897-1901
Finogenova K, Bonnet J, Poepsel S, Schafer IB, Finkl K, Schmid K, Litz C, Strauss M,
Benda C, Muller J (2020) Structural basis for PRC2 decoding of active histone
methylation marks H3K36me2/3. *Elife* 9
Flemming W (1880) Beiträge zur Kenntniss der Zelle und Ihrer Lebenserscheinungen.
Archiv für mikroskopische Anatomie 18: 151-259
Fukuda Y, Laugks U, Lucic V, Baumeister W, Danev R (2015) Electron cryotomography
of vitrified cells with a Volta phase plate. *J Struct Biol* 190: 143-154
Gan L, Ladinsky MS, Jensen GJ (2013) Chromatin in a marine picoeukaryote is a
disordered assemblage of nucleosomes. *Chromosoma* 122: 377-386
Garcia-Saez I, Menoni H, Boopathi R, Shukla MS, Soueidan L, Noirclerc-Savoye M, Le
Roy A, Skoufias DA, Bednar J, Hamiche A *et al* (2018) Structure of an H1-Bound
6-Nucleosome Array Reveals an Untwisted Two-Start Chromatin Fiber
Conformation. *Mol Cell* 72: 902-915 e907
Geiss CP, Keramisanou D, Sekulic N, Scheffer MP, Black BE, Frangakis AS (2014)
CENP-A arrays are more condensed than canonical arrays at low ionic strength.
Biophys J 106: 875-882
Gibson BA, Blaukopf C, Lou T, Chen L, Doolittle LK, Finkelstein I, Narlikar GJ, Gerlich
DW, Rosen MK (2023) In diverse conditions, intrinsic chromatin condensates
have liquid-like material properties. *Proc Natl Acad Sci U S A* 120: e2218085120
Gibson BA, Doolittle LK, Schneider MWG, Jensen LE, Gamarra N, Henry L, Gerlich
DW, Redding S, Rosen MK (2019) Organization of Chromatin by Intrinsic and
Regulated Phase Separation. *Cell* 179: 470-484 e421
Grau D, Zhang Y, Lee CH, Valencia-Sanchez M, Zhang J, Wang M, Holder M, Svetlov
V, Tan D, Nudler E *et al* (2021) Structures of monomeric and dimeric
PRC2:EZH1 reveal flexible modules involved in chromatin compaction. *Nat*
Commun 12: 714
Han HM, Huebinger J, Grabenbauer M (2012) Self-pressurized rapid freezing (SPRF)
as a simple fixation method for cryo-electron microscopy of vitreous sections. *J*
Struct Biol 178: 84-87
Han Y, Reyes AA, Malik S, He Y (2020) Cryo-EM structure of SWI/SNF complex bound
to a nucleosome. *Nature* 579: 452-455
Harapin J, Bormel M, Sapra KT, Brunner D, Kaech A, Medalia O (2015) Structural
analysis of multicellular organisms with cryo-electron tomography. *Nature*
methods 12: 634-636
Harlen KM, Churchman LS (2017) The code and beyond: transcription regulation by the
RNA polymerase II carboxy-terminal domain. *Nat Rev Mol Cell Biol* 18: 263-273
Hayashida M, Phengchat R, Malac M, Harada K, Akashi T, Ohmido N, Fukui K (2021)
Higher-Order Structure of Human Chromosomes Observed by Electron
Diffraction and Electron Tomography. *Microsc Microanal* 27: 149-155

Hayles MF, Stokes DJ, Phifer D, Findlay KC (2007) A technique for improved focused ion beam milling of cryo-prepared life science specimens. *J Microsc* 226: 263-269

He S, Wu Z, Tian Y, Yu Z, Yu J, Wang X, Li J, Liu B, Xu Y (2020) Structure of nucleosome-bound human BAF complex. *Science* 367: 875-881

Henneman B, van Emmerik C, van Ingen H, Dame RT (2018) Structure and function of archaeal histones. *PLoS Genet* 14: e1007582

Heumann JM, 2023. PEET. University of Colorado Boulder.

Heumann JM, Hoenger A, Mastronarde DN (2011) Clustering and variance maps for cryo-electron tomography using wedge-masked differences. *J Struct Biol* 175: 288-299

Hewish DR, Burgoyne LA (1973) Chromatin sub-structure. The digestion of chromatin DNA at regularly spaced sites by a nuclear deoxyribonuclease. *Biochem Biophys Res Commun* 52: 504-510

Heymann BJ (2022) Bsoft: Image Processing for Structural Biology. *Bio Protoc* 12: e4393

Hou Z, Nightingale F, Zhu Y, MacGregor-Chatwin C, Zhang P (2023) Structure of native chromatin fibres revealed by Cryo-ET in situ. *Nat Commun* 14: 6324

Huxley HE, Zubay G (1961) Preferential staining of nucleic acid-containing structures for electron microscopy. *J Biophys Biochem Cytol* 11: 273-296

Itoh Y, Woods EJ, Minami K, Maeshima K, Collepardo-Guevara R (2021) Liquid-like chromatin in the cell: What can we learn from imaging and computational modeling? *Current opinion in structural biology* 71: 123-135

Iudin A, Korir PK, Salavert-Torres J, Kleywegt GJ, Patwardhan A (2016) EMPIAR: a public archive for raw electron microscopy image data. *Nature methods* 13: 387-388

Jentink N, Purnell C, Kable B, Swulius MT, Grigoryev SA (2023) Cryoelectron tomography reveals the multiplex anatomy of condensed native chromatin and its unfolding by histone citrullination. *Mol Cell* 83: 3236-3252 e3237

Jentoft IMA, Bauerlein FJB, Welp LM, Cooper BH, Petrovic A, So C, Penir SM, Politi AZ, Horokhovskyi Y, Takala I *et al* (2023) Mammalian oocytes store proteins for the early embryo on cytoplasmic lattices. *Cell* 186: 5308-5327 e5325

Kamasaki T, O'Toole E, Kita S, Osumi M, Usukura J, McIntosh JR, Goshima G (2013) Augmin-dependent microtubule nucleation at microtubule walls in the spindle. *J Cell Biol* 202: 25-33

Kasinath V, Beck C, Sauer P, Poepsel S, Kosmatka J, Faini M, Toso D, Aebersold R, Nogales E (2021) JARID2 and AEBP2 regulate PRC2 in the presence of H2AK119ub1 and other histone modifications. *Science* 371

Kaufmann SH (1989) Induction of endonucleolytic DNA cleavage in human acute myelogenous leukemia cells by etoposide, camptothecin, and other cytotoxic anticancer drugs: a cautionary note. *Cancer Res* 49: 5870-5878

Kaufmann SH, Desnoyers S, Ottaviano Y, Davidson NE, Poirier GG (1993) Specific proteolytic cleavage of poly(ADP-ribose) polymerase: an early marker of chemotherapy-induced apoptosis. *Cancer Res* 53: 3976-3985

Kellenberger E (1987) The Response of Biological Macromolecules and Supramolecular Structures to the Physics of Specimen Cryopreparation. In:

Cryotechniques in Biological Electron Microscopy, Steinbrecht R.A., Zierold K. (eds.) pp. 35-63. Springer Berlin Heidelberg: Berlin, Heidelberg

Khatter H, Myasnikov AG, Natchiar SK, Klaholz BP (2015) Structure of the human 80S ribosome. *Nature* 520: 640-645

Kimanius D, Forsberg BO, Scheres SH, Lindahl E (2016) Accelerated cryo-EM structure determination with parallelisation using GPUs in RELION-2. *Elife* 5

Komarnitsky P, Cho EJ, Buratowski S (2000) Different phosphorylated forms of RNA polymerase II and associated mRNA processing factors during transcription. *Genes Dev* 14: 2452-2460

Kornberg RD (1974) Chromatin structure: a repeating unit of histones and DNA. *Science* 184: 868-871

Kremer JR, Mastronarde DN, McIntosh JR (1996) Computer visualization of three-dimensional image data using IMOD. *J Struct Biol* 116: 71-76

Kujirai T, Ehara H, Fujino Y, Shirouzu M, Sekine SI, Kurumizaka H (2018) Structural basis of the nucleosome transition during RNA polymerase II passage. *Science* 362: 595-598

Ladinsky MS, Pierson JM, McIntosh JR (2006) Vitreous cryo-sectioning of cells facilitated by a micromanipulator. *J Microsc* 224: 129-134

Langmore JP, Paulson JR (1983) Low angle x-ray diffraction studies of chromatin structure in vivo and in isolated nuclei and metaphase chromosomes. *J Cell Biol* 96: 1120-1131

Lau B, Cheng J, Flemming D, La Venuta G, Berninghausen O, Beckmann R, Hurt E (2021) Structure of the Maturing 90S Pre-ribosome in Association with the RNA Exosome. *Mol Cell* 81: 293-303 e294

Lewis TS, Sokolova V, Jung H, Ng H, Tan D (2021) Structural basis of chromatin regulation by histone variant H2A.Z. *Nucleic Acids Res* 49: 11379-11391

Li Y, Eshein A, Virk RKA, Eid A, Wu W, Frederick J, VanDerway D, Gladstein S, Huang K, Shim AR *et al* (2021) Nanoscale chromatin imaging and analysis platform bridges 4D chromatin organization with molecular function. *Sci Adv* 7

Liu X, Li M, Xia X, Li X, Chen Z (2017) Mechanism of chromatin remodelling revealed by the Snf2-nucleosome structure. *Nature* 544: 440-445

Liu Y, Zhang Y, Xue H, Cao M, Bai G, Mu Z, Yao Y, Sun S, Fang D, Huang J (2021) Cryo-EM structure of SETD2/Set2 methyltransferase bound to a nucleosome containing oncohistone mutations. *Cell Discov* 7: 32

Liu Y, Zhou K, Zhang N, Wei H, Tan YZ, Zhang Z, Carragher B, Potter CS, D'Arcy S, Luger K (2020) FACT caught in the act of manipulating the nucleosome. *Nature* 577: 426-431

Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ (1997) Crystal structure of the nucleosome core particle at 2.8 A resolution. *Nature* 389: 251-260

Ma OX, Chong WG, Lee JKE, Cai S, Siebert CA, Howe A, Zhang P, Shi J, Surana U, Gan L (2022) Cryo-ET detects bundled triple helices but not ladders in meiotic budding yeast. *PLoS One* 17: e0266035

Maeshima K, Rogge R, Tamura S, Joti Y, Hikima T, Szerlong H, Krause C, Herman J, Seidel E, DeLuca J *et al* (2016) Nucleosomal arrays self-assemble into supramolecular globular structures lacking 30-nm fibers. *EMBO J* 35: 1115-1132

Mahamid J, Pfeffer S, Schaffer M, Villa E, Danev R, Cuellar LK, Forster F, Hyman AA,
Plitzko JM, Baumeister W (2016) Visualizing the molecular sociology at the HeLa
cell nuclear periphery. *Science* 351: 969-972

Marko M, Hsieh C, Moberlychan W, Mannella CA, Frank J (2006) Focused ion beam
milling of vitreous water: prospects for an alternative to cryo-ultramicrotomy of
frozen-hydrated biological samples. *J Microsc* 222: 42-47

Mashtalir N, Suzuki H, Farrell DP, Sankar A, Luo J, Filipovski M, D'Avino AR, St Pierre
R, Valencia AM, Onikubo T *et al* (2020) A Structural Model of the Endogenous
Human BAF Complex Informs Disease Mechanisms. *Cell* 183: 802-817 e824

Mastronarde DN (1997) Dual-axis tomography: an approach with alignment methods
that preserve resolution. *J Struct Biol* 120: 343-352

Mastronarde DN (2005) Automated electron microscope tomography using robust
prediction of specimen movements. *J Struct Biol* 152: 36-51

Mastronarde DN, Held SR (2017) Automated tilt series alignment and tomographic
reconstruction in IMOD. *J Struct Biol* 197: 102-113

McDowall AW, Smith JM, Dubochet J (1986) Cryo-electron microscopy of vitrified
chromosomes in situ. *EMBO J* 5: 1395-1402

Medeiros JM, Bock D, Weiss GL, Kooger R, Wepf RA, Pilhofer M (2018) Robust
workflow and instrumentation for cryo-focused ion beam milling of samples for
electron cryotomography. *Ultramicroscopy* 190: 1-11

Michael AK, Grand RS, Isbel L, Cavadini S, Kozicka Z, Kempf G, Bunker RD, Schenk
AD, Graff-Meyer A, Pathare GR *et al* (2020) Mechanisms of OCT4-SOX2 motif
readout on nucleosomes. *Science* 368: 1460-1465

Miron E, Oldenkamp R, Brown JM, Pinto DMS, Xu CS, Faria AR, Shaban HA, Rhodes
JDP, Innocent C, de Ornellas S *et al* (2020) Chromatin arranges in chains of
mesoscale domains with nanoscale functional topography independent of
cohesin. *Sci Adv* 6

Nagashima R, Hibino K, Ashwin SS, Babokhov M, Fujishiro S, Imai R, Nozaki T,
Tamura S, Tani T, Kimura H *et al* (2019) Single nucleosome imaging reveals
loose genome chromatin networks via active RNA polymerase II. *J Cell Biol* 218:
1511-1530

Ng CT, Deng L, Chen C, Lim HH, Shi J, Surana U, Gan L (2019) Electron
cryotomography analysis of Dam1C/DASH at the kinetochore-spindle interface in
situ. *J Cell Biol* 218: 455–473

Ng CT, Gan L (2020) Investigating eukaryotic cells with cryo-ET. *Mol Biol Cell* 31: 87-
100

Ng CT, Ladinsky MS, Gan L (2020) Serial Cryomicrotomy of Saccharomyces cerevisiae
for Serial Electron Cryotomography. *Bio Protoc* 10: e3831

Nicastro D, Schwartz C, Pierson J, Gaudette R, Porter ME, McIntosh JR (2006) The
molecular architecture of axonemes revealed by cryoelectron tomography.
Science 313: 944-948

Nishimura M, Takizawa Y, Nozawa K, Kurumizaka H (2022) Structural basis for p53
binding to its nucleosomal target DNA sequence. *PNAS Nexus* 1: pgac177

Nishino Y, Eltsov M, Joti Y, Ito K, Takata H, Takahashi Y, Hihara S, Frangakis AS,
Imamoto N, Ishikawa T *et al* (2012) Human mitotic chromosomes consist

predominantly of irregularly folded nucleosome fibres without a 30-nm chromatin
structure. *EMBO J* 31: 1644-1653
Nodelman IM, Das S, Faustino AM, Fried SD, Bowman GD, Armache JP (2022)
Nucleosome recognition and DNA distortion by the Chd1 remodeler in a
nucleotide-free state. *Nat Struct Mol Biol* 29: 121-129
Nozaki T, Imai R, Tanbo M, Nagashima R, Tamura S, Tani T, Joti Y, Tomita M, Hibino
K, Kanemaki MT *et al* (2017) Dynamic Organization of Chromatin Domains
Revealed by Super-Resolution Live-Cell Imaging. *Mol Cell* 67: 282-293 e287
Ohta S, Bukowski-Wills JC, Sanchez-Pulido L, Alves Fde L, Wood L, Chen ZA, Platani
M, Fischer L, Hudson DF, Ponting CP *et al* (2010) The protein composition of
mitotic chromosomes determined using multiclassifier combinatorial proteomics.
Cell 142: 810-821
Olins AL, Olins DE (1974) Spheroid chromatin units (v bodies). *Science* 183: 330-332
Oliva A, Rosebrock A, Ferrezuelo F, Pyne S, Chen H, Skiena S, Futcher B,
Leatherwood J (2005) The cell cycle-regulated genes of Schizosaccharomyces
pombe. *PLoS Biol* 3: e225
Otterstrom J, Castells-Garcia A, Vicario C, Gomez-Garcia PA, Cosma MP, Lakadamyali
M (2019) Super-resolution microscopy reveals how histone tail acetylation affects
DNA compaction within nucleosomes in vivo. *Nucleic Acids Res* 47: 8470-8484
Ou HD, Phan S, Deerinck TJ, Thor A, Ellisman MH, O'Shea CC (2017) ChromEMT:
Visualizing 3D chromatin structure and compaction in interphase and mitotic
cells. *Science* 357
Pacheco-Fiallos B, Vorlander MK, Riabov-Bassat D, Fin L, O'Reilly FJ, Ayala FI,
Schellhaas U, Rappsilber J, Plaschka C (2023) mRNA recognition and packaging
by the human transcription-export complex. *Nature* 616: 828-835
Palozola KC, Donahue G, Liu H, Grant GR, Becker JS, Cote A, Yu H, Raj A, Zaret KS
(2017) Mitotic transcription and waves of gene reactivation during mitotic exit.
Science 358: 119-122
Patel AB, Moore CM, Greber BJ, Luo J, Zukin SA, Ranish J, Nogales E (2019)
Architecture of the chromatin remodeler RSC and insights into its nucleosome
engagement. *Elife* 8
Peng X, Karuturi RK, Miller LD, Lin K, Jia Y, Kondu P, Wang L, Wong LS, Liu ET,
Balasubramanian MK *et al* (2005) Identification of cell cycle-regulated genes in
fission yeast. *Mol Biol Cell* 16: 1026-1042
Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE
(2004) UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* 25: 1605-1612
Poepsel S, Kasinath V, Nogales E (2018) Cryo-EM structures of PRC2 simultaneously
engaged with two functionally distinct nucleosomes. *Nat Struct Mol Biol* 25: 154-
162
Prescott DM, Bender MA (1962) Synthesis of RNA and protein during mitosis in
mammalian tissue culture cells. *Exp Cell Res* 26: 260-268
Remnant L, Kochanova NY, Reid C, Cisneros-Soberanis F, Earnshaw WC (2021) The
intrinsically disorderly story of Ki-67. *Open Biol* 11: 210120

Ricci MA, Manzo C, Garcia-Parajo MF, Lakadamyali M, Cosma MP (2015) Chromatin
fibers are formed by heterogeneous groups of nucleosomes in vivo. *Cell* 160:
1145-1158
Rigort A, Bauerlein FJ, Leis A, Gruska M, Hoffmann C, Laugks T, Bohm U, Eibauer M,
Gnaegi H, Baumeister W *et al* (2010) Micromachining tools and correlative
approaches for cellular cryo-electron tomography. *J Struct Biol* 172: 169-179
Ruan W, Venkatachalam G, Sobota RM, Chen L, Wang LC, Jacobson A, Paramasivam
K, Surana U (2019) Resistance to anti-microtubule drug-induced cell death is
determined by regulation of BimEL expression. *Oncogene* 38: 4352-4365
Rustici G, Mata J, Kivinen K, Lio P, Penkett CJ, Burns G, Hayles J, Brazma A, Nurse P,
Bahler J (2004) Periodic gene expression program of the fission yeast cell cycle.
Nat Genet 36: 809-817
Sanulli S, Trnka MJ, Dharmarajan V, Tibble RW, Pascal BD, Burlingame AL, Griffin PR,
Gross JD, Narlikar GJ (2019) HP1 reshapes nucleosome core to promote phase
separation of heterochromatin. *Nature* 575: 390-394
Schaffer M, Engel BD, Laugks T, Mahamid J, Plitzko JM, Baumeister W (2015) Cryo-
focused Ion Beam Sample Preparation for Imaging Vitreous Cells by Cryo-
electron Tomography. *Bio Protoc* 5
Schalch T, Duda S, Sargent DF, Richmond TJ (2005) X-ray structure of a
tetranucleosome and its implications for the chromatin fibre. *Nature* 436: 138-141
Scheffer MP, Eltsov M, Bednar J, Frangakis AS (2012) Nucleosomes stacked with
aligned dyad axes are found in native compact chromatin in vitro. *J Struct Biol*
178: 207-214
Scheffer MP, Eltsov M, Frangakis AS (2011) Evidence for short-range helical order in
the 30-nm chromatin fibers of erythrocyte nuclei. *Proc Natl Acad Sci U S A* 108:
16992-16997
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for
biological-image analysis. *Nature methods* 9: 676-682
Scott SJ, Suvarna KS, D'Avino PP (2020) Synchronization of human retinal pigment
epithelial-1 cells in mitosis. *J Cell Sci* 133
Sedat J, McDonald A, Cang H, Lucas J, Arigovindan M, Kam Z, Murre C, Elbaum M
(2022a) A proposed unified interphase nucleus chromosome structure:
Preliminary preponderance of evidence. *Proc Natl Acad Sci U S A* 119:
e2119101119
Sedat J, McDonald A, Kasler H, Verdin E, Cang H, Arigovindan M, Murre C, Elbaum M
(2022b) A proposed unified mitotic chromosome architecture. *Proc Natl Acad Sci*
U S A 119: e2119107119
Simpson RT (1978) Structure of the chromatosome, a chromatin particle containing 160
base pairs of DNA and all the histones. *Biochemistry* 17: 5524-5531
Soman A, Wong SY, Korolev N, Surya W, Lattmann S, Vogirala VK, Chen Q,
Berezhnoy NV, van Noort J, Rhodes D *et al* (2022) Columnar structure of human
telomeric chromatin. *Nature* 609: 1048-1055
Song F, Chen P, Sun D, Wang M, Dong L, Liang D, Xu RM, Zhu P, Li G (2014) Cryo-
EM study of the chromatin fiber reveals a double helix twisted by
tetranucleosomal units. *Science* 344: 376-380

Spellman PT, Sherlock G, Zhang MQ, Iyer VR, Anders K, Eisen MB, Brown PO, Botstein D, Futcher B (1998) Comprehensive identification of cell cycle-regulated genes of the yeast Saccharomyces cerevisiae by microarray hybridization. *Mol Biol Cell* 9: 3273-3297

Stanek D, Fox AH (2017) Nuclear bodies: news insights into structure and function. *Current opinion in cell biology* 46: 94-101

Stenstrom L, Mahdessian D, Gnann C, Cesnik AJ, Ouyang W, Leonetti MD, Uhlen M, Cuylen-Haering S, Thul PJ, Lundberg E (2020) Mapping the nucleolar proteome reveals a spatiotemporal organization related to intrinsic protein disorder. *Mol Syst Biol* 16: e9469

Strickfaden H, Tolsma TO, Sharma A, Underhill DA, Hansen JC, Hendzel MJ (2020) Condensed Chromatin Behaves like a Solid on the Mesoscale In Vitro and in Living Cells. *Cell* 183: 1772-1784 e1713

Studer D, Klein A, Iacovache I, Gnaegi H, Zuber B (2014) A new tool based on two micromanipulators facilitates the handling of ultrathin cryosection ribbons. *J Struct Biol* 185: 125-128

Takizawa Y, Ho CH, Tachiwana H, Matsunami H, Kobayashi W, Suzuki M, Arimura Y, Hori T, Fukagawa T, Ohi MD *et al* (2020) Cryo-EM Structures of Centromeric Tri-nucleosomes Containing a Central CENP-A Nucleosome. *Structure* 28: 44-53 e44

Tan ZY, Cai S, Noble AJ, Chen JK, Shi J, Gan L (2023) Heterogeneous non-canonical nucleosomes predominate in yeast cells in situ. *Elife* 12

Tan ZY, Cai S, Paithankar SA, Nie X, Shi J, Gan L (2024) Macromolecular and cytological changes in fission yeast G0 nuclei. *bioRxiv*

Tanaka H, Takizawa Y, Takaku M, Kato D, Kumagawa Y, Grimm SA, Wade PA, Kurumizaka H (2020) Interaction of the pioneer transcription factor GATA3 with nucleosomes. *Nat Commun* 11: 4136

Tang G, Peng L, Baldwin PR, Mann DS, Jiang W, Rees I, Ludtke SJ (2007) EMAN2: an extensible image processing suite for electron microscopy. *J Struct Biol* 157: 38-46

Taylor JH (1960) Nucleic acid synthesis in relation to the cell division cycle. *Ann N Y Acad Sci* 90: 409-421

Teves SS, An L, Hansen AS, Xie L, Darzacq X, Tjian R (2016) A dynamic mode of mitotic bookmarking by transcription factors. *Elife* 5

Tivol WF, Briegel A, Jensen GJ (2008) An improved cryogen for plunge freezing. *Microsc Microanal* 14: 375-379

Villa E, Schaffer M, Plitzko JM, Baumeister W (2013) Opening windows into the cell: focused-ion-beam milling for cryo-electron tomography. *Current opinion in structural biology* 23: 771-777

Wang H, Schilbach S, Ninov M, Urlaub H, Cramer P (2023) Structures of transcription preinitiation complex engaged with the +1 nucleosome. *Nat Struct Mol Biol* 30: 226-232

Wang H, Xiong L, Cramer P (2021a) Structures and implications of TBP-nucleosome complexes. *Proc Natl Acad Sci U S A* 118

Wang J, Hu C, Chen X, Li Y, Sun J, Czajkowsky DM, Shao Z (2022a) Single-Molecule Micromanipulation and Super-Resolution Imaging Resolve Nanodomains

Underlying Chromatin Folding in Mitotic Chromosomes. *ACS Nano* 16: 8030-
8039

Wang L, Yu J, Yu Z, Wang Q, Li W, Ren Y, Chen Z, He S, Xu Y (2022b) Structure of
nucleosome-bound human PBAF complex. *Nat Commun* 13: 7644

Wang S, Vogirala VK, Soman A, Berezhnoy NV, Liu ZB, Wong ASW, Korolev N, Su CJ,
Sandin S, Nordenskiold L (2021b) Linker histone defines structure and self-
association behaviour of the 177 bp human chromatosome. *Sci Rep* 11: 380

Weis F, Hagen WJH, Schorb M, Mattei S (2021) Strategies for Optimization of
Cryogenic Electron Tomography Data Acquisition. *J Vis Exp*

Willhoft O, Ghoneim M, Lin CL, Chua EYD, Wilkinson M, Chaban Y, Ayala R,
McCormack EA, Ocloo L, Rueda DS *et al* (2018) Structure and dynamics of the
yeast SWR1-nucleosome complex. *Science* 362

Xiang W, Roberti MJ, Heriche JK, Huet S, Alexander S, Ellenberg J (2018) Correlative
live and super-resolution imaging reveals the dynamic structure of replication
domains. *J Cell Biol* 217: 1973-1984

Xu J, Ma H, Jin J, Uttam S, Fu R, Huang Y, Liu Y (2018) Super-Resolution Imaging of
Higher-Order Chromatin Structures at Different Epigenomic States in Single
Mammalian Cells. *Cell Rep* 24: 873-882

Yakovlev S, Downing KH (2011) Freezing in sealed capillaries for preparation of frozen
hydratedsections. *J Microsc* 244: 235-247

Yan L, Wu H, Li X, Gao N, Chen Z (2019) Structures of the ISWI-nucleosome complex
reveal a conserved mechanism of chromatin remodeling. *Nat Struct Mol Biol* 26:
258-266

Zhang H, Li Y, Liu Y, Li D, Wang L, Song K, Bao K, Zhu P (2023) A method for restoring
signals and revealing individual macromolecule states in cryo-ET, REST. *Nat*
Commun 14: 2937

Zhang M, Diaz-Celis C, Onoa B, Canari-Chumpitaz C, Requejo KI, Liu J, Vien M,
Nogales E, Ren G, Bustamante C (2022) Molecular organization of the early
stages of nucleosome phase separation visualized by cryo-electron tomography.
Mol Cell

Zhao W, Jensen GJ (2022) In situ architecture of human kinetochore-microtubule
interface visualized by cryo-electron tomography. *bioRxiv*

Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA (2017)
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. *Nature methods* 14: 331-332

Zhou BR, Feng H, Kale S, Fox T, Khant H, de Val N, Ghirlando R, Panchenko AR, Bai
Y (2021) Distinct Structures and Dynamics of Chromatosomes with Different
Human Linker Histone Isoforms. *Mol Cell* 81: 166-182 e166

Zhou BR, Jiang J, Feng H, Ghirlando R, Xiao TS, Bai Y (2015) Structural Mechanisms
of Nucleosome Recognition by Linker Histones. *Mol Cell* 59: 628-638

Zhou BR, Jiang J, Ghirlando R, Norouzi D, Sathish Yadav KN, Feng H, Wang R, Zhang
P, Zhurkin V, Bai Y (2018) Revisit of Reconstituted 30-nm Nucleosome Arrays
Reveals an Ensemble of Dynamic Structures. *J Mol Biol* 430: 3093-3110

Zhou K, Gaullier G, Luger K (2019) Nucleosome structure and dynamics are coming of
age. *Nat Struct Mol Biol* 26: 3-13

Zhou K, Gebala M, Woods D, Sundararajan K, Edwards G, Krzizike D, Wereszczynski
J, Straight AF, Luger K (2022) CENP-N promotes the compaction of centromeric
chromatin. *Nat Struct Mol Biol* 29: 403-413
Zhu L, Brangwynne CP (2015) Nuclear bodies: the emerging biophysics of
nucleoplasmic phases. *Current opinion in cell biology* 34: 23-30
Zivanov J, Nakane T, Forsberg BO, Kimanius D, Hagen WJ, Lindahl E, Scheres SH
(2018) New tools for automated high-resolution cryo-EM structure determination
in RELION-3. *Elife* 7
1925
1926

1927

1928 **Figure 1. Metaphase chromosomes are depleted of RNA polymerase II.**

1929 Differential interference contrast (DIC) images of representative G1 and metaphase

1930 cells that were stained to detect DNA (stained with DAPI) and immunofluorescent

1931 detection of initiating RNA polymerase II (RNAPII) phosphorylated at serine 5 of the

1932 RPB1 subunit's C-terminal tail heptad repeats (RPB1-S5P). (A) G1 and metaphase cells

1933 were incubated in DMEM/F12 medium for 1 minute before fixation. (B) G1 and

1934 metaphase cells were incubated in DMEM/F12 medium containing 9% DMSO for 1

1935 minute before fixation. (C) G1 and metaphase cells were incubated in DMEM/F12

1936 medium containing 9% glycerol for 1 minute before fixation. In the two middle columns,

1937 the DAPI and immunofluorescence signals are shown in inverted contrast for clarity.

1938

Figure 2. Cryo-ET of G1 chromatin *in situ*.
(A) Cryotomographic slice (10 nm) of the perinuclear region. The nuclear envelope is oblique to the milling direction and is therefore not visible. Its approximate center is indicated by the dashed line. **(B)** Cryotomographic slice (10 nm) of a region deeper inside the nucleus. **(C)** Enlargement (4-fold) of one region boxed in panel A, showing a perinuclear chromatin domain. **(D)** Enlargement of a megacomplex-rich nucleoplasm region boxed in panel A. (**E**) Enlargement of the chromatin domain boxed in panel B. **(F)** Enlargement (4-fold) of a "dense irregular body" in panel B. In all insets, nucleosomes are indicated by black arrowheads and megacomplexes are indicated by white arrowheads. These images are from cryotomograms denoised by CryoCARE (Buchholz *et al.*, 2019). Non-denoised versions are shown in Figure S37.

1950
1951

Figure 3. Subtomogram analysis of G1 nucleosomes.
(**A**) Subtomogram averages of mononucleosomes in G1 cells, shown in the (top to bottom) disc, disc, gyre, and side views, respectively. They are ordered (left to right) from those having the shortest to the longest ordered linker DNA. Classes marked with a magenta asterisk have noticeably asymmetric linker DNA length. (**B**) Subtomogram average of ordered stacked dinucleosomes. (**C**) Subtomogram averages of nucleosomes with disordered gyre-proximal densities; to show this extra density, these averages are rendered at a lower contour level than those of panels A and B. Some of the images in panels A – C are reproduced from Figure S15; the structures are also shown in Movie S1. At the bottom of panels A – C, the numbers of subtomograms that belong to each class is shown. (**D**) Remapped model of nucleosomes (blue) in the chromatin domains, preribosomes (light blue), and cytoplasmic ribosomes (yellow). The remapped model includes nucleosome classes shown in panels A – C. To facilitate the visualization of the nucleosomes, the proximal densities are hidden in the remapped model by a combination of higher contour level and the UCSF Chimera "hide dust" tool. The remapped model spans the entire thickness of the lamella (~90 nm). The left insets show 3-fold enlargements of the regions boxed on the left and right inset shows a cryotomographic slice (10 nm) of the same remapped region. The circular gap in the lower-right group of nucleosomes is volume occupied by a nuclear pore complex (approximate position indicated by a 100-nm dotted circle). See also Movie S2. The cryotomographic slices in panel D are from cryotomograms denoised by Cryo-CARE (Buchholz *et al.*, 2019). This figure is reproduced with non-denoised cryotomographic slices in Figure S38.

1975

Figure 4. Cryo-ET of metaphase chromatin *in situ*.

(A and B) Cryotomographic slice (10 nm) overview of portions of two metaphase chromosomes. (**C and D**) Enlargements (4-fold) of two regions boxed in panel A. (**E and F**) Enlargements (4-fold) of two regions boxed in panel B. **Panel E** shows an enlargement of a region of the chromosome boundary. **Panel F** shows an enlargement of microtubules embedded in the chromosome. In all the insets, the nucleosome-like densities (double-gyre motifs and top-views) are indicated by arrowheads. Note that there are stacked nucleosomes in panels C and E. Microtubules (MT) and ribosomes (R) are also indicated in the insets. These images are from cryotomograms denoised by Cryo-CARE (Buchholz *et al.*, 2019). Non-denoised versions are shown in Figure S39.

1986
Figure 5. Subtomogram analysis of metaphase nucleosomes.
(**A**) Subtomogram averages of mononucleosomes in metaphase cells, shown in the (top
to bottom) disc, disc, gyre, and side views, respectively. They are ordered (left to right)
from those having the shortest to the longest ordered linker DNA. The class marked

with a magenta asterisk has noticeably asymmetric linker DNA length. (**B**)
Subtomogram average of ordered stacked dinucleosomes. (**C**) Subtomogram averages
of nucleosomes with disordered gyre-proximal densities; to show this extra density,
these averages are rendered at a lower contour level than those of panels A and B.
Some of the images in panels A – C are reproduced from Figure S31; the structures are
also shown in Movie S3. At the bottom of panels A – C, the numbers of subtomograms
that belong to each class is shown. (**D**) Remapped model of nucleosomes (blue) and
ribosomes (yellow). The remapped model includes nucleosome classes shown in
panels A – C. Nucleosomes with gyre-proximal densities are not rendered in the
remapped model. The remapped volume contains all Z slices of the reconstructed
lamella (~140 nm thick). The left insets show 3-fold enlargements of the regions boxed
on the left and the right insets show cryotomographic slices (10 nm) of the same
remapped region. See also Movie S4. The cryotomographic slices in panel D are from
cryotomograms denoised by Cryo-CARE (Buchholz *et al.*, 2019). This figure is
reproduced with non-denoised cryotomographic slices in Figure S40.

2006

Figure 6. Example of partial overlaps between subtomogram class averages.
The panels here show remapped nucleosomes and a 10-nm cryotomographic slice from a region of a (A) G1 and (B) metaphase cell cryotomogram. In the upper panels, the remapped mononucleosomes (Group 1, blue), ordered stacked dinucleosomes (Group 2, green) and nucleosomes with gyre-proximal densities (Group 3, magenta) are rendered separately; and combined in the lower left panel. The remapped models shown here were generated using the nucleosome classes shown in Figures 3 and 5, panels A – C. To facilitate the visualization of the remapped densities, the UCSF Chimera "hide dust" tool was used to hide the small noisy densities. Note that all the nucleosomes in the remapped models here are rendered at a lower contour level to display the extra densities. Comparison of the remapped model with the cryotomographic slice from the same region show that some of the extra densities are indeed located over nucleosomes with side views. Black arrowheads indicate extra densities that partially overlap with a neighboring remapped nucleosome. White arrowheads indicate extra densities that do not appear to overlap any remapped nucleosome. Gray arrowheads indicate an overlap between an extra density with a tomographic slice feature that resembles a nucleosome side view. Possible explanations are that these densities are not nucleosomes or that they are nucleosomes but were misclassified. The cryotomographic slices shown here are from cryotomograms that were denoised using cryo-CARE. This figure is reproduced with non-denoised cryotomographic slices in Figure S41.

2028

Figure 7. Comparison of G1 and metaphase nucleosomes.
Crystal structures of the nucleosome core particle (PDB 1AOI) and the chromatosome
(PDB 5NL0) docked into the *in situ* nucleosome subtomogram averages (transparent
blue) with shorter and longer ordered linker DNA, respectively, for (**A**) G1 and (**B**)
metaphase nucleosomes. For the mononucleosome and chromatosomes averages, the
subtomograms from group 1 were pooled as either the shorter or longer linker
nucleosomes (separately for G1 and metaphase), then jointly refined. The DNA is
rendered dark blue and gold while the histones are rendered gold. The histone N-
terminal tails have been omitted for clarity. For the model containing the chromatosome,
the linker histone H1 was included, though the resolution of the density map is
insufficient to resolve the linker histone. The nucleosome density envelopes in the right
column appear larger than in the other columns because a lower contour level is used
to visualize the gyre-proximal densities. Dashed black lines and numbers indicate the

2042   super-helical locations. They also have poorer definition than in earlier figures because

2043   multiple classes are superposed here. The red dots are the approximate centers of

2044   masses of gyre-proximal densities.