Epi-microRNA mediated metabolic reprogramming ensures affinity maturation

Rinako Nakagawa[1*], Miriam Llorian[2], Sunita Varsani-Brown[3], Probir Chakravarty[2], Jeannie M.

Camarillo[4], David Barry[5], Roger George[6], Neil P. Blackledge[7], Graham Duddy[3], Neil L.

Kelleher[4], Robert J. Klose[7], Martin Turner[8] and Dinis P. Calado[1*]

1.   Immunity and Cancer Laboratory, Francis Crick Institute, London, NW1 1AT, UK

2.   Bioinformatics and Biostatistics Laboratory, Francis Crick Institute, London, NW1 1AT,

UK

3.   Genetic Modification Service Laboratory, Francis Crick Institute, London, NW1 1AT,

UK

4.   Department of Chemistry, Molecular Biosciences and the National Resource for

Translational and Developmental Proteomics, Northwestern University, Evanston, IL,

60208, USA

5.   Advanced Light Microscopy Laboratory, Francis Crick Institute, London, NW1 1AT,

UK

6.   Structural Biology Laboratory, Francis Crick Institute, London, NW1 1AT, UK

7.   Department of Biochemistry, University of Oxford, Oxford, OX1 3QU, UK

8.   Immunology Programme, The Babraham Institute, Cambridge, CB22 3AT, UK

* Correspondence; these authors contributed equally to this work

Abstract

To increase antibody affinity against pathogens, positively selected GC-B cells initiate cell

division in the light zone (LZ) of germinal centres (GCs). Among those, higher-affinity clones

migrate to the dark zone (DZ) and vigorously proliferate by relying on oxidative

phosphorylation (OXPHOS). However, it remains unknown how positively selected GC-B cells

adapt their metabolism for cell division in the glycolysis-dominant, cell cycle arrest-inducing,

hypoxic LZ microenvironment. Here, we show that microRNA (miR)-155 mediates metabolic

reprogramming during positive selection to protect high-affinity clones. Transcriptome

examination and mass spectrometry analysis revealed that miR-155 regulates H3K36me2 levels

by directly repressing hypoxia-induced histone lysine demethylase, *Kdm2a*. This is

indispensable for enhancing OXPHOS through optimizing the expression of vital nuclear

mitochondrial genes under hypoxia. The miR-155-*Kdm2a* interaction is crucial to prevent

excessive production of reactive oxygen species and apoptosis. Thus, miR-155-mediated

epigenetic regulation promotes mitochondrial fitness in high-affinity clones, ensuring their

expansion and consequently affinity maturation.

Main

Germinal centres (GCs) transiently form in B cell follicles of secondary lymphoid organs upon

infection or vaccination. GCs are indispensable sites for increasing the affinity of antibodies, i.e.,

for the production of high-affinity antibody-secreting plasma cells that are critical for protecting

hosts from pathogens. The process by which antibodies increase affinity to the antigen is called

affinity maturation [1,2] and occurs in GCs through a bidirectional migration cycle between two

regions, the light zone (LZ) and the dark zone (DZ) [3]. In the LZ, GC-B cells undergo antigen-

driven selection in response to signals from B cell receptor (BCR) engagement and T follicular

helper cells (TFHs) [4]. Positively selected LZ GC-B cells induce *Myc*, a critical cell cycle

regulator required for GC maintenance, and the transient c-MYC expression marks "licensed"

GC-B cells [5,6]. Upon receiving positive selection signals, GC-B cells initiate cell division in the

LZ. Among these cells, clones with higher-affinity further vigorously proliferate after their

transition to the DZ for the clonal expansion [7].

Mitochondrial oxidative phosphorylation (OXPHOS) is essential for GC-B cells to proliferate in

the DZ [8,9]. In contrast, quiescent GC-B cells are reliant on glycolysis [10]. This is concordant with

recent reports showing that the LZ, where predominantly quiescent cells reside, represents a

hypoxic microenvironment [10-12]. Hypoxia promotes glycolysis [13] and suppresses energy

consuming processes, such as cell cycle progression and translation [14-16]. Therefore, cells under

hypoxic conditions generally arrest the cell cycle. During positive selection, GC-B cells

substantially change gene expression profiles to ensure their proliferation in the LZ before

migrating to the DZ [4,7,17-19]. To initiate cell division, positively selected GC-B cells need to

undergo a metabolic reprogramming from glycolysis-dominant metabolism to OXPHOS

without overburdening mitochondria with oxidative stress in the hypoxic LZ microenvironment.

However, such mechanisms remain unknown.

Here, we show that microRNA (miR)-155 functions as an epigenetic regulator that is essential

for metabolic reprogramming in LZ GC-B cells during positive selection. Through miR-155-

mediated counteraction of hypoxia, mitochondrial remodelling within cMyc[+] GC-B cells was

achieved exclusively in the LZ. In high-affinity clones among these positively selected GC-B

cells, miR-155 directly supressed the expression of hypoxia-induced lysine demethylase 2a

(*Kdm2a*), a histone modifier involved in transcriptional repression through its preferential

demethylation of di-methylated lysine 36 of histone H3 (H3K36me2) [20,21]. MiR-155-mediated

repression of *Kdm2a* regulated H3K36me2 levels at promoters and transcription start sites

(TSSs), which subsequently facilitated optimal expression of key nuclear mitochondrial genes in

cMyc[+] LZ GC-B cells. OXPHOS was enhanced under hypoxic conditions through this

epigenetic regulation. The interaction between miR-155 and *Kdm2a* was required to prevent

GC-B cells from excessive production of reactive oxygen species (ROS) and subsequent

apoptosis. Hence, miR-155 promotes mitochondrial fitness in high-affinity LZ GC-B cells

during positive selection, which allows these cells to endure rapid proliferation upon transition

to the DZ. In this way, high-affinity B cell clones can undergo clonal expansion in the DZ, thus

ensuring affinity maturation.

Mitochondrial ROS production is exacerbated by *Mir155*-deficiency in GC-B cells

OXPHOS is required to support the robust proliferation of DZ GC-B cells particularly with

high-affinity BCRs [9]. Mitochondrial ROS are produced during normal OXPHOS activity at the

electron transport chain on the inner mitochondrial membrane [22]. However, defective

mitochondria drive overproduction of ROS, which can activate apoptotic pathways [23]. MiR-155

is predominantly expressed in cMyc$^+$ GC-B cells and is known to protect cMyc$^+$ GC-B cells

from apoptosis, particularly those with high-affinity BCRs [24]. Therefore, we hypothesized that

miR-155 prevents cell death, in part, by limiting excessive production of mitochondrial ROS.

To test the hypothesis, we utilized the SW$_{HEL}$ adoptive transfer system which allowed us to

investigate miR-155 functions in GC-B cells in a B cell-intrinsic manner. In this system, B cells

from *Mir155*-sufficient or *Mir155*-deficient SW$_{HEL}$ *Myc$^{gfp/gfp}$ Aicda-cre*-h*CD2* mice were

transferred into CD45.1$^+$ congenic recipient mice. SW$_{HEL}$ mice express the hen egg lysozyme

(HEL)-specific immunoglobulin heavy chain knock-in and light chain transgene (HyHEL10

BCR) [25]. *Myc$^{gfp/gfp}$* reporter mice express a GFP-cMYC fusion protein from the endogenous *Myc*

locus [26]. *Aicda-cre*-h*CD2* transgenic mice carry a *Cre* gene linked with a truncated human *CD2*

cDNA via an internal ribosome entry site (IRES) under the control of *Aicda* promoter [27]. In

these mice, positively selected cMyc$^+$ GC-B cells were identified by GFP fluorescence and

hCD2 surface expression. Recipient mice that received transferred B cells were challenged with

HEL3$^\times$ (**Fig. 1a**). Donor GC-B cells (CD45.1$^-$ CD45.2$^+$ B220$^+$ hCD2$^+$ CD38^{lo} CD95$^+$) were

gated as described in **Extended Data Fig. 1a**, and by coupling DZ/LZ distinguishing markers

(CXCR4/CD86) to cMyc-GFP expression. Then, GC-B cells were subdivided into four

consecutive subpopulations: cMyc$^-$ LZ, cMyc$^+$ LZ, cMyc$^-$ DZ and cMyc$^+$ DZ [7]. MiR-155 is

mostly co-expressed with cMyc in GC-B cells, and the expression of both cMyc and miR-155

changes as follows: highest levels in cMyc$^+$ LZ GC-B cells, intermediate levels in cMyc$^+$ DZ

GC-B cells and followed by negligible levels in cMyc⁻ DZ and cMyc⁻ LZ GC-B cells [5,6,24]

(**Extended Data Fig. 1b**). Lack of miR-155 did not impact the expression of cMyc in GC-B

cells, hence the mean fluorescent intensity (MFI) of cMyc-GFP was comparable between

wildtype (hereafter called as "WT") and *Mir155*-knock out (hereafter called as "KO") cells

(**Extended Data Fig. 1c**). However, the number of cMyc⁺ LZ and cMyc⁺ DZ GC-B cells were

significantly reduced in the absence of miR-155 (**Extended Data Fig. 1d**). The MitoSOX-based

flow cytometric assay for the detection of mitochondrial ROS revealed that in the absence of

miR-155, GC-B cell subpopulations, including cMyc⁺ LZ, cMyc⁺ DZ and cMyc⁻ DZ, exhibited

a significantly higher percentage of MitoSOX⁺ cells compared to their respective WT cells (**Fig.**

1b). Additionally, we observed a concurrent increase in the proportion of active-caspase 3⁺ cells

within the cMyc⁺ LZ, cMyc⁺ DZ and cMyc⁻ DZ subpopulations of the KO mice compared to the

WT mice (**Fig. 1c**). These findings indicate that miR-155 plays a role in mitigating

mitochondrial ROS levels in GC-B cells.

MiR-155 is required for promoting mitochondrial fitness in GC-B cells

To gain a comprehensive understanding of miR-155-mediated regulation of mitochondrial

functions and dynamics of OXPHOS activity during the migration cycle of GC-B cells, we

conducted single cell (sc) RNA-seq on both WT and KO GC-B cells and dissected GC-B cells

into distinct subpopulations. Unsupervised clustering identified nine distinct clusters: LZ

signature genes [9] were expressed in clusters 0, 1, 2, 5 and 8, and clusters 0 and 1 predominantly

represented cMyc⁻ LZ GC-B cells due to their enrichment in the G1 phase of the cell cycle

(**Extended Data Fig. 1e-g**). Cluster 8 showed enrichment in pre-memory B cell (pre-MBC)-

associated genes, such as *Ccr6*, *Cd3*8 and *Hhex* [28,29]), indicating its representation of pre-MBC

cells (**Supplementary Table 1**). DZ signature genes [9] were expressed in clusters 3, 4, 5, 6 and 7

(**Extended Data Fig. 1g**), which correlated strongly with cell cycle progression, i.e., cluster 6; S

phase, cluster 4; S→G2/M phase, cluster 7; G2/M phase and cluster 3; G2/M→G1 phase

(**Extended Data Fig. 1e**). The order of the clusters likely reflects a temporal sequence of DZ

GC-B cells as clusters 6 → 4 → 7 → 3.

To establish more specific relationships between the clusters we employed partition-based graph

abstraction (PAGA) [30]. The PAGA graph revealed that LZ clusters 0, 1, 2 and 8 formed a

closely interconnected group, while DZ clusters 3, 4, 6 and 7, characterized by high expression

of *Cxcr4*, formed a distinct group (**Fig. 1d** and **Extended Data Fig. 1h**). Cluster 5, which

exhibited higher mitochondrial content but lower gene counts than the other clusters, did not

belong to either LZ or DZ groups (**Fig. 1d**, **Extended Data Fig. 1h,i**). Notably, this cluster

displayed a significant decrease in rRNA levels (**Extended Data Fig. 1j**), which is indicative of

early apoptosis [31]. Therefore, we excluded cluster 5 from further analysis. The high expression

of *Myc* in clusters 2 (LZ) and 6 (DZ) in the PAGA graph corresponded to the enrichment of *Myc*

signature genes [5,6] observed in these clusters, as shown in the UMAP (**Fig. 1d** and **Extended**

Data Fig. 1g). The increased *Myc* expression in cluster 8 (pre-MBCs) in **Fig. 1d** is consistent

with our previous work suggesting that cMyc-induced positively selected GC-B cells can give

rise to MBCs [7,32]. In the PAGA graph, clusters 0 (LZ) and 3 (DZ) predominantly connect the LZ

and DZ groups (**Fig. 1d** and **Extended Data Fig. 1h**), indicating that cluster 0 represents LZ B

cells soon after migration back from the DZ. The temporal and spatial dynamics of the clusters

are summarized in **Fig. 1e**.

Using these temporally linked GC-B cell clusters, we investigated the changes in mitochondrial

functions in GC-B cells in the absence of miR-155 by calculating the gene signature scores for

"mitochondrion" and "OXPHOS". We found these to be significantly lower in cluster 2 (cluster

enriched with cMyc$^+$ LZ GC-B cells) and all the DZ clusters (clusters 6, 4, 7 and 3) of KO cells

than those of WT cells (**Fig. 1f**). These results suggest that the absence of miR-155 led to

impaired mitochondrial functions in cMyc$^+$ LZ GC-B cells (cluster 2) and cMyc$^+$ DZ GC-B

cells (cluster 6), and that such impairment was still observed in cMyc$^-$ DZ GC-B cells (clusters 4,

7 and 3) that no longer express miR-155 in WT mice. Thus, miR-155 plays a critical role in

maintaining mitochondrial integrity in cMyc$^+$ GC-B cells and likely impacts the bioenergetic

functions in GC-B cells.

To further clarify the regulatory role of miR-155 in LZ GC-B cells, we carried out bulk RNA-

seq analysis on WT and KO cMyc$^+$ LZ GC-B cells. Gene set enrichment analysis (GSEA) on

RNA-seq data revealed the top eight enriched biological pathways in WT cMyc$^+$ LZ GC-B cells

compared to KO cMyc$^+$ LZ GC-B cells were related to "ribosome", "translation" and

"mitochondria", either alone or in combination (**Extended Data Fig. 1k**). Nearly half of the

significantly increased genes in the top three pathways ("ribosomal subunit", "Structural

constituent of ribosome" and "ribosome") encoded mitochondrial ribosomal proteins

(**Supplementary Table 2**). Mitochondrial ribosomal proteins are encoded by nuclear genes and

are subsequently transported to the mitochondria following their synthesis in the cytoplasm [33].

The Cytoscape network analysis of GSEA results revealed an enrichment of pathways involved

in mitochondrial protein transport and mitochondrial functions, particularly electron transport, in

WT cMyc$^+$ LZ GC-B cells compared to their KO counterparts (**Fig. 1g**). This finding indicates

that these pathways are compromised or defective in KO cMyc$^+$ LZ GC-B cells. In line with the

enrichment map, genes associated with "OXPHOS", and "Mitochondrial translation" exhibited a

significant enrichment in WT cMyc$^+$ LZ GC-B cells compared to KO cMyc$^+$ LZ GC-B cells,

suggesting impaired functionality of these pathways in KO cMyc$^+$ LZ GC-B cells (**Fig. 1g**).

Collectively, our findings suggest that mitochondrial fitness is exclusively established in the LZ

during positive selection and that miR-155 expressed in the LZ is critical for maintaining

mitochondrial bioenergetic functions in GC-B cells.

MiR-155 regulates H3K36me2 levels in dividing LZ GC-B cells

To visualize the enriched pathways in KO cells, we generated a Cytoscape enrichment map

using RNA-seq data from cMyc[+] LZ GC-B cells. The enrichment map revealed that the gene

signature "Demethylation" was the only pathway significantly enriched in KO cMyc[+] LZ GC-B

cells when compared to WT cells (**Fig. 2a**). The signature "Demethylation" consisted of four

pathways: "Histone demethylase activity", "Protein demethylation", "Demethylase activity",

and "Demethylation" (**Fig. 2a**). Therefore, we investigated whether there were any alterations in

histone post-translational modifications (PTMs) in KO GC-B cells compared to WT GC-B cells.

To this end, we employed a liquid chromatography with tandem mass spectrometry (LC-

MS/MS) technique that allows unbiased identification and quantitative profiling of histone

PTMs [34,35]. This method successfully quantified the abundance of histone PTMs in both WT and

KO LZ GC-B cells, including methylation and acetylation on histone H1.4, H3 and H4. The

analysis showed a significant decrease in di-methylation of histone H3.1 at lysine 36 (K36me2)

and an increase in unmethylated histone H3 at lysine 23 (K23me0) in KO LZ GC-B cells

compared to WT LZ GC-B cells (**Fig. 2b** and **Extended Data Fig. 2a**). We also observed a

decrease in acetylation of H3 at K23 (K23ac) (**Extended Data Fig. 2a**), which is linked to

transcriptional activation [36]. Acetylation of H3 at K23 can be catalysed by cyclic adenosine

monophosphate response element binding protein binding protein (CREBBP), monocytic

leukemia zinc finger protein (MOZ) complex and its paralog monocytic leukemia zinc finger

related factor (MORF) complex [36,37]. However, there were no differentially expressed genes

(DEGs) in the bulk RNA-seq dataset between KO and WT cMyc[+] LZ-GC-B cells for any genes

encoding protein products that are part of these complexes (**Supplementary Table 2**). Therefore,

it was unlikely that the absence of miR-155 was directly responsible for the decreased H3K23ac

levels in KO LZ GC-B cells. Because the reduction of H3K23ac could lead to increased levels

of unmethylated H3K23, the elevation in H3K23me0 levels appears to be not directly caused by

the lack of miR-155 either. Thus, the decreased levels of K36me2 on histone H3.1 appeared to

be the sole altered histone methylation directly impacted by the absence of miR-155-mediated

regulation in KO cMyc[+] LZ GC-B cells.

The reduction of K36me2 levels was specific to DNA replication-dependent histone H3.1 and

was not observed in DNA replication-independent histone H3.3 (**Fig. 2b**). This suggests that cell

division is required for the depletion of H3K36me2. Previous studies have indicated that the loss

of H3K36me2 is associated with a decrease in acetylation of histone H3 at K27 (K27ac) [38].

There was decreased K27ac on histone H3.1 in KO LZ GC-B cells relative to WT cells, while

no such change was observed on histone H3.3 (**Fig. 2b**). The decrease in K36me2 on histone

H3.1 may lead to increased levels of monomethylated K36 (K36me1) on histone H3.1. While no

elevation of K36me1 was detected on histone H3.1 in KO LZ GC-B cells relative to their WT

cells, a significant increase in this histone PTM was observed in KO DZ GC-B cells compared

to their WT counterparts (**Fig. 2b** and **Extended Data Fig. 2b**). This could be attributed to the

migration of dividing LZ GC-B cells to the DZ, where chromatin marks could potentially be

deposited on newly synthesized histones. Apart from K36me1, no other changes were observed

in the histone PTMs at K36 of histone H3.1 in DZ GC-B cells between WT and KO cells

(**Extended Data Fig. 2b**). These findings suggest that miR-155 regulates the levels of K36me2

of H3.1 during cell division in LZ GC-B cells. Given that LZ GC-B cells are generally quiescent,

these dividing cells likely represent cMyc[+] LZ GC-B cells [7].

To examine the effect of miR-155-mediated regulation on histone H3K36me2 levels in cMyc$^+$

GC-B cells, we performed immunostaining of spleen sections from SRBC-immunised, *Mir155-*

sufficient and *Mir155*-deficient mice. Within the GL7-positive GC areas, a correlation in co-

expression of MYC and H3K36me2 was observed in both groups to some extents (**Fig. 2c,d**).

However, the Pearson's correlation coefficients with 95% confidence interval showed no

overlap between the values of WT and KO groups (r = 0.574 – 0.596 and 0.344 – 0.384, for WT

and KO, respectively), indicating a stronger correlation in the WT group than the KO group (**Fig.**

2d). To investigate the difference in H3K36me2 levels between WT and KO MYC$^+$ GC-B cells,

we compared the relative mean intensity of H3K36me2 in cells within the top 5% of MYC

positive cells from the total cell population shown in **Fig. 2d**. The analysis revealed a significant

reduction in H3K36me2 levels in KO MYC$^+$ GC-B cells compared to the corresponding WT

cells (**Fig. 2e**), which is consistent with the results derived from the MS analysis (**Fig. 2b**). We

also performed immunofluorescence-based correlation analysis between MYC and H3K36me1

or H3K36me3. However, the correlation in co-expression of MYC and these histone PTMs

exhibited only minor variations when comparing KO and WT MYC$^+$ GC-B cells (**Extended**

Data Fig. 2c,d). These results suggest that the co-expression pattern between MYC and

H3K36me1 or H3K36me3 remains largely unchanged, unlike the substantial changes observed

in H3K36me2. Collectively, these results provide compelling evidence for the crucial role of

miR-155 as an epigenetic modulator in histone PTMs. Notably, miR-155 predominantly

regulates the levels of H3K36me2 in dividing cMyc$^+$ LZ GC-B cells, potentially this may

influence mitochondrial functions in GC-B cells.

MiR-155 primarily controls *Kdm2a* expression in high-affinity cMyc$^+$ LZ GC-B cells

Demethylation of H3K36me2 can be catalysed by the lysine demethylases (KDMs), *Kdm2a*,

Kdm2b, *Kdm4a*, *Kdm4b* and *Kdm4c* [39]. We examined whether any of mRNAs encoding these

enzymes was differentially expressed between WT and KO cMyc$^+$ LZ GC-B cells. Expression

of *Kdm2a* was significantly higher in KO cMyc$^+$ LZ GC-B cells compared to WT cMyc$^+$ LZ

GC-B cells, while the others (*Kdm2b*, *Kdm4b* and *Kdm4c*) exhibited similar expression levels

between the two groups (**Fig. 3a**). In contrast, the expression of *Kdm4a* was found to be

decreased in KO cMyc$^+$ LZ GC-B cells compared to WT cMyc$^+$ LZ GC-B cells (**Fig. 3a** and

Supplementary Table 2). Next, we examined whether *Kdm2a* mRNA is predicted as a target of

miR-155. By utilizing miR prediction tools, we identified *Kdm2a* as one of the 18 genes that

were predicted to be targets of miR-155 and showed differential expression between WT and

KO cMyc$^+$ LZ GC-B cells (**Fig. 3a** and **Supplementary Table 2**). *Kdm2a* has one 8-mer miR-

155-binding site at 3' UTR that is conserved across multiple species, including human, mouse,

and rat (**Fig. 3b**).

To investigate the relationship between KDM2A and miR-155, we examined the co-expression

of KDM2A and MYC by immunostaining tissue sections derived from *Mir155*-sufficient and

Mir155-deficient mice using antibodies against IgD, MYC and KDM2A. This revealed a strong

correlation in co-expression of these two proteins in both groups (r = 0.775 – 0.788 and 0.720 –

0.740, for WT and KO, respectively) (**Fig. 3c,d**). Therefore, we concluded that KDM2A is

mainly expressed in positively selected cMyc$^+$ GC-B cells. To examine the degree and cell

subpopulations expressing KDM2A, we performed flow cytometric analysis of GC-B cells

derived from SW$_{HEL}$ adoptive transfer system. In WT mice, the analysis revealed a ~1.3-fold

increase in KDM2A expression in cMyc$^+$ LZ GC-B cells compared to cMyc$^-$ LZ GC-B cells

(**Fig. 3e**), indicating that positive selection signals can induce its expression. The highest

expression of KDM2A was observed in cMyc$^+$ DZ GC-B cells. A similar expression pattern was

observed in KO GC-B cell subpopulations. In all GC-B cell subpopulations, KDM2A

expression was significantly higher in KO cells compared to their WT counterparts (**Fig. 3e** and

Extended Data Fig. 3b).

The SW$_{HEL}$ system enables the distinction between high and low affinity B cells following

HEL$^{3\times}$ immunisation. When comparing high and low affinity clones within each WT GC-B cell

subpopulation, there were no significant differences in the level of KDM2A (**Fig. 3f**). In contrast,

the lack of miR-155 led to a substantial increase of KDM2A expression in high-affinity clones

compared to low-affinity clones within the cMyc$^+$ LZ GC-B cell subpopulation. As a result,

there was a marked elevation in KDM2A expression in high-affinity clones of KO cells

compared to WT cells (**Fig. 3f**). Collectively, these results suggest that KDM2A is preferentially

induced in high-affinity clones within the cMyc$^+$ LZ GC-B cell subpopulation during positive

selection, and therefore, miR-155 is required to dampen KDM2A expression in these cells.

Hypoxia induces KDM2A in LZ GC-B cells

Hypoxia induces *Kdm2a* expression at both transcript and protein levels [40,41]. Given that

previous studies have shown that the LZ of GCs is hypoxic [10-12], we hypothesized that miR-155

counteracts the hypoxia-induced increase of KDM2A expression: i.e., in the absence of miR-155,

KDM2A expression is enhanced under hypoxic conditions. Since the concept that the LZ has a

more hypoxic microenvironment than the DZ remains a subject of controversy [8,42], we

quantified the degree of hypoxia that GC-B cells were experiencing using the hypoxia detecting

dye, MAR (see methods). Flow cytometric analysis revealed that a bimodal distribution of MAR

fluorescence intensity in non-GC-B cells. This potentially reflects their location within the tissue,

as it is generally considered that regions adjacent to blood vessels are less hypoxic, whereas

areas distant from blood vessels experience more hypoxia [43]. Here, LZ GC-B cells exhibited

higher MAR fluorescence intensity than DZ GC-B cells (**Fig. 4a**), indicating greater levels of

hypoxia, in line with previous studies [10-12].

To examine the effect of hypoxia on KDM2A expression in B cells, we cultured splenic B cells

in the presence of anti-IgM, anti-CD40 and IL-4, which mimicked the combination of positive

selection signals derived from BCR and T cell help. These cells were cultured in either normal

oxygen concentration (21%, normoxic) or low oxygen concentration (1%, hypoxic), and

KDM2A expression was examined by flow cytometry. Following a two-day activation with

positive selection mimics under normoxic conditions, both WT and KO B cells showed

increased KDM2A expression (**Fig. 4b**). However, there was no significant difference in

KDM2A levels between WT and KO cells under normoxic conditions (**Fig. 4b**). Under hypoxic

conditions, the expression of KDM2A was markedly increased, resulting in higher expression of

KDM2A in both WT and KO B cells compared to normoxic conditions (**Fig. 4b**). By contrast,

in hypoxic cultures, KDM2A expression was ~1.6-fold greater in the KO B cells than WT B

cells (**Fig. 4b**). This difference in expression was attributed to the increased expression of miR-

155 under hypoxic conditions (**Extended Data Fig. 4a**), consistent with previous reports [44,45].

Therefore, hypoxia was necessary for observing differential levels of KDM2A between KO and

WT *in vitro* cultured B cells. These findings suggest that miR-155 exerts a strong repressive

effect on *Kdm2a* primarily under hypoxic conditions.

The regulatory activity of miR-155 is revealed under hypoxia

To examine the impact of hypoxia on cell division, we cultured CellTraceViolet (CTV)-labelled

B cells from *Mir155*-sufficient and *Mir155*-deficient mice with positive selection mimics in

either normoxic or hypoxic conditions. As expected, hypoxia suppressed cell proliferation

regardless of genotype (**Fig. 4c**). However, whereas both WT and KO cells showed similar

levels of division in normoxic cultures, miR-155 KO B cells exhibited significantly less division

than WT B cells under hypoxic conditions (**Fig. 4c**). The results highlight the critical role of

miR-155 for cell proliferation under hypoxic conditions.

To investigate the influence of hypoxia on mitochondrial ROS production, we used non-

fluorescent MitoTracker Orange CM-H$_2$ TMRos that becomes fluorescent upon oxidization.

After a two-day culture period under either hypoxic or normoxic conditions, we stained WT and

KO B cells with MitoTracker dyes. WT B cells produced lower amounts of mitochondrial ROS

under hypoxic conditions (**Fig. 4d,e**). In contrast, hypoxic KO cultures had significantly higher

amounts of mitochondrial ROS than normoxic KO cultures (**Fig. 4d,e**), indicating mitochondrial

dysfunction in these cells. Consequently, KO B cells exhibited a significantly increased ROS

production compared to WT B cells under hypoxic conditions, while no such difference was

observed under normoxic conditions (**Fig. 4d,e**). To further investigate mitochondrial integrity

in KO B cells, we assessed energy metabolic states coupled to mitochondrial functions. To this

end, we measured the abundance of adenosine triphosphate (ATP) produced through metabolic

processes. After two days of culturing WT and KO B cells under either normoxic or hypoxic

conditions, we further cultured the cells in the following media supplemented with either

glucose, glucose plus oligomycin, galactose, or galactose plus oligomycin. To directly compare

energy metabolic capacities of cells, all cells were cultured in hypoxic conditions for 2 hours

using the indicated media. Cells can use both glycolysis and OXPHOS to produce ATP when

cultured in glucose-supplemented medium, while in medium supplemented with glucose and

oligomycin, an inhibitor of OXPHOS, cells rely solely on glycolysis for ATP production. Under

hypoxic conditions, both WT and KO cells exhibited a reduction of ATP abundance when

cultured in glucose-supplemented medium, compared to their normoxic counterparts (**Extended**

Data Fig. 4b). This result is consistent with previous reports showing reduced energy

production under hypoxia [46,47]. When B cells were cultured in medium supplemented with both

glucose and oligomycin, the relative abundance of ATP was similar to corresponding cultures

supplemented with glucose alone across all cell types (**Fig. 4f**), indicating that the cells were

able to survive by relying on their glycolytic capacity without the need for OXPHOS, in

accordance with the well-known "Crabtree effect" [48,49]. In contrast, cells cultured in galactose-

supplemented medium must use OXPHOS to generate ATP [50]. In this culture condition, we

observed a reduction in ATP abundance in both WT and KO B cells under hypoxia compared to

normoxia.

Furthermore, while ATP production through OXPHOS was compromised in hypoxic cultures of

both WT and KO cells, the impairment was more severe in KO B cells (**Fig. 4f**). ATP

production was abrogated when WT or KO B cells were cultured in medium containing

galactose and oligomycin regardless of oxygen tension (**Fig. 4f**). These results suggest that miR-

155 plays a critical role in the metabolic reprogramming of activated B cells by regulating

OXPHOS under hypoxic conditions. Collectively, the results demonstrate that the regulatory

activity of miR-155 in modulating mitochondrial function and energy metabolism in activated B

cells is manifested in a hypoxic environment.

MiR-155 optimises nuclear mitochondrial gene expression through KDM2A-mediated

control of H3K36me2 levels

To investigate how *Mir155*-deficiency impacts genome-wide binding of KDM2A and

H3K36me2 distribution, we performed ChIP-seq assays using hypoxic B cell cultures. The

genomic distribution of KDM2A and H3K36me2 peaks showed significant differences between

WT and KO B cells (**Fig. 5a**). In KO B cells compared to WT B cells, there was a proportional

increase in KDM2A peaks within the promoter/transcriptional start site (TSS) regions (WT =

26.5%, KO = 34.5%), accompanied by a concurrent reduction of H3K36me2 peaks in the same

regions (WT = 4.59%, KO = 3.77%) (**Fig. 5a**). Consistent with a previous report showing that

KDM2A preferentially binds to CGIs islands (CGIs) primarily located at gene promoters [51], we

detected significant enrichment of KDM2A peaks specifically at the TSS-proximal regions of

genes containing CGIs (**Fig. 5b**). Furthermore, we observed a concurrent depletion in

H3K36me2 signals within these regions of CGI containing genes in both WT and KO B cells

(**Fig. 5b**). To identify distinctive patterns of KDM2A binding and H3K36me2 marks, we

examined the peak positions in relation to the nearest TSS (**Fig. 5c**). These plots revealed that a

higher density of KDM2A peaks in KO B cells at TSS-proximal regions compared to WT B

cells (**Fig. 5c**). Accordingly, H3K36me2 peak density was reduced in KO B cells relative to WT

B cells at these same regions and in the intragenic regions proximal to the TSS (**Fig. 5c**). Our

findings indicate that the absence of miR-155 leads to an elevated expression of KDM2A, which

in turn triggers enhanced binding of KDM2A to TSS-proximal regions and a concomitant

depletion of H3K36me2 in those regions as well as in the intragenic regions.

To elucidate the biological functions of genes bound by KDM2A, we investigated the KDM2A

target genes that displayed differential binding between WT and KO B cells and identified

"WT-enriched" and "KO-enriched" KDM2A peaks (**Supplementary Table 3**). Functional

annotation clustering analysis with DAVID [52] showed that genes with "WT-enriched" or "KO-

enriched" Kdm2a peaks were strongly associated with the regulation of mitochondrial functions

(**Fig. 5d**). These results suggest that KDM2A plays a role in regulating the expression of nuclear

mitochondrial genes during hypoxia. Among the genes with "WT-enriched" and "KO-enriched"

KDM2A peaks, we identified 102 and 30 genes encoding nuclear mitochondrial proteins listed

in MitoCarta 3.0 [53], respectively. Bulk RNA-seq analysis showed that five mitochondrial genes

with "KO-enriched" KDM2A peaks at promoter/TSS regions and two mitochondrial ribosomal

proteins with "KO-enriched" KDM2A peaks at the first intron, which corresponds to the

intragenic regions proximal to the TSS, were reduced in their expression in KO cMyc$^+$ LZ GC-

B cells compared to WT cMyc$^+$ LZ GC-B cells (**Fig. 5e** and **Supplementary Table 3**). The

nuclear mitochondrial genes with "KO-enriched" KDM2A peaks exhibited a concomitant

decrease in H3K36me2 levels (**Extended Data Fig. 5a**). These findings suggested that the

differential binding of KDM2A between WT and KO B cells affected the expression of nuclear

mitochondrial genes. Furthermore, among the five genes with "KO-enriched" KDM2A peaks

located in the promoter/TSS regions, *Ndufv3* and *Atp5l* are known to play an essential role in

OXPHOS [53]. Thus, the miR-155-*Kdm2a* axis has a critical function in regulating OXPHOS

coupled with mitochondrial bioenergetic functions in activated B cells under hypoxic conditions.

The miR-155-Kdm2a interaction regulates mitochondrial ROS and apoptosis in GC-B

cells

To validate the regulatory interaction of the miR-155-*Kdm2a* axis, we generated mice with a

mutated miR-155 binding site in the 3'UTR of *Kdm2a* (**Fig. 6a**). B cell development was

normal in homozygous mutant (hereafter called "MUT") mice (**Extended Data Fig. 6a**). WT

and MUT mice were immunised with SRBCs, and KDM2A expression in GC-B cells was

assessed by flow cytometry. In LZ GC-B cells of MUT mice, KDM2A expression exhibited a

~1.2-fold increase relative to their WT littermates (**Fig. 6b**). No significant difference in

KDM2A expression was observed in DZ GC-B cells. These results indicate that miR-155 limits

Kdm2a expression in LZ GC-B cells through its binding site on Kdm2a mRNA. To examine B

cell-intrinsic effects of the mutation, we performed adoptive transfer experiments by transferring

B cells isolated from the spleens of either WT or MUT mice into µMT mice, which lacks mature

B cells [54]. The mice were then immunised with SRBCs. Seven days later, all recipient mice gave

rise to GC-B cells, but the total number of DZ GC-B cells was reduced by ~29% in the MUT

background compared to that of WT, while the number of LZ GC-B cells was similar between

WT and MUT mice (**Fig. 6c** and **Extended Data Fig. 6b,c**).

We investigated the impact of the 3'UTR mutation on mitochondrial ROS and apoptosis in GC-

B cells by transferring B cells into μMT mice. The proportion of MitoSOX$^+$ cells was higher in

both LZ and DZ GC-B cells of MUT mice compared to WT (**Fig. 6d**). Moreover, the percentage

of active-Caspase3$^+$ cells were increased in DZ GC-B cells of MUT mice, when compared to

WT mice (**Fig. 6e**). The elevated apoptosis in GC-B cells from MUT mice compared to WT

mice provides an explanation for the decreased number of DZ GC-B cells depicted in **Fig. 6c**.

These results largely mirror the observations in *Mir155*-deficient mice. Moreover, the

overexpression of KDM2A in GC-B cells through lentiviral transduction resulted in an

increased percentage of MitoSOX$^+$ DZ GC-B cells compared to control cells (**Extended Data**

Fig. 6d). Taken together, these findings underscore the crucial role of miR-155 in repressing

Kdm2a to counteract the effects of hypoxia during the positive selection process.

Discussion

We demonstrated that miR-155 critically regulates H3K36me2 levels by directly repressing

histone demethylase *Kdm2a* during GC positive selection. This repression enabled GC-B cells

to reprogram their metabolism through the remodelling of mitochondria and to initiate cell

division in the hypoxic LZ microenvironment. Maintaining mitochondrial fitness during positive

selection in the LZ was essential for higher-affinity GC-B cells to undergo clonal expansion in

the DZ, consequently ensuring affinity maturation in GC responses. In the absence of miR-155,

dysfunctional mitochondria resulted in increased ROS production and subsequent apoptosis in

GC-B cells, which explains the compromised affinity maturation in *miR-155*-KO mice [24,55,56].

While *Mir155*-defcient mice show a compromised proliferation of GC-B cells, *in vitro* cultured

KO B cells exhibit normal DNA synthesis and cell proliferation rates regardless of stimuli [55-57].

Thus, it was unclear why these *in vitro* cultured B cells cannot recapitulate the phenotype

observed *in vivo*. In this study, we demonstrated that essential function of miR-155 is revealed

under hypoxia *in vitro* and reproduced the known *in vivo* phenotypes observed in KO B cells.

We have identified *Kdm2a* as a critical target gene of miR-155, which is induced under hypoxic

conditions. *Kdm2a* exhibits context-dependent effects on proliferation [20,21] and may act as a

negative regulator of proliferation in GC-B cells. Therefore, miR-155-mediated *Kdm2a*

repression enabled GC-B cells to initiate cell division under hypoxic conditions in the LZ during

positive selection.

Our findings demonstrated the critical role of miR-155 in facilitating the metabolic adaptation of

positively selected GC-B cells for cell division while mitigating hypoxia-driven stress in the LZ.

Our previous studies have shown that high-affinity clones display greater proliferative activity

than low-affinity clones within cMyc$^+$ LZ GC-B cell subpopulation [7]. Indeed, miR-155 exerted

its most potent suppressive effect on *Kdm2a* especially within the high-affinity clones of the

cMyc[+] LZ GC-B cell subpopulation (**Fig. 3f**). In agreement with this, *Mir155*-deficient mice

exhibit a selective loss of high-affinity GC-B cells due to apoptosis [24,55,56]. In the absence of

miR-155, cMyc[+] LZ GC-B cells with higher levels of KDM2A experience detrimental effects

such as increased ROS production due to dysfunctional mitochondrial and subsequent cell death

(**Fig. 1b,c** and **Extended Data Fig. 6d,e**). Despite the downregulation of miR-155 expression in

cMyc[+] DZ GC-B cells and its absence in cMyc[-] DZ GC-B cells, there was a persistent increase

in ROS production in these cells when miR-155 was absent. This may be because DZ GC-B

cells prioritize energy use for mitosis rather than mitochondrial repair. Therefore, GC-B cells

appear to be incapable of restoring mitochondrial function after migration to the DZ. The

findings underscore the importance of the miR-155-*Kdm2a* regulatory axis in the LZ, which

promoted metabolic reprogramming and gradual enhancement of OXPHOS in positively

selected GC-B cells. As a result, positively selected GC-B cells could cope with the higher rate

of cell division in the DZ by utilizing ATP provided by OXPHOS in remodelled mitochondria.

These results are consistent with a previous observation that GC-B cells with higher-affinity

require increased OXPHOS activity to undergo extensive proliferation [9].

KDM2A expression was upregulated >1.5-fold in *Mir155*-KO mice compared to WT mice in

both cMyc[+] and cMyc[-] DZ GC-B cells (**Fig. 3e**). However, the expression level of KDM2A in

DZ GC-B cells of MUT mice was comparable to that of WT mice. Firstly, there is evidence for

a 6-mer miR-155 binding site in the 3'UTR of mouse *Kdm2a* [58]. Since this 6-mer binding site is

intact in MUT mice, miR-155 may partially repress *Kdm2a* transcripts through this region,

inhibiting the KDM2A elevation in MUT mice. Alternatively, other miR-155 target(s) could

contribute to the induction of KDM2A. *Hif1a* can be directly targeted by miR-155 [58] and

potentially influences the expression of KDM2A because KDM2A may be induced partially in a

HIF-1α-dependent manner under hypoxic conditions [40]. Given that *Hif1a* is repressed by miR-

155 in a similar manner in both WT and MUT mice, the lower HIF-1α expression in MUT B

cells compared to KO B cells may limit the efficiency of KDM2A induction in MUT mice.

MiRs generally have multiple targets. We identified 18 mRNAs that are predicted miR-155

targets in cMyc[+] LZ GC-B cells and exhibit increased expression in the absence of miR-155

(**Extended Data Fig. 3a**). One of the molecules, *Jarid2* was previously discovered as a miR-

155 target in positively selected GC-B cells, and its overexpression has been shown to increase

apoptosis in LZ GC-B cells [24]. JARID2 functions as an accessory component of the Polycomb

Repressive Complex 2 (PRC2), which catalyses H3K27me3 [59], thereby epigenetically regulating

gene expression. Therefore, the modulation of PRC2 activity by JARID2 potentially contributes

to elevated apoptosis in these cells. Another molecule, F-Box Only Protein 11 (*Fbxo11*), was

also among the identified 18 genes, and is a part of a multi-subunit E3 ubiquitin-ligase protein

complex that catalyze the ubiquitination of proteins destined for proteasomal degradation. GC-B

cell-specific deletion of *Fbxo11* increases the number of GC-B cells, particularly DZ GC-B cells,

suggesting that this molecule is associated with GC-B cells proliferation [60]. Therefore, the

phenotypic alterations observed in *Mir155*-deficient mice may result from the cumulative effects

of several causal targets acting together with *Kdm2a*.

Collectively, our findings demonstrate that miR-155 acts as an epigenetic regulator to limit the

activity of *Kdm2a* on target genes, thereby maintaining appropriate levels of H3K36me2. The

precise control of H3K36me2 levels during positive selection is vital for the optimal expression

of nuclear mitochondrial genes in GC-B cells under hypoxic conditions, thus facilitating

mitochondrial remodelling. The resultant metabolic reprogramming is indispensable for the

survival and robust proliferation of GC-B cells within the DZ. Our findings provide insights that

could pave the way for the development of novel therapeutics targeting histone PTMs to control

GC responses in the context of vaccination and lymphomagenesis.

Main references

Berek, C., Berger, A. & Apel, M. Maturation of the immune response in germinal centers. *Cell* **67**, 1121-1129 (1991).

Jacob, J., Kelsoe, G., Rajewsky, K. & Weiss, U. Intraclonal generation of antibody mutants in germinal centres. *Nature* **354**, 389-392 (1991). https://doi.org:10.1038/354389a0

Kepler, T. B. & Perelson, A. S. Cyclic re-entry of germinal center B cells and the efficiency of affinity maturation. *Immunol Today* **14**, 412-415 (1993). https://doi.org:0167-5699(93)90145-B [pii]10.1016/0167-5699(93)90145-B

Nakagawa, R. & Calado, D. P. Positive Selection in the Light Zone of Germinal Centers. *Front Immunol* **12**, 661678 (2021). https://doi.org:10.3389/fimmu.2021.661678

Calado, D. P. *et al.* The cell-cycle regulator c-Myc is essential for the formation and maintenance of germinal centers. *Nat Immunol* **13**, 1092-1100 (2012). https://doi.org:10.1038/ni.2418 ni.2418 [pii]

Dominguez-Sola, D. *et al.* The proto-oncogene MYC is required for selection in the germinal center and cyclic reentry. *Nat Immunol* **13**, 1083-1091 (2012). https://doi.org:10.1038/ni.2428 ni.2428 [pii]

Nakagawa, R. *et al.* Permissive selection followed by affinity-based proliferation of GC light zone B cells dictates cell fate and ensures clonal breadth. *Proc Natl Acad Sci U S A* **118** (2021). https://doi.org:10.1073/pnas.2016425118

Weisel, F. J. *et al.* Germinal center B cells selectively oxidize fatty acids for energy while conducting minimal glycolysis. *Nat Immunol* **21**, 331-342 (2020). https://doi.org:10.1038/s41590-020-0598-4

Chen, D. *et al.* Coupled analysis of transcriptome and BCR mutations reveals role of OXPHOS in affinity maturation. *Nat Immunol* **22**, 904-913 (2021). https://doi.org:10.1038/s41590-021-00936-y

Jellusova, J. *et al.* Gsk3 is a metabolic checkpoint regulator in B cells. *Nat Immunol* **18**, 303-312 (2017). https://doi.org:10.1038/ni.3664

Cho, S. H. *et al.* Germinal centre hypoxia and regulation of antibody qualities by a hypoxia response system. *Nature* **537**, 234-238 (2016). https://doi.org:10.1038/nature19334

Li, L. *et al.* Regulation of humoral immune response by HIF-1alpha-dependent metabolic reprogramming of the germinal center reaction. *Cell Immunol* **367**, 104409 (2021). https://doi.org:10.1016/j.cellimm.2021.104409

Kierans, S. J. & Taylor, C. T. Regulation of glycolysis by the hypoxia-inducible factor (HIF): implications for cellular physiology. *J Physiol* **599**, 23-37 (2021). https://doi.org:10.1113/JP280572

Goda, N. *et al.* Hypoxia-inducible factor 1alpha is essential for cell cycle arrest during hypoxia. *Mol Cell Biol* **23**, 359-369 (2003). https://doi.org:10.1128/MCB.23.1.359-369.2003

Koshiji, M. *et al.* HIF-1alpha induces cell cycle arrest by functionally counteracting Myc. *EMBO J* **23**, 1949-1956 (2004). https://doi.org:10.1038/sj.emboj.7600196

Liu, L. *et al.* Hypoxia-induced energy stress regulates mRNA translation and cell growth. *Mol Cell* **21**, 521-531 (2006). https://doi.org:10.1016/j.molcel.2006.01.010

Ersching, J. *et al.* Germinal Center Selection and Affinity Maturation Require Dynamic Regulation of mTORC1 Kinase. *Immunity* **46**, 1045-1058 e1046 (2017). https://doi.org:10.1016/j.immuni.2017.06.005

Finkin, S., Hartweger, H., Oliveira, T. Y., Kara, E. E. & Nussenzweig, M. C. Protein Amounts of the MYC Transcription Factor Determine Germinal Center B Cell Division Capacity. *Immunity* **51**, 324-336 e325 (2019). https://doi.org:10.1016/j.immuni.2019.06.013

Mayer, C. T. *et al.* The microanatomic segregation of selection by apoptosis in the germinal center. *Science* **358** (2017). https://doi.org:10.1126/science.aao2602

Kawakami, E., Tokunaga, A., Ozawa, M., Sakamoto, R. & Yoshida, N. The histone demethylase Fbxl11/Kdm2a plays an essential role in embryonic development by repressing cell-cycle regulators. *Mech Dev* **135**, 31-42 (2015). https://doi.org:10.1016/j.mod.2014.10.001

Tanaka, Y. *et al.* JmjC enzyme KDM2A is a regulator of rRNA transcription in response to starvation. *EMBO J* **29**, 1510-1522 (2010). https://doi.org:10.1038/emboj.2010.56

Esposito, L. A., Melov, S., Panov, A., Cottrell, B. A. & Wallace, D. C. Mitochondrial disease in mouse results in increased oxidative stress. *Proc Natl Acad Sci U S A* **96**, 4820-4825 (1999). https://doi.org:10.1073/pnas.96.9.4820

Redza-Dutordoir, M. & Averill-Bates, D. A. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim Biophys Acta* **1863**, 2977-2992 (2016). https://doi.org:10.1016/j.bbamcr.2016.09.012

Nakagawa, R. *et al.* MicroRNA-155 controls affinity-based selection by protecting c-MYC[+] B cells from apoptosis. *J Clin Invest* **126**, 377-388 (2016). https://doi.org:10.1172/JCI82914

Phan, T. G. *et al.* B cell receptor-independent stimuli trigger immunoglobulin (Ig) class switch recombination and production of IgG autoantibodies by anergic self-reactive B cells. *J Exp Med* **197**, 845-860 (2003). https://doi.org:10.1084/jem.20022144 jem.20022144 [pii]

Huang, C. Y., Bredemeyer, A. L., Walker, L. M., Bassing, C. H. & Sleckman, B. P. Dynamic regulation of c-Myc proto-oncogene expression during lymphocyte development revealed by a GFP-c-Myc knock-in mouse. *Eur J Immunol* **38**, 342-349 (2008). https://doi.org:10.1002/eji.200737972

Kwon, K. *et al.* Instructive role of the transcription factor E2A in early B lymphopoiesis and germinal center B cell development. *Immunity* **28**, 751-762 (2008). https://doi.org:10.1016/j.immuni.2008.04.014 S1074-7613(08)00231-8 [pii]

Laidlaw, B. J., Duan, L., Xu, Y., Vazquez, S. E. & Cyster, J. G. The transcription factor Hhex cooperates with the corepressor Tle3 to promote memory B cell development. *Nat Immunol* **21**, 1082-1093 (2020). https://doi.org:10.1038/s41590-020-0713-6

Suan, D. *et al.* CCR6 Defines Memory B Cell Precursors in Mouse and Human Germinal Centers, Revealing Light-Zone Location and Predominant Low Antigen Affinity. *Immunity* **47**, 1142-1153 e1144 (2017). https://doi.org:10.1016/j.immuni.2017.11.022

Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome Biol* **20**, 59 (2019). https://doi.org:10.1186/s13059-019-1663-x

Mroczek, S. & Kufel, J. Apoptotic signals induce specific degradation of ribosomal RNA in yeast. *Nucleic Acids Res* **36**, 2874-2888 (2008). https://doi.org:10.1093/nar/gkm1100

Toboso-Navasa, A. *et al.* Restriction of memory B cell differentiation at the germinal center B cell positive selection stage. *J Exp Med* **217** (2020). https://doi.org:10.1084/jem.20191933

Huang, G., Li, H. & Zhang, H. Abnormal Expression of Mitochondrial Ribosomal Proteins and Their Encoding Genes with Cell Apoptosis and Diseases. *Int J Mol Sci* **21** (2020). https://doi.org:10.3390/ijms21228879

Camarillo, J. M. *et al.* Coupling Fluorescence-Activated Cell Sorting and Targeted Analysis of Histone Modification Profiles in Primary Human Leukocytes. *J Am Soc Mass Spectrom* **30**, 2526-2534 (2019). https://doi.org:10.1007/s13361-019-02255-x

Garcia, B. A. *et al.* Chemical derivatization of histones for facilitated analysis by mass spectrometry. *Nat Protoc* **2**, 933-938 (2007). https://doi.org:10.1038/nprot.2007.106

Zhao, Y. & Garcia, B. A. Comprehensive Catalog of Currently Documented Histone Modifications. *Cold Spring Harb Perspect Biol* **7**, a025064 (2015). https://doi.org:10.1101/cshperspect.a025064

Klein, B. J. *et al.* Histone H3K23-specific acetylation by MORF is coupled to H3K14 acylation. *Nat Commun* **10**, 4724 (2019). https://doi.org:10.1038/s41467-019-12551-5

Chen, H. *et al.* H3K36 dimethylation shapes the epigenetic interaction landscape by directing repressive chromatin modifications in embryonic stem cells. *Genome Res* **32**, 825-837 (2022). https://doi.org:10.1101/gr.276383.121

Tian, X. *et al.* Histone lysine-specific methyltransferases and demethylases in carcinogenesis: new targets for cancer therapy and prevention. *Curr Cancer Drug Targets* **13**, 558-579 (2013). https://doi.org:10.2174/1568009611313050007

Batie, M., Druker, J., D'Ignazio, L. & Rocha, S. KDM2 Family Members are Regulated by HIF-1 in Hypoxia. *Cells* **6** (2017). https://doi.org:10.3390/cells6010008

Xia, X. *et al.* Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc Natl Acad Sci U S A* **106**, 4260-4265 (2009). https://doi.org:10.1073/pnas.0810067106

Abbott, R. K. *et al.* Germinal Center Hypoxia Potentiates Immunoglobulin Class Switch Recombination. *J Immunol* **197**, 4014-4020 (2016). https://doi.org:10.4049/jimmunol.1601401

Zaidi, M., Fu, F., Cojocari, D., McKee, T. D. & Wouters, B. G. Quantitative Visualization of Hypoxia and Proliferation Gradients Within Histological Tissue Sections. *Front Bioeng Biotechnol* **7**, 397 (2019). https://doi.org:10.3389/fbioe.2019.00397

Bruning, U. *et al.* MicroRNA-155 promotes resolution of hypoxia-inducible factor 1alpha activity during prolonged hypoxia. *Mol Cell Biol* **31**, 4087-4096 (2011). https://doi.org:10.1128/MCB.01276-10

Wan, G. *et al.* Hypoxia-induced MIR155 is a potent autophagy inducer by targeting multiple players in the MTOR pathway. *Autophagy* **10**, 70-79 (2014). https://doi.org:10.4161/auto.26534

Cho, H. Y., Loreti, E., Shih, M. C. & Perata, P. Energy and sugar signaling during hypoxia. *New Phytol* **229**, 57-63 (2021). https://doi.org:10.1111/nph.16326

Papandreou, I., Cairns, R. A., Fontana, L., Lim, A. L. & Denko, N. C. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* **3**, 187-197 (2006). https://doi.org:10.1016/j.cmet.2006.01.012

Crabtree, H. G. Observations on the carbohydrate metabolism of tumours. *Biochem J* **23**, 536-545 (1929). https://doi.org:10.1042/bj0230536

Rodriguez-Enriquez, S., Juarez, O., Rodriguez-Zavala, J. S. & Moreno-Sanchez, R. Multisite control of the Crabtree effect in ascites hepatoma cells. *Eur J Biochem* **268**, 2512-2519 (2001). https://doi.org:10.1046/j.1432-1327.2001.02140.x

Marroquin, L. D., Hynes, J., Dykens, J. A., Jamieson, J. D. & Will, Y. Circumventing the Crabtree effect: replacing media glucose with galactose increases susceptibility of HepG2 cells to mitochondrial toxicants. *Toxicol Sci* **97**, 539-547 (2007). https://doi.org:10.1093/toxsci/kfm052

Blackledge, N. P. *et al.* CpG islands recruit a histone H3 lysine 36 demethylase. *Mol Cell* **38**, 179-190 (2010). https://doi.org:10.1016/j.molcel.2010.04.009

Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009). https://doi.org:10.1038/nprot.2008.211

Rath, S. *et al.* MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic Acids Res* **49**, D1541-D1547 (2021). https://doi.org:10.1093/nar/gkaa1011

Kitamura, D., Roes, J., Kuhn, R. & Rajewsky, K. A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin mu chain gene. *Nature* **350**, 423-426 (1991). https://doi.org:10.1038/350423a0

Thai, T. H. *et al.* Regulation of the germinal center response by microRNA-155. *Science* **316**, 604-608 (2007). https://doi.org:316/5824/604 [pii] 10.1126/science.1141229

Vigorito, E. *et al.* microRNA-155 regulates the generation of immunoglobulin class-switched plasma cells. *Immunity* **27**, 847-859 (2007). https://doi.org:S1074-7613(07)00503-1 [pii] 10.1016/j.immuni.2007.10.009

Rodriguez, A. *et al.* Requirement of bic/microRNA-155 for normal immune function. *Science* **316**, 608-611 (2007). https://doi.org:316/5824/608 [pii] 10.1126/science.1139253

Hsin, J. P., Lu, Y., Loeb, G. B., Leslie, C. S. & Rudensky, A. Y. The effect of cellular context on miR-155-mediated gene regulation in four major immune cell types. *Nat Immunol* **19**, 1137-1145 (2018). https://doi.org:10.1038/s41590-018-0208-x

Pasini, D. *et al.* JARID2 regulates binding of the Polycomb repressive complex 2 to target genes in ES cells. *Nature* **464**, 306-310 (2010). https://doi.org:10.1038/nature08788 nature08788 [pii]

Schneider, C. *et al.* FBXO11 inactivation leads to abnormal germinal-center formation and lymphoproliferative disease. *Blood* **128**, 660-666 (2016). https://doi.org:10.1182/blood-2015-11-684357

Figures and Figure legends

Fig. 1. MiR-155 is required for promoting mitochondrial fitness in GC-B cells

a, Experimental design for all the experiments for Fig. 1. **b**, Percentage of MitoSOX[+] GC-B

cells 5 days after HEL[3×]-SRBC immunisation. Pooled from three independent experiments (WT

n=12; KO n=12). Student's *t*-test. **c**, Percentage of active-Caspase 3[+] GC-B cells at 7 days after

HEL[3×]-SRBC immunisation. Pooled from two independent experiments (WT n=18; KO n=19)

Student's *t*-test. **d**, The PAGA algorithm generates a topology-preserving map of single cells

along with the sequential changes of gene expression. PAGA graphs were constructed using a k-

nearest neighbors' algorithm to show relationships between clusters. The Ball-and-stick

representation displays PAGA connectivity among GC-B cell clusters identified though scRNA-

seq. Ball size represents cluster size, and edge thickness indicates connectivity between clusters.

Normalized gene expression for *Cxcr4* or *Myc* is shown in the PAGA graph. **e**, Predicted relationship between the clusters identified in scRNA-seq of GC-B cells. **f**, Violin plots showing gene signature scores for "Mitochondrion" and "OXPHOS". The bar represents median value. Wilcoxon rank sum test, *p<0.05, ****p<0.00005, n.s., not significant. **g**, Enrichment map visualizing GSEA results of the bulk RNA-seq data from WT and KO cMyc$^+$ LZ GC-B cells. GC-B cells were sorted 5 days after HEL$^{3\times}$-SRBC immunisation. WT-enriched pathways (q<0.0001) were used. Nodes represent gene sets, and edges represent mutual overlap. Highly redundant gene sets were grouped as clusters (left). ES; enriched score, FDR; false discovery rate. Unless otherwise stated, mean ± SEM is indicated. *P<0.05, **p<0.01, *** p<0.001, ****p<0.0001.

1
Fig. 2. MiR-155 regulates H3K36me2 levels in dividing LZ GC-B cells

a, Enrichment map visualizing GSEA results of bulk RNA-seq data from WT and KO cMyc[+]

LZ GC-B cells. KO-enriched pathways (q<0.1) were used. **b**, The relative abundance of histone

PTMs, including histone H3.1 K36, H3.3 K36, H3.1 K27 and H3.3 K27 in LZ GC-B cells from

WT and KO mice. Donor-derived GC-B cells were sorted 5 days after HEL[3×]-SRBC

immunisation for analysis. Triplicate samples were used for both WT and KO backgrounds.

Mean ± SEM is indicated. Student's *t*-test. **c**, Splenic sections of C57BL/6 mice, in the presence

and absence of miR-155, seven days after SRBC immunisation were stained for GL7 (green),

MYC (red) and H3K36me2 (blue). Yellow wedges represent cells that are positive for MYC but

negative for KDM2A. Scale bar, 50 µm. Representative images from two experiments (WT n=4;

KO n=4). **d**, Scatter plots show the mean intensity (MI) of MYC and H3K36me2 per segmented

cell within GCs of spleen sections obtained from immunised WT and KO C57BL/6 mice. Cell

segmentation within GCs was performed using DAPI staining on the sections. The Pearson's

correlation coefficient indicates the co-expression relationship between MYC and H3K36me2.

1   The corresponding coefficient value (r) and a 95% confidence interval is displayed. **e**, Relative

2   H3K36me2 levels determined by calculating the ratio of H3K36me2 MI to MYC MI from **d**.

3   The top 5% of MYC intensity values, representing the highest MYC expressing cells, were

4   selected from each batch of two independent experiments. The bar represents mean. Student's *t*-

5   test. **p<0.01, ****p<0.0001, n.s., not significant.

Fig. 3. MiR-155 primarily controls KDM2A expression in high-affinity cMyc$^+$ LZ GC-B cells

a, Heat map showing the transcript levels of the enzymes involved in the demethylation of histone H3K36me2. The bulk RNA-seq dataset from WT and KO cMyc$^+$ LZ GC-B cells was used. FDR values are shown. **b**, Sequence alignment of a part of the 3'UTR of *Kdm2a*. The 8-mer miR-155 binding site is highlighted in a box. **c**, Immunofluorescence staining of splenic section from a WT mouse 7 days after SRBC immunisation, showing IgD (white), MYC (red) and KDM2A (blue). Scale bar, 50 µm. Representative section images from two experiments (WT n=4; KO n=4). **d**, Scatter plots show the MI of MYC and KDM2A within segmented GC cells in spleen sections from immunised WT and KO mice. The Pearson's correlation coefficient (r) is presented with a 95% confidence interval. **e**, Relative MFI of KDM2A in cells from four GC-B cell subpopulations. GC-B cells were derived from WT or KO donor B cells 7 days after HEL$^{3×}$-SRBC immunisation. Student's *t*-test. Pooled from two independent experiments (WT n=11; KO n=9). **f**, Relative MFI of KDM2A in cells from four GC-B cell subpopulations. GC-B cells were derived from WT or KO donor B cells 7 days after HEL$^{3×}$-SRBC immunisation.

1   Low- and high-affinity were determined based on $HEL^{3\times}$ binding within $Ig\kappa^+$ GC-B cells.

2   Student's *t*-test. Pooled from two independent experiments (WT n=11; KO n=9). Unless

3   otherwise stated, mean ± SEM is indicated. *P<0.05, **p<0.01, *** p<0.001. n.s., not

4   significant.

Fig. 4

Fig. 4. The regulatory activity of miR-155 is revealed under hypoxia

a, Representative flow cytometric histograms of hypoxia dye, MAR vs. relative cell number

(RCN) (left). Relative MFI of MAR (right) in GC-B and non-GC-B cells. GC-B and non-GC-B

cells were isolated from inguinal lymph nodes of C57BL/6 mice. One-way ANOVA. Pooled

from three independent experiments (n=6). **b**, KDM2A MFI of B cells from WT and KO

C57BL/6 mice cultured under normoxia (21% O_2) or hypoxia (1% O_2) for the indicated time

periods. Student's *t*-test. Pooled from three independent experiments (WT n=5; KO n=5). **c**,

Representative flow cytometric histograms of CTV vs. RCN at day 2 and 4 (top). CTV MFI of

B cells from WT and KO C57BL/6 mice cultured under normoxia or hypoxia for the indicated

time periods (bottom). Student's *t*-test. Pooled from five independent experiments (WT n=5; KO

n=5). **d**, Representative image of B cells from WT and KO C57BL/6 mice cultured under

normoxia or hypoxia for 2 days, followed by staining with MitoTracker Deep Red (Red) and

MitoTracker Orange CM-H$_2$ TMRos (green). DAPI (blue) was used as a nuclear counterstaining.

Scale bar, 10 µm. Representative images are shown. **e**, Quantification of relative MitoTracker

Orange CMTMRos levels. Each dot represents one cell. One-way ANOVA. Pooled from two

independent experiments (WT n=4; KO n=4). **f**, Relative ATP levels in B cells from WT and

KO C57BL/6 mice. The replated cells were cultured for 2 hours in hypoxia with indicated media

for ATP measurement. The luminescence values of each type of B cells cultured in media

containing glucose for 2 hours in hypoxia were used to calculate relative ATP levels. One-way

ANOVA. Pooled data from three experiments (WT n=3; KO n=3). Unless otherwise stated,

mean ± SEM is indicated. *P<0.05, **p<0.01, *** p<0.001, ****p<0.0001, n.s., not significant.

Fig. 5. MiR-155 optimizes nuclear mitochondrial gene expression through *Kdm2a-*

mediated control of H3K36me2 levels

a, Genomic distribution of KDM2A and H3K36me2 ChIP-seq peaks in B cells from WT and

KO mice cultured for 2 days in hypoxia is shown as pie charts. The average percentage of the

ChIP-seq triplicate samples is displayed. Promoter regions are defined as ± 2 kb of the

transcription start site (TSS). **b**, Metagene analysis and heatmaps of KDM2A or H3K36me2 for

CpG islands (CGIs) containing genes or genes without CGIs. Metagene analyses are displayed

in *± 10 kb* windows. A representative sample is shown. **c**, Plots showing the distribution of peak

distance to the nearest TSS. The average ChIP-seq read occupancy for KDM2A or H3K36me2

surrounding the TSS of CGIs containing genes. **d**, Analysis of genes with "WT-enriched" and

"KO-enriched" KDM2A peaks using the DAVID Functional Annotation Clustering Tool.

Pathways with a modified fisher exact p-value < 0.05 were included. **e**, A heat map showing

transcript levels of seven mitochondrial genes with "KO-enriched" KDM2A peaks at the

promoter/TSS regions or first introns near the TSS.

Fig. 6. The miR-155-*Kdm2a* interaction regulates mitochondrial ROS and apoptosis in GC-

B cells

a, Schematics representation of the 8-mer miR-155-binding site in the 3'UTR of the *Kdm2a*

(boxed) and the introduced base changes (red letters) to disrupt miR-155 binding in *Kdm2a*

mutant mice. **b**, Representative flow cytometric histograms showing KDM2A vs. RCN (top).

Relative MFI values of KDM2A in LZ and DZ GC-B cells are shown (bottom). GC-B cells

were derived from WT and MUT mice 7 days after SRBC immunisation. Student's *t*-test.

Pooled from three independent experiments (WT n=4; MUT n=4). **c**, Number of LZ and DZ

GC-B cells from WT and MUT mice within the µMT adoptive transfer system. Student's *t*-test.

Pooled from two independent experiments (WT n=7; MUT n=7). **d**, Representative flow

cytometric plots of MitoTracker DeepRed vs. MitoSOX in GC-B cells from WT and MUT mice

within the µMT adoptive transfer system (top). Percentage of MitoSOX$^+$ GC-B cells (bottom).

Student's *t*-test. Pooled from two independent experiments (WT n=7; MUT n=7). **e**,

Representative flow cytometric plots of B220 vs. active-Caspase 3 in GC-B cells from WT and

MUT mice within the µMT adoptive transfer system (top). Percentage of active-Caspase 3$^+$ GC-

B cells (bottom). Student's *t*-test. Pooled from two independent experiments (WT n=7; MUT

1   n=7). Unless otherwise stated, mean ± SEM is indicated. *P<0.05, **p<0.01, n.s., not

2   significant.

Methods

Mice

Mir155-deficient mice [57], SW$_{HEL}$ mice [25], *Myc*[gfp/gfp] mice [26], *Aicda*-cre-h*CD2* [27], µMT and

B6.SJL CD45.1 mice were maintained on C57BL/6J background. SW$_{HEL}$ mice were interbred

with *Myc*[gfp/gfp] and *Aicda*-cre-h*CD2* mice and were maintained either *Mir155*-sufficient or

Mir155-deficient background. B6.SJL CD45.1 mice were crossed with C57BL/6J mice , and

heterozygous mice were used. 6–15-week-old male and female mice were used. For the

generation of *Kdm2a* mutant mice, a single guide RNA (AAATTAAATGTAAGCATTAA) was

designed using CRISPOR [61] to cut in the 3'UTR of the *Kdm2a* gene (Position 3369 – 3376 of

Kdm2a   3'UTR,   ENST00000398645.2).   A   111bp   ssODN

(CTTATTTAGAGTCGATCTCCAATGTTGTGCTAAGATTTTTAAATTAAATGTAAGatgc

AtGGGAATGAGTCTTAGGCCACCAGCTGACACATTGGCATTATTTTGGGTCAG)   was

designed containing the "mir155-2" mutation. Zygotes were generated by superovulation of

C57BL/6J mice and electroporated with 50 µL of Cas9 protein / sgRNA / ssODN (final

concentrations 1.2 µM / 6 µM / 8 µM respectively). Both template and guide were added at the

1 cell stage. Electroporation of the embryos was performed using a Nepa21 electroporator

(Sonidel). Two males (1 heterozygous and 1 homozygous) were found to have sequence perfect

integration of the mir155-2 mutation via Sanger sequencing (Forward primer:

CGAGGACGTAGAACATGCGA, Reverse primer: GGGCAGGCACTTCCTAGTTT) and

were further bred to C57BL/6J yielding 2 heterozygous F1s. Q-PCR CNV analysis further

confirmed the presence of only one copy of the mir155-2 mutation. F1 mice were inbred to

generate homozygous mutant mice. All mice were bred and kept under specific-pathogen-free

conditions at the Francis Crick Institute Biological Research Facility in accordance with

guidelines set by UK Home Office and the Francis Crick Institute Ethical Review Panel.

Immunisation, treatments, and adoptive transfers

Mice were immunised intraperitoneally with 2×10^8 SRBCs. For HEL-specific GC responses,

splenic cells from donor mice were incubated with biotinylated antibodies: anti-CD43, anti-CD4

and anti-Ter-119, washed and incubated with anti-biotin microbeads (Miltenyi Biotec). B cells

were enriched by biotin negative selection using MACS LS Columns (Miltenyi Biotec). The

purity of recovered $CD19^+$ $B220^+$ B cells was >97%. 5×10^4 to 1×10^5 HEL-specific B cells

were transferred into B6.SJL $CD45.1^+$ hosts intravenously. Mice were immunised

intraperitoneally on the same day of cell transfer or one day later with 2×10^8 SRBCs

conjugated with $HEL^{3\times}$. The HyHEL10 BCR of SW_{HEL} mice exhibits a binding affinity 13,000-

fold lower to $HEL^{3\times}$ than to HEL. Immunisation with $HEL^{3\times}$ induces robust and persistent GC

responses at the expense of extrafollicular responses [62]. $HEL^{3\times}$ protein was harvested from the

supernatant of Chinese hamster ovary (CHO) cells expressing his-tagged $HEL^{3\times}$ [63] and was

purified using a His-Trap FF column (Cytiva) on AKTA Pure column chromatography system

(Cytiva). Eluted protein was concentrated to 0.5 ml (Vivaspin 15, 10,000 Da cutoff

concentrator) and applied to a S200 10/300 analytical size exclusion column (Cytiva). $HEL^{3\times}$

was found to eluted as a single, symmetrical peak and snap frozen at 1 mg/ml and stored at -

80 °C until required. For B cell transfer from wild type (*Kdm2a*-mutant littermates) or *Kdm2a*-

mutant mice, splenic B cells from donor mice were enriched by negative selection using MACS

LS Columns as described above. 2.5×10^7 to 3×10^7 B cells were transferred into μMT hosts

intravenously. Mice were immunised intraperitoneally on the same day of cell transfer with $2 \times$

10^8 SRBCs. To detect hypoxia, the hypoxia dye MAR was used. MAR is non-fluorescent

compound that is irreversibly converted into a fluorescent molecule by a reductase that is

activated in response to low oxygen concentrations. The conversion of MAR only occurs under

hypoxic conditions [64], and the fluorescence of MAR is not affected by increased oxygen

concentrations during tissue processing. Eight days after subcutaneous immunisation of

C57BL/6 mice with 50 µg of 4-Hydroxy-3-nitrophenyl acetyl hapten conjugated chicken

gamma globulin (NP-CGG) in Imject Alum Adjuvant (ThermoFisher Scientific), 4.3 µg MAR

hypoxia detecting probe (Goryo Chemical) in PBS per site or control DMSO in PBS was

subcutaneously injected into two sites around the immunised inguinal lymph node. Three hours

later, lymph nodes were harvested for flow cytometric analysis. Non-GC-B cells were gated as

$CD19^+ B220^+ CD38^+ CD138^- CD95^- GL7^-$.

Flow cytometry and cell sorting

Multicolour flow cytometry for analyses or for cell sorting was performed on LSR Fortessa or

FACSAria (BD Biosciences) and data were analysed using FlowJo v.10.8.1 software. PBS

supplemented with 2% FBS was used as flow cytometry buffer. Dead cells were excluded using

Zombie Fixable viability kit (BioLegend) during cell sorting. Harvested splenic B cells were

enriched by CD43, Ter119 and CD4 biotin negative selection using MACS LS Columns to

remove $CD43^+$ plasmablasts and non-B cells for analysis. For HEL-specific GC responses,

splenic B cells were enriched by CD45.1 negative selection using MACS LS Columns to

remove recipient-derived cells. Single-cell suspensions were prepared and stained for the

antibodies specific for cell surface markers. HEL-binding cells were stained with HEL (50

ng/ml), followed by HyHEL9 Alexa Fluor 647 [62]. For active-caspase 3 staining, after surface

marker staining, cells were fixed and permeabilized using a Fixation/Permeabilization Solution

Kit (BD Biosciences) and stained with anti-active caspase 3 antibody (BD Biosciences). For

KDM2A staining, cells were stained for surface marker antibodies, and were fixed and

permeabilized using a True-Nuclear Transcription Factor Buffer Set (BioLegend). The

permeabilized cells were stained with anti-KDM2A antibody, followed by donkey anti-rabbit

antibody staining. Cells stained with an anti-rabbit antibody, but no primary antibodies

determined the basal level of fluorescence intensity for KDM2A, and the basal fluorescence

intensity was subtracted from each MFI of KDM2A staining for the calculation of relative MFI.

For MitoSOX detection, cells were stained for surface marker antibodies. Then, they were

incubated in HBSS with Calcium and Magnesium (ThermoFisher scientific) supplemented with

2.5 µM MitoSOX Red (ThermoFisher Scientific) and 50 nM Mito Tracker Deep Red FM

(ThermoFisher Scientific) for 20 - 30 min and washed as described by the manufacturer before

analysis.

Antibodies

CD43, biotin (S7), BD Biosciences, 553269; CD4, biotin (GK1.5), BioLegend, 100404; Ter119,

biotin (TER-119), BioLegend, 116204; CD45.1, biotin (A20), ThermoFisher Sciences, 13-0453-

85; CD21/35, BV786 (7G6), BD Biosciences, 740894; CD23, FITC (B3B4), BD Biosciences,

553138; CD93, PE (AA4.1), ThermoFisher Sciences, 12-5892-82; CXCR4, PerCP-eFluor710

(2B11), ThermoFisher Sciences, 46-9991-82; CXCR4, APC (2B11), BD Biosciences, 558644;

CD86, FITC (GL1), BD Biosciences, 553691; CD86, PECy7 (GL1), BioLegend, 105014; CD86,

APCCy7 (GL1), BioLegend, 105030; CD86, BV605 (GL1), BioLegend, 105037; CD86, BV785

(GL1), BioLegend, 105043; CD95, PECy7 (Jo2), BD Biosciences, 557653; CD95, APC-R700

(Jo2), BD Biosciences, 565130; CD95, BV786 (Jo2), BD Biosciences, 740906; CD38, PerCP-

eFluor710 (90), ThermoFisher Sciences, 46-0381-82; CD38, PECy7 (90), BioLegend, 102718;

CD38, BV650 (90), BD Biosciences, 740489; GL7, Pacific blue (GL7), BioLegend, 144614;

GL7, Alexa Fluor 488 (GL7), ThermoFisher Sciences, 53-5902-82; human CD2, PECy7

(TS1/8), BioLegend, 309214; human CD2, APC (TS1/8), BioLegend, 309224; human CD2,

Alexa700 (TS1/8), BioLegend, 309228; CD45.2, BV605 (104), BD Biosciences, 563051;

CD45.2, BV786 (104), BD Biosciences, 563686; CD45.2, BUV395 (104), BD Biosciences,

564616; CD138, PE, (281-2), BioLegend, 142504; CD138, APC, (281-2), BioLegend, 142506;

CD138, APCCy7, (281-2), BioLegend, 142530; CD138, BV605, (281-2), BioLegend, 142531;

CD138, BV786, (281-2), BD Biosciences, 740880; B220, FITC (RA3-6B2), ThermoFisher

Sciences, 11-0452-82; B220, APCCy7 (RA3-6B2), ThermoFisher Sciences, 47-0452-82; B220,

BV421 (RA3-6B2), BioLegend, 103240; B220, BV650 (RA3-6B2), BioLegend, 103241; B220,

BUV787 (RA3-6B2), BD Biosciences, 612838; CD19, FITC (6D5), BioLegend, 115506; CD19,

BV605 (6D5), BioLegend, 115540; CD19, BV650 (6D5), BioLegend, 115541; IgM, BV605

(II/41), BD Biosciences, 743325; IgD, Alexa Fluor 647 (11-26c.2a), BioLegend, 405708; IgG1,

FITC (A85-1), BD Biosciences, 553443; IgG1, CF-594 (A85-1), BD Biosciences, 562559; Igk

light chain, BV421 (187.1), BD Biosciences, 562888; rabbit IgG, Alexa Fluor 647 (Poly4064),

BioLegend, 406414; rabbit IgG, PE (Poly4064), BioLegend, 406421; rabbit IgG, Alexa Fluor

488 (Poly4064), BioLegend, 406416; CD40, purified (HM-40), BioLegend, 102902; active

caspase 3, PE (C92-605), BD Biosciences, 550821; KDM2A, purified (EPR18602), Abcam,

ab191387; cMYC, Alexa Fluor 555 (9E10), ThermoFisher Sciences, MA1-980-A555; Histone

H3K36me2, purified (EPR16994(2)), Abcam, ab176921; mouse IgM F(ab')$_2$ fragment, Jackson

ImmunoResearch, 115-006-020; Streptavidin, PerCP-Cy5.5, BioLegend, 405214; Streptavidin,

PE, ThermoFisher Sciences, 12-4317-87; Streptavidin, BV650, BioLegend, 405231;

Streptavidin, BUV737, BD Biosciences, 612775.

B cell culture, proliferation assays and ATP measurement

Harvested splenic B cells from *Mir155*-sufficient or *Mir155*-deficient C57BL/6 mice were

enriched by CD43, Ter119 and CD4 biotin negative selection using MACS LS Columns.

Purified B cells were cultured with positive selection mimicking stimulants (1 µg/ml anti-CD40

antibody (HM40-3), 5 µg/ml AffiniPure F(ab')☐ Fragment Goat Anti-Mouse IgM (H+L)

(Jackson ImmunoResearch) and 4 ng/ml murine IL-4 (PeproTech) in B cell medium (RPMI

1640 GlutaMAX Supplement medium supplemented with 10 % fetal bovine serum, 1 % pen-

strep solution and 50 µM β-mercaptoethanol) at 37 °C in a humidified chamber containing 5%

CO_2 and either 21% O_2 (normoxia) or 1% O_2 (hypoxia). For CTV proliferation assays, purified

B cells were stained with 5µM CellTrace Violet (ThermoFisher Scientific) in PBS for 15 min at

RT and washed as described by the manufacturer. For ATP measurement, purified B cells were

cultured for 2 days as described above. On the measurement day, cells were washed twice in

PBS to remove glucose and resuspended in either glucose medium (DMEM no glucose medium

supplemented with 1 % pen-strep solution, 50 μM β-mercaptoethanol and 10 mM D-(+)-

glucose) or galactose medium (DMEM no glucose medium supplemented with 1 % pen-strep

solution, 50 μM β-mercaptoethanol and 10 mM galactose). After a thorough wash to remove

glucose in the medium, 1×10^5 cells were plated per well and were further cultured in the

presence or absence of 5 μM oligomycin A (Merck) at 37 °C in a humidified chamber

containing 5% CO_2 and 1% O_2 for 2 hours. All the cells were cultured in hypoxic conditions for

2 hours to directly compare ATP production ability under the same condition. ATP was

measured with Mitochondrial ToxGlo Assay kit (Promega) and followed the manufacturer's

protocol.

Immunofluorescence

Harvested spleens were embedded in OCT compound. 9 μm cryosections were fixed in ice cold

acetone for 3 min, followed by air dry for 20 min. The sections were blocked with blocking

buffer (0.3% Triton X-100, 1% BSA, 5% normal bovine serum and 1% rabbit serum in PBS)

and stained with the described antibodies for overnight at 4 C°. For MitoTracker staining of

cultured B cells, purified B cells were cultured for 2 days as described above and washed in PBS

twice. Cell suspension containing $2\text{-}3 \times 10^6$ cells/ml in PBS was placed on Shi-fix coverslips

(Shikhar Biotech) and incubated at 37 °C in a humidified chamber containing 5% CO_2 and 1%

O_2 for 30 - 40 min. The coverslips were then washed as instructed by the manufacturer. Cells

bound on the coverslips were stained with 83 nM MitoTracker Orange CM-H_2TMRos

(ThermoFisher Scientific) and 50 nM Mito Tracker Deep Red FM at 37 °C for 15 min, followed

by wash in warm PBS and fixation with 4% formaldehyde (ThermoFisher Scientific). Slides

were mounted with Prolong Gold Antifade Reagent with DAPI (ThermoFisher Scientific), and

images were acquired using LSM880 inverted confocal microscope (Zeiss) with 63× objective

and Zen Microscopy software (Zeiss).

Image analysis

For the detection of mitochondrial ROS on confocal images of in vitro cultured B cells, the

ratiometric cytoplasmic-to-nuclear intensity of mitochondrial ROS was quantified using a

custom-built CellProfiler (v.4.2.5) [65] pipeline. Uneven illumination was first corrected in the

Mito Tracker Deep Red FM and MitoTracker Orange CMTMRos channels by dividing the pixel

values by a polynomial fit to the intensity of the image pixels. Following smoothing with

median filtering, objects were then detected in the DAPI and Mito Tracker Deep Red FM

channels using minimum cross-entropy (Mito Tracker Deep Red FM) or Otsu (DAPI)

thresholding. All objects detected in the DAPI channel are then associated with corresponding

objects in the Mito Tracker Deep Red FM channel, i.e., any Mito Tracker Deep Red FM objects

without any corresponding DAPI objects are removed. The cytoplasm was then defined as the

area within Mito Tracker Deep Red FM objects not occupied by DAPI objects. The mean

intensity in the MitoTracker Orange CMTMRos channel was then measured within each

cytoplasm object, and this was divided by the background value, the mean intensity in the

MitoTracker Orange CMTMRos channel of each DAPI object to minimize batch effects. Cells

with incorrect segmentation and apoptotic cells in the images were manually removed. For the

analysis of co-expression of H3K36me2 / MYC and MYC / KDM2A in spleen sections was

quantified using a custom FIJI script [66]. First, the approximate centres of cells were detected by

identifying local extrema in eigenvalues of the Hessian tensor, calculated based on the IgD

channel using FeatureJ (https://imagescience.org/meijering/software/featurej/hessian/). Cells

were then fully segmented using these detected centres of cells as "markers" in a marker-

controlled watershed operation, implementing using MorphoLibJ [67], using a Gaussian- then

Sobel-filtered version of the IgD channel as input. The mean intensity of H3K36me2, MYC or KDM2A within the segmented cells was then quantified using ImageJ FIJI (v.2.9.0). All cells outside a manually defined region of interest were excluded.

Analysis of Histone PTMs

Cells were prepared for sorting by flow cytometry as described above. LZ or DZ GC-B cells (> 1×10^5) were directly sorted into 2 N H_2SO_4 as previously described [34]. After a 30 min incubation at room temperature, cellular debris was removed by centrifugation at $4,000 \times g$ for 5 min and histones were precipitated from the supernatant with trichloroacetic acid at a final concentration of 20% (v/v) overnight at 4 °C. Precipitated histones were pelleted with centrifugation at $10,000 \times g$ for 5 min at 4 °C, washed once with 0.1% HCl in acetone, then once with 100% acetone with centrifugation at $15,000 \times g$ for 5 min at 4 °C. Histones were briefly dried with SpeedVac concentrator (Eppendorf) and stored at -80 °C until derivatization. Histones were propionylated and digested according to Garcia *et al.* [35], with the modification of a single round of propionylation for 1 hr prior to and following digestion. Targeted LC-MS/MS was performed on a Thermo TSQ Quantiva (ThermoFisher Scientific). All injections were performed in technical triplicate. Raw data were analysed with Skyline (v.21.2) [68] according to published methods [34]. The relative abundance of each histone PTM was calculated from the total peak areas exported from Skyline. All the chemicals were purchased from ThermoFisher Sceintific.

RNA-seq and data analysis

Mir155-sufficient or *Mir155*-deficient cMyc-GFP[+] LZ GC-B cells were sorted by flow cytometry 5 days after HEL[3×]-SRBC immunisation. Total RNA was prepared with TRIzol Reagent (ThermoFisher Scientific) combined with RNeasy Mini Kit (QIAGEN) and processed

using a TruSeq Stranded mRNA Sample Prep kit (Illumina). RNA-sequencing was carried out

on the Illumina HiSeq 2500 platform and generated ~43 million, 75 bp paired-end reads per

sample. The RSEM package (v.1.2.31) [69] in conjunction with the bowtie2 alignment algorithm

(v.2.2.9) [70] was used for the mapping and subsequent gene-level counting of the sequenced reads

with respect to Ensembl mouse GRCm.38.86 version transcriptome. Normalization of raw count

data and differential expression analysis was performed with the DESeq2 package (v.1.24.1) [71]

within the R programming environment (v.3.6.0) [72]. Differentially expressed genes were defined

as those showing statistically significant differences between KO and WT sample groups (FDR

<0.05). Majority of *Mir155*-deficient donor mice were males, whereas majority of *Mir155*-

sufficient donor mice were female. Thus, the following 4 genes that are in the Y chromosome

and more enriched in KO than WT mice were removed from the list of DEGs

(ENSMUSG00000069045, ENSMUSG00000069049, ENSMUSG00000056673 and

ENSMUSG00000068457). Gene list ranked by the Wald statistic were used to look for pathway

and biological process enrichment using the Broad's GSEA software (v.2.1.0) with gene sets

from MSigDB (v6) [73]. GSEA results were visualized with EnrichmentMap (v3.3.4) [74] in

Cytoscape (v.3.7). We employed three miR target prediction tools, MiRDB [75], Diana [76] and

Target scan [77] for identifying miR-155 targets.

scRNA-seq and data analysis

Donor-specific *Mir155*-sufficient or *Mir155*-deficient GC-B cells were sorted 7 days after

HEL$^{3\times}$-SRBC immunisation. Libraries were prepared with Chromium Next GEM Single Cell 5'

Reagent Kit v2 (10× Genomics) and sequenced on the Illumina NovaSeq 6000 platform. Fastq

files were aligned and counted using CellRanger (v.6.1.2) and using as reference mm10-2020-A

(obtained from 10× Genomics). Matrices were imported into Seurat (v.4.0.5) [78], which was sued

for downstream analysis. After QC, cells with less than 1000 features and more than 10%

mitochondrial content were removed. Data was then normalized, scaled and variable features

were identified per cell. Samples were integrated using "CCA" by integrating in the following

order, first all replicates for each genotype and then by genotypes. DoubletFinder [79] was used to

predict doublets. Clusters identified as contaminating for expression of *Cd8a, Cd3e, Cd3d, Lyz2*

and *Gzma,* and cells labelled as doublets were removed from the analysis. Clusters were

manually annotated using differential expressed genes obtained from FindMarkers. Uniform

Manifold Approximation and Projection (UMAP) was used for visualization. VlnPlots and

FeaturePlots have been used to display gene expression values per cluster or overlay on UMAP.

The merged dataset of biological triplicates contained 25,358 and 24,087 GC-B cells for WT

and KO cells, respectively. Unsupervised clustering using Seurat identified nine distinct clusters,

each with their unique cell cycle status determined by the expression of genes associated with

different phases of the cell cycle (**Extended Data Fig. 1e,f**). For gene signatures, a score was

calculated using AddModuleScore prior to plotting. Statistical significance was obtained using

Wilcoxon test and multiple comparison correction was performed using the "fdr" procedure.

Differential gene expression between genotypes per cluster was carried out using a pseudobulk

approach, in which gene counts were aggregated per sample and the "pseudoBulkDGE"

function from the scran package [80] was used with the design ~genotype. Genes are considered

differentially expressed if FDR < 0.05. For trajectory analysis, Partition-based graph abstraction

PAGA [30] was used. The integrated Seurat object was converted into a h5 format, neighbours

were computed using 20 pcs and clusters detected using Seurat were used to compute the PAGA

graph. Mitochondrial gene list was downloaded from MitoCarta3.0 [53].

ChIP-seq and data analysis

WT and KO B cells were cultured under hypoxic conditions with positive selection mimicking

stimulants for 2 days. Cells were fixed for 1 hr in 2 mM EGS, followed by 15 min in 1%

Formaldehyde. Chromatin was sheared using Bioruptor Pico sonication device (Diagenode). An

aliquot of the chromatin was used to assess the size of the DNA fragments obtained by High

Sensitivity NGS Fragment Analysis Kit (Agilent) on Fragment Analyzer (Agilent). ChIP was

performed using IP-Star Compact Automated System (Diagenode). Chromatin corresponding to

13.5 µg and 1 µg were immunoprecipitated with KDM2A antibody (a gift from R. Klose,

University of Oxford) and H3K36me2 (Abcam). Chromatin corresponding to 1% was used as

Input. Libraries were prepared using MicroPlex Library Preparation Kit v3 (Diagenode) with 24

UDI for MicroPlex v3 (Diagenode). Generated libraries were purified using Agencourt AMPure

XP (Beckman Coulter) and quantified using Qubit 1× dsDNA HS Assay Kit (ThermoFisher

Scientific). The fragment size was analysed by High Sensitivity NGS Fragment Analysis Kit on

Fragment Analyzer. Sequencing was carried out on the Illumina NovaSeq 6000 platform. Read

alignment, library QC, peak detection and quantification was carried out using the nf-

core/chipseq pipeline (v.1.2.2), using "-profile crick --min_reps_consensus 2". Briefly, reads

were aligned using BWA [81] to GRCm38 using the genomes provided with the pipeline. Peaks

were called using MACS2 [82] and annotated using HOMER [83]. A list of consensus peaks is

generated as the union of peaks detected in at least two replicates per condition. Bigwigs

generated by the pipeline (scaled to 1 million mapped reads) were used to generate meta profile

plots with Deeptools "compute matrix" [84]. A table containing CpG island containing genes was

obtained from UCSC Table Browser [85]. Initially, an average of 26436 and 17586 KDM2A peaks

were identified for WT and KO cells, respectively. The peaks that only one of the triplicate

samples bound were removed from the subsequent analysis. After the exclusion of peaks, there

were 9129 KDM2A peaks that were bound by both WT and KO B cells. The average fold

change (fc) was calculated by dividing the average KDM2A peak signal by the input signal, and

then "WT-enriched" or "KO-enriched" peaks among the 9129 KDM2A peaks bound by both

WT and KO B cells were determined based on a fold change threshold (fc >3) and a ratio of

average fc between WT and KO cells ($>2^{0.23}$-fold enrichment). Peaks with an fc >3 from "peaks

bound only by WT" (5905) and "peaks bound only by KO" (1479) were also categorized as

"WT-enriched" and "KO-enriched" KDM2A peaks, respectively. After applying the selection

threshold, a total of 1403 genes with 1453 "WT-enriched KDM2A" peaks and 309 genes with

309 "KO-enriched KDM2A" peaks were identified (**Supplementary Table 3**). Among these,

73.6% and 78.0% of the "WT-enriched KDM2A" peaks and "KO-enriched KDM2A" peaks,

respectively are genes containing CpG island.

Statistical analysis

Statistical significance was determined by unpaired, two-tailed Student's t-test, Kruskal-Wallis

one-way analysis of variance (ANOVA) with multiple comparisons, or Wilcoxon rank sum test.

The statistical method used is indicated in the respective Fig. legends. Prism (v.9.4.1) software

(GraphPad) and R were used to calculate all statistical analyses.

Data and code availability

The scRNA-seq, bulk RNA-seq and ChIP-seq data have been deposited in the NCBU Gene

Expression Omnibus (GEO) database under accession number GSE230528, GSE231649,

GSE230527, respectively. Source data are provided with this paper.

Methods references

61 Concordet, J. P. & Haeussler, M. CRISPOR: intuitive guide selection for CRISPR/Cas9

genome editing experiments and screens. *Nucleic Acids Res* **46**, W242-W245 (2018).

https://doi.org:10.1093/nar/gky354

62 Paus, D. *et al.* Antigen recognition strength regulates the choice between extrafollicular

plasma cell and germinal center B cell differentiation. *J Exp Med* **203**, 1081-1091 (2006).

https://doi.org:jem.20060087 [pii]10.1084/jem.20060087

63 Long, Z., Phillips, B., Radtke, D., Meyer-Hermann, M. & Bannard, O. Competition for

refueling rather than cyclic reentry initiation evident in germinal centers. *Sci Immunol* **7**,

eabm0775 (2022). https://doi.org:10.1126/sciimmunol.abm0775

64 Piao, W. *et al.* Development of azo-based fluorescent probes to detect different levels of

hypoxia. *Angew Chem Int Ed Engl* **52**, 13028-13032 (2013).

https://doi.org:10.1002/anie.201305784

65 Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*

Bioinformatics **22**, 433 (2021). https://doi.org:10.1186/s12859-021-04344-9

66 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*

Methods **9**, 676-682 (2012). https://doi.org:10.1038/nmeth.2019

67 Legland, D., Arganda-Carreras, I. & Andrey, P. MorphoLibJ: integrated library and

plugins for mathematical morphology with ImageJ. *Bioinformatics* **32**, 3532-3534

(2016). https://doi.org:10.1093/bioinformatics/btw413

68 MacLean, B. *et al.* Skyline: an open source document editor for creating and analyzing

targeted proteomics experiments. *Bioinformatics* **26**, 966-968 (2010).

https://doi.org:10.1093/bioinformatics/btq054

Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011). https://doi.org:10.1186/1471-2105-12-323

Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012). https://doi.org:10.1038/nmeth.1923

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-0550-8

R Core Team. *R: A language and environment for statistical computing.*, https://www.R-project.org/. (2021).

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005). https://doi.org:10.1073/pnas.0506580102

Merico, D., Isserlin, R., Stueker, O., Emili, A. & Bader, G. D. Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. *PLoS One* **5**, e13984 (2010). https://doi.org:10.1371/journal.pone.0013984

Chen, Y. & Wang, X. miRDB: an online database for prediction of functional microRNA targets. *Nucleic Acids Res* **48**, D127-D131 (2020). https://doi.org:10.1093/nar/gkz757

Vlachos, I. S. *et al.* DIANA-TarBase v7.0: indexing more than half a million experimentally supported miRNA:mRNA interactions. *Nucleic Acids Res* **43**, D153-159 (2015). https://doi.org:10.1093/nar/gku1215

Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA target sites in mammalian mRNAs. *Elife* **4** (2015). https://doi.org:10.7554/eLife.05005

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015). https://doi.org:10.1038/nbt.3192

McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337 e324 (2019). https://doi.org:10.1016/j.cels.2019.03.003

Lun, A. T., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122 (2016). https://doi.org:10.12688/f1000research.9501.2

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009). https://doi.org:10.1093/bioinformatics/btp324

Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008). https://doi.org:10.1186/gb-2008-9-9-r137

Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010). https://doi.org:10.1016/j.molcel.2010.05.004

Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016). https://doi.org:10.1093/nar/gkw257

Karolchik, D. *et al.* The UCSC Table Browser data retrieval tool. *Nucleic Acids Res* **32**, D493-496 (2004). https://doi.org:10.1093/nar/gkh103

Acknowledgements

This work was supported by the Francis Crick Institute which receives core funding from Cancer Research UK Grants (grant CC2078), the UK Medical Research Council Grants (grant CC2078), and Wellcome Trust Grants (grant CC2078) to D.P.C., BBSRC (BBS/E/B/000C0407) to M.T., and the National Institute of General Medical Sciences of the National Institutes of Health (P41 GM108569) to N.L.K. We thank M. Busslinger (Research Institute of Molecular Pathology) for *Aicda-cre*-h*CD2* mice; B. P. Sleckman (University of Alabama at Birmingham) for *Myc*$^{gfp/gfp}$ mice; R. Brink (Garvan Institute of Medical Research/University of New South Wales) for SW$_{HEL}$ mice and HyHEL9 antibody; O. Bannard (University of Oxford) for providing CHO-his-tagged HEL$^{3\times}$ cells; the staff of Genomics Facility of the Babraham Institute for technical support; the staff of the Francis Crick Institute Animal Facility, the Flow Cytometry Facility, the Advanced Sequencing Facility, the Advanced Light Microscopy Facility, and the Experimental Histopathology Facility for technical support; and E. Vigorito and C. Vinuesa for critical review.

Author contributions

R.N. conceived and designed the study. M.L. and P.C performed bioinformatic analysis. S.V-B. and G.D. generated *Kdm2a* mutant mice. R.N., J.M.C. and R.G. performed experiments and analysed data. D.B. generated computational image analysis pipelines. N.P.B. and R.J.K. provided resources. N.L.K. provided supervision. M.T. interpreted data and critically revised the manuscript. R.N. and D.P.C. wrote the manuscript. All authors reviewed and edited the written manuscript.

Competing interest

The authors declare no direct competing financial or nonfinancial interests.

1   **Additional information**

2   Supplementary Information is available for this paper.

3   Correspondence and requests for materials should be addressed to Rinako Nakagawa

4   rinako.nakagawa@crick.ac.uk or Dinis Calado dinis.calado@crick.ac.uk

Extended data figures and Extended data figure legends

Extended Data Fig. 1 Gene expression patterns of miR-155-sufficient and miR-155-

difficient GC-B cell clusters

a, Flow cytometry gating strategy for isolating splenic donor GC-B cells 7 days after HEL$^{3\times}$-

SRBC immunisation. **b**, Schematics illustrating the relationship between four GC-B cell

subpopulations (cMyc^{-} LZ, cMyc^{+} LZ, cMyc^{+} DZ and cMyc^{-} DZ) and the expression patterns of

cMyc and miR-155. **c**, MFI values of cMyc-GFP in GC-B cells from WT and KO, 8 days after

HEL$^{3\times}$-SRBC immunisation. Each dot represents one mouse. Student's *t*-test. Pooled from two

experiments (WT n=11; KO n=11). **d**, Quantification of cMyc$^+$ LZ and DZ GC-B cell numbers

in WT and KO mice, 5 days after HEL$^{3×}$-SRBC immunisation. Student's *t*-test. Pooled from

four experiments (WT n=13; KO n=14). **e**, UMAP plots displaying merged scRNA-seq profiles

from WT and KO GC-B cells, with 9 clusters (left) and predicted cell cycle phases (right). **f**,

UMAP plots showing scRNA-seq profiles from WT or KO GC-B cells, with 9 identified

clusters. **g**, UMAP plots coloured by the normalized expression of signature genes associated

with LZ, DZ and Myc signature genes. **h**, Ball-and-stick representation of PAGA connectivity

among scRNA-seq GC-B cell clusters from WT or KO GC-B cells. **i**, UMAP plots coloured by

the number of genes detected in each cell or the percentage of mitochondrial genes. **j**, Violin

plots illustrating the expression levels of *Rps21* or *Rpl14* in the GC-B cell clusters. **k**, Top 8

GSEA gene signatures in WT vs. KO comparison using RNA-seq dataset of cMyc$^+$ LZ GC-B

cells. WT-enriched gene signatures are shown. FDR; false discovery rate, NEM; normalized

enrichment score. Unless otherwise stated, mean ± SEM is indicated. ****p<0.0001, n.s., not

significant.

Extended Data Fig. 2

Extended Data Fig. 2. Histone PTM states in *Mir155*-sufficient and *Mir155*-deficient GC-B cells

a, The relative abundance of histone PTMs in LZ GC-B cells from WT and KO mice. Donor-

derived GC-B cells were sorted 5 days after adoptive transfer for the analysis. **b**, The relative

abundance of histone PTMs, including histone H3.1 K36, H3.3 K36, H3.1 K27 and H3.3 K27,

in DZ GC-B cells from WT and KO mice. Donor-derived GC-B cells were sorted 5 days after

adoptive transfer for the analysis. Triplicate samples were used for both WT and KO

backgrounds. **c,d**, Immunofluorescence staining of splenic sections from WT mice 7 days after

SRBC immunisation, showing IgD (white), MYC (red) and H3K36me1 (**c**) or H3K36me3 (**d**)

(green). Scale bar, 20 µm. Scatter plots illustrating the expression of MYC and H3K36me1(**c**) or

H3K36me3 (**d**) in segmented GC cells within spleen sections from immunised WT and KO

mice. Pearson's correlation coefficient (r) indicates the co-expression between MYC and

H3K36me1 (**c**) or H3K36me3 (**d**). The corresponding coefficient value (r) and a 95%

confidence interval is displayed. Representative section images from two experiments (WT n=4;

KO n=4). Unless otherwise stated, mean ± SEM is indicated.

Extended Data Fig. 3

Extended Data Fig. 3. The histone demethylase *Kdm2a* is a target of miR-155

a, Heat map showing the transcript levels of the miR-155 targets. The miR-155 targets displaying a significant (FDR <0.05) and at least 1.3-fold increase in KO cMyc$^+$ LZ GC-B cells compared to WT are shown. The bulk RNA-seq dataset of WT and KO cMyc$^+$ LZ GC-B cells was used. **b**, Representative flow cytometric histograms showing KDM2A vs. relative cell number (RCN).

Extended Data Fig. 4

Extended Data Fig. 4. Induction of miR-155 by positive selection signals in normoxia and

hypoxia

a, MiR-155 expression levels in B cells cultured under normoxia and hypoxia. Each dot

represents one mouse sample. Student's *t*-test. Pooled data from two experiments (WT n=5; KO

n=5). **b**, Relative luminescence units (RLU) of B cells cultured in glucose containing media for

2 hours in hypoxia. One-way ANOVA. Pooled from three experiments (WT n=3; KO n=3).

Unless otherwise stated, mean ± SEM is indicated. *p<0.05, **p<0.01, ****p<0.0001, n.s., not

significant.

Extended Data Fig. 5

1
2 **Extended Data Fig. 5. The genes with KO-enriched KDM2A peaks showed a decrease in**

3 **H3K36me2 levels**

4 **a**, IGV snapshots showing KDM2A and H3K36me2 ChIP-seq in hypoxic cultured B cells from

5 WT and KO mice for representative nuclear mitochondrial genes with "KO-enriched KDM2A"

6 peaks. The interval scales for the y-axis are indicated at the top of each gene track.

Extended Data Fig. 6. Increased KDM2A expression exacerbates mitochondrial ROS in

the absence of miR-155

a, Total B lineage cell numbers in the spleen of WT and MUT mice. T1: CD19$^+$ B220$^+$ AA4.1$^+$

CD23$^-$ IgM$^+$; T2: CD19$^+$ B220$^+$ AA4.1$^+$ CD23$^+$ IgM$^+$; T3: CD19$^+$ B220$^+$ AA4.1$^+$ CD23$^+$ IgM^{low};

Follicular (Fo) B: CD19$^+$ B220$^+$ AA4.1$^-$ CD23$^+$ IgM^{low}; Marginal zone (MZ) B: CD19$^+$ B220$^+$

AA4.1$^-$ CD23^{low} IgM$^+$. Student's *t*-test. Pooled from two experiments (WT n=4; MUT n=4). **b**,

Representative flow cytometric plots showing the analysis of μMT mice transferred with or

without donor B cells from a WT mouse 6 days after SRBC immunisation. Recipient μMT mice

without donor B cells failed to generate GC-B cells. **c**, Representative flow cytometric plots

analysing μMT mice transferred with WT or MUT B cells 7 days after SRBC immunisation. **d**,

Activated SW$_{HEL}$ B cells were transduced with *Kdm2a* or control (empty) lentivirus. MFI of

KDM2A was analysed in activated B cells cultured for two days following transduction.

1   Representative of two independent experiments. Either control- or *Kdm2a*-transduced HEL-

2   activated donor SW$_{HEL}$ B cells were transferred into congenic mice. GC-B cells derived from

3   transduced donor cells were stained for MitoSOX four days after HEL$^{3\times}$-SRBC immunisation

4   (right). Student's *t*-test. Pooled from two experiments (Control n=10; *Kdm2a* n=10). Unless

5   otherwise stated, mean ± SEM is indicated. **p<0.01, n.s., not significant.

Extended Data Fig.7

Extended Data Fig. 7. Working model: miR-155-mediated epigenetic regulation control

mitochondrial fitness during GC-B cell positive selection

Following the reception of signals through BCR engagement and T cell help, LZ GC-B cells

undergo a positive selection process that induces cell division. Positively selected GC-B cells,

particularly high-affinity clones, migrate to the less hypoxic DZ and sustain their vigorous cell

division by relying on OXPHOS for energy production. To promote the dynamic transition of

GC-B cells from the hypoxic LZ to the DZ under hypoxia-induced stress, miR-155 plays a

crucial role by repressing the hypoxia-induced expression of KDM2A. This repression

mechanism enables precise regulation of H3K36me2 levels, leading to optimal expression of

nuclear mitochondrial genes. As a result, mitochondrial remodeling is facilitated, and OXPHOS

is enhanced in hypoxic conditions. Thus, the miR-155-*Kdm2a* interaction is indispensable for

1   promoting mitochondrial fitness during positively selection, which provides robust

2   mitochondrial in high-affinity clones to endure the high-rate of cell division in the DZ. This

3   mechanism allows high-affinity clones to undergo clonal expansion, thus ensuring the process of

4   affinity maturation.

Supplementary information

Supplementary Table 1. Identified cluster marker genes in scRNA-seq from total GC-B cells

of WT and KO.

Supplementary Table 2. Differentially expressed genes in bulk RNA-seq of WT and KO

cMyc$^+$ LZ GC-B cells.

Supplementary Table 3. Genes with "WT-enriched" and "KO-enriched" KDM2A peaks in

activated B cells under hypoxia.