Detection of prions from spiked and free-ranging carnivore feces

H. N. Inzalaco[a], E. E. Brandell[a], S.P. Wilson[b], M. Hunsaker[a], D. R. Stahler[c], K. Woelfel[d], D.

P. Walsh[e], T. Nordeen[b], D. J. Storm[f], S. S. Lichtenberg[g], W. C. Turner[h]

[a]Wisconsin Cooperative Wildlife Research Unit, Department of Forest and Wildlife Ecology,

University of Wisconsin, Madison, Madison, WI 53706, USA

[b]Nebraska Game and Parks Commission 2200 N 33rd St. P.O. Box 30370, Lincoln, NE 68503

[c]Yellowstone Center for Resources, Yellowstone National Park, WY 82190

[d]Wild and Free Wildlife Rehabilitation Program, 27264 MN-18 Garrison, MN 56450, USA

[e]U.S. Geological Survey, Montana Cooperative Wildlife Research Unit, University of Montana,

Missoula, MT, USA

[f]Wisconsin Department of Natural Resources, Eau Claire, WI 54701 USA

[g]Department of Veterinary and Biomedical Sciences, University of Minnesota, St. Paul, MN

55108, USA

[h]U.S. Geological Survey, Wisconsin Cooperative Wildlife Research Unit, Department of Forest

and Wildlife Ecology, University of Wisconsin – Madison, Madison, WI 53706, USA.

*Correspondence: inzalaco@wisc.edu

ABSTRACT

Chronic wasting disease (CWD) is a highly contagious, fatal neurodegenerative disease caused by infectious prions (PrP^{CWD}) affecting wild and captive cervids. Although experimental feeding studies have demonstrated prions in feces of crows (*Corvus brachyrhynchos*), coyotes (*Canis latrans*), and cougars (*Puma concolor*), the role of scavengers and predators in CWD epidemiology remains poorly understood. Here we applied the real-time quaking-induced conversion (RT-QuIC) assay to detect PrP^{CWD} in feces from cervid consumers, to advance surveillance approaches, which could be used to improve disease research and adaptive management of CWD. We assessed recovery and detection of PrP^{CWD} by experimental spiking of PrP^{CWD} into carnivore feces from 9 species sourced from CWD-free populations or captive facilities. We then applied this technique to detect PrP^{CWD} from feces of predators and scavengers in free-ranging populations. Our results demonstrate that spiked PrP^{CWD} is detectable from feces of free-ranging mammalian and avian carnivores using RT-QuIC. Results show that PrP^{CWD} acquired in natural settings is detectable in feces from free-ranging carnivores, and that PrP^{CWD} rates of detection in carnivore feces reflect relative prevalence estimates observed in the corresponding cervid populations. This study adapts an important diagnostic tool for CWD, allowing investigation of the epidemiology of CWD at the community-level.

INTRODUCTION

The breakdown of detritus is important for regulating the movement of energy through food webs[1], structuring ecological communities and ecosystem function[2], and can also influence the dynamics of disease systems[3, 4]. Obligate and/or facultative scavenger species participate in scavenging, or the consumption of carrion, as a resource acquisition strategy. The latter are often also predators, consuming live prey. Scavenging and predation processes have important influences on disease epidemiology through removal of infectious individuals or materials that could transmit pathogens to new hosts[5, 6]. Simultaneously, scavengers and predators may also spread infectious diseases by transporting the pathogen to other areas (through consumption, subsequent fecal deposition, or translocation of contaminated material)[7]. Determining whether predators and scavengers reduce or enhance disease transmission risk in the host species they consume is important for understanding the ecological context of infectious diseases in wildlife. The role of these consumers in disease dynamics will be affected by factors such as carnivore diversity, gut capacity, digestive rates and fecal deposition patterns, pathogen environmental stability, host-pathogen encounters, and infectious dose.

Chronic wasting disease (CWD) is a highly contagious fatal neurological disease caused by a misfolded prion protein (PrP^{CWD}) that affects several domestic and free-ranging cervid species. Currently, cervids are the only wildlife species monitored in routine CWD surveillance and the effects of other species and the environment on CWD dynamics are relatively unknown. For prion diseases, pathogen detection from abiotic and biotic environmental sources is technically challenging due to prions being devoid of genetic material. For years detection challenges limited the ability to address the sort of ecological questions readily possible for many other

infectious agents that contain nucleic acids, such as many bacterial pathogens with complex

disease ecology (e.g. *Bacillus anthracis[8]*, *Yersinia pestis[9]*, and *Francisella tularensis[10]*). Prion

detection assays need to overcome the sensitivity limitations of existing techniques to be

adaptable for detection of prions from a variety of ecological- and management-relevant

environmental samples. As such, an assay with this capacity could be used not only to begin

understanding how other species influence CWD processes but may also help to improve current

surveillance efforts.

Although the first observations of CWD were made in captive mule deer (*Odocoileus hemionus*)

and black-tailed deer (*Odocoileus hemionus columbianus*) in Colorado[11], the disease has since

spread broadly in cervid species across North America and Europe, including white-tailed deer

(*Odocoileus virginianus*, hereafter referred to as WTD), elk (*Cervus canadensis*), reindeer

(*Rangifer tarandus*), and moose (*Alces alces*)[12-15]. Through the disease course, PrP^{CWD}

accumulates in body fluids and tissues[16, 17] of presymptomatic and symptomatic CWD-infected

individuals, is shed in excreta and secreta[18-21], and ultimately results in neuroinvasion causing

clinical disease and death[22, 23]. Transmission of CWD can occur either directly (i.e., contact

between infectious and naive individuals) or indirectly through contact with contaminated

environments[24]. Given that PrP^{CWD} can remain stable and infectious for years in the

environment[25], deposition and accumulation of PrP^{CWD} in the environment from infected

carcasses or via shedding from infected individuals may lead to increased environmental disease

exposure risk[26]. Consumption of CWD-infected hosts, through predation or scavenging suggests

a role for carnivores in CWD epidemiology.

Coyotes (*Canis latrans*)[27], American crows (*Corvus brachyrhynchos*)[28], and cougars (*Puma*

concolor)[29] shed prions following ingestion of prion-infected material. Prion gut residency times

in carnivores appear to be fairly short, lasting $\leq$ 3 days in coyotes and cougars, to just several

hours in crows. Carnivores consuming contaminated cervid carrion may contribute to PrP^{CWD}

translocation and contamination in the environment[27, 28]. However, infectious PrP^{CWD} shed in

coyote feces was reduced relative to ingested material, implying feces-associated PrP^{CWD}

deposits may contain less infectious material compared to unconsumed carrion[27]. Baune *et al.*[29]

suggest that most of the prions ingested by cougars are eliminated or sequestered, supporting the

notion that predators may have a dilution effect on the CWD system. Further, a modeling study

from Brandell *et al.*[5] suggests that cougar and gray wolf (*Canis lupus*) predation pressure may

independently decrease CWD outbreak size and delay prevalence increases of deer and elk, and

this cleansing effect is amplified when predator selection for infected adults is greater than

uninfected juveniles. Similarly, Fisher *et al.*[30] recently reported predation by cougars may have

slowed the increase in prevalence in an area of high CWD prevalence.

Previous studies investigating prions in scavengers relied on detection methods that range from

mouse bioassays and antibody-based tools[28], to an ultra-sensitive *in vitro* protein amplification

assay, protein misfolding cyclic amplification (PMCA)[27]. Although mouse bioassay remains the

gold standard for definitively assessing the presence and titer of PrP^{CWD}, these assays can take

months to years to complete. Bioassays are also often used in conjunction with PMCA, which

offers the same level of sensitivity as mouse bioassay and is far more sensitive than antibody-

based detection tools[31, 32]. For example, in the study by Nichols *et al.*[27], PMCA was used to

characterize the fecal samples prior to use as inoculum in the mouse bioassay. However, PMCA

is costly and not adaptable for high-throughput diagnostics as it still requires the need for prion

disease susceptible brain tissue as a conversion substrate and can take up to two weeks to

produce a result. The development and application of an alternative ultrasensitive *in vitro* protein

amplification assay, the real-time quaking-induced conversion (RT-QuIC), has advanced high

throughput detection of PrP[CWD] from host tissues[33, 34] and various host secreta and excreta[19, 34-38]

without the use of animal tissue substrates. Currently, there has been only one report of the

application of RT-QuIC for PrP[CWD] detection from feces of cervid consumers (cougar), based on

captive animals in a controlled laboratory environment[29].

Here we evaluated the utility of RT-QuIC for detection of PrP[CWD] using fecal-spiking assays

from a suite of relevant predator and scavenger species. We assessed limits of PrP[CWD] recovery

and detection sensitivity from nine species relative to RT-QuIC characterized CWD-positive

brain tissue. We then used this assay to detect PrP[CWD] in feces samples from two free-ranging

carnivore species – coyotes and cougars. Our findings from field collected feces suggests that

this approach could be used to institute early surveillance of CWD, especially in locations or

jurisdictions neighboring CWD endemic zones that are considered areas of concern for

geographic spread or areas where hunter-harvest rates are low, and also to begin to unravel how

other species influence CWD dynamics and geographic spread.

MATERIALS AND METHODS

Fecal sample spiking assays. Individual fecal samples were handled separately, using fresh

disposable, single use nitrile gloves, and disposable weigh boats to prevent cross-contamination

of samples. Using CWD-negative fecal samples as determined by RT-QuIC described below

data analysis, spiking experiments were used to demonstrate recovery of PrP^{CWD} from feces of 9

different scavenger and predator species: gray wolf (hereafter wolf), cougar, coyote, American

crow (hereafter crow), American black bear (*Ursus americanus*, hereafter bear), raccoon

(*Procyon lotor*), common raven (*Corvus corax*, hereafter raven), red fox (*Vulpes vulpes*,

hereafter fox), and bald eagle (*Haliaeetus leucocephalus*, hereafter eagle; sources listed above

and in Table 1). Spiking assays were carried out by using 50 mg of each fecal sample combined

with 50 μL of a 10-fold dilution of a 10% (w/v) CWD-positive WTD brain homogenate (BH)

from the obex region of tagged WTD #5219 in the late stages of CWD (hereafter referred to as

the reference sample), provided by the Wisconsin Department of Natural Resources (WDNR).

The reference sample was prepared in 1X phosphate-buffered saline (PBS) to achieve a range of

final concentrations of 10^{-3} to 10^{-8} mg/mL of CWD-positive brain. CWD-positive and negative

BH were determined by RT-QuIC. For the negative control spike, 50 μL of a 10^{-3} mg/mL

dilution of a 10% (w/v) CWD-negative WTD BH was used. Each spike dilution was added to

feces and allowed to soak in for 24 hours, then placed in a 50 °C incubator without agitation for

16 h, to keep assay temperature conditions consistent with those used for RT-QuIC. Spiked fecal

samples were then prepared as described below.

Fecal sample extract preparation for RT-QuIC assay. Fecal samples (Table 1) were stored at

-80°C prior to preparation for use in the RT-QuIC assay. Individual fecal samples and

subsamples were handled separately, using fresh disposable/single use nitrile gloves, and

disposable weigh boats to prevent cross-contamination of samples. To prepare field collected

fecal samples for detection of PrP^{CWD}, each individual sample had one subsample collected from

three different areas/regions consisting of a slice from the middle and both ends of the sample,

totaling three biological replicates (50 mg each). A separate scalpel blade was used to cut each

section, that included surface material as well as inner material up to a depth of 1 cm to ensure

sufficient sampling of surface and inner portions. These 3 subsamples were then extracted

individually, for a total of 3 separate extractions, using 1 mL of sterile 1X PBS, then processed in

a thermomixer (1,400 rpm, 30 min, 25°C; Eppendorf ThermoMixer F1.5) followed by

centrifugation at 16,000 x g for 15 min. Supernatants (~750 µL) were collected, followed by

addition of 750 µL 1X PBS to each feces pellet and remaining buffer. Samples were vortexed,

thermomixed (1,400 rpm, 30 min, 25°C), and centrifuged again at 16,000 x g for 15 min, after

which an additional ~750 µL supernatant was removed and added to the first supernatant.

Resultant supernatants were then centrifuged again for further clarification at 16,000 x g for 15

min, followed by collection of 1 mL of clarified supernatant into a new sterile tube and 500 µL

of 23.1 mM sodium phosphotungstate hydrate (Na-PTA; Sigma-Aldrich, Cat. # 496626) was

added. Samples were incubated without agitation for 16 hours at 4°C, then centrifuged (4 °C, 30

min, 5000 x g). Sample supernatants were carefully removed from each tube and discarded.

Pellets were retained and washed with a 1:1 solution of 18 MΩ H_2O and 23.1 mM Na-PTA

followed by centrifugation (4 °C, 30 min, 5000 x g) and aspiration of the wash solution. Pellets

were resuspended in 30 µL of RT-QuIC sample buffer (0.1 g/mL sodium dodecyl sulfate in

phosphate-buffered saline with N-2 cell culture supplement; ThermoFisher, Waltham, MA) and

reconstituted using a Qsonica Q700 cup horn ultrasonicator (Amplitude 36 for 1 min.). A volume

of 2 µL of each reconstituted sample was used to seed each reaction well for 8 technical

replicates.

RT-QuIC Assay. The RT-QuIC *in vitro* prion amplification assay was performed as described

by Orru *et al.*[39], with sodium iodide as described by Metrick *et al.*[40] with minor modifications.

Briefly, 2 µL of sample extracts were added to a given well of a 96-well format optical-bottom

black microplate (Thermo Scientific, Fair Lawn, NJ, USA), each already containing 98 µL of

RT-QuIC reaction mixture (0.1 mg·mL^{-1} 90-231 recombinant hamster prion protein, produced as

previously described:[39], 300 mM sodium iodide, 20 mM sodium phosphate, 1.0 mM

ethylenediaminetetraacetic acid, and 10 µM thioflavin T). Microplate-compatible

spectrophotometers capable of heating, shaking, and fluorescence monitoring (BMG FLUOstar,

Cary, North Carolina) were used with the following instrument settings: 50 °C for spiked

samples double orbital pattern shaking at 700 rpm with 60-s shake / 60-s rest cycles, fluorescent

scans ($\lambda_{excitation}$ = 448 nm, $\lambda_{emission}$ = 482 nm) every 15 minutes, at a gain of 1,600, and a total run

time of 48 h.

Collection of fecal samples. Fecal samples included in this study were obtained from either the

field or from wildlife rehabilitation centers (Table 1). Individual fecal samples were collected

using sterile Whirl-Pak bags and sterile, single-use nitrile gloves (changed between sample

collects), then stored at -80 ℃ prior to analysis. Below we describe location information and

methods used to locate and collect fecal samples.

Northern Yellowstone ecosystem. Wolf, cougar, fox, coyote, and raven fecal samples included in

this study from the Northern Yellowstone ecosystem (NYE) were provided by the Yellowstone

Center for Resources, Yellowstone National Park, Wyoming, U.S.A. The NYE is a mountainous

region (ranging approximately 1500–2400 m) located in northwestern Wyoming, USA, and

south-central Montana, USA and is characterized primarily by lower elevation steppe and

grassland, and higher elevation coniferous forests, with relatively few wetland areas[41]. The NYE

is inhabited by several cervid prey species on which wolves and other predators prey[42]. CWD

had not been detected in the study area during sampling periods, nor had CWD been detected in

cervids within 30 kilometers of the study area. Fecal collection methods for wolf[43], followed by

fox, coyote, cougar, and raven from the NYE are described briefly. Feces from GPS radio

collared wolves (IACUC IMR YELL Smith wolves 2012) were collected in 2020 during early

winter (30 days: November 15-December 15) approximately 17 days following deposition.

Cougar, coyote, and fox fecal samples were located and collected from cougar GPS clusters near

cougar kill sites or by tracking fox and coyotes opportunistically during cougar cluster

investigations. Raven fecal samples were collected during capture, tagging, and handling of

ravens for a monitoring study following permits and animal handling protocols permitted by

state (Montana Scientific Collector's Permit 2022-020-W), federal (Master Banding Permit

22489), NPS (Yell-2022-SCI-8072), and University of Washington animal care and use

committee (Protocol 3077-01). All methods for feces collection were performed in accordance

with the relevant guidelines and regulations.

Southwest Wisconsin. As part of a multiyear CWD study centered in southwest Wisconsin, USA

the Wisconsin Department of Natural Resources (WDNR) captured and fit 763 deer with GPS

collars from 2017 to 2020 in Iowa, Grant, and Dane counties in southwestern Wisconsin,

following protocols approved by WDNR's Animal Care and Use Committee (Protocol: 16-

Storm-01). This region is characterized by high CWD prevalence[44]— an area where CWD was

first established in WI, has been estimated to have been present in the environment for over 20

years[45], and studied extensively in prior research[46-48]. Habitat in this area is characterized by

steep hills, forested ridges, deep river valleys, karst geology, and cold-water trout streams.

Elevations range from 184 to 524 m. Coyote are reported to leave numerous fecal deposits within

80 meters of carrion they are consuming[49], therefore fecal samples were collected by the field

crew opportunistically within 0-80 meters of collared WTD mortality sites from 2021-2022.

Deer GPS collars notified the field crew when the collar had been motionless for 4 or more

hours. Deer mortality sites were then identified by GPS points and/or VHF telemetry. Morality

sites were investigated to determine cause of death and those that had signs of predator activity

were searched for feces.

Northern Nebraska. The cougar fecal samples from the Pine Ridge and Niobrara Valley regions

in northern Nebraska, U.S.A. used in this study were provided by Nebraska Game and Parks

Commission (NGPC). Habitat in the Rine Ridge area is characterized by meadows, pine and

deciduous forests, steep buttes, small streams, and minor peaks (ranging approximately 900-1600

m). Habitat in the Niobrara Valley is characterized by steep hills, bluffs, pine forests and

canyons, boreal forests, grasslands, and the Niobrara River. As part of ongoing carnivore studies

in Cherry, Dawes, and Sheridan counties in Nebraska, U.S.A., cougar fecal samples were

collected by NGPC wildlife staff with the help of a trained detection dog and handler. While not

part of ongoing CWD surveillance efforts in Northwest Nebraska, cougar fecal samples included

in this study were collected within areas that are also designated big game research deer

management units (DMUs) where CWD has been detected.

Minnesota Wildlife Rehabilitation Center. Feces from adult bear, raccoon, and eagle were

collected during the month of September 2022 at the Minnesota Wild and Free Wildlife

Rehabilitation Center (MWF) in Garrison, Minnesota, USA. Animals that fecal samples were

collected from were originally found at different geographic locations prior to being brought to

WMF, varied in time post admittance to MWF, and had differing diets as appropriate for each

species. An admitted bear was found in Polk County, Minnesota, USA and the admitted raccoon

and eagle were found in Cass County, Minnesota, USA. Individuals whose feces were included

in this study were admitted between the months of May-July and were transported to MWF by

The Minnesota Department of Natural Resources (MNDNR). The bear diet consisted of dry dog

food supplemented with produce (i.e., apples, melon, corn, and assorted berries). The eagle diet

typically included fish and rodents (i.e., chipmunks, mice, or gophers). The raccoon diet

consisted of dry and/or wet dog food mixed with assorted produce (i.e., apples, melon, corn, and

assorted berries).

Dane County Humane Society Wildlife Center. Feces from a juvenile crow was collected during

the month of May 2022 at the Dane County Humane Society Wildlife Center in Madison,

Wisconsin, USA. The crow was admitted in early May and was fed a diet that consisted of eggs,

chicken, meal worms, seeds, nuts, and produce (i.e., apples, melon, corn, and assorted berries).

Data Analysis. Amyloid formation rate (AFR) data generated from the RT-QuIC assays were

analyzed and visualized using Jmp Pro 15 (SAS Institute, Cary, NC) and Prism 8 (GraphPad, San

Diego, CA). To apply a rigorous standard for distinguishing true positive samples from true

negatives, the AFR threshold times (i.e., the time at which amplification is determined to have

occurred in the RT-QuIC assay) were calculated by adding twenty times the standard deviation

of the relative fluorescence unit (RFU) values from cycles 3-14 to the mean of RFU values from

cycles 3-14. We previously applied this method to account for baseline fluorescence variation

amongst samples in determining if the sample was PrP^{CWD} positive[50]. Due to false seeding

observed in crow, bear, eagle, and raccoon negative control fecal samples, additional analysis

was required to distinguish true seeding from false seeding events for these species. To

accomplish this, empirical distributions of threshold times in hours of 24 replicates of unspiked

fecal samples were used to determine a threshold time that would yield a specificity $\geq$ 95% for

fecal samples from each species. These data were then used to determine a cycle end-time that

excluded $\geq$95% of the false seeding that occurred for each species. Cycle end-times for these

four feces were determined to be 20 h for crow (specificity of 95%), 24 h for bear (specificity

100%), 25 h for eagle (specificity 96%), and 17 h for raccoon (specificity 100%) (Supplementary

Fig. S1). All fecal samples analyzed by RT-QuIC were considered positive if a sample had at

least 3 out of 8 technical replicates (or half or more of 24 replicates) with seeding activity and

had AFR values that were significantly different from control AFR values based on statistical

analysis. Analyses for spiked samples were assessed with Dunnett's multiple comparison tests,

while for the surveillance samples, Kruskal-Wallis tests was used to distinguish which samples

had AFRs significantly different from the northern Yellowstone ecosystem (NYE) negative

control fecal sample.

Using AFR as a relative measure for prion concentration in a sample, we first evaluated our

ability to detect PrP^{CWD} from spiked fecal samples compared to the reference sample, and if the

AFR differed among carnivore species. We used a two-way (factorial) ANOVA to compare AFR

values between sample types (CWD-positive brain or spiked fecal samples from 9 different

species). We included an interaction between species and spike dilution; to assess if

detection/recovery in the different feces was sensitive to the concentration of the spike across the

10-fold dilution series. Significant interactions were determined using the Tukey HSD for

multiple comparisons.

Amyloid formation rates for field collected coyote and cougar fecal samples were found to be

non-normally distributed (Supplementary Figure S2). As such, the nonparametric Kruskal-Wallis

test was applied to compare individual fecal samples. If statistical differences were observed

across each individual fecal sample, then the non-parametric Steel-dewass post hoc test was used

to determine which individual samples differed from the negative control fecal sample.

RESULTS

Using RT-QuIC, our experimental spiking studies aimed to test extraction efficiency and

evaluate the presence of assay inhibitors through detection sensitivity of PrP^{CWD} in feces from

different scavenger and predator species using the reference sample as the source material for

spiking in all experiments (Fig. 1, see Table 1 for feces source information. Fecal samples for

each species were spiked with 10-fold dilutions of brain-derived PrP^{CWD} (Fig. 1). Variation in

assay sensitivity for the different dilutions of BH spike was observed across species. The limit of

relative PrP^{CWD} detection was significantly different from the negative controls in a species-

specific fashion. The most sensitive detection was from wolf feces, with 8/8 technical replicates

showing seeding activity (defined as having relative fluorescent units above the RFU threshold)

in feces spiked with only 10 ng of CWD-positive WTD BH. This was followed by eagle (6/8

seeding), which had a detection limit at 100 ng of brain-derived PrP^{CWD} (Fig. 1). Detection limits

were higher for other species, at a 10^3 ng spike for crow (6/8), bear (3/8), cougar (4/8), and raven

(8/8), a 10^4 ng spike for coyote (8/8) and raccoon (4/8) (Fig. 1), and a 10^5 ng spike for fox (8/8)

(Fig. 1).

The variable detection limits observed among species may be due to differences in inhibition of

the assay by the fecal material, the extraction process, or a combination of both. Overall, we

found differences across species do account for some of the variation seen in the magnitude of

AFRs for spiked feces when compared to dilutions of the reference sample on its own ($R^2 = 0.90$,

$F_{9, 420} = 155.46$, $p < 0.0001$). Mean AFRs for all species were significantly lower compared to

pure CWD-positive WTD BH and there were species differences in PrP^{CWD} recovery (Tukey

HSD post-hoc test for multiple comparisons, Supplementary Table S1, Fig. 2a). Spiked crow,

eagle, and wolf feces had significantly higher AFRs than the other species (Supplementary Table

S1, Fig. 2a). Cougar and fox feces had the lowest mean AFRs (Supplementary Table S1, Fig.

2a). AFR values from spiked raven did not differ significantly from wolf, bear, coyote, or

raccoon feces, however AFR values of bear, coyote, and raccoon fecal samples did differ from

the other species (Supplementary Table S1, Fig. 2a). AFRs for spiked fecal samples from all

species decreased across the dilution series ($F_{5,420} = 355.69$, $p < 0.0001$), with a significant

interaction between feces and spike dilution ($F_{45, 420}) = 9.36$, $p < 0.0001$). The magnitude of

differences in observed AFRs varied among species and by spike dilution, however there was a

general humped-shaped pattern observed for differences in recovery of PrP^{CWD} across many of

the species (Fig. 2b). The largest differences in recovery were observed at mid-level dilutions

rather than for the lowest and highest dilutions for all spiked feces samples, with an exception for

eagle and crow (Tukey HSD post-hoc tests, Fig. 2b, Supplementary Table S2).

Following these proof-of-concept spiking experiments, we then determined if we could detect

PrP^{CWD} in feces from free-ranging predators/scavengers from areas with CWD. We examined

coyote and cougar fecal samples collected from three distinct geographic regions where CWD

has been detected. In Iowa County, Wisconsin, six different coyote fecal samples were collected

within 80 meters of mortality sites for six different collared WTD in an area where CWD

prevalence is ~40% and 30% in adult males and females, respectively[51]. Of these WTD

carcasses, 5/6 were confirmed CWD-positive by RT-QuIC on tissues (i.e., brain or skin from ear

or belly; Fig. 3).

We also evaluated how variable PrP^{CWD} presence is across a single sample and how repeatable

our results for RT-QuIC were across plates by sampling and separately extracting and testing

three different regions of a given sample. RT-QuIC assays of three separate subsamples from

each coyote fecal sample (labeled as C1-C6) showed seeding activity in 8/8 of subsamples, with

exception of sample C3, where all eight technical replicates for the first and third extractions

showed seeding activity but had no seeding activity for the second extraction (Table 2). Coyote

feces AFRs were significantly greater than the controls (Kruskal-Wallis test: $X^2 = 106.29$,

p<0.0001, df = 6; Steel-dwass post hoc test: $p \le 0.00016$; Fig. 3; Table 2). Amyloid formation

rates of coyote feces were lower than those of WTD tissue from each mortality site but had

similar value ranges across each individual fecal sample and very little variation among technical

replicates of each scat subsample (Fig. 3).

Cougar fecal samples were collected from DMUs in the Niobrara Valley (2012: n=2; 2014: n=3)

and Pine Ridge area (2010: n=5; 2021: n=5) in Cherry, Dawes, and Sheridan Counties,

Nebraska, USA. AFRs of three Niobrara (one from 2012, and two from 2014) and three Pine

Ridge 2021 samples were significantly greater than the controls (Kruskal-Wallis test: X^2 =

255.25, p<0.0001, df = 15; Steel-dwass post hoc test: $p \leq$ 0.0226; Fig. 4). RT-QuIC assays of

three separate subsamples taken from each fecal sample showed variable seeding activity within

the 8 technical replicates, ranging from 2/8 to 8/8 (Table 3). Out of 24 replicates, at least half had

seeding activity for each of these six fecal samples (Table 3). The range of AFRs among these

fecal samples were variable, where values from the Pine Ridge 2021 samples were most elevated

compared to the three samples from Niobrara (Fig. 4). AFRs for the remaining samples were not

significantly greater than the controls (Steel-dwass post hoc test: $p \geq$ 0.05; Fig. 4). RT-QuIC

assays of three separate subsamples taken from each of these individual fecal samples also

showed variable seeding that ranged from 0/8 to 6/8 (Table 2). The total number of replicates

(out of 24) with seeding for each of these fecal samples was less than half (Table 3). Overall,

subsample replicate AFRs from cougar fecal samples (Fig. 4, Table 3) were more variable than

those seen from coyote feces (Fig. 3, Table 2).

Results of PrP^{CWD} occurrence in cougar feces from the 2010 (0/5) and 2021 (3/5) Pine Ridge

samples increase from 2010 to 2021 within this DMU[52] (Fig. 4, Table 3). Prevalence of CWD in

WTD for the Pine Ridge DMU was 7% (21/298; SE= 0.015) in 2010. Although host CWD

surveillance was not conducted in 2021 for the Pine Ridge DMU, surveillance in 2019 and 2022

estimated WTD CWD prevalence of 26% (69/264; SE= 0.027) and 27% (62/244; SE= 0.028),

respectively. The DMU where the Niobrara Valley cougar feces were collected has had host

CWD prevalence estimates of under 1% for the past 18 years. Estimated prevalence in 2012 was

0% (0/512), and although no surveillance was conducted from 2013-2016 for this DMU, 2018

prevalence was reported to be, 0.41% (1/243; SE= 0.004)[52]. Although a direct comparison of

PrP^{CWD} detection between cougar scat and WTD is difficult to make due to prevalence estimates

being very sensitive to small sample sizes[53], it appears that PrP^{CWD} can be detected in carnivore

scat even when CWD prevalence in WTD is very low.

Figure 1. Recovery and detection sensitivity of chronic wasting disease (CWD) prions (PrP^{CWD}) from spiking experiments of predator and scavenger feces by the real-time quaking-induced conversion (RT-QuIC) assay. Amyloid formation rates (AFRs) by RT-QuIC for predator and scavenger fecal samples spiked with 10-fold dilutions of 10% brain homogenate (from the obex region; BH) shown in nanograms (ng) from a CWD-positive white-tailed deer. Data points of AFRs ± standard deviation of eight technical replicates across each spiking dilution series in feces samples from crow, raven, eagle, raccoon, grey wolf, coyote, fox, cougar, and bear are shown. The dilution curve for the BH used as spiking material is shown for

393   comparison to PrP$^{\text{CWD}}$ recovery and detection sensitivity in feces. * = limit of detection

394   sensitivity of spiked material in each species. NC = negative control (feces without a BH spike).

 Figure 2. Differences in recovery and detection sensitivity of chronic wasting disease

(CWD) prions (PrP^{CWD}) from spiked carnivore feces. (a) Distribution of amyloid formations

rates (AFRs) across spiked fecal samples showing the median (solid black line) and first and

third quartiles (dotted black lines). CWD-positive brain homogenate (BH, the reference sample)

is shown in red, mammalian species in orange and avian species in purple. Letters show which

species are statistically different by Tukey HSD post-hoc test. **(b)** Illustration of the significant

least squares mean AFR value differences between 10-fold dilutions of CWD-positive brain

homogenate and recovered brain-derived PrP^{CWD} from spiked fecal samples by species and brain

spike dilutions using the Tukey HSD post-hoc test. For dilutions lacking a bar in the plots, there

was no significant difference in AFR between fecal samples and the reference sample.

411
412

Figure 3. The presence of chronic wasting disease (CWD) prions (PrP^CWD) in coyote fecal samples collected near collared white-tailed deer (WTD) mortality sites within Iowa County, Wisconsin, USA. Comparisons of real-time quaking-induced conversation (RT-QuIC) amyloid formation rates (AFR) of tissues (D1, obex; D2, obex; D3, belly skin; D4, ear pinna; D5, obex; D6, skin) from six different WTD carcasses (red) with six different coyote fecal samples found near each respective mortality site (blue). Samples are grouped by WTD mortality site ID (by number; D=deer and C=coyote from each site) on the x-axis. Data points show 8 and 24 replicates (3 separate extractions, 8 technical replicates each) of WTD tissue or coyote feces, respectively, with the median (solid black line) and first and third quartiles (dotted black lines). Negative control (NC; tan) represents 3 separate extraction results of 8 technical replicates each for coyote feces from 1 individual collected from the Northern Yellowstone Ecosystem.

428
429

Figure 4. Presence of chronic wasting disease (CWD) prions (PrP^{CWD}) in cougar fecal samples collected from two locations within Cherry, Dawes, and Sheridan Counties, Nebraska, USA. Real-time quaking-induced conversion (RT-QuIC) amyloid formation rates (AFR) from cougar feces collected from different individuals from the Niobrara Valley in 2012 (green) and 2014 (purple) and in 2010 (blue) and 2021 (orange) from the Pine Ridge region. Samples are grouped by individual cougar ID on the x-axis (F1.1 and F1.2 are two separate samples from the same individual). Data points shown are for 24 technical replicates (3 separate extractions of 8 technical replicates each), with the median (solid black line) and first and third quartiles (dotted black lines). Negative control (NC; tan) represents triplicate assay results for CWD- negative cougar feces from 1 individual collected from the Northern Yellowstone Ecosystem. * = positive for CWD based on Steel-dwass post-hoc test ($p \leq 0.05$).

Table 1. Species fecal sample collection location information, use, and storage conditions for

Species	Number of samples tested in triplicate	Year collected	Location of collection
Fecal samples used for RT-QuIC spiking assays			
Wolf	1	2020	Northern Yellowstone Ecosystem
Coyote	1	2022	Northern Yellowstone Ecosystem
Fox	1	2022	Northern Yellowstone Ecosystem
Raven	1	2022	Northern Yellowstone Ecosystem
Cougar	1	2022	Northern Yellowstone Ecosystem
Bear	1	2022	Minnesota Wild and Free Wildlife Rehabilitation Center (MWF) in Garrison, Minnesota.
Raccoon	1	2022	MWF
Eagle	1	2022	MWF
Crow	1	2022	Dane County Humane Society Wildlife Center, Madison, Wisconsin, U.S.A.
Fecal samples used for RT-QuIC diagnostics assays			
Coyote	6	2021-2022	Iowa County, Wisconsin, U.S.A.
Cougar	5	2012 & 2014	Niobrara Valley, Cherry County, Nebraska, U.S.A.
Cougar	10	2010 & 2021	Pine Ridge region, Dawes and Sheridan Counties, Nebraska, U.S.A.

samples included in this study. All samples were stored at –80°C.

Table 2. Presence or absence of chronic wasting disease (CWD) prions (PrP^{CWD}) from

subsamples of coyote (*Canis latrans*) fecal samples collected from within Iowa County,

Wisconsin, USA and one negative control (NC) coyote fecal sample collected from the Northern

Yellowstone Ecosystem (this separates the biological replicates shown together in Figure 3, to

note repeatability across subsamples) assessed by the real-time quaking-induced conversion (RT-

QuIC) assay. Values indicate the proportion of seeding activity for three separate extractions, 8

technical replicates each (24 total replicates) for each fecal sample.

Sample ID	Extraction 1	Extraction 2	Extraction 3	Total seeded reactions	PrP^{CWD} +/- for each extraction
C1	8/8	7/8	8/8	23/24	+/+/+
C2	8/8	8/8	8/8	24/24	+/+/+
C3	7/8	0/8	8/8	15/24	+/-/+
C4	8/8	8/8	8/8	24/24	+/+/+
C5	8/8	8/8	8/8	24/24	+/+/+
C6	8/8	8/8	8/8	24/24	+/+/+
NC	0/8	1/8	1/8	2/24	-/-/-

467 **Table 3.** Presence or absence of chronic wasting disease (CWD) prions (PrP^{CWD}) for fecal

468 samples from 15 different cougars (*Puma concolor*) from the Niobrara (N) of Pine Ridge (P)

469 deer management units within Cherry, Dawes, and Sheridan Counties, Nebraska, USA and one

470 negative control (NC) cougar fecal sample (collected from the Northern Yellowstone Ecosystem

Sample ID	Collection	Extraction	Extraction	Extraction	Total	PrP^{CWD}

471 (NYE), as depicted in Figure 4) assessed by the real-time quaking-induced conversion (RT-

472 QuIC) assay. Values indicate the proportion of seeding activity for three separate extractions, 8

473 technical replicates each (24 total replicates) for each fecal sample.

	Year/location	1	2	3	seeded reactions	+/- for each extraction
F1.1	2012/N	3/8	4/8	8/8	15/24	+/+/+
F1.2	2012/N	2/8	0/8	1/8	3/24	-/-/-
F2	2014/N	8/8	2/8	5/8	15/24	+/-/+
M1	2014/N	0/8	0/8	0/8	0/24	-/-/-
Unknown	2014/N	5/8	4/8	4/8	13/24	+/+/+
F3	2010/P	3/8	0/8	0/8	3/24	+/-/-
F4	2010/P	0/8	0/8	6/8	6/24	-/-/+
M2	2010/P	4/8	0/8	0/8	4/24	+/-/-
M3	2010/P	3/8	0/8	0/8	3/24	+/-/-
M4	2010/P	2/8	2/8	1/8	5/24	-/-/-
F5	2021/P	0/8	2/8	0/8	2/24	-/-/-
M5	2021/P	8/8	8/8	8/8	24/24	+/+/+
F6	2021/P	8/8	8/8	8/8	24/24	+/+/+
M6	2021/P	0/8	1/8	1/8	2/24	-/-/-
F7	2021/P	8/8	8/8	8/8	24/24	+/+/+
NC	2022/NYE	1/8	1/8	0/8	2/24	-/-/-

DISCUSSION

The role of sympatric wildlife in CWD epidemiology remains incompletely understood, but

cervid-consuming predators and scavengers may alter rates of disease spread through removal or

dispersal of PrP^{CWD} on the landscape. Determining the effects of cervid consumers on CWD

ecology and epidemiology could be facilitated with development of non-invasive and sensitive

methods with high throughput capacity for PrP^{CWD} detection. We presented results from

laboratory spiking experiments that CWD surveillance via fecal sampling is possible from a

range of mammalian and avian species. Such surveillance via fecal sampling could expand the

toolkit of available approaches, offering a non-invasive alternative to current surveillance

approaches, which typically requires animal capture. We then evaluated our methods using field-

collected feces from CWD endemic areas, determining that this approach is able to detect

PrP^{CWD} in the quantities present in coyote and cougar feces. Lack of tools that can be used to

detect the presence of PrP^{CWD} in abiotic and biotic environmental samples has hampered our

ability to address relevant ecological questions. This study directly addresses this need by

developing an important surveillance tool for CWD, allowing investigation of the epidemiology

of CWD at the community level.

The spiking experiments provided an assessment of how compatible RT-QuIC is with feces from

different cervid consumers. Limits of relative PrP^{CWD} detection ranged from 10^5 to 10 ng of

spiked material, depending on species and spike dilution (Fig. 1, Fig. 2a, b). These dilutions of

CWD-positive material are within the range of the relative PrP^{CWD} observed from feces of free-

ranging coyote and cougar. Comparing among carnivore species, wolf and eagle samples had the

most sensitive detection of brain derived PrP^{CWD} (Fig. 1), suggesting these species may be good

early sentinels of CWD in an area. However, an assay cutoff time of 25 hours had to be applied

to eagle results to remove false seeding events from unspiked samples, which could potentially

result in loss of sensitivity. This was also the case for bear, crow, and raccoon, where assay

cutoff times necessary to reduce false positives varied for each species. While these spiking

experiments provide a baseline of compatibility for feces from a range of carnivores with the

RT-QuIC assay, our study did not evaluate how intraspecific variation may affect the RT-QuIC

assay. It is also worth noting that feces from animals in rehabilitation centers needed an assay

cutoff time to reduce false seeding, whereas fecal samples from free-ranging animals did not

require an assay cutoff time to avoid false positives. Thus, dietary variation or other species- or

individual-specific differences may affect assay results. Further analyses of how differences in

diet and fecal composition affect assay sensitivity and specificity may improve our

understanding of assay strength and limitations.

Based on results of the spiking experiments, carnivore feces from certain cervid consumers could

be a useful non-invasive approach to CWD surveillance, possibly able to detect CWD when

prevalence rates in deer are very low. Our findings from both the Pine Ridge and Niobrara

Valley DMUs highlight the usefulness of incorporating carnivore feces in CWD surveillance

strategies. If predators select infected prey at a greater rate than uninfected prey, as has been

observed with cougars[54], predator feces may serve as early sentinels of CWD detection in areas

where CWD is not yet endemic or where areas where surveillance efforts are limited, as we have

demonstrated here with cougar feces. (Fig. 4, Table 3).

Differences in PrP^{CWD} recovery from feces spiked with CWD-positive WTD brain compared to

the reference sample followed a similar trend among carnivores based on AFR values. Smaller

differences were observed between the brain positive controls and spiked samples at the most

and least concentrated spike dilutions compared to larger differences in detection at the mid-

concentration dilutions (Fig. 2b). This hump-shaped variation in prion recovery across dilutions

may be a result of the extraction process used for the feces samples following spiking, where

some loss of seeding material likely occurred and was most sensitively observed at these spiking

dilutions for each feces sample. It is also possible that the variability in AFRs that occurred for

the 10^6 ng and 10 ng concentrations of pure CWD-positive WTD BH was similar enough to that

observed in many of the spiked fecal samples, that the differences were lower by comparison to

the less variable AFRs from all other dilutions of pure CWD-positive WTD BH (Fig. 1, Fig. 2b).

Additionally, the compounds that a given sample is composed of may influence the RT-QuIC

reaction itself, as has been demonstrated with the inhibitory effect of certain sample types[55, 56].

Thus, it remains possible that this trend may be due to similar fecal biochemical compositions

from each species that are resulting in similar recovery effects.

Detection of brain-derived PrP^{CWD} from wolf feces was the most sensitive across species, did not

require an assay cutoff time to distinguish true seeding from false seeding events, and could be

relatively easy to collect on a seasonal basis in more temperate CWD endemic areas such as the

upper Midwest. The sensitivity and specificity results from the spiking assay and ease of sample

collection suggest that wolf may be an optimal candidate for incorporating into CWD

surveillance strategies, within its range. However, there are other considerations when designing

fecal sampling programs in different environments, such as land access, effort required for

sample collection, and species density. From the selection of species evaluated here, most of the

mammal and avian species, with the exception of fox, may be candidates for further evaluation

by RT-QuIC. Although spiking assays for coyote and cougar feces revealed moderate sensitivity

compared to wolf (Fig 1), these species also did not require an assay cutoff time and exhibited

100% specificity, compared to ~90% specificity for wolf (1/8 false seeding rate; Fig. 1),

suggesting they may also be good candidates for CWD surveillance. Raven may also be a good

candidate due to smaller differences in detection of PrP^{CWD} compared to the reference sample,

however ease of collection of avian feces may limit utility for determining CWD occurrence in a

given area compared with mammalian feces. In this study, we readily had access to coyote and

cougar feces from areas where CWD is either endemic or has been detected at low prevalence.

We demonstrated detection of PrP^{CWD} using spiking assays for these two species, yet they were

not as sensitive as wolf samples. Because these larger scavenger and predator species tend to

consume larger amounts of biomass in one feeding—and thus, presumably more PrP^{CWD}—

detection of PrP^{CWD} from their feces may be more likely than fecal samples from avian species.

This suggests that analysis of mammalian carnivore feces may offer a way to advance CWD

surveillance strategies.

Detection of PrP^{CWD} within some of the individual field collected Nebraska cougar feces

subsamples was more variable than Wisconsin coyote subsamples (Fig. 3, Fig. 4, Table 2, Table

3). This may have been the result of differences in sample age, as increased sample age may

reduce detection by RT-QuIC for some sample types[57]. Wisconsin coyote feces were less than a

week old and had seeding ratios similar to the Pine Ridge 2021 cougar feces. Given that the more

recently collected Nebraska cougar feces (Pine Ridge 2021) had higher overall AFRs in samples

with PrP^{CWD} present and the most consistent detection among all other samples in that group

compared to the other groups of Nebraska cougar feces (Niobrara 2012, 2014; Pine Ridge 2010),

suggests that this variation could have been due to differences in sample age. In addition, lower

amounts of seeding material have been shown to result in more variable AFRs using RT-QuIC[31],

thus it is also possible that this variability in more recent Nebraska cougar fecal subsamples

could be the result of overall lower levels of PrP^{CWD} present in the sample compared to

Wisconsin coyote samples. Homogenizing or mixing the whole sample prior to RT-QuIC might

yield more consistent seeding from a given sample. However, because microparticles can lead to

complications in the RT-QuIC[39], we chose to initially evaluate subsampling of primary samples

rather than whole sample-homogenization in this study. Future studies comparing homogenized

vs. unhomogenized samples for a given species could help determine how homogenization of a

whole sample influences RT-QuIC reactions, recovery of PrP^{CWD}, and detection sensitivity and

specificity. Additionally, factors such as amount of biomass consumed, gut residence time of

PrP^{CWD} and time spent at carcass sites, or whether they are visiting a site repeatedly and feeding

over the course of several days are factors that warrant further investigation.

The results of the Wisconsin field collected coyote feces found at deer mortality sites

demonstrate that triplicate subsampling of each fecal sample was sufficient to overcome the

variability of sensitivity for a given sample and demonstrate PrP^{CWD} presence using the RT-

QuIC assay (Fig. 3, Table 2). Sample C3 had the most variable seeding activity and also had the

lowest average AFRs compared to all other Wisconsin coyote samples, suggesting that the

variability was likely due to lower overall levels of PrP^{CWD} present in this sample, a common

response of the RT-QuIC assay to samples with lower prion loads[31] (Fig. 3, Table 2). In addition,

relative loads of PrP^{CWD} of each WI coyote fecal sample were reduced compared to tissues from

the CWD-infected deer carcass they may have been scavenging (Fig. 3).

Because coyotes reportedly leave numerous scent marking feces within 1 to 80 m of the carrion

they are consuming, we find it reasonable to consider that some of these fecal samples came

from coyote that had scavenged on the respective carcass near the location their fecal samples

were collected[49]. It is also worth considering that since one of the coyote fecal samples was

collected on the periphery of a CWD-negative carcass site, —yet still had PrP^{CWD} present, some

of these individuals may have been feeding on more than one carcass. This makes it difficult to

interpret the reason for reduced loads of PrP^{CWD} in fecal samples compared to host tissue, to

ascertain if reduced PrP^{CWD} loads are an effect of digestion or an effect of mixed consumption of

different tissues from healthy and CWD-infected carcasses or other species in their diet. The

reduced prion loads in coyote feces compared to deer tissue could be due to different or mixed

dietary sources than just the sampled carcass, prion removal during digestion[27], a species or

sample type effect on prion recovery[58], or a loss of seeding material during sample extraction.

Additional studies assessing the relationship between CWD-infected biomass consumption and

field-collected predator and scavenger fecal samples could help decipher the role of these species

with respect to PrP^{CWD} environmental removal or deposition. Overall, these results support our

hypothesis that CWD surveillance using carnivore feces may provide a useful estimate of CWD

occurrence.

The developments we have reported here are critical steps in elucidating the roles of scavengers and predators in CWD epidemiology and for advancing and complimenting existing CWD surveillance strategies. Adding or altering ongoing surveillance efforts to include collection and analysis of predator and scavenger feces may call for additional funding, personnel, and logistical considerations by wildlife management agencies. However, CWD surveillance often relies upon hunter-harvest submission of tissues, followed by costly testing programs, typically funded by the state or province wildlife management agency. The ability to detect PrP^{CWD} in feces from cervid-consuming predators and scavengers may offer a cost-effective surveillance alternative for potentially early CWD detection and management action and could provide an efficient means to surveille at-risk areas neighboring CWD outbreak zones or areas where hunter-harvesting or hunter-submitted sampling is low, or where noninvasive approaches are desirable.

ACKNOWLEDGEMENTS. This work was supported by the Wisconsin Department of Natural Resources (37000-0000009433 and 37000-0000010649). We thank the Yellowstone Biological Technicians that helped collect fecal samples for this project, especially Connor Meyer, Nikki Tatton, Jeremy SunderRaj, and Jack Rabe. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

AUTHOR CONTRIBUTIONS. Conception and design of the work: H.N.I.; acquisition, analysis, and interpretation of the RT-QuIC data: H.N.I., W.C.T, S.S.L; writing the original draft: H.N.I., revision and edition of the final manuscript: H.N.I., E.E.B, S.P.W., M. H., D.R.S,

K.W., D.P.W., T.N., D.J.S., S.S.L., and W.C.T. All authors approved and agreed to the

submitted version of the present manuscript.

COMPETING INTERESTS. The authors declare no competing interests.

DATA AVAILABILITY. All data generated or analyzed during this study are included in this

published article.

ADDITIONAL INFORMATION. Supplementary information is available for this paper.

Correspondence and requests for materials should be addressed to inzalaco@wisc.edu

REFERENCES

[1] Wilson EE, Wolkovich EM: Scavenging: how carnivores and carrion structure communities. Trends in Ecology & Evolution 2011, 26:129-35.

[2] Barton PS, Cunningham SA, Lindenmayer DB, Manning AD: The role of carrion in maintaining biodiversity and ecological processes in terrestrial ecosystems. Oecologia 2013, 171:761-72.

[3] Houston DC, Cooper JE: The digestive tract of the whitebacked griffon vulture and its role in disease transmission among wild ungulates. J Wild Dis 1975, 11:306–13.

[4] Seabloom EW, Peace A, Asik L, Everett RA, Frenken T, González AL, Strauss AT, Van de Waal DB, White LA, Borer ET: Dead or alive: carbon as a currency to integrate disease and ecosystem ecology theory. Oikos 2023:e09880.

[5] Brandell EE, Cross PC, Smith DW, Rogers W, Galloway NL, MacNulty DR, Stahler DR, Treanor J, Hudson PJ: Examination of the interaction between age-specific predation and chronic disease in the Greater Yellowstone Ecosystem. Journal of Animal Ecology 2022.

[6] Ogada D, Torchin M, Kinnaird M, Ezenwa V: Effects of vulture declines on facultative scavengers and potential implications for mammalian disease transmission. Conservation biology 2012, 26:453-60.

[7] Vicente J, Carrasco R, Acevedo P, Montoro V, Gortazar C: Big game waste production: sanitary and ecological implications. Integrated Waste Management-Volume II: IntechOpen, 2011.

[8] Barandongo ZR, Dolfi AC, Bruce SA, Rysava K, Huang Y-H, Joel H, Hassim A, Kamath PL, van Heerden H, Turner WC: The persistence of time: the lifespan of Bacillus anthracis spores in environmental reservoirs. Res Microbiol 2023:104029.

[9] Cui Y, Schmid BV, Cao H, Dai X, Du Z, Ryan Easterday W, Fang H, Guo C, Huang S, Liu W: Evolutionary selection of biofilm-mediated extended phenotypes in Yersinia pestis in response to a fluctuating environment. Nature communications 2020, 11:281.

[10] Hennebique A, Gas F, Batina H, De Araujo C, Bizet K, Maurin M: Evaluation of the biotoxis qPCR detection kit for Francisella tularensis detection in clinical and environmental samples. J Clin Microbiol 2020, 59:e01434-20.

[11] Williams ES, Young S: Chronic wasting disease of captive mule deer: a spongiform encephalopathy. J Wildl Dis 1980, 16:89-98.

[12] Richards B: Distribution of Chronic Wasting disease in North America. The United States Geological Survey, National Wildlife Heath Center, 2023.

[13] Benestad SL, Mitchell G, Simmons M, Ytrehus B, Vikoren T: First case of chronic wasting disease in Europe in a Norwegian free-ranging reindeer. Vet Res 2016, 47:88.

[14] Vikoren T, Vage J, Madslien KI, Roed KH, Rolandsen CM, Tran L, Hopp P, Veiberg V, Heum M, Moldal T, Neves CGD, Handeland K, Ytrehus B, Kolbjornsen O, Wisloff H, Terland R, Saure B, Dessen KM, Svendsen SG, Nordvik BS, Benestad SL: First Detection of Chronic Wasting Disease in a Wild Red Deer (Cervus elaphus) in Europe. J Wildl Dis 2019, 55:970-2.

[15] Kim TY, Shon HJ, Joo YS, Mun UK, Kang KS, Lee YS: Additional cases of Chronic Wasting Disease in imported deer in Korea. J Vet Med Sci 2005, 67:753-9.

[16] Angers RC, Browning SR, Seward TS, Sigurdson CJ, Miller MW, Hoover EA, Telling GC: Prions in skeletal muscles of deer with chronic wasting disease. Science 2006, 311:1117.

[17] Angers RC, Seward TS, Napier D, Green M, Hoover E, Spraker T, O'Rourke K, Balachandran A, Telling GC: Chronic wasting disease prions in elk antler velvet. Emerg Infect Dis 2009, 15:696-703.

[18] Chen B, Morales R, Barria MA, Soto C: Estimating prion concentration in fluids and tissues by quantitative PMCA. Nat Methods 2010, 7:519-20.

[19] Henderson DM, Tennant JM, Haley NJ, Denkers ND, Mathiason CK, Hoover EA: Detection of chronic
wasting disease prion seeding activity in deer and elk feces by real-time quaking-induced conversion. J
Gen Virol 2017, 98:1953-62.
[20] Haley NJ, Seelig DM, Zabel MD, Telling GC, Hoover EA: Detection of CWD Prions in Urine and Saliva
of Deer by Transgenic Mouse Bioassay. Plos One 2009, 4.
[21] Mathiason CK, Hays SA, Powers J, Hayes-Klug J, Langenberg J, Dahmes SJ, Osborn DA, Miller KV,
Warren RJ, Mason GL, Hoover EA: Infectious prions in pre-clinical deer and transmission of chronic
wasting disease solely by environmental exposure. PLoS One 2009, 4:e5916.
[22] Moreno JA, Telling GC: Molecular Mechanisms of Chronic Wasting Disease Prion Propagation. Cold
Spring Harbor Perspectives in Medicine 2018, 8.
[23] Hoover CE, Davenport KA, Henderson DM, Denkers ND, Mathiason CK, Soto C, Zabel MD, Hoover
EA: Pathways of Prion Spread during Early Chronic Wasting Disease in Deer. J Virol 2017, 91.
[24] Miller MW, Williams ES, Hobbs NT, Wolfe LL: Environmental sources of prion transmission in mule
deer. Emerging infectious diseases 2004, 10:1003.
[25] Johnson CJ, Phillips KE, Schramm PT, McKenzie D, Aiken JM, Pedersen JA: Prions adhere to soil
minerals and remain infectious. PLoS Pathog 2006, 2:e32.
[26] Almberg ES, Cross PC, Johnson CJ, Heisey DM, Richards BJ: Modeling routes of chronic wasting
disease transmission: environmental prion persistence promotes deer population decline and extinction.
PLoS One 2011, 6:e19896.
[27] Nichols TA, Fischer JW, Spraker TR, Kong Q, VerCauteren KC: CWD prions remain infectious after
passage through the digestive system of coyotes (Canis latrans). Prion 2015, 9:367-75.
[28] VerCauteren KC, Pilon JL, Nash PB, Phillips GE, Fischer JW: Prion remains infectious after passage
through digestive system of American crows (Corvus brachyrhynchos). PLoS One 2012, 7:e45774.
[29] Baune C, Wolfe LL, Schott KC, Griffin KA, Hughson AG, Miller MW, Race B: Reduction of Chronic
Wasting Disease Prion Seeding Activity following Digestion by Mountain Lions. Msphere 2021,
6:e0081221.
[30] Fisher MC, Prioreschi RA, Wolfe LL, Runge JP, Griffin KA, Swanson HM, Miller MW: Apparent
stability masks underlying change in a mule deer herd with unmanaged chronic wasting disease.
Commun Biol 2022, 5:15.
[31] Wilham JM, Orru CD, Bessen RA, Atarashi R, Sano K, Race B, Meade-White KD, Taubner LM, Timmes
A, Caughey B: Rapid end-point quantitation of prion seeding activity with sensitivity comparable to
bioassays. PLoS Pathog 2010, 6:e1001217.
[32] Saa P, Castilla J, Soto C: Presymptomatic detection of prions in blood. Science 2006, 313:92-4.
[33] Haley NJ, Carver S, Hoon-Hanks LL, Henderson DM, Davenport KA, Bunting E, Gray S, Trindle B,
Galeota J, LeVan I: Detection of chronic wasting disease in the lymph nodes of free-ranging cervids by
real-time quaking-induced conversion. J Clin Microbiol 2014, 52:3237-43.
[34] Henderson DM, Davenport KA, Haley NJ, Denkers ND, Mathiason CK, Hoover EA: Quantitative
assessment of prion infectivity in tissues and body fluids by real-time quaking-induced conversion. J Gen
Virol 2015, 96:210-9.
[35] Haley NJ, Siepker C, Walter WD, Thomsen BV, Greenlee JJ, Lehmkuhl AD, Richt JA: Antemortem
detection of chronic wasting disease prions in nasal brush collections and rectal biopsy specimens from
white-tailed deer by real-time quaking-induced conversion. J Clin Microbiol 2016, 54:1108-16.
[36] Henderson DM, Denkers ND, Hoover CE, Garbino N, Mathiason CK, Hoover EA: Longitudinal
Detection of Prion Shedding in Saliva and Urine by Chronic Wasting Disease-Infected Deer by Real-Time
Quaking-Induced Conversion. J Virol 2015, 89:9338-47.
[37] Henderson DM, Manca M, Haley NJ, Denkers ND, Nalls AV, Mathiason CK, Caughey B, Hoover EA:
Rapid antemortem detection of CWD prions in deer saliva. PLoS One 2013, 8:e74377.

[38] Tennant JM, Li M, Henderson DM, Tyer ML, Denkers ND, Haley NJ, Mathiason CK, Hoover EA:
Shedding and stability of CWD prion seeding activity in cervid feces. PLoS One 2020, 15:e0227094.
[39] Orru CD, Groveman BR, Hughson AG, Manca M, Raymond LD, Raymond GJ, Campbell KJ, Anson KJ,
Kraus A, Caughey B: RT-QuIC Assays for Prion Disease Detection and Diagnostics. Methods Mol Biol
2017, 1658:185-203.
[40] Metrick MA, 2nd, do Carmo Ferreira N, Saijo E, Hughson AG, Kraus A, Orru C, Miller MW, Zanusso G,
Ghetti B, Vendruscolo M, Caughey B: Million-fold sensitivity enhancement in proteopathic seed
amplification assays for biospecimens by Hofmeister ion comparisons. Proc Natl Acad Sci U S A 2019.
[41] Houston DB: The northern Yellowstone elk : ecology and management. New York: Macmillan, 1982.
[42] Smith DW, Drummer TD, Murphy KM, Guernsey DS, Evans SB: Winter prey selection and estimation
of wolf kill rates in Yellowstone National Park, 1995–2000. The Journal of Wildlife Management 2004,
68:153-66.
[43] Brandell EE, Jackson MK, Cross PC, Piaggio AJ, Taylor DR, Smith DW, Boufana B, Stahler DR, Hudson
PJ: Evaluating noninvasive methods for estimating cestode prevalence in a wild carnivore population.
Plos one 2022, 17:e0277420.
[44] Joly DO, Ribic CA, Langenberg JA, Beheler K, Batha CA, Dhuey BJ, Rolley RE, Bartelt G, Van Deelen
TR, Samuel MD: Chronic wasting disease in free-ranging Wisconsin white-tailed deer. Emerging
infectious diseases 2003, 9:599.
[45] Wasserberg G, Osnas EE, Rolley RE, Samuel MD: Host culling as an adaptive management tool for
chronic wasting disease in white-tailed deer: a modelling study. Journal of Applied Ecology 2009, 46:457-
66.
[46] Jennelle CS, Samuel MD, Nolden CA, Berkley EA: Deer Carcass Decomposition and Potential
Scavenger Exposure to Chronic Wasting Disease. Journal of Wildlife Management 2009, 73:655-62.
[47] Skuldt LH, Mathews NE, Oyer AM: White-Tailed Deer Movements in a Chronic Wasting Disease Area
in South-Central Wisconsin. Journal of Wildlife Management 2008, 72:1156-60.
[48] Grear DA, Samuel MD, Langenberg JA, Keane D: Demographic patterns and harvest vulnerability of
chronic wasting disease infected white-tailed deer in Wisconsin. The Journal of Wildlife Management
2006, 70:546-53.
[49] Barja I, List R: The role of spatial distribution of faeces in coyote scent marking behaviour. Polish
Journal of Ecology 2014, 62:373-84.
[50] Inzalaco HN, Bravo-Risi F, Morales R, Walsh DP, Storm DJ, Pedersen JA, Turner WC, Lichtenberg SS:
Ticks harbor and excrete chronic wasting disease prions. Sci Rep 2022, Manuscript submitted.
[51] Wisconsin Department of Natural Resources: CWD prevalence in Wisconsin. 2023.
[52] Nebraska Game and Parks: Chronic Wasing Disease Testing Results. Lincoln, Nebraska
Nebraska.gov, 2022.
[53] Gregory R, Woolhouse M: Quantification of parasite aggregation: a simulation study. Acta Trop
1993, 54:131-9.
[54] Krumm CE, Conner MM, Hobbs NT, Hunter DO, Miller MW: Mountain lions prey selectively on
prion-infected mule deer. Biology Letters 2010, 6:209-11.
[55] Hoover CE, Davenport KA, Henderson DM, Zabel MD, Hoover EA: Endogenous Brain Lipids Inhibit
Prion Amyloid Formation In Vitro. J Virol 2017, 91.
[56] Davenport KA, Hoover CE, Denkers ND, Mathiason CK, Hoover EA: Modified Protein Misfolding
Cyclic Amplification Overcomes Real-Time Quaking-Induced Conversion Assay Inhibitors in Deer Saliva
To Detect Chronic Wasting Disease Prions. J Clin Microbiol 2018, 56.
[57] Li M, Schwabenlander MD, Rowden GR, Schefers JM, Jennelle CS, Carstensen M, Seelig D, Larsen
PA: RT-QuIC detection of CWD prion seeding activity in white-tailed deer muscle tissues. Scientific
Reports 2021, 11:16759.

[58] Tewari D, Fasnacht M, Ritzman M, Livengood J, Bower J, Lehmkuhl A, Nichols T, Hamberg A,
Brightbill K, Henderson D: Detection of chronic wasting disease in feces and recto-anal mucosal
associated lymphoid tissues with RT-QuIC in a naturally infected farmed white-tailed deer herd. Front
Vet Sci 2022, 9:959555.