Structure of Orthoreovirus RNA Chaperone σNS, a Component of Viral Replication Factories

Boyang Zhao,[1] Liya Hu,[2] Soni Kaundal,[2] Neetu Neetu, Christopher H. Lee,[3,4] Xayathed Somoulay,[3,4] Banumathi Sankaran[6], Gwen M. Taylor,[4,5] Terence S. Dermody,[3,4,5*] and B. V. Venkataram Prasad[1,2*]

[1]Department of Molecular Virology and Microbiology, Baylor College of Medicine

[2]Verna and Marrs Mclean Department of Biochemistry and Molecular Pharmacology, Baylor College of Medicine

[3]Department of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine

[4]Institute of Infection, Inflammation, and Immunity, UPMC Children's Hospital of Pittsburgh

[5]Department of Pediatrics, University of Pittsburgh School of Medicine

[6]Berkeley Center for Structural Biology, Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley Laboratory, Berkeley, CA, U.S.A.

*Corresponding authors (terence.dermody@chp.edu; vprasad@bcm.edu)

Abstract

The reovirus σNS RNA-binding protein is required for formation of intracellular compartments during viral infection that support viral genome replication and capsid assembly. Despite its functional importance, a mechanistic understanding of σNS is lacking. We conducted structural and biochemical analyses of an R6A mutant of σNS that forms dimers instead of the higher-order oligomers formed by wildtype (WT) σNS. The crystal structure of selenomethionine-substituted σNS-R6A reveals that the mutant protein forms a stable antiparallel dimer, with each subunit having a well-folded central core and a projecting N-terminal arm. The dimers interact with each other by inserting the N-terminal arms into a hydrophobic pocket of the neighboring dimers on either side to form a helical assembly that resembles filaments of WT σNS in complex with RNA observed using cryo-EM. The interior of the crystallographic helical assembly is positively charged and of appropriate diameter to bind RNA. The helical assembly is disrupted by bile acids, which bind to the same hydrophobic pocket as the N-terminal arm, as demonstrated in the crystal structure of σNS-R6A in complex with bile acid. This finding suggests that the N-terminal arm functions in conferring context-dependent oligomeric states of σNS, which is supported by the structure of σNS lacking the N-terminal arm. We discovered that σNS displays RNA helix destabilizing and annealing activities, likely essential for presenting mRNA to the viral RNA-dependent RNA polymerase for genome replication. The RNA chaperone activity is reduced by bile acids and abolished by N-terminal arm deletion, suggesting that the activity requires formation of σNS oligomers. Our studies provide structural and mechanistic insights into the function of σNS in reovirus replication.

Introduction

Most viruses that replicate in the cytoplasm of host cells form neoorganelles that serve as sites of viral genome replication and particle assembly. These highly specialized viral factory (VF) structures concentrate viral replication proteins and nucleic acids, limit activation of cell-intrinsic defenses, and coordinate release of progeny particles. Both viral and cellular proteins contribute to VF formation and orchestrate viral replication functions.

Mammalian orthoreoviruses (reoviruses) are members of the Orthoreovirus genus in the *Reoviridae* family. Reoviruses are classified into four serotypes based on sequence and antigenicity of the σ1 viral attachment protein and include prototype strains type 1 Lang, type 2 Jones, type 3 Dearing, and type 4 Ndelle [1]. While reovirus can infect many mammalian species, including humans, symptomatic infection develops only in neonatal mammals and human infants and children [2]. Infection by some reovirus strains appears to induce the development of celiac disease, a complex immune disorder characterized by an immune response to dietary gluten that can damage the intestinal lining and result in diarrhea and malabsorption [3]. Reovirus can induce apoptosis, necroptosis, and pyroptosis in tumor cells by activating programmed cell death pathways [4]. Because of these characteristics, reovirus was one of the first viruses used as an oncolytic agent [5].

Reovirus forms nonenveloped particles with an icosahedral capsid, 800 Å in diameter, consisting of two concentric protein layers and distinctive turret-like spikes projecting from the twelve icosahedral vertices. Encapsidated within the innermost protein layer is the viral genome, consisting of ten segments of linear double-stranded (ds) RNA, which encode eleven viral proteins. Eight of these proteins are structural and form the two capsid layers, turret, attachment protein, and an internally located RNA-dependent RNA polymerase. There are three

nonstructural proteins that function to promote viral replication. Reovirus replication occurs in the cytoplasm. After binding to cellular receptors, the virus particle enters many types of cells by clathrin-mediated endocytosis [6]. Within endosomes, proteolysis of outer-capsid proteins yields infectious subvirion particles, which penetrate endosomal membranes to release transcriptionally active viral cores into the cytoplasm [7]. Following transcription of dsRNA segments within the core interior, capped viral mRNAs are extruded through the turrets into the cytoplasm. The capped transcripts serve as templates for translation as well as synthesis of minus-strand genomic RNA by the viral polymerase to produce the dsRNA genome segments. Assortment of the genome segments into progeny core particles and addition of outer-capsid proteins occur inside VFs. Nonstructural proteins σNS and μNS are required to nucleate these compartments during infection.

In addition to its essential function in formation of viral replication factories, σNS recruits viral RNA [8] and both the eukaryotic translation initiation factor 3 subunit A (eIF3A) and the ribosomal subunit pS6R to enhance viral RNA translation [9]. Its association with the major stress granule effector protein, Ras-GAP SH3-binding protein 1 (G3BP1), disrupts stress granule formation, which relieves a block to reovirus replication [10]. σNS binds viral ssRNA nonspecifically [11], protects ssRNAs from degradation, and ferries viral mRNAs to VFs [12]. A mechanistic understanding of this multifunctional protein is lacking without its atomic-resolution structure. Our previous studies using cryo-EM show that σNS forms heterogeneous, higher-order oligomers and, when associated with ssRNA, it forms filamentous structures [12]. However, neither the unliganded nor the RNA-bound form of σNS are conducive to high-resolution structural studies.

We found that an arginine-to-alanine mutation at residue 6 in σNS (σNS-R6A) prevents RNA binding and formation of higher-order oligomers, thus facilitating σNS structural and biochemical characterization. From these analyses, we discovered that domain-swapping interactions of the flexible N-terminal arms of σNS dimers facilitate the association of dimers into filamentous structures that can bind ssRNA. Moreover, we found that such assemblies are required for an RNA chaperone activity of σNS, which we postulate is required for reovirus replication. Bile acids bind to the same hydrophobic pocket as the N-terminal arms and disrupt filament formation and impede RNA chaperone activity. Collectively, these findings provide mechanistic understanding of σNS function in reovirus replication and provide clues about dsRNA synthesis in a *Reoviridae* virus.

Results

R6A mutation disrupts σNS RNA binding and oligomerization

Initial attempts to crystalize wildtype (WT) σNS were unsuccessful due to binding of the protein to RNA, which leads to formation of higher-order oligomers that disassemble into heterogeneous populations during purification and crystallization. The N-terminal 38 residues of σNS include several amino acids required for σNS binding to RNA [12]. These residues are conserved in all reovirus σNS sequences reported to date [13]. We previously engineered σNS mutants incapable of binding RNA to assess the requirement of σNS RNA-binding activity in reovirus replication [8]. The σNS-R6A mutant does not distribute to VFs and fails to transport viral mRNA to these structures [8]. Based on its lack of RNA-binding capacity, we selected the σNS-R6A mutant for purification and crystallization. Expression and purification of σNS-R6A yielded pure and homogenous recombinant protein. Following anion-exchange chromatography, we compared

recombinant WT σNS and σNS-R6A for interactions with nucleic acid (**Fig. 1a**). Anion-

exchange chromatography of WT σNS showed two peaks in contrast to that of the σNS-R6A

mutant, which showed one peak, consistent with the failure of σNS-R6A to bind RNA. Size-

exclusion chromatography of WT σNS following anion-exchange chromatography, yielded an

oligomer of a size corresponding to a decamer (**Fig. 1b**) in contrast to σNS-R6A, which showed

a peak corresponding to a dimer (~80 kDa). Surprisingly, the selenomethionine (Se-Met)-

substituted σNS-R6A, which we purified for crystallization attempts, showed a size-exclusion

chromatography profile suggesting multimeric association with a peak corresponding to an

octamer.

σNS forms a stable core structure with a flexible N-terminal arm

To gain a mechanistic understanding of biological functions of σNS, we determined the structure

of the σNS-R6A mutant. Although we successfully obtained diffracting crystals of σNS-R6A,

determination of the structure by molecular replacement was not possible without suitable

phasing models. Accordingly, we proceeded with Se-Met-substituted σNS-R6A and determined

the structure using SAD phasing methods to a resolution of 3.0 Å (**Table 1**). Se-Met σNS-R6A

crystallized in the space group $P6_5$ with two molecules in the asymmetric unit. The crystal

structure of σNS-R6A reveals a stable core structure with a protruding N-terminal arm (**Fig. 2a**).

The core structure is formed by 13 β-strands and 19 α-helices connected by β- or γ-turns (**Fig.**

2b). The β-strands form four, antiparallel β-sheets on the surface of a globular central body

formed by α-helices. The N-terminal residues 4 to 11 form the first helix, which is connected to a

short strand by a seven-residue loop. A nine-residue β-hairpin follows the short strand and

connects to the first antiparallel sheet of σNS. This antiparallel sheet from residues 30 to 40

together with the N-terminus forms a cleft of about ~ 25 Å in width. There are several long

helices connected by β-turns from residues 245 to 335. The C-terminal region of σNS is

connected to helix 19 by a γ-turn and consists of four strands.

To define flexible regions of σNS, we conducted molecular dynamics simulations based

on the σNS-R6A structure. The simulations showed that the σNS N-terminus displays five-fold

greater flexibility relative to the other regions of the protein, as assessed by the root mean square

flexibility (RMSF) (**Fig. 2c**). The β-sheet connected to the N-terminal loop represents a second

region of flexibility, suggesting that σNS contains a flexible N-terminal region. Interestingly, the

addition of ssRNA into the simulation diminishes the RMSF of the N-terminus by about 25%.

σNS forms a stable dimer

The crystallographic asymmetric unit contains two σNS-R6A subunits that form an antiparallel

dimer related by non-crystallographic two-fold symmetry, with the N-terminal arms projecting

away at either end (**Fig. 3a**). The σNS-R6A dimer is stabilized by an interface with a calculated

buried surface area (BSA) of 1853 $Å^2$, indicating a biologically relevant oligomer that is stable in

solution. The σNS-R6A dimer is stabilized mainly by hydrogen bonds between multiple residues

(**Fig. 3b**). Since the dimer interface displays two-fold symmetry, equivalent clusters of residues

are formed on both sides of the interface. These clusters include hydrophobic residues (Gly 148

and Val 160) and polar residues (Arg 67, Arg 159, His 164, Asp 168, and Glu 243) (**Fig. 3c**),

typical of intersubunit interactions in stable homodimeric proteins [14]. These observations suggest

that σNS dimers with protruding N-terminal arms are the building blocks of higher-order

structures.

σNS dimers form a helical assembly by domain-swapping N-terminal arm interactions

The packing of the Se-Met σNS-R6A dimeric subunits in the crystals shows a helical

organization with a ~ 40 Å central cavity formed by domain-swapping interactions between the

N-terminal arms of the neighboring dimers (**Fig. 4a-d**). The overall diameter of the helical

assembly is ~ 150 Å. Interactions of σNS to form helical assemblies are consistent with previous

observations using cryo-EM, indicating that σNS forms filamentous structures in the presence of

RNA [12]. Such helical assemblies were not expected for the σNS-R6A mutant, as it does not bind

RNA and forms only dimers in solution. However, the size-exclusion chromatography profile of

Se-Met-substituted σNS-R6A clearly showed that it forms higher-order oligomers in solution as

observed with WT σNS (**Fig. 1b**). It is possible that increased hydrophobicity due to substitution

of a selenium atom for a sulfur atom[15] in Met 1 coupled with crystal-packing forces restrict the

flexibility of the N-terminal arms and allow this region of the protein to stabilize a multimeric

association.

Bile acid derivatives disrupt σNS filament formation

We observed that the native σNS-R6A crystals, without Se-Met substitution, crystallized in the

P4$_1$ space group instead of P6$_5$. Interestingly, native σNS-R6A crystallized only in conditions

with bile acid derivatives in contrast to Se-Met σNS-R6A, which crystallized without these

additives. Structure determination of native σNS-R6A, using the Se-Met σNS-R6A structure as a

molecular replacement model, showed that native σNS-R6A also forms an antiparallel dimer,

with two bile acid moieties bound to each subunit in a groove located near the σNS C-terminus

(**Fig. 5a**, sites 1 and 2). Binding of bile acid molecules is mediated by hydrophobic contacts with

residues Thr 107, Glu 110, Leu 111, Ser 114, Gly 203, Leu 204, Tyr 240, Glu 242, Ala 245, and

Glu 246 (**Fig. 5b**). Remarkably, in contrast to the Se-Met σNS-R6A structure, the N-terminal

arm is disordered and not observed in the native σNS-R6A structure. Instead of the distinct

helical assembly in the structure of Se-Met σNS-R6A, native σNS-R6A dimers crystallized with

bile acid show sheet-like packing (**Fig. 5c).** Further examination of the structure showed that one

of the bile acid molecules strikingly occupies the same location in the core domain as the N-

terminal arm required to form helical assemblies, suggesting that bile acid competes with the N-

terminal arm to disrupt the helical assembly formation observed in the Se-Met σNS-R6A

structure (**Fig. 5d-f**).

The σNS N-terminal arm is required for multimeric association

To provide additional evidence for the role of domain-swapping interactions of the N-terminal

arms in forming oligomeric assemblies, we engineered a deletion construct, σNS-ΔN17, by

truncating the N-terminal 17 residues of the protein. The σNS-ΔN17 mutant formed dimers in

solution and, interestingly, it also crystallized in the presence of bile acid derivatives. The σNS-

ΔN17 crystal structure determined using molecular replacement showed a dimer formed by a

central core identical to that in the structures of Se-Met or native σNS-R6A (**Fig. 5g**) with two

bound bile acid moieties. As expected, no helical packing was observed in the crystal structure.

Therefore, the structure of σNS-ΔN17 suggests that the N-terminal arm and the core of WT σNS

operate as independent structural units, and removing the N-terminal arm does not alter the

dimeric association of the core. Collectively, the three crystal structures provide strong evidence

that (i) WT σNS has a tendency to form dimers, (ii) the dimer is the basic unit of multimeric

assembly, and (iii) domain-swapping interactions of the flexible N-terminal arms are required to

link the dimeric units to form a multimer. Furthermore, the σNS-ΔN17 structure showed that bile

acid derivatives bind to the same location in σNS as the N-terminal arms in the σNS-R6A

structure (**Fig. 5g**), suggesting that this location in the σNS core domain has intrinsic affinity for

cholesterol-like molecules and that binding to such molecules disrupts filamentous helical

assemblies by competing with the binding of the N-terminal arms.

σNS oligomer exhibits RNA chaperone activity

The surface representation of the helical assembly in the Se-Met σNS-R6A structure shows a

negatively charged exterior (**Fig. 6a**) and a positively charged interior (**Fig. 6b**). A cutaway view

of the central tunnel of the helical assembly reveals a substantial number of positively charged

residues, suggesting that this region can bind RNA by electrostatic interactions. *In silico* docking

experiments demonstrated that the central tunnel can accommodate an RNA strand surrounded

by σNS dimers (**Fig. 6c-e**). This finding, coupled with previous studies showing that avian

reovirus σNS displays RNA-chaperone activity[16], prompted us to hypothesize that mammalian

orthoreovirus σNS is likewise an RNA chaperone. To test this hypothesis, we first examined

whether purified WT σNS and σNS-R6A disrupt dsRNA by incubating each of these σNS

proteins with an RNA-RNA duplex with an overhang of 6 nucleotides on either side. After

incubation, the reaction mixtures were treated with proteinase K and resolved by electrophoresis

in nondenaturing polyacrylamide gels. The RNA-RNA duplex migrated as a single prominent

band when σNS was not added to the reaction mixtures (**Fig. 6f**). However, following addition of

increasing concentrations of WT σNS, we observed a proportional increase in the formation of

ssRNA molecules. As an important control, formation of ssRNA was not observed in a similar

experiment using σNS-R6A (**Fig. 6g**). These results indicate that σNS catalyzes formation of

ssRNA molecules from dsRNA and that σNS displays RNA-destabilizing activity. We next

evaluated whether σNS is capable of annealing two ssRNA molecules to form dsRNA. In this

experiment, we observed a proportional increase in the formation of dsRNA molecules following

addition of increasing concentrations of WT σNS to the reaction mixtures (**Fig. 6h**), indicating

that σNS catalyzed formation of dsRNA molecules from ssRNA and thus possesses RNA-

annealing activity. Since the reaction buffer for these experiments contained no ATP, the RNA

chaperone activity of σNS is independent of energy input. These experiments demonstrate that

σNS is not a helicase but an RNA chaperone.

Binding of bile acid inhibits σNS chaperone activity

From our observation that bile acid binding disrupts formation of filamentous helical assemblies

of σNS, we hypothesized that bile acid binding would impede σNS RNA chaperone activity. To

test this hypothesis, we conducted RNA chaperone assays following incubation of σNS with

different components of bile acid derivatives, including taurocholic acid sodium salt hydrate

(TASAH), sodium glycocholate hydrate (SGAH), 3-[(3-cholamidopropyl)dimethylammonio]-1-

propanesulfonate (CHAPS), and 3-[(3-cholamidopropyl)dimethylammonio]-2-hydroxy-1-

propanesulfonate (CHAPSO). Each of these components reduced the RNA-destabilizing activity

of σNS to varying degrees (**Fig. 6h**), suggesting that the RNA chaperone activity of σNS requires

formation of filament-like assemblies. As anticipated, σNS RNA-binding capacity and oligomer

formation were abolished by removal of the N-terminal 17 residues and, concomitantly, the

σNS-ΔN17 mutant did not display RNA chaperone activity (**Fig. 6i**).

Discussion

VFs serve an essential function in the replication of many viruses, as these structures house viral

genome synthesis and assembly of progeny virions. In the case of mammalian orthoreoviruses,

two nonstructural proteins, σNS and μNS, initiate VF formation, and σNS recruits viral RNAs

into these compartments. Our crystallographic and biochemical studies reported here provide a

mechanistic understanding of how σNS forms filamentous assemblies by linking dimeric units

through domain-swapping interactions of the N-terminal arms and further how such a multimeric

association is required for the RNA-helix destabilizing and annealing activities of σNS. These

findings implicate σNS in unwinding viral RNA for loading into the viral RNA-dependent RNA

polymerase during genome replication and promoting interactions between the viral RNAs to

facilitate their selective encapsidation during virion assembly.

Higher-order oligomers and a flexible element–a common theme in VF formation?

Although distinct in its structure, σNS displays several functional and mechanistic similarities

with other dsRNA virus nonstructural proteins, such as rotavirus NSP2 and rice black-streaked

dwarf virus (RBSDV) P9-1. Like σNS, both NSP2 and P9-1 are implicated in formation of VFs.

The striking common structural feature of these proteins is the formation of oligomeric

assemblies requiring a flexible structural element. However, they differ in detail. Unlike the

formation of filamentous assemblies by σNS, rotavirus NSP2 forms a donut-shaped octamer [17,18].

Instead of the flexible N-terminal arm of σNS, which is responsible for dimer-dimer association

in filamentous assembly formation, NSP2 has a flexible C-terminal arm, which facilitates inter-

octamer association through domain-swapping interactions with neighboring NSP2 molecules [17].

Formation of higher-order NSP2 octamers is thought to be the basis for VF formation during

rotavirus replication [17,19]. RBSDV P9-1 also forms an octamer using a flexible C-terminal arm[20-

[22]. Like rotavirus NSP2, higher-order P9-1 octamers are required for formation of VFs, as

deletion of this sequence disrupts RBSDV VFs [22]. From our studies, deletion of the N-terminal

arm of σNS abrogates dimer-dimer interactions and precludes formation of higher-order

multimers. However, further experiments are required to assess the function of the flexible N-

terminal arm in the formation of functional VFs.

RNA chaperone activity–a flexible N-terminal arm

An important finding from our studies is that mammalian orthoreovirus σNS exhibits helix-

destabilizing and RNA-annealing activities without requiring a metal ion or ATP, which makes

this protein a *bona fide* RNA chaperone [24]. Similar RNA chaperone activity has been reported for

rotavirus NSP2 [25] and avian reovirus σNS [16]. The RNA duplex used in our experiments contains

overhangs at the ends of both strands, suggesting that passive strand displacement activity is

initiated after σNS binds to unduplexed overhang regions. The binding of σNS would then

prevent reannealing of the duplex and further separate the two strands. Using complementary

single-stranded RNAs, our studies show that σNS binds, unfolds, and releases these molecules to

facilitate formation of duplex RNA.

The RNA chaperone activity of σNS requires the flexible N-terminal arm with an

arginine at residue 6 to function in domain-swapping interactions, as the R6A substitution or

deletion of the N-terminal arm curtails this activity. In an analogous scenario, deletion of the

flexible C-terminal arm of NSP2 also significantly reduces RNA chaperone activity [26]. Thus, the

requirement for a flexible region in these proteins suggests that the mechanism is likely mediated

by disorder or entropy transfer. One possibility is that σNS uses the flexibility of the N-terminal

arm for binding RNA and subsequently releases the bound RNA. Such a possibility is consistent

with our observation that a single residue, Arg 6, in the N-terminal arm influences the RNA-

binding capacity of σNS [8]. The free N-terminal arms of the σNS multimer subunits (**Fig. 5e**)

likely exist in a dynamic equilibrium between a freely exposed open state, in which Arg 6 can

initiate binding to RNA, and a closed state, in which the N-terminal arm is engaged in domain-

swapping interactions for RNA release. Such dynamics also may allow the N-terminal arm to

recruit free σNS dimers to elongate the multimer and coat the RNA as a mechanism for

unwinding (**Fig. 7**). The dynamics and preference between open and closed states are likely

influenced by the presence of RNA and buffer conditions (pH and ionic strength). This idea is

consistent with our observation that in solution σNS exists as an octamer in the absence of RNA,

but in the presence of RNA, as observed in our previous cryo-EM images [12], σNS forms

filamentous structures of varying lengths.

Bile acid binding–tampering with the flexibility of the N-terminal arm

A serendipitous but fascinating finding from our study is that bile acid derivatives bind to the

same site in σNS as the N-terminal arm and impede the RNA chaperone activity of σNS (**Fig.**

5h). These findings are consistent with the proposed mechanism of σNS RNA chaperone

activity. First, the σNS crystal structure shows that the free N-terminal arm displaced by bile acid

is disordered, as the electron density for this region is absent in the presence of bile acid. This

observation is consistent with the disorder or entropy-transfer model of RNA chaperone activity

[27], in which entropy is transferred to bound RNA to facilitate RNA refolding. Second, our

finding that bile acid binding does not entirely abrogate RNA chaperone activity is consistent

with the idea that RNA modulates the kinetics of open and closed states of the N-terminal σNS

arm. It is possible that differences in the efficiency by which bile salts diminish RNA chaperone

activity are determined by differences in the capacity of these molecules to compete with the N-

terminal arm in the presence of RNA.

Role of RNA chaperone activity in reovirus replication

Our discovery that σNS displays RNA chaperone activity raises an important question about the

stage in viral replication at which this activity is required. While the function of σNS RNA-

binding activity in the recruitment of viral RNAs to VFs has been established [8], further studies

are needed to unequivocally determine the precise function of σNS RNA chaperone activity. As

proposed for RNA chaperones of single-stranded RNA viruses [27], σNS RNA chaperone activity

may be necessary for unfolding viral mRNA for translation, replication, packaging, or some

combination of these steps. In the case of dsRNA viruses like reoviruses and rotaviruses that

package multiple dsRNA segments, RNA chaperone activity is likely essential for proper

refolding of the kinetically trapped intermediate structures in capped viral mRNAs to allow these

RNAs to serve as templates for dsRNA synthesis by the viral polymerase and also for facilitating

interactions between specific viral RNAs for gene segment assortment and packaging.

Methods

Expression and purification of σNS constructs

Synthesized genes for WT σNS, σNS-R6A, and σNS-ΔN17 were subcloned into the bacterial

expression vector pET28 with an N-terminal His tag and a TEV protease cleavage site (Epoch

Life Science). *Escherichia coli* DE3 cells (Novagen) were transformed with the pET28 plasmid

and induced with 0.5 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) (Millipore Sigma)

when the optical density at 600 nm reached 0.6. *E. coli* B834 cells were incubated in Dream

medium supplemented to contain Dream Nutrient Mix and 10 mg/mL Se-Met. Cells were

resuspended in 50 mM Tris-HCl (pH 8.0), 300 mM NaCl, and 10 mM imidazole and

supplemented to contain a protease inhibitor cocktail (Roche). Cells were lysed using a

microfluidizer (Microfluidics), followed by removal of cell debris by centrifugation at 39,000 x g

at 4°C for 30 minutes. His-tagged σNS was loaded onto a Ni-NTA agarose column (Qiagen) and

eluted using a gradient of 50 mM Tris-HCl (pH 8.0), 300 mM NaCl, and 250 mM imidazole. The

eluted His-tagged σNS was concentrated using a 10-kDa centrifugal filter unit (Millipore),

dialyzed into 50 mM Tris-HCl (pH 8.0), 300 mM NaCl, and 10 mM imidazole, and incubated

with TEV protease at 4°C overnight. The cleaved protein mixture was reloaded onto a Ni-NTA

agarose column to remove the His-TEV protease and uncleaved His-σNS. Cleaved σNS was

subjected to ion-exchange chromatography using a Q-column (GE Healthcare) and eluted with

20 mM Tris-HCl (pH 8.0) and 1 M NaCl. Eluted σNS was separated using an S200 16/60-

increase column in 10 mM HEPES (pH 8.0) and 150 mM NaCl.

Crystallization, data processing, structure determination, and refinement

Freshly purified Se-Met σNS-R6A, native σNS-R6A, and σNS-ΔN17 were used for initial

crystallization screens immediately following size-exclusion chromatography. In each case,

fractions with σNS proteins were collected and concentrated to ~ 10 mg/mL in buffer containing

10 mM HEPES (pH 8.0) and 150 mM NaCl for crystallization. Crystals were propagated at 20°C

by hanging-drop vapor diffusion using a Mosquito crystallization robot (TTP LabTech) and

imaged using a Rock Imager (Formulatrix). Each drop contained 0.2 μL of the σNS protein and

0.2 μL of crystallization buffer. Se-Met σNS-R6A produced crystals in the condition containing

0.1 M MES monohydrate (pH 6.5) and 1.6 M magnesium sulfate heptahydrate (Hampton

Research), whereas native σNS-R6A and σNS-ΔN17 crystallized in the condition with 1.2 %

cholic acid derivatives mix, 0.1 M Buffer System 3 (pH 8.5), and 30 % Precipitant Mix 3

(Morpheus III, Molecular Dimensions). Crystals were transferred into a cryoprotectant solution

with 20% glycerol and flash-frozen in liquid nitrogen. X-ray diffraction data for σNS were

collected on beamline 5.0.1 at the Advanced Light Source at the Lawrence Berkeley National

Laboratory. Diffraction data were processed using the CCP4 software suite [28]. For Se-Met σNS-

R6A, phases and initial electron density maps were calculated by single-wavelength anomalous

diffraction (SAD) phasing using SHELX [29], and automated model building with experimental

phases was conducted using ARP/wARP [30] before iterative cycles of manual model building and

refinement using COOT[31] and PHENIX [32]. Further model building was accomplished using

COOT based on the difference maps [31] (see validation report for PDB id: 8TKA). The structures

of native σNS-R6A and σNS-ΔN17 were determined using molecular replacement with PHENIX

[32]. The bile acid binding in these structures was validated by omit maps (see validation reports

for PDB id: 8TL8 and 8TL1). Data collection and refinement statistics following the final

refinement cycle are provided in Supplementary Table 1. Interactions between σNS and colic

acid derivatives were analyzed using LigPlot+ v.2.1 [33]. Figures were prepared using Chimera [34].

RNA helix destabilizing assay

RNA helix destabilizing assays were conducted as described [35] with modifications. Various

concentrations of σNS, σNS-R6A, and σNS-ΔN17 (2.5 to 20 μM) and 0.5 μM dsRNA substrate

were incubated at 37°C for 1 h with 25 mM HEPES-KOH (pH 8.0), 100 mM NaCl, 2 mM

$MgCl_2$, 2 mM DTT, and 5 U RNase-OUT in a reaction volume of 20 μL. Reactions were

terminated by the addition of 5 U proteinase K for 15 min and 2.5 μL 5 × loading buffer (100

mM Tris–HCl [pH 7.5], 50% glycerol, and bromophenol blue) and electrophoresed in 4-20%

native polyacrylamide gels. Gels were scanned using a Typhoon 9200 imager (GE Healthcare).

The effect of bile acids on σNS RNA helix destabilizing activity was determined by incubating

20 and 40 μM of different components of bile acid derivatives, including TASAH, SGAH,

CHAPS, and CHAPSO, with 20 μM σNS for 1 h prior to initiation of the helix destabilization

assay.

RNA strand hybridization assay

Various concentrations of σNS (2.5 to 20 μM) were incubated with HEX-labeled and unlabeled

RNA strands (0.5 μM each strand) at 37°C for 1 h in 50 mM HEPES-KOH (pH 8.0), 2.5 mM

$MgCl_2$, 2 mM DTT, 0.01% BSA, and 5 U RNase-OUT in a reaction volume of 20 μl. Reactions

were terminated as described for RNA helix destabilizing assays. Two complementary stem-

loop-structured RNA strands were used for hybridization assays [35]. Reactions were resolved in 4-

20% native polyacrylamide gels and scanned using a Typhoon 9200 imager.

Acknowledgements

We are grateful to members of the Dermody and Prasad laboratories for many useful discussions.

This work was supported by U.S. Public Health Service award R01 AI032539, the Heinz

Endowments (T.S.D.), and grant Q1279 from Robert Welch Foundation (B.V.V.P). We

acknowledge the Advanced Light Source (8.2.2) (Berkeley, CA) for X-ray data collection. The

ALS-ENABLE beamlines are supported in part by U.S. Public Health Service award P30

GM124169. The Advanced Light Source is a Department of Energy Office of Science User

Facility under Contract No. DE-AC02-05CH11231. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

412 B.Z. and L.H. conceived, designed, and conducted experiments, analyzed data, contributed

413 materials and analytic tools, and drafted the paper. S.K., N.N., C.H.L., X.S., and G.M.T.

414 conceived and designed experiments, analyzed data, and contributed materials and analytic tools.

415 T.S.D. and B.V.V.P. conceived and designed experiments, analyzed the data, and drafted the

416 paper. All authors reviewed, critiqued, and provided comments on the manuscript.

Competing interests

419 The authors declare no competing interests.

Additional information

Supplementary information Supplementary Table 1.

Correspondence and requests for materials should be addressed to Terence S. Dermody or B. V.

Venkataram Prasad.

Figure Legends

Fig. 1 Purification of σNS-R6A dimer. Elution profiles of WT σNS and σNS-R6A after **a**

anion-exchange chromatography and **b** size-exclusion chromatography. A size shift is observed

with the different elution volumes for each peak.

Fig. 2 Crystal structure of σNS contains a flexible N-terminal arm. a Ribbon and surface

representations of the σNS crystal structure. The N-terminal arm is demarcated in a grey-dotted

frame. **b** Topologic representation of the σNS crystal structure generated by the PDBsum

database. Strands are colored in green with a light blue background, helices are colored in cyan,

and the connecting loops are colored in grey. **c** The RMSF graph of the σNS dimer and the σNS

dimer in complex with an RNA molecule after 100 nanoseconds of molecular dynamics

simulations.

Fig. 3 σNS dimer is the basic functional unit. a Ribbon (left) and surface representations

(right) of the σNS dimer with subunits colored in green and blue. The N-terminal arms are

demarcated in grey-dotted frames. **b** Ribbon representation of the σNS dimer highlighting

residues that mediate interactions at the dimer interface inside a black frame in blue and green. **c**

Amino acid residues from the subunits at the dimer interface are colored blue and green.

Hydrogen bond interactions are shown by red lines.

Fig. 4 Formation of σNS filaments. a Surface representation of the helical assembly formed by

σNS dimers using crystallographic P6_5 symmetry. Dimers are depicted as blue and green

subunits. A single helical turn (80 Å in length) is demarcated in the purple-dotted frame. **b** A 90°

rotation around the horizontal axis perpendicular to the filament shows the central tunnel. The

tunnel is 40 Å in width, and the total diameter of the filament is 150 Å. **c** Single helical turn

formed by six σNS dimers. Dimers are depicted in different colors. One pair of interacting

dimers (pink and cyan) are demarcated in a yellow frame. The location of the N-terminal arm of

the pink dimer is shown within the red frame, with a red arrow pointing to the (**e**) Ligplot of

interactions between N-terminal arm residues of the pink dimer and core domain residues of the

blue dimer. Residues involved in hydrophobic interactions are shown by brick-red spoked arcs,

and hydrogen bond interactions between the residues are indicated by green lines, with carbon,

oxygen, and nitrogen atoms depicted in black, red, and blue, respectively. Note the involvement

of the Se-Met residue at position 1 (Mse 1), indicated by an asterisk. **d** Interacting dimers (pink

and blue) corresponding to the yellow frame in **c** are shown separately to illustrate how the N-

terminal arms (in red frame) project away in opposite directions (thick red arrows) to chain-link

the neighboring dimers to form the helical assembly shown in **Fig. 1a.**

Fig. 5 Binding of bile acid derivatives disrupts σNS oligomerization and RNA-destabilizing

activity. a Electrostatic representation of σNS-R6A in complex with bile acid derivatives (S1

and S2), colored in green. The grey-dotted box demarcates the missing disordered N-terminal

arm. **b, c** Residues mediating bile-acid-derivative binding at S1 and S2 defined by LigPlot

following the same notations as in **Fig. 4e** for hydrophobic and hydrogen bond interactions. Bile

acid derivatives are shown in purple bonds with carbon and oxygen atoms in black and red,

respectively. **c** The higher-order structure formed by crystal packing of σNS-R6A when bile acid

derivatives are present. **d** The helical assembly as observed in the crystal structure of Se-Met

σNS-R6A with dimeric subunits shown in pink and blue. **e** Close-up view of the N-terminal arm

(pink) interacting with a groove in the neighboring subunit (blue). **f** In the structure of σNS-R6A

in complex with bile acid derivatives, the same groove is occupied by a bile acid replacing the N-

terminal arm, which becomes disordered. **g** Electrostatic representations σNS-ΔN17 in complex

with two bile acid derivatives (in green) bound at the same location as in the σNS-R6A structure

shown in **Fig. 5a**. The grey-dotted box demarcates the missing N-terminal arm.

Fig. 6 RNA chaperone activity of σNS. a Electrostatic representation of a σNS filament with

red indicating negatively charged residues and blue indicating positively charged residues. The

exterior of the filament is negatively charged. **b** Cutaway view of the filament. The interior of the

filament is positively charged. **c** A single dimer is demarcated in a black-dotted frame with its

relative position within the filament. **d** A 90-degree rotation around the z-axis from the previous

position shows a side-view of the dimer. A docked RNA molecule is colored in yellow. **e** A 90-

degree rotation around the z-axis from the previous position shows the positively charged pocket

of σNS that binds to a docked RNA molecule colored in yellow. The black-dotted frame depicts

the position of the dimer within the filament. **f** RNA-helix destabilizing activity of WT σNS.

Boiling of the duplex disrupts duplex formation. Increasing concentrations of WT σNS yield

more ssRNA products. Maltose-binding protein (MBP) and norovirus (NV) P41 serve as

negative and positive controls, respectively. **g** RNA-helix destabilizing activity of σNS-R6A.

Increasing concentrations of σNS-R6A yield only limited ssRNA products. **h** RNA-annealing

activity of WT σNS. Boiling two ssRNA molecules promotes strand annealing. Similar levels of

annealing activity are observed following incubation with the highest concentration of WT σNS.

i Effects of each component of the bile acid derivatives on the RNA-destabilizing activity of

σNS. TASAH, taurocholic acid sodium salt hydrate. SGAH, sodium glycocholate hydrate.

CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate. CHAPSO, 3-[(3-

cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate. **j** RNA-helix destabilizing

activity of σNSΔ17. Increasing concentrations of σNS Δ17 does not yield significant ssRNA

products.

Fig. 7 Model of σNS oligomerization, RNA binding, and RNA unfolding. The free N-

terminal arms of σNS oligomers exist in equilibrium between an open state that can initiate RNA

binding and a closed state that engages in domain-swapping interactions and RNA release from

the N-terminal arm. The dynamics of the N-terminal arms allow for recruitment of free σNS

dimers to elongating σNS oligomers to coat the RNA and enhance RNA unfolding.

References

1   Day, J. M. The diversity of the orthoreoviruses: molecular taxonomy and phylogentic
        divides. *Infection, genetics and evolution : journal of molecular epidemiology and
        evolutionary genetics in infectious diseases* **9**, 390-400 (2009).
        https://doi.org:10.1016/j.meegid.2009.01.011

2   Ouattara, L. A. *et al.* Novel human reovirus isolated from children with acute necrotizing
        encephalopathy. *Emerging infectious diseases* **17**, 1436-1444 (2011).
        https://doi.org:10.3201/eid1708.101528

3   Bouziat, R. *et al.* Reovirus infection triggers inflammatory responses to dietary antigens
        and development of celiac disease. *Science* **356**, 44-50 (2017).
        https://doi.org:10.1126/science.aah5298

4   DeAntoneo, C., Danthi, P. & Balachandran, S. Reovirus Activated Cell Death Pathways.
        Cells **11** (2022). https://doi.org:10.3390/cells11111757

5   Muller, L., Berkeley, R., Barr, T., Ilett, E. & Errington-Mais, F. Past, Present and Future
        of Oncolytic Reovirus. *Cancers (Basel)* **12** (2020).
        https://doi.org:10.3390/cancers12113219

6   Maginnis, M. S. *et al.* Beta1 integrin mediates internalization of mammalian reovirus. *J*
Virol **80**, 2760-2770 (2006). https://doi.org:10.1128/JVI.80.6.2760-2770.2006

7   Danthi, P. *et al.* From touchdown to transcription: the reovirus cell entry pathway.
Current topics in microbiology and immunology **343**, 91-119 (2010).
https://doi.org:10.1007/82_2010_32

8   Lee, C. H. *et al.* Reovirus Nonstructural Protein sigmaNS Recruits Viral RNA to
Replication Organelles. *mBio* **12**, e0140821 (2021). https://doi.org:10.1128/mBio.01408-
21

9   Desmet, E. A., Anguish, L. J. & Parker, J. S. Virus-mediated compartmentalization of the
host translational machinery. *mBio* **5**, e01463-01414 (2014).
https://doi.org:10.1128/mBio.01463-14

10   Choudhury, P., Bussiere, L. D. & Miller, C. L. Mammalian Orthoreovirus Factories
Modulate Stress Granule Protein Localization by Interaction with G3BP1. *J Virol* **91**
(2017). https://doi.org:10.1128/JVI.01298-17

11   Stamatos, N. M. & Gomatos, P. J. Binding to selected regions of reovirus mRNAs by a
nonstructural reovirus protein. *Proceedings of the National Academy of Sciences of the*
United States of America **79**, 3457-3461 (1982).

12   Zamora, P. F. *et al.* Reovirus Nonstructural Protein sigmaNS Acts as an RNA Stability
Factor Promoting Viral Genome Replication. *J Virol* **92** (2018).
https://doi.org:10.1128/JVI.00563-18

13   Goral, M. I., Mochow-Grundy, M. & Dermody, T. S. Sequence diversity within the
reovirus S3 gene: reoviruses evolve independently of host species, geographic locale, and
date of isolation. *Virology* **216**, 265-271 (1996). https://doi.org:10.1006/viro.1996.0059

14   Larsen, T. A., Olson, A. J. & Goodsell, D. S. Morphology of protein-protein interfaces.
Structure **6**, 421-427 (1998). https://doi.org:10.1016/s0969-2126(98)00044-6

15   Martinez, J. *et al.* Selenomethionine incorporation into amyloid sequences regulates
fibrillogenesis and toxicity. *PLoS One* **6**, e27999 (2011).
https://doi.org:10.1371/journal.pone.0027999

16   Borodavka, A., Ault, J., Stockley, P. G. & Tuma, R. Evidence that avian reovirus
sigmaNS is an RNA chaperone: implications for genome segment assortment. *Nucleic*
Acids Res **43**, 7044-7057 (2015). https://doi.org:10.1093/nar/gkv639

17   Hu, L. *et al.* Crystallographic analysis of rotavirus NSP2-RNA complex reveals specific
recognition of 5' GG sequence for RTPase activity. *J Virol* **86**, 10547-10557 (2012).
https://doi.org:10.1128/JVI.01201-12

Jayaram, H., Taraporewala, Z., Patton, J. T. & Prasad, B. V. Rotavirus protein involved in genome replication and packaging exhibits a HIT-like fold. *Nature* **417**, 311-315 (2002). https://doi.org:10.1038/417311a

Criglar, J. M. *et al.* Phosphorylation cascade regulates the formation and maturation of rotaviral replication factories. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E12015-E12023 (2018). https://doi.org:10.1073/pnas.1717944115

Akita, F. *et al.* Crystallographic analysis reveals octamerization of viroplasm matrix protein P9-1 of Rice black streaked dwarf virus. *J Virol* **86**, 746-756 (2012). https://doi.org:10.1128/JVI.00826-11

Li, X. *et al.* Characterization of the importance of terminal residues for southern rice black-streaked dwarf virus P9-1 viroplasm formations. *Protein Expr Purif* **111**, 98-104 (2015). https://doi.org:10.1016/j.pep.2015.04.003

Wu, J. *et al.* Viroplasm protein P9-1 of Rice black-streaked dwarf virus preferentially binds to single-stranded RNA in its octamer form, and the central interior structure formed by this octamer constitutes the major RNA binding site. *J Virol* **87**, 12885-12899 (2013). https://doi.org:10.1128/JVI.02264-13

Parker, J. S., Broering, T. J., Kim, J., Higgins, D. E. & Nibert, M. L. Reovirus core protein mu2 determines the filamentous morphology of viral inclusion bodies by interacting with and stabilizing microtubules. *J Virol* **76**, 4483-4496 (2002). https://doi.org:10.1128/jvi.76.9.4483-4496.2002

Semrad, K. Proteins with RNA chaperone activity: a world of diverse proteins with a common task-impediment of RNA misfolding. *Biochem Res Int* **2011**, 532908 (2011). https://doi.org:10.1155/2011/532908

Bravo, J. P. K. *et al.* Structural basis of rotavirus RNA chaperone displacement and RNA annealing. *Proceedings of the National Academy of Sciences of the United States of America* **118** (2021). https://doi.org:10.1073/pnas.2100198118

Coria, A. *et al.* Rotavirus RNA chaperone mediates global transcriptome-wide increase in RNA backbone flexibility. *Nucleic Acids Res* **50**, 10078-10092 (2022). https://doi.org:10.1093/nar/gkac738

Zuniga, S., Sola, I., Cruz, J. L. & Enjuanes, L. Role of RNA chaperones in virus replication. *Virus Res* **139**, 253-266 (2009). https://doi.org:10.1016/j.virusres.2008.06.015

Agirre, J. *et al.* The CCP4 suite: integrative software for macromolecular crystallography. *Acta Crystallogr D Struct Biol* **79**, 449-461 (2023). https://doi.org:10.1107/S2059798323003595

29 Uson, I. & Sheldrick, G. M. An introduction to experimental phasing of macromolecules
illustrated by SHELX; new autotracing features. *Acta Crystallogr D Struct Biol* **74**, 106-
116 (2018). https://doi.org:10.1107/S2059798317015121

30 Langer, G., Cohen, S. X., Lamzin, V. S. & Perrakis, A. Automated macromolecular
model building for X-ray crystallography using ARP/wARP version 7. *Nat Protoc* **3**,
1171-1179 (2008). https://doi.org:10.1038/nprot.2008.91

31 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501 (2010).
https://doi.org:10.1107/S0907444910007493

32 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
https://doi.org:10.1107/S0907444909052925

33 Laskowski, R. A. & Swindells, M. B. LigPlot+: multiple ligand-protein interaction
diagrams for drug discovery. *J Chem Inf Model* **51**, 2778-2786 (2011).
https://doi.org:10.1021/ci200227u

34 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612 (2004). https://doi.org:10.1002/jcc.20084

35 Yang, J. *et al.* A cypovirus VP5 displays the RNA chaperone-like activity that
destabilizes RNA helices and accelerates strand annealing. *Nucleic Acids Res* **42**, 2538-
2554 (2014). https://doi.org:10.1093/nar/gkt1256

Fig. 1 Purification of σNS-R6A dimer. Elution profiles of WT σNS and σNS-R6A after **a** anion-exchange chromatography and **b** size-exclusion chromatography. A size shift is observed with the different elution volumes for each peak.

Fig. 2 Crystal structure of σNS contains a flexible N-terminal arm. a Ribbon and surface representations of the σNS crystal structure. The N-terminal arm is demarcated in a grey-dotted frame. **b** Topologic representation of the σNS crystal structure generated by the PDBsum database. Strands are colored in green with a light blue background, helices are colored in cyan, and the connecting loops are colored in grey. **c** The RMSF graph of the σNS dimer and the σNS dimer in complex with an RNA molecule after 100 nanoseconds of molecular dynamics simulations.

Fig. 3 σNS dimer is the basic functional unit. a Ribbon (left) and surface representations (right) of the σNS dimer with subunits colored in green and blue. The N-terminal arms are demarcated in grey-dotted frames. **b** Ribbon representation of the σNS dimer highlighting residues that mediate interactions at the dimer interface inside a black frame in blue and green. **c** Amino acid residues from the subunits at the dimer interface are colored blue and green. Hydrogen bond interactions are shown by red lines.

Fig. 4 Formation of σNS filaments. a Surface representation of the helical assembly formed by σNS dimers using crystallographic P6₅ symmetry. Dimers are depicted as blue and green subunits. A single helical turn (80 Å in length) is demarcated in the purple-dotted frame. **b** A 90° rotation around the horizontal axis perpendicular to the filament shows the central tunnel. The tunnel is 40 Å in width, and the total diameter of the filament is 150 Å. **c** Single helical turn formed by six σNS dimers. Dimers are depicted in different colors. One pair of interacting dimers (pink and cyan) are demarcated in a yellow frame. The location of the N-terminal arm of the pink dimer is shown within the red frame, with a red arrow pointing to the (**e**) Ligplot of interactions between N-terminal arm residues of the pink dimer and core domain residues of the blue dimer. Residues involved in hydrophobic interactions are shown by brick-red spoked arcs, and hydrogen bond interactions between the residues are indicated by green lines, with carbon, oxygen, and nitrogen atoms depicted in black, red, and blue, respectively. Note the involvement of the Se-Met residue at position 1 (Mse 1), indicated by an asterisk. **d** Interacting dimers (pink and blue) corresponding to the yellow frame in **c** are shown separately to illustrate how the N-terminal arms (in red frame) project away in opposite directions (thick red arrows) to chain-link the neighboring dimers to form the helical assembly shown in **Fig. 1a.**

Fig. 5 Binding of bile acid derivatives disrupts σNS oligomerization and RNA-destabilizing activity. a Electrostatic representation of σNS-R6A in complex with bile acid derivatives (S1 and S2), colored in green. The grey-dotted box demarcates the missing disordered N-terminal arm. **b, c** Residues mediating bile-acid-derivative binding at S1 and S2 defined by LigPlot following the same notations as in **Fig. 4e** for hydrophobic and hydrogen bond interactions. Bile acid derivatives are shown in purple bonds with carbon and oxygen atoms in black and red, respectively. **c** The higher-order structure formed by crystal packing of σNS-R6A when bile acid derivatives are present. **d** The helical assembly as observed in the crystal structure of Se-Met σNS-R6A with dimeric subunits shown in pink and blue. **e** Close-up view of the N-terminal arm (pink) interacting with a groove in the neighboring subunit (blue). **f** In the structure of σNS-R6A in complex with bile acid derivatives, the same groove is occupied by a bile acid replacing the N-terminal arm, which becomes disordered. **g** Electrostatic representations σNS-ΔN17 in complex with two bile acid derivatives (in green) bound at the same location as in the σNS-R6A structure shown in **Fig. 5a**. The grey-dotted box demarcates the missing N-terminal arm.

Fig. 6 RNA chaperone activity of σNS. a Electrostatic representation of a σNS filament with red indicating negatively charged residues and blue indicating positively charged residues. The exterior of the filament is negatively charged. **b** Cutaway view of the filament. The interior of the filament is positively charged. **c** A single dimer is demarcated in a black-dotted frame with its relative position within the filament. **d** A 90-degree rotation around the z-axis from the previous position shows a side-view of the dimer. A docked RNA molecule is colored in yellow. **e** A 90-degree rotation around the z-axis from the previous position shows the positively charged pocket of σNS that binds to a docked RNA molecule colored in yellow. The black-dotted frame depicts the position of the dimer within the filament. **f** RNA-helix destabilizing activity of WT σNS. Boiling of the duplex disrupts duplex formation. Increasing concentrations of WT σNS yield more ssRNA products. Maltose-binding protein (MBP) and norovirus (NV) P41 serve as negative and positive controls, respectively. **g** RNA-helix destabilizing activity of σNS-R6A. Increasing concentrations of σNS-R6A yield only limited ssRNA products. **h** RNA-annealing activity of WT σNS. Boiling two ssRNA molecules promotes strand annealing. Similar levels of annealing activity are observed following incubation with the highest concentration of WT σNS. **i** Effects of each component of the bile acid derivatives on the RNA-destabilizing activity of σNS. TASAH, taurocholic acid sodium salt hydrate. SGAH, sodium glycocholate hydrate. CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate. CHAPSO, 3-[(3-cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate. **j** RNA-helix destabilizing activity of σNSΔ17. Increasing concentrations of σNS Δ17 does not yield significant ssRNA products.

Fig. 7 Model of σNS oligomerization, RNA binding, and RNA unfolding. The free N-terminal arms of σNS oligomers exist in equilibrium between an open state that can initiate RNA binding and a closed state that engages in domain-swapping interactions and RNA release from the N-terminal arm. The dynamics of the N-terminal arms allow for recruitment of free σNS dimers to elongating σNS oligomers to coat the RNA and enhance RNA unfolding.

Table 1: X-ray diffraction data and structure refinement statistics

PDB ID	8TKA	8TL8	8TL1
Wavelength (Å)	0.99	0.99	0.99
Resolution range (Å)	34.28 – 3.0 (3.10 - 3.0)	34.22 - 3.20 (3.31 -3.20)	34.52-3.16 (3.27 - 3.16)
Space group	P 6_5	P 4_1	C 1 2 1
Unit cell			
a, b, c (Å)	137.11, 137.11, 98.24	54.12, 54.12, 305.52	77.52, 76.80, 89.09
α, ββ, γγ (°)	90, 90, 120	90, 90, 90	90, 94.27, 90
Unique reflections	41331 (4152)	11801 (1150)	7841 (302)
Multiplicity	2.0 (2.0)	1.0 (1.0)	1.0 (1.0)
Completeness (%)	99.77 (99.67)	81.91 (82.38)	86.39 (33.82)
Mean I/σ(I)	21.07 (3.53)	18.04 (5.40)	10.09 (3.21)
Wilson B-factor (Å^2)	68.54	62.51	67.74
R-merge (%)	0.0693	0.0311	0.1319
R-work (%)	0.2446 (0.3333)	0.2240 (0.2944)	0.208 (0.2561)
R-free (%)	0.2937 (0.4157)	0.2434 (0.2995)	0.250 (0.2895)
Number of non-hydrogen atoms	5494	5337	2708
Macromolecules	5494	5205	2637
Ligands	0	132	66
Solvent	0	0	5
Protein residues	708	669	336
R.M.S. deviations			
Bond lengths (Å)	0.012	0.005	0.010
Bond angles (°)	1.60	1.06	1.36
Ramachandran			
Favored (%)	96.29	2.57	95.48
Allowed (%)	3.14	2.57	3.61
Disallowed (%)	0.57	0.00	0.90
Average B-factor (Å^2)	81.30	59.23	67.73
Macromolecules	81.30	58.98	67.10
Ligand		69.46	95.72
Solvent			30.00

Numbers in parentheses correspond to the highest resolution range.