Opsin gene expression plasticity and spectral sensitivity in male damselflies could mediate

female colour morph detection

Keywords: opsins, plasticity, female polymorphism, visual tuning, frequency-dependent

selection, spectral sensitivity

ABSTRACT

The visual systems of Odonata are characterized by many opsin genes, which form the

primary light-sensitive photopigments of the eye. Female-limited colour polymorphisms are

also common in Odonata, with one morph typically exhibiting male-like (androchrome)

colouration and one or two morphs exhibiting female-specific colouration (gynochromes).

These colour polymorphisms are thought to be maintained by frequency-dependent sexual

conflict, in which males form search images for certain morphs, causing disproportionate

mating harassment. Here, we investigate opsin sensitivity and gene expression plasticity in

mate-searching males of the damselfly *Ischnura elegans* during adult maturation and across

populations with different female morph frequencies. We find evidence for opsin-specific

plasticity in relative and proportional opsin mRNA expression, suggesting changes in opsin

regulation and visual sensitivity during sexual maturation. In particular, expression of the

long-wavelength sensitive opsin LWF2 changed over development and varied between

populations with different female morph frequencies. UV-Vis analyses indicate that short-

and long-wavelength opsins absorb wavelength of light between 350 and 650 nm. Assuming

opponency between photoreceptors with distinct short- and long-wavelength sensitivities,

these sensitivities suggest male spectral visual discrimination ability of andromorph and

gynomorph females. Overall, our results suggest that opsin sensitivity and expression

changes contribute to visual tuning that could impact conspecific discrimination.

INTRODUCTION

Opsins are G-protein coupled receptors that, when bound with a vitamin-A derived chromophore, form light sensitive visual pigments responsible for colour vision in animals (Terakita, 2005). Many insects possess three classes of opsin proteins: ultraviolet sensitive (UVS), short-wavelength sensitive (SWS), and long-wavelength sensitive (LWS) with maximum sensitivity to wavelengths (λ_{max}) around 350 nm, 440 nm, and 530 nm, respectively (Briscoe & Chittka, 2001; van der Kooi et al., 2021). Photoreceptor spectral sensitivity depends mainly on the absorption spectrum of expressed opsins, which can vary due to opsin gains or losses, structural changes, or expression level differences (Castiglione et al., 2023; Liénard et al., 2021; Sharkey et al., 2023). Spectral sensitivity can also be modified by screening and filtering mechanisms, which are relatively widespread in insects (van der Kooi et al., 2021). The visual system is also shaped by environmental factors, behaviours, and associated selection pressures. For example, loss of functional opsin genes correlates with nocturnal lifestyles in mammals (Jacobs, 2013), while opsin duplications in other species contribute to the regaining of spectral sensitivity (Rossetto et al., 2023; Sharkey et al., 2023). In several insect species, functional changes amongst duplicated opsins likely improves detection and/or discrimination of environmental and sexual signals (Liénard et al., 2021; McCulloch et al., 2017).

Opsin gene expression is typically not fixed, but changes plastically over individual ontogeny or in response to environmental factors and lightning conditions (Chang et al., 2021; Fuller et al., 2010; Shand et al., 2008). For example, changes in opsin expression have been associated with light exposure in honey bees and ants (Sasagawa et al., 2003; Yilmaz et al., 2016), and in "choosy" relative to "non-choosy" females of the butterfly *Bicyclus anynana* (Everett et al., 2012).

Damselflies and dragonflies (Insecta: Odonata) are colourful diurnal flying insects

that possess large conspicuous eyes comprising thousands of ommatidia (Fig. 1A). Odonates

inhabit a wide array of visual environments, having both aquatic and terrestrial life stages,

and they strongly rely on vision for a variety of behaviours including predator avoidance,

prey capture, and mate searching (Bybee et al., 2012; Corbet, 1999; Labhart & Nilsson,

1995). These insects are therefore excellent model organisms to study evolutionary and

functional aspects of colour vision. Like many other insects, odonates express UVS, SWS,

and LWS opsins in their eyes. However, duplications of the SWS and LWS opsin classes

have resulted in remarkable genetic diversity, with between 11 – 30 visual opsin genes

identified across several families (Futahashi et al., 2015). Electrophysiological recordings of

visual sensitivity in the eyes of some species indicate photoreceptor spectral curves ranging

from UV to red light (Bybee et al., 2012). While it is not clear how their abundant opsin

copies function in vision, visual stimuli are clearly important for odonate behaviour. For

example, comparisons of visual and olfactory cues suggested that vision is primarily used in

mate choice in several species of damselflies (Rebora et al., 2018; Winfrey & Fincke, 2017).

Colouration is also an important signal indicating female sexual maturity (Svensson et al.,

2020; Takahashi & Watanabe, 2011; Van Gossum et al., 2011; Willink et al., 2019). Males

can discriminate between immature and mature females, and they direct more mating

attempts to mature than immature females (Huang et al., 2014; Van Gossum et al., 2001a,

2011; Willink et al., 2019). Experimental manipulation of age-specific colouration in

Ischnura damselflies has further confirmed that vision is a primary discrimination cue

between immature and mature females (Willink et al., 2019).

In addition to colour changes associated with sexual maturation, many odonates

exhibit heritable (genetic) female-limited colour polymorphisms, with one female morph

having male-like colouration (androchrome females) and one or two morphs expressing

female-specific colouration (gynochrome females) (Blow et al., 2021; Fincke et al., 2005).

Observational and experimental field studies suggest that female colour polymorphisms

function to reduce the negative fitness effects of excessive male sexual harassment by

disrupting males' ability to efficiently form search images for mates (Fincke, 2004; Miller &

Fincke, 1999; Takahashi et al., 2014). According to the male mimicry hypothesis,

androchrome females have a negative frequency-dependent advantage in their male-like

similarity, which reduces male mating harassment (Robertson, 1985). Alternatively, although

not mutually exclusive, the learned mate recognition (LMR) hypothesis suggests that males

form a search image for the most common female morph in the population, allowing them to

better detect the most locally abundant female morph, resulting in a rare-morph advantage

(Fincke, 2004). The LMR hypothesis further predicts that the ratio of correct (e.g., mature

conspecific females) versus incorrect (e.g., other males, heterospecifics) male mate

recognition should increase with experience (Miller & Fincke, 2004).

There is partial empirical support for both the male mimicry and LMR hypotheses

and, as stated above, these two hypotheses are not mutually exclusive. In support of the LMR

hypothesis, males of the damselflies *Enallagma civile* and *Ischnura elegans* show increased

preference for androchrome or gynochrome females following previous exposure to these

morphs (Fincke et al., 2007; Miller & Fincke, 1999; Van Gossum et al., 2001b). Further,

naive *Enallagma* damselflies react sexually to both female morphs, but rarely to other males,

contrary to expectations of the male mimicry hypothesis (Fincke et al., 2007). However,

experiments in *Ischnura ramburi* revealed more male-male interactions in androchrome-

biased settings, consistent with both mistaken mate recognition due to male mimicry and the

formation of male search images for the common female phenotype (Gering, 2017).

Similarly, male mating harassment in *I. elegans* increases with increasing morph density for

gynochrome females, indicative of search-image formation, but not for androchrome females

(Gosden & Svensson, 2009).

This growing body of ecological, evolutionary, sensory, and physiological studies,

combined with the increasing availability of genomic resources (Willink et al., 2024), makes

Odonata an excellent system to investigate the molecular basis of colour vision. Here, we

focus on the well-studied Common Bluetail Damselfly (*Ischnura elegans*) and first

characterize opsin spectral sensitivity via heterologous expression in HEK293T cells.

Functional characterization of SWS and LWS rhodopsins show absorbance peaks (λ_{max}) in

the range of 406 – 419 nm and 531 – 548 nm, respectively. We next quantify variation in

opsin mRNA expression across adult male maturation, in populations with variable female

morph frequencies. Opsin expression changes are observed over adult male development,

with increased expression of specific opsin types that correlate with variation in local female

morph frequencies across populations. We next use visual modeling to predict the effect of

shifts in visual sensitivity on female morph detection and discrimination. These results

suggest that opsin expression plasticity may provide one mechanistic proximate link between

male vision, morph detection/discrimination, and the resulting frequency-dependent sexual

conflict, consistent with models for how these genetic polymorphisms are maintained in

natural populations (Fincke, 2004; Le Rouzic et al., 2015; Svensson et al., 2005). Further

experiments coupling behavioral differences in mate preferences will be interesting to link

our results of ontogenetic and plastic variation in male opsin expression profiles to colour

vision preferences and perception of female colour signals in this dynamic system strongly

shaped by visually guided male mating behaviours.

MATERIALS AND METHODS

<u>Study system</u>

Sexually mature male *I. elegans* are monomorphic, exhibiting blue body colouration

on the thorax. In contrast, female *I. elegans* are polymorphic, with sexually mature females

belonging to three distinct genetically determined colour morphs (Fig. 1B) (Willink et al.,

2020). Androchrome (male-like) females exhibit body colouration spectrally similar to male

colouration, while gynochrome females exhibit spectrally distinct green (*infuscans*) or brown

(*infuscans-obsoleta*) thorax colouration (Henze et al., 2019; Van Gossum et al., 2011).

Females exhibit ontogenetic changes in thorax and abdomen colouration (Fig. 1C), with male

I. elegans known to prefer body colouration of mature over immature females (Willink et al.,

2019). Apart from colour differences, androchrome and gynochrome females also differ in

resistance and tolerance to ectoparasites (Willink & Svensson, 2017) and in mating rates,

resistance to mating attempts, and fecundity (Gosden & Svensson, 2009).

<u>Quantification of female morph frequencies between sites and field collection of males</u>

We quantified female morph frequencies from eight sites in southern Sweden (Fig.

1D) as part of an ongoing long-term field population study of *I. elegans* (see Le Rouzic et al.,

2015 and Svensson et al., 2020 for general methodology). Studied sites have both low genetic

differentiation and high allelic diversity, indicating recent divergence and/or high gene flow

(Abbott et al., 2008). All study populations were visited and re-visited at regular intervals (1-

2 weeks) between 19 May – 1 August 2021 and 16 May – 31 July 2022. Including both

immature and mature females, all populations were trimorphic except for one (Höje å 6) in

2022, which had experienced large overall population declines that year (Table S1). The

frequency of each morph in the current study was calculated based on females that displayed

mature adult colouration.

Male *I. elegans* were caught from sampled populations between 28 June – 13 July

2021 and 9 June – 20 July 2022 using sweep nets (Table S2). For all collected individuals,

we recorded sexual maturity ("immature" vs. "mature") by evaluating wing stiffness (Corbet,

1999). We also recorded whether males were found *in copula* with a female or singly

("couple" vs "single"). In 2022, we recorded the morph of the female that was found in

copula with each collected male. In addition to immature and mature males, we also collected

males that had newly emerged from the aquatic nymph stage ("tenerals") from two field sites

and from a semi-naturalistic mesocosm experiment at the Stensoffa Ecological Field Station

at Lund University (Table S2). Teneral males are characterized by their extremely soft wing

and body tissues and lack of body pigment. Including teneral males allowed us to assess

opsin gene expression in males that have not had any significant visual experience or mated

with sexually mature females. All collected males were immediately euthanized by cutting

off their head using cleaned RNAse-free dissection tools, placed into individual vials filled

with RNA*later* (Ambion, Inc., Austin, TX, USA) at field sites, and then stored at -20°C until

RNA extraction.

<u>Functional expression in HEK293T cells and homology modelling</u>

For functional expression, we selected SWS and LWS visual opsin types whose

orthologs in *S. frequens* and *I. asiatica* have been shown to be expressed in the adult ventral

compound eye (SWb1, SWb2, LWA2, LWF1 – F4) and LWE1 to test for function of an

opsin whose ortholog in *I. asiatica* is primarily expressed in ocelli (Futahashi et al., 2015).

Homology modelling was performed for the SWb1 and LWF1 opsin amino acid sequences,

aligned against the invertebrate jumping spider rhodopsin crystal structure (PDB 6i9k)

(Varma et al., 2019). Full methods are presented in supplemental methods.

<u>Quantitative PCR analysis of *I. elegans* opsin expression levels</u>

We quantified whole head opsin mRNA expression via quantitative PCR (see

supplemental methods) in N = 87 male *I. elegans* (8 teneral, 36 immatures, 43 matures; Table

S2) for UV, SWb1, SWb2, LWA2, LWF1-F4 and LWE1 opsins. Hereafter, we use

abbreviated names for the opsins SWb1, SWb2, LWA2, and LWE1, which have been

shortened to SW1, SW2, LWA, and LWE, respectively.

Expression of opsin genes was calculated relative to two housekeeping genes and

calibrated against the UV opsin gene using the $(1 + E)^{-\Delta\Delta CT}$ method where E equals the PCR

efficiency for each primer (Livak & Schmittgen, 2001; Pfaffl, 2001). We also calculated the

proportion opsin expression relative to total opsin expression for opsins likely expressed in

the main compound eye in other odonatans by dividing relative expression for each

individual opsin by the sum of total opsin expression. Results of relative opsin expression

have been shown to indicate differential regulation of opsin genes, while proportional

measures of opsin expression allow to make inferences about colour vision (Fuller &

Claricoates, 2011).

<u>Visual modelling of female morph detection and discrimination</u>

To assess how observed differences in LWS opsin expression might impact detection

or discrimination of female morphs, we focused on the two most abundant female morphs

(androchrome and *infuscans*) and calculated just noticeable differences (JNDs) for three

parsimonious variations of Odonate visual systems. The first modelled visual system follows

ERG spectral sensitivity values from *Ischnura heterosticta* with SWS λ_{max} = 450 nm and

LWS λ_{max} = 525 nm (Huang et al., 2014). For the second and third visual models, based on *I.*

elegans opsin spectral sensitivity, we calculated JNDs with SWS λ_{max} = 413 nm, together

with LWS λ_{max} = 531 nm (model 2) or 543 nm (model 3). The LWS spectral sensitivities

correspond to maximal opsin absorbance spectra for *I. elegans* LWF1 and LWF2, for which

both relative and proportional gene expression decreased and increased, respectively, in

mature versus immature males (see results). All models included a fixed UVS spectral

sensitivity (λ_{max} =360 nm), consistent with published electroretinogram (ERG) data in other

Odonate species (Futahashi et al., 2015; Huang et al., 2014). Equations used in visual model

calculations are presented in supplemental methods.

<u>Statistical analysis</u>

We performed a Multivariate Analysis of Variance (MANOVA) to assess the effect

of maturity on log relative opsin gene expression of all eight opsin genes for male *I. elegans*.

Log relative opsin expression for each individual male was averaged between replicates to

maintain the assumption of independence.

We used linear mixed modeling to test the effects of population morph frequency,

maturity stage, and opsin type on log relative and proportion opsin expression in male *I.*

elegans, with site, male ID, year, and qPCR replicate as possible random effects. Model

selection followed methods in Zuur et al. (2009) to identify the random and fixed effects

included in each model. In total, we ran three main models. **Model 1** assessed changes in

relative opsin expression for the main effects of maturity stage and opsin type, including

opsins likely expressed in both the compound eyes and ocelli. **Model 2** assessed changes in

proportion opsin expression for opsins with likely expression in only the compound eye, with

the main effects of maturity stage and opsin type. **Model 3** tested the effects of maturity

stage, opsin type, population morph frequency and its squared component on relative opsin

expression. Detailed modelling methods are presented in supplemental methods.

We then assessed whether there were differences in opsin expression for males

captured *in copula* with androchrome or *infuscans* females for males captured in 2022.

However, model selection suggested no significant effect of the female morph that a male

was found mating with on relative opsin expression, possibly due to relatively small sample

sizes. We therefore do not discuss these results in the main text (see supplemental methods).

RESULTS

Female morph frequencies

The frequencies of the three mature female morphs differed between sites and years

(Fig. 1E-F). Androchrome females (A) were the majority female morph at all sites, and their

frequency ranged from 57.1 – 85.7% (mean ± sd: 72.6 ± 9.7%) in 2021 and 65.9 – 83.9%

(72.3 ± 5.8%) in 2022. The *infuscans* female morph (I) was the next most common morph,

ranging from 14.3 – 39.1% (24.1 ± 8.8%) in 2021 and 16.1 – 30.7 (24.9 ± 5.3%) in 2022.

Finally, the third morph (*infuscans-obsoleta*, O) were rare, making up only between 0.00 and

5.5% (3.3 ± 1.8%) of mature females in 2021 and between 0.00 and 7.8% (2.8 ± 2.8%) in

2022. Morph fluctuations across years is common in this species, although in many

populations morph frequencies are relatively stable over time and across many generations,

as a result of negative frequency-dependent selection (Le Rouzic et al., 2015; Svensson &

Abbott, 2005). Accordingly, only one of these eight populations differed significantly in

morph frequencies between these two years (Vombs Vattenverk). This population had a 1.7-

fold decrease in the frequency of I-females and a 10-fold decrease in the frequency of O-

females relative to A-females between the two years (see Table S3 for statistical comparisons

for all sites).

Opsin spectral sensitivity determined using HEK293 cell heterologous expression

We found that the SWS opsins SWb1 and SWb2 absorb maximally at λ_{max} of 406 and

419 nm (Fig. 2A-B). The LWS opsin types display maximal sensitivity between 530 and 550

nm with the following λ_{max}: LWA2 = 548 nm, LWF1 = 531 nm, LWF2 = 543 nm, LWF3 =

545 nm, LWF4 = 541 nm, LWE1 = 533 nm (Fig. 2C-H). Spectral sensitivity functions of

SWS and LWS opsins overlap with spectral reflectance of female thorax coloration of

androchrome and *infuscans* females (Fig. 3).

We ran homology modeling of SWb1 and LWF1 opsins against the spider rhodopsin

(PDB: 6i9k) and mapped sites interacting with the chromophore in PyMOL. Of the 71 amino

acid substitutions between SWb1 and SWb2 (81.8% aa identity), we identified 23 sites

predicted to be within 5Å of the binding pocket (Table S4). Among these, Y136 (TM3) and

Y294 (TM6) residues located at the top and bottom of the binding pocket, respectively, are

substituted by phenylalanine (F136 and F294) in SWb2 (Fig. S1A). For LWS opsins, which

share between 88.9 - 91.2% aa identity (i.e., 33 - 42 variant residues), we obtained the

predicted LWF1 opsin structure based on 6i9k and mapped 24 sites predicted to interact with

the cis-retinal, all of which are conserved for all four LWF opsins (Table S4; Fig. S1B). The

LWF2 protein shares 89.9-90.5% identity with LWF1-F4, translating to 36 to 38 residue

substitutions, and 67.4% identity (123 amino acid residue differences) with LWA2 (Fig.

S3B). LWA2 is more divergent in sequence, 67.5 - 68.7% aa identity with 119 - 124 aa

differences with LWF opsins. Of the 24 chromophore-interacting residues, its binding pocket

exhibits two potential spectral variant residues with LWF1 (A131G in TM1 and S319A in

TM7).

Opsin expression in teneral, immature, and mature males

We compared the expression levels of 2 SWS and 6 LWS opsins, relative to UV opsin

expression, across adult male developmental stages in *I. elegans* from eight sites. Our results

from the MANOVA showed a significant effect of male maturity stage on overall opsin gene

expression ($F_{16,146}$ = 3.2, *p* < 0.0001; Fig. 4). To further assess how relative and proportion

opsin expression differed between teneral, immature, and mature male *I. elegans*, we

performed mixed modeling with maturity stage, opsin type, and their interaction as main

effects.

For relative opsin expression (model 1), we found a significant interaction between

maturity stage and opsin (Table 1, Tables S5 – S6) with increasing differences in relative

expression from teneral males (Fig. 5A) to immature (Fig. 5B) and mature (Fig. 5C) males.

For teneral males, highest relative opsin expression was for LWF4 (Fig. 5A), while LWF2

has the highest relative expression for immature and mature males (Fig. 5B, C). Results also

show that, for immature and mature males, the opsin LWE has significantly lower relative

expression relative to other opsin types, consistent with expression levels observed in S.

frequens and *I. asiatica* (Futahashi et al., 2015).

Comparing teneral, immature, and mature males, results showed that expression

changes over male maturation are not parallel across opsins, but that there were opsin-

specific changes (Fig. 6). Increases in relative expression occurred for the SW1 (teneral –

mature: t = -2.7, df = 396, p = 0.02) and LWF2 opsins (teneral – immature: t = -5.1 df = 401,

$p < 0.0001$; teneral – mature: t = -6.9, df = 396, $p < 0.0001$; immature – mature: t = -3.0, df =

419, p = 0.009). Conversely, decreases in relative expression occurred for the LWF1 (teneral

– immature: t = 4.7, df = 402, $p < 0.0001$; teneral – mature: t = 6.1, df = 399, $p < 0.0001$;

immature – mature: t = 2.4, df = 433, p = 0.04) and LWE opsins (teneral – immature: t = 3.4,

df = 401, p = 0.002; teneral – mature: t = 3.9, df = 400, p = 0.0004). For results for each

independent population, see supplemental materials (Table S7, Fig. S2).

When analyzing proportional opsin expression (model 2), we again found a

significant interaction between maturity stages and opsin type (Table 2). For all LWS opsin

types, with the exception of LWF1, we found a significant difference between teneral males

and both immature and mature males, but not between immature and mature males (Table

S8). For LWF1, there was a significant difference between all maturity stages tested (teneral

– immature: t = 12.4, df = 84, *p* < 0.001; teneral – mature: t = 15.2, df = 84, *p* < 0.001;

immature – mature: t = 4.4, df = 84, *p* < 0.001). Relative and proportional opsin expression

were thus largely consistent, with a decrease in LWF1 expression and increase in LWF2

expression over maturity, although differences in proportional LWF2 expression were not

significant between immature and mature males (t = 2.1, df = 84, *p* = 0.10), as with relative

expression (Fig. 7). LWF4 expression, which did not significantly differ in relative

expression over development (Fig. 6), made up the largest proportion expressed for teneral

males (0.523 ± 0.01). In immature and mature males, LWF2 made up the largest proportion

opsin expressed, accounting for 0.524 ± 0.03 and 0.599 ± 0.02 of expression, respectively

(Fig. 7).

Opsin expression across populations with different female morph frequencies

Comparing variation in opsin expression in immature and mature males including the

effect of local female morph frequency (model 3), we find a significant interaction between

opsin type, maturity stage, and the squared component of androchrome frequency (Table 3).

Plotting log relative expression for each opsin type revealed a quadratic relationship between

LWF2 expression and andromorph frequency in mature, but not in immature, males. We

observed the highest LWF2 expression in populations with the lowest and highest proportion

of androchrome females. In contrast, androchrome frequency had little effect on relative

expression for immature or mature males for the other opsins tested (Fig. 8).

Visual modelling of female morph detection and discrimination

Comparing the results of visual models using ERG data from *I. heterosticta*, *I.*

elegans LWF1 opsin sensitivity, and *I. elegans* LWF2 opsin sensitivity, we found that the

LWF2 visual system had larger just-noticeable differences (JNDs) for discriminating

androchrome and *infuscans* color morphs (ΔJND$_{LWF2-ERG}$ = 0.14, ΔJND$_{LWF2-LWF1}$ = 0.41) and

for detecting androchrome females against brown and green vegetation (brown: ΔJND$_{LWF2-ERG}$

$_{ERG}$ = 0.53, ΔJND$_{LWF2-LWF1}$ = 0.56; green: ΔJND$_{LWF2-ERG}$ = 0.14, ΔJND$_{LWF2-LWF1}$ = 0.36).

Modelling with parameters from the *I. heterosticta* ERG visual system led to larger JNDs for

detection of *infuscans* females against brown vegetation (ΔJND$_{LWF1-ERG}$ = -0.16, ΔJND$_{LWF2-}$

$_{ERG}$ = -0.17) (Table S9). The three tested visual systems performed similarly well for

discrimination of *infuscans* females against green vegetation (ΔJND$_{LWF2-ERG}$ = 0.02,

ΔJND$_{LWF2-LWF1}$ = 0.07, ΔJND$_{LWF1-ERG}$ = -0.05).

DISCUSSION

Vision is an important sensory modality in Odonata and the prevalence of female-

limited colour polymorphisms makes the damselfly genus *Ischnura* a powerful model system

to study how and why polymorphisms emerge and persist over both micro- and

macroevolutionary time scales (Blow et al., 2021; Le Rouzic et al., 2015; Willink et al.,

2024). Here, we combined ecological, molecular, and functional approaches to investigate

opsin gene expression in male *I. elegans* of different maturation stages and across

populations with varying female morph frequencies. Our measures of *in vitro* opsin

absorbance evidence that the photosensitive opsin receptors forming the peripheral visual

system of *I. elegans* perceive wavelengths that overlap with reflectance intensity (brightness)

and chromatic body colouration cues of the two most abundant female morphs, namely

androchrome and *infuscans*, from field sites (Fig. 3). Further, recurrent directional changes in

opsin expression over male ontogeny, particularly during sexual maturation (Figs. 4 - 7), as

well as changing male gene expression profiles in response to local female morph frequencies

(Fig. 8) suggest a role for opsin expression plasticity in the *Ischnura* system.

Vision is used for many tasks in Odonates, and ecological factors might select for

additional differences that may not necessarily be attributable to conspecific search and

morph discrimination. The spatial organization of yet unknown ommatidial types possibly

expressing a subset of the multiple SWS and LWS opsins across the retina may also influence

how male *I. elegans* perceive conspecific body cues. However, *I. elegans* is equipped with

numerous functional SWS and LWS visual opsins activated maximally by wavelengths of

light highly reflected by female morphs (Fig. 3). In light of known behavioural male

preference for female colour morphs and dynamically changing morph frequencies (Van

Gossum et al., 1999, 2001b), opsin-based spectral sensitivity across short and long

wavelengths likely contributes to visual detection and discrimination of local female morphs.

Overlapping opsin spectral sensitivities likely confer broad wavelength discrimination

A total of 12 expressed visual opsin genes have been identified in the damselfly

Ischnura asiatica (Futahashi et al., 2015), a gene repertoire that we find conserved in the

genome of *I. elegans* (Fig. S3A). From the *I. elegans* opsin gene set, we quantitatively

assessed whole head adult expression levels of 2 SWS (SWb1, SWb2) and 5 LWS (LWA,

LWF1-F4) opsins for which orthologs in *I. asiatica* and *Sympetrum frequens* are primarily

expressed in the ventral eye (Futahashi et al 2015), a region implicated in conspecific search

(Lancer et al., 2020). For comparison, we also quantified expression levels of *I. elegans*

LWE1 opsin, which is restricted to ocelli in other odonates (Futahashi et al 2015). The *I.*

elegans LWE1 opsin, which is sensitive to green wavelength light (Fig. 2), consistently

exhibited the lowest monitored expression levels across opsin types (Fig. 3), suggesting a

restricted ocelli-specific expression pattern and function. In contrast, SWS and the other

LWS opsins exhibited higher baseline expression levels compared to LWE1, as observed in *I.*

asiatica, therefore supporting a role as visual opsins in the main compound eye.

The SWS opsins confer sensitivity to light in the violet-blue spectrum (λ_{max} = 406 nm

and 419 nm) and the five visual LWS opsins show broad sensitivity to green wavelengths of

light, with λ_{max} between 530 nm and 540 – 550 nm (Fig. 2). This forms a key component

defining the genetic basis of light perception and functionality of multiple opsin duplicates,

as part of several complementary mechanisms that certainly contribute to modulate visual

sensitivity and capture a remarkably broad light spectrum from ultraviolet to long-

wavelengths above 600 nm. While achromatic vision of intensity related cues can be directly

modulated by light capture of single opsins (Kelber, 2005), chromatic discrimination involves

at least two distinct but overlapping sensitivities in a given ommatidial unit (Kelber &

Osorio, 2010; Lind et al., 2017). ERG recordings from other odonates show broad sensitivity

of the green photoreceptor type which may indicate the presence of multiple opsin genes

expressed within a single photoreceptor (Futahashi, 2016; Huang et al., 2014). Irrespective of

how many LWS opsins may be expressed in a single photoreceptor or ommatidium, overlap

between short- (UV and violet-blue) and at least one long-wavelength opsin sensitivities

could provide fine-tuned trichromatic discrimination capacity across a broad spectral range in

I. elegans (Figs. 2, 3), with additional filtering mechanisms common in arthropods potentially

further increasing spectral channels (Cronin et al., 2001; Stavenga & Arikawa, 2011).

Our results show a high degree of overlap in the recorded sensitivities of the LWS

opsins, which suggest that changes in expression of any single opsin might not have a large

impact on spectral sensitivity if several LWS opsins are co-expressed. This also suggests that

there may be no opponency unless combining LWS opsins with 10-20 nm differences

between peak spectral sensitivity (Kelber et al., 2003). However, substantial increases and

decreases in opsin expression levels for opsin types with λ_{max} differences greater than 10 nm

(e.g., LWF1 λ_{max} = 531 nm, and LWF2 λ_{max} = 543 nm), may affect the underlying

photoreceptor spectral visual sensitivity if these specific opsins are also spatially segregated

across the retina.

Spatial differences in opsin expression patterns have been identified in other insects

(Rister & Desplan, 2011) including other species of Odonata (Futahashi et al., 2015; Labhart

& Nilsson, 1995), which suggests that *I. elegans* LWS opsin duplicates could have

regionalized expression patterns throughout the retina. For example, the eyes of the dragonfly

Hemicordulia tau seem to have three short-wavelength sensitive zones (i.e., the dorsal acute

zone, the dorsal rim area, and the frontal acute zone) and display ventral spectral band

patterning maximally sensitive to either short-, middle-, or long-wavelengths (Lancer et al.,

2020). Likewise, ERG recordings from the dragonfly *S. frequens* show regional differences in

spectral sensitivity coinciding with dorso-ventral opsin expression levels and broad green

spectral sensitivity (Futahashi et al., 2015). Future analysis of LWS opsin spatial distribution

via gene-specific antibody staining or *in situ* hybridization, along with *in vivo* ERG

quantification of visual sensitivity, will be valuable next steps to determine how opsin spatial

distribution and co-expression patterns may contribute to finely tune visual spectral

sensitivity in *I. elegans.*

Candidate spectral tuning sites between SWS and LWS opsin duplicates

Homology modeling indicated that the residues predicted to interact with the retinal

chromophore are conserved between all tested LWF opsins. It is therefore possible that

variant residues at positions outside the canonical binding pocket play a role in fine spectral

tuning, causing the observed narrow spectral shifts.

For SWS opsins, we mapped two strong candidate spectral tuning sites at positions

136 and 294 between *I. elegans* duplicate SWb1 (λ_{max} 406 nm) and SWb2 (λ_{max} 419 nm)

opsins (Fig. S1). A comparison of duplicate SWb opsins across five *Ischnura* species (see SI

dataset 2), *I. cervula*, *I. verticalis*, *I. hastata* and *I. asiatica,* shows that Y294F (H6) is present

only in *I. cervula* whereas Y136F (H3) is conserved across all species. Although we have not

tested the effects on these mutations in the present study, repeated homologous Y/F changes

have been functionally shown to cause convergent spectral shifts in SWS insect opsins

(Frentiu et al., 2015; Liénard et al., 2021; Wakakuwa et al., 2010). Accumulated evidence for

the role of Y to F substitutions in providing longer wavelength-sensitive SWS invertebrate

opsins is consistent with the gain of F136 in *I. elegans* SWSb2, possibly underlying the

change in paralog SWS sensitivity (Fig. 2). The conserved Y136F substitution presumably

also influences spectral tuning across other *Ischnura* SWb1 and SWb2 opsin paralogs,

although possibly to varying degrees, as species-specific variation in spectral sensitivity can

be expected between orthologous native opsins, as observed with the orthologous opsins of *S.*

frequens: LWA2 (λ_{max} 557 nm) and LWD1 (λ_{max} 542 nm) (Liénard et al., 2022) compared to

the *Ischnura* orthologs (LWA2 λ_{max} 548 nm; LWD1 λ_{max} 541 nm) characterized here.

Opsin expression plasticity as a potential mechanism underlying morph detection and

discrimination

In this system with rapidly changing female morph frequencies, males are expected to

rapidly develop plastic search images, which may be shaped by local frequencies of female

morphs and their intrinsic fecundities (Fincke, 2004; Gosden & Svensson, 2009; Verzijden et

al., 2012). The visual system and peripheral opsin receptors are known major evolutionary

targets (Dungan & Chang, 2022; Hauzman et al., 2021; Macias-Muñoz et al., 2016; Schott et

al., 2024) with the potential to fine tune wavelength light capture and colour vision, in

particular in altering opsin gene evolution, function and expression. There is also increasing

empirical evidence for a key role of phenotypic plasticity in evolution, including in shaping

the development of both male and female mate preferences (Verzijden et al., 2012; West-

Eberhard, 2003). Such learned mate preferences may impact sexual selection, sexual conflict,

the maintenance of sexually selected polymorphisms, population divergence, and ultimately

speciation (Svensson et al., 2010; Verzijden et al., 2012; Westerman et al., 2012).

The damselfly *I. elegans* is characterized by visually guided mate-searching males

who are exposed to female colour polymorphisms and a continuum of androchrome female

frequencies across local populations (Fig. 1). This system allowed us to explore male visual

system response dynamics quantitatively in natural field settings. Quantification of opsin

mRNA levels throughout adult *I. elegans* maturation at eight sampling sites over two flying

seasons revealed consistent plasticity for specific opsin genes over adult male development.

In particular, we identified opsin-specific directional changes in relative expression for

LWF1 and LWF2 and proportional changes in the same opsin genes, in addition to the other

LWS opsin types tested, over male ontogeny across several populations (Figs. 4 - 7). This

suggests that males might experience age-specific selection pressures on these vision genes.

Successful visual mate identification has also been predicted to vary in response to local

female morph abundances and correlate with increased sexual experience (Fincke, 2004;

Miller & Fincke, 2004). Since opsin expression profiles of adult males were consistent with a

gradual increase (i.e., LWF2) and decrease (i.e., LWF1) in opsins with distinct spectral

sensitivities, these changes may contribute, at least partly, in fine-tuning the male visual

system for mate detection and mate discrimination during sexual maturation.

Moreover, sexually immature teneral males are inexperienced with visually guided

behaviors compared to immature and mature adult males. In line with this, we found that

opsin gene expression levels, for SWS and LWS visual opsin types, are more homogenous in

teneral whole head tissues, whereas variation in relative expression increased in immature

and mature male *I. elegans* (Fig. 5). Major directional expression changes were found at the

metapopulation level, including changes that reliably depend on maturity stage, and target

opsin type, with notable decreases in LWF1 and increases in LWF2 expression (Fig. 6). The

relationship between LWF1 and LWF2 expression appeared to be conserved across all eight

sampled populations, and statistical signal is recovered at the population level for LWF1 and

LWF2 in several of the individual sites (Fig. S2), despite lower sampling power. Taken

together, these differences in relative expression are consistent with opsin-specific regulation

during ontogenic development and male sexual maturation (Fig. 6), suggesting that vision

changes in immature and mature adult males compared to teneral males.

Proportional opsin expression also differed between developmental stages, with

significantly more abundant LWF2 opsin expression and significantly less abundant

expression of other LWS opsin types in immature and mature males relative to teneral males

(Fig. 7). Visual modelling, informed by *in vitro* measurements of opsin sensitivity, indicated

that visual systems with maximal LWS opsin sensitivity set as the LWF2 λ_{max} (i.e., 543 nm)

would likely outperform chromatic visual systems modelled with shorter LWS λ_{max} values

(i.e., LWF1 and ERG λ_{max}) in discriminating between thorax coloration of androchrome and

infuscans females, and for detecting androchromes against vegetation. Additionally, males

may respond to achromatic cues, with a nearly 2-fold difference in the overall brightness of

androchrome and *infuscans* females at the descending midpoint limb of absorbance of LWS

opsins around 600 nm (Fig. 3). These results suggest that abundantly expressed LWS opsins

sensitive to longer wavelengths, can provide an ecological advantage for specific visual tasks

involved in conspecific detection and likely discrimination between brightness and/or

chromatic cues.

For mature males across all sites, the LWF2 opsin expression pattern exhibited a

strong positive quadratic relationship ($\alpha > 1$), meaning that LWF2 expression was

significantly greater in populations with either very low (<0.6) or very high (>0.8) female

androchrome frequencies (Fig. 8). We established that the long wavelength shifted spectral

sensitivity of LWF2 can, in principle, improve chromatic discrimination of androchrome and

infuscans females and improve detection of androchrome females against vegetation, which

could benefit males in populations were androchrome females are more or less abundant

compared to other sites (although still the majority morph in all populations studied here).

Further behavioural experiments will be important to causally establish if and how female

morph frequencies might impact male behavioural choice.

Links between opsin expression and behaviour, while weak in some cases (Fuller et

al., 2010; Wright et al., 2020) have been established in several biological systems (Sakai et

al., 2018; Seehausen et al., 2008). Such studies include female guppies from low-predator

populations which have been shown to prefer males with more orange/red colouration, a

preference linked to higher expression of opsins sensitive to orange and red wavelengths

compared to females from high-predator populations (Sandkam et al., 2015). Future work

directly assessing the effect of mating outcomes on specific opsin expression profiles will

provide valuable insights into the function of opsin plasticity in mate detection, mate choice

and female morph preference in Odonata.

CONCLUSION

Our study has revealed that the visual system of *I. elegans* is characterized by

multiple SWS and LWS opsin genes, and that their absorbance spectra confer broad spectral

sensitivity. Future studies should further assess regional photoreceptor spectral sensitivity,

particularly in the ventral eye, which will reveal the level of functional redundancy and

chromacity. We found striking variation in opsin gene expression profiles between male

maturation stages and across populations with varying female morph frequencies.

Specifically, the long-wavelength sensitive LWF2 opsin, showed expression profiles that

consistently increased over male ontogeny, and showed expression changes that correlated

with local female morph frequencies encountered by mate-searching mature male *I. elegans*.

Conversely, the relative and proportional expression of another long-wavelength sensitive

opsin, LWF1, decreased over male ontogeny. Functional analyses revealed that the range of

SWS and LWS spectral sensitivity of *I. elegans* likely allows males to discriminate

androchrome and *infuscans* female body reflectance, with visual modeling indicating that

long-wavelength sensitivity characterized by LWF2 would improve detection of

andromorphs against vegetation and discrimination between the two most common morph

types. Other behavioural and physiological differences between androchrome and

gynochrome females, such as aggression or fecundity differences, or ecological factors such

as local lightning environments that are not necessarily related to mating behaviours may also

play a role in modulating male opsin gene expression profiles. Overall, our new results

suggest that directional changes in opsin expression may reflect visual plasticity that might

aid males in mate recognition, morph detection and/or discrimination. Future work should

aim to complement these findings from natural populations with experimental manipulations

to determine if there is a casual role in mate choice in modulating expression profiles.

REFERENCES

Abbott, J. K., Bensch, S., Gosden, T. P., & Svensson, E. I. (2008). Patterns of

differentiation in a colour polymorphism and in neutral markers reveal rapid genetic

changes in natural damselfly populations. *Molecular Ecology*, *17*(6), 1597–1604.

https://doi.org/10.1111/j.1365-294X.2007.03641.x

Blow, R., Willink, B., & Svensson, E. I. (2021). A molecular phylogeny of forktail

damselflies (genus *Ischnura*) reveals a dynamic macroevolutionary history of female

colour polymorphisms. *Molecular Phylogenetics and Evolution*, *160*, 107134.

https://doi.org/10.1016/j.ympev.2021.107134

Briscoe, A. D., & Chittka, L. (2001). The evolution of color vision in insects. *Annual Review*

of Entomology, *46*(1), 471–510. https://doi.org/10.1146/annurev.ento.46.1.471

Bybee, S. M., Johnson, K. K., Gering, E. J., Whiting, M. F., & Crandall, K. A. (2012). All the

better to see you with: A review of odonate color vision with transcriptomic insight

into the odonate eye. *Organisms Diversity & Evolution*, *12*(3), 241–250.

https://doi.org/10.1007/s13127-012-0090-6

Castiglione, G. M., Hauser, F. E., Van Nynatten, A., & Chang, B. S. W. (2023). Adaptation of

antarctic icefish vision to extreme environments. *Molecular Biology and Evolution*,

40(4), msad030. https://doi.org/10.1093/molbev/msad030

Chang, C.-H., Catchen, J., Moran, R. L., Rivera-Colón, A. G., Wang, Y.-C., & Fuller, R. C.

(2021). Sequence analysis and ontogenetic expression patterns of cone opsin genes in

the bluefin killifish (*Lucania goodei*). *Journal of Heredity*, *112*(4), 357–366.

https://doi.org/10.1093/jhered/esab017

Corbet, P. S. (1999). *Dragonflies: Behavior and Ecology of Odonata.* Cornell Univ. Press,

Ithaca, NY. 829pp.

Cronin, T. W., Caldwell, R. L., & Marshall, J. (2001). Tunable colour vision in a mantis

shrimp. *Nature*, *411*(6837), Article 6837. https://doi.org/10.1038/35079184

Dungan, S. Z., & Chang, B. S. W. (2022). Ancient whale rhodopsin reconstructs dim-light

vision over a major evolutionary transition: Implications for ancestral diving behavior.

Proceedings of the National Academy of Sciences, *119*(27), e2118145119.

https://doi.org/10.1073/pnas.2118145119

Everett, A., Tong, X., Briscoe, A. D., & Monteiro, A. (2012). Phenotypic plasticity in opsin

expression in a butterfly compound eye complements sex role reversal. *BMC*

Evolutionary Biology, *12*(232). https://doi.org/10.1186/1471-2148-12-232

Fincke, O. M. (2004). Polymorphic signals of harassed female odonates and the males that

learn them support a novel frequency-dependent model. *Animal Behaviour*, *67*(5),

833–845. https://doi.org/10.1016/j.anbehav.2003.04.017

Fincke, O. M., Fargevieille, A., & Schultz, T. D. (2007). Lack of innate preference for morph

and species identity in mate-searching Enallagma damselflies. *Behavioral Ecology*

and Sociobiology, *61*(7), 1121–1131. https://doi.org/10.1007/s00265-006-0345-3

Fincke, O. M., Jödicke, R., Paulson, D. R., & Schultz, T. D. (2005). The evolution and

frequency of female color morphs in Holarctic Odonata: Why are male-like females

typically the minority? *International Journal of Odonatology*, *8*(2), 183–212.

https://doi.org/10.1080/13887890.2005.9748252

Frentiu, F. D., Yuan, F., Savage, W. K., Bernard, G. D., Mullen, S. P., & Briscoe, A. D.

(2015). Opsin clines in butterflies suggest novel roles for insect photopigments.

Molecular Biology and Evolution, *32*(2), 368–379.

https://doi.org/10.1093/molbev/msu304

Fuller, R. C., & Claricoates, K. M. (2011). Rapid light-induced shifts in opsin expression:

Finding new opsins, discerning mechanisms of change, and implications for visual

sensitivity. *Molecular Ecology*, *20*(16), 3321–3335. https://doi.org/10.1111/j.1365-

294X.2011.05180.x

Fuller, R. C., Noa, L. A., & Strellner, R. S. (2010). Teasing apart the many effects of lighting

environment on opsin expression and foraging preference in bluefin killifish. *The*

American Naturalist, *176*(1), 1–13. https://doi.org/10.1086/652994

Futahashi, R. (2016). Color vision and color formation in dragonflies. *Current Opinion in*

Insect Science, *17*, 32–39. https://doi.org/10.1016/j.cois.2016.05.014

Futahashi, R., Kawahara-Miki, R., Kinoshita, M., Yoshitake, K., Yajima, S., Arikawa, K., &

Fukatsu, T. (2015). Extraordinary diversity of visual opsin genes in dragonflies.

Proceedings of the National Academy of Sciences, *112*(11), E1247–E1256.

https://doi.org/10.1073/pnas.1424670112

Gering, E. J. (2017). Male-mimicking females increase male-male interactions, and decrease

male survival and condition in a female-polymorphic damselfly. *Evolution*, *71*(5),

1390–1396. https://doi.org/10.1111/evo.13221

Gosden, T. P., & Svensson, E. I. (2009). Density☐dependent male mating harassment, female

resistance, and male mimicry. *The American Naturalist*, *173*(6), 709–721.

https://doi.org/10.1086/598491

Hauzman, E., Pierotti, M. E. R., Bhattacharyya, N., Tashiro, J. H., Yovanovich, C. A. M.,

Campos, P. F., Ventura, D. F., & Chang, B. S. W. (2021). Simultaneous expression of

UV and violet SWS1 opsins expands the visual palette in a group of freshwater

snakes. *Molecular Biology and Evolution*, *38*(12), 5225–5240.

https://doi.org/10.1093/molbev/msab285

Henze, M. J., Lind, O., Wilts, B. D., & Kelber, A. (2019). Pterin-pigmented nanospheres

create the colours of the polymorphic damselfly *Ischnura elegans*. *Journal of The*

Royal Society Interface, *16*(153), 20180785. https://doi.org/10.1098/rsif.2018.0785

Huang, S., Chiou, T., Marshall, J., & Reinhard, J. (2014). Spectral sensitivities and color

signals in a polymorphic damselfly. *PLOS ONE*, *9*(1), e87972.

https://doi.org/10.1371/journal.pone.0087972

Jacobs, G. H. (2013). Losses of functional opsin genes, short-wavelength cone

photopigments, and color vision—A significant trend in the evolution of mammalian

vision. *Visual Neuroscience*, *30*(1–2), 39–53.

https://doi.org/10.1017/S0952523812000429

Kelber, A. (2005). Alternative use of chromatic and achromatic cues in a hawkmoth.

Proceedings of the Royal Society B: Biological Sciences, *272*(1577), 2143–2147.

https://doi.org/10.1098/rspb.2005.3207

Kelber, A., & Osorio, D. (2010). From spectral information to animal colour vision:

Experiments and concepts. *Proceedings of the Royal Society B: Biological Sciences*,

277(1688), 1617–1625. https://doi.org/10.1098/rspb.2009.2118

Kelber, A., Vorobyev, M., & Osorio, D. (2003). Animal colour vision-behavioural tests and

physiological concepts. *Biol Rev*, *78*, 81–118.

Labhart, T., & Nilsson, D.-E. (1995). The dorsal eye of the dragonfly *Sympetrum*:

Specializations for prey detection against the blue sky. *Journal of Comparative*

Physiology A, *176*(4), 437–453. https://doi.org/10.1007/BF00196410

Lancer, B. H., Evans, B. J. E., & Wiederman, S. D. (2020). The visual neuroecology of

anisoptera. *Current Opinion in Insect Science*, *42*, 14–22.

https://doi.org/10.1016/j.cois.2020.07.002

Le Rouzic, A., Hansen, T. F., Gosden, T. P., & Svensson, E. I. (2015). Evolutionary time-

series analysis reveals the signature of frequency-dependent selection on a female

mating polymorphism. *The American Naturalist*, *185*(6), E182–E196.

https://doi.org/10.1086/680982

Liénard, M. A., Bernard, G. D., Allen, A., Lassance, J.-M., Song, S., Childers, R. R., Yu, N.,

Ye, D., Stephenson, A., Valencia-Montoya, W. A., Salzman, S., Whitaker, M. R. L.,

Calonje, M., Zhang, F., & Pierce, N. E. (2021). The evolution of red color vision is

linked to coordinated rhodopsin tuning in lycaenid butterflies. *Proceedings of the*

National Academy of Sciences, *118*(6), e2008986118.

https://doi.org/10.1073/pnas.2008986118

Liénard, M. A., Valencia-Montoya, W. A., & Pierce, N. E. (2022). Molecular advances to

study the function, evolution and spectral tuning of arthropod visual opsins.

Philosophical Transactions of the Royal Society B: Biological Sciences, *377*(1862),

20210279. https://doi.org/10.1098/rstb.2021.0279

Lind, O., Henze, M. J., Kelber, A., & Osorio, D. (2017). Coevolution of coloration and colour

vision? *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1724), 20160338. https://doi.org/10.1098/rstb.2016.0338

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-

time quantitative PCR and the 2−ΔΔct method. *Methods*, *25*(4), 402–408.

https://doi.org/10.1006/meth.2001.1262

Macias-Muñoz, A., Smith, G., Monteiro, A., & Briscoe, A. D. (2016). Transcriptome-wide

differential gene expression in *Bicyclus anynana* butterflies: Female vision-related

genes are more plastic. *Molecular Biology and Evolution*, *33*(1), 79–92.

https://doi.org/10.1093/molbev/msv197

McCulloch, K. J., Yuan, F., Zhen, Y., Aardema, M. L., Smith, G., Llorente-Bousquets, J.,

Andolfatto, P., & Briscoe, A. D. (2017). Sexual dimorphism and retinal mosaic

diversification following the evolution of a violet receptor in butterflies. *Molecular*

Biology and Evolution, *34*(9), 2271–2284. https://doi.org/10.1093/molbev/msx163

Miller, M. N., & Fincke, O. M. (1999). Cues for mate recognition and the effect of prior

experience on mate recognition in *Enallagma* damselflies. *Journal of Insect Behavior*,

12(6), 801–814. https://doi.org/10.1023/A:1020957110842

Miller, M. N., & Fincke, O. M. (2004). Mistakes in sexual recognition among sympatric

Zygoptera vary with time of day and color morphism (Odonata: Coenagrionidae).

International Journal of Odonatology, *7*(3), 471–491.

https://doi.org/10.1080/13887890.2004.9748233

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT–

PCR. *Nucleic Acids Research*, *29*(9), e45. https://doi.org/10.1093/nar/29.9.e45

Rebora, M., Frati, F., Piersanti, S., Salerno, G., Selvaggini, R., & Fincke, O. M. (2018). Field

tests of multiple sensory cues in sex recognition and harassment of a colour

polymorphic damselfly. *Animal Behaviour*, *136*, 127–136.

https://doi.org/10.1016/j.anbehav.2017.12.015

Rister, J., & Desplan, C. (2011). The retinal mosaics of opsin expression in invertebrates and

vertebrates. *Developmental Neurobiology*, *71*(12), 1212–1226.

https://doi.org/10.1002/dneu.20905

Robertson, H. M. (1985). Female dimorphism and mating behaviour in a damselfly, *Ischnura*

ramburi: Females mimicking males. *Animal Behaviour*, *33*(3), 805–809.

https://doi.org/10.1016/S0003-3472(85)80013-0

Rossetto, I. H., Sanders, K. L., Simões, B. F., Van Cao, N., & Ludington, A. J. (2023).

Functional duplication of the short-wavelength-sensitive opsin in sea snakes:

Evidence for reexpanded color sensitivity following ancestral regression. *Genome*

Biology and Evolution, *15*(7), evad107. https://doi.org/10.1093/gbe/evad107

Sakai, Y., Kawamura, S., & Kawata, M. (2018). Genetic and plastic variation in opsin gene

expression, light sensitivity, and female response to visual signals in the guppy.

Proceedings of the National Academy of Sciences, *115*(48), 12247–12252.

https://doi.org/10.1073/pnas.1706730115

Sandkam, B., Young, C. M., & Breden, F. (2015). Beauty in the eyes of the beholders: Colour

vision is tuned to mate preference in the Trinidadian guppy (*Poecilia reticulata*).

Molecular Ecology, *24*(3), 596–609. https://doi.org/10.1111/mec.13058

Sasagawa, H., Narita, R., Kitagawa, Y., & Kadowaki, T. (2003). The expression of genes

encoding visual components is regulated by a circadian clock, light environment and

age in the honeybee (*Apis mellifera*). *European Journal of Neuroscience*, *17*(5), 963–970. https://doi.org/10.1046/j.1460-9568.2003.02528.x

Schott, R. K., Fujita, M. K., Streicher, J. W., Gower, D. J., Thomas, K. N., Loew, E. R., Bamba Kaya, A. G., Bittencourt-Silva, G. B., Guillherme Becker, C., Cisneros-Heredia, D., Clulow, S., Davila, M., Firneno, T. J., Jr, Haddad, C. F. B., Janssenswillen, S., Labisko, J., Maddock, S. T., Mahony, M., Martins, R. A., … Bell, R. C. (2024). Diversity and evolution of frog visual opsins: Spectral tuning and adaptation to distinct light environments. *Molecular Biology and Evolution*, *41*(4), msae049. https://doi.org/10.1093/molbev/msae049

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., van der Sluijs, I., Schneider, M. V., Maan, M. E., Tachida, H., Imai, H., & Okada, N. (2008). Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), Article 7213. https://doi.org/10.1038/nature07285

Shand, J., Davies, W. L., Thomas, N., Balmer, L., Cowing, J. A., Pointer, M., Carvalho, L. S., Trezise, A. E. O., Collin, S. P., Beazley, L. D., & Hunt, D. M. (2008). The influence of ontogeny and light environment on the expression of visual pigment opsins in the retina of the black bream, *Acanthopagrus butcheri*. *Journal of Experimental Biology*, *211*(9), 1495–1503. https://doi.org/10.1242/jeb.012047

Sharkey, C. R., Blanco, J., Lord, N. P., & Wardill, T. J. (2023). Jewel beetle opsin duplication and divergence is the mechanism for diverse spectral sensitivities. *Molecular Biology and Evolution*, *40*(2), msad023. https://doi.org/10.1093/molbev/msad023

Stavenga, D. G., & Arikawa, K. (2011). Photoreceptor spectral sensitivities of the Small White butterfly *Pieris rapae crucivora* interpreted with optical modeling. *Journal of Comparative Physiology A*, *197*(4), 373–385. https://doi.org/10.1007/s00359-010-0622-5

Svensson, E. I., & Abbott, J. (2005). Evolutionary dynamics and population biology of a

polymorphic insect. *Journal of Evolutionary Biology*, *18*(6), 1503–1514.

https://doi.org/10.1111/j.1420-9101.2005.00946.x

Svensson, E. I., Abbott, J., & Härdling, R. (2005). Female polymorphism, frequency

dependence, and rapid evolutionary dynamics in natural populations. *The American*

Naturalist, *165*(5), 567–576. https://doi.org/10.1086/429278

Svensson, E. I., Eroukhmanoff, F., Karlsson, K., Runemark, A., & Brodin, A. (2010). A role

for learning in population divergence of mate preferences. *Evolution*, *64*(11), 3101–

3113. https://doi.org/10.1111/j.1558-5646.2010.01085.x

Svensson, E. I., Willink, B., Duryea, M. C., & Lancaster, L. T. (2020). Temperature drives

pre-reproductive selection and shapes the biogeography of a female polymorphism.

Ecology Letters, *23*(1), 149–159. https://doi.org/10.1111/ele.13417

Takahashi, Y., Kagawa, K., Svensson, E. I., & Kawata, M. (2014). Evolution of increased

phenotypic diversity enhances population performance by reducing sexual harassment

in damselflies. *Nature Communications*, *5*(1), Article 1.

https://doi.org/10.1038/ncomms5468

Takahashi, Y., & Watanabe, M. (2011). Male mate choice based on ontogenetic colour

changes of females in the damselfly *Ischnura senegalensis*. *Journal of Ethology*,

29(2), 293–299. https://doi.org/10.1007/s10164-010-0257-6

Terakita, A. (2005). The opsins. *Genome Biology*, *6*(3), 213. https://doi.org/10.1186/gb-2005-

6-3-213

van der Kooi, C. J., Stavenga, D. G., Arikawa, K., Belusic, G., & Kelber, A. (2021).

Evolution of insect colour vision – from spectral sensitivity to visual ecology. *Annual*

Review of Entomology, *66*. https://doi.org/DOI 10.1146/annurev-ento-061720-

071644.

Van Gossum, H., Bots, J., Van Heusden, J., Hammers, M., Huyghe, K., & Morehouse, N. I.

(2011). Reflectance spectra and mating patterns support intraspecific mimicry in the

colour polymorphic damselfly *Ischnura elegans*. *Evolutionary Ecology*, *25*(1), 139–

154. https://doi.org/10.1007/s10682-010-9388-z

Van Gossum, H., Stoks, R., & De Bruyn, L. (2001a). Discriminative mate choice in relation

with female maturation in *Ischnura elegans* (Odonata: Coenagrionidae). *International*

Journal of Odonatology, *4*(1), 83–91.

https://doi.org/10.1080/13887890.2001.9748161

Van Gossum, H., Stoks, R., & De Bruyn, L. (2001b). Reversible frequency–dependent

switches in male mate choice. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *268*(1462), 83–85. https://doi.org/10.1098/rspb.2000.1333

Van Gossum, H., Stoks, R., Matthysen, E., Valck, F., & De bruyn, L. (1999). Male choice for

female colour morphs in *Ischnura elegans* (Odonata, Coenagrionidae): Testing the

hypotheses. *Animal Behaviour*, *57*(6), 1229–1232.

https://doi.org/10.1006/anbe.1999.1100

Varma, N., Mutt, E., Mühle, J., Panneels, V., Terakita, A., Deupi, X., Nogly, P., Schertler, G.

F. X., & Lesca, E. (2019). Crystal structure of jumping spider rhodopsin-1 as a light

sensitive GPCR. *Proceedings of the National Academy of Sciences*, *116*(29), 14547–

14556. https://doi.org/10.1073/pnas.1902192116

Verzijden, M. N., ten Cate, C., Servedio, M. R., Kozak, G. M., Boughman, J. W., &

Svensson, E. I. (2012). The impact of learning on sexual selection and speciation.

Trends in Ecology & Evolution, *27*(9), 511–519.

https://doi.org/10.1016/j.tree.2012.05.007

Wakakuwa, M., Terakita, A., Koyanagi, M., Stavenga, D. G., Shichida, Y., & Arikawa, K.

(2010). Evolution and mechanism of spectral tuning of blue-absorbing visual

768   pigments in butterflies. *PLOS ONE*, *5*(11), e15015.

769   https://doi.org/10.1371/journal.pone.0015015

770 West-Eberhard, M. J. (2003). *Developmental Plasticity and Evolution*. Oxford University

771   Press.

772 Westerman, E. L., Hodgins-Davis, A., Dinwiddie, A., & Monteiro, A. (2012). Biased learning

773   affects mate choice in a butterfly. *Proceedings of the National Academy of Sciences*,

774   *109*(27), 10948–10953. https://doi.org/10.1073/pnas.1118378109

775 Willink, B., Duryea, M. C., Wheat, C., & Svensson, E. I. (2020). Changes in gene expression

776   during female reproductive development in a color polymorphic insect. *Evolution*,

777   *74*(6), 1063–1081. https://doi.org/10.1111/evo.13979

778 Willink, B., Duryea, M., & Svensson, E. (2019). Macroevolutionary origin and adaptive

779   function of a polymorphic female signal involved in sexual conflict. *The American*

780   *Naturalist*, *194*. https://doi.org/10.1086/705294

781 Willink, B., & Svensson, E. I. (2017). Intra- and intersexual differences in parasite resistance

782   and female fitness tolerance in a polymorphic insect. *Proceedings of the Royal Society*

783   *B: Biological Sciences*, *284*(1847), 20162407. https://doi.org/10.1098/rspb.2016.2407

784 Willink, B., Tunström, K., Nilén, S., Chikhi, R., Lemane, T., Takahashi, M., Takahashi, Y.,

785   Svensson, E. I., & Wheat, C. W. (2024). The genomics and evolution of inter-sexual

786   mimicry and female-limited polymorphisms in damselflies. *Nature Ecology &*

787   *Evolution*, *8*(1), 83–97. https://doi.org/10.1038/s41559-023-02243-1

788 Winfrey, C., & Fincke, O. M. (2017). Role of visual and non-visual cues in damselfly mate

789   recognition. *International Journal of Odonatology*, *20*(1), 43–52.

790   https://doi.org/10.1080/13887890.2017.1297259

791 Wright, D. S., van Eijk, R., Schuart, L., Seehausen, O., Groothuis, T. G. G., & Maan, M. E.

792   (2020). Testing sensory drive speciation in cichlid fish: Linking light conditions to

opsin expression, opsin genotype and female mate preference. *Journal of*

Evolutionary Biology, *33*(4), 422–434. https://doi.org/10.1111/jeb.13577

Yilmaz, A., Lindenberg, A., Albert, S., Grübel, K., Spaethe, J., Rössler, W., & Groh, C.

(2016). Age-related and light-induced plasticity in opsin gene expression and in

primary and secondary visual centers of the nectar-feeding ant *Camponotus rufipes*.

Developmental Neurobiology, *76*(9), 1041–1057. https://doi.org/10.1002/dneu.22374

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects*

models and extensions in ecology with R. Springer. https://doi.org/10.1007/978-0-387-

87458-6

TABLES & FIGURES

Table 1. Final model and model output for model 1, showing the effect of maturity stage and opsin type on log relative opsin expression in teneral, immature, and mature male *I. elegans*. Significant differences are shown in italics.

log Relative Opsin Expression ~ Maturity × Opsin + (1|ID) +(1|Replicate) + (1|Year)			
	df	F	*P*
Maturity	2	0.2	0.82
Opsin	7	292.3	*< 0.0001*
Maturity × Opsin	14	12.3	*< 0.0001*

Table 2. Final model and model output for model 2, showing the effect of maturity stage and opsin type on proportion opsin expression in teneral, immature, and mature male *I. elegans*. Significant differences are shown in italics.

Proportion Opsin Expression ~ Maturity × Opsin + (1|ID)			
	df	χ^2	*P*
Maturity	2	20.7	*< 0.0001*
Opsin	6	8877.3	*< 0.0001*
Maturity × Opsin	12	437.5	*< 0.0001*

Table 3. Final model and model output for model 3, showing the effect of the proportion of androchrome females in a population, opsin type, and maturity stage on log relative opsin expression for immature and mature male *I. elegans*. Significant differences are shown in italics.

log Relative Opsin Expression ~ Opsin × Maturity × (Prop. Androchrome + Prop. Androchrome^2) + (1|ID) + (1|Replicate) + (1|Year)			
	df	F	*P*
Opsin	7	18.7	*< 0.0001*
Maturity	1	1.3	0.27
Prop. Androchrome	1	0.2	0.66
Prop. Androchrome^2	1	0.2	0.67
Opsin × Maturity	7	3.1	*< 0.01*
Opsin × Prop. Androchrome	7	17.1	*< 0.0001*
Opsin × Prop. Androchrome^2	7	17.3	*< 0.0001*
Maturity × Prop. Androchrome	1	1.3	0.26
Maturity × Prop. Androchrome^2	1	1.3	0.25
Opsin × Maturity × Prop. Androchrome	7	3.3	*< 0.01*
Opsin × Maturity × Prop. Androchrome^2	7	3.6	*< 0.001*

Figure 1. Head and eyes, colour polymorphism, and female morph frequencies from eight populations in southern Sweden for the damselfly *Ischnura elegans*. (A) Head of male *I. elegans*, showing the large compound eyes and three dorsally situated ocelli. Box shows magnified view of the compound eye depicting individual ommatidial units. (B) Mature male *I. elegans* and the three female-limited colour morphs in their sexual mature colour phases. (C) Females undergo ontogenetic colour changes in the distal abdomen segments (left photo) and thorax (right photo) between immature and sexually mature adult stages. (D) Sampling locations of *I. elegans* used in the current study with inset showing the region sampled. (E) Female morph frequencies per site in 2021 and 2022. Numbers in parentheses above each bar indicate sample size. (B, C) modified from Willink et al. (2020). A = androchrome, I = *infuscans*, O = *infuscans-obsoleta*.

Figure 2. Functional expression of *Ischnura elegans* short-wavelength (SWS) and long-wavelength (LWS) opsins. Ultraviolet-visible (UV-VIS) absorbance analyses of dark-adapted rhodopsin visual pigments reconstituted and purified from HEK293T cell cultures in the dark in the presence of 11-*cis*-retinal. The black dots represent mean absorbances at a given wavelength. (A) SWb1 (n = 7), (B) SWb2 (n = 3), (C) LWA2 (n = 2), (D) LWF1 (n = 2), I LWF2 (n = 5), (F) LWF3 (n = 6), (G) LWF4 (n = 1), and opsins expressed in the ocelli: (H) LWE1 (n = 6), where n is the number of measurements of protein aliquots with active rhodopsin complexes. (I) Combined absorbances for SWS and LWS opsins, excluding LWE1, which is expressed in the ocelli in other odonates. Absorbance at 380 nm in A and B is due to residual unbound *cis*-retinal. Relative absorbance data are fitted to a visual template with polynomial function analyses computed in R to obtain the best estimates following 1000 bootstrap analysis of lambda max for each rhodopsin (Liénard et al., 2022). Confidence intervals are indicated in parentheses.

Figure 3. **Visual opsin sensitivity and female body reflectance.** Absorbance spectra of *I. elegans* SWS and LWS opsins characterized in this study are represented by coloured visual fit curves and overlapped with (A) androchrome and (B) *infuscans* female body reflectance (shaded gray areas). Spectral sensitivity functions correspond to differences in achromatic (brightness) and chromatic cues (body colour reflectance), assuming opponency between SWS and LWS opsins, of androchrome and *infuscans* females. SW1, SW2 and LWA labels correspond to SWb1, SWb2 and LWA2 opsins, respectively. LWE1, which is primarily expressed in the ocelli of other odonates, is not included. Body reflectance data from Henze et al. (2019).

Figure 4. Linear discriminate analysis showing the relationship between relative expression of eight opsins expressed in teneral, immature, and mature male *I. elegans*. The percent of variation described by each linear discriminate factors (LD1 and LD2) are shown in parenthesis. Distance between points across axes indicates differences in overall relative opsin expression between maturity stages, with results showing the largest differences between teneral and mature males, with immature males intermediate to the two groups.

Figure 5. Relative opsin expression for (A) teneral, (B) immature, and (C) mature male
I. elegans **across 8 sites**. Bars in box and whisker plots show medians, boxes indicate upper
and lower quartiles, whiskers show sample minima and maxima, and open circles show
outliers (representing 5.2 and 6.4 % for immature and mature treatments respectively).
Expression data are normalized against housekeeping genes and calibrated against UVS opsin
expression levels. Letters above each bar show significant differences between opsin types
within each maturity level following Tukey corrections for multiple comparisons. Test
statistics and p-values for all comparisons are presented in Table S5. Single population
analyses are presented in Fig. S2.

Figure 6. Relative male opsin expression across eight *I. elegans* populations for seven opsins expressed in the compound eye (SW1 – LWF4) and one low-expressed opsin likely expressed in the ocelli (LWE). Teneral males are shown in light blue, immature adult males in medium blue, and mature males in dark blue. Bars in box and whisker plots show medians, boxes indicate upper and lower quartiles, whiskers show sample minima and maxima, and open circles show outliers (representing 5.2 and 6.4 % for immature and mature treatments respectively). Expression data are normalized against housekeeping genes and calibrated against UVS opsin expression levels. Single asterisks indicate comparisons that are significant at alpha < 0.05 and double asterisks represent comparisons that are significant at alpha < 0.01 following Tukey corrections for multiple comparisons within opsin types. Test statistics and p-values for all comparisons are presented in Table S6; single population analyses are presented in Fig. S2.

Figure 7. Average proportion expression of visual opsins in teneral, immature, and mature male *I. elegans*. Proportion expression for 2 short-wavelength sensitive (SWS) and five long-wavelength sensitive (LWS) opsins.

Figure 8. Opsin expression changes relative to proportions of androchrome females. Log relative opsin expression in immature (open circles, dashed line) and mature (black triangle, solid line) male *I. elegans* from populations with different proportions of androchrome females. Grey shading around trend lines indicates 95% confidence intervals.