Opsin gene expression plasticity and spectral sensitivity as mechanisms for search image

formation in mate-searching male damselflies

Natalie S. Roberts[1], Erik I. Svensson[2], Marjorie A. Liénard[1]

[1]Department of Biology, Lund Vision Group, Lund University, Sölvegatan 35B, 223 62 Lund,

Sweden

[2]Department of Biology, Evolution Ecology Unit, Lund University, Sölvegatan 37, 223 62

Lund, Sweden

ABSTRACT

Odonata (damselflies and dragonflies) are a largely diurnal, colourful, and strongly visually

guided group of insects with visual systems influenced by many opsin genes that form the

primary light-sensitive eye photopigments. Heritable (genetic) female-limited colour

polymorphisms are also common in Odonata, with one female morph typically exhibiting

male-like (androchrome) colouration and one or two morphs exhibiting female-specific

colouration (gynochromes). In the Common Bluetail Damselfly (*Ischnura elegans*),

androchrome females express blue body colouration similar to that of mature males while

gynochrome females express either green or red-brown colouration. This and other colour

polymorphisms in damselflies are thought to be maintained by male mating harassment and

frequency-dependent sexual conflict, in which males form search images for certain morphs

which suffer disproportionally from mating harassment. Here, we investigate the role of opsin

sensitivity and expression plasticity in visual mate detection in *I. elegans* by quantifying

relative opsin mRNA expression over adult maturation in populations with different female

morph frequencies in southern Sweden. We find evidence for opsin-specific plasticity in

relative and proportion opsin expression, suggesting changes in opsin regulation and visual

sensitivity over adult maturation. Furthermore, the relative expression of the long-wavelength

sensitive opsin LWF2 changed in response to female morph frequencies. The highest relative

expression levels were found in populations with either a high or low proportion of

androchrome females. *In vitro* results indicate that long-wavelength sensitive opsins in *I.*

elegans provide a good visual match to the colouration of green gynomorph females and could

in principle confer male colour discrimination between female morphs. We discuss these

results in relation to frequency dependent selection, male sensory adaptations, plastic search

images and mate searching costs. We suggest that opsin gene expression could play an

important role in male search image formation of suitable mates.

INTRODUCTION

Opsins are G-protein coupled receptors that, when bound with a vitamin-A derived

chromophore, form a light sensitive visual pigment that is responsible for colour vision in

animals (Terakita, 2005). Many insects possess three classes of opsin proteins: ultraviolet

sensitive (UVS), short-wavelength sensitive (SWS), and long-wavelength sensitive (LWS) that

form visual pigments with maximum sensitivity to wavelengths (λ_{max}) around 350 nm, 440 nm,

and 530 nm, respectively (Briscoe & Chittka, 2001; van der Kooi et al., 2021). Differences in

spectral sensitivity are caused, in part, by changes in opsin expression, such as opsin gains or

losses, sequence changes, or expression level differences (Castiglione et al., 2023; Liénard et

al., 2021; Shand et al., 2008; Sharkey et al., 2023). In some cases, changes to the visual system

are also shaped by environmental factors, behaviour, and associated selection pressures. For

example, loss of functional opsin genes correlates with nocturnal lifestyles in mammals

(Jacobs, 2013), while opsin duplications in other species are found to contribute to the

regaining of spectral sensitivity (Rossetto et al., 2023; Sharkey et al., 2023) and, for several

insect species, functional changes amongst duplicated opsins likely improves detection and/or

discrimination of environmental and sexual signals (Liénard et al., 2021; McCulloch et al.,

2017, 2022).

In many cases, opsin gene expression is not fixed, but changes plastically over

individual ontogeny (Carleton et al., 2016; Chang et al., 2021; Yilmaz et al., 2016). Opsin gene

expression profiles also change in in response to environmental pressures (Fuller et al., 2010;

Shand et al., 2008; Wright et al., 2020). For example, changes in opsin expression is associated

with light exposure in honey bees and ants (Sasagawa et al., 2003; Yilmaz et al., 2016) and in

"choosy" relative to "non-choosy" females of the butterfly *Bicyclus anynana* (Everett et al.,

2012). In several species of damselfish (Pomacentridae), opsin gene expression plasticity tunes

visual sensitivity to local environmental lighting conditions for individuals inhabiting shallow

(< 4 m) versus deep (> 10 m) water (Stieb et al., 2016).

Odonata (damselflies and dragonflies) is an insect order with many colourful species

and these insects are also characterized by their large and complex eyes (Corbet, 1999;

Córdoba-Aguilar, 2023). Odonates inhabit a wide array of visual environments, having both

aquatic and terrestrial life stages, and they use vision for a variety of behaviours, including

prey capture and mate detection (Combes et al., 2013; Corbet, 1999; Córdoba-Aguilar, 2023).

These features make these insects excellent model organisms to study evolutionary and

functional aspects of colour vision. Like many other insects, odonates express UVS, SWS, and

LWS opsins in their eyes; however, duplications of the SWS and LWS opsin classes have

resulted in a remarkable genetic diversity, with between 11 – 30 visual opsin genes identified

across several families (Futahashi et al., 2015). Recordings of visual sensitivity in the eyes of

some species indicate spectral sensitivity ranging from UV to red light (Horridge, 1969; Huang

et al., 2014; Meinertzhagen et al., 1983; Yang & Osorio, 1991). While it is not clear how their

abundant opsin copies function in vision, visual stimuli are clearly important for odonate

behaviour (Combes et al., 2013; Corbet, 1999; Miller & Fincke, 1999). For example,

comparisons of visual and olfactory cues suggest that vision is primarily used in mate choice

in several species of damselflies (Rebora et al., 2018; Winfrey & Fincke, 2017). Colouration

is also an important signal indicating female sexual maturity (Bybee et al., 2012; Gosden &

Svensson, 2009; Huang et al., 2014; Svensson et al., 2020; Takahashi & Watanabe, 2011; Van

Gossum et al., 2011; Willink et al., 2019, 2020). Males can discriminate between immature

and mature females, and they direct more mating attempts to mature than immature females

(Huang et al., 2014; Van Gossum et al., 2001a, 2011; Willink et al., 2019). Experimental

manipulation of age-specific colouration in *Ischnura* damselflies also suggest that vision is the

primary cue in males discrimination between immature and mature females (Willink et al.,

2019).

In addition to colour changes associated with sexual maturation, many odonates also

exhibit heritable (genetic) female-limited colour polymorphisms, with one female morph

having male-like colouration (androchrome females) and one or two morphs expressing

female-specific colouration (gynochrome females) (Blow et al., 2021; Fincke et al., 2005).

Evidence suggests that female colour polymorphisms function to reduce the negative fitness

effects of excessive male sexual harassment by disrupting males' ability to efficiently form

search images for mates (Fincke, 2004; Miller & Fincke, 1999; Takahashi et al., 2014).

According to the male mimicry hypothesis, androchrome females have a negative frequency-

dependent advantage in their male-like similarity, which reduces male mating harassment

(Robertson, 1985). Alternatively, although not mutually exclusive, according to the learned

mate recognition (LMR) hypothesis, males form search images for the most common female

morph in the population (Fincke, 2004). The LMR hypothesis further predicts that the ratio of

correct (e.g., mature conspecific females) versus incorrect (e.g., other males, heterospecifics)

identification of mates should increase with experience. There is partial empirical support for

both the male mimicry hypothesis and LMR and, as stated above, these two hypotheses are not

mutually exclusive (Gosden & Svensson, 2009; Svensson et al., 2005, 2020). Empirical

evidence has shown that males show increased preferences for the most abundant female

morph in their population (Takahashi & Watanabe, 2010) or increased preference following

exposure to specific female morphs (Miller & Fincke, 1999; Van Gossum et al., 2001b; but see

Van Gossum et al. 2011). Moreover, male mate preferences become increasingly non-random

with increasing density of certain morphs (Gosden & Svensson, 2009). Female fecundity is

also higher in populations with more balanced morph frequencies, presumably because females

are then expected to suffer from less per-capita harassment than females in populations with

biased morph frequencies as, in the latter case, males should more easily form search images

for a single morph (Takahashi et al., 2014). In androchrome-biased populations, there is also

an increasing number of male-male interactions, consistent with mistaken mate recognition due

to male mimicry and the formation of male search images for the common female phenotype

(Gering, 2017). Finally, analyses of time-series of morph frequency fluctuations and population

genetic modeling have revealed a clear signature of negative frequency-dependent selection,

where the fitness of a given female morph decreases with its frequency in the population (Le

Rouzic et al., 2015; Svensson et al., 2005; Takahashi et al., 2010).

This growing body of ecological, evolutionary, sensory, and physiological studies,

paired with the increasing availability of genomic resources, makes Odonata an excellent

system to investigate the molecular basis of colour vision. Here, we characterize opsin spectral

sensitivity via heterologous expression and quantify variation in opsin mRNA expression

across adult male maturation from populations with variable female morph frequencies in the

Common Bluetail Damselfly (*Ischnura elegans*). Functional characterization of 2 SWS and 5

LWS rhodopsins found in the ventral eye show absorbance peaks (λ_{max}) in the range of 406 –

419 nm and 531 – 548 nm, respectively. We also demonstrate opsin expression changes over

adult male development, with increased expression of specific opsin types in response to

changes in local female morph frequencies. These novel results suggest that the visual system

of male *I. elegans* can rapidly and plastically respond to changes in local morph frequencies

that, alongside opsin sensitivities, may aid in mate detection and discrimination. Our results

further suggest that opsin expression plasticity may provide a mechanistic proximate link

between male mate preferences, mating harassment, and the resulting frequency-dependent

sexual conflict, consistent with models for how these polymorphisms are maintained (Fincke,

2004; Le Rouzic et al., 2015; Svensson et al., 2005). Plastic male opsin expression profiles

could therefore help to regulate colour vision and perception of female colour signals in this

dynamic system strongly shaped by visually guided male mating behaviours.

MATERIALS AND METHODS

<u>Study system</u>

The Common Bluetail Damselfly (*Ischnura elegans*) is found broadly across Europe,

with the northern end of its range limit in Central Sweden (Dijkstra & Lewington, 2006;

Dudaniec et al., 2018). Sexually mature male *I. elegans* are monomorphic, exhibiting blue body

colouration on the thorax. In contrast, female *I. elegans* are polymorphic, with sexually mature

females belonging to three distinct genetically determined colour morphs (Fig. 1A) (Willink et

al., 2020). Androchrome (male-like) females exhibit body colouration spectrally similar to

male colouration while gynochrome females exhibit spectrally distinct green (infuscans) or

brown (infuscans-obsoleta; hereafter referred to as obsoleta) thorax colouration (Henze et al.,

2019; Van Gossum et al., 2011). Females exhibit ontogenetic changes in thorax and abdomen

colouration (Fig. 1B), with male *I. elegans* preferring body colouration of mature over

immature females (Willink et al., 2019). Apart from colour differences, androchrome and

gynochrome females also differ in mating rates, parasite resistance, and tolerance to parasites

(Willink & Svensson, 2017) as well as resistance to mating attempts, and fecundity (Gosden &

Svensson, 2009).

<u>Quantification of female morph frequencies between sites and field collection of males</u>

We quantified female morph frequencies from eight sites in southern Sweden (Fig. 1C)

as part of an ongoing long-term field population study of *I. elegans* (see Le Rouzic et al. 2015

and Svensson et al. 2020 for general methodology and details about this long-term study). All

study populations were visited and re-visited at regular intervals (1-2 weeks) between 19 May

– 1 August 2021 and 16 May – 31 July 2022. Taking into account both immature and mature

females, all populations were trimorphic except for one (Höje å 6) in 2022, which had

experienced large overall population declines that year (Table S1). However, since male *I.*

elegans are known to prefer colouration of mature over immature females (Willink et al.,

2019), we only considered females that displayed mature adult colouration to calculate the

relative abundance of each morph within the current study. The frequencies of the three mature

female morphs differed between sites and years (Fig. 1D-E). Androchrome females comprised

the majority of females across all sites, ranging from 57.1– 85.7% (mean ± sd: 72.6 ± 9.7%) of

females in 2021 and 65.9 – 83.9% (72.3 ± 5.8%) in 2022. Infuscans females were the next most

common morph, ranging from 14.3 – 39.1% (24.1 ± 8.8%) of females in 2021 and 16.1 – 30.7

(24.9 ± 5.3%) in 2022. Mature obsoleta females were rare, making up 0.00 – 5.5% (3.3 ± 1.8%)

of females in 2021 and 0.00 – 7.8% (2.8 ± 2.8%) in 2022.

Male *I. elegans* used for opsin quantification were caught from these populations

between 28 June – 13 July 2021 and 9 June – 20 July 2022 using sweep nets (Table S2). For

all collected individuals, we recorded sexual maturity ("immature" vs. "mature") by evaluating

wing stiffness (Corbet, 1999). We also recorded whether males were found in copula with a

female or singly ("couple" vs "single") and, in 2022, we recorded the morph of the female that

was found in copula with each collected male. In addition to immature and mature males, we

also collected males that had newly emerged from the aquatic nymph stage ("tenerals") from

two field sites and from a semi-naturalistic mesocosm experiment at the Stensoffa Field Station

at Lund University (Table S2). Teneral males are characterized by their extremely soft wing

and body tissues and lack of body pigment. These teneral males allowed us to assess opsin

gene expression in males that have not had any significant visual experience and have not

mated with sexually mature females. All collected males were immediately euthanized by

cutting off their head using cleaned RNAse-free dissection tools, placed into individual vials

filled with RNA*later* (Ambion, Inc., Austin, TX, USA) at field sites, and then stored at -20°C

until RNA extraction.

Opsin heterologous expression

We retrieved visual opsin open reading frame (ORF) sequences for *I. elegans* (UV,

SWb1, SWb2, LWA1, LWA2, LWC1, LWD1, LWE1, LWF1, LWF2, LWF3, and LWF 4, see

Supplementary File S1) by mining whole genome data from the genome assembly

iolscEleg1.1. using complete sequences previously reported from the closely related species

Ischnura asiatica (Futahashi et al., 2015) as our reference set. Opsin transcript and protein

sequences were predicted with AUGUSTUS (Keller et al., 2011) followed by sequence

annotation in HMMER v3.1b2 (http://hmmer.org). Opsin ORFs flanked with suitable

restriction sites for subcloning in the pcDNA5-FLAG-T2A-mruby2 expression cassette were

synthesized for human codon optimization, overexpressed by transient transfection following

the PaSHE procedure (Liénard et al., 2021, 2022) followed by purification and ultraviolet-

visible spectroscopy. Briefly, we first performed small scale transfections to verify that each

final pcDNA5-Opsin construct produced detectable protein levels (Fig. S1). For this,

HEK293T cells were seeded at a density of 0.6×10^6 cells in DMEM medium (Gibco) in 6-

well plates. The next day cells were transfected with a mixture of 4 ug plasmid DNA and a 1:3 ratio of PEI (1mg/mL) in Opti-Mem (Gibco). Forty-eight hours later, cells were visualized under a fluorescent microscope to verify expression of mRuby2 and then harvested in cold D-PBS (Sigma-Aldrich). Cells were resuspended in cold Ripa buffer supplemented with 1% n-Dodecyl-ß-D-maltoside and complete ethylenediaminetetraacetic acid (EDTA)-free protein inhibitors (Sigma-Aldrich), and incubated at 4°C for 1 hour, followed by western blot analysis.

For large scale opsin purifications, fifteen HEK293T cell dishes containing 2×10^6 cells were transfected with 24 ug DNA and 72 uL PEI each. Five micromolar 11,*cis*-retinal was added 6hr post-transfection with fresh culture medium. Cells were collected under dim red light 48h post-transfection and decanted in cold Hepes wash buffer (3 mM MgCl2, 140 mM NaCl, 50 mM Hepes, EDTA-free protein inhibitors) followed by incubation at 4°C for 1 hour in presence of 40 µM 11,*cis*-retinal. Cell pellets were collected by ultracentrifugation and membrane proteins were solubilized for 1hr at 4°C in ice-cold extraction buffer (3 mM MgCl2, 140 mM NaCl, 50 mM Hepes, 20% glycerol vol/vol, 1% n-dodecyl β-D- maltoside, complete EDTA-free protein inhibitors) and ultracentrifuged. The crude extract (supernatant) was incubated in the dark with FLAG-resin on a rotator at 10 rpm until the next day. The crude extract containing resin-bound FLAG-epitope rhodopsin complexes was loaded onto a Pierce centrifuge purification column, washed with 3x 3 mL wash buffer (3 mM MgCl2, 140 mM NaCl, 50 mM Hepes, 20% glycerol vol/vol, 0.1% n-dodecyl β-D- maltoside) and eluted using FLAG-peptide. The eluate was concentrated to 100-120 microliters using an Amicon Ultra-2 10 kDa Amicon centrifugal filter for 50 min at 4°C and UV-visible absorption spectra (200 to 800 nm) of dark-adapted purified proteins were measured in the dark from 1.5-µL aliquots using a NanoDrop 2000/2000c UV-VIS spectrophotometer (Thermo Fisher). For each rhodopsin, we obtained estimates of lambda max through nonlinear least-square fitting to the absorbance data according to a visual template (Govardovskii et al., 2000) and performed 1000

bootstrap replication to calculate lambda max predictions and confidence intervals (Liénard et

al., 2022) in R v.3.6.6.(*R Core Team*, 2021) using the packages rsample and tidymodels (Frick

et al., 2022; Kuhn & Wickham, 2020).

<u>Quantitative PCR analysis of *I. elegans* opsin expression levels</u>

We extracted RNA from the heads of N = 87 male *I. elegans* (Table S2) using a

Monarch Total RNA Miniprep Kit protocol (Thermo Fisher Scientific, Waltham, MA, USA)

following the manufacturer's instructions and including a genomic DNA clean-up. cDNA was

synthesized for each sample using a GoScript™ Reverse Transcriptase kit (Promega, Madison,

WI, USA). The *Ischnura*-specific RL14 and GADPH gene sequences were annotated as

described above for opsin sequences and used as reference for mRNA quantification.

For quantitative PCR analysis, two sets of gene-specific primers were designed in

Geneious Prime v.2022.91 (Biomatters, Auckland, New Zealand) for both housekeeping genes

and all opsins expressed in the adult ventral eye (UV, SWb1, SWb2, LWA2, LWF1 – F4) and

the ocelli opsins (LWD1 and LWE1). We did not include LWA1 or LWC1 since these opsin

types are primarily expressed in larval tissues in other Odonate species (Futahashi et al., 2015).

We tested the efficiency of designed primers by generating a standard curve following a 4-fold

dilution of *I. elegans* cDNA and selected one primer pair for each opsin based on amplification

efficiency (Table S3). For the ocelli-specific opsins, we focused further analysis on the LWE1

opsin due to more consistent amplification rates relative to LWD1. The threshold detection

cycle (Ct) values of selected primers were plotted against the log of cDNA dilutions and the

efficiency percentage values were assessed to ensure that they were within acceptable ranges

(94 – 117%) and R^2 (0.95 – 1.00). PCR amplicons were run by electrophoresis on a 2% agarose

gel to ensure the absence of primer-dimers and PCR samples for each primer pair-opsin target

were purified using Exonuclease (EXO) and Shrimp alkaline phosphatase (SAP) (NEB) prior

to Sanger Sequencing to confirm gene-specific amplification. Hereafter, we use abbreviated

names for the opsins SWb1, SWb2, LWA2, and LWE1, which have been shortened to SW1,

SW2, LWA, and LWE, respectively.

qPCR was performed in a CFX384 Real-Time System (BioRad Laboratories, Inc.,

Hercules, CA, USA). Reaction mixes were prepared that contained 1x SYBR Green, 1uM

primer mix, and 0.1 ng cDNA. The program was designed as follows: denaturation at 95°C for

2min, followed by 39 cycles at 95°C for 10s and 60°C for 10s, with melting curve analysis

performed from 65.5°C to 89.5°C at 0.6°C increments to confirm the specificity of PCR

products. Two independent technical replicates were processed for each sample and each

sample was run in triplicate within each technical replicate. We averaged triplicates within

each technical replicate for a total of n = 2 Ct values per male sample.

We calculated expression of opsin genes relative to the housekeeping genes and

normalized against the UV opsin gene using the $(1 + E)^{-\Delta\Delta CT}$ method where E equals the PCR

efficiency for each primer (Livak & Schmittgen, 2001; Pfaffl, 2001). We also calculated the

proportion opsin expression relative to total opsin expression for opsins by dividing relative

expression for each individual opsin by the sum of total opsin expression, excluding the ocelli

opsin LWE. Results of relative opsin expression indicated differential regulation of opsin

genes, while proportional measures of opsin expression are suggested to be better for making

inferences about colour vision (Fuller & Claricoates, 2011). For proportion opsin expression,

we focused only on opsins expressed in the compound eye since this is where colour

discrimination primarily occurs within the insect visual system.

Statistical analysis

We performed a Multivariate Analysis of Variance (MANOVA) to assess the effect of

maturity on log relative opsin gene expression of all eight opsin genes for male *I. elegans*. Log

relative opsin expression for each individual male was averaged between replicates to maintain

the assumption of independence. We assessed the dataset for outliers and compared result

outputs performed before and after their removal to determine whether this affected results.

The inclusion of outliers did not qualitatively change the general results, and we therefore

report results with outliers included. Further analysis to account for repeated measures on the

full dataset were performed using linear mixed modelling (see below).

General protocol for model selection

We used linear mixed modeling using the lme4 package (Bates et al., 2015) in R v 4.1.1

to test the effects of population morph frequency and its squared component, maturity level,

and opsin type on log relative opsin expression in male *I. elegans*, with site, male ID, year, and

qPCR replicate as random effects. Model selection followed methods in Zuur (Zuur et al.,

2009), in which we first selected the random effect structure for the model by comparing

Akaike information criterions (AICs) between models fitted using restricted maximum

likelihood (REML) estimation that included all possible fixed effects and relevant interactions.

Following selection of the random effect structure, we used backward model selection with

maximum likelihood (ML) estimation to determine the optimal fixed effect structure by testing

the significance of each interaction or term. After selection of the optimal random and fixed

effect structures, the final model was run using REML estimation and the model output and

results of post-hoc analysis were obtained using the lmerTest and emmeans packages

(Kuznetsova et al., 2017; Lenth, 2022), respectively.

Opsin mRNA expression in teneral, immature, and mature males across sites

We tested the effect of maturity stage (factor with 3 levels: teneral, immature, mature),

opsin type (factor with 8 levels: SW1, SW2, LWA, LWF1 – LWF4, LWE), and their interaction

on log relative opsin expression. Model selection indicated a significant interaction between

maturity level and opsin type (likelihood ratio test: χ^2 (14) = 163.9, *p* < 0.0001). The final

model therefore included year, site, and male ID as random effects with fixed effects of

maturity stage, opsin type, and their interaction to explain log relative opsin expression in male

I. elegans.

To investigate how proportional opsin gene expression changes over development, we

again used maturity stage, opsin type, and their interaction as fixed effects, but now only

focused on opsins found in the compound eye (factor with 7 levels: SW1, SW2, LWA, LWF1

– LWF4). Data exploration suggested that opsin type violated the homogeneity of variance

assumption for this dataset. We therefore allowed for differing variance structure in the term

opsin using the varIdent function in the nlme package (Pinheiro et al., 2021). We included only

male ID as a random effect since inclusion of additional random effects did not improve model

AIC. Model selection showed a significant interaction between opsin type and maturity stage

(likelihood ratio test: χ^2 (12) = 268.7, *p* < 0.0001), making the final model the interaction of

maturity stage and opsin type on proportion opsin expression.

Preliminary analysis suggested differences in relative opsin expression between teneral

males from the two field populations and those from outdoor mesocosm tanks at Stensoffa

Field Station (Table S4). Despite these differences, the results with and without males from the

mesocosm tanks were largely comparable. We therefore retained the teneral males from the

mesocosm tanks in our main analyses. Results of log relative expression and proportion opsin

expression without the mesocosm males are presented in Supplementary Material (Tables S5-

S7, Fig. S2).

Opsin mRNA expression in populations with different female morph structure

To assess the effect of local female morph frequency on male opsin expression, we used

the frequency of androchrome females for each site and year to describe local population morph

frequency. We used the proportion of androchrome females within a site for each year

separately, instead of generating a site average, since a primary aim of the current study was to

assess if opsin expression changes in response to current local conditions. Therefore,

"population morph frequency" refers to the local morph frequency in a given year, with two

population morph frequencies per site.

We followed the same modeling procedure to test the effect of maturity stage, opsin

type, the proportion of androchrome females, and their interactions on relative opsin

expression. We included the proportion of androchrome females also as a quadratic term to

account for any non-linear effect of androchrome frequencies. In this analysis, we focused only

on immature and mature males since data for teneral males was not available for most sites.

Backward model selection of fixed effect structure indicated that the three-way interaction

between opsin type, maturity stage, and androchrome proportion was significant (likelihood

ratio test: $\chi 2$ (14) = 43.9, $p < 0.0001$). The final model therefore accounted for a quadratic

relationship between androchrome proportion, opsin type, maturity stage, and their interaction

as fixed effects with year, male ID, and replicate as random effects.

Opsin expression in response to mating outcomes

We recorded the female morph of all males collected in copula in 2022 to assess

whether relative opsin expression differed between males collected in copula with

androchrome or infuscans females. The selected random effect structure included male ID and

replicate. We considered maturity level, opsin type, and female morph (factor with 2 levels:

androchrome, infuscans), and their interactions as fixed effects. Backwards model selection

resulted in a final model with opsin as the only fixed effect, suggesting that there is no

significant effect of female morph type on a males' relative opsin expression in the current

study. However, it is possible this is due to relatively small sample size, with n = 8 males

collected in copula with an androchrome female and n = 7 males collected in copula with an

infuscans female.

RESULTS

Opsin functional expression and spectral sensitivity

To determine the spectral sensitivity of opsins expressed in the ventral eye and ocelli,

we first subcloned codon-optimized individual opsin open reading frames for two short-

wavelength opsins (SW1 and SW2), as well as seven long-wavelength opsins including five

expressed in the ventral eye (LWA, LWF1-F4) and two ocelli-specific opsins (LWE and LWD)

(Futahashi et al., 2015) in a recently developed expression cassette (Liénard et al., 2021). Small

scale assays were performed to verify transcriptional activity via fluorescent microscopy of a

co-transcribed mRuby marker, and via western-blot visualization (Fig. S1). We then transiently

expressed each opsin construct in large scale HEK293T cell cultures. Active rhodopsin

complexes were reconstituted by addition of 11, *cis*-retinal in the dark and purified to obtain

dark adapted rhodopsin eluates prior to ultraviolet-visible spectroscopy analyses. First, we

found that the SWS opsins SWb1 and SWb2 absorb maximally at λ_{max} of 406 and 419 nm (Fig.

2A-B). We then assayed the seven LWS opsins and measured their individual absorbance

spectrum (Fig. 2C-I). The LW opsin types expressed in the ventral eye display maximal

sensitivity to approximately 530 – 550 nm with the following λ_{max}: LWA2 = 548 nm, LWF1 =

531 nm, LWF2 = 543 nm, LWF3 = 545 nm, and LWF4 = 541 nm (Fig. 2C-G). Finally, we find

that the ocelli-specific opsins, LWD1 and LWE1 absorb maximally at 541 nm and 533 nm,

respectively (Fig. 2H-I).

We ran homology modeling of SWb1 against the spider rhodopsin (PDB: 6i9k) and

mapped sites interacting with the chromophore in Pymol. Of the 71 amino acid substitutions

between SWb1 and SWb2 (81.8% aa identity), we identified 23 sites predicted to be within 5Å

of the binding pocket (Table S8). Among these, Y136 (TM3) and Y294 (TM6) residues located

at the top and bottom of the binding pocket, respectively, are substituted by phenylalanine

(F136 and F294) in SW2 (Fig. S3A). For LWS opsins, which share between 88.9 - 91.2% aa

identity (i.e., 33 - 42 variant residues), we obtained the predicted LWF1 opsin structure based

on 6i9k and mapped 24 sites predicted to interact with the cis-retinal, all of which are conserved

for all four LWF opsins (Table S8; Fig. S3B). LWA2 is more divergent in sequence, 67.5 -

68.7% aa identity with 119 - 124 aa differences with LWF opsins. Of the 24 chromophore-

interacting residues, its binding pocket exhibits two potential spectral variant residues with

LWF1 (A131G in TM1 and S319A in TM7).

Finally, sensitivity of the SWS and LWS opsins expressed in the ventral compound

provides sensitivity to body reflectance of androchrome and infuscans females (Fig. 3).

Particularly, sensitivity of the LWA and LWF2 – LWF4 opsins provides a close

correspondence to infuscans female thorax colouration (Fig. 3B).

Opsin expression in teneral, immature, and mature males

Results from the MANOVA show a significant effect of male maturity stage on overall

opsin gene expression ($F_{16,146} = 3.2$, $p < 0.0001$; Fig. 4). To further assess how relative and

proportion opsin expression differed between teneral, immature, and mature male *I. elegans*,

we performed mixed modeling with maturity stage, opsin type, and their interaction as main

effects.

There was a significant effect of opsin type and maturity stage × opsin interaction on

log relative opsin expression in male *I. elegans* (Table 1, Tables S9-S10). For teneral males,

relative opsin expression was lowest for the SW1, SW2, and the ocelli LWE opsin, increased

in LWA – LWF3 opsins, and was significantly highest for LWF4 (Fig. 5A). In immature males,

the ocelli specific LWE opsin had significantly lower relative expression than the opsins found

within the compound eye, as expected. Among visual opsins, SW2 and LWF1 opsins had the

lowest relative expression, followed by the SW1 and LWF3 opsins, then LWA, LWF4, and

finally, LWF2 which was expressed at significantly higher levels (Fig. 5B). In mature males,

relative expression of LWE is also significantly lower than the opsins within the compound

eye, followed by increasing expression for LWF1, SW2, SW1 and LWF3, LWA, LWF4 and

LWF2 (Fig. 5C).

We next compared relative opsin expression between teneral, immature, and mature

males, pooling sites, to assess whether developmental changes affect relative opsin expression

uniformly or whether there is an opsin-specific effect on relative expression, suggestive of

differential plasticity in opsin expression. We found that relative opsin expression changed

over sexual maturation in an opsin-specific manner (Fig. 6). There was a significant increase

in relative expression of the SW1 opsin between teneral and mature males (teneral – mature: t

= -2.7, df = 396, p = 0.02) and between all maturity categories for relative LWF2 expression

(teneral – immature: t = -5.1 df = 401, p < 0.0001; teneral – mature: t = -6.9, df = 396, p <

0.0001; immature – mature: t = -3.0, df = 419, p = 0.009). In contrast, we saw decreases in

opsin expression over maturity between all maturity categories for relative LWF1 expression

(teneral – immature: t = 4.7, df = 402, p < 0.0001; teneral – mature: t = 6.1, df = 399, p <

0.0001; immature – mature: t = 2.4, df = 433, p = 0.04) and between teneral males with

immature and mature adult males for relative LWE expression (teneral – immature: t = 3.4, df

= 401, p = 0.002; teneral – mature: t = 3.9, df = 400, p = 0.0004).

To decouple the effect of LWF1 and LWF2 per site and assess the variation in the

dataset, we further assessed differences over maturity separately for each site (Table S11, Fig.

S4). While there were some sites that did not show a significant difference in relative

expression over maturation, three sites, Bunkeflostrand, Ilstorp, and Flackarp showed a

significant interaction between maturity stage and opsins found in the ventral eye. Differences

between this analysis and the large comparative metapopulation analysis likely arise due to

reduced sample size in each site, but nevertheless highlight consistent results for three

independent sites, corroborating the robustness of the pooled analysis that supports specific

opsin expression changes over maturity.

Comparing proportional opsin expression levels, we again observe a significant

interaction between maturity stage and opsin type (Table 2). For teneral males, LWF4

expression accounts for over half of opsin expression in the compound eye (0.523 ± 0.01) while

in immature and mature males, LWF2 expression makes up the largest proportion of

expression, accounting for 0.524 ± 0.03 of expression in immature and 0.599 ± 0.02 of

expression in mature males (Fig. 7). Post-hoc analysis show that proportional expression levels

differed significantly between all LWS opsins for teneral and immature males and between all

LWS and the SW2 opsin for teneral and mature males. There were also significant differences

between immature and mature males in proportional expression of the SW2 and LWF1 opsins

(Table S12).

Opsin expression in populations with different female morph frequencies

We performed mixed modeling to assess whether relative opsin expression differs

between immature and mature male *I. elegans* that had experienced different population morph

frequencies (here measured as proportion of androchrome females). There was a significant

effects of opsin type, maturity stage, frequency of androchrome females and their interaction

on log relative opsin expression (Table 3). For immature males, the proportion of androchrome

females in a population had a relative weak linear effect on relative opsin expression. However,

in mature males, there was a significant quadratic relationship of the proportion of

androchrome females on relative LWF2 expression. The highest LWF2 expression in mature

males were found in the populations with the lowest and highest proportion of androchrome

females (Fig. 8).

DISCUSSION

Vision is an important sensory modality in Odonata, functioning in a wide array of

behaviours ranging from flight stabilization to mate choice (Rebora et al., 2018; Wang et al.,

2013; Winfrey & Fincke, 2017). The prevalence of female-limited colour polymorphisms in

Odonata have made these insects powerful model systems to study how and why

polymorphisms emerge and persist over micro- and macroevolutionary time scales (Blow et

al., 2021; Le Rouzic et al., 2015; Willink et al., in press). Here, we combined ecological,

molecular, and functional approaches to investigate the peripheral visual system in male *I.*

elegans of different maturation stages. Specifically, we investigated the role of opsin

sensitivities, opsin gene regulation, and proportional opsin expression over male ontogeny and

across populations with different female morph frequencies. Our results suggest that the

peripheral visual system of *I. elegans* is sensitive to body colouration of the two most abundant

female morphs from our sampled sites (Fig. 3). Further, plastic changes in opsin expression

over male ontogeny, particularly during sexual maturation (Figs. 4 - 7), as well as changing

male opsin expression profiles in response to local female morph frequencies (Fig. 8) suggest

a role of opsin expression in male mate detection and/or mate preference. Taken together, these

novel results suggest that tuning of the peripheral visual system in response to local morph

abundances could provide a mechanistic link between male search image formation and the

negative-frequency dependent selection that maintains female polymorphism within the tested

sites (Fincke, 2004; Le Rouzic et al., 2015; Svensson et al., 2005).

Overlapping opsin spectral sensitivities confer broad colour vision in the ventral eye

The many opsin duplications present in Odonata are fairly unique among insects. Other insect species typically express only three – four opsins, while over 30 visual opsins genes have been identified in some species of dragonflies and 12 visual opsin genes have been identified from the damselfly *Ischnura asiatica* (Futahashi et al., 2015). Here, we confirm the expression of 10 opsin genes that have been predicted to be expressed in adult ventral eyes or ocelli from the heads in adult male *I. elegans*. These opsins include 1 UVS, 2 SWS, and 5 LWS opsins that are primarily expressed in the compound eye and 2 LWS opsins that are primarily expressed in ocelli. Our functional analyses of reconstituted active rhodopsin pigments followed by visual template fit for all five ventrally expressed LWS opsins show broad sensitivity to green wavelengths of light, produced from opsins with λ_{max} between 530 nm and 540 – 550 nm (Fig. 2). The observed overlap in long-wavelength sensitivities provides the ventral eye in principle with spectral discrimination capacity ranging from 450 nm to 640 nm (Figs. 2 - 3). In addition, functional data of SWS opsins shows two peaks of sensitivity in blue wavelengths of light, with λ_{max} of 406 and 419 nm, which in principle allows *I. elegans* damselflies to readily distinguish wavelengths in the ultraviolet – blue spectrum from 350 nm to circa 490 nm (Figs. 2 - 3). These results demonstrate that spectral sensitivities derived from the multiple SWS and LWS opsin types known to be expressed in the ventral eye of Odonata, and in particular *Ischnura* species, allows for capture of a remarkably broad light spectrum from short ultraviolet to long-wavelengths above 650 nm.

Given our findings of developmental regulation of opsin expression and overlapping spectral sensitivities of LWS opsins expressed in the ventral eye, and previously identified differences in expression patterns between the dorsal and ventral eye in several Odonate species (Futahashi et al., 2015; Labhart & Nilsson, 1995), it is likely that the multiple tandem

LWS duplicate opsins in *I. elegans* exhibit some degree of spatial regionalization in expression

patterns throughout the retina. Regional retinal patterning and spectral mosaics occur in a

number of insects (Rister & Desplan, 2011) and opsin co-expression in other insects are

hypothesized to provide broadband spectral sensitivities (reviewed in Arikawa, 2017; van der

Kooi et al., 2021; Wernet et al., 2015). In Odonata, the eyes of the dragonfly *Hemicordulia*

tau have three short-wavelength sensitive zones (i.e., the dorsal acute zone, the dorsal rim area

and the frontal acute zone) and display ventral spectral band patterning maximally sensitive to

either short-, middle-, or long-wavelengths (Lancer et al., 2020). Electroretinogram (ERG)

recordings in the ventral eye region showed that the dragonfly *Sympetrum frequens* exhibit a

wide variety of sensitivities from UV to red (Futahashi et al., 2015). Similarly, early ERG

recordings from *Ischnura heterosticta* show UV and blue-sensitive photoreceptor types with

peak sensitivities (λ_{max}) at 360 nm and 450 nm, respectively, and a green-sensitive

photoreceptor type (λ_{max} = 525 nm) with broad sensitivity to light between 520 – 540 nm and

decreasing sensitivity up to 650 nm (Huang et al., 2014). It is therefore also possible that the

broad sensitivity to green and red light recorded from other odonate species results from several

ommatidial types expressing combinations of LWS opsins. Analysis of opsin expression via

gene-specific antibody staining or *in situ* hybridization, along with *in vivo* ERG quantification

of visual sensitivity, will be valuable next steps to determine how opsin spatial distribution and

co-expression patterns contribute to overall visual spectral sensitivity in *I. elegans.*

Finally, we mapped two strong candidate spectral tuning sites at positions 177 and 300

between *I. elegans* duplicate SWb1 (λ_{max} 406 nm) and b2 (λ_{max} 419 nm) opsins (Fig. S3). A

comparison of duplicate SWb opsins across five *Ischnura* species, *I. cervula*, *I. verticalis*, *I.*

hastata and *I. asiatica,* shows that Y294F (H6) is present only in *I. cervula* whereas Y136F

(H3) is conserved across all species. Although we have not tested the effects on these mutations

in the present study, repeated homologous Y/F changes have been functionally shown to cause

convergent spectral shifts in SWS insect opsins, including a 5nm shift to longer wavelengths caused by Y195F in *Limenitis* Blue butterfly opsins (Frentiu et al., 2015), a 4nm shift towards longer wavelengths caused by Y177F between *Pieris* violet and blue opsins (Wakakuwa et al., 2010), a 25nm shift towards long wavelength (Y177F) in combination with A116S in *Eumaeus* blue opsin duplicates (Liénard et al., 2021), and a 3-nm shift to shorter wavelengths involving the reverse substitution F285Y in bruprestid beetle UV opsins (Sharkey et al., 2023). Accumulated evidence for the role of Y to F substitutions in providing longer wavelength-sensitive SWS invertebrate opsins is consistent with the gain of F136 in *I. elegans* SWb2, possibly underlying the change in paralog SWS sensitivity (Fig. 2). The conserved Y136F substitution presumably also influences spectral tuning across other *Ischnura* SWb1 and b2 opsin paralogs, although possibly to varying degrees, as species-specific variation in spectral sensitivity can be expected between orthologous native rhodopsins, as observed with the orthologous opsins of *Sympetrum frequens*: LWA2 (λ_{max} 557 nm) and LWD1 (λ_{max} 542 nm) (Liénard et al., 2022) compared to the *Ischnura* orthologs (LWA2 λ_{max} 548 nm; LWD1 λ_{max} 541 nm) characterized here.

Opsin expression plasticity as a mechanism underlying learned male mate choice

One major goal of this study was to investigate the potential role of peripheral visual system tuning in a species characterized by dynamic, female-limited polymorphisms and sexual conflict. In such systems with rapidly changing female morph frequencies, males are expected to rapidly respond to the challenges associated with mate detection by developing plastic search images shaped by the local frequencies of female morphs and their intrinsic fecundities (Fincke, 2004; Gosden & Svensson, 2009; Verzijden et al., 2012). There is increasing interest in the role of plasticity in evolution (West-Eberhard, 2003), including plasticity in mate preferences of both males and females (Verzijden et al., 2012). Such plastic

or learned mate preferences can have profound consequences for sexual selection, sexual

conflict, the maintenance of sexually selected polymorphisms, population divergence, and

speciation (Svensson et al., 2010; Verzijden et al., 2012; Westerman et al., 2012). Our results

suggest that opsin gene expression profiles could provide a proximate, mechanistic link

between learned or plastic mate preferences, the visual system, and ultimate evolutionary

theories about polymorphism maintenance.

In this study, we quantified opsin mRNA levels across adult *I. elegans* maturation from

eight sites across two years, allowing for opsin expression to be quantified across a continuum

of androchrome female frequencies and over male ontogony. Patterns of opsin plasticity over

adult male development and opsin expression profiles of adult males were consistent in some

ways with predictions of the LMR hypothesis, namely that increased sexual experience should

tune the visual system for successful mate identification and in response to local female morph

abundances; however, our results suggest a more complicated effect of local female morph

abundances on opsin expression.

We identified opsin-specific plasticity in relative expression over male ontogeny (Figs.

4 - 7), consistent with age-specific selection pressures likely acting on males within each

maturity stage. Teneral males, which are sexually immature, are not expected to express opsin

genes important in mate detection, under the assumption that such expression is costly. In line

with this assumption, we found no significant differences in relative expression of SWS opsins

and few significant differences between LWS opsins for teneral males (Fig. 5A). In contrast,

there were many differences in relative expression between SWS and LWS opsin classes for

sexually immature and sexually mature males (Fig. 5B, C), which could indicate gradual tuning

of the visual system corresponding to the increasing relevance of sexual signals in mate-

searching males. Between maturity stages, significant increases in relative expression of the

SW1 and LWF2 opsins occurred over maturation, while LWF1 and the ocelli opsin LWE

showed significant decreases in relative expression (Fig. 6), consistent with opsin-specific

regulation over male sexual maturation. Further, proportional opsin expression, indicative of

changes in visual sensitivity, differed between developmental stages, with significant increases

in proportion expression of the LWF2 opsin in immature and mature males relative to teneral

males (Fig. 7). *In vitro* measurements of opsin sensitivity show that the LWF2 opsin provides

the ability to perceive wavelengths of light reflected by infuscans females for a finely tuned

visual match to infuscans body colouration (Fig. 3). Lastly, assessing the effect of local female

morph abundances on relative opsin gene expression suggest that mature males, which are

likely to have had more sexual encounters than immature and teneral males, have opsin

expression profiles well-suited to detect the relatively more common infuscans females in

populations with low frequencies of androchrome females (Fig. 8).

585   These results are broadly in agreement with the predictions of the LMR hypothesis.

However, unexpectedly, LWF2 expression was also increased in mature males in settings with

high frequencies of androchrome females, with no evidence that SWS opsins were significantly

upregulated in androchrome-biased settings (Fig. 8). Our previous research has revealed that

androchrome females have the lowest mating rates of the three female morphs in *I. elegans*

(Gosden & Svensson, 2009; Willink et al., 2019). This could increase mating competition

between males in populations with high frequencies of androchrome females. Experiments in

Ischnura ramburii have revealed that males exposed to higher frequencies of androchrome

females experience more antagonistic male-male interactions, had lower body condition, and

higher mortality rates than males from gynochrome-biased populations (Gering, 2017). Given

these potential costs of increased androchrome encounters and the intrinsic fecundity

differences between female morphs, with as much as 30% higher relative fecundity estimated

for infuscans females (Svensson & Abbott, 2005; Willink & Svensson, 2017), male *I. elegans*

from high androchrome populations might gain direct fitness benefits by tuning their visual

system to be better able to identify infuscans females.

Overall, our results suggest opsin expression can be tuned in the male visual system in

I. elegans in a manner consistent with a role in mate choice; however, we would ideally link

the expression differences documented in the current study to behavioural differences in mate

identification or preferences. While the link between opsin expression and behaviour is weak

in some systems (Fuller et al., 2010; Wright et al., 2020), in other cases, differences in opsin

expression does correspond with behaviour (Sakai et al., 2018; Seehausen et al., 2008). For

example, female guppies from low-predator populations prefer males with more orange/red

colouration and have higher expression of opsins that are sensitive to orange and red

wavelengths compared to females from high-predator populations (Sandkam et al., 2015).

Future work directly assessing the effect of specific opsin expression profiles on mating

outcomes will provide valuable insights into the function of opsin plasticity in mate choice and

the maintenance of female polymorphism across Odonata.

CONCLUSION

The results of the current study show that opsin gene expression, specifically the long-

wavelength sensitive LWF2 opsin, is plastic over male ontogeny and changes in response to

local female morph frequencies in mate-searching male *I. elegans*. Functional

analyses revealed that the range of spectral sensitivity of *I. elegans* long-wavelength opsins

closely corresponds to infuscans female body reflectance, with subsequent opsin-specific

expression changes likely contributing to efficiently detect infuscans females over the male-

mimicking androchrome females. Males that likely encounter proportionally more infuscans

females showed increased relative expression of LWF2, suggestive of a mechanism at the

peripheral visual system level to improve detection of common female morphs. In contrast,

higher relative LWF2 expression in males from high androchrome populations does not support

visual tuning for identification of common female morphs. In this case, other behavioural and

physiological differences between androchrome and gynochrome females (e.g., aggression or

fecundity differences) may incur additional costs or benefits that could explain observed

patterns of opsin gene expression in the current study. Future work exploring how these

dynamic patterns of opsin expression correspond to differences in male mating behaviour have

the potential to reveal the proximate mechanisms underlying morph detection and mate choice.

ACKNOWLEDGMENTS

We would like to thank E. Kärrnäs, K. Pehrson, S. Nilén, A.S. Torres Lara, and S. Haley

for assistance in the field, B. Willink for providing access to *I. elegans* transcriptomic data, and

J.-M. Lassance for his help annotating *I. elegans* opsin genes from genomic data. We would

also like to thank S. Nilén for providing access to scripts and to generate sampling maps. This

work was supported by grants from the Swedish Research Council (VR 2020-0517 to M.A.L.

and VR2020-03123 to E.I.S.), Carl Trygger Foundation (CTS20:248) to M.A.L, Erik Philip-

Sörensens Stiftelse (G2022-007) to E.I.S, and Stiftelsen Längmanska Kulturfonden (BA21-

1192) to N.S.R.

DATA ACCESSIBILITY

Datafiles and code associated all the analyses in this paper will be uploaded to the Dryad Digital

Data Repository upon acceptance.

AUTHOR CONTRIBUTIONS

All authors contributed to conception of the experimental design. M.A.L performed

opsin heterologous expression and analysis of opsin function sensitivity data, N.S.R collected

field data and performed qPCR and statistical analyses with support from E.I.S and M.A.L.

N.S.R wrote the manuscript with support from E.I.S and M.A.L. All authors reviewed the final

manuscript.

REFERENCES

Arikawa, K. (2017). The eyes and vision of butterflies. *J Physiol*, *595*(16), 5457–5464.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Blow, R., Willink, B., & Svensson, E. I. (2021). A molecular phylogeny of forktail

damselflies (genus *Ischnura*) reveals a dynamic macroevolutionary history of female

colour polymorphisms. *Molecular Phylogenetics and Evolution*, *160*, 107134.

https://doi.org/10.1016/j.ympev.2021.107134

Briscoe, A. D., & Chittka, L. (2001). The evolution of color vision in insects. *Annual Review*

of Entomology, *46*(1), 471–510. https://doi.org/10.1146/annurev.ento.46.1.471

Bybee, S. M., Johnson, K. K., Gering, E. J., Whiting, M. F., & Crandall, K. A. (2012). All the

better to see you with: A review of odonate color vision with transcriptomic insight

into the odonate eye. *Organisms Diversity & Evolution*, *12*(3), 241–250.

https://doi.org/10.1007/s13127-012-0090-6

Carleton, K. L., Dalton, B. E., Escobar-Camacho, D., & Nandamuri, S. P. (2016). Proximate

and ultimate causes of variable visual sensitivities: Insights from cichlid fish

radiations. *Genesis*, *54*(6), 299–325. https://doi.org/10.1002/dvg.22940

Castiglione, G. M., Hauser, F. E., Van Nynatten, A., & Chang, B. S. W. (2023). Adaptation of

antarctic icefish vision to extreme environments. *Molecular Biology and Evolution*,

40(4), msad030. https://doi.org/10.1093/molbev/msad030

Chang, C.-H., Catchen, J., Moran, R. L., Rivera-Colón, A. G., Wang, Y.-C., & Fuller, R. C.

(2021). Sequence analysis and ontogenetic expression patterns of cone opsin genes in

the bluefin killifish (*Lucania goodei*). *Journal of Heredity*, *112*(4), 357–366.

https://doi.org/10.1093/jhered/esab017

Combes, S. A., Salcedo, M. K., Pandit, M. M., & Iwasaki, J. M. (2013). Capture success and

efficiency of dragonflies pursuing different types of prey. *Integrative and*

Comparative Biology, *53*(5), 787–798. https://doi.org/10.1093/icb/ict072

Corbet, P. S. (1999). *Dragonflies: Behavior and Ecology of Odonata.* Cornell Univ. Press,

Ithaca, NY. 829pp.

Córdoba-Aguilar, A. (2023). *Dragonflies and Damselflies: Model Organisms for Ecological*

and Evolutionary Research. Oxford University Press.

Dijkstra, K. D., & Lewington, R. (2006). *Field Guide to the Dragonflies of Britain and*

Europe. Bloomsbury Publishing.

Dudaniec, R. Y., Yong, C. J., Lancaster, L. T., Svensson, E. I., & Hansson, B. (2018).

Signatures of local adaptation along environmental gradients in a range-expanding

damselfly (Ischnura elegans). *Molecular Ecology*, *27*(11), 2576–2593.

https://doi.org/10.1111/mec.14709

Everett, A., Tong, X., Briscoe, A. D., & Monteiro, A. (2012). Phenotypic plasticity in opsin

expression in a butterfly compound eye complements sex role reversal. *BMC*

Evolutionary Biology, *12*(232). https://doi.org/10.1186/1471-2148-12-232

Fincke, O. M. (2004). Polymorphic signals of harassed female odonates and the males that

learn them support a novel frequency-dependent model. *Animal Behaviour*, *67*(5),

833–845. https://doi.org/10.1016/j.anbehav.2003.04.017

Fincke, O. M., Jödicke, R., Paulson, D. R., & Schultz, T. D. (2005). The evolution and

frequency of female color morphs in Holarctic Odonata: Why are male-like females

typically the minority? *International Journal of Odonatology*, *8*(2), 183–212.

https://doi.org/10.1080/13887890.2005.9748252

Frentiu, F. D., Yuan, F., Savage, W. K., Bernard, G. D., Mullen, S. P., & Briscoe, A. D.

(2015). Opsin clines in butterflies suggest novel roles for insect photopigments.

Molecular Biology and Evolution, *32*(2), 368–379.

https://doi.org/10.1093/molbev/msu304

Frick, H., Chow, F., Kuhn, M., Mahoney, M., Silge, J., & Wickham, H. (2022). *rsample:*

General Resampling Infrastructure. https://rsample.tidymodels.org,

https://github.com/tidymodels/rsample.

Fuller, R. C., & Claricoates, K. M. (2011). Rapid light-induced shifts in opsin expression:

Finding new opsins, discerning mechanisms of change, and implications for visual

sensitivity. *Molecular Ecology*, *20*(16), 3321–3335. https://doi.org/10.1111/j.1365-

294X.2011.05180.x

Fuller, R. C., Noa, L. A., & Strellner, R. S. (2010). Teasing apart the many effects of lighting

environment on opsin expression and foraging preference in bluefin killifish. *The*

American Naturalist, *176*(1), 1–13. https://doi.org/10.1086/652994

Futahashi, R., Kawahara-Miki, R., Kinoshita, M., Yoshitake, K., Yajima, S., Arikawa, K., &

Fukatsu, T. (2015). Extraordinary diversity of visual opsin genes in dragonflies.

Proceedings of the National Academy of Sciences, *112*(11), E1247–E1256.

https://doi.org/10.1073/pnas.1424670112

Gering, E. J. (2017). Male-mimicking females increase male-male interactions, and decrease

male survival and condition in a female-polymorphic damselfly. *Evolution*, *71*(5),

1390–1396. https://doi.org/10.1111/evo.13221

Gosden, T. P., & Svensson, E. I. (2009). Density-dependent male mating harassment, female

resistance, and male mimicry. *The American Naturalist*, *173*(6), 709–721.

https://doi.org/10.1086/598491

Govardovskii, V. I., Fyhrquist, N., Reuter, T., Kuzmin, D. G., & Donner, K. (2000). In search

of the visual pigment template. *Visual Neuroscience*, *17*(4), 509–528.

https://doi.org/10.1017/S0952523800174036

Henze, M. J., Lind, O., Wilts, B. D., & Kelber, A. (2019). Pterin-pigmented nanospheres

create the colours of the polymorphic damselfly <i>Ischnura elegans<i/>. *Journal of*

The Royal Society Interface, *16*(153), 20180785.

https://doi.org/10.1098/rsif.2018.0785

Horridge, G. A. (1969). Unit studies on the retina of dragonflies. *Zeitschrift Für*

Vergleichende Physiologie, *62*(1), 1–37. https://doi.org/10.1007/BF00298040

Huang, S., Chiou, T., Marshall, J., & Reinhard, J. (2014). Spectral sensitivities and color

signals in a polymorphic damselfly. *PLOS ONE*, *9*(1), e87972.

https://doi.org/10.1371/journal.pone.0087972

Jacobs, G. H. (2013). Losses of functional opsin genes, short-wavelength cone

photopigments, and color vision—A significant trend in the evolution of mammalian

vision. *Visual Neuroscience*, *30*(1–2), 39–53.

https://doi.org/10.1017/S0952523812000429

Keller, O., Kollmar, M., Stanke, M., & Waack, S. (2011). A novel hybrid gene prediction

method employing protein multiple sequence alignments. *Bioinformatics*, *27*(6), 757–

763. https://doi.org/10.1093/bioinformatics/btr010

Kuhn, M., & Wickham, H. (2020). *Tidymodels: A collection of packages for modeling and*

machine learning using tidyverse principles.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). LmerTest package: Tests in

linear mixed effects models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Labhart, T., & Nilsson, D.-E. (1995). The dorsal eye of the dragonfly *Sympetrum*:

Specializations for prey detection against the blue sky. *Journal of Comparative*

Physiology A, *176*(4), 437–453. https://doi.org/10.1007/BF00196410

Lancer, B. H., Evans, B. J. E., & Wiederman, S. D. (2020). The visual neuroecology of

anisoptera. *Current Opinion in Insect Science*, *42*, 14–22.

https://doi.org/10.1016/j.cois.2020.07.002

Le Rouzic, A., Hansen, T. F., Gosden, T. P., & Svensson, E. I. (2015). Evolutionary time-

series analysis reveals the signature of frequency-dependent selection on a female

mating polymorphism. *The American Naturalist*, *185*(6), E182–E196.

https://doi.org/10.1086/680982

Lenth, R. V. (2022). *Emmeans: Estimated Marginal Means, aka Least-Squares Means. R*

package version 1.8.2. https://CRAN.R-project.org/package=emmeans

Liénard, M. A., Bernard, G. D., Allen, A., Lassance, J.-M., Song, S., Childers, R. R., Yu, N.,

Ye, D., Stephenson, A., Valencia-Montoya, W. A., Salzman, S., Whitaker, M. R. L.,

Calonje, M., Zhang, F., & Pierce, N. E. (2021). The evolution of red color vision is

linked to coordinated rhodopsin tuning in lycaenid butterflies. *Proceedings of the*

National Academy of Sciences, *118*(6), e2008986118.

https://doi.org/10.1073/pnas.2008986118

Liénard, M. A., Valencia-Montoya, W. A., & Pierce, N. E. (2022). Molecular advances to

study the function, evolution and spectral tuning of arthropod visual opsins.

Philosophical Transactions of the Royal Society B: Biological Sciences, *377*(1862),

20210279. https://doi.org/10.1098/rstb.2021.0279

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using

real-time quantitative PCR and the 2−ΔΔct method. *Methods*, *25*(4), 402–408.

https://doi.org/10.1006/meth.2001.1262

McCulloch, K. J., Macias-Muñoz, A., Mortazavi, A., & Briscoe, A. D. (2022). Multiple

mechanisms of photoreceptor spectral tuning in heliconius butterflies. *Molecular*

Biology and Evolution, *39*(4), msac067. https://doi.org/10.1093/molbev/msac067

McCulloch, K. J., Yuan, F., Zhen, Y., Aardema, M. L., Smith, G., Llorente-Bousquets, J.,

Andolfatto, P., & Briscoe, A. D. (2017). Sexual dimorphism and retinal mosaic

diversification following the evolution of a violet receptor in butterflies. *Molecular*

Biology and Evolution, *34*(9), 2271–2284. https://doi.org/10.1093/molbev/msx163

Meinertzhagen, I. A., Menzel, R., & Kahle, G. (1983). The identification of spectral receptor

types in the retina and lamina of the dragonfly *Sympetrum rubicundulum*. *Journal of*

Comparative Physiology, *151*(3), 295–310. https://doi.org/10.1007/BF00623906

Miller, M. N., & Fincke, O. M. (1999). Cues for mate recognition and the effect of prior

experience on mate recognition in *Enallagma* damselflies. *Journal of Insect Behavior*,

12(6), 801–814. https://doi.org/10.1023/A:1020957110842

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT–

PCR. *Nucleic Acids Research*, *29*(9), e45. https://doi.org/10.1093/nar/29.9.e45

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2021). *nlme: Linear and*

Nonlinear Mixed Effects Models. https://CRAN.R-project.org/package=nlme

R Core Team. (2021). R Foundation for Statistical Computing. URL https://www.R-

project.org/

Rebora, M., Frati, F., Piersanti, S., Salerno, G., Selvaggini, R., & Fincke, O. M. (2018). Field tests of multiple sensory cues in sex recognition and harassment of a colour polymorphic damselfly. *Animal Behaviour*, *136*, 127–136. https://doi.org/10.1016/j.anbehav.2017.12.015

Rister, J., & Desplan, C. (2011). The retinal mosaics of opsin expression in invertebrates and vertebrates. *Developmental Neurobiology*, *71*(12), 1212–1226. https://doi.org/10.1002/dneu.20905

Robertson, H. M. (1985). Female dimorphism and mating behaviour in a damselfly, *Ischnura ramburi*: Females mimicking males. *Animal Behaviour*, *33*(3), 805–809. https://doi.org/10.1016/S0003-3472(85)80013-0

Rossetto, I. H., Sanders, K. L., Simões, B. F., Van Cao, N., & Ludington, A. J. (2023). Functional duplication of the short-wavelength-sensitive opsin in sea snakes: Evidence for reexpanded color sensitivity following ancestral regression. *Genome Biology and Evolution*, *15*(7), evad107. https://doi.org/10.1093/gbe/evad107

Sakai, Y., Kawamura, S., & Kawata, M. (2018). Genetic and plastic variation in opsin gene expression, light sensitivity, and female response to visual signals in the guppy. *Proceedings of the National Academy of Sciences*, *115*(48), 12247–12252. https://doi.org/10.1073/pnas.1706730115

Sandkam, B., Young, C. M., & Breden, F. (2015). Beauty in the eyes of the beholders: Colour vision is tuned to mate preference in the Trinidadian guppy (*Poecilia reticulata*). *Molecular Ecology*, *24*(3), 596–609. https://doi.org/10.1111/mec.13058

Sasagawa, H., Narita, R., Kitagawa, Y., & Kadowaki, T. (2003). The expression of genes encoding visual components is regulated by a circadian clock, light environment and age in the honeybee (*Apis mellifera*). *European Journal of Neuroscience*, *17*(5), 963–970. https://doi.org/10.1046/j.1460-9568.2003.02528.x

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., van

der Sluijs, I., Schneider, M. V., Maan, M. E., Tachida, H., Imai, H., & Okada, N.

(2008). Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), Article

7213. https://doi.org/10.1038/nature07285

Shand, J., Davies, W. L., Thomas, N., Balmer, L., Cowing, J. A., Pointer, M., Carvalho, L. S.,

Trezise, A. E. O., Collin, S. P., Beazley, L. D., & Hunt, D. M. (2008). The influence of

ontogeny and light environment on the expression of visual pigment opsins in the

retina of the black bream, *Acanthopagrus butcheri*. *Journal of Experimental Biology*,

211(9), 1495–1503. https://doi.org/10.1242/jeb.012047

Sharkey, C. R., Blanco, J., Lord, N. P., & Wardill, T. J. (2023). Jewel beetle opsin duplication

and divergence is the mechanism for diverse spectral sensitivities. *Molecular Biology*

and Evolution, *40*(2), msad023. https://doi.org/10.1093/molbev/msad023

Stieb, S. M., Carleton, K. L., Cortesi, F., Marshall, N. J., & Salzburger, W. (2016). Depth-

dependent plasticity in opsin gene expression varies between damselfish

(Pomacentridae) species. *Molecular Ecology*, *25*(15), 3645–3661.

https://doi.org/10.1111/mec.13712

Svensson, E. I., & Abbott, J. (2005). Evolutionary dynamics and population biology of a

polymorphic insect. *Journal of Evolutionary Biology*, *18*(6), 1503–1514.

https://doi.org/10.1111/j.1420-9101.2005.00946.x

Svensson, E. I., Abbott, J., & Härdling, R. (2005). Female polymorphism, frequency

dependence, and rapid evolutionary dynamics in natural populations. *The American*

Naturalist, *165*(5), 567–576. https://doi.org/10.1086/429278

Svensson, E. I., Eroukhmanoff, F., Karlsson, K., Runemark, A., & Brodin, A. (2010). A role

for learning in population divergence of mate preferences. *Evolution*, *64*(11), 3101–

3113. https://doi.org/10.1111/j.1558-5646.2010.01085.x

Svensson, E. I., Willink, B., Duryea, M. C., & Lancaster, L. T. (2020). Temperature drives

pre-reproductive selection and shapes the biogeography of a female polymorphism.

Ecology Letters, *23*(1), 149–159. https://doi.org/10.1111/ele.13417

Takahashi, Y., Kagawa, K., Svensson, E. I., & Kawata, M. (2014). Evolution of increased

phenotypic diversity enhances population performance by reducing sexual harassment

in damselflies. *Nature Communications*, *5*(1), Article 1.

https://doi.org/10.1038/ncomms5468

Takahashi, Y., & Watanabe, M. (2010). Female reproductive success is affected by selective

male harassment in the damselfly *Ischnura senegalensis*. *Animal Behaviour*, *79*(1),

211–216. https://doi.org/10.1016/j.anbehav.2009.10.032

Takahashi, Y., & Watanabe, M. (2011). Male mate choice based on ontogenetic colour

changes of females in the damselfly *Ischnura senegalensis*. *Journal of Ethology*,

29(2), 293–299. https://doi.org/10.1007/s10164-010-0257-6

Takahashi, Y., Yoshimura, J., Morita, S., & Watanabe, M. (2010). Negative frequency-

dependent selection in female color polymorphism of a damselfly. *Evolution*, *64*(12),

3620–3628. https://doi.org/10.1111/j.1558-5646.2010.01083.x

Terakita, A. (2005). The opsins. *Genome Biology*, *6*(3), 213. https://doi.org/10.1186/gb-2005-

6-3-213

van der Kooi, C. J., Stavenga, D. G., Arikawa, K., Belusic, G., & Kelber, A. (2021).

Evolution of insect colour vision – from spectral sensitivity to visual ecology. *Annual*

Review of Entomology, *66*. https://doi.org/DOI 10.1146/annurev-ento-061720-

071644.

Van Gossum, H., Bots, J., Van Heusden, J., Hammers, M., Huyghe, K., & Morehouse, N. I.

(2011). Reflectance spectra and mating patterns support intraspecific mimicry in the

colour polymorphic damselfly *Ischnura elegans*. *Evolutionary Ecology*, *25*(1), 139–

154. https://doi.org/10.1007/s10682-010-9388-z

Van Gossum, H., Stoks, R., & De Bruyn, L. (2001a). Discriminative mate choice in relation

with female maturation in *Ischnura elegans* (Odonata: Coenagrionidae). *International*

Journal of Odonatology, *4*(1), 83–91.

https://doi.org/10.1080/13887890.2001.9748161

Van Gossum, H., Stoks, R., & De Bruyn, L. (2001b). Reversible frequency–dependent

switches in male mate choice. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences, *268*(1462), 83–85. https://doi.org/10.1098/rspb.2000.1333

Verzijden, M. N., ten Cate, C., Servedio, M. R., Kozak, G. M., Boughman, J. W., &

Svensson, E. I. (2012). The impact of learning on sexual selection and speciation.

Trends in Ecology & Evolution, *27*(9), 511–519.

https://doi.org/10.1016/j.tree.2012.05.007

Wakakuwa, M., Terakita, A., Koyanagi, M., Stavenga, D. G., Shichida, Y., & Arikawa, K.

(2010). Evolution and mechanism of spectral tuning of blue-absorbing visual

pigments in butterflies. *PLOS ONE*, *5*(11), e15015.

https://doi.org/10.1371/journal.pone.0015015

Wang, B., Xiao, J.-H., Bian, S.-N., Niu, L.-M., Murphy, R. W., & Huang, D.-W. (2013).

Evolution and expression plasticity of opsin genes in a fig pollinator, *Ceratosolen*

solmsi. *PLOS ONE*, *8*(1), e53907. https://doi.org/10.1371/journal.pone.0053907

Wernet, M. F., Perry, M. W., & Desplan, C. (2015). The evolutionary diversity of insect

retinal mosaics: Common design principles and emerging molecular logic. *Trends in*

Genetics, *31*(6), 316–328. https://doi.org/10.1016/j.tig.2015.04.006

West-Eberhard, M. J. (2003). *Developmental Plasticity and Evolution*. Oxford University

Press.

Westerman, E. L., Hodgins-Davis, A., Dinwiddie, A., & Monteiro, A. (2012). Biased learning

affects mate choice in a butterfly. *Proceedings of the National Academy of Sciences*,

109(27), 10948–10953. https://doi.org/10.1073/pnas.1118378109

Willink, B., Duryea, M. C., Wheat, C., & Svensson, E. I. (2020). Changes in gene expression

during female reproductive development in a color polymorphic insect. *Evolution*,

74(6), 1063–1081. https://doi.org/10.1111/evo.13979

Willink, B., Duryea, M., & Svensson, E. (2019). Macroevolutionary origin and adaptive

function of a polymorphic female signal involved in sexual conflict. *The American*

Naturalist, *194*. https://doi.org/10.1086/705294

Willink, B., & Svensson, E. I. (2017). Intra- and intersexual differences in parasite resistance

and female fitness tolerance in a polymorphic insect. *Proceedings of the Royal Society*

B: Biological Sciences, *284*(1847), 20162407. https://doi.org/10.1098/rspb.2016.2407

Willink, B., Tunström, K., Nilén, S., Chikhi, R., Lemane, T., Takahashi, M., Takahashi, Y.,

Svensson, E. I., & Wheat, C. W. (in press). The genomics and evolution of inter-

sexual mimicry and female-limited polymorphisms in damselflies. *Nature Ecology &*

Evolution, 2023.03.27.532508. https://doi.org/10.1101/2023.03.27.532508

Winfrey, C., & Fincke, O. M. (2017). Role of visual and non-visual cues in damselfly mate

recognition. *International Journal of Odonatology*, *20*(1), 43–52.

https://doi.org/10.1080/13887890.2017.1297259

Wright, D. S., van Eijk, R., Schuart, L., Seehausen, O., Groothuis, T. G. G., & Maan, M. E.

(2020). Testing sensory drive speciation in cichlid fish: Linking light conditions to

opsin expression, opsin genotype and female mate preference. *Journal of*

Evolutionary Biology, *33*(4), 422–434. https://doi.org/10.1111/jeb.13577

Yang, E.-C., & Osorio, D. (1991). Spectral sensitivities of photoreceptors and lamina

monopolar cells in the dragonfly, *Hemicordulia tau*. *Journal of Comparative*

Physiology A, *169*(6), 663–669. https://doi.org/10.1007/BF00194895

Yilmaz, A., Lindenberg, A., Albert, S., Grübel, K., Spaethe, J., Rössler, W., & Groh, C.

(2016). Age-related and light-induced plasticity in opsin gene expression and in

primary and secondary visual centers of the nectar-feeding ant *Camponotus rufipes*.

Developmental Neurobiology, *76*(9), 1041–1057. https://doi.org/10.1002/dneu.22374

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects*

models and extensions in ecology with R. Springer. https://doi.org/10.1007/978-0-387-

87458-6

TABLES & FIGURES

Table 1. Final model and model output showing the effect of maturity stage and opsin type on log relative opsin expression in teneral, immature, and mature male *I. elegans*. Significant differences are shown in italics.

log Relative Opsin Expression ~ Maturity × Opsin + (1|ID) +(1|Replicate) + (1|Year)			
	df	F	*P*
Maturity	2	0.2	0.82
Opsin	7	292.3	*< 0.0001*
Maturity × Opsin	14	12.3	*< 0.0001*

Table 2. Final model and model output showing the effect of maturity stage and opsin type on proportion opsin expression in teneral, immature, and mature male *I. elegans*. Significant differences are shown in italics.

Proportion Opsin Expression ~ Maturity × Opsin + (1|ID)			
	df	F	*P*
Maturity	2	10.8	*< 0.0001*
Opsin	6	415.6	*< 0.0001*
Maturity × Opsin	12	29.8	*< 0.0001*

Table 3. Final model and model output showing the effect of the proportion of androchrome females in a population, opsin type, and maturity stage on log relative opsin expression for immature and mature male *I. elegans*. Significant differences are shown in italics.

log Relative Opsin Expression ~ Opsin × Maturity × (Prop. Androchrome + Prop. Androchrome^2) + (1|ID) + (1|Replicate) + (1|Year)			
	df	F	*P*
Opsin	7	18.7	*< 0.0001*
Maturity	1	1.3	0.27
Prop. Androchrome	1	0.2	0.66
Prop. Androchrome^2	1	0.2	0.67
Opsin × Maturity	7	3.1	*< 0.01*
Opsin × Prop. Androchrome	7	17.1	*< 0.0001*
Opsin × Prop. Androchrome^2	7	17.3	*< 0.0001*
Maturity × Prop. Androchrome	1	1.3	0.26
Maturity × Prop. Androchrome^2	1	1.3	0.25
Opsin × Maturity × Prop. Androchrome	7	3.3	*< 0.01*
Opsin × Maturity × Prop. Androchrome^2	7	3.6	*< 0.001*

Figure 1. Colour polymorphism and female morph frequencies in the damselfly *I. elegans* from eight populations from sites in the region of Skåne, southern Sweden for the damselfly *Ischnura elegans*. (A) Mature male *I. elegans* and the three female-limited colour morphs in their sexual mature colour phases. (B) Females undergo ontogenetic colour changes in the distal abdomen segments (left photo) and thorax (right photo) between immature and sexually mature adult stages. A = androchrome, I = infuscans, O = infuscans-obsoleta. (C) Sampling locations of *I. elegans* used in the current study with inset showing the area of southern Sweden sampled. (D) Female morph frequencies per population in 2021 and I 2022. Numbers in parentheses above each bar indicate sample size. (A, B) modified from Willink et al. (2020).

Figure 2. Functional expression of *Ischnura elegans* short-wavelength (SW) and long-wavelength (LW) opsins. Ultraviolet-visible (UV-VIS) absorbance analyses of dark-adapted rhodopsin visual pigments reconstituted and purified from HEK293T cell cultures in the dark in the presence of 11-*cis*-retinal. The black dots represent mean absorbances at a given wavelength. (A) SWb1 (n = 7), (B) SWb2 (n = 3), (C) LWA2 (n = 2), (D) LWF1 (n = 2), I LWF2 (n = 5), (F) LWF3 (n = 6), (G) LWF4 (n = 1), and opsins expressed in the ocelli: (H) LWD1 (n = 3) and (I) LWE1 (n = 6), where n is the number of measurements of protein aliquots with active rhodopsin complexes. Absorbance at 380 nm in A and B is due to residual unbound *cis*-retinal. Relative absorbance data are fitted to a visual template with polynomial function analyses computed in R to obtain the best estimates following 1000 bootstrap analysis of lambda max for each rhodopsin (Liénard et al., 2022). Confidence intervals are indicated in parentheses.

Figure 3. Absorbance spectra of opsins and female body reflectance in *I. elegans*. Sensitivity of opsins expressed in the ventral compound eye of male *I. elegans* are represented by coloured visual fit curves. Body reflectance for (A) androchrome and (B) infuscans female body colouration is represented by the shaded gray area. Body reflectance data from Henze et al. (2019).

Figure 4. Linear discriminate analysis showing the relationship between relative expression of eight opsins expressed within the ventral eye or ocelli of teneral, immature, and mature male *I. elegans* for the first two linear discriminate factors (LD1 and LD2; percentages shown on the x- and y -axes).

Figure 5. Relative opsin expression for (A) teneral, (B) immature, and (C) mature male *I. elegans*. Bars in box and whisker plots show medians, boxes indicate upper and lower quartiles, whiskers show sample minima and maxima, and open circles show outliers. Letters above each bar show significant differences between opsin types within each maturity level following Tukey corrections for multiple comparisons. Test statistics and p-values for all comparisons are presented in Table S8.

Figure 6. Relative opsin expression across seven opsins expressed primarily in the ventral compound eye (SW1 – LWF4) and one opsin primarily expressed in the ocelli (LWE) for male *I. elegans.* Teneral males are shown in light blue, immature adult males in medium blue, and mature males in dark blue. Bars in box and whisker plots show medians, boxes indicate upper and lower quartiles, whiskers show sample minima and maxima, and open circles show outliers. Single asterisks indicate comparisons that are significant at alpha < 0.05 and double asterisks represent comparisons that are significant at alpha < 0.01 following Tukey corrections for multiple comparisons within opsin types. Test statistics and p-values for all comparisons are presented in Table S9.

Figure 7. Average proportion expression of opsins found in the ventral compound eye of teneral, immature, and mature male *I. elegans*.

Figure 8. Log relative opsin expression in immature (open circles, dashed line) and mature (black triangle, solid line) males *I. elegans* from populations with different proportions of androchrome females. The opsins SW1, SW2, LWA, and LWF1 – LWF4 are expressed in the compound eye and the opsin LWE is expressed primarily in ocelli. Grey shading around trend lines indicates 95% confidence intervals.