Transcriptomics elucidates metabolic regulation and functional promoters in the basidiomycete red yeast *Xanthophyllomyces dendrorhous* CBS 6938

Emma E. Tobin[1,†], Joseph H. Collins[1,†], Celeste B. Marsan[1], Gillian T. Nadeau[1], Kim Mori[1], Anna Lipzen[2], Stephen Mondo[2], Igor V. Grigoriev[2,3], Eric M. Young[1]*

[1] Life Sciences and Bioengineering Center, Department of Chemical Engineering, Worcester Polytechnic Institute, Worcester, MA

[2] U.S. Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

[3] Plant and Microbial Biology, University of California Berkeley, Berkeley, CA 94720

[†] Equal contribution

* To whom correspondence should be addressed. emyoung@wpi.edu

ABSTRACT

Genomics has become the primary way to explore microbial diversity, because genetic

tools are currently difficult to develop in non-model organisms. Here, we demonstrate that -omics

can be leveraged to accelerate genetic tool development for the basidiomycete yeast

Xanthophyllomyces dendrorhous CBS 6938, the sole biotechnologically relevant organism in the

Tremellomycete family. First, we sequence the genome. Then, we perform transcriptomics under

a variety of conditions, focusing on light and oxidative stress. This data not only reveals novel

photobiology and metabolic regulation, it also allows derivation of constitutive and regulated gene

expression parts. Our analysis of *X. dendrorhous* photobiology shows for the first time that a

complex system of white-collar and cryptochrome homologs mediate response to ultraviolet light

(UV). Our analysis of metabolic regulation shows that UV activates DNA repair, aromatic amino

acid and carotenoid biosynthesis and represses central carbon metabolism and the fungal-like

apoptotic pathway. Thus, *X. dendrorhous* shows a dynamic response toward biosynthetic

pathways for light-absorbing compounds and survival and away from energy production. We then

define a modular cloning system, including antibiotic selections, integration sites, and reporter

genes, and use the transcriptomics to derive strong constitutive and regulated promoters. Notably,

we discover a novel promoter from a hypothetical gene that has 9-fold activation upon UV

exposure. Thus, -omics-to-parts workflows can simultaneously provide useful genomic data and

advance genetic tools for non-model microbes, particularly those without a closely related model

organism. This approach will be broadly useful in current efforts to engineer diverse microbes.

INTRODUCTION

Yeasts are a diverse group of fungi characterized by unicellular growth, which is useful in industrial fermentations. The group is not monophyletic, rather the yeast state has emerged several times across the fungal kingdom. Therefore, yeasts have vastly different genome architectures, metabolic capabilities, and stress tolerances that are not only due to the ecological niches to which they are adapted, but also due to their different lineages. The impact this biodiversity might have on industrial biotechnology has only just begun to be explored.[1] However, the different lineages limit the transferability of genomics and genetic tools between yeasts, particularly for the basidiomycete yeasts that are not closely related to any model organism.[2] Here, we demonstrate how modern genomics and transcriptomics can be leveraged to simultaneously gain new insight into nonconventional yeast biology and derive a collection of genetic parts for future studies. We demonstrate this for the basidiomycete yeast *Xanthophyllomyces dendrorhous,* the only biotechnologically relevant organism from the Tremellomycete family of Basidiomycota.[3]

Historically, the model ascomycete *Saccharomyces cerevisiae* has been the host of choice for yeast metabolic engineering.[4] While valuable for converting hexoses to ethanol, engineering *S. cerevisiae* to consume other carbon sources and produce other economically valuable compounds can require extensive reprogramming, which imposes metabolic and gene expression burden.[5] This is particularly true of tolerance, which can be due to many genes interacting in currently unpredictable ways. This has led to a growing body of literature that investigates other yeasts, termed nonconventional simply because they are not *S. cerevisiae*, as potential hosts for metabolic engineering since they are already adapted to consume desired carbon sources, produce desired classes of molecules, and tolerate different fermentation conditions, thus less burden is incurred when engineering them.[6,7] Methods for developing genetic tools are needed for these organisms.

Developing genomics and genetic tools in nonconventional yeasts is challenging. This is

because a high-quality genome, gene regulatory data, species-specific DNA parts for genetic

manipulation, and efficient transformation procedures are often missing. The process of

developing these elements, termed "onboarding," can be accelerated for nonconventional

ascomycete yeasts closely related to *S. cerevisiae* because the genomics and genetics are

similar, permitting transfer of similar promoter and plasmid designs. Onboarding organisms from

relatively unknown classes, particularly basidiomycete yeasts, does not have this advantage

because there is not a closely related model organism with a complete, contiguous genome and

well-developed genetics.

Xanthophyllomyces dendrorhous is a nonconventional yeast that has gained attention in

the aquaculture and nutraceutical industries due to its natural ability to produce astaxanthin, a

vibrant red pigment with potent antioxidant properties. It is currently the only Tremellomycete with

biotechnology applications. While gene knock-out and upregulation techniques have been

employed to modify *X. dendrorhous,* particularly strain CBS 6938, reprogramming this yeast

remains a challenge because several key tools in the onboarding toolkit are missing, including a

high-quality genome sequence, a modular gene expression parts collection, and efficient

transformation methods.[6] Furthermore, astaxanthin is only overproduced in response to light and

oxidative stress, which complicates and limits the scale of bioprocesses.[7–9]

Here, we demonstrate an omics-driven onboarding workflow including genome

sequencing, transcriptomics, high efficiency transformation, and modular gene expression parts

collection development for *X. dendrorhous*. This workflow generates new insights into

basidiomycete photobiology and metabolism, constitutive promoters, light-inducible promoters,

and efficient genome integration. Therefore, this "omics-to-parts" approach can rapidly deliver a

suite of data and tools for onboarding nonconventional yeast that have no closely related model

host.

RESULTS

High quality genome of *X. dendrorhous* CBS 6938

A high-quality CBS 6938 genome was obtained through Prymetime sequencing (Figure

1a).[10] Prymetime assembled the genome into sixteen contigs, and relative lengths of the contigs

are depicted in the figure. Subsequent benchmarking universal single-copy ortholog (BUSCO)

analysis revealed that the CBS 6938 genome contains 94.0% complete and single copy BUSCOs

(Figure 1b). This genome sequence was the foundation for the subsequent work.

Foundational genetic tools for onboarding *X. dendrorhous*

In addition to obtaining an accurate genome sequence, we performed a set of experiments

to make *X. dendrorhous* easier to engineer. We first examined possible selection markers. Rather

than seek to create synthetic auxotrophy, which has a variety of metabolic impacts,[11] we screened

for antibiotic sensitivity. We measured *X. dendrorhous* sensitivity to several common antifungals

via minimum inhibitory concentration (MIC) assay (Methods). The most effective antifungal was

found to be nourseothricin with 11.6% cell survival at a concentration of 10 µg/mL (Figure 2a).

We then designed a modular cloning system for rapidly assembling *X. dendrorhous*

genetic elements (Figure 2b). This system uses the same enzymes and scars as previously

published modular cloning systems.[12] There are two significant differences in this red yeast

modular cloning system. The first is that the *lacZ* element in the destination plasmids is replaced

with the lethal gene *ccdb* to shift from blue/white screening of correct clones to a system where

only correct clones are able to grow on a selection plate. The second is that the Level 2 (L2)

integrative plasmid has homology arms compatible with the *X. dendrorhous* genome. We used

our genome sequence to select homology arms flanking that *crtYB* gene, the gene that catalyzes

production of the colorful carotenoid pigments. This creates a facile red/white screening method

for correctly integrated clones.

We then designed an initial integration cassette using the known glutamate

dehydrogenase promoter (P_{gdh}) to drive expression of a codon-optimized *nat* gene which confers

nourseothricin resistance. We used this initial integration cassette to optimize the efficiency of

transformation. The method we found to be most efficient is described in Methods. In brief, it

consists of the culture of CBS 6938 cells until the end of exponential growth phase is reached,

combination with the integrative pathway, and application of an electric shock (Figure 2c). Ultra-

high-performance liquid-chromatography (UPLC) was then used to demonstrate successful *crtYB*

knock-out through measurement of astaxanthin. Astaxanthin peaks were detected in wild type but

not knock-out extractions (Figure 2d). Thus, with the design of the integration cassette and

optimization of the transformation protocol we are able to achieve efficient and targeted

integration of DNA into the *X. dendrorhous* genome using homologous recombination.

We then established a luminescent reporter, which is common in fungal engineering, for

measuring gene expression strength. We cloned the NanoLuc enzyme[13] into our modular cloning

scheme, integrated it into the genome, and measured expression via luciferase assay (Figure 2e).

NanoLuc-producing cells created approximately 1,000-fold greater luminescence compared to

the *crtYB* knockout strain, and nearly 10,000-fold greater luminescence compared to CBS 6938.

The higher luminescence observed from the *crtYB* knockout strain may be attributed to biological

autoluminescence, a phenomenon that has been found to be directly related to oxidative stress.[14]

Together with the genome sequence, these experiments enabled the development of

advanced genetic tools for *X. dendrorhous*.

Confirming that astaxanthin biosynthesis responds to light through transcriptional regulation

We next sought to more fully understand *X. dendrorhous* biology by examining genetic response to light and oxidative stress. While previous work has shown that visible and ultraviolet (UV) light increases carotenogenesis in *X. dendrorhous*, specific genes and mechanisms involved in this regulation have not been investigated. Therefore, we first sought to confirm that the carotenogenesis pathway was transcriptionally regulated. Before conducting transcriptomics, we used the genome sequence to design quantitative polymerase chain reaction (qPCR) probes to target genes in the terpenoid (*MVK*, *PMVK*, *MVD*, *IDI*, *FPS*, and *crtE*) and carotenogenic (*crtYB*, *crtI*, and *crtS*) pathways. Cells were grown under either dark or UV-light conditions, RT-qPCR was performed, and log2 fold-change was calculated (Methods). Upstream genes *MVK*, *PMVK*, *MVD*, *IDI*, *FPS*, and *crtE* exhibited little to no expression response. However, *crtYB*, *crtI*, and *crtS* expression was upregulated by UV light exposure. Notably, the *crtI* gene was the most impacted with log2 fold-changes of 0.90 (Figure 3a). These results demonstrate that light, particularly UV, transcriptionally upregulates carotenogenesis, not the upstream terpenoid pathway, in *X. dendrorhous*.

Transcriptomics to elucidate *X. dendrorhous* photobiology

We then designed a transcriptomics experiment to investigate the genome-wide response of *X. dendrorhous* across the light spectrum including UV, blue, green, and red light. We included a positive control condition of oxidative stress by hydrogen peroxide exposure, and growth of cells in the dark was used as a negative control (Methods) (Figure 4a). We extracted the RNA, which was then sequenced and processed by the DOE Joint Genome Institute through their Community Science Program. We defined a noteworthy log2 fold-change as any greater than 0.50.

The data corroborated our previous evidence that exposure to UV light leads to

upregulated expression of crtI with a log2 fold-change of 0.90. Further, the upstream terpenoid

pathway again had no significant changes, indicating upstream terpenoid synthesis to the

diterpenoid intermediate, geranylgeranyl pyrophosphate, was unaffected by light. In contrast,

oxidative stress upregulated genes in both pathways (*MVK*, *PMVK*, *IDI*, *FPS*, *crtYB*, and *crtI*),

with *IDI* having the greatest increase (log2 fold-change of 1.79) (Figure 4b). Induction of oxidative

stress may be a method to overcome the bottleneck that *IDI* imposes on the mevalonate

pathway.[15] Overall, the results corroborate the previous evidence that UV light upregulates the

final three steps of carotenogenesis in *X. dendrorhous* and show that oxidative stress broadly

upregulates terpenoid biosynthesis.

We then looked for possible light response systems in *X. dendrorhous.* There are several

known classes of fungal light receptors, particularly white collar and cryptochrome proteins in

ascomycete and two receptors in basidiomycetes, BLUF, which senses to blue light, and

rhodopsin, which senses green light.[16] We searched our *X. dendrorhous* CBS 6938 genome with

BLAST for putative fungal photoreceptor proteins within the BLUF, rhodopsin, white collar, and

cryptochrome families. Photoreceptor proteins predicted to be in the same family were

differentiated with arbitrarily assigned numbers. We found that the putative cryptochromes

increased transcription most dramatically in UV and blue light exposure. To our knowledge, this

is the first evidence of cryptochrome light response systems in basidiomycetes. Additionally,

putative white collar and rhodopsin genes increased expression. Analysis of the oxidative stress

positive control corroborated that these genes were responding specifically to light because most

putative photoreceptors were downregulated or unaffected by hydrogen peroxide (Figure 4c).

Finally, we filtered the whole genome transcriptomic data for the genes most up- or

downregulated by UV light. We identified ten upregulated and six downregulated genes with log2

fold-changes of 1.50 or greater (Figure 4d). These genes were from different functional classes,

from cell survival to energy storage. Endonuclease III, MSS4-like protein, and short-chain

dehydrogenases are directly involved in response to UV stress through DNA repair, cell survival

through cell cycle regulation and actin cytoskeleton formation, and induction of light-regulated

pathways, respectively.[17–19] These functions collectively correspond to an expected stress

response from UV exposure.[20] Conidiation-specific protein 6, a protein involved in the

reproductive cycle, is also upregulated after blue-light exposure in *Neurospora crassa*.[21]

Unexpectedly, UV light upregulates expression of glycogenin glucosyltransferase, an enzyme that

initiates glycogen nucleation. Downregulated genes were found to be involved in transcriptional

regulation, fungal apoptotic-like cell death, and the electron transport chain. Also of note, four of

the ten upregulated genes were predicted to result in hypothetical proteins, indicating that our

understanding of light response mechanisms is incomplete. Genes most up- or downregulated in

response to hydrogen peroxide exposure were found to not be affected by light treatments, further

supporting the notion of separate response pathways (Figure 4e). In summary, it appears that *X.*

dendrorhous activates DNA repair, survival, and reproductive mechanisms while downregulating

programmed cell death. The metabolic response is complex, with some pathways upregulated

and others downregulated.

To more clearly analyze the metabolic response, genes were clustered into functional

groups using the ERGO™ Bioinformatics platform from Igenbio. Upregulation of functional groups

is depicted in Figure 5a. Generally, DNA repair and amino acid metabolic functions were

upregulated. Interestingly, aromatic amino acid degradation and modification were upregulated

by both UV light and hydrogen peroxide, but amino acid uptake was only upregulated in UV-

exposed cells. Furthermore, electron transport, glycolysis, gluconeogenesis, and carbohydrate

metabolism were downregulated (Figure 5b). These results support the general trends observed

in the analysis of individual genes. We synthesized all of these observations into a qualitative

description of the CBS 6938 metabolic shift in response to UV (Figure 5c). With this view, the

transcriptional response indicates that *X. dendrorhous* likely redirects flux from central carbon
metabolism to carotenoids and aromatic amino acids upon UV exposure. Aromatic amino acids
absorb UV and are also precursors to other UV light absorbing molecules like melanin and
mycosporine compounds.[22–24] Yet, the activation of aromatic amino acid pathways in response to
UV has not been previously observed in basidiomycetes. These results indicate that the response
to UV in *X. dendrorhous* is more extensive and complex than acting on carotenogenesis alone.
Furthermore, the significant fraction of hypothetical proteins hints at additional unknown
mechanisms that also contribute to the response to UV exposure.

Transcriptomics-driven part selection allows simultaneous derivation of constitutive and
regulated genetic parts

We then leveraged the transcriptomics data to derive functional genetic elements for use
in our established modular cloning scheme. Since the data covered a variety of conditions, we
reasoned that it should be possible to derive putative constitutive and regulated promoters. We
identified strong constitutive promoters by sorting each transcriptomics dataset by fragments per
kilobase of exon per million mapped fragments (FPKM) count, choosing the genes that had the
highest and consistent FPKM counts across conditions (Figure 6a), and using our genome
sequence to obtain the sequences upstream of those genes. We also obtained the sequences
upstream of the genes in Figure 4c that were the most activated or repressed by UV light (Figure
6a). Promoter lengths were chosen with the assistance of an analysis conducted by JGI. These
putative promoters were then cloned into our modular cloning system to drive expression of
nanoluciferase and integrated into the genome (Figure 6b).

We measured expression of each construct via a luciferase assay in the dark and under
UV exposure (Methods). As Figure 6c shows, the constitutive promoters result in high
luminescence that does not change between the dark and the light. In contrast, the putative UV

activated promoters showed increased luminescence in response to UV (Figure 6c). However,

most of these increases were not large. The two strongest inducible promoters were found to be

hypothetical proteins HP3 and HP2 with log2 fold-changes of 3.01 and 1.47, respectively. This

means that the strongest UV-inducible promoter results in a 9-fold change in gene expression.

The putative UV repressed promoters did not show any significant repression. We reasoned that

the long half-life of luciferase was confounding measurement of repression. Thus, we designed a

qPCR experiment, and the results indicated no change. Therefore, it is possible our promoter

lengths were insufficient to capture the regulatory elements in these promoters.

Taken together, our results show that transcriptomics driven part discovery can

consistently derive strong constitutive promoters, but inconsistently derive regulated promoters

due to a variety of possible factors. Even so, we were able to discover two promoters that can be

activated by UV. Thus, our part discovery efforts generated a collection of constitutive and

regulated promoters that are compatible with our modular cloning workflow for *X. dendrorhous*

CBS 6938.

DISCUSSION

In this study, a promising nonconventional yeast was subjected to a streamlined set of

genomics and genetics experiments that transformed it into a powerful platform for genetic

engineering. This was done by obtaining a high-quality genome, increasing transformation

efficiency, developing advanced modular genetic tools, measuring gene regulation, and deriving

a genetic parts collection from transcriptomic data. Now, it is possible to interrogate the unique

metabolism and photobiology of *X. dendrorhous* in a similar manner to other nonconventional

yeasts.

Our analysis of *X. dendrorhous* photobiology revealed a number of novel insights into the

biological mechanisms of basidiomycete light response. We found evidence of multiple different

fungal light response systems, including the first indication of cryptochromes in basidiomycetes.

We determined for the first time that light transcriptionally regulates the carotenoid, or

tetraterpenoid, pathway but not the upstream terpenoid pathway. We learned that *X. dendrorhous*

engages a variety of survival responses in response to UV, including activation of DNA repair and

reproduction pathways and repression of the fungal apoptotic-like pathway. We also observed a

complex transcriptional response in metabolism, where central carbon metabolism is repressed

in favor of aromatic amino acid and tetraterpenoid biosynthesis. Taken together, these data paint

a picture of an extensive and coordinated response to light. This complex system is likely key to

survival of *X. dendrorhous* in its ecological niche of fallen trees at high altitudes, which would

include extended periods of intense UV exposure.

It is interesting to note that our genome integration approach results in efficient on-target

insertion of constructs via homologous recombination (HR). Our findings support the results of

others that *X. dendrorhous* CBS 6938 has an active HR pathway, which is incredibly beneficial

for future strain engineering efforts. This challenges the dogma that nonconventional yeasts favor

the non-homologous end joining (NHEJ) mechanism, and argues that other yeasts besides *S.*

cerevisiae favor HR.[25,26]

Leveraging transcriptomics allowed us to derive constitutive and regulated gene

expression parts. Transcriptomics has been previously used to identify regulated promoters in

other yeasts. We built on this prior evidence and coupled it with a modular cloning standard to

create an -omics-to-parts workflow that can yield needed constitutive and inducible promoters for

construction of genetic devices. This -omics-to-parts workflow can be generalized to derive

genetic parts collections for a number of nonconventional microbes.

Development of novel production platform hosts requires an understanding of genome

composition and genetic regulation, which also is essential for genetic tool and genetic part

development. This work demonstrates how an integrated genomics and genetics approach can

simultaneously deliver new biological understanding and onboard a nonconventional organism,

particularly a nonconventional yeast with no closely related model organism. This integrated

approach, consisting of a small number of experiments, promises to accelerate organism

onboarding efforts.

 MATERIAL AND METHODS

Strains and media

The *X. dendrorhous* type strain CBS 6938 (ATCC 96594) was cultured at 21 °C in YPD

(YEP, Sunrise Science, 1877-1KG plus 20 g/L glucose, Alfa Aesar, A16828) media for all

experiments. For all light plate apparatus experiments, the *X. dendrorhous* seed cultures were

shielded from any external light by wrapping the flasks with aluminium foil. *X. dendrorhou*s

transformants were selected using YPD with 30 mg/L nourseothricin (Jena Bioscience, AB-101-

10ML). Solid media plates were made using 20 g/L agar (Sunrise Science, 1910-1KG). Golden

gate (Type-IIS) cloning was conducted using chemically competent *Escherichia coli* DH5α cells

(NEB, C2987H). Construction of destination vectors with the toxic selection marker ccdB was

done with chemically competent One Shot ccdB Survival 2 T1R *E. coli* cells (Invitrogen, A10460).

Both *E. coli* strains were grown in 25 g/L LB Miller broth (Fisher Scientific, BP1426-2) at 37°C.

Antibiotic selection was carried out using 100 mg/L carbenicillin (Alfa Aesar, J61949), 25 mg/L

chloramphenicol (Alfa Aesar, B20841), and 50 mg/L kanamycin (Alfa Aesar, J61272) for level 0,

level 1, and level 2 assemblies, respectively.

gDNA isolation and sequencing

Isolation of gDNA from *X. dendrorhous* and next-generation sequencing was performed

as described by Collins et al.[10]

Construction of a light plate apparatus for various light wavelengths

The light plate apparatus (LPA) was assembled following a user's manual published by

the Tabor lab at Rice University.[27] The LPA is an instrument capable of shining two individual LED

lights on cell cultures in a 24-well plate. It is comprised of a 3D-printed shell surrounding a

soldering board with 48 LED light sockets oriented below a 24-well plate with a clear bottom

(AWLS-324042, Arctic White). The LPA allows each culture to be exposed to two unique LED

lights without disrupting neighboring cultures.

RNA isolation from the light plate apparatus

LPA cultures were started in 2 mL of YPD media at OD_{600} = 1 and shaken for 4 hours at

200 rpm while exposed to one red (λ = 660 nm), green (λ = 565 nm), blue (λ = 470 nm), white

(visible light spectrum), or UV LED light (λ = 400 nm), or no LED light. For the transcriptome

sequencing experiments, the hydrogen peroxide condition replaced the white light condition. The

hydrogen peroxide condition consisted of YPD media with hydrogen peroxide at a concentration

of 10 mM and had no LED light. 1 mL of *X. dendrorhous* culture was taken forward for RNA

isolation, which used a cell homogenization and TRIzol-based method. *X. dendrorhous* cells were

first resuspended in 200 µL of cell lysis buffer (0.5 M sodium acetate, 5% SDS, 1 mM EDTA) and

transferred to pre-filled tubes with 400-micron zirconium beads (OPS diagnostics, PFMB 400-

100-34). The cells were homogenized using a FastPrep-24 machine from MP Biomedical for two

cycles at 6 m/s for 30 seconds each, with a 1-minute incubation on ice between each cycle. 800

µL of Trizol reagent (Invitrogen, 15596026) was then added to the cell lysate and incubated on

ice for 10 minutes, followed by two more cycles on the FastPrep machine. On ice, 100 µL of 1–

bromo–3–chloropropane (Sigma Aldrich, B9673-200ML) was added to the cell lysate, inverted to

mix, and incubated for 5 minutes. The tubes were spun down at 4°C for 14,000 x g for 15 minutes.

400 µL of supernatant was transferred to 400 µL of 100% ethanol. The RNA was purified using

the Monarch Total RNA Miniprep Kit (NEB, T2010S) following the supplier's instructions.

qRT-PCR evaluation of carotenogenesis pathway activity in response to light exposure

RNA samples were converted to cDNA using the Protoscript II First Strand cDNA

Synthetsis Kit (NEB, E6560S) following the supplier's standard protocol instructions. RT-qPCR

was then performed using IDT's PrimeTime Gene Expression Master Mix (Integrated DNA

Technologies Inc., Skokie, Illinois, 1055772) and PrimeTime qPCR pre-mixed assay with probes

and primers. Sequences can be found in Supplementary Table 2. Probes targeting genes-of-

interest were labeled with FAM fluorophores and probes targeting the actin housekeeping gene

were labeled with HEX fluorophores. Primers were designed on Benchling (https://benchling.com/)

and optimal probe placement was determined using PrimerQuest

(https://www.idtdna.com/PrimerQuest). The QuantStudio 6 Flex system was used, and

instructions provided by IDT for the standard cycling protocol were followed.

RNA transcriptome sequencing and differential gene expression analysis

cDNA prep and sequencing protocol JGI method for cDNA preparation and sequencing.

Raw FASTQ file reads were filtered and trimmed using the Joint Genome Institute (JGI) QC

pipeline.[28] Filtered reads from each library were aligned to the reference genome using HISAT

version 0.1.4-beta.[29] featureCounts was used to generate the raw gene counts file using gff3

annotations.[30] Raw gene counts were used to evaluate the level of correlation between biological

replicates using Pearson's correlation and determine which replicates would be used in the DGE

analysis. DESeq2 (version 1.10.0) was subsequently used to determine which genes were

differentially expressed between pairs of conditions.[31] The parameter used to call a gene

differentially expressed between conditions was a P-value < 0.05.

Gene set enrichment analysis

Annotated *X. dendrorhous* CBS 6938 genome and gene count table from mapped

transcriptomic reads were uploaded to ERGO 2.04 (Igenbio, inc).[32] Genes were functionally

enriched according to ERGO groups using GAGE5.[33]

Polymerase Chain Reaction (PCR)

All PCRs were done using Q5 2X Master Mix (NEB, M0492L). Primers were designed on

Benchling (https://benchling.com/) and the NEB Tm Calculator (https://tmcalculator.neb.com/). All

primers were ordered from IDT. PCR reactions closely followed NEB instructions. Briefly,

reactions were done in 50 µL total volume; 25 µL Q5 Master Mix, 2.5 µL of each primer, X µL of

template DNA (1 ng plasmid DNA or 100 ng genomic DNA), and 20-X µL nuclease free water

(VWR 02-0201-0500). Reactions were run on a thermocycler with the following settings:

1. 98°C for 30 sec

2. 30 PCR cycles:

(a) 98°C for 10 sec

(b) annealing temp. for 15 sec

(c) 72°C for 20 sec per kbp

3. 72°C for 2 min

4. 10°C hold

Gibson cloning

Gibson assembly was used to construct destination vectors, which were designed for

TypeIIS-based cloning. Reactions closely followed instructions from the NEBuilder HiFi DNA

Assembly Master Mix (NEB, E2621S). Fragments were amplified with PCR and designed to have

20-30 bp overlaps. The DpnI enzyme was used to digest template plasmid or *X. dendrorhous*

genomic DNA according to the manufacturer's instructions (NEB, R0176S). The amplified

fragments were purified using the Zymo Clean & Concentrator Kit (Zymo, D4005), followed by

dilution to 0.2 pmols for 2-3 fragments or 0.5 pmols for 4 or more fragments. The fragments, 10

µL of HiFi master mix, and nuclease free water were combined in a PCR tube for a 20 µL total

reaction volume. The mixture was run on a thermocycler at 50°C for 60 minutes, with a hold at

10°C.

TypeIIS Cloning

A modular, hierarchical TypeIIS cloning scheme was used construct genetic designs for

integration into *X. dendrorhous*. This scheme consisted of three levels; transcriptional parts (level

0), transcription units (level 1), and integrative pathways (level 2). Cloning reactions were based

on the enzymes BbsI (Thermo Scientific, ER1011) or BsaI (NEB, R3733). Cloning reactions

consisted of 1 µL of N parts, 1 µL of BsaI or BbsI, 1 µ L of 10X Ligase Buffer, 0.4 µL of T4 DNA

ligase (Promega, M1794), and 7.9 - N µL of nuclease free water, where N is the number of parts.

Reactions were run on a thermocycler at 37°C for 5 hours, 50 °C for 15 minutes, 80 °C for 20

minutes, and a hold at 10°C.

Parts and strain design

All genetic parts used in this study can be found in Supplementary Table 1.

Minimum Inhibitory Concentration Assay

X. dendrorhous cultures were diluted with YPD to an OD_{600} = 1.0 and 100 µL was pipetted

into columns 2-11 of a 96-well plate (Corning, 3596). Column 11 served as a positive control of

only X. dendrorhous culture. A negative control of 100 µL YPD was pipetted into column 12. The

antifungals hygromycin (ThermoFisher, 10687010), zeocin (Jena Bioscience, AB-103S),

geneticin (ThermoFisher, 10131-035), and nourseothricin (Jena Bioscience, AB-101L) were

chosen for the experiment. Stocks of each antifungal were made in concentrations of 5120 µg/mL

and 7680 µg/mL. Three replicates of antifungal concentration were made by pipetting 200 µL of

an antifungal into three wells of column 1. A multichannel pipette was used to perform a serial

dilution to column 10. At column 10, 100 µL of the culture and antibiotic mixture was pipetted and

discarded, leaving 100 µL of mixture in every well. The 96-well plates were incubated at 21°C for

3 days. The OD_{600} was measured on a BioTek Synergy H1 plate reader with column 12 as a

blank. Outliers were calculated using an interquartile range and were excluded in survival percent

calculations.

***X. dendrorhous* transformation**

Integrative pathway DNA for transformation was excised from level 2 vectors with a BsaI

digestion. This digestion was comprised of 40 µL plasmid DNA, 5 µL 10X CutSmart Buffer, 2 µL

BsaI, and 3 µL nuclease free water. The reaction was run at 37°C for 10 hours, 80°C for 20

minutes, and a hold at 10°C. The resulting DNA fragments were then purified with the Zymo Clean

& Concentrator Kit and were ready for transformation. *X. dendrorhous* transformations were

based on an electroporation method described by Visser et. al.[34] *X. dendrorhous* was first

streaked on a YPD agar plate and grown at 21°C for approximately 2 days. A single colony was

used to inoculate 50 mL YPD media in a 125 mL Erlenmeyer flask. The cells were shaken at 21°C

and 200 rpm for another 2 days. These cells were then used to inoculate 200 mL of YPD in a 1 L

Erlenmeyer flask at OD_{600} = 0.02. The cells were grown at 21°C and 200 rpm until reaching OD_{600}

= 1.2 (approximately 20 hours). From here, the cells were pelleted at 1500 x g for 5 min and

resuspended in 25 mL of freshly made 50 mM potassium phosphate buffer (pH=7.0, Sigma-

Aldrich, P8281-100G & P9791-100G) with 25 mM diothiothreitol (Acros Organics, 426380500).

The cells were incubated at room temperature for 15 minutes, and then pelleted at 4°C for 5

minutes at 1500 x g. The cells were then washed with 25 mL of ice cold STM buffer (pH = 7.5,

270 mM sucrose, Millipore Sigma, 1.07651.5000, 10 mM Tris HCl, Alfa Aesar, J67233, 1 mM

MgCl$_2$, VWR, E525-100ML) and spun down again at 4°C for 5 minutes at 1500 x g. This wash

step was repeated and followed by resuspension of the pellet in 500 μL of STM buffer. Now

electrocompetent, the cells were divided into 60 μL aliquots and kept on ice until electroporation.

10 μL of transforming DNA was mixed with a 60 μL aliquot and transferred to an ice cold 0.2 cm

electroporation cuvette (ThermoFisher, 21-237-2). Electroporation was performed with a Gene

Pulser electroporator (Bio-Rad, 1998.018.1, 1998.018.2, & 1998.018.3) at 0.8 kV, 1000 ohms,

and 25 μF. Immediately following the electric pulse, 500 μL of ice cold YPD was added to the

cuvette. This mixture was then transferred to 4 mL of YPD in a 14 mL Falcon tube and grown

overnight on a rotating drum. The next morning, the cells were spun down for 5 minutes at 1500

x g, resuspended in 500 mL of filtered water, and spread onto selection plates. The plates were

incubated at 21°C until colonies appeared, which typically occurred in 2-3 days.

Astaxanthin extraction from wild-type and *crtYB* knockouts

X. dendrorhous CBS 6938 wild-type and *crtYB* knockout strains were grown for 3 days in

the dark at 21°C and 200 rpm. 1 mL of culture was taken forward and centrifuged at 10,850 x g

for 10 minutes. The cell pellet was washed with Milli-Q filtered water twice before resuspension

in 1 mL of acetone (SigmaAldrich, 34850-1L). Cell and acetone mixtures were poured into pre-

filled tubes of 400-micron zirconium beads (OPS Diagnostics, PFMB 400-100-34), then

transferred to snap-cap microcentrifuge tubes (Eppendorf, 022363743). The Bullet Blender Storm

Pro tissue homogenizer (NextAdvance, BT24M) was used at 12 m/s for 5 minutes to mechanically

lyse the cells. Afterward, tubes were centrifuged for 10,850 x g for 10 minutes. 200 μL of

supernatant was pipetted into a nylon syringeless filter (Whatman, UN203NPENYL) and

transferred to a 2 mL chromatography vial (Agilent, 5181-3376) with a glass vial insert (5183-

2085) and crimp top cap (Agilent, 8010-0051).

Ultra-high performance liquid chromatography detection of astaxanthin

430 UPLC analysis was performed using a modified method from Bohoyo-Gil et al.[35]

431 Modifications include an injection volume of 10 µL, usage of a Shimadzu Nexcol C18 1.8 µm

432 column (Shimadzu, 220-91394-03), and analysis on the Nexera Series UPLC (Shimadzu; RF-

433 20AXS, RID-20A, SCL-40, DGU-403, DGU-405, CTO-40C, SPD-M40, C-40 LPGE, LC-40D XS,

434 SIL-40C XS).

Bioluminescence detection assay

LPA cultures were started in 2 mL of YPD media at OD_{600} = 1 and shaken for 24 hours at

200 rpm. *X. dendrorhous* was exposed to either a UV (λ = 400 nm) or dark (no LED light) condition.

Bioluminescence was detected using the Nano-Glo Luciferase Assay System (Promega, N1110).

Assay buffer and assay reagent were mixed in a 50:1 ratio to create to assay solution. Aliquots of

50 µL of *X. dendrorhous* cells were mixed with 50 µL assay solution in a 96-well plate (Corning,

3904). Bioluminescence was measured on a BioTek Synergy H1 plate reader.

FUNDING

This project is supported by a National Science Foundation CAREER award to EMY,

award number 1944046. EET is supported by a National Science Foundation Graduate Research

Fellowship.

The work (proposal 10.46936/10.25585/60001246) conducted by the U.S. Department of

Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility,

is supported by the Office of Science of the U.S. The Department of Energy operated under

Contract No. DE-AC02-05CH11231.

Development of Prymetime was supported by the Office of the Director of National

Intelligence (ODNI), Intelligence Advanced Research Projects Activity (IARPA) under Finding

Engineering Linked Indicators (FELIX) program contract #N66001-18-C-4507. The views and

conclusions contained herein are those of the authors and should not be interpreted as

necessarily representing the official policies, either expressed or implied, of ODNI, IARPA, or the

U.S. Government. The U.S. Government is authorized to reproduce and distribute reprints for

governmental purposes notwithstanding any copyright annotation therein.

AUTHOR CONTRIBUTIONS

EET, JHC, and EMY conceived of the study. EET and JHC performed all experiments and analysis save the sequencing and initial processing of RNA-seq data. CBM and GTN constructed the light plate apparatus (LPA). KM acquired microscope images of red yeast. AL, SM, and IG performed transcriptomic sequencing and initial processing of RNA-seq data. EET and EMY wrote the manuscript.

*Emma E. Tobin and Joseph H. Collins contributed equally.

458   **CONFLICT OF INTEREST**

459   The authors declare that there are no conflicts of interest regarding this research study.

 FIGURES

a.

Xanthophyllomyces dendrorhous
CBS 6938

b.

462 Figure 1. **Overview of *X. dendrorhous* genome sequencing and quality analysis. (a)**
463 schematic of Prymetime next generation sequencing workflow. Illumina and Nanopore reads are
464 combined for a higher quality genome. **(b)** BUSCO analysis of the *X. dendrorhous* genome.

Figure 2. **"Onboarding" process of *X. dendrorhous*. (a)** Minimum inhibitory concentration assay data from testing geneticin, zeocin, hygromycin, and nourseothricin against *X. dendrorhous*. **(b)** Cloning workflow adapted from CIDAR for homologous recombination into the *crtYB* site. **(c)** Depiction of high-efficiency transformation procedure. **(d)** Overlapped chromatograms illustrating the lack of astaxanthin at a retention time of 0.998. Wild type is shown in red and the *crtYB* knockout is shown in black. **(e)** Comparison of luminescence between two strong constitutive promoters for *act* and *adh4*, the *crtYb* knockout, and CBS 6938 wild type.

Figure 3. **Preliminary investigation into ultraviolet light regulation through RT-qPCR. (a)** RT-qPCR results for enzymatic steps in the mevalonate and carotenogenesis pathway. Upstream steps are not affected by UV exposure, but downstream genes *crtYB*, *crtI*, and *crtS* are considerably induced. **(b)** RNA isolation process for transcriptomic sequencing. Six treatments were implemented including four light wavelengths, a positive control (H_2O_2), and a negative control (dark).

a.

b.

Carotenogenesis genes

c.

Putative photoreceptor genes

d.

Top UV differentially expressed genes

e.

Top H₂O₂ differentially expressed genes

Figure 4. **Heatmaps illustrating the response of gene expression to various light treatments**
and a general oxidative stressor, H₂O₂. Blue represents induction and red represents
repression. (a) Regulation of mevalonate and carotenogenesis genes by light and H₂O₂. **(b)**
Regulation of putative photoreceptors. Numbers are arbitrarily assigned. **(c)** Top ten UV induced
and repressed genes. **(d)** Top ten H₂O₂ induced and repressed genes.

Figure 5. **Transcriptomic analyses clustered by ERGO functional groups, and therefore**
metabolic processes. (a) Overall up- and downregulation of metabolic processes by UV light or
H_2O_2. **(b)** Summary of metabolic processes regulated by UV light. Generally, metabolic flux is
being drawn from central carbon metabolism to synthesis of aromatic amino acids.

492
Figure 6. **Comparison of transcriptome-derived promoters to predicted expression through**
FPKM values. (a) Putative **c**onsitutive promoters are shown. **(b)** Putative nducible promoters are
shown. **(c)** Putative epressible promoters are shown. **(d)** Depiction of putative inducible promoters
with log2 fold-change. P_{HP3} has the greatest induction to UV light at a log2 fold-change of 3.01.
(e) Depiction of putative repressible promoters with log2 fold-change. None were found to be
repressible.

REFERENCES

(1)   Boekhout, T.; Amend, A. S.; El Baidouri, F.; Gabaldón, T.; Geml, J.; Mittelbach, M.; Robert, V.; Tan, C. S.; Turchetti, B.; Vu, D.; Wang, Q. M.; Yurkov, A. Trends in Yeast Diversity Discovery. *Fungal Diversity*. 2022. https://doi.org/10.1007/s13225-021-00494-6.

(2)   Wagner, J. M.; Alper, H. S. Synthetic Biology and Molecular Genetics in Non-Conventional Yeasts: Current Tools and Future Advances. *Fungal Genetics and Biology* **2016**, *89*. https://doi.org/10.1016/j.fgb.2015.12.001.

(3)   Bellora, N.; Moliné, M.; David-Palma, M.; Coelho, M. A.; Hittinger, C. T.; Sampaio, J. P.; Gonçalves, P.; Libkind, D. Comparative Genomics Provides New Insights into the Diversity, Physiology, and Sexuality of the Only Industrially Exploited Tremellomycete: Phaffia Rhodozyma. *BMC Genomics* **2016**, *17* (1). https://doi.org/10.1186/s12864-016-3244-7.

(4)   Nielsen, J. Yeast Systems Biology: Model Organism and Cell Factory. *Biotechnology Journal*. 2019. https://doi.org/10.1002/biot.201800421.

(5)   Li, W.; Cui, L.; Mai, J.; Shi, T. Q.; Lin, L.; Zhang, Z. G.; Ledesma-Amaro, R.; Dong, W.; Ji, X. J. Advances in Metabolic Engineering Paving the Way for the Efficient Biosynthesis of Terpenes in Yeasts. *Journal of Agricultural and Food Chemistry*. 2022. https://doi.org/10.1021/acs.jafc.2c03917.

(6)   Torres-Haro, A.; Verdín, J.; Kirchmayr, M. R.; Arellano-Plaza, M. Metabolic Engineering for High Yield Synthesis of Astaxanthin in Xanthophyllomyces Dendrorhous. *Microbial Cell Factories*. 2021. https://doi.org/10.1186/s12934-021-01664-6.

(7)   Rodríguez-Sáiz, M.; De La Fuente, J. L.; Barredo, J. L. Xanthophyllomyces Dendrorhous for the Industrial Production of Astaxanthin. *Applied Microbiology and Biotechnology*. 2010. https://doi.org/10.1007/s00253-010-2814-x.

(8)   Tropea, A.; Gervasi, T.; Melito, M. R.; Curto, A. Lo; Curto, R. Lo. Does the Light Influence

Astaxanthin Production in Xanthophyllomyces Dendrorhous? *Nat Prod Res* **2013**, *27* (7).

https://doi.org/10.1080/14786419.2012.688045.

(9)   Vázquez, M. Effect of the Light on Carotenoid Profiles of Xanthophyllomyces Dendrorhous

Strains (Formerly Phaffia Rhodozyma). *Food Technol Biotechnol* **2001**, *39* (2).

(10)   Collins, J. H.; Keating, K. W.; Jones, T. R.; Balaji, S.; Marsan, C. B.; Çomo, M.; Newlon, Z.

J.; Mitchell, T.; Bartley, B.; Adler, A.; Roehner, N.; Young, E. M. Engineered Yeast

Genomes Accurately Assembled from Pure and Mixed Samples. *Nat Commun* **2021**, *12*

(1). https://doi.org/10.1038/s41467-021-21656-9.

(11)   Mülleder, M.; Capuano, F.; Pir, P.; Christen, S.; Sauer, U.; Oliver, S. G.; Ralser, M. A

Prototrophic Deletion Mutant Collection for Yeast Metabolomics and Systems Biology.

Nature Biotechnology. 2012. https://doi.org/10.1038/nbt.2442.

(12)   Iverson, S. V.; Haddock, T. L.; Beal, J.; Densmore, D. M. CIDAR MoClo: Improved MoClo

Assembly Standard and New E. Coli Part Library Enable Rapid Combinatorial Design for

Synthetic and Traditional Biology. *ACS Synth Biol* **2016**, *5* (1).

https://doi.org/10.1021/acssynbio.5b00124.

(13)   England, C. G.; Ehlerding, E. B.; Cai, W. NanoLuc: A Small Luciferase Is Brightening Up

the Field of Bioluminescence. *Bioconjugate Chemistry*. American Chemical Society May

18, 2016, pp 1175–1187. https://doi.org/10.1021/acs.bioconjchem.6b00112.

(14)   Bereta, M.; Teplan, M.; Chafai, D. E.; Radil, R.; Cifra, M. Biological Autoluminescence as

a Noninvasive Monitoring Tool for Chemical and Physical Modulation of Oxidation in Yeast

Cell Culture. *Sci Rep* **2021**, *11* (1). https://doi.org/10.1038/s41598-020-79668-2.

(15)   Shin, J.; South, E. J.; Dunlop, M. J. Transcriptional Tuning of Mevalonate Pathway
545   Enzymes to Identify the Impact on Limonene Production in Escherichia Coli. *ACS Omega*
546   **2022**, *7* (22). https://doi.org/10.1021/acsomega.2c00483.

547   (16)   Yu, Z.; Fischer, R. Light Sensing and Responses in Fungi. *Nature Reviews Microbiology*.
548   2019. https://doi.org/10.1038/s41579-018-0109-x.

549   (17)   Desrivières, S.; Cooke, F. T.; Parker, P. J.; Hall, M. N. MSS4, a Phosphatidylinositol-4-
550   Phosphate 5-Kinase Required for Organization of the Actin Cytoskeleton in
551   Saccharomyces Cerevisiae. *Journal of Biological Chemistry* **1998**, *273* (25), 15787–15793.
552   https://doi.org/10.1074/jbc.273.25.15787.

553   (18)   Serafini, D. M.; Schellhorn, H. E. Endonuclease III and Endonuclease IV Protect
554   Escherichia Coli from the Lethal and Mutagenic Effects of Near-UV Irradiation. *Can J*
555   *Microbiol* **1999**, *45* (7). https://doi.org/10.1139/cjm-45-7-632.

556   (19)   Bruchez, J. J. P.; Eberle, J.; Kohler, W.; Kruft, V.; Radford, A.; Russo, V. E. A. Bli-4, a Gene
557   That Is Rapidly Induced by Blue Light, Encodes a Novel Mitochondrial, Short Chain Alcohol
558   Dehydrogenase-like Protein in Neurospora Crassa. *Molecular and General Genetics* **1996**,
559   *252* (3). https://doi.org/10.1007/BF02173767.

560   (20)   Sinha, R. P.; Häder, D. P. UV-Induced DNA Damage and Repair: A Review. *Photochemical*
561   *and Photobiological Sciences* **2002**, *1* (4), 225–236. https://doi.org/10.1039/b201230h.

562   (21)   Olmedo, M.; Ruger-Herreros, C.; Luque, E. M.; Corrochano, L. M. A Complex
563   Photoreceptor System Mediates the Regulation by Light of the Conidiation Genes Con-10
564   and Con-6 in Neurospora Crassa. *Fungal Genetics and Biology* **2010**, *47* (4), 352–363.
565   https://doi.org/10.1016/j.fgb.2009.11.004.

566   (22)   Llewellyn, C. A.; Greig, C.; Silkina, A.; Kultschar, B.; Hitchings, M. D.; Farnham, G.
567   Mycosporine-like Amino Acid and Aromatic Amino Acid Transcriptome Response to UV

and Far-Red Light in the Cyanobacterium Chlorogloeopsis Fritschii PCC 6912. *Sci Rep*
2020, *10* (1). https://doi.org/10.1038/s41598-020-77402-6.

(23)   Gilchrest, B. A.; Park, H. Y.; Eller, M. S.; Yaar, M. Echanisms of Ultraviolet Light-Induced
Pigmentation. *Photochemistry and Photobiology*. American Society for Photobiology 1996,
pp 1–10. https://doi.org/10.1111/j.1751-1097.1996.tb02988.x.

(24)   Bhatia, S.; Garg, A.; Sharma, K.; Kumar, S.; Sharma, A.; Purohit, A. P. Mycosporine and
Mycosporine-like Amino Acids: A Paramount Tool against Ultra Violet Irradiation.
Pharmacognosy Reviews. 2011. https://doi.org/10.4103/0973-7847.91107.

(25)   Schwartz, C.; Shabbir-Hussain, M.; Frogue, K.; Blenner, M.; Wheeldon, I. Standardized
Markerless Gene Integration for Pathway Engineering in Yarrowia Lipolytica. *ACS Synth
Biol* **2017**, *6* (3). https://doi.org/10.1021/acssynbio.6b00285.

(26)   Schwarzhans, J. P.; Wibberg, D.; Winkler, A.; Luttermann, T.; Kalinowski, J.; Friehs, K.
Non-Canonical Integration Events in Pichia Pastoris Encountered during Standard
Transformation   Analysed   with   Genome   Sequencing.   *Sci   Rep*   **2016**,   *6*.
https://doi.org/10.1038/srep38952.

(27)   Gerhardt, K. P.; Olson, E. J.; Castillo-Hair, S. M.; Hartsough, L. A.; Landry, B. P.; Ekness,
F.; Yokoo, R.; Gomez, E. J.; Ramakrishnan, P.; Suh, J.; Savage, D. F.; Tabor, J. J. An
Open-Hardware Platform for Optogenetics and Photobiology. *Sci Rep* **2016**, *6*.
https://doi.org/10.1038/srep35363.

(28)   Clum, A.; Huntemann, M.; Bushnell, B.; Foster, B.; Foster, B.; Roux, S.; Hajek, P. P.;
Varghese, N.; Mukherjee, S.; Reddy, T. B. K.; Daum, C.; Yoshinaga, Y.; O'Malley, R.;
Seshadri, R.; Kyrpides, N. C.; Eloe-Fadrosh, E. A.; Chen, I.-M. A.; Copeland, A.; Ivanova,
N.   N.   DOE   JGI   Metagenome   Workflow.   *mSystems*   **2021**,   *6*   (3).
https://doi.org/10.1128/msystems.00804-20.

(29)   Kim, D.; Langmead, B.; Salzberg, S. L. HISAT: A Fast Spliced Aligner with Low Memory

593   Requirements. *Nat Methods* **2015**, *12* (4). https://doi.org/10.1038/nmeth.3317.

594   (30)   Liao, Y.; Smyth, G. K.; Shi, W. FeatureCounts: An Efficient General Purpose Program for

595   Assigning Sequence Reads to Genomic Features. *Bioinformatics* **2014**, *30* (7).

596   https://doi.org/10.1093/bioinformatics/btt656.

597   (31)   Love, M. I.; Huber, W.; Anders, S. Moderated Estimation of Fold Change and Dispersion

598   for RNA-Seq Data with DESeq2. *Genome Biol* **2014**, *15* (12).

599   https://doi.org/10.1186/s13059-014-0550-8.

600   (32)   Overbeek, R.; Larsen, N.; Walunas, T.; D' Souza, M.; Pusch, G.; Selkov, E.; Liolios, K.;

601   Joukov, V.; Kaznadzey, D.; Anderson, I.; Bhattacharyya, A.; Burd, H.; Gardner, W.; Hanke,

602   P.; Kapatral, V.; Mikhailova, N.; Vasieva, O.; Osterman, A.; Vonstein, V.; Fonstein, M.;

603   Ivanova, N.; Kyrpides, N. The ErgoTM Genome Analysis and Discovery System. *Nucleic*

604   *Acids Research*. 2003. https://doi.org/10.1093/nar/gkg148.

605   (33)   Luo, W.; Friedman, M. S.; Shedden, K.; Hankenson, K. D.; Woolf, P. J. GAGE: Generally

606   Applicable Gene Set Enrichment for Pathway Analysis. *BMC Bioinformatics* **2009**, *10*.

607   https://doi.org/10.1186/1471-2105-10-161.

608   (34)   Visser, H.; Van Ooyen, A. J. J.; Verdoes, J. C. Metabolic Engineering of the Astaxanthin-

609   Biosynthetic Pathway of Xanthophyllomyces Dendrorhous. *FEMS Yeast Research*. 2003.

610   https://doi.org/10.1016/S1567-1356(03)00158-2.

611   (35)   Bohoyo-Gil, D.; Dominguez-Valhondo, D.; García-Parra, J. J.; González-Gómez, D.

612   UHPLC as a Suitable Methodology for the Analysis of Carotenoids in Food Matrix.

613   *European Food Research and Technology* **2012**, *235* (6). https://doi.org/10.1007/s00217-

614   012-1838-0.

615