Neural substrates of cold nociception in *Drosophila* larva

Atit A. Patel[1]*, Albert Cardona[2,3,4], and Daniel N. Cox[1]*

[1]Neuroscience Institute, Georgia State University, Atlanta, GA, USA

[2]HHMI Janelia Research Campus, Ashburn, VA, USA

[3]MRC Laboratory of Molecular Biology, Cambridge, UK

[4]Department of Physiology, Development, and Neuroscience, University of Cambridge, UK

* Correspondence:
Corresponding Authors
apatel221@gsu.edu; dcox18@gsu.edu

Keywords: cold nociception, thermosensation, somatosensory circuits, *Drosophila*.

Abstract

Metazoans detect and differentiate between innocuous (non-painful) and/or noxious (harmful) environmental cues using primary sensory neurons, which serve as the first node in a neural network that computes stimulus specific behaviors to either navigate away from injury-causing conditions or to perform protective behaviors that mitigate extensive injury. The ability of an animal to detect and respond to various sensory stimuli depends upon molecular diversity in the primary sensors and the underlying neural circuitry responsible for the relevant behavioral action selection. Recent studies in *Drosophila* larvae have revealed that somatosensory class III multidendritic (CIII md) neurons function as multimodal sensors regulating distinct behavioral responses to innocuous mechanical and nociceptive thermal stimuli. Recent advances in circuit bases of behavior have identified and functionally validated *Drosophila* larval somatosensory circuitry involved in innocuous (mechanical) and noxious (heat and mechanical) cues. However, central processing of cold nociceptive cues remained unexplored. We implicate multisensory integrators (Basins), premotor (Down-and-Back) and projection (A09e and TePns) neurons as neural substrates required for cold-evoked behavioral and calcium responses. Neural silencing of cell types downstream of CIII md neurons led to significant reductions in cold-evoked behaviors and neural co-activation of CIII md neurons plus additional cell types facilitated larval contraction (CT) responses. Further, we demonstrate that optogenetic activation of CIII md neurons evokes calcium increases in these neurons. Finally, we characterize the premotor to motor neuron network underlying cold-evoked CT and delineate the muscular basis of CT response. Collectively, we demonstrate how *Drosophila* larvae process cold stimuli through

34 functionally diverse somatosensory circuitry responsible for generating stimulus-specific
35 behaviors.
36
37
38 **Introduction**
39
Metazoans detect innocuous and/or noxious environmental cues and appropriately
generate relevant behavioral responses. There is a large diversity in types of nervous systems
from relatively simple nerve nets to highly complex centralized organs dedicated for processing
and executing behavioral commands. The ability of an organism to sense and respond to
environmental cues is based on the underlying neural architecture and its connectivity to muscle
groups for generating behavioral responses. Understanding the neural substrates underlying the
execution of how behavioral commands are generated in the nervous system, spanning from the
sensory neuron input to motor neuron output, constitutes one of the key areas of research in
contemporary neuroscience.
Drosophila melanogaster is among the premier model organisms for studying molecular,
cellular and circuit bases of behaviors in modern neuroscience driven by advances in electron
microscopy (EM) connectomics, ability to perform high-throughput behavioral screens,
characterization of stereotyped behavioral phenotypes, the ability to perform cell-type specific
manipulations and genetic accessibility (for review see (Eschbach & Zlatic, 2020). Another
critical benefit of using *Drosophila* to study neural connectivity at the synaptic level is its
relatively small, but complex CNS, where there are only ~10,000 neurons in the *Drosophila*
larval CNS and ~100,000 neurons in adult *Drosophila melanogaster* brain compared to 86
billion neurons in the human brain (Herculano-Houzel, 2009; Scheffer et al., 2020). Highly
detailed serial section transmission electron microscopy (ssTEM) whole brain volumes with
synaptic level resolution have been obtained for both *Drosophila* larval (Ohyama et al., 2015)
and adult (Zheng et al., 2018) brains, where neurons were reconstructed using a collaborative
web-based software, CATMAID (Saalfeld et al., 2009; Schneider-Mizell et al., 2016). Concerted
efforts from many laboratories have made tremendous advances in reconstructing the *Drosophila*
larval connectome at synaptic level resolution and there has been great progress in mapping out
select connectomes at the EM level (Clark et al., 2018; Eschbach & Zlatic, 2020; Kohsaka et al.,
2017). Connectomes have been further validated using behavioral and functional imaging studies
for olfaction in combination with learning and memory (Berck et al., 2016; Eichler et al., 2017;
Eschbach et al., 2020; Saumweber et al., 2018), feeding (Miroschnikow et al., 2018; Schlegel et
al., 2016), visual processing (Larderet et al., 2017), locomotion (Carreira-Rosario et al., 2018;
Fushiki et al., 2016; Heckscher et al., 2015; Hiramoto et al., 2021; Kohsaka et al., 2019; Zarin et
al., 2019; Zwart et al., 2016), chemotaxis (Tastekin et al., 2018), thermosensation (Hernandez-
Nunez et al., 2021), mechanosensation (Jovanic et al., 2016; Jovanic et al., 2019; Masson et al.,
2020), nociceptive modalities (Burgos et al., 2018; Gerhard et al., 2017; Hu et al., 2017;
Imambocus et al., 2022; Kaneko et al., 2017; Ohyama et al., 2015; Takagi et al., 2017), among

others (Andrade et al., 2019; Huckesfeld et al., 2021; Imura et al., 2020; Mark et al., 2021;
Valdes-Aleman et al., 2021; Winding et al., 2023).
We are particularly interested in how larval somatosensation functions through peripheral
sensory neurons located along the body wall just below the larval cuticle. Larval somatosensory
neurons are comprised of type I, mono-ciliated dendrites, (external sensory (es) and chordotonal
(Ch) neurons) and type II, bipolar dendritic (td and bd) and highly branched multidendritic (md)
neurons (classes I-IV, referred to as CI-IV md). Functional and behavioral roles of these sensory
neurons include proprioception (CI md & Ch) (Caldwell et al., 2003; He et al., 2018; Vaadia et
al., 2018), heat thermoreception (CIV md) (Babcock et al., 2009; Babcock et al., 2011; Im et al.,
2018; Im et al., 2015; Ohyama et al., 2015; Tracey et al., 2003), cold thermoreception (Ch, CII
md, & CIII md) (Maksymchuk et al., 2022; Maksymchuk et al., 2023; Turner et al., 2016; Turner
et al., 2018), chemoreception (CIV md) (Himmel et al., 2019; Lopez-Bellido et al., 2019) and
mechanosensation (Ch, CII md, CIII md, & CIV md) (Hu et al., 2017; Hu et al., 2020; Jovanic et
al., 2019; Kaneko et al., 2017; Masson et al., 2020; Ohyama et al., 2015; Scholz et al., 2015;
Tsubouchi et al., 2012; Yan et al., 2012; Yoshino et al., 2017). Recent studies have unraveled
circuit bases underlying thermo- (heat), chemo- and mechanosensation (Burgos et al., 2018;
Hernandez-Nunez et al., 2021; Hu et al., 2017; Hu et al., 2020; Imambocus et al., 2022; Jovanic
et al., 2016; Jovanic et al., 2019; Kaneko et al., 2017; Lopez-Bellido et al., 2019; Masson et al.,
2020; Ohyama et al., 2015; Yoshino et al., 2017), however, circuit bases of noxious cold
thermosensory evoked behaviors are yet to be described.
Neural circuitry downstream of select somatosensory neurons has been elucidated both
physiologically and behaviorally in the context of nociceptive stimuli that evoke characteristic
rolling behavior in *Drosophila* larvae. Specifically, Ch and CIV md neurons signal through
multisensory integrator neurons (Basins) via ascending pathways to command neurons for
mechanical and nociceptive stimulation that synergistically impact larval rolling escape
responses (Ohyama et al., 2015). CIV md neuron mediated nociceptive escape behaviors also
function through premotor neurons such as Down-and-Back (DnB) and medial clusters of CIV
md interneurons (mCSI) (Burgos et al., 2018; Yoshino et al., 2017). Additionally, A08n
projection neurons receive input from CIV md neuron and the peptidergic neuron DP-ilp7 to
integrate multisensory inputs from CII, CIII and CIV md neurons (Imambocus et al., 2022).
A08n and DP-ilp7 neurons are specifically required for nociceptive mechanical and chemical
sensing but not for nociceptive thermosensation (Hu et al., 2017; Imambocus et al., 2022;
Kaneko et al., 2017; Tenedini et al., 2019; Vogelstein et al., 2014). Lastly, Ch and CIII md
neuronal connectivity, through ascending projection neurons A09e and TePns as well as
premotor neuron Chair-1, are required for anemotaxis and innocuous mechanical sensing
(Jovanic et al., 2019; Masson et al., 2020). While CIII md neuron connectivity to select second
order neurons has been published (Masson et al., 2020), the functional and behavioral roles of
these circuit components in the context of cold nociception remain unexplored. We hypothesized
that the noxious cold sensitive somatosensory nociceptive circuit functions through shared
circuitry amongst other somatosensory modalities.

Analyses of cold sensitive circuitry downstream of CIII md neurons elucidated dual
pathways for signal transduction: (1) CIII md neurons directly signal through premotor neurons
that modulate motor activity and (2) CIII md neurons also function via multisensory integrators
and projections neurons to convey somatosensory information to higher order brain regions.
Somatosensory neurons (Ch and CIII md) are required for cold-evoked responses but not CIV
md neurons (Turner et al., 2016; Turner et al., 2018). Based on EM connectivity, we performed
an unbiased behavioral screen of neuronal types that are synaptically connected to CIII md
neurons. Multisensory integration neurons, Basins 1-4, are required for cold nociception,
facilitate CIII md-evoked larval contraction (CT) responses, and select Basins exhibit cold-
evoked increases in calcium. Premotor neurons, DnB and mCSI, are required and facilitate CIII
md neuron mediated responses, whereas Chair-1 premotor neurons are required for cold
nociception but do not facilitate CIII md neuron mediated responses. Additionally, nociceptive
projection neurons A09e and TePns also integrate cold stimuli but only A09e neurons enhance
CIII md neuron mediated behavioral responses. Lastly, cold-evoked muscle activation is
strongest in abdominal segments (2-4). Individual muscle groups show distinct stimulus-evoked
responses, where dorsal longitudinal/oblique (DL & DO) muscles have sustained activation,
lateral transverse (LT) muscles exhibit increasing response during stimulus exposures, and
ventral longitudinal/oblique (VL & VO) muscles exhibit declining response. Collectively, we
functionally validate cold sensitive circuitry in *Drosophila* larvae from thermosensation, central
processing, to identifying muscles responsible for stimulus-evoked behavioral response.
134
135   **Results**
136
Peripheral sensory neurons share synaptic partners
Recent work on competitive interactions and behavioral transitions in *Drosophila* larvae
has reported CIII md and Ch connectomes in the context of larval mechanosensation (Masson et
al., 2020). We performed a comparative circuit analysis of somatosensory (Ch, CIII md and CIV
md) neurons using an online repository hosted by LMB Cambridge, which contains a ssTEM
volume of *Drosophila melanogaster* first instar larval central nervous system and a repository of
neural reconstructions (https://neurophyla.mrc-lmb.cam.ac.uk/). Here, we report and
contextualize relevant CIII md neuron downstream neurons, whose functional roles in cold
nociception were evaluated. The meta-analyses of previously published literature on *Drosophila*
larval somatosensory connectomes revealed several common post-synaptic partners that are
shared amongst Ch, CIII md and CIV md neurons (**Figure 1A**). CIII md neurons are upstream of
multisensory integration neurons, pre-motor neurons and projection neurons. Class III md
neurons are synaptically connected to multisensory integration neurons (Basin-2, -3 & -4). The
downstream pathway from Basins includes connectivity to A00c neurons and a polysynaptic
pathway to the rolling command neuron (Goro) via A05q neurons. CIII md neurons are also
connected to various pre-motor neurons, which are ventral nerve cord localized neurons
providing synaptic input to motor neurons, including DnB, A02n, Chair-1, A01d3, A02f, A02h

and A27k, and projection neurons including A05q, A09e, TePn05, A08n, A09o, A02o, and
dILP7. The CIII md neuron connectome analyses reveal broad second order connectivity and
complex interconnectivity amongst second order neurons (**Figure 1A, Figure 1-figure
supplement 1**). Functional and behavioral roles of CIII md neuron second order interneurons in
cold nociception remain unexplored and below we address this knowledge gap.
159
160
161   **Functional analysis of somatosensory neurons in cold nociception**
162   Due to shared post-synaptic neural connectivity of Ch, CIII md and CIV md neurons
(**Figure 1A, Figure 1-figure supplement 1**), we first assessed functional roles of these sensory
neurons in cold nociception by investigating: (1) stimulus-evoked calcium responses in the
Drosophila larval ventral nerve cord; (2) cold-evoked calcium responses of these sensory
neurons; (3) necessity and sufficiency of these sensory neurons in cold-evoked behaviors; and
(4) examining whether co-activating multiple sensory neurons facilitates CIII-mediated
behavioral response in *Drosophila* larvae.
169
170   1. <u>CaMPARI analyses reveal distinct neuropil activation patterns by sensory stimuli</u>
171   *Drosophila* larvae have distinct behavioral responses to various sensory stimuli including
touch, heat or cold. Exposure to noxious cold temperatures (≤10°C) evokes highly stereotyped
head and tail withdrawal towards the center of the animal, termed here as contraction (CT)
response (Patel et al., 2022; Turner et al., 2016). The cold-evoked CT response is defined as at
least 10% reduction in larval surface area (**Figure 1-figure supplement 2**). Innocuous
mechanical stimuli evoke a suite of behaviors including pausing, turning, head withdrawal,
and/or reverse locomotion. Noxious heat exposure leads to a corkscrew body roll escape
response (Babcock et al., 2009; Babcock et al., 2011; Im et al., 2015; Ohyama et al., 2015;
Tracey et al., 2003). Previous work on stimulus-evoked changes in neural ensembles in larval
zebrafish (*Danio rerio*) used the genetically encoded calcium integrator CaMPARI to reveal
distinct CNS neural activation patterns in response noxious heat or cold exposure (Fosque et al.,
2015). CaMPARI affords great spatial resolution as stimulus-evoked neural responses are
captured from a freely moving animal.
184   With CaMPARI and pan-neural imaging of ventral nerve cord neurons, we visualized
central representations of somatosensory stimuli across sensory modalities including innocuous
mechanical, noxious cold or noxious heat, which are primarily detected via Ch, CIII md, & CIV
md neurons, respectively. We pan-neuronally expressed CaMPARI using $R57C10^{GAL4}$ (Jenett et
al., 2012; Pfeiffer et al., 2008) and *Drosophila* larvae were simultaneously exposed to diverse
sensory stimuli and photoconverting light. Post-hoc imaging of intact *Drosophila* larvae ventral
nerve cord revealed relatively little neural activity when larvae are not presented with any
sensory stimulus (**Figure 1-figure supplement 3**). However, upon innocuous mechanical (gentle
touch) stimulation, there is a marked increase in neural activation as reported by $F_{red/green\ LUT}$
(**Figure 1-figure supplement 3**). *Drosophila* larvae exposed to noxious heat experience a large,

spatially broad increase in neural activity. Incidentally, similar neuropil regions appear to be
activated by both innocuous touch and noxious heat, albeit at different levels (**Figure 1-figure**
supplement 3). Lastly, noxious cold exposure leads to robust neural activation medially, in the
neuropil, however, cell bodies seem to have lower activation levels compared to touch or heat
stimulations (**Figure 1-figure supplement 3**). These experiments measuring pan-neuronal
ventral nerve cord activity in response to sensory stimuli would be ideal for comparative neural
activation analyses, were the spatial resolution of the optical microscope sufficient.
Acknowledging the limitations of the approach, we heretofore focus instead on single-cell type
specific genetic driver lines for identified neurons that can be assigned to known neurons in the
connectome,
204
2. <u>CIII md somatosensory neurons exhibit robust Ca^{2+} responses to cold</u>
Somatosensory neurons function as primary sensors of external stimuli. Ch and CIII md
neurons have previously been reported to exhibit cold-evoked Ca^{2+}increases (Turner et al., 2016;
Turner et al., 2018) whereas CIV md neurons are weakly sensitive to cold stimuli (Turner et al.,
2016). Furthermore, previous studies have addressed the potential confounds of cold-induced
muscle contraction on cold-induced electrical activity of CIII md neurons, where
electrophysiological recordings performed on de-muscled larval fillets revealed that CIII md
neural activity is not dependent upon muscles in response to cold. Exposure to noxious cold
stimuli results in temperature-dependent increases in CIII neuron electrical activity consisting of
both bursting and tonic firing (Himmel et al., 2021; Himmel et al., 2022; Himmel et al., 2023;
Maksymchuk et al., 2022; Maksymchuk et al., 2023; Patel et al., 2022). Cold sensitivity of
sensory neurons (Ch, CIII md, & CIV md) was assessed by selectively expressing the
CaMPARI2 Ca^{2+} integrator in sensory neurons using cell-type specific driver lines (**Figure 1-**
figure supplement 4A). CaMPARI2 signal, as assessed by F_{red}/F_{green} ratios at the cell body,
reveal all three neuron subtypes (Ch, CIII md and CIV md) have significantly higher response
upon cold exposure compared to no stimulus controls (**Figure 1B, Figure 1-figure supplement**
4B). Unsurprisingly, as high-threshold nociceptors, CIII md or CIV md neurons present
relatively low responses upon photoconverting light exposure *sans* cold (**Figure 1B, Figure 1-**
figure supplement 4B). However, mechanosensitive Ch neurons exhibit relatively high
responses upon exposure to only photoconverting light but no stimulus indicative of high
baseline neuronal activity (**Figure 1B, Figure 1-figure supplement 4B**).
Similar to the analysis at cell bodies, in the dendrites of Ch neurons we observe marked
increases in CaMPARI2 response to photoconverting light (control), however, upon cold
exposure (experimental condition) there is a further increase in CaMPARI2 response (**Figure**
1C). Sholl intensity analysis (see methods) likewise reveals that CIII md neurons present
relatively low CaMPARI2 response in control conditions, however, there is a robust cold-evoked
increase in CaMPARI2 response throughout the dendrites (**Figure 1C**). Interestingly, CIV md
neurons also exhibited significant increases in CaMPARI2 response upon cold exposure in the
cell body (**Figure 1B**). However, there is no change in distal dendritic CIV md neuron

CaMPARI2 response between cold and no stimulus conditions (**Figure 1C**). CIII md have robust
cold-evoked Ca^{2+} responses compared to Ch and CIV md neurons, which have relatively low
cold-evoked Ca^{2+} increases (**Figure 1B, C**). Collectively, these data validate previous findings
that identified CIII md neurons as the most sensitive to noxious cold temperatures both at the cell
body and dendrites, and thus we focused our study on the neural circuitry postsynaptic to CIII
md neurons.
240
3.   <u>CIII md somatosensory neurons are necessary and sufficient for the cold response</u>
We assessed the necessity of somatosensory neurons in noxious cold-evoked behavioral
responses (**Figure 1-figure supplement 2**) by expressing tetanus toxin light chain, which
inhibits neurotransmitter release (Sweeney et al., 1995). Based on Ca^{2+} imaging, we expected
both Ch and CIII md neurons may be necessary for cold-evoked CT responses. Neural silencing
of Ch or CIII md neurons led to significant reductions in cold-evoked CT responses compared to
either *w1118* (parental line) or *Empty*^{GAL4} controls (a GAL4 construct lacking the promotor
sequence; **Figure 1D**). Instantaneous behavioral response curves also indicate lower cold
sensitivity when either Ch or CIII md neurons are silenced. However, inhibiting neurotransmitter
release in CIV md neurons alone did not result in significant reductions in cold-evoked CT
responses (**Figure 1D**). To further assess the requirement of CIII md neuron activity for cold-
evoked behaviors, we simultaneously measured the delay from the onset of CIII md cold-evoked
Ca^{2+} increases to evoked mouth hook retraction. Upon neural silencing of CIII md neurons, there
is a significant delay in mouth hook retraction compared to controls suggesting that cold-evoked
head withdrawal requires sensory perception (**Figure 1-figure supplement 4C**). Among the
three somatosensory neuron types tested, neural silencing of CIII md neurons resulted in the
strongest impairment in cold-evoked behavioral response.
Next, we evaluated whether neural activation of sensory neurons via optogenetics would
be sufficient to elicit the CT behavioral response (**Figure 1-figure supplement 5A**). First, we
tested the efficacy of various optogenetic actuators in eliciting CIII-evoked CT responses using
either blue light actuators (ChR2, ChR2-H134R, and ChETA) or red-shifted actuator
(CsChrimson) and two independent CIII md *GAL4* drivers were used (*19-12*^{GAL4} and
GMR83B04^{GAL4}) (**Figure 1-figure supplement 6 A-F**) (Berndt et al., 2011; Boyden et al., 2005;
Gunaydin et al., 2010; Klapoetke et al., 2014). Collectively, using either blue or red-shifted
optogenetic actuators third instar *Drosophila* larvae exhibited light-evoked CT responses (**Figure
1-figure supplement 6**). We selected ChETA for the majority of the experiments due to its high
spike fidelity and lack of plateau potential (Gunaydin et al., 2010). We expressed ChETA in
individual sensory neuron subtypes and assessed evoked behavioral responses (Gunaydin et al.,
2010). When assessing evoked responses, we first analyzed CT responses as measured by
changes in surface area (**Figure 1-figure supplement 5B**). Only neural activation of CIII md
neurons led to CT responses in *Drosophila* larvae (**Figure 1E**) consistent with our previously
published work (Turner et al., 2016). Additionally, we analyzed larval mobility, which refers to
changes in larval postures as measured by changes in occupied space (**Figure 1-figure**

supplement 5C). Upon neural activation of CIII md neurons, there is a large increase in
Drosophila larval immobility compared to controls (**Figure 1-figure supplement 7A-C**).
However, there was no difference in immobility when either Ch or CIV md neurons were
optogenetically activated compared to control (**Figure 1-figure supplement 7A-C**). Of the
somatosensory neurons tested, only CIII md neurons are sufficient to elicit the CT response.
279
4. <u>Effect of co-activating CIII md somatosensory neurons plus additional somatosensory</u>
<u>neuron classes</u>
Both Ch and CIV md neurons share common first order post-synaptic partners with CIII
md neurons and all three somatosensory present cold-evoked increases in calcium levels.
However, neither Ch nor CIV md are sufficient to elicit a CT response. To further clarify the
roles of Ch and CIV md in cold-evoked behaviors, we simultaneously activated CIII md neurons
plus either Ch or CIV md neurons. We expected that co-activation of multiple sensory neuron
subtypes would facilitate optogenetically-evoked CT responses. Optogenetic activation of CIII
md neurons using two CIII driver lines (*19-12^{GAL4}* and *83B04^{GAL4}*) led to sustained increases in
instantaneous CT responses compared to control, where only one driver line (*19-12^{GAL4}*) was
used to activate CIII md neurons (**Figure 1F**). The activation of CIII md neurons concurrently
using two driver lines (expressed in the same cell type, CIII md) led to a significant increase in
immobility compared to single *GAL4* driver-mediated activation of these neurons (**Figure 1-**
figure supplement 7D-F), suggesting a single GAL4 line does not exhaust the dynamic range of
the CT response. Co-activation of CIII md and Ch neurons, which are cold sensitive and required
for cold-evoked CT responses, led to a subtle initial increase in instantaneous CT response
compared to CIII activation alone, however, the initial increase in evoked CT response was
quickly reduced to well below control (**Figure 1F**). Therefore, the co-activation of two
somatosensory neuron types, Ch and CIII md, elicited a CT response that varied along the
temporal axis relative to the activation of CIII md alone. *Drosophila* larval immobility was
reduced during co-activation of Ch and CIII md neurons compared to CIII md neuron activation
alone suggesting that Ch neurons do not facilitate CIII md neuron mediated CT responses
(**Figure 1-figure supplement 7D-F**). CIV md and CIII md neurons share a large proportion of
common second order interneuron connectivity including multisensory integration neurons,
premotor and ascending neurons. We predicted that CIV md and CIII md neuron co-activation
might potentiate CT responses. Interestingly, simultaneous activation of CIII and CIV md
neurons led to small but insignificant reductions in instantaneous and peak CT responses
compared to CIII md neuron activation alone (**Figure 1F**). Similarly, co-activation of CIII md
and CIV md neurons did not alter larval immobility compared to only CIII md neuron activation
(**Figure 1-figure supplement 7D-F**). Ch neurons are cold sensitive and required for cold
nociceptive behaviors but do not facilitate CIII md neuron-evoked behavioral responses (Figure
1B-F) (Turner et al., 2018). Meanwhile, CIV md neurons are modestly cold sensitive but are not
required for cold nociception and do not facilitate CIII md neuron-evoked CT responses.

Collectively, CIII md neurons have the strongest cold-evoked calcium response, are required for
cold-evoked behavioral response and are sufficient for the CT response.
315
316
317   5. CT response in *Drosophila* larvae throughout development
318
Synaptic connectivity was mapped using first instar larvae, however, thus far we assessed
cold- or CIII md-evoked larval behavioral responses in third instar larvae. Previous work
assessing how synaptic connectivity scales between first and third instar larvae revealed that
there is a five-fold increase in the number of synaptic connections and the size of neurons
thus conserving overall connectivity as measured by percent of input synapses during larval
development (Gerhard et al., 2017). Similarly, larval muscle cell size and sarcomere number
scale linearly with the larval body size, while maintaining average sarcomere length
(Balakrishnan et al., 2020; Demontis & Perrimon, 2009). Muscle contractile force analyses
showed that total larval contractile force generation is not dependent on the size or the
orientation of third instar larvae (Ormerod et al., 2022). We investigated how CT behavior
changes throughout larval development by assessing both cold- and CIII md-evoked
responses. First, cold-evoked larval CT magnitude and duration scale linearly across
development (**Figure 1G, H**). *Drosophila* larvae with silenced CIII md neurons exhibited
significantly reduced CT duration and magnitude compared to age matched controls (**Figure
1G, H**). Next, we evaluated how CIII md neuron-evoked CT behavior develops in
Drosophila larvae. Similar to cold-evoked CT responses, CIII md neuron-evoked CT
behavior also shows a linear trend in CT duration and magnitude (**Figure 1G, H**). Lastly, we
assessed the similarity between cold- and CIII md-evoked CT responses throughout larval
development. For second instar larvae, there was a 20% change increase in CT duration from
first instar larvae for both cold and CIII md activation (**Figure 1H**). For third instar larvae,
there was a greater increase in CT duration from first instar larvae for cold-evoked compared
to CIII-activation (**Figure 1H**). Meanwhile, developmental increases in CT magnitude
between cold- and CIII md neuron activation are nearly identical (**Figure 1H**). Collectively,
cold-evoked CT response scales linearly throughout development, where CIII md neurons are
necessary and sufficient for cold nociceptive responses.
344

Figure 1: CIII md neurons are primary cold sensors and share common post-synaptic
partners with mechanosensory and nociceptive Ch and CIV md neurons.
(**A**) Sensory neuron second order connectome analyses. Heatmap plot of synaptic connections
between sensory neuron subtypes including chordotonal (Ch), class III (CIII) md, class IV (CIV)
md neurons and previously published centrally located neurons. Sensory neurons (SN) Ch, CIII
md, and CIV md were analyzed. Multisensory integrator (MSI) neurons include Basin-1, -2, -3,
and -4. Neurons downstream of MSI (Post-MSI) include A00c, A05q and Goro. Premotor
neurons (PMNs) include Chair-1 (A10a), Down and Back (DnB), A02m and A02n. Lastly,
sensory neurons are also connected to several command/projection neurons including A02o,
A09e, A10j, A09o, TePn04 and TePn05. Synaptic connectivity data was extracted from
Neurophyla LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-C**) Calcium imaging
of sensory neurons including chordotonal (*IAV^{GAL4}*), class III md (*19-12^{GAL4}*) and class IV md
(*ppk^{GAL4}*) neurons using CaMPARI2. There were three conditions for sensory neurons: No
photoconversion (PC) control (no stimulus and no photoconversion), photoconversion control
(photoconversion and no stimulus) and stimulus (Stim) condition (photoconversion and 6°C
stimulus). CaMPARI2 data are reported area normalized intensity ratios for F_{red}/F_{green} (mean ±
SEM). Average N for each cell type and each condition is n=32. (**B**) CaMPARI2 response
measured at the cell body for each neuron type. (**C**) Sholl intensity analysis performed using
custom FIJI scripts. CaMPARI2 response is measured radially away from the center of the soma.
(**D**) Cold-evoked responses of third instar *Drosophila* larva. Sensory neurons Ch, CIII md or CIV
md neurons were silenced by inhibiting neurotransmitter release via cell type specific expression
of tetanus toxin (TNT). (**D, left**) Instantaneous %CT over time. The heatmap on top represents
change in temperature over time. (**D, right**) Cumulative %CT response for a duration of 5
seconds. Controls include *w1118* and *Empty^{GAL4}>TNT*. Significant stars: turquoise stars represent
comparison to *w1118* and purple stars represent comparison to *Empty^{GAL4}>TNT*. Average n=72.
(**E**) Neural activation of sensory neurons via cell type specific expression of ChETA relative to
Empty^{GAL4}>ChETA control. *Empty^{GAL4}* n=35. Ch n=20, CIV n=20. & CIII n=143. (**E, left**)
Instantaneous %CT over time. Blue bar represents optogenetic neural activation. (**E, right**) Peak
%CT. (**F**) Neural co-activation of sensory neurons and CIII md neurons. Here each condition
represents expression of *ChETA* in CIII md neurons and plus Ch (via *IAV-GAL4*), CIII (via
R83B04^{GAL4}) or CIV md neurons (via *ppk^{GAL4}*). (**F, left**) Instantaneous %CT over time. Blue bar
represents optogenetic neural activation. (**F, right**) Peak %CT response during optogenetic
stimulation. CIII n=143 and average experimental n=50 Significant differences indicated via
asterisks, where *p<0.05, ***p<0.001, and ****p<0.0001. (**G, left**) Cold-evoked CT behavioral
responses throughout larval development. In controls (*w^{1118}*), cold-evoked CT responses are
stronger as the larvae develop. CT magnitude is significantly greater between 1st v. 3rd instars
and 2nd v. 3rd instars. CT duration is significantly higher between 1st v. 3rd instars. Silencing CIII
md (*19-12^{GAL4}*) using TNT leads to significantly reduced cold-evoked responses as measured via
CT duration and magnitude across development when compared to age matched *w^{1118}* controls.
For each condition average n = 82 larvae. (**G, right**) CIII md neuron activation via optogenetics

reveals that CT responses increase as the animal develops. No ATR controls do not exhibit any
CT responses as measured by CT duration and magnitude. Activating CIII md (*19-12^{GAL4}*) using
ChR2-H134R leads to significantly increased optogenetic-evoked responses as measured via CT
duration and magnitude across development when compared to age matched no ATR controls.
Drosophila larvae with ATR show significant increase in CT magnitude between 1st v. 3rd instars
and 2nd v. 3rd instars. Additionally, CT duration is significantly higher between 1st v. 2nd instars
and 1st v. 3rd instars. For each condition average n = 27 larvae (**H**) Comparison between cold-
and CIII-evoked CT responses throughout development, where the CT duration and magnitude
are normalized to 1st instar larvae. (**H, left**) For 2nd instar larvae both cold- and CIII-evoked CT
duration increases by ~20% compared to 1st instar larvae. For 3rd instar larvae both cold- and
CIII-evoked CT duration increases by ~40% and ~30%, respectively, compared to 1st instar
larvae. (**H, right**) For 2nd instar larvae both cold- and CIII-evoked CT magnitude increases by
~13% and ~10%, respectively, compared to 1st instar larvae. For 3rd instar larvae both cold- and
CIII-evoked CT magnitude increases by ~32% compared to 1st instar larvae.

Figure 1-figure supplement 1: Sensory neuron connectivity matrix.
Synaptic connectivity matrix for neurons whose role in cold nociception was assessed. The
number in brackets indicates the total number of neurons analyzed for each cell-type. (**A**)
Absolute number of synaptic connections between pre- and post-synaptic neurons. (**B**)
Connectivity represented as proportion of synaptic input to the post-synaptic neurons.

Figure 1-figure supplement 2: *Drosophila* larval cold plate assay and quantitative analysis.

(**A**) A schematic of cold plate assay. Briefly, 3rd instar *Drosophila* larvae are plated on a thin

metal plate, then we expose the larvae to noxious temperature by transferring the plate onto pre-

chilled Peltier cold plate. (**B**) Behavior videos are automatically processed using custom macros

in Fiji, where behavioral videos are cropped, and background is removed to improve quantitative

analysis. (**C**) Larval surface is area is measured using Noldus Ethovision. The following larval

cold-evoked contraction (CT) behavioral metrics are calculated using r: Instantaneous CT%,

cumulative CT%, CT duration and CT magnitude. We define CT behavioral response as a

reduction in surface area less than -10%.

Figure 1-figure supplement 3: Stimulus-evoked calcium responses of *Drosophila* larval ventral nerve cord.

Representative images of *Drosophila* larval *in vivo* intact animal ventral nerve cord calcium responses assessed via pan-neural CaMPARI expression. Freely moving larvae were exposed to innocuous touch, noxious heat (45°C) or noxious cold (6°C) stimulus for 20 seconds and simultaneously exposed to 20 seconds of photo-converting light. Z-stacks of ventral nerve cord are shown with 2µm steps. Briefly, all animals were from the same batch, imaging was conducted using the same settings, and larvae were exposed to the same levels of photo-converting light. For better visualization, F_{red}/F_{green} CaMPARI fluorescence is reported, and images were pseudo-colored for enhancing signal-to-noise ratio, where highest intensity represented as white and lowest intensity with dark blue. Scale bar represents 50µm.

Figure 1-figure supplement 4: Somatosensory neural dendritic morphology and representative images of sensory neurons expressing CaMPARI2.
(**A**) Representative images of somatosensory neurons (Ch, CIII md and CIV md). Note, all images have separate magnification. Each image's scale bar represents 50µm. Drosophila larva and larval brain graphic was created with BioRender.com. (**B**) Confocal images of chordotonal (Ch: *IAV^{GAL4}*), class III md (CIII: *19-12^{GAL4}*) and class IV md (CIV: *ppk^{GAL4}*) neurons expressing CaMPARI2. There were three conditions for sensory neurons: No photoconversion control (no stimulus (Stim) and no photoconversion (PC)), photoconversion control (photoconversion and no stimulus) and stimulus condition (photoconversion and 6°C stimulus). Top row shows merge of F_{red} and F_{green}, second row is F_{green}, third row is F_{red} and last row contains F_{redLUT}, which is pseudo-colored with highest intensity being white and lowest intensity being dark blue (color

scale bar on bottom right). Cell body and dendrites are outlined in white for Ch and cell body are
outlined in white for CIII and CIV neurons. Scale bar represents 10μm. (**C**) CIII md
neurotransmitter release is required for cold-evoked mouth hook retraction in *Drosophila* larvae.
Using epifluorescence microscopy, CIII md neuron cold-evoked increases in Ca^{2+} levels (*19-*
12>GCaMP6m ± TNT) were assessed and the delay in mouth hook retraction was measured.
Compared to controls, there is a significant delay in mouth retraction, when CIII md neurons are
silenced with tetanus toxin expression. Average n=12 with three stimulations per animal.

Figure 1-figure supplement 5: *Drosophila* **larval neural activation assays using**
optogenetics. To perform neural activation experiments, we created a custom built optogenetic
experimental setup, which has a very high spatial resolution and signal to noise ratio by using
principles of dark field illumination. (**A**) Schematic of custom optogenetic rig, where stimulus
and video recording are controlled via computer using Noldus Ehtovision. Individual *Drosophila*
larva are plated on clear glass plate and illuminated from below with white light. Neuron

activating blue light is also delivered from the below. Individual behavioral videos are
automatically cropped and stabilized using custom Fiji macros. (**B-C**) We measured to two
variables *Drosophila* larval surface area and mobility. (**B**) Top, image stills from various
timepoints before and during optogenetic stimulation of cold sensitive CIII md neurons resulting
in contraction (CT) behavioral response. Bottom, percent change in area over time of an
individual animal. During baseline, larval locomotion and turns results brief changes in surface
area (±2-4%). Whereas upon neural stimulation, there is a distinct lasting reduction in surface
area of less than -10%. (**C**) Larval mobility refers to changes in larval postures as measured by
changes in occupied space. Since the amount of larval mobility is a function of various intrinsic
and extrinsic factors, we normalized larval mobility to baseline period, where there was no
neural activation. Top, conceptual framework of how larval mobility is measured. Original
videos are motion stabilized in the XY axis and thresholded. Next, we perform image
calculations (Mobility=$\text{Frame}_{\text{timepoint 2}}$ – $\text{Frame}_{\text{timepoint 1}}$) to get just the red shaded portions
denoting changes in postural locations (red arrowheads). *Drosophila* larval time series mobility
data are reported as percent change in mobility, time spent being immobile and percent of
immobile animals, or for a genotype percent of animals that are immobile over time, immobility
is defined as -25% or more reduction in mobility.

Figure 1-figure supplement 6: Optogenetic activation of CIII md neurons.
(**A**) Activation of CIII md neurons using different optogenetic actuators including *CsChrimson,*
ChETA, ChR2 and *ChR2-H134R*. Activation of CIII md neurons using either *19-12^{GAL4}* or
83B04^{GAL4} leads to a contraction response in third instar *Drosophila* larvae. Optogenetic
activation of CIII md neurons (*19-12^{GAL4}*) using *CsChrimson, ChETA, ChR2* or *ChR2-H134R*
leads to CT responses. Each optogenetic actuator has unique impacts neuronal action potential
firing rates leading to slightly different CT responses, however, CIII md neuron activation leads
to CT behavioral response. (**B-F**) Represents instantaneous CT percentage responses of no ATR
control and with ATR conditions for each of the actuators. (**B-E**) 470nm light was used to
activate CIII md neurons. (**F**) CsChrimson is very sensitive to minimal ambient white light
exposure. Here, the experiments were conducted in near darkness conditions and thus no pre-
stimulus data are recorded. The larval behavior was only recorded upon optogenetic light
(617nm) exposure. (**B**) *19-12^{GAL4}>ChETA* average n=27. (**C**) *19-12^{GAL4}> ChR2-H134R* average
n=37. (**D**) *19-12^{GAL4}>ChR2* average n=29. (**E**) *83B04^{GAL4}>ChETA* average n=30. (**F**) *19-
12^{GAL4}>CsChrimson* average n=33.

Figure 1-figure supplement 7: ***Drosophila* larval mobility pipeline and effects of neural**
activation and co-activation of sensory neurons on larval mobility (**A-F**) *Drosophila* larval
mobility following somatosensory neuron optogenetic activation (**A-C**) and CIII md neuron plus
co-activation of Ch, additional CIII *GAL4* driver ($83B04^{GAL4}$) or CIV md neurons (**D-F**). (**A, D**)
Instantaneous percent immobility. Blue bar represents optogenetic neural activation. (**B, E**)
Average percent change in mobility for each genotype, where greater percent immobility results
in larger changes in average mobility. (**C, F**) Immobility duration in seconds during stimulation.
Neural activation: *Empty^{GAL4'}* n=35. Ch n=20, CIV n=20. & CIII n=143. Neural co-activation:
CIII n=143 and average experimental n=50.Significant differences indicated via asterisks, where
****p<0.0001.

Multisensory integrators are required and facilitate larval contraction response

Basin interneurons function as multisensory integrators receiving convergent inputs from mechano-, chemo- and thermo-sensitive peripheral sensory neurons (**Figure 2A, Figure 2-figure supplement 1A**). Previous behavioral and functional studies have revealed that Basin interneurons are required for nociceptive escape responses mediated by Ch and CIV md neurons (Ohyama et al., 2015). Both Ch and CIV md neurons play roles in CIII md mediated behavioral responses either in cold nociception and/or can detect cold stimuli. We hypothesized that Basin interneuron function is required downstream of cold nociceptive somatosensory neurons. We evaluated whether Basin interneurons are necessary and sufficient for cold nociception, exhibit cold-evoked increases in calcium response and function downstream of CIII md neurons.

Silencing Basin interneurons reduces cold-evoked CT

Basin interneurons receive somatosensory cues from CIII md neurons, thus we predicted that inhibiting neurotransmitter release from Basin neurons will result in impaired cold-evoked behaviors (**Figure 2A**). We assessed the requirement of all Basin neurons using "pan-" Basin driver lines and also assessed roles of individual Basin neurons using subtype specific driver lines. Neural silencing of all Basin (1-4) neurons, using $R72F11^{GAL4}$, led to significant reductions in cold-evoked CT responses compared to genetic background (w^{1118}) and $Empty^{GAL4'}$ controls (**Figure 2B**). *Drosophila* larvae with impaired Basin (1-4) neuronal function had significant reductions in CT duration, magnitude and cumulative %CT response compared to controls (**Figure 2C-E**). CIII neurons have differential connectivity to individual Basin subtypes (**Figure 2A**). Therefore, we assessed requirement of Basin-1, -2 or -4 neurons, for which there are previously validated independent driver lines (Jovanic et al., 2016; Ohyama et al., 2015). Tetanus toxin mediated neural silencing of Basin-2 or -4 led to significant reductions in cold-evoked CT response, as measured by cumulative %CT response, CT duration and CT magnitude, compared to controls (**Figure 2B-E**). CIII md neurons do not synapse onto Basin-1 neurons according to the EM-mapped connectome (**Figure 2A**) (Masson et al., 2020), however, Basin-1 & -2 neurons share synaptic connectivity to both feedback and feedforward GABAergic interneurons, both of which are downstream of sensory neurons (Jovanic et al., 2016). Furthermore, Basin-1 neural activation leads to depolarizations in Basin-2 through GABAergic disinhibitory pathway (Jovanic et al., 2016). Therefore, we expected that inhibiting neurotransmitter release in Basin-1 neurons would result in reduced cold-evoked responses. Neural silencing of Basin-1 neurons resulted in modest, but significant reductions in cold-evoked responses (**Figure 2B-E**). Impaired Basin neuron signaling results in at least 25% reduction in cold-evoked responses with the strongest reductions for all-Basin, Basin-2 or Basin-4 driver lines (**Figure 2J**).

Co-activation of Basin interneurons and CIII md somatosensory neurons enhances CT

Next, we evaluated whether neural activation of Basin neurons would impair or elicit a CT response. Optogenetic activation of Basin neurons, either using all Basin or individual Basin

driver lines, did not elicit a CT response (**Figure 2-figure supplement 2**). But the simultaneous

co-activation of CIII md and Basin neurons led to sustained increases in CT responses compared

to controls, where only CIII md neurons were activated, across multiple behavioral metrics

(**Figure 2F-I**). Coactivation did not result in a change in larval immobility (**Figure 2-figure**

supplement 1B-D), whereas activation of all Basins led to significantly greater immobility

(**Figure 2-figure supplement 1B-D**). These results indicate that Basin neurons are not sufficient

for the CT response, but that the combined activation of CIII md and Basins not only suffices but

also elicits an even stronger CT response than activating CIII md neurons alone.

To parse the contribution of individual Basin neuronal subtypes, we next assessed the

role of Basin-1, -2 or -4, and all four Basin together, in either facilitating or suppressing CT

responses using our co-activation paradigm (**Figure 2F-I**). Co-activation of Basin-1 or Basin -2

with CIII md neurons led to an enhanced CT response compared to controls across all measures

of behavioral response including instantaneous %CT response, peak %CT response, CT duration

and CT magnitude (**Figure 2F-I**), with a subsequent significant increase in larval immobility

(**Figure 2-figure supplement 1B-D**). Basin-2 plus CIII md neuron co-activation led to strong

facilitation of CT responses, but surprisingly resulted in significantly reduced immobility

(**Figure 2-figure supplement 1B-D**). Unlike for all other Basin neurons tested, Basin-4 and CIII

md neuron co-activation led to a suppression of CT response, where peak instantaneous %CT

response was similar to controls, however, there was a rapid reduction in instantaneous %CT

response compared to controls (**Figure 2F-I**). Both CT duration and CT magnitude were

significantly impaired for Basin-4 plus CIII md neuron co-activation compared to controls

(**Figure 2H, I**), which also showed significantly lower immobility (**Figure 2-figure supplement**

1B-D). Interestingly, dual activation of CIII md and all Basin neurons led to weaker CT

enhancement compared to co-activation of either Basin-1 or -2 coupled with CIII md neurons

(**Figure 2J**), consistent with the finding that co-activation with Basin-4 reduced the CT response.

Collectively, second-order Basin neurons are required for cold-evoked responses and specifically

Basin-1 and Basin-2 are able to enhance CIII md neuron-evoked behavioral responses.

<u>CaMPARI reveals Basin-2- and Basin-4 are activated in CT responses</u>

To further explore how Basin interneurons function in cold nociception, we sought to

investigate cold-evoked Ca^{2+} responses of Basin neurons. Since somatosensory neurons are

cholinergic (Salvaterra & Kitamoto, 2001), we expected that Basin neurons postsynaptic to CIII

md neurons will exhibit cold-evoked increases in Ca^{2+}. Post-hoc imaging of evoked CaMPARI2

fluorescence revealed that Basin neurons have significantly higher F_{red}/F_{green} ratios compared to

control, as assessed by all-Basin driver line (**Figure 3A**). We further investigated Ca^{2+} responses

in greater detail using individual driver lines for Basin-1, -2 or -4. Basin-1 neurons are weakly

required for cold-evoked CT responses (**Figure 2J**) and coherently do not exhibit cold-evoked

increases in Ca^{2+} response (**Figure 3B**). In contrast, Basin-2 and -4 neurons both exhibit

significant increases in Ca^{2+} responses compared to their respective controls (**Figure 3C-D**).

Collectively, Basin-2 and -4 neuron subtypes that are required for cold-evoked CT responses
(**Figure 2J**), also exhibit cold-evoked increases in Ca^{2+}.
619

<u>CIII md neurons and Basin-2 and -4 neurons are functionally connected</u>
Basin-2 and -4 are postsynaptic to CIII md neurons, are required for cold nociception and
have cold-evoked increases in Ca^{2+}. Next, we assessed whether CIII md neurons and Basin-2 or -
4 neurons are functionally connected. From the EM-reconstructed connectome we predicted that
the activation of CIII md neuron will result in increased Ca^{2+} levels of Basin-2 or -4. As
expected, optogenetic activation of CIII md neurons led to significant increases in Ca^{2+} levels in
Basin-2 neurons, and repeated stimulation of CIII md neurons did not lead to sensitization of
Basin-2 Ca^{2+} responses (**Figure 3E, F**). Previously, it was shown that Ch and CIII md neurons
could elicit Ca^{2+} response in Basin-4 neurons (Kaneko et al., 2017), however, in this previous
study the authors were unable to determine which of the two sensory neuron cell types led to
increases in the Basin-4 Ca^{2+} response (Kaneko et al., 2017). Here, we show that upon
specifically activating CIII md neurons, Basin-4 neurons have large, rapid increases in cytosolic
Ca^{2+} followed by quick return to baseline levels. In contrast to Basin-2, Basin-4 neurons showed
reduced Ca^{2+} responses upon repetitive activations of CIII md neurons (**Figure 3G, H**).
Therefore, CIII md neuron activation is differentially encoded by Basin-2 and -4 neurons, where
upon initial stimulation Basin-4 neurons have a much larger CIII md neuron-evoked increase in
Ca^{2+} levels but exhibit habituation compared to Basin-2 neurons, which show consistent Ca^{2+}
upon repetitive stimulations. Collectively, our data demonstrate that Basin neurons are required
for cold-evoked behaviors and Basin-2 and -4 neurons functionally operate downstream of CIII
md neurons (**Figure 3-figure supplement 1**).

Figure 2: Multisensory integrator second order neurons are required for cold-evoked behaviors and facilitate CIII-evoked behaviors.

(**A**) Basins (1-4) receive inputs from sensory neurons (Ch, CIII md and CIV md), Basins, premotor neuron Down and Back (DnB) and command-like/projection neuron A02o and TePn05. Heatmap plot of pre-synaptic connections to Basins. Synaptic connectivity data was extracted from Neurophyla LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-E**) Cold-evoked responses of third instar *Drosophila*. Basin (1-4) neurons were silenced by inhibiting neurotransmitter release via cell type specific expression of tetanus toxin (TNT), where All Basin (*R72F11^{GAL4}*), Basin-1 (*R20B01^{GAL4}*), Basin-2 (*SS00739^{splitGAL4}*) and Basin-4 (*SS00740^{splitGAL4}*). (**B**) Instantaneous %CT over time. The heatmap on top represents change in temperature over time. (**C**) Cumulative %CT response for a duration of 5 seconds. (**D**) CT duration in seconds. (**E**) CT magnitude as average percent change in area for the duration of stimulus. Controls include w^{1118}, *Empty^{GAL4}'* (*w;;attP2*) or *Empty^{GAL4}''>TNT* (*w;attP40;attP2*). Significant differences were compared to each *GAL4's* respective controls dependent on insertion sites. Significant stars: turquoise stars represent comparison to *w1118*, purple stars represent comparison to *Empty^{GAL4}'>TNT* and sea green stars represent comparison to *Empty^{GAL4}''>TNT*. Average n = 68. (**F-I**) Neural co-activation of Basin neurons and CIII md neurons. Here each condition represents expression of ChETA in CIII md neurons and plus Basin (1-4) neurons. (**F**) Instantaneous %CT over time. Blue bar represents optogenetic neural activation. (**G**) Peak %CT response during optogenetic stimulation. (**H**) CT duration in seconds during optogenetic stimulation. (**I**) CT magnitude as average percent change in area for the duration of stimulus. Significant stars: purple stars represent comparison to CIII md + *Empty^{GAL4}'>ChETA*. Empty^{GAL4}' n=143 and experimental condition average n =49. (**J**) Overall percent change from control for either neural silencing or neural co-activation. The metrics for neural silencing include cumulative %CT, CT magnitude, and CT duration. The following metrics were used to calculate percent for neural co-activation: cumulative %CT, peak %CT, CT duration and magnitude. Significant differences indicated via asterisks, where *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

 Figure 2-figure supplement 1: Neural reconstructions and larval mobility for multisensory
integrator neurons.
(**A**) Individual Basin neuron subtypes (magenta) and CIII md neuron axons (blue). Neural
reconstruction data was extracted from Neurophyla LMB Cambridge. (https://neurophyla.mrc-
lmb.cam.ac.uk/). (**B-D**) *Drosophila* larval mobility for Basin plus CIII md neuron optogenetic
co-activation. (**B**) Instantaneous percent immobility. Blue bar represents optogenetic neural
activation. (**C**) Average percent change in mobility for each genotype, where greater percent
immobility results in larger changes in average mobility. (**D**) Immobility duration in seconds
during stimulation. *Empty$^{GAL4'}$* n=143 and experimental condition n =49. Comparisons to *CIII md*
+ Empty$^{GAL4'}$>ChETA. Significant differences indicated via asterisks, where **p<0.01, and
****p<0.0001.

699
700

701
Figure 2-figure supplement 2: Optogenetic activation of individual neuronal cell-types.
Heatmap represents optogenetically-evoked instantaneous contraction (CT) proportions of
Drosophila larvae. Individual neuronal cell-types were optogenetically activated using cell-type
specific expression of *ChETA*. Black arrow indicates optogenetic activation of primary cold
somatosensory CIII md neurons, which are the only the genotype with high proportions CT.
Average n= 31.

Figure 3: Cold- and CIII- evoked calcium responses of Basin neurons.

(**A-D**) Ca^{2+} responses of Basin neurons upon cold exposure vs. controls (room temperature). Neural responses (CaMPARI2) of Basin neuron cell bodies were analyzed using the following cell type driver lines (**A**) All Basin ($R72F11^{GAL4}$) average n=197, (**B**) Basin-1 ($R20B01^{GAL4}$) average n=27, (**C**) Basin-2 ($SS00739^{splitGAL4}$) average n=119 and (**D**) Basin-4 ($SS00740^{splitGAL4}$) average n=46. CaMPARI2 fluorescence ratio is reported as F_{red}/F_{green}. We report the data as individual datapoints, where the red line represents mean, and hybrid plots (boxplot & violin) for visualizing the distribution and quartiles of data. Significant stars represent p<0.05, where comparisons were made to their respective no stimulus controls. (**E-H**) To functionally assess CIII md neuron to Basin-2 or Basin-4 connectivity, we optogenetically activated CIII md neurons ($83B04^{lexA}>CsChrimson$) and visualized changes in evoked Ca^{2+} using $Basin-2^{splitGAL4}$ or $Basin-4^{splitGAL4}>GCaMP6m$. Control: No all *trans*-retinal (ATR) supplemented diet, which is required for optogenetic stimulation in *Drosophila*. Orange bars indicate optogenetic stimulation. (**E,G**) Basin-2 and Basin-4 changes in GCaMP reported as $\Delta F/F_{prestimulus}$, where prestimulus refers to 15 seconds prior to optogenetic stimulation. (**F,H**) Maximum Basin-2 and Basin-4

735 neuronal responses (ΔF/F$_{prestimulus}$) upon optogenetic stimulation. Average n for each genotype
736 was 13. Comparisons made to relevant controls and significant differences indicated via
737 asterisks, where *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

738
Figure 3-figure supplement 1: Summary of behavioral and functional roles of multisensory
integrators in cold nociception.
Blue arrows indicate strength of synaptic connectivity between the CIII md neurons and second
order neurons. In the behavioral analysis column, the arrow direction indicates reduction or
enhancement of CT response and shading indicates magnitude of change from control. In neural
activity column: NC denotes no significant change in Ca^{2+} response, the arrow direction
indicates reduction or enhancement in evoked Ca^{2+} response and shading indicates magnitude of
change from control. Empty spaces indicate experimental analyses were not performed.
747
748 **Multisensory integrators function independently of Goro pathway for cold nociception**
749
Basin neurons innervate a set of projection neurons (A05q and A00c), which are
upstream of a command neuron (Goro) that is responsible for initiating CIV md neuron-mediated
nociceptive escape behaviors (**Figure 4A, Figure 4-figure supplement 1A**) (Ohyama et al.,
2015). We set out to test whether cold-evoked behavioral responses mediated by CIII md and
Basin neurons function through A00c, A05q, and/or Goro neurons. We first assessed whether
these neurons are required for cold-evoked behavioral responses (**Figure 4B-E**). Neural silencing
of A05q neurons via tetanus toxin led to mild, yet significant reductions in cold-evoked
responses, when compared to w^{1118} genetic control (**Figure 4B-E**). Neural silencing of A00c
neurons resulted in significantly lower cold-evoked cumulative CT response compared to w^{1118}
(**Figure 4C**). There were also significant reductions in CT duration and CT magnitude when
A00c neurons were silenced compared to both controls (**Figure 4D, E**). In contrast, the Goro
command neuron for nociceptive rolling behavior is not required for cold-evoked CT responses
(**Figure 4B-E**). Next, we assessed whether neural activation of these neurons led to evoked CT

responses. Like Basin neurons, single cell-type activation of A00c, A05q, or Goro neurons did
not lead to CT behavior (**Figure 2-figure supplement 2**). Similarly, optogenetic co-activation of
CIII md neurons and A05q or Goro neurons did not lead to significant facilitation of the CT
response (**Figure 4F-I**). However, simultaneously co-activating A00c and CIII md neurons led to
significant increases in CT duration and CT magnitude (**Figure 4H, I**). There were no changes in
Drosophila larval immobility upon co-activation of CIII md plus A00c, A05q or Goro neurons
(**Figure 4 - figure supplement 1B-D**). Only co-activation of A00c neurons leads to notable
enhancement of CIII md mediated CT response, whereas neural silencing of A00c or A05q
neurons led to greater than 25% impairment in cold-evoked behavioral responses (**Figure 4J**). In
conclusion, the Basin to Goro polysynaptic pathway does not significantly contribute to the cold-
evoked CT response.

Figure 4: Projection neurons downstream of Basins and sensory neurons function in cold-
evoked responses.
(**A**) A00c and A05q primarily receive inputs from Basins and premotor neuron Down and Back
(DnB). Goro neurons primarily receive inputs from A05q neurons. Heatmap plot of pre-synaptic
connections to downstream neurons. Synaptic connectivity data was extracted from Neurophyla
LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-E**) Cold-evoked responses of
third instar *Drosophila*. We used cell-type specific driver lines for downstream neurons to drive
expression of tetanus toxin (TNT): A00c (*R71A10^{GAL4}*), A05q (*R47D0 ^{GAL4}*) and Goro
(*R69F06^{GAL4}*). (**B**) Instantaneous %CT over time. The heatmap on top represents change in
temperature over time. (**C**) Cumulative %CT response for a duration of 5 seconds. (**D**) CT
duration in seconds. (**E**) CT magnitude as average percent change in area for the duration of
stimulus. Controls include *w^{1118}* and *Empty^{GAL4}' (w;;attP2)*. For each genotype average n=64.
Significant stars: turquoise stars represent comparison to *w^{1118}* and purple stars represent
comparison to *Empty^{GAL4}'>TNT*. (**F-I**) Neural co-activation of downstream neurons and CIII md
neurons. Here each condition represents expression of ChETA in CIII md neurons plus A00c,
A05q or Goro neurons. (**F**) Instantaneous %CT over time. Blue bar represents optogenetic neural
activation. (**G**) Peak %CT response during optogenetic stimulation. (**H**) CT duration in seconds
during optogenetic stimulation. (**I**) CT magnitude as average percent change in area for the
duration of stimulus. Significant purple stars represent comparison to *CIII md +*
Empty^{GAL4}'>ChETA. Empty^{GAL4} n=143 and experimental condition average n =33. (**J**) Overall
percent change from control for either neural silencing or neural co-activation. The metrics for
neural silencing include cumulative %CT, CT magnitude, and CT duration. The following
metrics were used to calculate percent for neural co-activation: cumulative %CT, peak %CT, CT
duration and magnitude. Significant differences indicated via asterisks, where *p<0.05,
**p<0.01, and ***p<0.001.

Figure 4-figure supplement 1: Neural reconstructions and larval mobility for A00c, A05q, and Goro neurons.

(**A**) Individual interneuron subtypes (magenta) and CIII md neuron axons (blue). Neural reconstruction data was extracted from Neurophyla LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-D**) *Drosophila* larval mobility for A00c, A05q or Goro neurons plus CIII md neuron optogenetic coactivation. (**B**) Instantaneous percent immobility. Blue bar represents optogenetic neural activation. (**C**) Average percent change in mobility for each genotype, where greater percent immobility results in larger changes in average mobility. (**D**) Immobility duration in seconds during stimulation. Significant stars represent $p<0.05$, where stars represent comparison to CIII md + *Empty*$^{GAL4'}$>*ChETA*. Empty^{GAL4} n=143 and experimental condition average n =33.

823
824 **Premotor neurons function downstream of CIII md neurons to mediate cold nociceptive**
825 **responses**
826
Select *Drosophila* larval premotor neurons were previously implicated in CIV-mediated
nociceptive escape responses. Specifically, DnB premotor neurons are involved in noxious
thermal stimulus-evoked c-bending and rolling behavior (Burgos et al., 2018; Lopez-Bellido et
al., 2019). *Drosophila* larvae also roll in response to activation of mCSI premotor neurons, which
are synaptically connected to CIV md neurons and predicted to be A02m/n neurons from EM
connectomes (Lopez-Bellido et al., 2019; Yoshino et al., 2017). Additionally, Chair-1 (A10a)
premotor neurons have been implicated in anemotaxis (Jovanic et al., 2019). Collectively, these
nocifensive premotor neurons primarily receive inputs from both primary sensory neurons (CIII
& CIV md) and multisensory integrators (Basin-2 & -4) (**Figure 5A, Figure 5-figure**
supplement 1A). CIII md neurons have additional synaptic connectivity with intersegmental
feedback circuitry involved in locomotion, where Ifb-fwd (A01d3) is active during forward
locomotion and Ifb-bwd (A27k) is active during backwards locomotion (Kohsaka et al., 2019).
The intersegmental feedback circuit neurons (A01d3 and A27k) are interconnected via A02e and
A02g (Kohsaka et al., 2019). GABAergic A31k premotor neuron is downstream of A01d3 and is
responsible for inhibiting motor activity causing relaxation of the larval body. Lastly, A02f has
been implicated in delaying nocifensive behavioral responses (Garner, 2020). We predicted that
select premotor neurons are required for cold-evoked responses and function downstream of CIII
md neurons in a stimulus-specific manner.
Inhibition of neural transmission via cell type specific expression of tetanus toxin in
individual premotor neurons led to reduced cold-evoked responses in *Drosophila* larvae (**Figure**
5B-E). Silencing Chair-1 neurons resulted in the strongest reduction of cold-evoked CT
responses, where instantaneous %CT was the lowest of all premotor neuron subtypes tested
(**Figure 5B**). CT duration, magnitude and cumulative percent response were all significantly
reduced compared to controls (**Figure 5B-E**). Impairment in Chair-1 neuronal signaling leads to
75% reduction from controls in cold-evoked behaviors (**Figure 5J**). We silenced DnB neurons
using two independent cell-type specific driver lines (DnB' (*IT4051^{GAL4}*) & DnB''(*IT412^{GAL4}*)),
where both resulted in reduced instantaneous %CT response, along with significant reductions in
CT duration and magnitude compared to controls (**Figure 5B-E**). Silencing mCSI (*R94B10^{GAL4}*)
neurons also resulted in significantly reduced cold-evoked CT responses compared to controls
(**Figure 5B-E**). DnB or mCSI inhibition of neurotransmitter release leads to approximately 50%
reduction in cold-evoked CT responses (**Figure 5J**). Furthermore, we silenced premotor neurons
previously implicated in locomotion (A01d3, A27k, A02e, A02g, A31k) or delaying nocifensive
responses (A02f), this did not lead to significant changes in cold-evoked CT responses (**Figure**
5-figure supplement 2). Collectively, the previously implicated nocifensive premotor neurons
are required for cold-evoked behavioral responses.

Based on EM connectivity and neural silencing experiments, we predicted that activation
of these premotor neurons would be sufficient for *Drosophila* larval CT response. First, we
found that activation of premotor neurons alone did not evoke CT response (**Figure 2-figure**
supplement 2). Next, we performed optogenetic co-activation of CIII md and premotor neuron
subtypes. Chair-1 neuron co-activation with CIII md neuron did not have an effect on CT
responses (**Figure 5F-I**), however, there was a reduction in larval immobility, where average
mobility was increased and duration of immobility was reduced compared to when only CIII md
neurons are activated (**Figure 5-figure supplement 1B-D**). Simultaneous activation of CIII md
neurons and DnB or mCSI both led to significant increases in CT responses, as measured by
peak CT response, CT duration or magnitude, compared to CIII md neuron activation alone
(**Figure 5F-I**). DnB and CIII md neuron co-activation led to short lived, subtle but insignificant
increases in larval immobility (**Figure 5-figure supplement 1B-D**). Co-activation of mCSI and
CIII md neurons resulted in mild reductions in larval immobility (**Figure 5-figure supplement**
1B-D). These data reveal DnB and mCSI neuronal activity enhances CIII md neuron mediated
CT responses.
Since these premotor neurons are postsynaptic to CIII md neurons and their activity is
required for proper cold-evoked behaviors, we predicted that either activation of CIII md neurons
or exposure to cold would cause increases in Ca^{2+}. To test this prediction, we selectively
expressed the Ca^{2+} integrator CaMPARI2 in premotor neurons. Unexpectedly, there was no
change in cold-evoked Ca^{2+} levels in mCSI or Chair-1 neurons (**Figure 6A-B**). Noxious cold
exposure did however lead to significant Ca^{2+} increases in DnB neurons (**Figure 6C-D**). To
further assess how CIII md neuronal activity affects DnB function, we optogenetically activated
CIII md neurons and assessed evoked Ca^{2+} levels of DnB using GCaMP6. Interestingly,
Drosophila larvae raised without all *trans*-retinal, a requisite light sensitive cofactor for
optogenetic experiments, also had a mild light-evoked increase in Ca^{2+} (**Figure 6E, F**). CIII md
neuron optogenetic activation led to significant increases in DnB Ca^{2+} levels that slowly returned
to baseline levels (**Figure 6E, F**). Upon repeated CIII md neuron activations, DnB neurons
exhibit a blunted Ca^{2+} response relative to initial stimulation (**Figure 6E, F**). Taken together we
find that premotor neuronal function is required for cold nociception and premotor neuron
activity can facilitate CIII md neuron mediated CT responses (**Figure 6-figure supplement 1**).

903
904
Figure 5: Premotor neurons downstream of sensory neurons and Basin neurons are
required for cold-evoked responses.
(**A**) Chair-1 (A10a), A02m/n (predicted to be mCSI neurons) and Down and Back (DnB,
A09l) primarily receive inputs from Basins, CIII md and CIV md neurons. Heatmap plot of pre-
synaptic connections to premotor neurons. Synaptic connectivity data was extracted from
Neurophyla LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-E**) Cold-evoked
responses of third instar *Drosophila*. Premotor neurons were silenced by inhibiting
neurotransmitter release via cell type specific expression of tetanus toxin (TNT), where Chair-1
(*SS00911^{splitGAL4}*), DnB' (*IT4015^{GAL4}*), DnB'' (*IT412^{GAL4}*) and mCSI (*R94B10^{GAL4}*). (**B**)
Instantaneous %CT over time. The heatmap on top represents change in temperature over time.
(**C**) Cumulative %CT response for a duration of 5 seconds. (**D**) CT duration in seconds. (**E**) CT
magnitude as average percent change in area for the duration of stimulus Controls include *w^{1118}*,
Empty^{GAL4}' (*w;;attP2*) or *Empty$^{GAL4''}$>TNT* (*w;attP40;attP2*). Significant differences were
compared to each *GAL4's* respective controls dependent on insertion sites. Significant stars:
turquoise stars represent comparison to *w^{1118}*, purple stars represent comparison to
Empty^{GAL4}'>TNT and sea green stars represent comparison to *Empty$^{GAL4''}$>TNT*. For each
genotype average n=71. (**F-I**) Neural co-activation of premotor neurons and CIII md neurons.
Here each condition represents expression of ChETA in CIII md neurons plus premotor neurons.
(**F**) Instantaneous %CT over time. Blue bar represents optogenetic neural activation. (**G**) Peak
%CT response during optogenetic stimulation. (**H**) CT duration in seconds during optogenetic
stimulation. (**I**) CT magnitude as average percent change in area for the duration of stimulus.
Empty^{GAL4}' n=143 and experimental condition average n =35. Significant stars represent p<0.05,
where purple stars represent comparison to *CIII md + Empty^{GAL4}'>ChETA*. (**J**) Overall percent
change from control for either neural silencing or neural co-activation. The metrics for neural
silencing include cumulative %CT, CT magnitude, and CT duration. The following metrics were
used to calculate percent for neural co-activation: cumulative %CT, peak %CT, CT duration and
magnitude. Significant differences indicated via asterisks, where *p<0.05, **p<0.01,
***p<0.001, and ****p<0.0001.

Figure 5-figure supplement 1: Neural reconstructions and larval mobility for premotor
neurons.
(**A**) Individual premotor neuron subtypes (magenta/green) and CIII md neuronal axons (blue).
Neural reconstruction data was extracted from Neurophyla LMB Cambridge.
(https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-D**) *Drosophila* larval mobility observed with
premotor neurons plus CIII md neuron optogenetic coactivation. (**B**) Instantaneous percent
immobility. Blue bar represents optogenetic neural activation. (**C**) Average percent change in
mobility for each genotype, where greater percent immobility results in larger changes in average
mobility. (**D**) Immobility duration in seconds during stimulation. *Empty$^{GAL4'}$* n=143 and

experimental condition average n =35. Comparison to *CIII md + Empty$^{GAL4'}$>ChETA*.
Significant differences indicated via asterisks, where *$p<0.05$, and ****$p<0.0001$.

Figure 5-figure supplement 2: Premotor network neurons are not required for cold-evoked
responses. (A-D) Cold-evoked responses of third instar *Drosophila* larvae. Premotor neurons
were silenced by inhibiting neurotransmitter release via cell type specific expression of tetanus
toxin (TNT), using these driver lines A01d3 (*SS02065^{splitGAL4}*), A27k (*SS026694^{splitGAL4}*), A02e
(*R70C01^{GAL4}*), A02f (*SS01792^{GAL4}*), A02g (*R36G02^{GAL4}*), and A31k (*SS04399^{splitGAL4}*). **(A)**
Instantaneous %CT over time. The heatmap on top represents change in temperature over time.
(B) Cumulative %CT response for a duration of 5 seconds. **(C)** CT duration in seconds. **(D)** CT
magnitude is an average percent change in area for the duration of stimulus. Controls include
w^{1118}, Empty^{GAL4}' (w;;attP2) or *Empty$^{GAL4''}$*>TNT (w;attP40;attP2). Significant differences were
compared to each GAL4's respective controls dependent on insertion sites. There were no
significant differences detected. Therefore, the following premotor neurons (A01d3, A27k,

A02e, A02f, A02g, and A31k) are not required for cold-evoked CT responses. For each genotype
average n=68.

Figure 6: Cold- and CIII- evoked calcium responses of premotor neurons.

(**A-D**) Ca^{2+} responses of premotor neurons upon cold exposure vs. controls (room temperature).

Neural responses (CaMPARI2) of premotor neuron cell bodies were analyzed using the

following cell type driver lines (**A**) mCSI (*R94B10*[GAL4]) average n=116, (**B**) Chair-1

(*SS00911*[splitGAL4]) average n=66, (**C**) DnB' (*IT4015*[GAL4]) average n=250, and (**D**) DnB''

(*IT412*[GAL4]) average n=245, CaMPARI2 fluorescence ratio is reported as F_{red}/F_{green}. We report

the data as individual datapoints, where the red line represents mean, and hybrid plots (boxplot &

violin) for visualizing the distribution and quartiles of data. Significant stars represent p<0.05,

where comparisons were made to their respective no stimulus controls. (**E-F**) To functionally
assess CIII md neuron to DnB connectivity, we optogenetically activated CIII md neurons
(*83B04^{lexA}>CsChrimson*) and visualized changes in evoked Ca^{2+} using *DnB-GAL4>GCaMP6m*.
Control: No ATR supplemented diet, which is required for optogenetic stimulation in
Drosophila. Orange bars indicate optogenetic stimulation. (**E**) DnB changes in GCaMP reported
as $\Delta F/F_{prestimulus}$, where prestimulus refers to 15 seconds prior to optogenetic stimulation. (**F**)
Maximum DnB neuronal responses ($\Delta F/F_{prestimulus}$) upon optogenetic stimulation. Average n=10.
Significant differences indicated via asterisks, where *p<0.05, **p<0.01, and ****p<0.0001.
990

991
992
993
Figure 6-figure supplement 1: Summary of behavioral and functional roles of premotor
neurons in cold nociception.
Blue arrows indicate strength of synaptic connectivity between the CIII md neurons and second
order neurons. In the behavioral analysis column, the arrow direction indicates reduction or
enhancement of CT response and shading indicates magnitude of change from control. In neural
activity column: NC denotes no significant change in Ca^{2+} response, the arrow direction
indicates reduction or enhancement in evoked Ca^{2+} response and shading indicates magnitude of
change from control. Empty spaces indicate experimental analyses were not performed.

1002
1003
1004
1005 **Ascending interneurons are required for cold nociceptive responses**

1006
Sensory, second order multisensory integration neurons (Basins) and a premotor neuron
(DnB) have further direct synaptic connectivity to a set of projection neurons including A09e,
A08n, A02o, dILP7 and *R61A01^{GAL4}* labeled neurons (labels: A10j, A09o, TePn04, TePn05) that
have previously been implicated in anemotaxis, mechanosensory or chemosensory evoked
behavioral responses (Hu et al., 2017; Jovanic et al., 2019; Kaneko et al., 2017; Masson et al.,
2020; Takagi et al., 2017; Vogelstein et al., 2014). Briefly, A09e receives synaptic inputs from

CIII md, CIV md, A08n, DnB, and TePn05 (**Figure 7A, Figure 7-figure supplement 1A**). A09e
and CIII md neurons are both required for *Drosophila* larval anemotaxis (Jovanic et al., 2019;
Masson et al., 2020). A08n primarily receives inputs from CIV md neurons and very few inputs
from CIII md neurons (**Figure 7A, Figure 7-figure supplement 1A**). However, *Drosophila*
third instar larval synaptic connectivity visualization using GFP reconstituted across synaptic
partners (GRASP) revealed that A08n are not synaptic partners of CIII md neurons (Kaneko et
al., 2017). A08n neurons function downstream of CIV md neurons for noxious chemical and
mechanical nociception (Hu et al., 2017; Kaneko et al., 2017) and neural activation of Ch and
CIII md neurons does not lead to activation of A08n neurons (Tenedini et al., 2019). *R61A01*^{GAL4}
labeled neurons receive inputs from Ch, CIII md, CIV md, Basins, DnB, and A09e (**Figure 7A,**
Figure 7-figure supplement 1A). Neurons labeled by *R61A01*^{GAL4} have been implicated in
mechanosensation (Masson et al., 2020). CIII md neurons synapse onto two additional neurons
that function in processing mechanosensitive information A02o and dILP7. A02o neurons are
command-like neurons required for directional avoidant responses to noxious touch (Takagi et
al., 2017), whereas dILP7 neurons are required for noxious mechanical and/or UV/blue light
avoidance using short neuropeptide F or Ilp7, respectively (Hu et al., 2017; Imambocus et al.,
2022). CIII md to dILP7 synaptic connectivity is observed in the first instar larval EM
connectome, however, assessing synapse connectivity using activity-dependent GRASP in third
instar larvae, there was no detectable connectivity (Hu et al., 2017). We assessed whether the
following projection neurons A09e, A08n, neurons in the expression pattern of the GAL4 line
R61A01 (A10j, A09o, TePn04, TePn05), A02o and dILP7 neurons are required for cold-evoked
behaviors and function in conjunction with CIII md neurons for generating CT behavioral
response.
Neural silencing of projection neurons using cell specific expression of tetanus toxin led
to impairments in cold-evoked behaviors. Neurotransmission inhibition in projection neurons
(A09e and *R61A01*) that receive strong connectivity from CIII md neurons led to significant
reductions in all the CT behavioral metrics that we analyzed (**Figure 7B-E**). As expected, A08n
neurons, which primarily receive inputs form CIV md neurons, are not required for cold-evoked
CT response (**Figure 7B-E**). Whereas neural silencing of A09e or *R61A01* neurons led to greater
than 50% reduction in cold-evoked responses from controls (**Figure 7J**). Furthermore, neural
silencing of dILP7 led to a mild but significant reduction in cold-evoked CT responses, whereas
A02o neurons were dispensable for cold nociception (**Figure 7A, Figure 7-figure supplement**
2).
Next, we assessed whether these projection neurons were able to elicit a CT response
upon neural stimulation. Optogenetic stimulation of projection neurons alone did not lead to any
CT response (**Figure 2-figure supplement 2**). Simultaneous co-activation of CIII md and
R61A01^{GAL4} or A08n neurons led to mild but statistically insignificant increases in CT responses
(**Figure 7F-I, J**). Co-activation of A09e and CIII md neurons led to facilitation of CT behavior,
where compared to controls there were significant increases across all behavioral metrics that we
analyzed (**Figure 7F, G-I**). There were mild, statistically insignificant, reductions in larval

immobility when these projection neurons were co-activated with CIII md neurons, however,

they were significant for A08n (**Figure 7-figure supplement 1B-D**). A09e, which has receives

substantial inputs from CIII md neuron, was sufficient to facilitate CIII md neuron-evoked

behavioral responses (**Figure 7J**).

Our neural silencing analyses suggest that A09e and *R61A01*^{GAL4} (A10j, A09o, TePn04,

TePn05) neurons are required for cold-evoked responses. We hypothesized that these neurons

are cold sensitive and function downstream of CIII md neurons. We predicted that these neurons

would have increases in Ca^{2+} upon cold stimulation or optogenetic activation of CIII md

neurons. A09e neurons present indeed significant increases in Ca^{2+} levels upon cold stimulation,

as measured by cell type-specific expression of CaMPARI2 (**Figure 8A**). Optogenetic activation

of CIII md neurons led to significant increases in evoked Ca^{2+} levels of A09e neurons (**Figure**

8B, C). Multiple stimulations of CIII md neurons did lead to slightly lower levels of evoked Ca^{2+}

but overall Ca^{2+} response was largely similar between stimulations (**Figure 8B, C**). For

R61A01^{GAL4} Ca^{2+} imaging experiments, we restricted our analyses to TePn04 and TePn05

neurons, which were reliably identifiable in an intact *Drosophila* larval preparation. There were

significant increases in CaMPARI2 response of TePn04/05 neurons upon cold stimulation

(**Figure 8D**). Optogenetic stimulation of CIII md neurons led to strong increases in Ca^{2+} levels of

TePn04/05 neurons, however, Ca^{2+} levels rapidly returned to baseline levels (**Figure 8E, F**).

Repeated stimulations of CIII md neurons led to a blunted Ca^{2+} response in TePn04/05 neurons

following the initial stimulation (**Figure 8E, F**). Combined, these data indicate A09e and

R61A01^{GAL4} neurons function downstream of CIII md neurons for cold nociception (**Figure 8-**

figure supplement 1), indicating that ascending neurons relay cold somatosensation to the brain.

1075

1076

1077

1078

1079

Figure 7: Projection neurons downstream of CIII md neurons are required for cold-evoked

responses.

(**A**) A09e, A09o, A10j, TePn04, TePn05, A08n, dILP7 and A02o primarily receive

inputs from CIII md, CIV md, Basin-1, Basin-2 and DnB neurons. Heatmap plot of pre-synaptic

connections to projection neurons. Synaptic connectivity data was extracted from Neurophyla

LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-E**) Cold-evoked responses of

third instar *Drosophila*. Projection neurons were silenced by inhibiting neurotransmitter release

via cell type specific expression of tetanus toxin (TNT), where A09e (*SS00878^{splitGAL4}*),

R61A01^{GAL4} (labels A09o, A10j, TePn04, and TePn05) and A08n (*R82E12^{GAL4}*). (**B**)

Instantaneous %CT over time. The heatmap on top represents change in temperature over time.

(**C**) Cumulative %CT response for a duration of 5 seconds. (**D**) CT duration in seconds. (**E**) CT

magnitude as average percent change in area for the duration of stimulus. Controls include *w^{1118}*,

Empty^{GAL4}' (*w;;attP2*) or *Empty$^{GAL4''}$>TNT* (*w;attP40;attP2*). Significant differences were

compared to each *GAL4's* respective controls dependent on insertion sites. Significant stars:

turquoise stars represent comparison to *w^{1118}*, purple stars represent comparison to

Empty^{GAL4}'>TNT and sea green stars represent comparison to *Empty$^{GAL4''}$>TNT*. For each

genotype average n=72. (**F-I**) Neural co-activation of projection neurons and CIII md neurons.

Here each condition represents expression of ChETA in CIII md neurons plus projection

neurons. (**F**) Instantaneous %CT over time. Blue bar represents optogenetic neural activation.

(**G**) Peak %CT response during optogenetic stimulation. (**H**) CT duration in seconds during

optogenetic stimulation. (**I**) CT magnitude as average percent change in area for the duration of

stimulus. *Empty^{GAL4}'* n=143 and experimental condition average n =35. Significant purple stars

represent comparison to *CIII md + Empty^{GAL4}'>ChETA*. (**J**) Overall percent change from control

for either neural silencing or neural co-activation. The metrics for neural silencing include

cumulative %CT, CT magnitude, and CT duration. The following metrics were used to calculate

percent for neural co-activation: cumulative %CT, peak %CT, CT duration and magnitude.

Significant differences indicated via asterisks, where *$p<0.05$, **$p<0.01$, and ***$p<0.001$.

1107

Figure 7-figure supplement 1: Neural reconstructions and larval mobility for projection
neurons.
(**A**) Individual projection neuron subtypes (magenta) and CIII md neuron axons (blue). Neural
reconstruction data was extracted from Neurophyla LMB Cambridge. (https://neurophyla.mrc-
lmb.cam.ac.uk/). (**B-D**) *Drosophila* larval mobility observed with projection neurons plus CIII
md neuron optogenetic coactivation. (**B**) Instantaneous percent immobility. Blue bar represents
optogenetic neural activation. (**C**) Average percent change in mobility for each genotype, where
greater percent immobility results in larger changes in average mobility. (**D**) Immobility duration
in seconds during stimulation. Significant stars represent p<0.05, where stars represent
comparison to *CIII md + Empty$^{GAL4'}$>ChETA*. *Empty$^{GAL4'}$* n=143 and experimental condition
average n =35. Significant differences indicated via asterisks, where **p<0.01.
1119
1120

1121

Figure 7-figure supplement 2: dLIP7 and A02o neuronal roles in cold nociception.
(**A-D**) Cold-evoked responses of third instar *Drosophila*. Projection neurons were silenced by
inhibiting neurotransmitter release via cell type specific expression of tetanus toxin (TNT) using
driver lines for dILP7 (ILP7^{GAL4}) and A02o (MB120b^{splitGAL4}). (**B**) Instantaneous %CT over time.
The heatmap on top represents change in temperature over time. (**C**) Cumulative %CT response
for a duration of 5 seconds. (**D**) CT duration in seconds. (**E**) CT magnitude as average percent

1128 change in area for the duration of stimulus. Controls include w^{1118}, $Empty^{GAL4}$' (w;;attP2) or
1129 $Empty^{GAL4}$''>TNT (w;attP40;attP2). Significant differences were compared to each GAL4's
1130 respective controls dependent on insertion sites. For each genotype average n=73. Significant
1131 differences indicated via asterisks, where *p<0.05 and **p<0.01.
1132
1133

Figure 8: Cold- and CIII- evoked calcium responses of projection neurons.
Neural responses of A09e ($SS00878^{GAL4}$) (**A-C**) and terminally located TePns (-04, -05) were
analyzed using $R61A01^{GAL4}$(**D-F**). (**A, D**) Ca^{2+} responses of projection neurons upon cold
exposure vs. controls (room temperature). Cold-evoked neural responses (CaMPARI2) of
projection neuron cell bodies were analyzed for (**A**) A09e (n=27) and (**D**) TePns04, & TePn05
(n=42). CaMPARI2 fluorescence ratio is reported as F_{red}/F_{green}. We report the data as individual
datapoints, where the red line represents mean, and hybrid plots (boxplot & violin) for
visualizing the distribution and quartiles of data. Significant stars represent p<0.05, where
comparisons were made to their respective no stimulus controls. (**B-C, E-F**) To assess, if A09e
or TePns functions downstream of CIII md neurons, we optogenetically activated CIII md
neurons ($83B04^{lexA}$>CsChrimson) and visualized changes in evoked Ca^{2+} using projection
neuron specific GAL4>GCaMP6m. Control: No ATR supplemented diet, which is required for
optogenetic stimulation in *Drosophila*. Orange bars indicate optogenetic stimulation. (**B, E**)
Changes in GCaMP reported as $\Delta F/F_{prestimulus}$, where prestimulus refers to 15 seconds prior to
optogenetic stimulation. (**C, F**) Maximum neuronal responses ($\Delta F/F_{prestimulus}$) upon optogenetic

stimulation. A09e average n=22. TePns average n=8. Significant differences indicated via
asterisks, where *p<0.05, ***p<0.001, and ****p<0.0001.
1151

Figure 8-figure supplement 1: Summary of behavioral and functional roles of projection
neurons in cold nociception.
Blue arrows indicate strength of synaptic connectivity between the CIII md neurons and second
order neurons. In the behavioral analysis column, the arrow direction indicates reduction or
enhancement of CT response and shading indicates magnitude of change from control. In neural
activity column: NC denotes no significant change in Ca^{2+} response, the arrow direction
indicates reduction or enhancement in evoked Ca^{2+} response and shading indicates magnitude of
change from control. Empty spaces indicate experimental analyses were not performed.
1160

1161   **Comparative analyses of cold sensitive neurons**

1162

Thus far, data were presented in logical groups based on their previously known cell-
types and functions, however, these neurons function in an interconnected network and
behavioral and functional imaging data must be assessed collectively to study circuit function.
As discussed in previous sections, of all the cell types we tested only optogenetic activation of
CIII md neurons alone is sufficient to elicit CT responses (**Figure 2-figure supplement 2**).
Neural co-activation of CIII md neurons plus additional cell types resulted in marked increases in
CT responses compared to only CIII md neuron activation (**Figure 9A**). Neural silencing of cell
types downstream of CIII md neurons led to significant reductions in cold-evoked CT (**Figure**
9B). To understand how these behavioral phenotypes are interrelated, we performed t-distributed
stochastic neighbor embedding (t-SNE) analysis on both neural silencing and co-activation
behavioral datasets. We identified five different clusters that exhibit varying impacts on
behavioral phenotypes upon either neural co-activation or silencing (**Figure 9C, D**). The first
group clusters together with the control genotype and includes CIV, A05q, Goro, and A08n
neurons. This group on average had less than 25% change in behavioral phenotypes in either
neural co-activation or silencing experiments (**Figure 9C, D**). Basin-1 and DnB neurons
clustered together, where on average they had 49% enhancement of CT response upon neural co-
activation and 45% reduction in cold-evoked CT response upon neural silencing (**Figure 9C, D**).

R61A01 and CIII md neurons formed a cluster, where neural silencing resulted in nearly 60%
reduction cold-evoked behavior and 20% enhancement of CIII md neuron-evoked behaviors
upon neural co-activation (**Figure 9C, D**). A group of neurons including Chair-1, Basin-4 and
Ch were required for cold-evoked behavioral responses, however, these neurons did not facilitate
CIII evoked CT responses (**Figure 9C, D**). In the last group, neural silencing of all Basins,
Basin-2, mCSI, A00c or A09e led to an overall 69% reduction in cold-evoked behaviors and
34% enhancement in CIII evoked CT response (**Figure 9C, D**). Additionally, we assessed how
CIII md neuron synaptic connectivity to post-synaptic neurons could inform whether CIII md
post-synaptic neurons are required for cold nociception (**Figure 9E**). Generally, neural silencing
of highly connected CIII md post-synaptic neurons exhibited greater requirement in cold
nociception, however, there are a few notable exceptions (A02f & A02o), where greater
connectivity did not lead to significant impairment in cold sensitivity (**Figure 9E**).
Analyses of *Drosophila* larval behavioral phenotypes revealed distinct roles for
downstream neurons in cold nociceptive circuitry. Mapping synaptic connectivity of a particular
circuit is only the first step in understanding how individual behaviors arise. Here, we draw
attention to select CIII md neuron first-order neurons (Basins (-2 & -4), DnB, TePns and A09e)
that showed robust requirements for cold nociceptive behaviors and had functional connectivity
to CIII md neurons (**Figure 9F**). TePns synapse onto Basin-4 and form reciprocal connections to
Basin-2, DnB, and A09e (**Figure 9F**). DnB further synapses onto Basin-4 and A09e. Within this
circuit motif one might predict that A09e neurons function as the master integrators, where they
might be computing synaptic information from various sources (**Figure 9F**). Consistent with EM
connectivity, A09e neurons have the highest CIII md neuron-evoked Ca^{2+} responses of the tested
cell types (**Figure 9G, H**). Compared to Basin-4 neurons, Basin-2 neurons receive greater
synaptic input from CIII md neurons (**Figure 9F**). However, CIII md neuron activation leads to
nearly twice as much Ca^{2+} response in Basin-4 neurons than Basin-2 (**Figure 9G, H**). DnB
neurons receive the second highest synaptic input from CIII md neuron and have the second
highest CIII md neuron-evoked Ca^{2+} response (**Figure 9G, H**). TePns receive significantly lower
synaptic input compared to Basin-2 neurons, however, they both exhibit similar levels of CIII md
neuron-evoked Ca^{2+} response (**Figure 9G, H**). Comparative analyses reveal that synaptic
connectivity is informative about functional neural activity, however, additional molecular and
functional studies are required to fully understand roles of these neurons in cold nociception.
Collectively, our findings on neural substrates of cold nociception indicate that second order
multisensory integration by Basin neurons, select pre-motor neurons and projection neurons are
preferentially activated in a stimulus specific manner to elicit appropriate behavioral responses.

Figure 9: Dimensional reduction analysis of *Drosophila* larval behavioral responses and synaptic connectivity informs functional connectivity assessed via Ca^{2+} imaging. (**A-B**) Instantaneous CT proportions for all genotypes in this study. (**A**) Neural co-activation experiments, where CIII plus additional neuronal types were simultaneously optogenetically activated. Controls for optogenetic experiments were tested with or without ATR supplement and include the following conditions: background strain (*w1118*), background strain crossed to *UAS-ChETA*, and *Empty^{GAL4}* crossed to *UAS-ChETA*. Blue bar represents optogenetic stimulation. (**B**) Neurotransmitter release inhibition of individual neuronal types using cell-type specific expression of TnT. (**C-D**) t-distributed stochastic neighbor embedding (t-SNE) analysis of all neuronal subtypes role in both cold nociception (neural silencing data) and CIII md

neuron-evoked CT facilitation (co-activation data). (**C**) 2D plot of t-SNE analysis, where post-
hoc clustering analysis based on "Euclidian complete" method revealed 5 unique groups. The
following percent change from control (*Empty^{GAL4}*) data were included in the analysis: For neural
co-activation (peak % CT response, cumulative % CT response, average % change in area, and
CT duration) and for neural silencing (cumulative % CT response, average % change in area, and
CT duration). (**D**) Average percent change from control for each cluster in '**C**' across all neural
co-activation or neural silencing metrics. (**E**) Synaptic connectivity informs cold-evoked
behavioral responses. Here, for each neuron-type, the percent synaptic input from CIII md
neurons is plotted against cold-evoked CT % upon neuron silencing. There is a negative
correlation between greater synaptic connectivity and lower % CT response. (**F-H**) Analyses of
connectivity upon select circuit components and comparative CIII md neuron-evoked calcium
responses in post-synaptic neurons. (**F**) Proportion of synaptic inputs amongst neurons are
plotted. A09e neurons integrate responses from multiple pathways originating from CIII md
neurons. Network map created using Cytoscape (Shannon et al., 2003). (**G-H**) CIII md neuron-
evoked calcium responses in post-synaptic neurons. (**G**) $\Delta F/F_{baseline}$ over time. (**H**) Average max
$\Delta F/F_{baseline}$. Averages $\pm$ SEM of all trials are plotted. Significant differences indicated via
asterisks, where *p<0.05, **p<0.001, and ****p<0.0001.
1246
1247

1248   **Neuromuscular basis of cold-evoked CT in *Drosophila* larval**
1249

Sensory perception is the first node for interfacing with the external environment, which
is processed through various pathways in the brain eventually leading to motor commands that
generate stimulus specific behavioral responses. Thus far, we have implicated CIII md neurons
as the primary cold sensitive neurons, which transduce thermal cues to select multisensory
integration neurons, premotor neurons and ascending neurons. However, it remains unexplored
how the sensory and central processing of cold leads to CT responses. To this end, assessing CIII
md neuron's premotor neuron network revealed that collectively CIII md premotor neurons have
connectivity to 5 of the 6 major muscle groups in a *Drosophila* larva hemi-segment (**Figure
10A**). Next, we assessed how individual *Drosophila* larva muscular segments respond to noxious
cold. Analysis of muscle localized Ca^{2+} responses revealed that all larval segments have cold-
evoked increases in Ca^{2+} (**Figure 10B-D)**. Interestingly, the strongest cold-evoked response is
present in abdominal segments (A2-A5), whereas anterior (T1-A1) and posterior (A6-A8)
segments have marginally smaller cold-evoked Ca^{2+} responses (**Figure 10B,C**). Similar to cold-
evoked CT behavior, where the head and tail contract towards the center of the animal, muscle
segments in anterior and posterior reach peak activity sooner than the central segments by
average of 4 seconds (**Figure 10B'**). Evaluating individual larval cold-evoked Ca^{2+} levels
revealed segmental responses that can be attributed to previously identified cold-evoked
behavioral responses (*i.e.* CT and head and/or tail raises) (**Figure 10D)** (Turner et al., 2016;
Turner et al., 2018).

Furthermore, we assessed whether impaired circuit function would lead to reduced cold-
evoked muscle responses. Firstly, while maintaining sensory perception, silencing motor neuron
activity using GtACR1 resulted in significant reductions in cold-evoked Ca^{2+} levels compared to
controls (**Figure 10-figure supplement 1A-E**). Secondly, application of ethyl ether also resulted
in significant reductions in cold-evoked Ca^{2+} activity in postsynaptic densities (PSDs) in larval
muscles (**Figure 10-figure supplement 1F-H**).

To achieve greater granularity of how muscles respond to noxious cold, we quantitatively
assessed how individual muscles respond to changes in temperature. Based on electrical
stimulations of abdominal motor nerves, larval body wall muscles have equivalent force
generation capabilities for dorsal, lateral or ventral muscles (Ormerod et al., 2022). Dorsal
longitudinal and oblique (DL and DO) muscles respond acutely and maintain high cold-evoked
Ca^{2+} levels throughout the stimulus period (**Figure 10E-I**). Meanwhile, lateral transverse (LT)
muscles exhibit a gradually increasing cold-evoked Ca^{2+} levels reaching a peak response at the
end of the cold stimulus (**Figure 10E-I**). Lastly, ventral lateral and oblique (VL and VO)
muscles respond most robustly to the onset of cold stimulation which gradually declines over the
course of the stimulation, in contrast to DL and DO muscles, which maintain their levels of Ca^{2+}
activation throughout the stimulus (**Figure 10E-I**). Both dorsal (DO & DL) and ventral (VO &
VL) muscles reach peak response ~10 seconds before lateral muscles (**Figure 10I**). Thus,
Drosophila larvae exhibit stimulus specific muscle activation patterns that are consistent with
cold-evoked behavioral responses arising from sensory processing of temperature changes.
1289

1290

Figure 10: Muscular basis of *Drosophila* larval cold-evoked CT at segmental and individual muscle level.

(**A**) Motor neurons (MN) receiving synaptic input from premotor neurons that are downstream of CIII md neurons. Heatmap plot of pre-synaptic connections to motor neurons. Synaptic connectivity data was extracted from Neurophyla LMB Cambridge. (https://neurophyla.mrc-lmb.cam.ac.uk/). (**B-D**) Analysis of cold-evoked Ca^{2+} responses of muscles in individual segments measured by expressing *jRCaMP1a* using *Mef2^GAL4^*. Larval graphic created using BioRender.com. (**B**) Average $\Delta F/F_{prestimulus}$ of individual larval segments across time. (**B'**) Average cold-evoked Ca^{2+} response in the anterior (T1-A1), central (A2-A5), posterior (A6-A8) segments. Dotted lines mark max $\Delta F/F_{prestimulus}$ and time to max $\Delta F/F_{prestimulus.}$ (**C**) Average $\Delta F/F_{prestimulus}$ of individual larval segments during cold exposure. (**D**) Individual larval average $\Delta F/F_{prestimulus}$ for each larval segments during cold exposure. Individual larval responses exhibit a variety of cold-evoked increases in Ca^{2+}, similar to the range of observed micro-behaviors that lead to CT. Average n= 16. (**E-H**) Analysis of cold-evoked Ca^{2+} of individual muscle cells in *Drosophila* larvae as measured by expressing *jRCaMP1a* using *Mef2^GAL4^* and laser confocal microscopy. (**E**) Schematic of individual muscle cells in a larval hemi-segment from an external view. Muscles that are numerically labeled were analyzed; many ventrally located muscles could not reliably be analyzed. (**F**) Average $\Delta F/F_{prestimulus}$ of individual spatial muscle groups as defined in (Zarin et al., 2019). All muscle groups show cold-evoked Ca^{2+} increase, however, the dynamics of each muscle group are varied. Dorsal longitudinal (DL) and Dorsal oblique (DO) both have strong cold-evoked Ca^{2+} response that is largely stable throughout the stimulus period. Both DL and DO muscles have a peak average $\Delta F/ F_{prestimulus}$ about halfway through the stimulus. Lateral Transverse (LT) muscles show an ever increasing cold-evoked Ca^{2+} response that peaks near the end of the stimulus. Lastly, Ventral Longitudinal (VL) and Ventral Oblique (VO) have cold-evoked Ca^{2+} increase that is highest at the onset of stimulus and decays gradually throughout the stimulus period. (**G**) Cold-evoked average $\Delta F/F_{prestimulus}$ of individual muscles across time. (**H**) Cold-evoked average max $\Delta F/F_{prestimulus}$ of individual muscles across time. Average n= 16. (**I**) Heatmap representation of average time max $\Delta F/F_{prestimulus}$ for individual muscles.

1321

Figure 10 -figure supplement 1: Motor neuron silencing or pharmacological anesthesia results in significant reductions in cold-evoked muscle activity

(A-E) Cold-evoked muscle Ca^{2+} responses with functional motor neurons and optogenetically silenced motor neurons. (**A-B**) Heatmap of average $\Delta F/F_{prestimulus}$ GCaMP6f responses in segments T1-A4. Segments A5-A8 were not analyzed due to high autofluorescence of the digestive tract. (**C-D**) Heatmap of average $\Delta F/F_{prestimulus}$ GCaMP6f responses individual larvae and segments during the stimulus period. When motor neurons are silenced, nearly all analyzed segments have the similar cold-evoked responses compared to controls, where we certain segments reaching very high average $\Delta F/F_{prestimulus}$. (**E**) Average muscle average $\Delta F/F_{prestimulus}$ GCaMP6f response comparison for each segment between control and silenced motor neurons. Average n= 13 larvae. Genotype: 44H10-GCaMP6f, D42>GtACR1 ± ATR. (**F-H**) Muscle postsynaptic density (PSD) Ca^{2+} responses were assessed using SynapGCaMP6f (MHC-CD8-GCaMP6f-Sh) in *Drosophila* larval segment T3/A1. In controls, there is a robust cold-evoked increase in Ca^{2+} levels, however, upon ethyl ether application, there is a significant

1336 reduction in evoked Ca^{2+} levels. (**F**) Average $\Delta F/F_{prestimulus}$ over time for all stimulations. (**G**)
1337 Average $\Delta F/F_{prestimulus}$ during stimulus of all quantified PSDs for each animal across three
1338 stimulations. (**H**) Average $\Delta F/F_{prestimulus}$ of individual PSDs across three stimulations. Average
1339 n=3 larvae with average PSDs n=171.
1340
1341 **Discussion**
1342
Environmental stimuli are detected by peripheral sensory neurons, which transduce
relevant cues to downstream central circuitry responsible for processing multisensory input and
generating stimulus-relevant behavioral responses. The study of the diverse molecular and
cellular mechanisms involved in sensory discrimination have yielded key insights into how
animals interact with their environment (Arnadóttir et al., 2011; Bandell & Patapoutian, 2009;
Chrispin et al., 2013; Corfas & Vosshall, 2015; Coste et al., 2010; Derby et al., 2016; Dietrich et
al., 2022; Fowler & Montell, 2013; Freeman & Dahanukar, 2015; Himmel et al., 2017; Leung &
Montell, 2017; Montell, 2021; Wu et al., 2018; Xiao & Xu, 2021). However, how animals
distinguish between innocuous and noxious stimuli at molecular, cellular and circuit level
remains an important question in modern neuroscience (Bushnell et al., 1985; Dannhauser et al.,
2020; Himmel et al., 2023; Imambocus et al., 2022; Moehring et al., 2018; Patel et al., 2022;
Turner et al., 2016; Ward et al., 1996).
Here, we assessed the downstream circuitry of a multimodal sensory (CIII md) neuron
that detects both innocuous mechanical and noxious cold temperatures. In CIII md neurons,
detection of innocuous cues occurs via low threshold activation and noxious cues are detected
via high threshold activation leading to stimulus-relevant behaviors (Turner et al., 2016).
Transduction of nociceptive cold stimulus is predominantly mediated by multimodal Ch and CIII
md neurons. However, nociceptive peripheral sensory neurons, CIII md and CIV md neurons,
function through shared downstream neural circuitry. We describe the behavioral and functional
requirements of multisensory integration (Basins) neurons, premotor (DnB and mCSI) neurons
and projection (A09e and TePns) neurons in noxious cold-evoked behavioral responses. We
identified circuit components that play a role in amplifying noxious cues and differentiating
between opposing noxious (heat versus cold) stimuli-evoked behaviors at various levels of
sensory processing. Finally, we characterize how stimulus-evoked muscle contractions lead to
cold nociceptive behavioral response. Our findings provide key insights into how environmental
cues are processed by multiplexed networks for multisensory integration and decision making.
1369
1370 **Basin-1 neurons function across sensory modalities as gain modulators for noxious stimuli-**
1371 **evoked escape responses.**
1372
Accurately responding to potentially harmful stimuli by appropriately executing
energetically expensive escape behaviors is critical for animal survival. Incorrectly performing
escape responses in absence of dangerous stimuli can have long-term detrimental consequences.

Therefore, the need for neural mechanisms responsible for integrating various noxious and
innocuous sensory stimuli for accurately executing stimulus-appropriate escape responses. Here,
we focus on the role of Basin neurons, known to mediate various escape responses (Jovanic et
al., 2016; Masson et al., 2020; Ohyama et al., 2015).
EM connectome analyses revealed that somatosensory (Ch, CIII md and CIV md)
neurons all synapse onto multisensory integrating Basin neurons, and that amongst the Basins
subtypes, Basin-1 neurons are presynaptic to Basin-2,-3 and -4 neurons (Masson et al., 2020;
Ohyama et al., 2015). On the basis of the known connectome, we analyzed the function of Basin
neurons in cold-evoked escape behaviors. Our findings indicate that optogenetic activation of
Basin-1 is not sufficient to evoke noxious cold-evoked behavioral response (**Figure 2 –**
Supplement 2) and it was known that thermogenetic activation of Basin-1 is also not sufficient
for evoked rolling escape (Ohyama et al., 2015). Therefore, Basin-1 neuron activity alone is not
sufficient to elicit a noxious stimuli-evoked behavioral response. However, due to intricate
connectivity amongst the Basins, the total integrated output of multiple Basin neurons may lead
to threshold activation of noxious stimulus-evoked escape responses. And indeed, thermogenetic
co-activation assays found that Basin-1 neural activity could facilitate a Basin-4 -mediated
rolling response (Ohyama et al., 2015), and also the activation of Basin-1 neurons leads to the
activation of Basin-2 neurons through lateral disinhibition (Jovanic et al., 2016). Here, we found
that co-activation of Basin-1 plus CIII md neurons led to strong-evoked CT responses (**Figure**
2). Taken together, these studies concluded that Basin-1 facilitates the activation of other Basin
neurons, and our results also provide additional support for the role of Basin-1 neurons in fine-
tuning behavioral outcomes and enhancing cumulative output of multisensory integration Basin
neurons leading stimulus specific escape responses.
1399
1400 **Multisensory integration neurons drive behavioral selection downstream of sensory**
1401 **neurons.**
1402
In order to evaluate the functional and behavioral roles of neurons within a circuit, a
comprehensive analysis of both upstream and downstream synaptic connectivity is necessary.
Basin-2 and -4 neurons integrate different amounts of synaptic input from somatosensory
neurons, other Basin neurons, feedforward/back inhibitory neurons, and projection neurons.
Basin-4 neurons receive greater excitatory synaptic input compared to Basin-2 neurons, which
have greater connectivity with local inhibitory neurons (Jovanic et al., 2016; Masson et al., 2020;
Ohyama et al., 2015). Larger inhibitory connectivity of Basin-2 neurons likely arises from much
broader downstream connectivity including several projection/descending neurons (A00c, A02o,
A05q, A10j, TePn04), premotor neurons (A02m, A02n, Chair-1), descending feedback inhibitory
neuron (SeIN128), local feedback neurons (Handle-a/-b), whereas Basin-4 downstream
connectivity is restricted to limited set of neurons (A00c, A02o, Basin-3, Chair-1) (Zhu et al.,
2024). These synaptic level differences in multisensory integration neurons underlie noxious
stimuli-specific evoked responses and suggest that CIII md neuron mediated responses primarily

function through Basin-2 neurons and CIV md neuron mediated environmental cues are
processed via Basin-4 neurons.
1418
Chemical nociception in *Drosophila* larvae primarily functions via CIV md neurons.
Among the md neurons, CIII md neurons are the least sensitive to noxious chemical stimulus
(Lopez-Bellido et al., 2019). Analyses of Basin neuron requirement in chemical nociception
revealed that all Basins are necessary for chemical-evoked rolling, where comparative analyses
of Basin subtypes revealed that chemical nociception is primarily mediated through Basin-4 with
weaker behavioral phenotypes of Basin-1 or -2 (Lopez-Bellido et al., 2019). Our work shows
that inhibition of neurotransmission in Basin-2 or Basin-4 neurons leads to significant reductions
in cold nociceptive responses, where across the analyzed behavioral metrics Basin-2 had stronger
deficits in cold-evoked responses (**Figure 2**). Furthermore, neural co-activation analyses
revealed that Basin-2 plus CIII md led to enhanced CT response, whereas Basin-4 plus CIII md
co-activation did not facilitate but rather suppressed the CT response (**Figure 2**). Basin-2 neural
activity likely has cascading effects on functions of downstream neurons (i.e. TePns, A02m/n,
Chair-1, & A00c), which we also found to be required for cold nociception. Behavioral
assessment of multisensory integration neurons supports the notion that CIII md neuron sensory
input is processed primarily by Basin-2 and CIV md neuron sensory cues are processed via
Basin-4 neurons.
1435
Despite the accumulating evidence that cold-evoked responses are primarily mediated by
Basin-2, our Ca^{2+} analyses revealed that compared to Basin-2, Basin-4 neurons have much
greater cold- or CIII md neuron- evoked Ca^{2+} responses. There are a few possible explanations
for greater Basin-4 activity levels. Firstly, greater Basin-4 Ca^{+2} activity could be due to greater
synaptic input from projection neurons (TePn04/05) and premotor neurons (DnB), which are
both required for cold nociception and exhibit cold- or CIII md neuron-evoked Ca^{2+} responses
(**Figure 7, Figure 8 – Supplement 1**) (Masson et al., 2020). Secondly, reduced Basin-2 Ca^{2+}
responses could be due to greater connectivity with feedback/feedforward inhibitory local
neurons (Jovanic et al., 2016; Masson et al., 2020) or descending feedback neuron (SeIN128)
(Zhu et al., 2024). Descending inhibitory, SeIN128, neuron functions in transitioning from
rolling to fast crawling response in *Drosophila* larvae (Zhu et al., 2024). SeIN128 receives
synaptic input from Basin-2 but not Basin-4, while providing feedback inhibition to Basins (-1, -
2, -3), premotor neurons (A02m/n, Chair-1 & DnB) and projection neurons (A00c, A02o, &
A05q) (Ohyama et al., 2015; Zhu et al., 2024). Thus, lower cold-/CIII md neuron-evoked Basin-
2 neural activity could be due to greater feedback inhibitory input compared to Basin-4.
Comparative Ca^{2+} imaging or electrophysiological analyses can further delineate the precise
interplay between Basin-2 and -4 neurons in integrating sensory information across
somatosensory modalities leading to distinct behavioral responses. However, to date, synaptic
connectivity and behavioral analyses collectively indicate an initial sensory discrimination node
amongst the Basin neurons, where downstream processing of CIII mediated sensory input is

conveyed through Basin-2 and its downstream connectivity, meanwhile CIV somatosensory
information is transduced through Basin-4 neuronal pathway.
1458

1459 **Differential roles of pre-motor neurons lead to behavioral selection.**

1460

Animal locomotive behavioral responses are implemented through neurosecretory
systems and muscle contractions evoked by motor neuron activity. Motor neurons integrate a
conglomerate of neural impulses from premotor neurons, leading to appropriate muscle
activation patterns to implement specific behavioral responses (Bellardita & Kiehn, 2015;
Berkowitz et al., 2010; Carreira-Rosario et al., 2018; Green & Soffe, 1996; Huang & Zarin,
2022; Liao & Fetcho, 2008; Talpalar et al., 2013; Zarin et al., 2019). Recent work in premotor
and motor neuron connectomics has provided evidence for both labeled line and combinatorial
connectivity between promotor and motor neurons that give rise to co-active motor neurons
states leading to selective muscle group activation (Cooney et al., 2023; Huang & Zarin, 2022;
Zarin et al., 2019). Various *Drosophila* larval premotor neuron subtypes and/or motor pools have
been implicated in locomotion, nociception, and innocuous mechanosensation (Burgos et al.,
2018; Cooney et al., 2023; Huang & Zarin, 2022; Kohsaka et al., 2017; Kohsaka et al., 2014;
Kohsaka et al., 2019; Yoshino et al., 2017; Zarin et al., 2019). Neural reconstruction efforts in
Drosophila larvae have revealed that somatosensory (CIII md and CIV md) neurons are not
directly connected to motor neurons, but rather feed into polysynaptic pathways leading to motor
neurons via premotor neurons (such as the DnB, mCSI, & Chair-1 neurons studied here)
(Cooney et al., 2023; Jovanic et al., 2019; Ohyama et al., 2015; Winding et al., 2023; Yoshino et
al., 2017). Premotor neurons (DnB, mCSI, and Chair-1) that receive synaptic input from CIII md
neurons are required for cold-evoked responses. However, only premotor neurons previously
implicated in nociceptive escape responses, DnB and mCSI, can facilitate CIII-evoked CT
responses (Burgos et al., 2018; Lopez-Bellido et al., 2019; Yoshino et al., 2017). Meanwhile,
Chair-1 neurons, which are required for innocuous mechanosensation (Jovanic et al., 2019), do
not facilitate CIII md neuron mediated CT responses. We showed, with Ca^{2+} imaging, that Chair-
1 neurons are not activated upon cold exposure. It remains unclear whether Chair-1 neuronal
function is sufficient to drive specific sensory evoked behaviors or requires a combinatorial pre-
motor neuronal activity for generating evoked behavioral responses. Given the data, we speculate
that Chair-1 neurons likely gate innocuous-to-noxious stimulus-evoked behaviors; however,
additional functional studies are required to tease out how Chair-1 neurons function in behavioral
selection.

1490

1491 **Heterogeneity in thermosensory processing reflects variety of thermal stimuli.**

1492

1493   Animals detect changes in temperature through a variety of sensors that occupy and
1494 function in different spatiotemporal timescales. Thermosensory systems function to detect
1495 changes at the epidermal, brain and visceral levels, and these broad thermosensory inputs must

be integrated to generate appropriate behavioral responses (Nakamura, 2018). The underlying neural circuitry for thermosensation differs based on body plan and type of thermosensation. In mice, cool and warm sensory signals, which are detected by TRP channels, are transmitted to the spinal dorsal horn, where distinct cool/warm pathways relay epidermal changes in temperature to thalamocortical regions for perception and discrimination (Nakamura, 2018). In *Drosophila* larvae and adults, thermotaxis occurs through temperature detection via dorsal organ ganglion (DOG) and antennal thermoreceptors, respectively (Frank et al., 2015; Gallio et al., 2011; Hernandez-Nunez et al., 2021; Klein et al., 2015; Liu et al., 2003). Investigations of circuit bases of thermotaxis have revealed different strategies for larval and adult systems. Adult antennal thermoreceptors have distinct hot or cold sensors, which project to the proximal antennal lobe (PAL), where hot and cold information is segregated into discrete regions (Gallio et al., 2011; Macpherson et al., 2015). Thermal coding properties of thermosensory projections neurons (tPNs), whose dendrites are located in PAL, include both slow/fast adapting tPNs and broadly/narrowly tuned tPNs (Frank et al., 2017). *Drosophila* larval thermosensory DOG contains both warm- and cool- sensing cells that are connected to individual cool or warm projection neurons (Hernandez-Nunez et al., 2021). DOG thermosensory information is further integrated by integration projection neurons (iPNs), which receive inputs from both warm and cool projection neurons (Hernandez-Nunez et al., 2021). The *Drosophila* larval noxious thermosensory system has distinct hot (CIV md) or cold (CIII md) sensing neurons (Babcock et al., 2009; Tracey et al., 2003; Turner et al., 2016). Larval somatosensory thermal inputs mediated by CIII md neurons are heavily integrated by second order neurons (Basins, DnB, A09e, etc.), where population coding leads to cold-evoked behavioral response. Noxious cold stimulus is encoded downstream of CIII md neurons by ascending neurons (A09e and TePns) that allow for systems-level integration of multiple thermal inputs including those from cool/warm sensitive larval DOG neural circuits and cold/heat information from CIII/CIV md neurons, respectively. Thermal information from the larval DOG and somatosensory neurons converges onto the mushroom body, where cool or warm projection neurons downstream of DOG and feed-forward projection neurons downstream of ascending neurons (A09e) synapse onto the same mushroom body output neuron (Eichler et al., 2017; Hernandez-Nunez et al., 2021; Winding et al., 2023). Thus far, ascending pathways into the brain have been characterized for larval cold nociception, however, descending output neurons responsible for cold-evoked behaviors remains unidentified. Diversity in processing thermosensory stimuli reflects heterogeneity at molecular, cellular, and circuit levels that collectively function to maintain proper thermal homeostasis and ultimately behavioral action selection. The *Drosophila* larval thermosensory system is well-suited to dissect mechanisms underlying central nervous system integration of varying spatial and functional thermosensory information by utilizing various molecules, sensory systems, and interconnected circuits for appropriate behavioral responses.

Muscle activity patterns underpin behavioral responses

Animals interact with the external environment by detecting external cues via sensory
systems that are transduced to CNS, where motor commands are generated to elicit stimulus
relevant muscle activity patterns leading to behavioral responses. *Drosophila* larval body wall
muscle have unique behavior relevant muscle activity patterns for locomotion and nociceptive
responses (rolling and CT) (Cooney et al., 2023; Heckscher et al., 2012; Liu et al., 2023; Zarin et
al., 2019, This work). In forward locomotion, posterior segments contract before anterior
segments generating a forward contraction wave, where the longitudinal (dorsal & ventral)
muscles contract first during the wave phase and lateral muscles contract during the interwave
interval (Heckscher et al., 2012; Liu et al., 2023; Zarin et al., 2019). Exposure to nociceptive
heat, mechanical or chemical cues larvae execute a corkscrew rolling response (Babcock et al.,
2009; Hu et al., 2017; Hu et al., 2020; Hwang et al., 2012; Im et al., 2018; Im et al., 2015;
Kaneko et al., 2017; Lopez-Bellido et al., 2019; Ohyama et al., 2013; Ohyama et al., 2015;
Robertson et al., 2013; Tracey et al., 2003). In rolling response, multiple abdominal segments
respond synchronously for executing a roll (Cooney et al., 2023). Furthermore, within a hemi-
segment, the muscle activation pattern moves circumferentially during the roll durations, where
DL, DO, and VL muscles are active during the bend phase and LT, VO, & VA muscles
transition the larval hemi-segment into a stretch phase completing a roll (Cooney et al., 2023). In
cold-evoked CT responses, anterior and posterior segments reach peak activation before central
segments. Within a hemi-segment individual muscles groups have unique activation profiles,
where upon cold exposure both dorsal and ventral muscles reach peak response before lateral
muscles. Thus, we can conclude the following. For both Cold-evoked CT and locomotion,
longitudinal (dorsal and ventral) muscles contract before lateral muscles, whereas rolling
response has a mixed activation pattern. The larval segments responds uniquely and dependent
on the behavior being generated. In forward locomotion had a propagating wave of activation
from posterior to anterior, rolling has a synchronous activation of segments and in CT response,
anterior/posterior segments contract before central. The underlying circuitry responsible for
generating these three unique behaviors differs based on sensory, central and motor processing
modules.
1564
1565 **Methods**
1566
Fly Strains
All *Drosophila melanogaster* strains used in this study are listed in (**Table 1**). All
Drosophila reagents were maintained on standard cornmeal-molasses-agar diet in 12:12 hour
light-dark cycle at ~22°C. All experimental crosses were raised in 12:12 hour light-dark cycle at
29°C, unless otherwise stated. We used two separate CIII md neuron driver lines in this
study:*19-12^{GAL4}*, *R83B04^{GAL4}* and *R83B04^{lexA}*.
1573
Molecular cloning and transgenic generation of *CIII^{lexA}*

GMR83B04^{GAL4} was characterized as CIII md neuron driver using both optogenetics and
visualization using membrane markers (Galindo et al., 2023; Himmel et al., 2023; Patel et al.,
2022). The R83B04 enhancer containing entry vector was a gift from FlyLight team at Janelia
Research Campus, Ashburn, VA. We performed Gateway cloning to insert R83B04 enhancer
upstream of lexA. LR reaction was performed using R83B04 containing entry vector and lexA
containing (pBPLexA::p65Uw) destination vector (Addgene: 26231). Transgenic fly generation
was conducted by GenetiVision. *R83B04*^{lexA} was inserted in VK20 docking site using PhiC31
integrase mediated transformation.
1583
EM connectomics
All data for synaptic connectivity, circuit diagrams and wire frame projections of neural
cell types were extracted from Neurophyla LMB Cambridge (https://neurophyla.mrc-
lmb.cam.ac.uk/). Relevant primary literature sources are cited within the main text. Sensory
neurons from abdominal segments 1-4 were analyzed and for the remaining cell types, all
reconstructed neurons were included (date accessed: March 04, 2022).
1590
Cold Plate Assay
We assessed requirements of neurons downstream from peripheral sensory neurons in
cold-evoked responses using the cold plate assay (Patel & Cox, 2017; Patel et al., 2022; Turner
et al., 2016). *GAL4* drivers for select neuronal subtypes were used to express the light chain of
tetanus toxin (*UAS-TNT*). We used two sets of controls: *w1118*, genetic background control, and
Empty^{GAL4}, which contains a GAL4 construct but no regulatory promoter (*GAL4* only control).
Genetic crosses were raised on standard cornmeal-molasses-agar diet and at 29°C. To assess
cold-evoked behavioral responses of age matched *Drosophila* third instar larvae, we exposed the
ventral surface to noxious cold (10°C) temperatures. Briefly, using a brush we remove third
instar larvae from food and place them on wet Kimwipe. Food debris is removed passively by
allowing the larvae to freely locomote on wet Kimwipe. We place 6-8 larvae on a thin black
metal plate that is subsequently placed on a pre-chilled (10°C) Peltier plate TE technologies
Peltier plate (CP-031, TC-48-20, RS-100-12). Larval responses are recorded from above using
Nikon DSLR (D5300). Changes in larval surface area were extracted using FIJI and Noldus
Ethovision XT (https://github.com/CoxLabGSU/CaMPARI-intensity-and-cold-plate-assay-
analysis/branches) (Patel et al., 2022). Next, we isolated individual larva from each video,
removed background and used Ethovision to measure larval surface area. Using custom built r
scripts, we compiled data from each larva and each genotype. Utilizing r, we calculated percent
change in larval surface area (Area change= (Area$_N$ – Average_Area$_{baseline}$)/
Average_Area$_{baseline}$*100). From the percent change in area dataset, we report three behavioral
metrics: Average change in area, which is average percent change in area for the stimulus
duration. We defined cold-evoked CT response as change in area of -10% or less for at least 0.5
consecutive seconds. CT duration, time spent at or below -10% change in area. Lastly, percent

CT response, which is cumulative percent of animals that CT for at least 0.5 consecutive
seconds.
Statistical analysis: We performed the following statistical tests for all cold plate assay
data analysis. %CT response: Fisher's exact with Benjamini-Hochberg for multiple comparison.
We used r for performing comparisons of percent behavior response between genotypes, we used
Benjamini-Hochberg multiple comparison correction. CT duration: Kruskal-Wallis with
Benjamini, Krieger and Yekutieli for multiple comparisons. CT duration data are not normal. CT
magnitude: One-way ANOVA with Holm-Šídák's for multiple comparisons.
1622
Neural activation via optogenetics
We performed two types of neural activations to assess sufficiency for CT response:
single downstream neural activation and co-activation, where we simultaneously activated CIII
md neurons and individual downstream neurons. For optogenetic experiments in *Drosophila*, a
light sensitive co-factor all *trans*-retinal (ATR) is required. For all conditions, all adult animals in
the genetic cross were placed in ATR (1500μM) supplemented food and subsequently F1
progeny were also raised in food containing ATR and raised in dark. For control condition, we
used an *Empty*^{GAL4} containing GAL4 construct but no regulatory promoter. Optogenetic
experiments were conducted using a similar setup as previously described (Patel et al., 2022).
Briefly, we used principles of dark field microscopy to enhance signal to noise ratio and capture
high resolution larval videos. We created a custom dark field stage, where a Canon DSLR T3i
camera captures video from above. Neural activation is performed by two blue led lights that are
controlled remotely using the Noldus control box (Thorlabs: DC4100, DC4100-hub, and two
M470L3-C4 led light. Noldus: mini-IO box). Larval behaviors are directly captured using
Noldus Ethovision XT software, which also controls blue led activation. All optogenetic
experiments were conducted in a dimly lit room. Third instar *Drosophila* larva were removed
from food plug and placed onto wet Kimwipe, where larval locomotion allowed for passive
removal of food debris. We lightly sprayed water onto a thin glass plate, then placed a single
larva for optogenetic stimulation. The glass plate was manually moved on XY-axis to keep larva
in the field of view. The following stimulus paradigm was used: 5 seconds of baseline (light off)
and 5 seconds of neural activation (blue led lights on).
We performed video processing and behavioral analysis using FIJI and data compilation
and analysis using r (https://github.com/CoxLabGSU/Drosophila_larval_optogenetic_analysis-
area_and_mobility). In order to analyze *Drosophila* larval behavioral responses, we first
automatically stabilized (XY axis) and then measured changes in larval surface area and mobility
(described below).
Raw videos from Noldus Ethovision XT were uncompressed using video-to-video
convertor (https://www.videotovideo.org/). The following steps were scripted in FIJI macro
language for automatic video processing and data acquisition. For increasing processing speed,
the uncompressed videos were automatically cropped to dimensions to contain all of the larva's
movement. Next, using a pre-determined threshold, we created a mask followed by background

removal using erode, remove outlier and dilate functions. We then used the 'Analyze Particles'
function to obtain XY coordinates of the larva in each frame. Larval movements were stabilized
using XY coordinates and the 'Translate' function was used to create a highly stabilized video.
Next, we measured larval surface area using automatic thresholding 'Huang method' and
'Analyze Particles' to obtain area. We define *Drosophila* larval mobility as changes in occupied
pixels between two frames. We used our stabilized larval video to measure larval mobility,
where larval peristaltic movements (XY displacement) are not captured. However, changes in
occupied pixels resulting from turning and head sweeps are captured. Specifically, larval
mobility was measured by subtracting thresholded larva in each frame from the previous frame
(Raw mobility = Thresholded larva$_{\text{Frame N}}$ - Thresholded larva$_{\text{Frame N-1}}$).

Data compilation and analysis was performed in r using custom scripts. Optogenetically-
evoked changes in behavior were analyzed independently for mobility and changes in area. For
each larva, we calculated percent change in area (Area change= (Area$_N$ – Average_Area$_{\text{baseline}}$)/
Average_Area$_{\text{baseline}}$*100). We measured CT duration as the amount of time the larva has -10%
or lower change in area. We also measure CT magnitude by analyzing average change area for
stimulus duration. We report percent instantaneous CT over time as percent of animals that are at
or below -10% change in area. Additionally, we report peak %CT from percent instantaneous CT
dataset for each genotype. For analysis of mobility, we calculated percent change in mobility
(Mobility change= (Mobility$_N$ – Average_ Mobility$_{\text{baseline}}$)/ Average_ Mobility$_{\text{baseline}}$*100). We
report average percent change in mobility during stimulus for each genotype. We also plot
percent immobility, which is calculated by the percent of animals with -25% or more reduction
in mobility. Immobility duration is calculated based on amount of time individual animals spend
at or below -25% mobility.

Statistical analysis: We performed the following statistical tests for all behavioral
optogenetic data analysis. %CT peak response: Fisher's exact with Benjamini-Hochberg for
multiple comparison. We used r for performing comparisons of percent behavior response
between genotypes, we used Benjamini-Hochberg multiple comparison correction.
CT/immobility duration: Kruskal-Wallis with Benjamini, Krieger and Yekutieli for multiple
comparisons. CT duration data are not normal. CT/immobility magnitude: One-way ANOVA
with Holm-Šídák's for multiple comparisons.

1684

1685

1686   **CT behavior across life stages - Cold Plate Assay**

1687

For assessing cold-evoked larval behavior across life stages, we developed a specialized
behavior apparatus that improved to signal to noise ratios, allowed for similar stimulus
magnitude between trial and across larval life stages, and an improved assay throughput. The
apparatus consists of VWR circulating water chiller (model: MX7LR20), two aquarium mini-
pumps (model: AD20P-1230D), custom 3D-printed water chamber with glass 1 inch glass walls,
a remote control to activate the aquarium pumps (Syantek model: BHZ0320U-RF), three-way

water valve (Malida model: 607569660968), iPad screen protector (Ailun model:
B0BJQ24MK7), Canon R6 with RF50mm F1.8 and RF85mm F2 macro lens and led lights for
illumination from all sides (Daybetter model: FLSL-DB-501215RGBUS). The larvae were
removed and cleaned as described above, then placed on a thin transparent glass plate (screen
protector). The glass plate bearing larvae is placed on a 3D-printed water chamber, then pre-
chilled water is piped into the chamber using mini-aquarium pumps. After the water touches the
bottom of the glass plate a three-way water valve is turned off and the chilled water remains in
the 3D-printed water chamber providing cold stimulation to the larvae through the glass surface.
At the end of the trial, the water valve is opened to drain the water into the VWR water chiller.
For first and second instar larval assays, a 85mm RF macro lens was used at different distances
to increase magnification. A 50mm RF lens was used to image third instar larval cold-evoked
responses.
1706
To quantitatively analyze the larval responses, we used a slightly modified versions of
FIJI and r macros as described above. In the FIJI macro, individual larvae were extracted into
their individual video file and the background was removed. A technical challenge arose in the
new assay, where each larva gets the cold stimulus at slightly different time point as the water
filling up in the water chamber. There is a minor but consistent change in brightness/contrast for
each larva, when the water below the glass plate touches the bottom of the glass plate. Using
Noldus Ethovision, changes in brightness/contrast are quantified and then using r, the time of
stimulus delivery is obtained. After determining the first frame automatically in r, the analysis
for various cold-evoked CT response metric are obtained using the same r code as described
above.
1717
1718   Statistical analysis: We performed the following statistical tests for all behavioral data
1719   analysis. CT magnitude and duration: One-way ANOVA with Holm-Šídák's for multiple
1720   comparisons.
1721
1722
CT behavior across life stages – Optogenetic Assay
To evaluate CIII-evoked CT responses across life stages, we utilized optogenetic neural
activation of CIII md neurons using *ChR2-H134R* ± ATR. In order to maintain consistent blue-
light based neural activation, we used Canon R6 camera mounted on AxioScope V16 Zeiss
microscope for visualizing *Drosophila* larvae at life stage specific magnification while applying
oblique blue light exposure (Thorlabs: DC4100, DC4100-hub, and one M470L3-C4 led light).
We recorded larva responses without blue light (pre-stimulus) for 5 seconds and 5 seconds of
blue light exposure. To quantitatively analyze larval responses, we used a similar method as
described above for neural activation.
1732

1733   Statistical analysis: We performed the following statistical tests for all behavioral
1734 optogenetic data analysis. CT magnitude and duration: One-way ANOVA with Holm-Šídák's for
1735 multiple comparisons.
1736
1737 **Cold-evoked Ca^{2+} responses of muscles in *Drosophila* larvae**
1738
For visualizing cold-evoked Ca^{2+} responses of larval muscles in each segment, we
expressed *jRCaMP1a* in *Drosophila* larval muscles using *mef2^{GAL4}* and motor neuron silencing
using GtACR1 with muscle Ca^{2+} imaging. Cold stimulus was delivered using CherryTemp
temperature controller (Cherry Biotech: Cherry Temp dual channel temperature control), which
allows for rapid changes in temperature. To analyze *Drosophila* larval cold-evoked segmental
response, we used AxioScope V16 Zeiss microscope with a Canon R6 camera mounted. *In vivo*
jRCaMP1a responses were observed in third instar larva, where the larva was mounted with
dorsal midline centered on the top. Using FIJI, changes in jRCaMP1a fluorescence intensities in
each segment were quantified with manually drawn regions of interest (ROI). We report changes
in jRCaMP1a fluorescence as ΔF/F =(F-F$_{prestimulus}$)/ F$_{prestimulus}$*100 and average ΔF/F for each
larval segment.
1750
Drosophila larva's individual muscles cold-evoked Ca^{2+} responses were visualized using
mef2^{GAL4}> jRCaMP1a and Zeiss LSM 780 confocal microscope with EC Plan-Neofluar 10x
objective. Time-lapse acquisition was acquired at 500.11µm x 500.11µm x 307.2ms using
561nm laser wavelength. Three different imaging planes (dorsal, lateral, & ventral) were used to
capture the majority of the muscles of a hemi-segment. Using FIJI, changes in jRCaMP1a
fluorescence intensities in each muscle were quantified with manually drawn ROIs. We report
changes in jRCaMP1a fluorescence as ΔF/F =(F-F$_{prestimulus}$)/ F$_{prestimulus}$*100 and maximum ΔF/F
for each muscle. No statistical analyses were performed.
1759
1760   The following temperature exposure paradigm was used for both segmental and
1761 individual muscle cold-evoked Ca^{2+} response assays: Pre-stimulus (25°C for 45 seconds),
1762 temperature ramp down, steady state cold (5°C for 15 seconds), temperature ramp up, and post-
1763 stimulus (25°C for 40 seconds). Temperature ramp duration 10 seconds.
1764
We used the above-mentioned Cherry Temp. dual channel temperature control system,
Canon R6 and Zeiss V16 to visualize cold-evoked Ca^{2+} responses for assessing delay in mouth
hook retraction (10°C stimulus) and ethyl ether pharmacological application and muscle post-
synaptic density Ca^{2+} imaging (5°C stimulus). We used the following temperature exposure
paradigm: pre-stimulus (20 seconds), 10 second ramp down, steady state cold (10 seconds),
temperature ramp up, baseline (10 seconds), temperature ramp down, steady state cold (10
seconds), temperature ramp up, baseline (10 seconds), temperature ramp down, steady state cold
(10 seconds), temperature ramp up, and baseline (20 seconds). The temperature ramp duration

was10 seconds. For muscle Ca^{2+} fluorescence analyses, we used $\Delta F/F = (F-F_{prestimulus})/$
$F_{prestimulus}*100$. For assessing, CIII md Ca^{2+} increase and delay in mouth hook retraction, we used
$\Delta F/F = (F-F_{prestimulus})/ F_{prestimulus}*100$ for CIII md Ca^{2+} levels and visually determined the time at
which the mouth retracted.

1777

1778 Statistical analysis: Delay in mouth hook retraction - Mann-Whitney test. For motor
1779 neuron silencing assay - Two-way ANOVA with Holm-Šídák's for multiple comparisons. For
1780 pharmacology assay - Mann-Whitney test for non-parametric data and Welch's t-test for
1781 parametric data.

1782

1783

1784

CaMPARI Imaging
Post-synaptic neuron CaMPARI2 imaging: For assessing cold-evoked Ca^{2+} responses of sensory
neurons and CIII md neuron downstream neurons, we utilized CaMPARI2, which upon
photoconverting light and high intracellular Ca^{2+} stably photoconverts fluorescence from green
to red. We performed the cold plate assay as described above. For stimulus condition, the Peltier
plate was set to noxious cold (6°C) temperature and for control condition, the Peltier plate was
turned off (room temperature). We placed individual third instar larvae onto the Peltier plate and
simultaneously exposed the animal to photoconverting light for 20 seconds. CaMPARI2
fluorescence was imaged in live, intact larvae via confocal microscopy. Zeiss LSM 780 Axio
examiner microscope, Plan-Apochromat 20x objective, and excitation wavelengths of 561nm
and 488nm were used to image larval ventral nerve cord or peripheral sensory neurons. We
mounted live intact larva onto microscope slide and immobilized the larva by placing a
coverslip, as previously described (Im et al., 2018; Patel & Cox, 2017; Patel et al., 2022; Turner
et al., 2016). Three dimensional CaMPARI2 fluorescence in ventral nerve cord localized
downstream neurons was imaged at 607.28µm x 607.28µm (XY resolution) and 2µm z-slices.
Regions of interests were identified and area normalized fluorescence intensity for red & green
signals were obtained via Imaris 9.5 software. CaMPARI responses are reported as ratio of
F_{red}/F_{green}. Statistical comparisons: Parametric data – Welch's t-test and non-parametric data –
Mann-Whitney test.

1804

PNS CAMPARI2 imaging: Sensory neuron CaMPARI2 responses were analyzed using custom
FIJI macros that automatically detected cell bodies and sholl intensity analyses were performed
using semi-automated custom FIJI macros (https://github.com/CoxLabGSU/CaMPARI-intensity-
and-cold-plate-assay-analysis/branches) (Patel et al., 2022).
Cell body analysis – We created a set of three sequential macros that draw ROIs around
the cell body, user verification of the ROIs and lastly quantification of fluorescence intensities.
The first custom FIJI script generates maximum intensity projections of z-stacks, image masks
were created by thresholding (Moments method) GFP signal, and next, background and dendritic

branches removed using erode and dilate functions. At the end of background clearing only cell
bodies remain, where Analyze particle function is used to draw ROIs around soma. The second
FIJI script is used for manual verification of each ROI and manually redrawing any incorrect
ROIs. Lastly, upon ROI verification, area normalized F_{red} and F_{green} intensities are quantified. As
previously described (Fosque et al., 2015; Im et al., 2018; Patel & Cox, 2017; Patel et al., 2022;
Turner et al., 2016), we report evoked photoconverted CaMPARI signal as F_{red}/F_{green} ratio.
Statistical comparisons: Non-parametric data – Mann-Whitney test.
Sholl intensity analysis – Sholl intensity analysis was performed using a set of two
custom FIJI scripts as previously described (Patel et al., 2022). Briefly, we first perform
background clearing by manually thresholding the GFP signal and select all branches and soma
of neuron of interest using the "Wand Tool" in FIJI and then a mask of neuron of interest is
created. The second FIJI script draws five-pixel wide radial ROIs at a single pixel interval, here
only the dendrites and soma from the neuron of interest are selected at each radial interval. After
Sholl ROIs are drawn, area and area normalized F_{red} and F_{green} fluorescence intensities are
extracted for each radial step away from the soma. Similar to CaMPARI cell body analysis, we
CaMPARI2 signal as F_{red}/F_{green} ratios away from the soma.
Ventral nerve cord CaMPARI imaging: We utilized Pan-neural ($R57C10^{GAL4}>CaMPARI$)
driver to visualize ventral nerve cord Ca^{2+} responses to various stimuli including innocuous
touch, noxious heat (45°C) and noxious cold (6°C). Stimulus and photoconverting light were
delivered for 20 seconds. Whole ventral nerve cord was imaged at 607.28µm x 607.28µm (XY
resolution) and 2µm z-slices. The rest of the stimulus delivery and imaging was similar to
previously described CaMPARI experiments. No statistical analyses were performed.
1835
CIII activation and second order neuron GCaMP imaging
CIII md neuron-evoked responses in downstream neurons were evaluated by using
optogenetics and GCaMP. We expressed *lexAop-CsCrimson* in CIII md neurons using
$R83B04^{lexA}$ and used downstream neuron specific *GAL4* to drive expression of *GCaMP6m*. For
experimental condition, all adult animals in the genetic cross were reared in ATR (1500µM)
supplemented food and subsequently F1 progeny were also raised in food containing ATR. For
control condition, adult flies and F1 progeny were raised in standard cornmeal-molasses-agar
diet. Both control and experimental crosses were reared in 24hr dark. We mounted live intact
third instar larva in between microscope slide and a coverslip. Larval ventral nerve cord and cells
of interest were located using epifluorescence on Zeiss LSM 780 confocal microscope. Larval
GCaMP6 responses were allowed to return to baseline for at least 2 minutes. Time-lapse
acquisition was imaged at 250.06µm x 250.06µm x 307.2ms using 488nm laser wavelength. CIII
md neural activation was performed using two oblique 617nm leds (Thorlabs M617F2 and
M79L01) that were manually operated using Thorlabs led controller (Thorlabs DC4100 and
DC4100-hub). We performed three sequential neural activations using the following paradigm
was used: Baseline light off (30 seconds)-> neural activation (617nm for 15 seconds)-> light off
(30 seconds)-> neural activation (617nm for 15 seconds)-> light off (30 seconds)-> neural

activation (617nm for 15 seconds)-> light off (30 seconds). Time-lapse videos were stabilized
using Stack reg – Rigid transformation in FIJI (Linkert et al., 2010; Schindelin et al., 2012;
Thevenaz et al., 1998). Regions of interest were manually drawn and area normalized GCaMP
fluorescence over time was exported. We report changes in GCaMP6m fluorescence as $\Delta F/F$
$=(F-F_{prestimulus})/ F_{prestimulus}*100$ and max $\Delta F/F$ for each of the three neural activation epochs.
Statistical comparisons: Parametric data – Welch's t-test and non-parametric data – Mann-
Whitney test.

Statistics and data visualization

Statistical analyses were performed using r (Fisher's exact test) and GraphPad Prism. All
graphical visualization of the data were created using Prism GraphPad. Details on specific
statistical tests are listed in respective methods section.

1864

1865

1866 *Table 1: Drosophila melanogaster strains used in this study*

Designation	Nomenclature	Source or reference	Identifier	Insertion site
w1118		BDSC	3605	
CaMPARI2	*UAS-CaMPARI2*	BDSC	78317	attP5
ChETA	*UAS-ChETA::YFP*	BDSC	36495	attP2
ChR2	*UAS-ChR2*	BDSC	9681	Chr 3
ChR2-H134R	*UAS-H134R-ChR2*	BDSC	28995	Chr 2
CsChrimson	*LexAop2-CsChrimson.tdTomato*	BDSC	82183	VK00005
CsChrimson	*UAS-IVS-CsChrimson.mCherry*	BDSC	82180	VK00005
GCaMP6m	*UAS-GCaMP6m*	BDSC	42748	attP40
Pan-neural CaMPARI	*R57C10^{GAL4}, UAS-CaMPARI*	BDSC	58763	VK00040/VK00020
RCaMP	*UAS-jRCaMP1a*	BDSC	63792	VK00005
SynapGCaMP6f	*Mhc-SynapGCaMP6f*	BDSC	67739	Chr 3
Muscle GCaMP + GtACR1	*w;;44H10::GCaMP6f, UAS-GtACR1*	Gift from Zarin Lab		
TnT	*UAS-TeTxLC.tnt-E2*	BDSC	28837	Chr 2
Ch	*IAV - GAL4*	BDSC	52273	Chr 3
CIII	*19-12 - GAL4*			Chr 3
CIII	*R83B04 - GAL4*	BDSC	41309	attP2
CIII	*R83B04 - lexA*	This Study		
CIII>ChETA	*19-12^{GAL4}, UAS-ChETA::YFP*	BDSC	36495	Chr 3,attP2
CIV	*ppk - GAL4*	BDSC	32079	Chr 3

Mef2	*Mef2-GAL4*	BDSC	27390	Chr 3
Empty[GAL4']	*w;;attP2*	BDSC	68384	attP2
Empty[GAL4'']	*w;attP40;attP2*	Gift from Rubin Lab		attP40;attP2
A00c	*R71A10 - GAL4*	BDSC	39562	attP2
A01d3	*SS02065 - splitGAL4*	Gift from Nose Lab		attP40;attP2
A02e	*R70C01 - GAL4*	BDSC	39520	attP2
A02f	*SS01792 - GAL4*	Gift from Zlatic & Ohyama Labs		attP40;attP2
A02g	*R36G02 - GAL4*	BDSC	49939	attP2
A02o	*MB120B - splitGAL4*	Gift from Nose Lab		attP40;attP2
A05q	*R47D07 - GAL4*	BDSC	50304	attP2
A08n	*R82E12 - GAL4*	BDSC	40153	attP2
A09e	*SS00878 - splitGAL4*	Gift from Zlatic Lab		attP40;attP2
A27k	*SS026694 - splitGAL4*	Gift from Nose Lab		attP40;attP2
A31k	*SS04399 - splitGAL4*	Gift from Zarin Lab		attP40;attP2
All Basin	*R72F11 - GAL4*	BDSC	39786	attP2
Basin-1	*R20B01 - GAL4*	BDSC	48877	attP2
Basin-2	*SS00739 - splitGAL4*	Gift from Zlatic Lab		attP40;attP2
Basin-4	*SS00740 - splitGAL4*	Gift from Zlatic Lab		attP40;attP2
Chair-1	*SS00911 - splitGAL4*	Gift from Zlatic Lab		attP40;attP2
dILP7	*ILP7 – GAL4*	Gift from Jan Lab		
DnB'	*IT4051 - GAL4*	Gift from Grueber Lab		Chr 3
DnB''	*IT412 - GAL4*	BDSC	63300	Chr 3
Goro	*R69F06 - GAL4*	BDSC	39497	attP2
mCSI	*R94B10 - GAL4*	BDSC	41325	attP2
R61A01/TePns	*R61A01 - GAL4*	BDSC	39269	attP2
D42	*D42 – GAL4*	BDSC	8816	Chr 3

1867

Data availability statement

1869 The original contributions presented in the study are included in the article/Supplementary
1870 material, further inquiries can be directed to the corresponding author.

1871

Conflict of Interest

1873 The authors declare that the research was conducted in the absence of any commercial or
1874 financial relationships that could be construed as a potential conflict of interest.
1875

Author Contributions

1877 Conceptualization: AAP and DNC. Methodology: AAP, and DNC. Cold-plate assays: AAP;
1878 Optogenetics, AAP; Calcium imaging: AAP. Statistics and other formal analyses: AAP. EM
1879 Connectome analysis: AAP and AC. Writing – Original Draft: AAP and DNC; Writing – Review
1880 & Editing: AAP, AC and DNC. Visualization: AAP. Supervision: DNC. Funding acquisition:
1881 DNC.
1882

Funding

1884 This work was supported by NIH R01 NS115209 (to DNC). AAP was supported by a Brains &
1885 Behavior Fellowship, a 2CI Neurogenomics Fellowship, and a Kenneth W. and Georganne F.
1886 Honeycutt Fellowship from Georgia State University. AC thanks the Wellcome Trust (award
1887 205038/Z/16/Z and 205038/A/16/Z) and the HHMI Janelia Resarch Campus for funding.
1888

Acknowledgments

We thank members of Cox Lab and Michael J. Galko (MD Anderson Cancer Center) for critical
comments on the manuscript. We thank Shatabdi Bhattacharjee for sharing insightful feedback
on the manuscript. We thank the Janelia Visiting Scientist program hosted by HHMI Janelia
Research Campus for providing critical training in EM connectomics. We thank members of the
Drosophila research community for sharing reagents: Gerald M. Rubin, Marta Zlatic, Tomoko
Ohyama, Aref Zarin, Akinao Nose, Yuh Nung Jan, Lily Jan and Wesley Grueber. We
acknowledge the Georgia State University Imaging Core Facility for training and instrument
support associated with this work.
1898

References
1900 Andrade, I. V., Riebli, N., Nguyen, B. M., Omoto, J. J., Cardona, A., & Hartenstein, V. (2019,
1901   Feb 4). Developmentally Arrested Precursors of Pontine Neurons Establish an Embryonic

Blueprint of the Drosophila Central Complex. *Curr Biol, 29*(3), 412-425 e413. https://doi.org/10.1016/j.cub.2018.12.012

Arnadóttir, J., O'Hagan, R., Chen, Y., Goodman, M. B., & Chalfie, M. (2011, 2011/8//). The DEG/ENaC protein MEC-10 regulates the transduction channel complex in Caenorhabditis elegans touch receptor neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 31*(35), 12695-12704. https://doi.org/10.1523/JNEUROSCI.4580-10.2011

Babcock, D. T., Landry, C., & Galko, M. J. (2009, 2009/5//). Cytokine signaling mediates UV-induced nociceptive sensitization in Drosophila larvae. *Current biology : CB, 19*(10), 799-806. https://doi.org/10.1016/j.cub.2009.03.062

Babcock, Daniel T., Shi, S., Jo, J., Shaw, M., Gutstein, Howard B., & Galko, Michael J. (2011, 2011/9//). Hedgehog Signaling Regulates Nociceptive Sensitization. *Current Biology, 21*(18), 1525-1533. https://doi.org/10.1016/j.cub.2011.08.020

Balakrishnan, M., Yu, S. F., Chin, S. M., Soffar, D. B., Windner, S. E., Goode, B. L., & Baylies, M. K. (2020, 2020/07/21/). Cofilin Loss in Drosophila Muscles Contributes to Muscle Weakness through Defective Sarcomerogenesis during Muscle Growth. *Cell Reports, 32*(3), 107893. https://doi.org/https://doi.org/10.1016/j.celrep.2020.107893

Bandell, M., & Patapoutian, A. (2009, 2009/12//). Itching for insight. *Cell, 139*(7), 1224-1226. https://doi.org/10.1016/j.cell.2009.12.012

Bellardita, C., & Kiehn, O. (2015, Jun 1). Phenotypic characterization of speed-associated gait changes in mice reveals modular organization of locomotor networks. *Curr Biol, 25*(11), 1426-1436. https://doi.org/10.1016/j.cub.2015.04.005

Berck, M. E., Khandelwal, A., Claus, L., Hernandez-Nunez, L., Si, G., Tabone, C. J., Li, F., Truman, J. W., Fetter, R. D., Louis, M., Samuel, A. D. T., & Cardona, A. (2016). The wiring diagram of a glomerular olfactory system. https://doi.org/10.7554/eLife.14859.001

Berkowitz, A., Roberts, A., & Soffe, S. R. (2010). Roles for multifunctional and specialized spinal interneurons during motor pattern generation in tadpoles, zebrafish larvae, and turtles. *Front Behav Neurosci, 4*, 36. https://doi.org/10.3389/fnbeh.2010.00036

Berndt, A., Schoenenberger, P., Mattis, J., Tye, K. M., Deisseroth, K., Hegemann, P., & Oertner, T. G. (2011, May 3). High-efficiency channelrhodopsins for fast neuronal stimulation at low light levels. *Proceedings of the National Academy of Sciences of the United States of America, 108*(18), 7595-7600. https://doi.org/10.1073/pnas.1017210108

Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G., & Deisseroth, K. (2005, Sep). Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci, 8*(9), 1263-1268. https://doi.org/10.1038/nn1525

Burgos, A., Honjo, K., Ohyama, T., Qian, C. S., Shin, G. J.-e., Gohl, D. M., Silies, M., Tracey, W. D., Zlatic, M., Cardona, A., & Grueber, W. B. (2018, 2018/3//). Nociceptive interneurons control modular motor pathways to promote escape behavior in Drosophila. *eLife, 7*. https://doi.org/10.7554/eLife.26016

Bushnell, M. C., Duncan, G. H., Dubner, R., Jones, R. L., & Maixner, W. (1985, May). Attentional influences on noxious and innocuous cutaneous heat detection in humans and monkeys. *J Neurosci, 5*(5), 1103-1110. https://doi.org/10.1523/JNEUROSCI.05-05-01103.1985

Caldwell, J. C., Miller, M. M., Wing, S., Soll, D. R., & Eberl, D. F. (2003, 2003/12/23). Dynamic analysis of larval locomotion in Drosophila chordotonal organ mutants. *Proceedings of the National Academy of Sciences, 100*(26), 16053-16058. https://doi.org/10.1073/pnas.2535546100

Carreira-Rosario, A., Zarin, A. A., Clark, M. Q., Manning, L., Fetter, R. D., Cardona, A., & Doe, C. Q. (2018, 2018/8//). MDN brain descending neurons coordinately activate backward and inhibit forward locomotion. *eLife, 7*. https://doi.org/10.7554/eLife.38554

Chrispin, J., Martin, S. S., Hasan, R. K., Joshi, P. H., Minder, C. M., McEvoy, J. W., Kohli, P., Johnson, A. E., Wang, L., Blaha, M. J., & Blumenthal, R. S. (2013, Sep). Landmark lipid-lowering trials in the primary prevention of cardiovascular disease. *Clin Cardiol, 36*(9), 516-523. https://doi.org/10.1002/clc.22147

Clark, M. Q., Zarin, A. A., Carreira-Rosario, A., & Doe, C. Q. (2018, 2018/12//). Neural circuits driving larval locomotion in Drosophila. *Neural Development, 13*(1), 6-6. https://doi.org/10.1186/s13064-018-0103-z

Cooney, P. C., Li, W., Huang, Y., Hormigo, R., Tabachnik, T., Hillman, E. M. C., Grueber, W. B., & Zarin, A. A. (2023). Neuromuscular Basis of Drosophila larval escape behavior. *bioRxiv*, 2023.2002.2001.526733. https://doi.org/10.1101/2023.02.01.526733

Corfas, R. A., & Vosshall, L. B. (2015, 2015/12//). The cation channel TRPA1 tunes mosquito thermotaxis to host temperatures. *eLife, 4*. https://doi.org/10.7554/eLife.11750

Coste, B., Mathur, J., Schmidt, M., Earley, T. J., Ranade, S., Petrus, M. J., Dubin, A. E., & Patapoutian, A. (2010, 2010/10//). Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. *Science (New York, N.Y.), 330*(6000), 55-60. https://doi.org/10.1126/science.1193270

Dannhauser, S., Lux, T. J., Hu, C., Selcho, M., Chen, J. T., Ehmann, N., Sachidanandan, D., Stopp, S., Pauls, D., Pawlak, M., Langenhan, T., Soba, P., Rittner, H. L., & Kittel, R. J. (2020, Sep 30). Antinociceptive modulation by the adhesion GPCR CIRL promotes mechanosensory signal discrimination. *eLife, 9*. https://doi.org/10.7554/eLife.56738

Demontis, F., & Perrimon, N. (2009, Mar). Integration of Insulin receptor/Foxo signaling and dMyc activity during muscle growth regulates body size in Drosophila. *Development, 136*(6), 983-993. https://doi.org/10.1242/dev.027466

Derby, C. D., Kozma, M. T., Senatore, A., & Schmidt, M. (2016, Jun). Molecular Mechanisms of Reception and Perireception in Crustacean Chemoreception: A Comparative Review. *Chem Senses, 41*(5), 381-398. https://doi.org/10.1093/chemse/bjw057

Dietrich, S., Company, C., Song, K., Lowenstein, E. D., Riedel, L., Birchmeier, C., Gargiulo, G., & Zampieri, N. (2022, Nov 11). Molecular identity of proprioceptor subtypes innervating different muscle groups in mice. *Nat Commun, 13*(1), 6867. https://doi.org/10.1038/s41467-022-34589-8

Eichler, K., Li, F., Litwin-Kumar, A., Park, Y., Andrade, I., Schneider-Mizell, C. M., Saumweber, T., Huser, A., Eschbach, C., Gerber, B., Fetter, R. D., Truman, J. W., Priebe, C. E., Abbott, L. F., Thum, A. S., Zlatic, M., & Cardona, A. (2017, 2017/8//). The complete connectome of a learning and memory centre in an insect brain. *Nature, 548*(7666), 175-182. https://doi.org/10.1038/nature23455

Eschbach, C., Fushiki, A., Winding, M., Schneider-Mizell, C. M., Shao, M., Arruda, R., Eichler, K., Valdes-Aleman, J., Ohyama, T., Thum, A. S., Gerber, B., Fetter, R. D., Truman, J. W., Litwin-Kumar, A., Cardona, A., & Zlatic, M. (2020, Apr). Recurrent architecture for adaptive regulation of learning in the insect brain. *Nat Neurosci, 23*(4), 544-555. https://doi.org/10.1038/s41593-020-0607-9

Eschbach, C., & Zlatic, M. (2020, Dec). Useful road maps: studying Drosophila larva's central nervous system with the help of connectomics. *Curr Opin Neurobiol, 65*, 129-137. https://doi.org/10.1016/j.conb.2020.09.008

Fosque, B. F., Sun, Y., Dana, H., Yang, C. T., Ohyama, T., Tadross, M. R., Patel, R., Zlatic, M., Kim, D. S., Ahrens, M. B., Jayaraman, V., Looger, L. L., & Schreiter, E. R. (2015, 2015/2//). Labeling of active neural circuits in vivo with designed calcium integrators. *Science, 347*(6223), 755-760. https://doi.org/10.1126/science.1260922

Fowler, M. A., & Montell, C. (2013, 2013/3//). Drosophila TRP channels and animal behavior. *Life sciences, 92*(8-9), 394-403. https://doi.org/10.1016/j.lfs.2012.07.029

Frank, D. D., Enjin, A., Jouandet, G. C., Zaharieva, E. E., Para, A., Stensmyr, M. C., & Gallio, M. (2017, 2017/8//). Early Integration of Temperature and Humidity Stimuli in the Drosophila Brain. *Current Biology, 27*(15), 2381-2388.e2384. https://doi.org/10.1016/j.cub.2017.06.077

Frank, D. D., Jouandet, G. C., Kearney, P. J., Macpherson, L. J., & Gallio, M. (2015, 2015/3//). Temperature representation in the Drosophila brain. *Nature, 519*(7543), 358-361. https://doi.org/10.1038/nature14284

Freeman, E. G., & Dahanukar, A. (2015). Molecular neurobiology of Drosophila taste. *Current Opinion in Neurobiology, 34*, 140-148. https://doi.org/10.1016/j.conb.2015.06.001

Fushiki, A., Zwart, M. F., Kohsaka, H., Fetter, R. D., Cardona, A., & Nose, A. (2016, 2016/2//). A circuit mechanism for the propagation of waves of muscle contraction in Drosophila. *eLife, 5*. https://doi.org/10.7554/eLife.13253

Galindo, S. E., Wood, A. J., Cooney, P. C., Hammond, L. A., & Grueber, W. B. (2023, Mar 1). Axon-axon interactions determine modality-specific wiring and subcellular synaptic specificity in a somatosensory circuit. *Development, 150*(5). https://doi.org/10.1242/dev.199832

Gallio, M., Ofstad, T. A., Macpherson, L. J., Wang, J. W., & Zuker, C. S. (2011, Feb 18). The coding of temperature in the Drosophila brain. *Cell, 144*(4), 614-624. https://doi.org/10.1016/j.cell.2011.01.028

Garner, A. (2020). *Identification of neural premotor substrates affecting escape behaviours in Drosophila larvae* [Master's thesis, McGill University]. https://escholarship.mcgill.ca/concern/theses/th83m403v

Gerhard, S., Andrade, I., Fetter, R. D., Cardona, A., & Schneider-Mizell, C. M. (2017, 2017/10//). Conserved neural circuit structure across Drosophila larval development revealed by comparative connectomics. *eLife, 6*. https://doi.org/10.7554/eLife.29089

Green, C. S., & Soffe, S. R. (1996, 1996/02/01). Transitions between two different motor patterns in Xenopus embryos. *Journal of Comparative Physiology A, 178*(2), 279-291. https://doi.org/10.1007/BF00188169

Gunaydin, L. A., Yizhar, O., Berndt, A., Sohal, V. S., Deisseroth, K., & Hegemann, P. (2010, 2010/3//). Ultrafast optogenetic control. *Nature Neuroscience, 13*(3), 387-392. https://doi.org/10.1038/nn.2495

He, L., Gulyanon, S., Skanata, M. M., Karagyozov, D., Heckscher, E., Krieg, M., Tschekpenakis, G., Gershow, M., & Tracey, D. (2018, 2018/11//). Proprioceptive neurons of larval Drosophila melanogaster show direction selective activity requiring the mechanosensory channel TMC. *bioRxiv*, 463216-463216. https://doi.org/10.1101/463216

Heckscher, E. S., Lockery, S. R., & Doe, C. Q. (2012). Characterization of *Drosophila* Larval Crawling at the Level of Organism, Segment, and Somatic Body Wall Musculature. *The Journal of Neuroscience, 32*(36), 12460. https://doi.org/10.1523/JNEUROSCI.0222-12.2012

Heckscher, Ellie S., Zarin, Aref A., Faumont, S., Clark, Matthew Q., Manning, L., Fushiki, A., Schneider-Mizell, Casey M., Fetter, Richard D., Truman, James W., Zwart, Maarten F., Landgraf, M., Cardona, A., Lockery, Shawn R., & Doe, Chris Q. (2015, 2015/10//). Even-Skipped+ Interneurons Are Core Components of a Sensorimotor Circuit that

Maintains Left-Right Symmetric Muscle Contraction Amplitude. *Neuron, 88*(2), 314-329. https://doi.org/10.1016/J.NEURON.2015.09.009

Herculano-Houzel, S. (2009). The human brain in numbers: a linearly scaled-up primate brain. *Front Hum Neurosci, 3*, 31. https://doi.org/10.3389/neuro.09.031.2009

Hernandez-Nunez, L., Chen, A., Budelli, G., Berck, M. E., Richter, V., Rist, A., Thum, A. S., Cardona, A., Klein, M., Garrity, P., & Samuel, A. D. T. (2021). Synchronous and opponent thermosensors use flexible cross-inhibition to orchestrate thermal homeostasis. *Science Advances, 7*(35), eabg6707. https://doi.org/doi:10.1126/sciadv.abg6707

Himmel, N. J., Letcher, J. M., Sakurai, A., Gray, T. R., Benson, M. N., & Cox, D. N. (2019, Nov 11). Drosophila menthol sensitivity and the Precambrian origins of transient receptor potential-dependent chemosensation. *Philos Trans R Soc Lond B Biol Sci, 374*(1785), 20190369. https://doi.org/10.1098/rstb.2019.0369

Himmel, N. J., Letcher, J. M., Sakurai, A., Gray, T. R., Benson, M. N., Donaldson, K. J., & Cox, D. N. (2021, Jun 25). Identification of a neural basis for cold acclimation in Drosophila larvae. *iScience, 24*(6), 102657. https://doi.org/10.1016/j.isci.2021.102657

Himmel, N. J., Patel, A. A., & Cox, D. N. (2017). *Invertebrate Nociception* (Vol. 1). Oxford University Press. https://doi.org/10.1093/acrefore/9780190264086.013.166

Himmel, N. J., Sakurai, A., Donaldson, K. J., & Cox, D. N. (2022, Sep 16). Protocols for measuring cold-evoked neural activity and cold tolerance in Drosophila larvae following fictive cold acclimation. *STAR Protoc, 3*(3), 101510. https://doi.org/10.1016/j.xpro.2022.101510

Himmel, N. J., Sakurai, A., Patel, A. A., Bhattacharjee, S., Letcher, J. M., Benson, M. N., Gray, T. R., Cymbalyuk, G. S., & Cox, D. N. (2023, Jan 23). Chloride-dependent mechanisms of multimodal sensory discrimination and nociceptive sensitization in Drosophila. *eLife, 12*. https://doi.org/10.7554/eLife.76863

Hiramoto, A., Jonaitis, J., Niki, S., Kohsaka, H., Fetter, R. D., Cardona, A., Pulver, S. R., & Nose, A. (2021, May 19). Regulation of coordinated muscular relaxation in Drosophila larvae by a pattern-regulating intersegmental circuit. *Nat Commun, 12*(1), 2943. https://doi.org/10.1038/s41467-021-23273-y

Hu, C., Petersen, M., Hoyer, N., Spitzweck, B., Tenedini, F., Wang, D., Gruschka, A., Burchardt, L. S., Szpotowicz, E., Schweizer, M., Guntur, A. R., Yang, C.-H., & Soba, P. (2017, 2017/8//). Sensory integration and neuromodulatory feedback facilitate Drosophila mechanonociceptive behavior. *Nature Neuroscience, 20*(8), 1085-1095. https://doi.org/10.1038/nn.4580

Hu, Y., Wang, C., Yang, L., Pan, G., Liu, H., Yu, G., & Ye, B. (2020, Dec 21). A Neural Basis for Categorizing Sensory Stimuli to Enhance Decision Accuracy. *Curr Biol, 30*(24), 4896-4909 e4896. https://doi.org/10.1016/j.cub.2020.09.045

Huang, Y., & Zarin, A. A. (2022). A premotor microcircuit to generate behavior-specific muscle activation patterns in Drosophila larvae. *bioRxiv*, 2022.2008.2018.504452. https://doi.org/10.1101/2022.08.18.504452

Huckesfeld, S., Schlegel, P., Miroschnikow, A., Schoofs, A., Zinke, I., Haubrich, A. N., Schneider-Mizell, C. M., Truman, J. W., Fetter, R. D., Cardona, A., & Pankratz, M. J. (2021, Jun 4). Unveiling the sensory and interneuronal pathways of the neuroendocrine connectome in Drosophila. *eLife, 10*. https://doi.org/10.7554/eLife.65745

Hwang, R. Y., Stearns, N. A., & Tracey, W. D. (2012, 2012/1//). The Ankyrin Repeat Domain of the TRPA Protein Painless Is Important for Thermal Nociception but Not Mechanical Nociception. *PLoS One, 7*(1), e30090-e30090. https://doi.org/10.1371/journal.pone.0030090

Im, S. H., Patel, A. A., Cox, D. N., & Galko, M. J. (2018, 2018/5//). <i>Drosophila</i> Insulin receptor regulates the persistence of injury-induced nociceptive sensitization. *Disease Models & Mechanisms, 11*(5), dmm034231-dmm034231. https://doi.org/10.1242/dmm.034231

Im, S. H., Takle, K., Jo, J., Babcock, D. T., Ma, Z., Xiang, Y., & Galko, M. J. (2015). Tachykinin acts upstream of autocrine Hedgehog signaling during nociceptive sensitization in Drosophila. *eLife, 4*. https://doi.org/10.7554/eLife.10735

Imambocus, B. N., Zhou, F., Formozov, A., Wittich, A., Tenedini, F. M., Hu, C., Sauter, K., Macarenhas Varela, E., Heredia, F., Casimiro, A. P., Macedo, A., Schlegel, P., Yang, C. H., Miguel-Aliaga, I., Wiegert, J. S., Pankratz, M. J., Gontijo, A. M., Cardona, A., & Soba, P. (2022, Jan 10). A neuropeptidergic circuit gates selective escape behavior of Drosophila larvae. *Curr Biol, 32*(1), 149-163 e148. https://doi.org/10.1016/j.cub.2021.10.069

Imura, E., Shimada-Niwa, Y., Nishimura, T., Huckesfeld, S., Schlegel, P., Ohhara, Y., Kondo, S., Tanimoto, H., Cardona, A., Pankratz, M. J., & Niwa, R. (2020, Jun 8). The Corazonin-PTTH Neuronal Axis Controls Systemic Body Growth by Regulating Basal Ecdysteroid Biosynthesis in Drosophila melanogaster. *Curr Biol, 30*(11), 2156-2165 e2155. https://doi.org/10.1016/j.cub.2020.03.050

Jenett, A., Rubin, Gerald M., Ngo, T.-T. B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, Barret D., Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, Joanna H., Peng, H., Trautman, Eric T., Svirskas, Robert R., Myers, Eugene W., Iwinski, Zbigniew R., Aso, Y., DePasquale, Gina M., Enos, A., Hulamm, P., Lam, Shing Chun B., Li, H.-H., Laverty, Todd R., Long, F., Qu, L., Murphy, Sean D., Rokicki, K., Safford, T., Shaw, K., Simpson, Julie H., Sowell, A., Tae, S., Yu, Y., & Zugates, Christopher T.

(2012, 2012/10//). A GAL4-Driver Line Resource for Drosophila Neurobiology. *Cell Reports, 2*(4), 991-1001. https://doi.org/10.1016/j.celrep.2012.09.011

Jovanic, T., Schneider-Mizell, C. M., Shao, M., Masson, J.-B., Denisov, G., Fetter, R. D., Mensh, B. D., Truman, J. W., Cardona, A., & Zlatic, M. (2016, 2016/10//). Competitive Disinhibition Mediates Behavioral Choice and Sequences in Drosophila. *Cell, 167*(3), 858-870.e819. https://doi.org/10.1016/j.cell.2016.09.009

Jovanic, T., Winding, M., Cardona, A., Truman, J. W., Gershow, M., & Zlatic, M. (2019, Feb 18). Neural Substrates of Drosophila Larval Anemotaxis. *Curr Biol, 29*(4), 554-566 e554. https://doi.org/10.1016/j.cub.2019.01.009

Kaneko, T., Macara, A. M., Li, R., Hu, Y., Iwasaki, K., Dunnings, Z., Firestone, E., Horvatic, S., Guntur, A., Shafer, O. T., Yang, C.-H., Zhou, J., & Ye, B. (2017, 2017/8//). Serotonergic Modulation Enables Pathway-Specific Plasticity in a Developing Sensory Circuit in Drosophila. *Neuron, 95*(3), 623-638.e624. https://doi.org/10.1016/j.neuron.2017.06.034

Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y. K., Morimoto, T. K., Chuong, A. S., Carpenter, E. J., Tian, Z., Wang, J., Xie, Y., Yan, Z., Zhang, Y., Chow, B. Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton, M., Wong, G. K.-S., & Boyden, E. S. (2014, 2014/3//). Independent optical excitation of distinct neural populations. *Nature Methods, 11*(3), 338-346. https://doi.org/10.1038/nmeth.2836

Klein, M., Afonso, B., Vonner, A. J., Hernandez-Nunez, L., Berck, M., Tabone, C. J., Kane, E. A., Pieribone, V. A., Nitabach, M. N., Cardona, A., Zlatic, M., Sprecher, S. G., Gershow, M., Garrity, P. A., & Samuel, A. D. (2015, Jan 13). Sensory determinants of behavioral dynamics in Drosophila thermotaxis. *Proc Natl Acad Sci U S A, 112*(2), E220-229. https://doi.org/10.1073/pnas.1416212112

Kohsaka, H., Guertin, P. A., & Nose, A. (2017, 2017/5//). Neural Circuits Underlying Fly Larval Locomotion. *Current Pharmaceutical Design, 23*(12), 1722-1733. https://doi.org/10.2174/1381612822666161208120835

Kohsaka, H., Takasu, E., Morimoto, T., & Nose, A. (2014, 2014/11//). A Group of Segmental Premotor Interneurons Regulates the Speed of Axial Locomotion in Drosophila Larvae. *Current Biology, 24*(22), 2632-2642. https://doi.org/10.1016/j.cub.2014.09.026

Kohsaka, H., Zwart, M. F., Fushiki, A., Fetter, R. D., Truman, J. W., Cardona, A., & Nose, A. (2019, 2019/06/14). Regulation of forward and backward locomotion through intersegmental feedback circuits in Drosophila larvae. *Nature Communications, 10*(1), 2654. https://doi.org/10.1038/s41467-019-10695-y

Larderet, I., Fritsch, P. M. J., Gendre, N., Larisa Neagu-Maier, G., Fetter, R. D., Schneider-Mizell, C. M., Truman, J. W., Zlatic, M., Cardona, A., & Sprecher, S. G. (2017).

Organization of the Drosophila larval visual circuit.
https://doi.org/10.7554/eLife.28387.001

Leung, N. Y., & Montell, C. (2017, Oct 6). Unconventional Roles of Opsins. *Annu Rev Cell Dev Biol, 33*, 241-264. https://doi.org/10.1146/annurev-cellbio-100616-060432

Liao, J. C., & Fetcho, J. R. (2008, Nov 26). Shared versus specialized glycinergic spinal interneurons in axial motor circuits of larval zebrafish. *J Neurosci, 28*(48), 12982-12992. https://doi.org/10.1523/JNEUROSCI.3330-08.2008

Liu, L., Yermolaieva, O., Johnson, W. A., Abboud, F. M., & Welsh, M. J. (2003, Mar). Identification and function of thermosensory neurons in Drosophila larvae. *Nat Neurosci, 6*(3), 267-273. https://doi.org/10.1038/nn1009

Liu, Y., Hasegawa, E., Nose, A., Zwart, M. F., & Kohsaka, H. (2023, 2023/08/08). Synchronous multi-segmental activity between metachronal waves controls locomotion speed in Drosophila larvae. *eLife, 12*, e83328. https://doi.org/10.7554/eLife.83328

Lopez-Bellido, R., Himmel, N. J., Gutstein, H. B., Cox, D. N., & Galko, M. J. (2019, Nov 11). An assay for chemical nociception in Drosophila larvae. *Philos Trans R Soc Lond B Biol Sci, 374*(1785), 20190282. https://doi.org/10.1098/rstb.2019.0282

Macpherson, L. J., Zaharieva, E. E., Kearney, P. J., Alpert, M. H., Lin, T.-Y., Turan, Z., Lee, C.-H., & Gallio, M. (2015, 2015/12//). Dynamic labelling of neural connections in multiple colours by trans-synaptic fluorescence complementation. *Nature Communications, 6*(1), 10024-10024. https://doi.org/10.1038/ncomms10024

Maksymchuk, N., Sakurai, A., Cox, D. N., & Cymbalyuk, G. (2022). Transient and Steady-State Properties of Drosophila Sensory Neurons Coding Noxious Cold Temperature [Original Research]. *Frontiers in Cellular Neuroscience, 16*. https://www.frontiersin.org/articles/10.3389/fncel.2022.831803

Maksymchuk, N., Sakurai, A., Cox, D. N., & Cymbalyuk, G. S. (2023, Sep 27). Cold-Temperature Coding with Bursting and Spiking Based on TRP Channel Dynamics in Drosophila Larva Sensory Neurons. *Int J Mol Sci, 24*(19). https://doi.org/10.3390/ijms241914638

Mark, B., Lai, S. L., Zarin, A. A., Manning, L., Pollington, H. Q., Litwin-Kumar, A., Cardona, A., Truman, J. W., & Doe, C. Q. (2021, May 11). A developmental framework linking neurogenesis and circuit formation in the Drosophila CNS. *eLife, 10*. https://doi.org/10.7554/eLife.67510

Masson, J. B., Laurent, F., Cardona, A., Barre, C., Skatchkovsky, N., Zlatic, M., & Jovanic, T. (2020, Feb). Identifying neural substrates of competitive interactions and sequence transitions during mechanosensory responses in Drosophila. *PLoS Genet, 16*(2), e1008589. https://doi.org/10.1371/journal.pgen.1008589

2266

2267 Miroschnikow, A., Schlegel, P., Schoofs, A., Hueckesfeld, S., Li, F., Schneider-Mizell, C. M.,
2268   Fetter, R. D., Truman, J., Cardona, A., & Pankratz, M. J. (2018, 2018/7//). The Feeding
2269   Connectome: Convergence of Monosynaptic and Polysynaptic Sensory Paths onto
2270   Common Motor Outputs. *bioRxiv*, 368878-368878. https://doi.org/10.1101/368878

2271

2272 Moehring, F., Cowie, A. M., Menzel, A. D., Weyer, A. D., Grzybowski, M., Arzua, T., Geurts,
2273   A. M., Palygin, O., & Stucky, C. L. (2018, Jan 16). Keratinocytes mediate innocuous and
2274   noxious touch via ATP-P2X4 signaling. *eLife, 7*. https://doi.org/10.7554/eLife.31684

2275

2276 Montell, C. (2021, Mar 3). Drosophila sensory receptors-a set of molecular Swiss Army Knives.
2277   *Genetics, 217*(1), 1-34. https://doi.org/10.1093/genetics/iyaa011

2278

2279 Nakamura, K. (2018). Afferent pathways for autonomic and shivering thermoeffectors. *Handb*
2280   *Clin Neurol, 156*, 263-279. https://doi.org/10.1016/B978-0-444-63912-7.00016-3

2281

2282 Ohyama, T., Jovanic, T., Denisov, G., Dang, T. C., Hoffmann, D., Kerr, R. A., & Zlatic, M.
2283   (2013, 2013/8//). High-Throughput Analysis of Stimulus-Evoked Behaviors in
2284   Drosophila Larva Reveals Multiple Modality-Specific Escape Strategies. *PLoS One, 8*(8),
2285   e71706-e71706. https://doi.org/10.1371/journal.pone.0071706

2286

2287 Ohyama, T., Schneider-Mizell, C. M., Fetter, R. D., Aleman, J. V., Franconville, R., Rivera-
2288   Alba, M., Mensh, B. D., Branson, K. M., Simpson, J. H., Truman, J. W., Cardona, A., &
2289   Zlatic, M. (2015, 2015/4//). A multilevel multimodal circuit enhances action selection in
2290   Drosophila. *Nature, 520*(7549), 633-639. https://doi.org/10.1038/nature14297

2291

2292 Ormerod, K. A.-O., Scibelli, A. E., & Littleton, J. A.-O. (2022). Regulation of excitation-
2293   contraction coupling at the Drosophila neuromuscular junction. *The Journal of*
2294   *Physiology, 600 (2)*(349–372). https://doi.org/https://doi.org/10.1113/JP282092

2295

2296 Patel, A., & Cox, D. (2017). Behavioral and Functional Assays for Investigating Mechanisms of
2297   Noxious Cold Detection and Multimodal Sensory Processing in Drosophila Larvae. *BIO-*
2298   *PROTOCOL, 7*(13). https://doi.org/10.21769/BioProtoc.2388

2299

2300 Patel, A. A., Sakurai, A., Himmel, N. J., & Cox, D. N. (2022). Modality specific roles for
2301   metabotropic GABAergic signaling and calcium induced calcium release mechanisms in
2302   regulating cold nociception. *bioRxiv*, 2022.2005.2012.491647.
2303   https://doi.org/10.1101/2022.05.12.491647

2304

2305 Pfeiffer, B. D., Jenett, A., Hammonds, A. S., Ngo, T. T., Misra, S., Murphy, C., Scully, A.,
2306   Carlson, J. W., Wan, K. H., Laverty, T. R., Mungall, C., Svirskas, R., Kadonaga, J. T.,
2307   Doe, C. Q., Eisen, M. B., Celniker, S. E., & Rubin, G. M. (2008, Jul 15). Tools for
2308   neuroanatomy and neurogenetics in Drosophila. *Proc Natl Acad Sci U S A, 105*(28),
2309   9715-9720. https://doi.org/10.1073/pnas.0803697105

2310

Robertson, J. L., Tsubouchi, A., & Tracey, W. D. (2013). Larval defense against attack from parasitoid wasps requires nociceptive neurons. *PLoS One, 8*(10), e78704-e78704. https://doi.org/10.1371/journal.pone.0078704

Saalfeld, S., Cardona, A., Hartenstein, V., & Tomancak, P. (2009, Aug 1). CATMAID: collaborative annotation toolkit for massive amounts of image data. *Bioinformatics, 25*(15), 1984-1986. https://doi.org/10.1093/bioinformatics/btp266

Salvaterra, P. M., & Kitamoto, T. (2001, Aug). Drosophila cholinergic neurons and processes visualized with Gal4/UAS-GFP. *Brain Res Gene Expr Patterns, 1*(1), 73-82. https://doi.org/10.1016/s1567-133x(01)00011-4

Saumweber, T., Rohwedder, A., Schleyer, M., Eichler, K., Chen, Y.-c., Aso, Y., Cardona, A., Eschbach, C., Kobler, O., Voigt, A., Durairaja, A., Mancini, N., Zlatic, M., Truman, J. W., Thum, A. S., & Gerber, B. (2018, 2018/12//). Functional architecture of reward learning in mushroom body extrinsic neurons of larval Drosophila. *Nature Communications, 9*(1), 1104-1104. https://doi.org/10.1038/s41467-018-03130-1

Scheffer, L. K., Xu, C. S., Januszewski, M., Lu, Z., Takemura, S. Y., Hayworth, K. J., Huang, G. B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., Clements, J., Hubbard, P. M., Katz, W. T., Umayam, L., Zhao, T., Ackerman, D., Blakely, T., Bogovic, J., Dolafi, T., Kainmueller, D., Kawase, T., Khairy, K. A., Leavitt, L., Li, P. H., Lindsey, L., Neubarth, N., Olbris, D. J., Otsuna, H., Trautman, E. T., Ito, M., Bates, A. S., Goldammer, J., Wolff, T., Svirskas, R., Schlegel, P., Neace, E., Knecht, C. J., Alvarado, C. X., Bailey, D. A., Ballinger, S., Borycz, J. A., Canino, B. S., Cheatham, N., Cook, M., Dreher, M., Duclos, O., Eubanks, B., Fairbanks, K., Finley, S., Forknall, N., Francis, A., Hopkins, G. P., Joyce, E. M., Kim, S., Kirk, N. A., Kovalyak, J., Lauchie, S. A., Lohff, A., Maldonado, C., Manley, E. A., McLin, S., Mooney, C., Ndama, M., Ogundeyi, O., Okeoma, N., Ordish, C., Padilla, N., Patrick, C. M., Paterson, T., Phillips, E. E., Phillips, E. M., Rampally, N., Ribeiro, C., Robertson, M. K., Rymer, J. T., Ryan, S. M., Sammons, M., Scott, A. K., Scott, A. L., Shinomiya, A., Smith, C., Smith, K., Smith, N. L., Sobeski, M. A., Suleiman, A., Swift, J., Takemura, S., Talebi, I., Tarnogorska, D., Tenshaw, E., Tokhi, T., Walsh, J. J., Yang, T., Horne, J. A., Li, F., Parekh, R., Rivlin, P. K., Jayaraman, V., Costa, M., Jefferis, G. S., Ito, K., Saalfeld, S., George, R., Meinertzhagen, I. A., Rubin, G. M., Hess, H. F., Jain, V., & Plaza, S. M. (2020, Sep 7). A connectome and analysis of the adult Drosophila central brain. *eLife, 9*. https://doi.org/10.7554/eLife.57443

Schlegel, P., Texada, M. J., Miroschnikow, A., Schoofs, A., Hückesfeld, S., Peters, M., Schneider-Mizell, C. M., Lacin, H., Li, F., Fetter, R. D., Truman, J. W., Cardona, A., & Pankratz, M. J. (2016, 2016/11//). Synaptic transmission parallels neuromodulation in a central food-intake circuit. *eLife, 5*. https://doi.org/10.7554/eLife.16799

Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., Champion, A., Midgley, F. M., Fetter, R. D., Saalfeld, S., & Cardona, A. (2016,

2356   2016/3//). Quantitative neuroanatomy for connectomics in Drosophila. *eLife, 5*.
2357   https://doi.org/10.7554/eLife.12059
2358
2359 Scholz, N., Gehring, J., Guan, C., Ljaschenko, D., Fischer, R., Lakshmanan, V., Kittel, R. J., &
2360   Langenhan, T. (2015, 2015/05/12/). The Adhesion GPCR Latrophilin/CIRL Shapes
2361   Mechanosensation. *Cell Reports, 11*(6), 866-874.
2362   https://doi.org/https://doi.org/10.1016/j.celrep.2015.04.008
2363
2364 Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N.,
2365   Schwikowski, B., & Ideker, T. (2003, Nov). Cytoscape: a software environment for
2366   integrated models of biomolecular interaction networks. *Genome Res, 13*(11), 2498-2504.
2367   https://doi.org/10.1101/gr.1239303
2368
2369 Sweeney, S. T., Broadie, K., Keane, J., Niemann, H., & O'Kane, C. J. (1995, Feb). Targeted
2370   expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic
2371   transmission and causes behavioral defects. *Neuron, 14*(2), 341-351.
2372   https://doi.org/10.1016/0896-6273(95)90290-2
2373
2374 Takagi, S., Cocanougher, B. T., Niki, S., Miyamoto, D., Kohsaka, H., Kazama, H., Fetter, R. D.,
2375   Truman, J. W., Zlatic, M., Cardona, A., & Nose, A. (2017, 2017/12//). Divergent
2376   Connectivity of Homologous Command-like Neurons Mediates Segment-Specific Touch
2377   Responses in Drosophila. *Neuron, 96*(6), 1373-1387.e1376.
2378   https://doi.org/10.1016/j.neuron.2017.10.030
2379
2380 Talpalar, A. E., Bouvier, J., Borgius, L., Fortin, G., Pierani, A., & Kiehn, O. (2013, Aug 1).
2381   Dual-mode operation of neuronal networks involved in left-right alternation. *Nature,
2382   500*(7460), 85-88. https://doi.org/10.1038/nature12286
2383
2384 Tastekin, I., Khandelwal, A., Tadres, D., Fessner, N. D., Truman, J. W., Zlatic, M., Cardona, A.,
2385   & Louis, M. (2018, Nov 22). Sensorimotor pathway controlling stopping behavior during
2386   chemotaxis in the Drosophila melanogaster larva. *eLife, 7*.
2387   https://doi.org/10.7554/eLife.38740
2388
2389 Tenedini, F. M., Sáez González, M., Hu, C., Pedersen, L. H., Petruzzi, M. M., Spitzweck, B.,
2390   Wang, D., Richter, M., Petersen, M., Szpotowicz, E., Schweizer, M., Sigrist, S. J.,
2391   Calderon de Anda, F., & Soba, P. (2019, 2019/08/05). Maintenance of cell type-specific
2392   connectivity and circuit function requires Tao kinase. *Nature Communications, 10*(1),
2393   3506. https://doi.org/10.1038/s41467-019-11408-1
2394
2395 Tracey, W. D., Wilson, R. I., Laurent, G., & Benzer, S. (2003, 2003/4//). painless, a Drosophila
2396   gene essential for nociception. *Cell, 113*(2), 261-273.
2397   http://www.ncbi.nlm.nih.gov/pubmed/12705873
2398
2399 Tsubouchi, A., Caldwell, J. C., & Tracey, W. D. (2012, 2012/11//). Dendritic filopodia, Ripped
2400   Pocket, NOMPC, and NMDARs contribute to the sense of touch in Drosophila larvae.
2401   *Current biology : CB, 22*(22), 2124-2134. https://doi.org/10.1016/j.cub.2012.09.019

Turner, H. N., Armengol, K., Patel, A. A., Himmel, N. J., Sullivan, L., Iyer, S. C., Bhattacharya, S., Iyer, E. P. R., Landry, C., Galko, M. J., & Cox, D. N. (2016, Dec 5). The TRP Channels Pkd2, NompC, and Trpm Act in Cold-Sensing Neurons to Mediate Unique Aversive Behaviors to Noxious Cold in Drosophila. *Curr Biol, 26*(23), 3116-3128. https://doi.org/10.1016/j.cub.2016.09.038

Turner, H. N., Patel, A. A., Cox, D. N., & Galko, M. J. (2018). Injury-induced cold sensitization in Drosophila larvae involves behavioral shifts that require the TRP channel Brv1. *PLoS One, 13*(12), e0209577. https://doi.org/10.1371/journal.pone.0209577

Vaadia, R., Li, W., Voleti, V., Singhania, A., Hillman, E. M. C., & Grueber, W. B. (2018, 2018/11//). Characterization of proprioceptive system dynamics in behaving Drosophila larvae using high-speed volumetric microscopy. *bioRxiv*, 467274-467274. https://doi.org/10.1101/467274

Valdes-Aleman, J., Fetter, R. D., Sales, E. C., Heckman, E. L., Venkatasubramanian, L., Doe, C. Q., Landgraf, M., Cardona, A., & Zlatic, M. (2021). Comparative Connectomics Reveals How Partner Identity, Location, and Activity Specify Synaptic Connectivity in Drosophila. *Neuron, 109*(1), 105-122.e107. https://doi.org/10.1016/j.neuron.2020.10.004

Vogelstein, J. T., Park, Y., Ohyama, T., Kerr, R. A., Truman, J. W., Priebe, C. E., & Zlatic, M. (2014, 2014/4//). Discovery of Brainwide Neural-Behavioral Maps via Multiscale Unsupervised Structure Learning. *Science, 344*(6182), 386-392. https://doi.org/10.1126/science.1250298

Ward, L., Wright, E., & McMahon, S. B. (1996, Jan). A comparison of the effects of noxious and innocuous counterstimuli on experimentally induced itch and pain. *Pain, 64*(1), 129-138. https://doi.org/10.1016/0304-3959(95)00080-1

Winding, M., Pedigo, B. D., Barnes, C. L., Patsolic, H. G., Park, Y., Kazimiers, T., Fushiki, A., Andrade, I. V., Khandelwal, A., Valdes-Aleman, J., Li, F., Randel, N., Barsotti, E., Correia, A., Fetter, R. D., Hartenstein, V., Priebe, C. E., Vogelstein, J. T., Cardona, A., & Zlatic, M. (2023, Mar 10). The connectome of an insect brain. *Science, 379*(6636), eadd9330. https://doi.org/10.1126/science.add9330

Wu, P., Arris, D., Grayson, M., Hung, C. N., & Ruparel, S. (2018). Characterization of sensory neuronal subtypes innervating mouse tongue. *PLoS One, 13*(11), e0207069. https://doi.org/10.1371/journal.pone.0207069

Xiao, R., & Xu, X. Z. S. (2021, Feb 10). Temperature Sensation: From Molecular Thermosensors to Neural Circuits and Coding Principles. *Annu Rev Physiol, 83*, 205-230. https://doi.org/10.1146/annurev-physiol-031220-095215

2447 Yan, Z., Zhang, W., He, Y., Gorczyca, D., Xiang, Y., Cheng, L. E., Meltzer, S., Jan, L. Y., &
2448   Jan, Y. N. (2012, 2012/12//). Drosophila NOMPC is a mechanotransduction channel
2449   subunit for gentle-touch sensation. *Nature, 493*(7431), 221-225.
2450   https://doi.org/10.1038/nature11685
2451
2452 Yoshino, J., Morikawa, R. K., Hasegawa, E., & Emoto, K. (2017, 2017/8//). Neural Circuitry that
2453   Evokes Escape Behavior upon Activation of Nociceptive Sensory Neurons in Drosophila
2454   Larvae. *Current Biology, 27*(16), 2499-2504.e2493.
2455   https://doi.org/10.1016/j.cub.2017.06.068
2456
2457 Zarin, A. A., Mark, B., Cardona, A., Litwin-Kumar, A., & Doe, C. Q. (2019, Dec 23). A
2458   multilayer circuit architecture for the generation of distinct locomotor behaviors in
2459   Drosophila. *eLife, 8*. https://doi.org/10.7554/eLife.51781
2460
Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., Torrens, O.,
Price, J., Fisher, C. B., Sharifi, N., Calle-Schuler, S. A., Kmecova, L., Ali, I. J., Karsh, B.,
Trautman, E. T., Bogovic, J. A., Hanslovsky, P., Jefferis, G., Kazhdan, M., Khairy, K.,
Saalfeld, S., Fetter, R. D., & Bock, D. D. (2018, Jul 26). A Complete Electron
Microscopy Volume of the Brain of Adult Drosophila melanogaster. *Cell, 174*(3), 730-
743 e722. https://doi.org/10.1016/j.cell.2018.06.019
2467
2468 Zhu, J., Boivin, J. C., Garner, A., Ning, J., Zhao, Y. Q., & Ohyama, T. (2024, Aug 28). Feedback
2469   inhibition by a descending GABAergic neuron regulates timing of escape behavior in
2470   Drosophila larvae. *eLife, 13*. https://doi.org/10.7554/eLife.93978
2471
2472 Zwart, M. F., Pulver, S. R., Truman, J. W., Fushiki, A., Fetter, R. D., Cardona, A., & Landgraf,
2473   M. (2016, 2016/8//). Selective Inhibition Mediates the Sequential Recruitment of Motor
2474   Pools. *Neuron, 91*(4), 944-944. https://doi.org/10.1016/j.neuron.2016.07.041
2475
2476