1 **Synergistic effects of the next generation insecticide flupyradifurone with a fungal**
2 **pathogen**
3

4 Daniel Schläppi[1*], Adam Al-Hashemi[1], Vaneeza Wasif[1], Florent Masson[1], Nathalie Stroeymeyt[1]

5 [1] School of Biological Sciences, University of Bristol, Life Sciences Building, 24 Tyndall Avenue, BS8 1TQ
6 Bristol, UK

7 *Correspondence:* d.schlaeppi@bristol.ac.uk

8
9

Summary

The global insect decline is largely driven by agricultural pesticides, whose detrimental effects are

further amplified by synergistic interactions with other stressors. Whilst there is increasing evidence of

adverse impacts of the next-generation insecticide flupyradifurone (FPF), initially marketed as safe for

pollinators, its effects on non-pollinators and potential interactions with other stressors remain poorly

understood. Here we investigate the impacts of chronic exposure to FPF alone and in combination with

the entomopathogenic fungus *Metarhizium brunneum* in a non-target arthropod: the ant *Lasius niger*.

Our results show that FPF concentrations upwards of 100 ppm increased worker mortality. Furthermore,

while sublethal ($\leq$ 50 ppm) FPF exposure alone did not affect survival, it increased susceptibility to

fungal infection, indicating a synergy between the two stressors. This new report of a synergistic

interaction between FPF and pathogens raises further concerns about the impact of novel pesticides and

could have important implications for insect conservation.

Keywords: ants (Hymenoptera: Formicidae), chronic exposure, field realistic*, Metarhizium*, pesticide,

stressor interaction, sublethal effects

Introduction

Insects are important indicators for ecosystem health and provide vital ecosystem services such as pollination, pest control, and nutrient cycling[1,2]. However, global insect biodiversity and abundance are being lost at alarming rates[3,4]. Recent research identified agrochemicals such as neonicotinoid insecticides as a major driver of these declines[5,6]. The detrimental impacts of neonicotinoids on non-target organisms[7-10] led to restrictions in the use of clothianidin, thiamethoxam, and imidacloprid in the European Union. Following the neonicotinoid ban, new pesticides were introduced to the market, including the butenolide insecticide flupyradifurone (FPF). Although structurally different, neonicotinoids and FPF target the same receptor and both disrupt the central nervous system of insects, making them highly effective against a broad range of pests[11,12]. Upon introduction, FPF was advertised as "bee safe" because of its ecotoxicological profile. However, recent evidence has uncovered that FPF does pose a risk to pollinators, as exposure to environmentally relevant doses of FPF leads to detrimental sublethal effects, including impaired cognitive abilities, motion coordination deficits, hyperactivity and apathy, as well as impacts on flight and foraging behaviours[13-17]. Such sublethal impacts frequently elude detection in risk assessments which are typically based on acute toxicity assays.

A major concern associated with the use of agrochemicals are their interactions with other stressors, as synergistic interactions may amplify their environmental significance. Traditional agrochemicals have been demonstrated to interact synergistically with a variety of stressors such as other pesticides, pathogens, malnutrition, or climate change, leading to increased host mortality[18-22]. By contrast, there have been conflicting reports regarding interactions between FPF and other pesticides[23-25], and so far, there is no evidence for an interaction between FPF and disease[26,27]. Furthermore, the great majority of studies examining the direct and indirect impacts of FPF have focused on bees, whilst other non-target insects such as ants have been overlooked, in spite of their ubiquity and the essential ecosystem services they provide[28]. Here we aimed to address these gaps by (i) investigating the direct effects of FPF exposure on ant survival and (ii) testing for potential indirect effects via interaction with disease susceptibility. To do so, we first determined the susceptibility of the black garden ant *Lasius niger* to exposure to a range of FPF concentrations. We then used exposure to field-realistic, sublethal

FPF doses in combination with the fungal pathogen *Metarhizium brunneum* to investigate potential

interactive effects between the two stressors.

Methods

L. niger colonies were initiated using newly mated queens collected in Berlin (July 2021). The

colonies were raised at 25°C and 65% humidity, with a 12h day/night cycle, *ad libitum* supply of water

and honey water (15% mass fraction of honey) and weekly provision of *Drosophila hydei*.

Flupyradifurone susceptibility test

We sampled 120 workers from each of four stock colonies and randomly split them into eight

subsets of fifteen workers (N=480 ants across 32 subsets). Each subset was kept in a separate Petri-dish

($\varnothing$ = 50 mm) with fluon-coated walls and sealed with parafilm. The floor of each Petri-dish was covered

with a thin layer of plaster of Paris to which 1 mL of water was added to maintain humidity. After one

day of acclimatisation, each colony's subsets were pseudo-randomly allocated to the control treatment

or one of seven FPF concentration treatments. Although field-realistic exposure doses remain to be

determined for ants[9,29], data provided by the Environmental Protection Agency (EPA) showed residues

up to 36 ppm in flowers and up to 4.3 ppm in nectar[30]. Hence, we considered 5 ppm to be a conservative

estimate for field-realistic dose and designed our experimental concentration range around that value.

Fresh FPF feeding solutions were prepared on each feeding day by mixing different proportions of a

30% honey water solution, a 1000 ppm FPF stock solution and MilliQ water to produce 8 different

solutions with final concentrations of 15% honey water each and 0 (control), 0.5, 1, 5, 10, 50, 100 or

500 ppm FPF (Sigma-Aldrich PESTANAL® analytical standard, 99.5% purity). At the start of the

experiment and every 3 days thereafter, each subset received a fresh FPF feeding solution provided in

a 200 μL Eppendorf tube closed with cotton wool. Survival was checked daily for two weeks by

counting and removing dead ants. All survival checks were conducted blind to treatment.

Flupyradifurone and fungus synergy test

To test for potential interactive effects between FPF and pathogens, ants were chronically exposed

to sublethal FPF doses for ten days, and then subsequently challenged with a fungal pathogen. We used

the generalist entomopathogenic fungus *Metarhizium brunneum* (strain MA275, KVL 03-143).

M. brunneum is common in soil environments and naturally infects *L. niger*, which display a variety of

physiological and behavioural defences against it[31-33]. A preliminary test was conducted to confirm that

FPF does not directly interfere with fungal germination (supplementary Fig. 1).

We sampled 132 workers from each of five stock colonies and distributed them evenly across 3

separate Petri-dishes, resulting in a total of 15 Petri-dishes containing 44 ants each (total ants N = 660).

The three Petri-dishes from each colony were pseudo-randomly assigned to one of three FPF treatment

groups (0, 5 and 50 ppm of FPF, that is, control, low, and high sublethal doses as determined in

experiment 1). Feeding solutions were freshly prepared and provided every 3 days as described above.

After ten days, 40 workers from each Petri dish were randomly selected and split into two groups of 20,

which were randomly allocated to two pathogen treatments (fungus-exposed and sham-exposed). This

resulted in a full factorial design involving six treatment groups (three FPF treatments $\times$ two pathogen

treatments) with 5 replicates each (one per stock colony). Accordingly, the experiment was organised

into 5 successive experimental blocks, each involving a single stock colony and one replicate for every

treatment. In each block, all sham- and fungal exposures were performed in a single session lasting less

than two hours. The 20 ants of each Petri-dish were exposed consecutively, while the order in which

the Petri-dishes for all treatments were exposed was varied pseudo-randomly across all blocks to

mitigate temporal effects.

Fungal and sham exposures were performed using standard procedures[33]: *M. brunneum* was

cultured on Sabouraud-Dextrose Agar (SDA) plates at 24°C until sporulation. Conidiospores were

harvested in 0.05% Triton-X100, washed twice and adjusted to 10^9 spores/mL in 0.05% Triton-X100

to create a spore stock suspension. The stock was diluted by half with 0.05% Triton-X100 resulting in

a fungal suspension with a final spore concentration of $5x10^8$ corresponding to the median lethal dose

LD50 (unpublished data). Ants were placed on ice, then exposed to 0.3 μL of the treatment solution

(fungal suspension or sham solution of 0.05% Triton X-100 only) pipetted onto their gaster. They were

then left to dry on filter paper for 1 minute and subsequently transferred individually into individual

Petri-dishes ($\varnothing$ = 35 mm, N = 600) with fluon coated walls and a water tube (200 μL Eppendorf tubes

closed with cotton wool). These Petri-dishes were then sealed with parafilm and placed in the incubator

for two weeks with daily survival checks conducted blind to treatment.

Statistical Analyses

All statistical analyses were performed in using R v4.2.2[34]. Survival analyses were performed using

Cox proportional hazards mixed-effect models using R packages *survival*[35] and *coxme*[36]. For the FPF

susceptibility test treatment group was used as a categorical fixed effect and Petri-dish and stock colony

as random effects. For the FPF and fungus synergy test, fungal exposure group, FPF concentration and

their one-way interaction were used as categorical fixed effects and Petri-dish, stock colony and

exposure block as random effects. Fixed effect significance was tested using two-tailed Wald chi-square

tests. *Post-hoc* contrasts, defined using package *multcomp*[37], were used for pairwise comparisons of

experimental groups. P-values were corrected for multiple comparisons using the Benjamini-Hochberg

(BH) method.

Results

Flupyradifurone susceptibility test

FPF concentration had a significant effect on ant mortality (Fig. 1; Cox proportional hazards mixed

effect model, effect of FPF concentration: $\chi^2 = 31.67$, df = 7, p < 0.001). *Post-hoc* pairwise comparisons

revealed that concentrations of 100 ppm and above led to a significant increase in ant mortality

compared to the control (100 ppm vs. 0 ppm: z = 2.81, p=0.017; 500 ppm vs. 0 ppm: z = 4.97, p <

0.001). By contrast, concentrations of 50 ppm and below did not increase mortality compared to the

control (z < 2.08; p > 0.09 in all pairwise comparisons); these were used as sublethal concentrations in

experiment 2. The absence of lethal effects in concentrations of 50 ppm and below cannot be attributed

to the ants restricting their food intake to limit their exposure to the pesticide, as feeding volumes were

only found to decrease at the highest FPF concentration tested (500 ppm, see supplementary Fig. 2).

Flupyradifurone and fungus synergy test

Challenge with the pathogenic fungus *M. brunneum* increased ant mortality in all three FPF

exposure conditions (Fig. 2a; Cox proportional hazards mixed effect model, effect of fungal exposure:

$\chi^2 = 57.49$, df = 1, p < 0.001; pairwise comparisons of fungus-treated vs. sham-treated ants with BH

correction, no FPF: Hazard Ratio (HR) = 1.69, z = 2.36, p = 0.028; FPF 5 ppm: HR = 3.56, z = 5.04, p

< 0.001; FPF 50 ppm: HR = 3.96, z = 5.94, p < 0.001). Furthermore, although FPF alone did not affect

ant survival ($\chi^2 = 3.84$, df = 2, p = 0.15), there was a significant interaction between FPF treatment and

fungal exposure ($\chi^2 = 8.29$, df = 2, p = 0.016). More specifically, prior sublethal exposure to FPF

significantly increased the fungus-induced mortality compared to the control (Fig. 2b; pairwise

comparisons of Hazard Ratios with BH correction, control vs. FPF 5ppm: z = 2.22, p = 0.032; control

vs. FPF 50ppm: z = 2.65, p = 0.016; FPF 5ppm vs. 50ppm: z = 0.31, p = 0.76). This indicates that

infections with *M. brunneum* are even more likely to be lethal if the ants have previously been

chronically exposed to a sublethal FPF dose (Fig. 2b).

Discussion

This study assessed the direct effect of chronic FPF exposure on the survival of black garden ants

(*L. niger*), revealing increased mortality at concentrations of 100 ppm and above. Furthermore, we

demonstrated an indirect effect of FPF on ants at lower, sublethal concentrations ($\leq$ 50 ppm)*,* as the

challenge with *M. brunneum* was more likely to be lethal following chronic exposure to sublethal FPF

doses, indicating a synergistic interaction between the two stressors. Overall, the results of this study

raise further concerns about the long-term impacts of novel pesticides on the health of insects.

While chronic exposure to FPF concentrations upwards of 100 ppm resulted in increased worker

mortality, concentrations of 50 ppm or lower showed comparable mortality rates to the control group

and were thus classified as sublethal. However, the direct impact of FPF on ants in the field remains

unclear as the exact exposure of ants to FPF in their natural environment has yet to be quantified.

Residues in pollen and nectar are considered to be the main exposure routes for bees and their reported

residue values are as high as 68 ppm for pollen and 21.8 ppm for extrafloral nectaries[38]. However,

studies of honey bees typically use 4 ppm as field-realistic doses because nectar collected by returning

foragers contained 4.3 ppm of FPF[16,30]. Our sublethal concentrations for ants (0.5 - 50 ppm) cover a

comparable range, suggesting that FPF residue values in the field may not increase ant mortality via

direct toxicity. Nonetheless, future studies should quantify true field-realistic exposure for ant colonies

over the course of a year, based on a combination of all relevant acute and chronic exposure routes,

which includes contact with spray droplets, contaminated soils, foliage, and water, as well as foodborne

exposure[9,29,39].

At sublethal concentrations, FPF exposure alone did not affect ant survival, but it increased

mortality when combined with *M. brunneum* exposure, indicating that FPF may induce indirect

sublethal effects in non-target hosts at field-realistic concentrations via a synergistic interaction with

pathogens. This contrasts with previous studies in bees, which did not detect FPF-induced changes in

survival rates to fungal or viral pathogens[26,27,40]. However, our findings are consistent with findings

from traditional neonicotinoids in ants, as imidacloprid was shown to increase the susceptibility of ants

to infection by the fungus *Beauveria bassiana*[41].

The synergistic effect between FPF and *M. brunneum* suggests that FPF may affect the behavioural

or biochemical immune response of the ants. However, as we did not record the ants' behaviour or

measure their immune function, we cannot identify the underlying mechanism for this interaction. On

one hand, FPF could cause a direct immune suppression or trigger a resource allocation trade-off

between immune functions and detoxification[42], as has been shown in honey bees exposed to

neonicotinoids[18,43,44]. On the other hand, as FPF disrupts the function of the central nervous system, it

could also interfere with individual and collective behaviours that mitigate the risk of fungal infection

in ants, such as self-grooming, allo-grooming or chemical disinfection[31]. Future studies involving entire

colonies will be required to determine whether FPF impairs colony-wide social immunity mechanisms

that protect colonies against epidemics and how individual level effects impact the colony as a whole[45-

47].

Overall, this study contributes to the understanding of stressor interactions by providing evidence

for a synergistic effect between FPF and pathogens in a non-target organism, and it highlights the

ecotoxicological risk posed by novel insecticides. Our findings are particularly concerning as they

suggest that FPF may pose a comparable threat to beneficial insects as banned neonicotinoids, despite

its initial perception of relative safety for bees. This stresses the importance of considering sublethal,

long-term effects as well as interactive effects between multiple stressors when assessing the risks of

agrochemicals, rather than only evaluating direct mortality at low doses. Furthermore, our findings

revealed a synergistic interaction between FPF and pathogens in ants, which contrasts with the absence

of such an effect in previous studies on honey bees. This shows that the use of bees as surrogates for

other non-target organisms may be inadequate when evaluating the risk of agrochemicals, and suggests

that other economically and ecologically important arthropods such as ants should be included

alongside bees as representative model organisms.

Limitations of the study

This study demonstrates the synergistic effects of FPF and fungal pathogens in a controlled

laboratory setting, which may not account for the full range of environmental factors that ants encounter

in natural settings. Additionally, the study focuses on *L. niger* ants and one exemplary fungal model

pathogen, hence generalizations to other species should be made with caution. The absence of

behavioural and immunological assessments leaves the underlying mechanisms behind the observed

synergy speculative and further research is required to explore these aspects.

Acknowledgements

199   We would like to thank Prof Nicolai Vitt Meyling from University of Copenhagen for providing

200   *M. brunneum*. Financial support was provided to DS by the Swiss National Science Foundation

201   (P2BEP3_195575 & P500PB_206883) and to N.S. by the European Research Council (ERC Starting

202   Grant 'DISEASE', no. 802628).

Author contributions

204   D.S. conceived the study and designed it with N.S. and F.M; D.S., V.W., and A. AH. conducted

205   the research and performed the experiments with inputs from N.S. & F.M.; N.S. provided laboratory

206   space and materials; D.S., V.W., and A. AH. analysed the data with inputs from N.S.; D.S. wrote the

207   manuscript based on a first draft by A. AH, with contributions from all authors. All authors edited and

208   approved the manuscript.

Declaration of interests

210   The authors declare no conflict of interest.

Data accessibility statement

The supplementary files, raw data and code supporting the results are archived on figshare and can be accessed via DOI: 10.6084/m9.figshare.25782330

Supplemental information titles and legends

Document S1. containing details on the antifungal assay including supplementary Fig. 1 and details on the quantification of food uptake and supplementary Fig. 2.

References

1. Elizalde, L., Arbetman, M., Arnan, X., Eggleton, P., Leal, I. R., Lescano, M. N., Saez A., Werenkraut, V., and Pirk, G. I. (2020). The ecosystem services provided by social insects: traits, management tools and knowledge gaps. *Biological Reviews*, *95*(5), 1418-1441. https://doi.org/10.1111/brv.12616

2. Losey, J. E., and Vaughan, M. (2006). The economic value of ecological services provided by insects. *Bioscience*, *56*(4), 311-323. https://doi.org/10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2

3. Cardoso, P., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C. S., Gaigher, R., Habel, J. C., Hallmann, C. A., et al. (2020). Scientists' warning to humanity on insect extinctions. *Biological Conservation*, *242*, 108426. https://doi.org/10.1016/j.biocon.2020.108426

4. Wagner, D. L. (2020). Insect declines in the anthropocene. *Annual Review of Entomology*, *65*, 457-480. https://doi.org/10.1146/annurev-ento-011019-025151

5. Goulson, D., Nicholls, E., Botías, C., and Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, *347*(6229), 1255957. https://doi.org/10.1126/science.1255957

6. Sánchez-Bayo, F., and Wyckhuys, K. A. (2019). Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation*, *232*, 8-27. https://doi.org/10.1016/j.biocon.2019.01.020

7. Desneux, N., Decourtye, A., and Delpuech, J.-M. (2007). The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology*, *52*, 81-106. https://doi.org/10.1146/annurev.ento.52.110405.091440

8. Pisa, L. W., Amaral-Rogers, V., Belzunces, L. P., Bonmatin, J.-M., Downs, C. A., Goulson, D., Kreutzweiser, D. P., Krupke, C., Liess, M., McField M., et al. (2015). Effects of neonicotinoids and fipronil on non-target invertebrates. *Environmental Science and Pollution Research*, *22*(1), 68-102. https://doi.org/10.1007/s11356-014-3471-x

9. Schläppi, D., Stroeymeyt, N., and Neumann, P. (2021). Unintentional effects of neonicotinoids in ants (Hymenoptera: Formicidae). *Myrmecological News*, *31*, 181-184. https://doi.org/10.25849/myrmecol.news_031:181

10. Stanley, J., and Preetha, G. (2016). *Pesticide toxicity to non-target organisms*. Springer. https://doi.org/10.1007/978-94-017-7752-0

11. Nauen, R., Jeschke, P., Velten, R., Beck, M. E., Ebbinghaus-Kintscher, U., Thielert, W., Wölfel, K., Haas, M., Kunz K., and Raupach, G. (2015). Flupyradifurone: a brief profile of a new butenolide insecticide. *Pest Management Science*, *71*(6), 850-862. https://doi.org/10.1002/ps.3932

12. Tomizawa, M., and Casida, J. E. (2005). Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annual Review of Pharmacology and Toxicology*, *45*, 247-268. https://doi.org/10.1146/annurev.pharmtox.45.120403.095930

13. Hesselbach, H., and Scheiner, R. (2019). The novel pesticide flupyradifurone (Sivanto) affects honeybee motor abilities. *Ecotoxicology*, *28*(3), 354-366. https://doi.org/10.1007/s10646-019-02028-y

14. Hesselbach, H., Seeger, J., Schilcher, F., Ankenbrand, M., and Scheiner, R. (2020). Chronic exposure to the pesticide flupyradifurone can lead to premature onset of foraging in honeybees *Apis mellifera*. *Journal of Applied Ecology*, *57*(3), 609-618. https://doi.org/10.1111/1365-2664.13555

15. Siviter, H., and Muth, F. (2020). Do novel insecticides pose a threat to beneficial insects? *Proceedings of the Royal Society B*, *287*(1935), 20201265. https://doi.org/10.1098/rspb.2020.1265

16. Siviter, H., and Muth, F. (2022). Exposure to the novel insecticide flupyradifurone impairs bumblebee feeding motivation, learning, and memory retention. *Environmental Pollution*, *307*, 119575. https://doi.org/10.1016/j.envpol.2022.119575

17. Tosi, S., Nieh, J. C., Brandt, A., Colli, M., Fourrier, J., Giffard, H., Hernández-López, J., Malagnini, V., Williams G. R., and Simon-Delso, N. (2021). Long-term field-realistic exposure to a next-generation pesticide, flupyradifurone, impairs honey bee behaviour and survival. *Communications Biology*, *4*(1), 805. https://doi.org/10.1038/s42003-021-02336-2

18. Doublet, V., Labarussias, M., de Miranda, J. R., Moritz, R. F., and Paxton, R. J. (2015). Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. *Environmental Microbiology*, *17*(4), 969-983. https://doi.org/10.1111/1462-2920.12426

19. Feldhaar, H., and Otti, O. (2020). Pollutants and their interaction with diseases of social Hymenoptera. *Insects*, *11*(3), 153. https://doi.org/10.3390/insects11030153

20. Kaunisto, S., Ferguson, L. V., and Sinclair, B. J. (2016). Can we predict the effects of multiple stressors on insects in a changing climate? *Current Opinion in Insect Science*, 17, 55-61. https://doi.org/10.1016/j.cois.2016.07.001

21. Siviter, H., Bailes, E. J., Martin, C. D., Oliver, T. R., Koricheva, J., Leadbeater, E., and Brown, M. J. (2021). Agrochemicals interact synergistically to increase bee mortality. *Nature*, *569*, pages 389–392. https://doi.org/10.1038/s41586-021-03787-7

22. Tosi, S., Nieh, J. C., Sgolastra, F., Cabbri, R., and Medrzycki, P. (2017). Neonicotinoid pesticides and nutritional stress synergistically reduce survival in honey bees. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1869), 20171711. https://doi.org/10.1098/rspb.2017.1711

23. Al Naggar, Y., Singavarapu, B., Paxton, R. J., and Wubet, T. (2022). Bees under interactive stressors: the novel insecticides flupyradifurone and sulfoxaflor along with the fungicide azoxystrobin disrupt the gut microbiota of honey bees and increase opportunistic bacterial pathogens. *Science of the Total Environment*, *849*, 157941. https://doi.org/10.1016/j.scitotenv.2022.15794

24. Bava, R., Lupia, C., Castagna, F., Ruga, S., Nucera, S., Carresi, C. et at. (2024). Interaction of flupyradifurone and deltamethrin, two pesticides commonly used for plant pest control, in honeybees. *Animals*, *14*(6), 851. https://doi.org/10.3390/ani14060851

25. Tosi, S., and Nieh, J. C. (2019). Lethal and sublethal synergistic effects of a new systemic pesticide, flupyradifurone (Sivanto®), on honeybees. *Proceedings of the Royal Society B*, *286*(1900), 20190433. https://doi.org/10.1098/rspb.2019.0433

26. Al Naggar, Y., and Baer, B. (2019). Consequences of a short time exposure to a sublethal dose of Flupyradifurone (Sivanto) pesticide early in life on survival and immunity in the honeybee (*Apis mellifera*). *Scientific Reports*, *9*(1), 19753. https://doi.org/10.1038/s41598-019-56224-1

27. Al Naggar, Y., and Paxton, R. J. (2021). The novel insecticides flupyradifurone and sulfoxaflor do not act synergistically with viral pathogens in reducing honey bee (*Apis mellifera*) survival but sulfoxaflor modulates host immunocompetence. *Microbial Biotechnology*, *14*(1), 227-240. https://doi.org/10.1111/1751-7915.13673

28. Del Toro, I., Ribbons, R. R., and Pelini, S. L. (2012). The little things that run the world revisited: a review of ant-mediated ecosystem services and disservices (Hymenoptera: Formicidae). *Myrmecological News*, *17*, 133-146.

29. Pohl, M., Hommen, U., Eilebrecht, S., Schäfers, C., Gadau, J., and Otto, M. (2024). Ants are no bees–Gaps in the assessment of relevant exposure routes to pesticides and plant incorporated protectants. *Environmental Chemistry and Ecotoxicology*, *6*, 71-80. https://doi.org/10.1016/j.enceco.2024.02.001

30. Carleton, J. (2014) *Environmental Fate and Ecological Risk Assessment for Foliar, Soil Drench, and Seed Treatment Uses of the New Insecticide Flupyradifurone* (BYI 02960). United States Environmental Protection Agency, Washington, DC, USA.

31. Cremer, S., Pull, C. D., and Fuerst, M. A. (2018). Social immunity: emergence and evolution of colony-level disease protection. *Annual Review of Entomology*, *63*, 105-123. https://doi.org/10.1146/annurev-ento-020117-043110

32. Schluns, H., and Crozier, R. H. (2009). Molecular and chemical immune defenses in ants (Hymenoptera:
Formicidae). *Myrmecological News*, *12*, 237-249.

33. Stroeymeyt, N., Grasse, A. V., Crespi, A., Mersch, D. P., Cremer, S., and Keller, L. (2018). Social
network plasticity decreases disease transmission in a eusocial insect. *Science*, *362*(6417), 941-945.
https://doi.org/10.1126/science.aat4793

34. R Core Team (2020). *R: A language and environment for statistical computing*. Version 3.6.3, R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

35. Therneau, T. (2022). A Package for Survival Analysis in R. Version 3.4-0. https://CRAN.R-
project.org/package=survival

36. Therneau, T. (2022). Coxme: Mixed Effects Cox Models. Version 2.2-18.1. https://CRAN.R-
project.org/package=coxme

37. Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous inference in general parametric
models. *Biometrical Journal*, *50*(3), 346–363. https://doi.org/10.1002/bimj.200810425

38. Glaberman, S., and White, K. (2014). Environmental fate and ecological risk assessment for foliar, soil
drench, and seed treatment uses of the new insecticide flupyradifurone (BYI 02960). *U.S. Environmental*
Protection Agency (EPA), Office of Pesticide Programs, Environmental Fate and Effects Division EFED

39. Schläppi, D., Kettler, N., Straub, L., Glauser, G., and Neumann, P. (2020). Long-term effects of
neonicotinoid insecticides on ants. *Communications Biology*, *3*(1), 335. https://doi.org/10.1038/s42003-
020-1066-2

40. Harwood, G. P., Prayugo, V., and Dolezal, A. G. (2022). Butenolide insecticide flupyradifurone affects
honey bee worker antiviral immunity and survival. *Frontiers in Insect Science*, *2*, 907555.
https://doi.org/10.3389/finsc.2022.907555

41. Santos, A. V., de Oliveira, B. L., and Samuels, R. I. (2007). Selection of entomopathogenic fungi for use
in combination with sub-lethal doses of imidacloprid: perspectives for the control of the leaf-cutting ant
Atta sexdens rubropilosa Forel (Hymenoptera: Formicidae). *Mycopathologia*, *163*(4), 233-240.
https://doi.org/10.1007/s11046-007-9009-8

42. Collison, E., Hird, H., Cresswell, J., and Tyler, C. (2016). Interactive effects of pesticide exposure and
pathogen infection on bee health–a critical analysis. *Biological Reviews*, *91*(4), 1006-1019.
https://doi.org/10.1111/brv.12206

43. Sánchez-Bayo, F., Goulson, D., Pennacchio, F., Nazzi, F., Goka, K., and Desneux, N. (2016). Are bee
diseases linked to pesticides?—A brief review. *Environment International*, *89*, 7-11.
https://doi.org/10.1016/j.envint.2016.01.009

44. T O'Neal, S., Anderson, T. D., and Wu-Smart, J. Y. (2018). Interactions between pesticides and pathogen
susceptibility in honey bees. *Current Opinion in Insect Science*, *26*, 57-62.
https://doi.org/10.1016/j.cois.2018.01.006

45. Crall, J. D., Switzer, C. M., Oppenheimer, R. L., Ford Versypt, A. N., Dey, B., Brown, A., Eyster, M.,
Guérin, C., Pierce, N. E., Combes S. A., and de Bivort B. L. (2018). Neonicotinoid exposure disrupts
bumblebee nest behavior, social networks, and thermoregulation. *Science*, *362*(6415), 683-686.
https://doi.org/10.1126/science.aat1598

46. Gill, R. J., Ramos-Rodriguez, O., and Raine, N. E. (2012). Combined pesticide exposure severely affects
individual-and colony-level traits in bees. *Nature*, *491*(7422), 105-108.
https://doi.org/10.1038/nature11585

47. Schläppi, D., Kettler, N., Glauser, G., Straub, L., Yañez, O., and Neumann, P. (2021). Varying impact of
neonicotinoid insecticide and acute bee paralysis virus across castes and colonies of black garden ants,
Lasius niger (Hymenoptera: Formicidae). *Scientific reports*, *11*(1), 20500.
https://doi.org/10.1038/s41598-021-98406-w

Figure 1: Flupyradifurone susceptibility. Lines represent the proportion of *L. niger* workers surviving as a

function of time during chronical exposure to different concentrations of FPF (total sample size: N=467). Shaded

areas represent the 95% confidence interval around each line. Concentration had a significant effect on survival

(Cox proportional hazards model; $\chi^2 = 31.699$, df = 7, p < 0.001). Results of pairwise comparisons are indicated

by letters in the legend (same letter: p > 0.05; different letter: p ≤ 0.05 in *post hoc* tests with BH correction).

Figure 2: Interactive effects between Flupyradifurone and *M. brunneum*. a) Proportion of *L. niger* workers

surviving as a function of time following exposure to *M. brunneum* spores (yellow lines) or to a sham solution

(blue lines) after ten days chronic exposure to different concentrations of flupyradifurone (0, 5, and 50 ppm;

dashdotted, dashed and solid lines, respectively). Shaded areas represent the 95% confidence interval around each

curve. b) Hazard Ratio of fungus-treated ants relative to sham-treated ants in the three FPF exposure conditions.

Bars and whiskers represent model estimates and standard errors, respectively. Cox proportional hazards mixed

effect model, interaction FPF exposure $\times$ fungal exposure: $\chi^2 = 8.22$, df = 2, p = 0.016. Results of pairwise

comparisons are indicated by letters (same letter: p > 0.05; different letter: p ≤ 0.05; *post hoc* contrast with BH

correction).

Supplementary Materials - Synergistic effects of the next generation insecticide flupyradifurone with a fungal pathogen

Antifungal assay: Metarhizium brunneum growth inhibition test

Flupyradifurone acts by specifically impairing the nervous system of insects (Nauen et al., 2015) and thus, no antifungal activity is expected from this agrochemical. Nonetheless, any results of combined exposure treatments would be difficult to interpret or even invalidated if flupyradifurone did inhibit the growth of *M. brunneum.* To test the ability of flupyradifurone to inhibit *M. brunneum* germination we used a disk-diffusion assay as described in Balouiri et al. (2016). Briefly, SDA plates were inoculated with 50 µL of 10^9/mL *M. brunneum* conidiospore suspension. Then, filter paper disks (6 mm, Cytiva Life Sciences) were soaked with 15 µL of flupyradifurone solutions (10, 100, 1000 ppm), distilled water (0 ppm; negative control) or 5% sodium hypochlorite (positive control) and put on the agar using forceps. Plates were sealed, incubated for 48h at 24°C and photographed to measure the growth inhibition diameter. Even at the highest flupyradifurone dose no grow inhibition zone was visible, suggesting that *M. brunneum* germinated normally. Our findings are in line with findings for neonicotinoids, which have been used successfully in studies with fungal pathogens where no negative effects on conidia germination, conidia production and vegetative growth of Metarhizium fungi were detected (Cramer, 2020; Neves et al., 2001, Santos et al., 2007).

Supplementary Figure 1. Antifungal assay. Disk diffusion method to test for germination inhibition of (a) flupyradifurone at 0, 10, 100 and 1000 ppm and (b) sodium hypochlorite (positive control).

Quantification of food uptake

Methods

Pesticide contamination can modulate feeding motivation and food uptake, as shown for honey bees (Kessler et al., 2015; Siviter & Muth, 2022; Wu et al., 2021). Thus, to determine whether disparities in survival may be due to differences in food consumption, we quantified honey water uptake over a day at different FPF concentrations. For this purpose, we quantified fluorescence of homogenized ants that had access to a feeding solution containing fluorescein dye (Jové et al., 2020). The feeding solutions were prepared as described in the main manuscript, but with a fraction of water replaced by fluorescein sodium salt (Sigma-Aldrich) stock solution at 2% in MiliQ water. Four treatment solutions with final concentrations of 15% honey water, 0.01% fluorescein and varying concentrations of FPF (0, 5, 50 or 500 ppm, that is control, low, mid, and high treatment) were prepared. Workers were sampled from 5 source colonies and split into 4 subsets per colony in a stratified random way. Subsets were kept in petri dishes in the incubator for three days without food. Subsequently, each subset was assigned pseudo-randomly one of the four treatments groups. Final sample size was 106, 108, 105 and 102 workers in the control, low, mid and high treatment respectively. Workers had access to the solution for 24h before being freeze killed at -80°C for 30 min. Samples were then homogenized individually in racked collection tubes (Qiagen) containing 100 μL of phosphate-buffered saline (PBS) buffer and a 2 mm glass bead each. Homogenization was performed using a tissue lyser at 30 Hz for two runs of 1 minute with a plate rotation between the two runs. Following 2 minutes of centrifugation at 3000 rpm, 20 μL of the supernatant and 80 μL PBS buffer were transferred into 96-well plates. Fluorescence was measured on a SpectraMax iD5 plate reader with excitation at 485 nm and emission at 535 nm. Each measure was averaged from 4 reading points 1.5 mm apart, with 400 ms integration time and a gain of 500 volt.

Quantification was done using a standard curve, which was obtained using an aliquot of the control treatment feeding solution containing 0.01% fluorescein and 0 FPF, which was treated identically to the meals that were served to the ants (same light and temperature conditions throughout the duration of the experiment, and subsequent freezing). Reference standard curves were prepared from the retained

aliquot by making a serial dilution for 8 solutions containing 5, 2.5, 1.25, 0.625, 0.3125, 0.15625,

0.078125, or 0 μL of 0.01% fluorescein treatment solution mixed with PBS to a final total volume of

100 μL. An unfed ant treated identically to the samples was used as a blank. To calculate the meal

volumes, mean fluorescence value of blanks was subtracted from all other samples and consumed

volume was extrapolated from the standard curves.

Statistical analyses were performed in R. A generalized linear mixed model (GLMM) with a

negative binomial distribution was used to test for differences in meal size depending on FPF

concentration. Treatment was included as fixed effect and petri dishes, colonies as well a block as

random effects. Post hoc testing p-values were adjusted using the Benjamini-Hochberg method to

correct for multiple comparisons.

Results

The estimated consumed volume of honey water over 24 hours, was significantly different among

treatments (χ^2 = 7.853, df = 3, p = 0.049; supplementary figure 2). Pairwise comparisons using Tukey

adjustment indicated that ants in the treatment with the highest FPF concentration (500 ppm) consumed

lower volumes (0.06 ± 0.07 μL) compared to ants in the low treatment (0.17 ± 0.14 μL; p = 0.048; fig

2). The remaining pairwise comparisons with the control and the mid treatment (0.159 ± 0.14 μL and

0.15 ± 0.14 μL respectively) were not significant.

In contrast to honey bees, which respond to field-realistic concentrations of FPF by a decreased

food intake and erratic foraging behaviour (Hesselbach et al., 2020; Wu et al., 2021), we found that ants

do not alter their food uptake if FPF is present in food. Even at 500 ppm the food uptake was comparable

to the controls. Consequently, our results suggest that ants are not repelled by sublethal concentrations

of FPF, i.e. below 50 ppm. The absence of a repellent effect implies that the ants are prone to collect

contaminated food from the environment and sharing it among nestmates, with potentially severe

consequences for the colony. The insecticide is likely to reach all members of the colony, including the

queen. Even though the queen might have some protection via the colony or potentially superior

detoxification compared to workers (Schläppi et al., 2020), she will still get exposed repeatedly over

extended periods and thus might face a trade-off between detoxification and reproduction (Schwenke

et al., 2016).

References

Balouiri, M., Sadiki, M., & Ibnsouda, S. K. (2016). Methods for in vitro evaluating antimicrobial activity: A
review. Journal of pharmaceutical analysis, 6(2), 71-79. doi:10.1016/j.jpha.2015.11.005

Cramer, L. (2020) The synergistic effect of pesticides on the fitness of the ant species Cardiocondyla obscurior.
Magister thesis, Karl-Franzens-Universität Graz.

Hesselbach, H., Seeger, J., Schilcher, F., Ankenbrand, M., & Scheiner, R. (2020). Chronic exposure to the
pesticide flupyradifurone can lead to premature onset of foraging in honeybees Apis mellifera. Journal
of Applied Ecology, 57(3), 609-618. doi: 10.1111/1365-2664.13555

Jové, V., Venkataraman, K., Gabel, T. M., & Duvall, L. B. (2020). Feeding and quantifying animal-derived
blood and artificial meals in Aedes aegypti mosquitoes. JoVE (Journal of Visualized Experiments),
(164), e61835. doi: 10.3791/61835

Kessler, S. C., Tiedeken, E. J., Simcock, K. L., Derveau, S., Mitchell, J., Softley, S., ... & Wright, G. A. (2015).
Bees prefer foods containing neonicotinoid pesticides. Nature, 521(7550), 74-76. doi:
10.1038/nature14414

Nauen, R., Jeschke, P., Velten, R., Beck, M. E., Ebbinghaus-Kintscher, U., Thielert, W., ... & Raupach, G.
(2015). Flupyradifurone: a brief profile of a new butenolide insecticide. Pest management
science, 71(6), 850-862.

Neves, P. M., Hirose, E., Tchujo, P. T. & Moino JR, A. (2001). Compatibilidade de fungos entomopatogênicos
com inseticidas nicotinóides. Neotropical Entomology, 30, 263-268. doi:10.1590/S1519-
566X2001000200009

Santos, A. V., de Oliveira, B. L., & Samuels, R. I. (2007). Selection of entomopathogenic fungi for use in
combination with sub-lethal doses of imidacloprid: perspectives for the control of the leaf-cutting ant
Atta sexdens rubropilosa Forel (Hymenoptera: Formicidae). Mycopathologia, 163, 233-240.
doi:10.1007/s11046-007-9009-8

Schläppi, D., Kettler, N., Straub, L., Glauser, G., & Neumann, P. (2020). Long-term effects of neonicotinoid
insecticides on ants. Communications Biology, 3(1), 335. doi: 10.1038/s42003-020-1066-2

Schwenke, R. A., Lazzaro, B. P., & Wolfner, M. F. (2016). Reproduction–immunity trade-offs in insects. Annual
review of entomology, 61, 239-256. doi: 10.1146/annurev-ento-010715-023924

Siviter, H., & Muth, F. (2022). Exposure to the novel insecticide flupyradifurone impairs bumblebee feeding
motivation, learning, and memory retention. Environmental Pollution, 307, 119575. doi:
10.1016/j.envpol.2022.119575

Wu, Y. Y., Pasberg, P., Diao, Q. Y., & Nieh, J. C. (2021). Flupyradifurone reduces nectar consumption and
foraging but does not alter honey bee recruitment dancing. Ecotoxicology and Environmental
Safety, 207, 111268.