# Astonishing diversity and multifaceted biological connections of Type IV restriction-modification systems

Ryan T. Bell[1,*], Jonas Juozapaitis[2], Inga Songailiene[2], Harutyun Sahakyan[1], Toma Liegute[2], Kira S. Makarova[1], Yuri I. Wolf[1], Virginijus Siksnys[2], Eugene V. Koonin[1,*]

[1]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD 20894, USA

[2]Institute of Biotechnology, Life Sciences Center, Vilnius University, Vilnius, Lithuania

*Correspondence to: Ryan T. Bell (ryan.bell@nih.gov) or Eugene V. Koonin (koonin@ncbi.nlm.nih.gov)

## Abstract

A comprehensive census of McrBC systems, a major form of prokaryotic Type IV restriction-modification (RM), followed by phylogenetic analysis, reveals their enormous abundance in diverse bacteria and archaea and a plethora of genomic associations. We focus on a previously uncharacterized branch, which we denote CoCoNuTs (coiled-coil nuclease tandems) for their salient features, the presence of extensive coiled-coil structures and tandem nucleases. The CoCoNuTs alone show extraordinary variety, with 3 distinct types and multiple subtypes. All CoCoNuTs contain domains predicted to interact with translation system components, such as OB-folds resembling the SmpB protein that binds bacterial tmRNA, YTH-like domains that might recognize methylated tmRNA, tRNA, or rRNA, and RNA-binding Hsp70 chaperone homologs, along with RNases, such as HEPN domains, all suggesting that the CoCoNuTs target RNA. Many CoCoNuTs may also target DNA, via McrC nuclease homologs. Additional restriction systems, such as Type I RM and Druantia Type III, are frequently encoded in the same predicted superoperons. In many of these superoperons, CoCoNuTs are likely regulated by cyclic nucleotides, possibly, RNA fragments with cyclic termini, that bind associated CARF (CRISPR-Associated Rossmann Fold) domains. We experimentally demonstrate the ability of Type II CoCoNuTs to protect *E. coli* against specific phages and plasmids.

## Introduction

All organisms are subject to an incessant barrage of genetic parasites, such as viruses and
transposons. Over billions of years, the continuous arms race between hosts and parasites drove the
evolution of immense, intricately interconnected networks of diverse defense systems and pathways [1-
6]. In particular, in the last few years, targeted searches for defense systems in prokaryotes, typically
capitalizing on the presence of variable genomic defense islands, have dramatically expanded their
known diversity and led to the discovery of a plethora of biological conflict strategies and mechanisms
[2, 4, 7, 8].

One of the most ancient and common forms of defense against mobile genetic elements (MGE)
is the targeted restriction of nucleic acids. Since the initial discovery of this activity among strains of
bacteria resistant to certain viruses, myriad forms of recognition and degradation of nucleic acids have
been described, in virtually all life forms. Characterization of the most prominent of these systems, such
as restriction-modification (RM), RNA interference (RNAi), and particularly CRISPR-Cas (<u>c</u>lustered
<u>r</u>egularly <u>i</u>nterspaced <u>s</u>hort <u>p</u>alindromic <u>r</u>epeats-<u>C</u>RISPR <u>as</u>sociated genes), has led to the development
of a profusion of highly effective experimental and therapeutic techniques, in particular, genome editing
and engineering [9-15].

Historically, the two-component McrBC (<u>m</u>odified <u>c</u>ytosine <u>r</u>estriction) system was the first form
of restriction to be described, although the mechanism remained obscure for decades, and for a time,
this system was referred to as RglB (<u>r</u>estriction of <u>gl</u>ucose<u>l</u>ess phages) due to its ability to restrict T-even
phage DNA which contained hydroxymethylcytosine, but not glucosylated bases [16-19]. Today, the
prototypical McrBC system, native to *E. coil* K-12, is considered a Type IV (modification-dependent)
restriction system that degrades DNA containing methylcytosine (5mC) or hydroxymethylcytosine
(5hmC), with a degree of sequence context specificity [20, 21].

Type IV restriction enzymes contain at least two components: 1) a dedicated specificity domain
that recognizes modified DNA, and 2) an endonuclease domain that cleaves the target [9, 22]. In the
well-characterized example from *E. coli* K-12, McrB harbors an N-terminal DUF(domain of unknown
function)3578 that recognizes methylcytosine. We denote this domain, as its function is not unknown,
as ADAM (<u>a d</u>omain with an <u>a</u>ffinity for <u>m</u>ethylcytosine). The ADAM domain is fused to a GTPase domain
of the AAA+ ATPase superfamily, the only known GTPase in this clade, which is believed to translocate
DNA (Figure 1 A, B, C) [20, 23-25]. McrC, encoded by a separate gene in the same operon, contains an N-
terminal DUF2357 domain, which interacts with the GTPase domain in McrB and stimulates its activity,
and is fused to a PD-(D/E)xK superfamily endonuclease (Figure 1 A, B, D) [20, 26].

Our recent analysis of the modified base-binding EVE (named for Protein Data Bank (PDB)
structural identifier 2eve) domain superfamily demonstrated how the distribution of the EVE-like
domains connects the elaborate eukaryotic RNA regulation and RNA interference-related epigenetic
silencing pathways to largely uncharacterized prokaryotic antiphage restriction systems [4]. EVE
superfamily domains, which in eukaryotes recognize modified DNA or RNA as part of mRNA maturation
or epigenetic silencing functions, are often fused to McrB-like GTPases in prokaryotes, and indeed, these
are the most frequently occurring EVE-containing fusion proteins [4]. These observations motivated us
to conduct a comprehensive computational search for McrBC systems, followed by a census of all
associated domains, to chart the vast and diverse population of antiviral specificity modules, vital for

prokaryotic defense, that also provided important source material during the evolution of central
signature features of eukaryotic cells. Here we present the results of this census and describe an
extraordinary, not previously appreciated variety of domain architectures of the McrBC family of Type IV
restriction systems. In particular, we focus on a major McrBC branch that we denote coiled-coil nuclease
tandem (CoCoNuT) systems, which we explore in detail and report proof of principle experiments
demonstrating the anti-MGE defense mediated by the CoCoNuTs.

## Materials and Methods

*Comprehensive identification and phylogenetic and genomic neighborhood analysis of McrB and*
*McrC proteins*

The comprehensive search for McrB and McrC proteins was seeded with publicly available
multiple sequence alignments COG1401 (McrB GTPase domain), AAA_5 (The branch of AAA+ ATPases
containing the McrB GTPase), COG4268 (McrC), PF10117 (McrBC), COG1700 (McrC PD-D/ExK nuclease
domain), PF04411 (McrC PD-D/ExK nuclease domain), PF09823 (McrC N-terminal DUF2357). Additional
alignments and individual queries were derived from data from our previous work on the EVE domain
family [4]. All alignments were clustered, and each sub-alignment or individual query sequence was used
to produce a position-specific scoring matrix (PSSM). These PSSMs were used as PSI-BLAST queries
against the non-redundant (nr) NCBI database (*E*-value ≤ 10) [27]. Genome neighborhoods for the hits
were generated by downloading their gene sequence, coordinates, and directional information from
GenBank, as well as for 10 genes on each side of the hit. Domains in these genes were identified using
PSI-BLAST against alignments of domains in the NCBI Conserved Domain Database (CDD) and Pfam (*E*-
value 0.001). Some genes were additionally analyzed with HHpred [28]. Then, these neighborhoods
were filtered for the presence of a COG1401 hit (McrB GTPase domain), or hits to both an McrB "alias"
(MoxR, AAA_5, COG4127, DUF4357, Smc, WEMBL, Myosin_tail_1, DUF3578, EVE, Mrr_N, pfam01878)
and an McrC "alias" (McrBC, McrC, PF09823, DUF2357, COG1700, PDDEXK_7, RE_LlaJI). These aliases
were determined from a preliminary manual investigation of the data using HHpred [29].  Several of the
McrB aliases are not specific to McrB, and instead are domains commonly fused to McrB GTPase
homologs, or are larger families, such as AAA_5, that contain McrB homologs. We found that many *bona*
*fide* McrB homologs, validated by HHpred, produced hits not to COG1401 but rather to these other
domains, so we made an effort to retain them.
The McrB candidates identified by this filtering process were clustered to a similarity threshold
of 0.5 with MMseqs2 [30], after which the sequences in each cluster were aligned with MUSCLE [31].
Next, profile-to-profile similarity scores between all clusters were calculated with HHsearch [32].
Clusters with high similarity, defined as a pairwise score to self-score ratio > 0.1, were aligned to each
other with HHalign [33]. This procedure was performed for a total of 3 iterations. The alignments
resulting from this protocol were analyzed with HHpred to remove false positives, after which the
GTPase domain sequences were extracted manually using HHpred, and the alignments were used as
queries for a second round of PSI-BLAST against the nr NCBI database as described above. At this stage,
the abundance of the CoCoNuT and CoCoPALM (coiled-coil and pilus assembly linked to McrBC, see
below) types of McrB GTPases had become apparent, therefore, results of targeted searches for these
subtypes were included in the pool of hits from the second round of PSI-BLAST. Genome neighborhoods
were generated for these hits and filtered for aliases as described above. Further filtering of the data,
which did not pass this initial filter, involved relaxing the criteria to include neighborhoods with only one
hit to an McrB or McrC alias, but with a gene adjacent to the hit (within 90 nucleotides), oriented in the

| 114 | same direction as the hit, and encoding a protein of sufficient size (> 200 aa for McrB, > 150 aa for McrC) |
| 115 | to be the undetected McrB or McrC component. Afterward, we filtered the remaining data to retain |
| 116 | genome islands with no McrBC aliases but with PSI-BLAST hits in operonic association with genes of |
| 117 | sufficient size, as described above, to be the other McrBC component. These data, which contained |
| 118 | many false positives but captured many rare variants, were then clustered and analyzed as described |
| 119 | above to remove false positives. Next, an automated procedure was developed to excise the GTPase |
| 120 | domain sequences using the manual alignments generated during the first phase of the search as a |
| 121 | reference. These GTPase sequences were further analyzed by clustering and HHpred to remove false |
| 122 | positives. Definitive validation by pairing McrB homologs with their respective McrC homologs was also |
| 123 | used to corroborate their identification. Occasionally, the McrB and McrC homologs were separated by |
| 124 | intervening genes, or the operon order was reversed, and consideration of those possibilities allowed |
| 125 | the validation of many additional systems. The pairing process was complicated by and drew our |
| 126 | attention to the frequent occurrence of multiple copies of McrBC systems in the same islands that may |
| 127 | function cooperatively. Finally, the rigorously validated set of McrBC pairs, supplemented only with |
| 128 | orphans manually annotated as McrBC components using HHpred, were used for our phylogenetics. The |
| 129 | final alignments of GTPase and DUF2357 domains were produced using the iterative alignment |
| 130 | procedure described above for 10 iterations. Approximately-maximum-likelihood trees were built with |
| 131 | the FastTree program [34] from representative sequences following clustering to a 0.9 similarity |
| 132 | threshold with MMseqs2. The phyletic distribution was calculated by assigning a weight to the lineage of |
| 133 | each sequence that was inversely proportional to its cluster size (1/n, where n=the cluster size). |
| 134 | |
| 135 | *Preliminary phylogenetic analysis of CoCoNuT CnuH helicases* |
| 136 | |
| 137 | A comprehensive search and phylogenetic analysis of this family was beyond the scope of this |
| 138 | work, but to determine the relationships between the CoCoNuTs and the rest of the UPF1-like helicases, |
| 139 | we used the following procedure. We retrieved the best 2000 hits in each of two searches with UPF1 |
| 140 | and CoCoNuT helicases as queries against both a database containing predicted proteins from 24,757 |
| 141 | completely sequenced prokaryotic genomes downloaded from the NCBI GenBank in November 2021 |
| 142 | and a database containing 72 representative eukaryotic genomes that were downloaded from the NCBI |
| 143 | GenBank in June 2020. Next, we combined all proteins in four searches, made a nonredundant set, and |
| 144 | annotated them using CDD profiles, as described above. Then, we aligned them with MUSCLE v5 [35], |
| 145 | constructed an approximately-maximum-likelihood tree with FastTree, and mapped the annotations |
| 146 | onto the tree. Genome neighborhoods were generated for these hits, as described above. |

*Structural modeling with AlphaFold2 and searches for related folds*

We predicted the structures described in this work using AF2 v2.2.0 with local installations of
complete databases required for AF2 [36].  We used only protein models with average predicted local
distance difference test (pLDDT) scores ≥ 80 for further analysis [37]. We used many of these models as
queries to search for structurally similar proteins using DALI v5 against the Protein Data Bank (PDB) and
using FoldSeek against the AlphaFold/UniProt50 v4, AlphaFold/Swiss-Prot v4, AlphaFold/Proteome v4,
and PDB100 2201222 databases [38, 39]. Structure visualizations and comparisons were performed with
ChimeraX [40] and the RCSB PDB website [41].

*Gene cloning*

Synthetic gene fragments coding CoCoNuT gene operons with 100 nt upstream regions were
ordered commercially from "Twist Bioscience." Gene fragments were designed with ~30 nucleotide

complementary ends. pACYC184 vector was amplified by PCR, generating ~30 nucleotide ends
complementary to the operon ends. Fragments were assembled using Gibson assembly and
transformed into *E. coli* DH5α. The Type II CoCoNuT TerY-PP2C-STK only variant was constructed by
whole plasmid PCR retaining only the gene triad of interest, followed by blunt-end ligation and
transformation. Plasmids were purified by silica spin columns and sequenced.

*Phage plaque assay*

Bacteriophages were amplified in the *E. coli* MG1655 strain. 0.2-1 ml of *E. coli* night cultures
(depending on culture density) were diluted with Luria-Bertani (LB) medium to 2 ml volume and mixed
with 2 ml of hot agar-LB (1.5% agar, 60 °C). 4 ml of soft agar with bacteria was immediately spread on
LB-agar Petri dishes supplemented with 35 μg/ml chloramphenicol and air-dried for 20 min. 3.5 μl of 10-
fold phage dilutions were plated on soft agar, air-dried for 20 min, and incubated for 16 h at 30-37° C,
depending on the phage used.

*Phage challenge in liquid medium*

100 μl of *E. coli* night cultures grown with 1 % glucose were added to 4 ml of fresh LB medium
and incubated at 37 °C in a shaker for 5 h. Bacteria were diluted to $OD_{600}$ 0.5 and kept on ice while
preparing a 96-well flat-bottom plate. To limit condensation, the inside of the 96-well plate lid was
impregnated with 20% ethanol 0.05% Triton X-100 solution and dried on a paper towel. In the plate, 70
μl of LB medium and 20 μl of diluted bacteria were mixed. Just before measuring, 10 μl of phage or LB
medium for control was added to 90 μl of diluted cells. The final $OD_{600}$ was 0.1, and phage was added at
MOI 0.05. The dried lid was replaced, sides of the plate were sealed with parafilm tape and immediately
transferred to the plate reader. Cultures were incubated at 37°C at 200 rpm and $OD_{600}$ was measured
every 5 min. For all dilutions and cultures, the same stock of LB medium supplemented with 35 μg/ml
chloramphenicol was used. For phage λ-vir, 5 mM of $CaCl_2$ was added to soft agar.

*CFU measurement after phage infection*

100 μl of *E. coli* night cultures grown with 1% glucose were added to 4 ml of fresh LB medium
supplemented with 35 μg/ml chloramphenicol and incubated at 37 °C in a shaker for 5 h. Bacteria were
diluted to $OD_{600}$ 0.1, supplemented with 5 mM of $CaCl_2$, phage λ-vir added at MOI 0.05, and incubated at
37°C at 200 rpm. After 4 hours of incubation, an equal volume of cultures was transferred to a 1.5 ml
tube and centrifuged for 5 minutes at 12000 rpm on a table centrifuge. The supernatant containing
phages was removed by pipette and bacteria were resuspended in the same volume of LB medium
without $CaCl_2$. Serial 10-fold dilutions were performed and 3.5 μl of diluted cultures were plated on LB
agar, air-dried for 20 minutes, and incubated for 16 h at 37° C. CFU/ml was determined by counting the
colonies at the lowest visible dilution, taking into account the dilution and 3.5 μl volume of the plated
drop. All LB media were supplemented with chloramphenicol.

*Plasmid interference assay*

*E. coli* MG1655 strain cells were pre-transformed with pACYC184 plasmids (chloramphenicol
resistance, Cm) harboring each wild-type CoCoNuT defense system variant or control empty pACYC184
vector. CoCoNuT cells were cultivated in LB medium overnight supplemented with 2% glucose and
chloramphenicol 30 μg/ml at 37°C, 200 rpm. The next day, flasks with 50 ml of liquid LB medium and 30
μl/ml chloramphenicol were inoculated with cells from overnight culture at 37°C, 200 rpm until they

reached $OD_{600}$~ 0.6. The same number of cells for each variant (based on OD) were taken and
centrifuged for 10 min, 3000 rpm, next washed using 10 ml deionized water 4 times, resuspended in
1500 µl of deionized water, and used for electroporation using Thermo Scientific™ Electroporation
Cuvettes, 1mm. 100 ng of different potential target plasmids bearing distinct *ori* sequences
(Supplementary Table S3) were electroporated into the prepared cells with or without CoCoNuT. Cells
were immediately transferred to Eppendorfs with fresh LB medium and left for 60 min at 37°C 200 rpm
for recovery. Next, the cells were serially diluted in a 96-well plate with LB medium and 5 µl were
spotted on a selection medium with chloramphenicol and a selection medium containing additional
antibiotic for the selection of co-electroporated plasmid. Plates were incubated at 37°C for 16h.

## Results and Discussion
209

### *Comprehensive census of McrBC systems*
210

Our initial search for McrBC systems consisted of PSI-BLAST runs against the non-redundant
protein sequence database at the NCBI using as queries position-specific scoring matrices (PSSMs)
derived from publicly available alignments of McrB and McrC homologs, or our EVE-McrB data [4].
Although a branch of McrB GTPase homologs has been described in animals, these are highly divergent
in function, and no associated McrC homologs have been reported [23]. Therefore, we excluded
eukaryotic sequences from our analysis to focus on the composition and contextual connections of
prokaryotic McrBC systems. Through this preliminary search, we obtained a set of approximately 12,500
McrBC homologs, which was validated using HHpred and several other criteria, including the presence of
the McrB GTPase signature motif NxxD (where x is any amino acid), which distinguishes McrB from
closely related AAA+ ATPase families such as MoxR (Figure 1 B) [23, 26, 42, 43]. Using alignments of the
McrB GTPase domains from this set, we conducted a second iteration of searches and validation,
increasing the number of McrBC systems to nearly 34,000, distributed broadly among prokaryotes
(Figure 1 E).

**Figure 1: Genetic organization, signature sequence motifs, structural models, and phyletic distribution of McrB GTPases detected in this work**

A) McrBC is a two-component restriction system with each component typically (except for extremely rare gene fusions) encoded by a separate gene expressed as a single operon, depicted here and in subsequent figures as arrows pointing in the direction of transcription. In most cases, McrB is the upstream gene in the operon. B) In the prototypical *E.coli* K-12 McrBC system, McrB contains an N-terminal methylcytosine-binding domain, ADAM/DUF3578, fused to a GTPase of the AAA+ ATPase clade [21]. This GTPase contains the Walker A and Walker B motifs that are conserved in P-loop NTPases as well as a signature NxxD motif, all of which are required for GTP hydrolysis [24, 26, 44]. McrC consists of a PD-DxK nuclease and an N-terminal DUF2357 domain with a stalk-like extension comprising a helical bundle that interacts with and activates individual McrB GTPases while they are assembled into hexamers [24, 26]. C) AlphaFold2 structural model of *E. coli* K-12 ADAM-McrB GTPase monomer. D) AlphaFold2 structural model of *E. coli* K-12 McrC monomer with DUF2357-PD-D/ExK architecture. The structures were visualized with ChimeraX [40]. E) Phyletic distribution of McrB GTPase homologs detected in this work, each found in a genomic island with distinct domain composition, clustered to a similarity threshold of 0.9, then assigned a weight inversely proportional to the cluster size (see Methods).

In the subsequent phase of analysis, we extracted GTPase domain sequences from the McrB homolog pool, leaving as a remainder the fused specificity domains that we intended to classify. Unexpectedly, we found that about 34% of the *mcrB* genes contained insertions into the GTPase coding sequence. These insertions likely encode specificity domains, both known and uncharacterized, typically located between the Walker A (P-loop) and Walker B motifs of the GTPase [23, 24, 26]. The previously identified specificity domains are usually fused at the N terminus, or rarely, at the C-terminus of McrB-like GTPases [21, 45].

We also observed that, although the McrC homologs are not the primary bearers of specificity
modules in McrBC systems, they can be fused to various additional domains, including those of the EVE
superfamily (Figure 2, Supplementary Figure S1). Many McrC homologs lack the typical PD-D/ExK
nuclease domain or contain an apparently inactivated version of this domain that lacks the signature
catalytic site, but the N-terminal DUF2357 domain, which interacts with McrB, is nearly always present.
To isolate the likely specificity domains, we first attempted to extract the two common domain
sequences from the pool of McrC homologs. Much like in the case of McrB GTPases, we identified
insertions in many DUF2357 sequences. Most of these insertions consisted of predicted coiled-coils, but
some contained beta-sheets predicted to fold into distinct domains. We defined the conserved regions
with the maximum precision possible using HHpred, coiled-coil prediction tools, and AlphaFold2 (AF2)
structural predictions [28, 36, 46-49].

Removing the variable inserts from the conserved McrBC domains allowed accurate,
comprehensive phylogenetic analysis of the McrB GTPase (Figure 2) and McrC DUF2357 (Supplementary
Figure S1) families. These two trees were generally topologically concordant and revealed several
distinct branches not previously recognized. The branch containing the prototypical McrBC system from
*E. coil* K-12 is characterized by frequent genomic association with Type I RM systems, usually with
unidirectional gene orientations and the potential of forming a single operon [50]. This branch and
others, which exhibit two particularly prevalent associations, with a DISARM-like antiphage system [51],
and, surprisingly for a Type IV restriction system, with predicted DNA methyltransferases, will be the
subject of a separate, forthcoming publication.

269                  **Figure 2: Phylogenetic tree of the McrB GTPases**

The major clades in the phylogenetic tree of the McrB GTPases are distinguished by the distinct versions of the Nx(xx)D
signature motif. The teal and yellow groups, with bootstrap support of 97%, have an NxD motif, whereas the blue, green, red,
and purple groups, with variable bootstrap support, have an NxxD motif, indicated by the arrows; the sequences in the smaller,
cyan clade, with 98% bootstrap support, have an NxxxD motif. Each of the differently colored groups is characterized by distinct
conserved genomic associations that are abundant within but not completely confined to the respective groups. This tree was

built from the representatives of 90% identity clusters of all validated homologs. Abbreviations of domains: McrB – McrB
GTPase domain; CoCo/CC – coiled-coil; MN – McrC N-terminal domain (DUF2357); CSD – cold shock domain; IG –
Immunoglobulin (IG)-like beta-sandwich domain; ZnR – zinc ribbon domain; SPB – SmpB-like domain; RTL – RNase toxin-like
domain; HEPN – HEPN family nuclease domain; OB – OB-fold domain; iPD-D/ExK – inactivated PD-D/ExK fold; Hsp70-like ATPase
– Hsp70-like NBD/SBD; HEAT – HEAT-like helical repeats; YprA – YprA-like helicase domain; DUF1998 - DUF1998 is often found
in or associated with helicases and contains four conserved, putatively metal ion-binding cysteine residues; SWI2/SNF2 –
SWI2/SNF2-family ATPase; PglX – PglX-like DNA methyltransferase; HsdR/M/S – Type I RM system restriction, methylation, and
specificity factors.

*A variant of the McrB GTPase signature motif distinguishes a large group of unusual McrBC systems*

A major branch of the McrBC family is characterized by a conserved deletion within the NxxD
GTPase signature motif, reducing it to NxD (Figure 1 B, Figure 2). No NxD motif McrB GTPase has been
characterized, but the extensive study of the NxxD motif offers clues to the potential impact of the motif
shortening. The asparagine (N) residue is strictly required for GTP binding and hydrolysis [44]. As this
residue is analogous to sensor-1 in ATP-hydrolyzing members of the AAA+ family, it can be predicted to
position a catalytic water molecule for nucleophilic attack on the γ-phosphate of an NTP [26, 43, 44, 52,
53]. The role of the conserved aspartate (D), however, was initially unclear. Given the specificity of McrB
for GTP, the motif was proposed to function similarly to the NKxD G4 box in conventional GTPases and
confer specificity for GTP by interacting with the guanine ring [44]. However, a D336N mutation in the *E.*
*coli* K-12 McrB GTPase NxxD motif did not alter binding specificity, unlike the corresponding mutation in
EF-Tu and several other GTPases containing the G4 box [44]. Instead, this substitution unexpectedly
inhibited GTP hydrolysis and prevented the stimulation of GTPase activity by McrC [44]. A recent
structural analysis has shown that this aspartate interacts with a conserved arginine/lysine residue in
McrC which, via a hydrogen-bonding network, resituates the NxxD motif in relation to its interface with
the Walker B motif such that, together, they optimally position a catalytic water to stimulate hydrolysis
[26]. Accordingly, the truncation of this motif might be expected to modulate the rate of hydrolysis, and
potentially compel functional association with only a subset of McrC homologs containing compensatory
mutations. Indeed, the NxD branch is characterized by many McrC homologs with unusual features, such
as predicted RNA-binding domains, that might not be compatible with conventional McrB GTPases from
the NxxD clade, which often occur in the same genomes (see below).

Many of these NxD GTPases are contextually associated with DNA methyltransferases, like the
NxxD GTPases, but are distinguished by additional complexity in the domains fused to the McrC
homologs and the frequent presence of two-component regulatory system genes in the same operon
(Figure 2). We also detected a small number of GTPases with an insertion in the signature motif (Figure
2), expanding it to NxxxD, which are methyltransferase-associated as well. This association is likely to be
ancestral, as it is found in all 3 branches of the McrB GTPase tree with different signature motif
variations.

The most notable feature of the NxD branch of GTPases is a large clade characterized by fusion
to long coiled-coil domains, a derived and distinctly different architecture from the small, modular DNA-
binding specificity domains typically found in canonical McrB homologs (Figure 2). The McrC homologs
associated with these coiled-coil McrB GTPase fusions also often lack the PD-D/ExK endonuclease
entirely or contain an AF2-predicted endonuclease fold with inactivating replacements of catalytic
residues (Supplementary Table S1). However, in the cases where the McrC nuclease is missing or likely

inactive, these systems are always encoded in close association, often with overlapping reading frames,
with HEPN (higher eukaryotes and prokaryotes nucleotide-binding) ribonuclease domains, or other
predicted nucleases (Figure 3, Supplementary Figure S2) [54]. Based on these features of domain
architecture and genomic context, we denote these uncharacterized McrBC-containing operons coiled-
coil nuclease tandem (CoCoNuT) systems.

The deepest branching group of the NxD GTPase systems is typified by a fusion of an Hsp70-like
nucleotide-binding domain and protein substrate-binding domain (NBD/SBD) to the McrB GTPase
domain (Figure 2, Supplementary Figure S2, Figure 3, Supplementary Table S1). Hsp70 is an ATP-
dependent protein chaperone that binds exposed hydrophobic peptides and facilitates protein folding
[55]. It also associates with AU-rich mRNA, an interaction involving both the ATPase and SBD [56-58].
These systems might be functionally related to the CoCoNuTs, given that an analogous unit is encoded
by a type of CoCoNuT system where similar domains are fused to the McrC homolog rather than to the
McrB GTPase homolog (see below). In another large clade of CoCoNuT-like systems, the GTPase is fused
to a domain homologous to FtsB, an essential bacterial cell division protein containing transmembrane
and coiled-coil helices (Supplementary Table S1) [59]. These enzymes form the core of rapidly evolving
protein secretion systems in Bacillota and Mycoplasmatota, likely regulated by EAL and HD-GYP domains
that bind cyclic di-GMP, and are presumably implicated in defense (Supplementary Figure S2) [60].
These unusual McrBC systems were primarily detected in metagenomic data obtained from the gut
microbiomes of ruminants and humans and might be involved in host-microbe interactions. They are
associated with signal peptidase family proteins likely to function as pilus assembly factors [47] and
usually contain coiled-coil domains fused to both the McrB and McrC homologs. Therefore, we denote
them as coiled-coil and pilus assembly linked to McrBC (CoCoPALM) systems (Supplementary Figure S2).
Here, we focus on the CoCoNuTs, whereas the CoCoPALMs and the rest of the NxD GTPase
methyltransferase-associated homologs will be explored in a separate, forthcoming publication.

**Figure 3: CoCoNuT system phylogeny and classification**

The figure shows the detailed phylogeny of McrB GTPases from CoCoNuT systems and their close relatives. All these GTPases possess an NxD GTPase motif rather than NxxD. CoCoNuT systems are abundant in Pseudomonadota and Terrabacteria, with the various subtypes generally concentrated in a more restricted range of species, as indicated. This tree was built from the representatives of 90% clustering of all validated homologs. Abbreviations of domains: McrB – McrB GTPase domain; CoCo/CC – coiled-coil; MN – McrC N-terminal domain (DUF2357); CSD – cold shock domain; YTH – YTH-like domain; IG – Immunoglobulin (IG)-like beta-sandwich domain; Hsp70 – Hsp70-like NBD/SBD; HEAT – HEAT-like helical repeats; ZnR – zinc ribbon domain; PYD – pyrin/CARD-like domain; SPB – SmpB-like domain; RTL – RNase toxin-like domain; HEPN – HEPN family nuclease domain; OB – OB-fold domain; iPD-D/ExK – inactivated PD-D/ExK fold; REC – Phosphoacceptor receiver-like domain; PLD – Phospholipase D-like nuclease domain; Vsr – very-short-patch-repair PD-D/ExK nuclease-like domain. Underneath each gene is a proposed protein name, with Cnu as an abbreviation for CoCoNuT.

CoCoNuT systems are extremely diverse and represented in a wide variety of bacteria, particularly Pseudomonadota and Bacillota, but are nearly absent in archaea (Figure 4).

**Figure 4: Phyletic distribution of CoCoNuTs**

The phyletic distribution of CnuB/McrB GTPases in CoCoNuT systems found in genomic islands with distinct domain
compositions. Most CoCoNuTs are found in either Bacillota or Pseudomonadota, with particular abundance in
Gammaproteobacteria. Type I-B and the related Pseudo-CoCoNuTs are restricted mainly to Bacillota. In contrast, the other
types are more common in Pseudomonadota, but can be found in a wide variety of bacteria. Type III-B and III-C are primarily
found in Alphaproteobacteria and Cyanobacteriota, respectively.

*Type I CoCoNuT systems*

We classified the CoCoNuTs into 3 types and 7 subtypes based on the GTPase domain phylogeny
and conserved genomic context (Supplementary Figure S2, Figure 3, Figure 4). Type I-A and Type I-C
systems consist of McrB and McrC homologs only (Figure 5), which we denote CnuB and CnuC (Figure 3,
Figure 4). Type I-B systems usually encode a separate coiled-coil protein, which we denote CnuA, in
addition to CnuB/McrB and CnuC/McrC, with no coiled-coil fused to the McrB GTPase domain in CnuB
(Figure 3, Figure 4, Supplementary Figure S2, Figure 5). Deeper branching Type I-B groups lack CnuA and
more closely resemble Type I-C in architecture, but the GTPase domains and N-terminal
oligonucleotide/oligosaccharide-binding (OB)-folds are more similar to the rest of Type I-B. CnuA is fused
at the N-terminus to a pyrin (PYD)/CARD-like helical domain and at the C-terminus to a REC domain
(Figure 5, Supplementary Table S1). Notably, in animals, the prototypical PYD domain of the PYRIN
protein is also fused to a long coiled-coil domain [61]. PYD and CARD (caspase activation and
recruitment domain) folds belong to the DEATH domain superfamily and are best characterized in the
context of innate immunity, inflammasome formation, and programmed cell death (PCD), where these
domains mediate the formation of multisubunit complexes via homotypic interactions [62]. An extensive
network of (putative) prokaryotic defense systems containing DEATH superfamily domains has recently
been discovered [8, 63]. The association of the PYD domain with CoCoNuTs suggests involvement in a
PCD/abortive infection-type response. The REC domains constitute one of the components of two-
component regulatory systems. They are targeted for phosphorylation by histidine kinases [64], which
could be a mechanism of Type I-B CoCoNuT regulation.

In contrast to Type II and Type III CnuB homologs, which contain only coiled-coils fused at their
N-termini, cold-shock domain (CSD)-like OB-folds are usually fused at the N-termini of Type I CoCoNuT
CnuB/McrB proteins, in addition to coiled-coils (except for the majority of Type I-B, where the coiled-
coils are encoded separately) (Supplementary Figure S2, Figure 3, Figure 5, Supplementary Table S1)
[65]. There is also considerable variation in the length of the coiled-coils in Type I-C that was not
observed in the other CoCoNuTs. Given that Type I-C is one of the deepest CoCoNuT branches, this
greater variation was not unexpected, mirroring what we observed in the other basal clade, Type I-B.
Often, in Type I-B and I-C, but not Type I-A, a YTH-like domain, a member of the modified base-binding
EVE superfamily, is present in CnuB as well, between the CSD and coiled-coil, or in Type I-B, between the
CSD and the GTPase domain (Supplementary Figure S2, Figure 3, Figure 5, Supplementary Table S1) [4,
66, 67].

The CnuC/McrC proteins in Type I CoCoNuTs all contain an immunoglobulin-like N-terminal
beta-sandwich domain of unknown function, not present in the *E. coli* K-12 McrC homolog, similar to a
wide range of folds from this superfamily with diverse roles (Figure 5) [68, 69]. It is also present in non-
CoCoNuT McrC homologs associated with McrB GTPase homologs in the NxD clade, implying its function
is not specific to the CoCoNuTs.

**Figure 5: Domain composition, operon organization, and AlphaFold2 structural predictions of components of the Type I CoCoNuT systems**

A) Type I CoCoNuT domain composition and operon organization. The arrows indicate the direction of transcription. B-D) High quality (Average pLDDT > 80), representative AlphaFold2 structural predictions for proteins in B) Type I-B CoCoNuT systems, C) Type I-A, and D) Type I-C CoCoNuT systems. Models were generated from representative sequences with the following GenBank accessions (see Supplementary Data for sequences and locus tags): ROR86958.1 (Type I-A CnuB), APL73566.1 (Type I-A CnuC), GED20858.1 (Type I-B CnuB), GED20857.1 (Type I-B CnuC), TKH01449.1 (Type I-B CnuA), GFD85286.1 (Type I-C CnuB), MBV0932851.1 (Type I-C CnuC). Abbreviations of domains: CSD – cold shock domain; YTH – YTH-like domain; CoCo – coiled-coil; IG – Immunoglobulin (IG)-like beta-sandwich domain; ZnR – zinc ribbon domain; PYD – pyrin/CARD-like domain; REC – Phosphoacceptor receiver-like domain. These structures were visualized with ChimeraX [40].

We also detected a close relative of Type I-B CoCoNuT in many *Bacillus* species, which we denoted Pseudo-CoCoNuT because it lost the separate coiled-coil protein CnuA. Pseudo-CoCoNuT occurs in genomic contexts suggesting a role in overcrowding-induced stress responses (Supplementary Figure S2, Figure 3, Supplementary Figure S3). We inferred this functional prediction from the fact that these islands encode a quorum-sensing hormone synthase, mechanosensitive ion channel, various transporters, antibiotic resistance and synthesis factors, and cell wall-related proteins. In many cases, we detected Type I-B CoCoNuT homologs in the extended neighborhoods of the Pseudo-CoCoNuTs, although only rarely in the immediate vicinity. Thus, Pseudo-CoCoNuTs might be derived duplicates of Type I-B CoCoNuTs that acquired a specialized but likely related functionality, perhaps still using the coiled-coil protein encoded by the Type I-B CoCoNuT.

*Type II and III CoCoNuT systems*

Type II and III CoCoNuT CnuB/McrB GTPase domains branch from within Type I-A and are encoded in a nearly completely conserved genomic association with a Superfamily 1 (SF1) helicase of the

UPF1-like clade, which we denote CnuH (Supplementary Figure S2, Figure 3, Figure 4, Supplementary
Table S1) [70, 71]. The UPF1-like family encompasses helicases with diverse functions acting on RNA and
single-stranded DNA (ssDNA) substrates, and notably, the prototypical UPF1 RNA helicase and its closest
relatives are highly conserved in eukaryotes, where they play a critical role in the nonsense-mediated
decay (NMD) RNA surveillance pathway [72, 73]. SF1 helicases are composed of two RecA-like domains
which together harbor a series of signature motifs required for the ATPase and helicase activities,
including the Walker A and Walker B motifs conserved in P-loop NTPases, which are located in the N-
terminal RecA-like domain [71]. Following the Walker A motif in all 4 Type II and Type III CoCoNuT
subtypes, the CnuH helicases contain a large helical insertion, with some of the helices predicted to form
coiled-coils (Figure 6). The Type II, Type III-A, and Type III-B CoCoNuT CnuH helicases contain an OB-fold
domain that, in Type II and Type III-A, is flanked by helices predicted to form a stalk-like helical extension
of the N-terminal RecA-like domain, a structural feature characteristic of the entire UPF1/DNA2-like
helicase family within SF1 [73-75] (Figure 6). DALI comparisons show that the CnuH OB-fold domain in
Type II CoCoNuTs is similar to the OB-fold domain in UPF1 and related RNA helicases SMUBP-2 and
SEN1, and this holds for Type III-A as well, although, in these systems, the best DALI hits are to
translation factor components such as EF-Tu domain II (Supplementary Table S1) [73, 76]. The Type III-B
CnuH OB-folds also match that of UPF1, albeit with lower statistical support (Supplementary Table S1).

In Type II and Type III-A CoCoNuTs, CnuH is fused at the N-terminus to a second OB-fold domain
similar to that of SmpB (small protein B) (Figure 6, Supplementary Figure S4, Supplementary Table S1),
which we denote SPB (SmpB-like). SmpB binds to SsrA RNA, also known as transfer-messenger RNA
(tmRNA), and is required for tmRNA to associate with and rescue stalled ribosomes [77-79]. These OB-
folds also show significant similarity to the tracrRNA/crRNA-interacting oligonucleotide-binding domains
(OBDs) of the Cas12 family CRISPR-associated endonucleases, although typically with weaker support
than to SmpB (Supplementary Figure S4) [80-82]. Conceivably, SmpB-like OB-folds were independently
recruited during the evolution of the complex CoCoNuTs and the transposon-encoded TnpB nucleases,
which also possess a similar OB-fold domain (WED) and are the ancestors of the Cas12 endonucleases
found in Type V CRISPR-Cas systems [83-85]. The N-terminal OB-folds in Type II and III-A CnuH homologs
also resemble prokaryotic HIRAN domains, which are uncharacterized, but have eukaryotic homologs
fused to helicases that bind ssDNA (Supplementary Figure S4) [86]. None of these comparisons is
without caveats. The SPB domain is around 20 amino acids shorter on average than SmpB itself, and a
helix that is conserved in SmpB orthologs and interacts with the tmRNA is absent in the CoCoNuT OB
folds (Supplementary Figure S4) [87]. However, two other structural elements of SmpB involved in
binding tmRNA are present (Supplementary Figure S4) [88]. The Cas12 family endonuclease OBDs are
interrupted by helical inserts, and thus, appear unlikely to share binding specificity and function with the
CoCoNuT domains [80-82]. Nevertheless, the structural similarity suggests an ancient evolutionary
relationship between the tmRNA system, the CoCoNuTs, and TnpBs/Type V CRISPR-Cas systems. The
tmRNA system itself is a type of RNA-guided restriction machinery, in that it recruits RNA degradation
factors such as RNase R to mRNAs bound to stalled ribosomes [89]. The CoCoNuT OB-folds are also
significantly structurally similar to HIRAN domains that, however, do not overlay with the CoCoNuT
domains any better than SmpB (Supplementary Figure S4), lack a small helix that SmpB and the
CoCoNuT domains share, and being DNA-binding, do not fit the other pieces of evidence we gathered
that all suggest an RNA-binding role for this domain in the CoCoNuTs (see below).

The Type II and Type III-A CoCoNuT CnuH helicases contain an additional domain, structurally
similar to the RelE/Colicin D RNase fold [90], which we denote RNase toxin-like (RTL), located between
the SPB OB-fold and the helicase (Figure 6, Supplementary Table S1). The proteins of this family are
ribosome-dependent toxins that cleave either mRNA or tRNA in the ribosomal A-site [91]. This domain
was identified by structural similarity search with the AF2 models, but no sequence conservation with
characterized members of this family was detected, leaving it uncertain whether the RTL domain is an
active nuclease. However, considerable divergence in sequence is not unusual in this toxin family [92,
93]. Notably, Type III-A CoCoNuTs contain another domain with the potential to associate with
ribosomes, the Hsp70-like NBD/SBD (Figure 6, Supplementary Table S1) [55, 94]. In Type III-B CoCoNuTs,
a small potential RNA-binding wHTH domain resembling the archaeal ssDNA-binding protein Sul7s is
fused to the CnuH N-termini (Figure 6, Supplementary Table S1). We did not detect additional domains
fused to CnuH in Type III-C systems (Figure 6).

We searched for homologs of CnuH (see Methods). Our observations of the contextual
associations, both of the CoCoNuTs and their relatives, showed that helicases of this large family are
typically encoded in operons with downstream genes coding for an elongated wHTH domain fused at its
C-terminus to a variety of effectors, generally, nucleases, which we denote CnuE (Figure 3, Figure 4). In
the CoCoNuTs, except for Types III-B and III-C, these CnuE effectors are HEPN ribonucleases, with one
HEPN domain in Type II and two in Type III-A (Figure 6, Supplementary Table S1). In Type III-B, the
effector is a Vsr (very-short-patch repair)-like PD-(D/E)xK family endonuclease [95] fused directly to the
helicase, with no wHTH domain present, whereas in Type III-C, a distorted version of the elongated
wHTH domain is fused to two Phospholipase D (PLD) family endonucleases (Figure 3, Figure, 4, Figure 6,
Supplementary Table S1). All these nucleases can degrade RNA, and some, such as HEPN, have been
found to cleave RNA exclusively [54, 96-99]. We also detected coiled-coils in the region between the
wHTH and effector domains in Type II, Type III-A, and Type III-C CoCoNuT CnuE homologs. Multimer
structural modeling with AlphaFold2 suggests that the wHTH domain in CnuE might interact with the
coiled-coil-containing helical insertion of CnuH, perhaps mediated by the coiled-coils in each protein, to
couple the ATP-driven helicase activity to the various nuclease effectors (Supplementary Figure S5). The
accuracy of this model notwithstanding, the fusion of the Vsr-like effector to CnuH in Type III-B
CoCoNuTs and the similarity of this system to the other CoCoNuT types strongly suggests that the CnuE
effector proteins in these systems form complexes with their respective CnuH helicases. An additional
factor in potential complexing by these proteins is the presence of coiled-coils in the associated
CnuB/McrB and CnuC/McrC homologs, which may interact with the coiled-coils in CnuH and CnuE. Type
III-B CoCoNuTs also code for a separate coiled-coil protein, which we denote CnuA, as it resembles the
CnuA protein encoded in Type I-B CoCoNuTs (Figure 3, Figure 4). However, it is distinguished from Type
I-B CnuA in containing no recognizable domains other than the coiled-coil (Figure 3, Figure 4). This also
can potentially interact with other coiled-coil proteins in the system.

**Figure 6: Domain composition, operon organization, and AlphaFold2 structural predictions for core protein components of Type II and III CoCoNuT systems**

A) Type II and III CoCoNuT domain composition and operon organization. The arrows indicate the direction of transcription. Type II and III-A systems very frequently contain TerY-P systems as well, but not invariably, and these are never found in Type III-B or III-C, thus, we do not consider them core components. B-D) High quality (Average pLDDT > 80), representative AlphaFold2 structural predictions for proteins in B) Type II CoCoNuT systems, C) Type II and III-A CoCoNuT systems, and D) Type III-A CoCoNuT systems. Models were generated from representative sequences with the following GenBank accessions (see Supplementary Data for sequences and locus tags): AMO81401.1 (Type II CnuB), AVE71177.1 (Type II CnuC), AMO81399.1 (Type II and III-A CoCoNuT CnuH), AVE71179.1 (Type II CnuE), ATV59464.1 (Type III-A CnuE), PSR44427.1 (Type III-A CnuB), NMY00740.1 (Type III-A CnuC). Abbreviations of domains: CSD – cold shock domain; CoCo – coiled-coil; IG – Immunoglobulin (IG)-like beta-sandwich domain; ZnR – zinc ribbon domain; SPB – SmpB-like domain; RTL – RNase toxin-like domain; HEPN – HEPN family nuclease domain; OB/stalk – OB-fold domain attached to a helical stalk-like extension of ATPase; Vsr – very-short-patch-repair PD-D/ExK nuclease-like domain; PLD – Phospholipase D family nuclease domain; HEAT – HEAT-like helical repeats. These structures were visualized with ChimeraX [40].

Type II and Type III-A CoCoNuTs, the most widespread varieties apart from Type I, also include conserved genes coding for a "TerY-P" triad. TerY-P consists of a TerY-like von Willebrand factor type A (VWA) domain, a protein phosphatase 2C-like enzyme, and a serine/threonine kinase (STK) fused at the C-terminus to a zinc ribbon (Figure 3, Figure 4, Supplementary Figure S2, Figure 6, Supplementary Table S1). TerY-P triads are involved in tellurite resistance, associated with various predicted DNA restriction and processing systems, and are predicted to function as a metal-sensing phosphorylation-dependent signaling switch [7]. In addition, TerY-P-like modules, in which the kinase is fused at the C-terminus to an OB-fold rather than a zinc ribbon, have been recently shown to function as stand-alone antiphage defense systems [2]. The OB-fold fusion suggests that this kinase interacts with an oligonucleotide and raises the possibility that the zinc ribbon, which occupies the same position in the CoCoNuTs, is also nucleic acid-binding. Almost all CoCoNuT systems containing *cnuHE* operons also encompass TerY-P,

with a few exceptions among Terrabacterial Type II systems and Myxococcal Type III-A systems, implying
an important contribution to their function (Supplementary Data). However, the complete absence of
the TerY-P module in Type III-B and Type III-C systems implies that when different nucleases and other
putative effectors fused to the helicase are present, TerY-P is dispensable for the CoCoNuT activity.
Therefore, we do not consider them to be core components of these systems. Below, we describe
experimental results showing that TerY-P in Type II CoCoNuT can indeed function independently to
protect *E. coli* from λ-vir infection partially, but that the other factors are required for full-scale survival.

Type II and Type III-A CoCoNuTs are generally similar, but a more detailed comparison reveals
substantial differences. The CnuE proteins in Type II contain one HEPN domain with the typical RxxxxH
RNase motif conserved in most cases, whereas those in Type III-A contain two HEPN domains, one with
the RxxxxH motif, and the other, closest to the C-terminus, with a shortened RxH motif. Furthermore,
Type III-A CnuB/McrB GTPase homologs often contain C-terminal HTH-domain fusions absent in Type II
(Figure 6). Finally, Type II CnuC/McrC homologs resemble Type I-A CnuCs, which contain N-terminal
immunoglobulin-like beta-sandwich domains, insertions of coiled-coils and a CSD-like OBD into
DUF2357, PD-D/ExK nucleases, and zinc ribbon domains, except that in Type II CnuCs, this domain
architecture underwent reductive evolution (Figure 5, Figure 6, Supplementary Table S1). In particular,
the beta-sandwich domain and coiled-coils are shorter, the CSD was lost, the nuclease domain was
inactivated, and in many cases, the number of Zn-binding CPxC motifs was reduced from three to two
(Figure 5, Figure 6, Supplementary Table S1). This degeneration pattern could indicate functional
replacement by the associated CnuH and CnuE proteins, often encoded in reading frames overlapping
with the start of the *cnuBC* operon.

By contrast, Type III-A CnuC/McrC homologs entirely lost the beta-sandwich domain, PD-D/ExK
nuclease, and zinc ribbon, but gained an Hsp70-like NBD/SBD unit similar to those that are fused to the
McrB GTPase in early branching members of the NxD clade, as well as a helical domain similar to the
HEAT repeat family, and in some cases, a second CSD (Figure 5, Figure 6, Supplementary Table S1). Most
Type III-A CnuCs contain a CSD and coiled-coils, and thus, resemble Type I-A, but the positioning of these
domains, which in Type I-A and Type II are inserted into the DUF2357 helix bundle, is not conserved in
Type III-A, where these domains are located outside DUF2357 (Figure 5, Figure 6). The further addition
of the Hsp70-like NBD/SBD and HEAT-like repeat domains accentuates the distinct organization of Type
III-A (Figure 5, Figure 6, Supplementary Table S1). Moreover, some *Pseudomonas* genomes encode both
Type II and Type III-A CoCoNuTs, implying distinct targets. As mentioned above, the Type II CnuH
helicases also differ from Type III-A in that their stalk-attached OBDs more closely resemble UPF1-like
helicases, whereas the OBDs in this position in Type III-A are more similar to EF-Tu domain II, a
divergence potentially linked to the number of HEPN domains (1 vs. 2) in the closely associated CnuE
proteins (Figure 6, Supplementary Table S1).

Hsp70-like NBD/SBD domains and HEAT-like repeats are fused to the CnuC/McrC N-terminal
DUF2357 domain in Type III-A CoCoNuT, but their homologs in Type III-B and III-C are encoded by a
separate gene. We denote these proteins CnuD and CnuCD, the latter for the CnuC-CnuD fusions in Type
III-A. The separation in Type III-B and III-C implies that fusion is not required for their functional
interaction with the CnuBC/McrBC systems (Figure 6, Supplementary Table S1). CnuD proteins
associated with both Type III-B and III-C contain predicted coiled-coils, suggesting that they might
interact with the large coiled-coil in the CnuB homologs, which would not be unexpected, given that
similar domains are fused to closely related McrB GTPases (Figure 3, Supplementary Figure S2). Because

Hsp70 homologs have an affinity for AU-rich RNA, and in some cases, such as the bacterial DnaK, C-rich RNA, and given the strong predictions of RNA interactions for the other components of the CoCoNuTs, we hypothesize that the CnuD homologs present in all Type III CoCoNuT subtypes and fused to the NxD McrB GTPase also bind RNA [56, 57]. As cited above, many Hsp70 homologs have been reported to associate with ribosomes [55, 94], which could also be true for the CoCoNuT CnuDs. Hsp70 ATP binding and hydrolysis allosterically regulate the affinity of Hsp70s for peptide and protein substrates, such that the ATP-bound state has low affinity for misfolded proteins, but the ADP-bound protein has high affinity, and ATP hydrolysis "traps" protein substrates by generating a high affinity ADP-bound complex, after initially contacting the substrates in the ATP-bound form [55]. To our knowledge, the effects of ATP binding and hydrolysis on RNA-binding by Hsp70 homologs were investigated in one study only, which reported that RNA binding was independent of nucleotide binding, ATP hydrolysis, and unfolded peptide interaction [56]. In the CnuD homologs found in the CoCoNuTs and fused to NxD McrB GTPases, Walker B-like motifs [100] are usually, but not invariably, conserved, whereas the sequences of the helical domains just adjacent to the motifs are more strongly constrained. Walker A-like motifs [101] are present but degenerate, with the DxGT(T/S) signature motif mutated, although the initial aspartate is conserved in all except the Type III-C CnuD homologs (Supplementary Figure S6). It appears likely that the CoCoNuT CnuD homologs bind ATP/ADP but hydrolyze ATP with extremely low efficiency, at best. Such properties are better compatible with RNA binding than unfolded protein binding or remodeling, suggesting that these domains target the respective systems to aberrant RNA.

Consistent with this prediction, Type II and III-A CoCoNuTs generally appear likely to target RNA rather than DNA, given that the respective operons encode HEPN RNases, typically the only recognizable nuclease in these systems. All CnuC/McrC homologs in Type II or III CoCoNuT systems lack the PD-D/ExK catalytic motif that is required for nuclease activity, although for Type II and Type III-B and III-C, but not Type III-A, structural modeling indicates that the inactivated nuclease domain was retained, likely for a nucleic acid-binding role (Figure 6, Supplementary Table S1). In Type III-A CoCoNuT, the nuclease domain was lost entirely and replaced by the Hsp70-like NBD/SBD domain with RNA-binding potential (Figure 6, Supplementary Table S1). Often, one or two CSDs, generally RNA-binding domains, although capable of binding ssDNA, are fused to Type III-A CnuC homologs as well (Figure 6, Supplementary Table S1) [102]. RNA targeting capability of Type III-B and III-C can perhaps be inferred from the similarity to Type III-A. Moreover, higher-order associations of Type II and Type III-A CoCoNuT systems with various DNA restriction systems suggest a two-pronged DNA and RNA restriction strategy reminiscent of Type III CRISPR-Cas (see below).

We suspect that RNA targeting is an ancestral feature of the CoCoNuT systems. Several observations are compatible with this hypothesis:

1) CSD-like OB-folds are pervasive in these systems, being present in the CnuB/McrB homologs of all Type I subtypes and in Type I-A and Type III-A CnuC/McrC homologs. As previously noted, these domains typically bind RNA, although they could bind ssDNA as well.

2) YTH-like domains are present in most Type I systems, particularly, in almost all early branching Type I-B and I-C systems, suggesting that the common ancestor of the CoCoNuTs contained such a domain. YTH domains in eukaryotes sense internal N6-methyladenosine (m6A) in mRNA [66, 67, 103].

3) Type II and III-B/C CoCoNuTs, which are likely to target RNA, given their genomic context
encoding HEPN proteins and Hsp70 NBD/SBD1 homologs, retain inactivated PD-DxK nuclease domains,
which suggests these domains contribute an affinity for RNA inherited from Type I-A CoCoNuTs. PD-
(D/E)xK nucleases are generally DNA-specific, however, some examples of RNase activity have been
reported [97, 98]. The inactivated PD-DxK domains might also bind DNA from which the target RNA is
transcribed.

*Type II CoCoNuT protects bacteria against plasmids with CloDF13 ori and λ-vir phage*

To probe whether CoCoNuTs can defend bacteria against MGE, we engineered heterologous *E.*
*coli* strains carrying Type I-A, Type II, or Type III-C CoCoNuT systems on a low-copy pACYC plasmid
(Figure 7 A) and challenged them with different phages and plasmids.

Initially, we monitored phage restriction in *E. coli* strains DH5α, MG1655, and BL21(DE3)
carrying CoCoNuT systems using a classical double-layer agar assay and a set of 22 *E. coli* phages
spanning five families: *Autographiviridae* T3, T7; *Demerecviridae* T5, Lau5; *Drexlerviridae* NBD2;
*Siphoviridae* λ-vir, Ervi20; *Myoviridae* T2, T4, T6, and other T-even-like phages (Supplementary Table S2).
This assay did not reveal phage restriction or plaque size decrease, indicating no phage defense activity
of CoCoNuT (Supplementary Figure S7 A, B). The Type I-A CoCoNuT expression in *E. coli* BL21(DE3)
triggered toxicity so that the bacterial lawn did not form in soft agar (Supplementary Figure S7 A, B).

Next, we tested the antiphage activity of the CoCoNuTs in *E. coli* MG1655 and BL21-AI strains in
a liquid medium (Figure 7 B, Supplementary Figure S7 C, D). In BL21-AI, Type II CoCoNuT increased
bacterial survival under λ-vir infection, while for Type I-A CoCoNuT, less pronounced bacterial survival
was observed (Figure 7 B, Supplementary Figure S7 D). In the MG1655 strain, only Type II CoCoNuT
enabled bacterial survival under phage infection (Figure 7 B, Supplementary Figure S7 C). Given that the
phage resistance phenotype was reliably replicated for Type II CoCoNuT and λ-vir phage in both strains
(Figure 7 B), we further focused on this CoCoNuT type.

As described above, the Type II CoCoNuT locus includes three genes homologous to a stand-
alone TerY-P defense system that has been shown to provide resistance against T3, T7, and φ-V1 phages
[2]. To probe whether the TerY-PP2C-STK genes from Type II CoCoNuT can provide antiphage activity,
we cloned only these genes in the same pACYC vector and used the *E. coli* BL21-AI strain to monitor cell
survival upon λ-vir infection in liquid medium (Figure 7 B, C, D). The cells showed increased survival
compared to control, but the effect was weaker compared to the full-length Type II CoCoNuT system,
indicating that TerY-PP2C-STK alone is insufficient to provide full-scale survival (Figure 7 B, D).
Furthermore, we compared the number of colony-forming units (CFU) in cultures without CoCoNuT,
with wild-type Type II CoCoNuT, or with the CoCoNuT TerY-PP2C-STK variant after 4 hours of λ-vir
infection (Figure 7 C, D). Cells carrying an empty vector did not yield any colonies, while the cells
carrying Type II CoCoNuT or CoCoNuT TerY-PP2C-STK survived under λ-vir infection, albeit to different
extents (Figure 7 D). Under the same conditions, full-length Type II CoCoNuT cells formed 100-fold more
CFUs than the TerY-PP2C-STK variant (Figure 7 D). In a separate set of experiments, we found that λ-vir
phage formed turbid plaques on Type II CoCoNuT bacteria lawn, indicating that a fraction of the *E. coli*
population survives during the infection (Supplementary Figure S7 E). This finding could explain why
initially we did not observe a classic phage resistance phenotype in the plaque assay (Supplementary
Figure S7 A, B).

We next tested whether heterologous *E. coli* strains carrying CoCoNuTs can provide plasmid
interference. In this experiment, we used *E. coli* MG1655 strains with CoCoNuTs or a control plasmid and
electroporated a set of plasmids with different *ori* sequences (Supplementary Table S3). We plated serial
dilutions of the cells on a single antibiotic to select for a CoCoNuT or on two antibiotics, to select for
both a CoCoNuT and an invading plasmid that carried a distinct antibiotic resistance gene (Figure 7 E).
When Type II CoCoNuT was co-electroporated with the pCDF plasmid containing CloDF13 *ori* sequence,
decreased cell viability and smaller colonies were observed, indicative of plasmid interference mediated
by the CoCoNuT (Figure 7 F). Plasmid interference was detected only for pCDF plasmids suggesting that
CoCoNuT can restrict plasmids carrying a specific *ori* sequence.

Taken together, our experimental results show that the Type II CoCoNuT system protects *E. coli*
against λ-vir infection in a liquid medium and impairs the retention of pCDF plasmid. Our findings further
suggest that similarly to the previously characterized TerY-P system, the TerY-P module of Type II
CoCoNuT has an antiphage activity of its own, but that other CoCoNuT components are required for the
full antiphage effect. Clearly, these are only the proof of principle experiments demonstrating that at
least some of the CoCoNuTs are bacterial defense systems. The basis of the CoCoNuT specificity, the
functionalities of the other types of CoCoNuTs, and the mechanistic details of the CoCoNuT activities
remain to be elucidated, but the additional bioinformatic analysis we describe below allows some
plausible and testable hypotheses to be proposed.

**Figure 7: Type II CoCoNuT protects *Escherichia coli* against plasmids and λ-vir phage**

A) Gene organization of the CoCoNuT systems selected for experimental analysis. In the case of the Type II CoCoNuT
system, for optimal gene synthesis, 5 bp were changed, preserving the amino acid sequence (see Methods). B) Phage λ-vir
challenge in a liquid medium for Type II CoCoNuT or TerY-PP2C-STK gene module from Type II CoCoNuT in different *E. coli*
strains. The cells were infected at MOI 0.05. Top – infection of *E. coli* MG1655 strain with empty vector or wild-type Type II
CoCoNuT system (5 independent colonies selected, each picked twice, for 10 replicates in total). Middle – infection of *E.
coli* BL21-AI strain with empty vector or with wild-type Type II CoCoNuT system (in the case of the Type II CoCoNuT variant,
4 independent colonies, each picked twice, for 8 replicates in total; otherwise, as described for the MG1655 strain).
Bottom – infection of *E. coli* BL21-AI strain with empty vector or wild-type TerY-PP2C-STK gene triad from Type II CoCoNuT
(3 independent replicates). The shaded area shows standard deviation. C) Schematic representation of the experimental
setup to evaluate cell viability upon λ-vir phage infection. Diluted overnight culture with control or CoCoNuT plasmids
were infected with λ-vir phage at MOI 0.05. After inoculation for 4 h, the phage was removed by centrifugation, and cells
were plated by serial dilutions to compare the viability of different variants. D) Viability of cells carrying Type II CoCoNuT or
TerY-PP2C-STK gene module after infection with λ-vir phage. Colony forming units per milliliter of culture (CFU/ml) were

calculated after 4 hours of λ-vir phage infection in *E. coli* cultures harboring vector control (no colonies formed), wild-type
Type II CoCoNuT system, or Type II CoCoNuT TerY-STK gene triad only. Three independent colonies were used for three
replicates; the black line indicates the median. E) Schematic representation of plasmid interference assay. Heterologous *E.*
*coli* cells harboring Type I-A, Type II, or Type III-C CoCoNuT systems in pACYC184 plasmid or empty pACYC184 were co-
transformed with plasmids containing different *ori* sequences (Supplementary Table S2), and cell viability was evaluated
on agar plates by performing serial dilutions. F) pCDF plasmid co-electroporation into heterologous *E. coli* hosts expressing
CoCoNuT systems. Cell viability was evaluated by serial dilutions in the presence of chloramphenicol to select CoCoNuT
systems in pACYC184 or in the presence of both chloramphenicol and streptomycin (for pACYC184 and co-transformed
pCDF). Type II CoCoNuT showed ~2-fold interference with pCDF. Data with other plasmids are shown in Supplementary
Figure S8.

*CoCoNuT, CARF, and RtcR: A convergence on tmRNA?*

Genomic neighborhoods of many Type II and Type III-A CoCoNuTs encompass complex operonic
associations with genes encoding several types of CARF (CRISPR-Associated Rossmann Fold) domain-
containing proteins. This connection seems to suggest multifarious regulation by cyclic (oligo)nucleotide
second messengers synthesized in response to viral infection and bound by CARF domains [104-106].
Activation of an effector, most often a nuclease, such as HEPN or PD-(D/E)xK, by a CARF bound to a
cyclic (oligo)nucleotide is a crucial mechanism of CBASS (cyclic oligonucleotide-based antiphage signaling
system) as well as Type III CRISPR-Cas systems [104-106]. These CARF-regulated enzymes generally
function as a fail-safe that eventually induces PCD/dormancy, if other antiphage defenses fail to bring an
infection under control, and are deactivated, typically through cleavage of the second messenger by a
RING nuclease, if other mechanisms succeed [104, 107-109].

The presence of CARFs could implicate the HEPN domains in these systems as PCD effectors that
would carry out non-specific RNA degradation in response to infection. Surprisingly, however, most of
these CARF domains showed the highest similarity to the CARF domain of RtcR, a sigma54
transcriptional coactivator of the RNA repair system RtcAB with a CARF-ATPase-HTH domain
architecture, suggesting an alternative functional prediction (Supplementary Figure S9, Supplementary
Table S1). Specifically, by analogy with RtcR, CoCoNuT-associated CARF domain-containing proteins
might bind (t)RNA fragments with cyclic phosphate ends [110, 111]. This interaction could activate
transcription initiation of downstream genes, in this case, genes encoding CoCoNuT components,
through binding an upstream activating sequence by the HTH domain fused to the CARF-ATPase C-
terminus [110, 111]. The manifold effects of tRNA-like fragments are only beginning to be appreciated.
Lately, it has been shown that bacterial anticodon nucleases, in response to infection and DNA
degradation by phages, generate tRNA fragments, likely a signal of infection and a defensive strategy to
slow down the translation of viral mRNA, and that phages can deploy tRNA repair enzymes and other
strategies to counteract this defense mechanism [112-115].

In the RtcAB system, tRNA fragments with 2', 3'-cyclic phosphate ends, which could be produced
as a consequence of stress-related damage and anticodon nuclease activity, bind to the CARF domain of
RtcR, leading to its oligomerization and activation of *rtcAB* operon expression [111, 116]. However, a
recent study found that, although the CARF domain of RtcR does bind 2', 3'-cyclic phosphate tRNA
fragments, full-length RtcR did not stably associate with these tRNA fragments that were insufficient for
RtcR to initiate transcription *in vitro* [110]. Thus, the exact identity of the ligands of these CARFs remains
uncertain, but a consensus that they bind 2' 3' cyclic phosphate RNA is emerging [110, 111]. RtcA is an

RNA cyclase that converts 3ʹ-phosphate RNA termini to 2ʹ, 3ʹ-cyclic phosphate and thus, in a feedback
loop, generates 2', 3'-cyclic phosphate RNA fragments that induce expression of the operon [111, 116,
117]. RtcB is an RNA ligase that joins 2', 3'-cyclic phosphate RNA termini to 5'-OH RNA fragments,
generating a new 5'-3' bond and, in many cases, reconstituting a functional tRNA [118]. In eukaryotes,
RtcB homologs have important roles in joining pre-tRNA halves after intron splicing and in the unfolded
protein response [119, 120], and in bacteria, compelling recent work has shown that the most frequent
target of RtcB is SsrA, the tmRNA [110].

The tmRNA, a unique hybrid molecule that combines a tRNA-like domain (TLD) with an internal
ORF in an mRNA-like region (MLR), might unexpectedly be key to cracking the CoCoNuTs and elucidating
their functions [78]. In all bacteria and some mitochondria, but not archaea or eukaryotes, where this
system does not appear to exist, tmRNA binds to the SmpB protein [78]. This interaction is essential for
targeting the complex to stalled ribosomes, a type of translation stress [78]. In response to these
irregular translation events, often caused by mRNAs missing a stop codon, tmRNA switches translation
by the stalled ribosome to the ORF in the tmRNA MLR, which encodes a small peptide tag [79].
Translation of this ORF fuses the tag to the C-terminus of the nascent polypeptide, marking it for
proteolysis, and releases the ribosome when it reaches the MLR stop codon [89].

In both Type II and III-A CoCoNuTs, the SmpB-like SPB OB-fold (Figure 6, Supplementary Figure
S4) fused at the N-termini of the CnuH helicases might bind tmRNA, which is targeted for repair by RtcB,
itself regulated by the same family of RtcR homologs as the CoCoNuT systems. Given this close kinship,
these CoCoNuTs might respond to the same stimuli as *rtcAB*, with which they are often present in the
same genomes. If both genes in the operon are activated, RtcA could provide RNA fragments with 2', 3'-
cyclic phosphate ends that would promote CoCoNuT transcription. Concurrently, RtcB could regenerate
tmRNA that might enable the localization of CoCoNuT RNA degradation complexes to mRNAs associated
with stalled ribosomes. A search against the database of predicted proteins from completely sequenced
prokaryotic genomes, described above, showed that 31% of those that encoded a Type II or III-A
CoCoNuT also contained a typical RtcRB locus (RtcA is not always present in these operons). Although
this co-occurrence was less common than expected, our search was not exhaustive, and furthermore,
we observed several cases when an RtcB homolog was encoded within or nearby the CoCoNuT operon
itself (Supplementary Figure S10). Regardless of the co-occurrence with RtcRAB, tmRNA is present in any
bacterium encoding CoCoNuTs because this RNA is (nearly) universal in bacteria, while missing in
archaea, which conspicuously lack CoCoNuTs as well.

We performed phylogenetic analysis of confidently identified ATPase domains of CnuH and
UPF1-like helicase homologs obtained by a search against the completely sequenced prokaryotic
genomes described above, as well as a search against a set of 72 representative eukaryotic genomes
that were downloaded from the NCBI GenBank in June 2020. The topology of the resulting phylogenetic
tree indicated that the CoCoNuT CnuH helicases are more closely related to the UPF1-like helicases than
other members of this family, such as DNA2, which is fused to an AddB-like nuclease and acts on ssDNA
[74] (Figure 8 A). As detailed above, in addition to the ATPase domain, these helicases also share OBDs
attached to stalk-like helical domains (Figure 6). In UPF1, this OBD interacts with RNA 3' ends [73]. These
domains in the CoCoNuT CnuHs are most structurally similar to either the corresponding domain in
UPF1 or to OB-fold domains of translation factors, such as EF-Tu and ribosomal protein L3
(Supplementary Table S1). The tmRNA forms a ribonucleoprotein complex with SmpB, EF-Tu, and
ribosomal protein S1 [77], and S1-like CSDs are present in the CnuC/McrC homologs of Type III-A

CoCoNuTs (Figure 6). Therefore, if the CoCoNuT helicases indeed bind tmRNA, the multiple OB-folds
structurally similar to tmRNA-binding domains could compensate for the absence of some of the
structural elements of SmpB in the CoCoNuT SPB domains. Additionally, the ATPase domain of UPF1
contains a helical insert, which is required for RNA binding, in the same position, following the Walker A
motif, as the much larger helical insert in the CoCoNuT CnuH helicases, which might also bind RNA
(Figure 6) [72].

Emerging from this phylogeny, and in accord with the hypothesis of tmRNA interaction with the
Type II and Type III-A CoCoNuT CnuH helicases, we observed clear associations between close relatives
of the CoCoNuTs and SlpA homologs that catalyze recombination involving tmRNA genes (Figure 8 B).
The SlpA proteins are P4 integrase homologs characterized as factors required for the excision of a P4-
like cryptic prophage, CP4-57, in *E. coli* K-12 [121]. This excision disrupts the tmRNA gene *ssrA* that
overlaps with the *attL* integration site and removes at least several kilobases of the downstream
sequence [121]. The presence of CnuH-like helicases encoded adjacent to *slpA* genes implies that
excision of the cryptic prophage would also delete the helicase gene while simultaneously inactivating
the tmRNA, establishing a direct connection between the functioning of the tmRNA system and
expression of the helicase. Encoded immediately upstream of these loci is the tmRNA-binding protein
SmpB (Figure 8 B). The CnuH-like helicases in these neighborhoods share many features with those
found in Type II and III-A CoCoNuTs, including the ATPase, the OBD/stalk domain, the helical insert
following the Walker A motif with coiled-coils, and the RelE family RNase toxin-like (RTL) domain. At the
N-termini, these helicases are fused to domains with a serine-threonine kinase (STK)-like fold
(Supplementary Table S1). These are similar to the kinase homology (KH) domains found in RNase L, a
eukaryotic antiviral HEPN superfamily endoribonuclease that, notably, is activated by oligoadenylate
second messengers [122, 123] (Figure 8 B, Supplementary Table S1). However, whereas the RNase L KH
domains are inactivated pseudokinases, we predict the domains in the CnuH homologs to be active
kinases [122]. Like their CoCoNuT CnuH relatives, these helicases occur in operons with CnuE-like
predicted effectors containing a wHTH domain fused to a PLD-type nuclease (Figure 8 B, Supplementary
Table S1). In some CnuH homologs, more distantly related to the CoCoNuTs, there is an additional N-
terminal NERD (nuclease-related domain) with a D-ExK nuclease motif, and two tandem kinase domains
instead of one (Figure 8 B). In the associated CnuE-like effector, the PLD nuclease is likely inactivated
(Figure 8 B, Supplementary Table S1). These CnuHE-like tandems, AlpA, and SlpA homologs are adjacent
to tyrosine tRNA operons that express RtT sRNAs rather than tmRNA genes (Figure 8 B). This genomic
context implies that these loci are distinct cryptic prophages that integrate into the RtT sRNA gene.
Intriguingly, RtT is another type of RNA that regulates a stress response, in this case, to starvation and
antibiotics, distinct from the tmRNA translation stress response mechanism [124-126]. These tmRNA
and RtT sRNA-associated CnuH-like helicases are among the most closely related to the CoCoNuTs in our
phylogenetic tree of the UPF1-like family (Figure 8 A, B).

**Figure 8: Type II and III-A CoCoNuT helicases are close relatives of UPF1-like RNA helicases, and CoCoNuT homologs are associated with SmpB/SsrA(tmRNA) genes**

A) Phylogenetic analysis of the UPF1-like helicase family and CnuH. B) Neighborhood analysis of the closest relatives of the CoCoNuT helicases shows association with SlpA prophage integrase homologs that integrate into tmRNA and RtT sRNA genes in *E. coli*. The broken line for *E. coli* NC_017633.1 indicates a region omitted for clarity that contains several recombination-related genes. C) Type I-B CoCoNuTs and the closely related Pseudo-CoCoNuTs can be found adjacent to tmRNA and SmpB genes in *Clostridium*. Abbreviations of domains: SPB – SmpB-like domain; RTL – RNase toxin-like domain; OB/stalk – OB-fold domain attached to a helical stalk-like extension of ATPase; STK – serine/threonine kinase-like domain; PLD – Phospholipase D family nuclease domain; SmpB – Small protein B; SsrA – tmRNA gene; SlpA – P4 integrase homolog; RtT – tRNA-derived ncRNA; AlpA – A prophage transcription factor that stimulates expression of SlpA homologs; Abi_2 – Abi-like HEPN domain-containing protein; BaeS – Signal transduction histidine kinase; TPR – Tetratricopeptide repeat protein; PAS – PAS fold signal transduction domain; iPLD? – predicted inactivated PLD nuclease fold; Ub-RnfH – RnfH family ubiquitin; START – START domain-containing protein. Ub-RnfH and START are frequently associated with SmpB/tmRNA in bacteria, suggesting they play a role in the tagging and proteolysis of targeted proteins, similar to how ubiquitin family proteins are involved in this process in eukaryotes [127].

These observations suggest an evolutionary scenario in which SmpB became fused at the N-terminus of a common ancestor of the Type II CoCoNuT CnuH helicases and their closest relatives, which still associate with SmpB (Figure 8 B). This hypothetical fusion protein could have retained the tmRNA-binding capacity of SmpB. The operon encoding this CnuH-like helicase and its associated CnuE-like effector, which we can infer to have been a HEPN domain-containing protein, could then have become associated with a Type I-A CoCoNuT system, likely also RNA-targeting, as well as a TerY-P operon, to give rise to the Type II CoCoNuTs, after reductive evolution of Type I-A CoCoNuT (see above). Conceivably, this combination could have enabled the highly efficient degradation of both aberrant mRNAs and their nascent proteins, an effective antiviral and general stress response, but perhaps one that comes with a substantial cost to the host. Regulation by RtcR homologs could have been selected to confer additional precision in deploying this potent mechanism. Type III CoCoNuTs and CoCoPALMs seem to have evolved subsequently from Type II CoCoNuTs through a complex evolutionary process. Thus, the CoCoNuT CnuH helicases appear to be but one branch of a large helicase family associated with RNA restriction systems, including NMD and tmRNA/SmpB/RNase R, many of which likely act at the level of translation and, in prokaryotes, often mobilize via prophages which integrate into tmRNA and tRNA genes.

Further support for this hypothesis seems to come from the contextual associations of Type I-A
CoCoNuTs in Gammaproteobacteria and Type I-B CoCoNuTs and Pseudo-CoCoNuTs in *Clostridium,* which
are often encoded near the tmRNA gene *ssrA* and SmpB (Figure 8 C, D, Supplementary Table S1). Type I-
A CoCoNuTs, which possess two CSD-like OB-folds, one each in CnuB/McrB and CnuC/McrC (Figure 3,
Figure 5), can be found associated not only with SmpB/tmRNA, but also with SlpA and AlpA homologs,
implying these are also capable of mobilization via prophages that target the tmRNA gene (Figure 8 C,
Supplementary Table 1). In two examples, the Type I-A CoCoNuT is also tightly associated with a Type I
RM system (Figure 8 C). The Type I-B and Pseudo-CoCoNuTs have contextual connections to additional
factors predicted to be involved in defensive and stress-related signaling, such as alarmone-interacting
phosphoribosyltransferases, PAS domain-containing proteins, and HEPN domain-containing Abi-like
proteins (Figure 8 D, Supplementary Table S1) [128-130]. The genomic environments of these Type I-A
and I-B/Pseudo-CoCoNuT systems suggest they might be involved in the tmRNA-directed degradation of
aberrant and viral mRNA (Figure 8 C, D). In Type I-A, the two CSD-like OB-folds might interact with
tmRNA, and in Type I-B, the CSD and YTH domains fused to the CnuB/McrB GTPase domains in these
systems might interact with modified tmRNA.

In light of these observations, the origins and roles of the CnuBC systems in Type II and III
CoCoNuTs, which have lost the OB-fold domains, begin to come into focus. Both the Type I-A CoCoNuTs
and the CnuHE-like systems that are the closest relatives of the Type II/III CoCoNuTs can be found as
apparent cargo of P4-like prophages that integrate into tmRNA genes (Figure 8 B, C). Therefore, it can be
confidently presumed that the CnuHE and Type I-A CoCoNuT operons could have come into association
with each other, adding credence to the scenario of the origins of Type II CoCoNuTs proposed above.
The reductive evolution of CnuBC in Type II CoCoNuT can be explained by the observation that both
CnuHE-like systems and Type I-A CoCoNuT systems associate with SmpB/tmRNA. If Type I-A CoCoNuTs
interact with tmRNA and can associate with CnuHE homologs, then, the capture of an SmpB-like tmRNA-
binding domain by the associated CnuHE system would make redundant the OB-fold domains of the
Type I-A CoCoNuT system. Moreover, the nuclease activity of CnuC could also have become dispensable
if the RNase activity of a HEPN domain in CnuE was now directed by the tmRNA interaction.

The present role of CnuBC in Type II CoCoNuTs can perhaps be predicted from domains that
remain: the inactivated PD-D/ExK nuclease, potentially descending from an ancestral Rnase, and the
coiled-coil fused to CnuB, which is much longer than the one fused to CnuB in Type I-A CoCoNuT. The
retention of these domains implies an RNA-binding function for Type II CnuC and a gain of function for
the coiled-coil in Type II CnuB, likely, complex formation with CnuHE, which allowed the reductive
evolution to take place. The role of the CnuB GTPase domain remains somewhat enigmatic. This domain
is believed to translocate DNA in canonical McrBC systems, therefore translocation of RNA by the
CoCoNuTs is a possibility [25, 44]. The GTPase domain also assembles into hexamers upon binding GTP,
and this could be the primary reason for the conservation of CnuBC in Type II CoCoNuT [131]. This
trigger for complex formation could be important for a system interacting with the translation
apparatus, which is also GTP-dependent [132-134]. Depletion of GTP by CnuBs associated with
ribosomes might also contribute to slowing translation elongation and allowing time for RNA restriction
and proteolysis to occur.

*Complex higher-order associations and functions of the CoCoNuT systems*

Although the tmRNA-associated CoCoNuT activity proposed in the preceding section would be a
distinct antivirus mechanism, there are intriguing parallels with Type III CRISPR-Cas systems, where the
Cas10-Csm-crRNA effector complex binds phage RNA complementary to the spacer of the crRNA,
triggering both restriction of phage DNA and indiscriminate cleavage of RNA [105]. The target RNA
recognition stimulates the production of cyclic oligoadenylate (cOA) signals by Cas10, which activate, via
CARF domain binding, PCD effectors, typically HEPN domain-containing proteins, such as Csm6, that
function as promiscuous RNases [54, 135]. One of the two outcomes can result from this cascade: the
infection can either be abrogated quickly due to the restriction of the virus DNA, which inhibits cOA
signaling via the depletion of viral RNA, along with the activity of RING nucleases, thus averting PCD, or
the continued presence of viral RNA stimulates cOA signaling until PCD or dormancy occurs, limiting the
spread of viruses to neighboring cells in the bacterial population [5, 107, 109].

In CoCoNuTs associated with RtcR homologs, a conceptually similar but mechanistically distinct
phenomenon might occur. Although many of these CoCoNuTs only contain an appended gene encoding
a CARF domain protein at the 5' end of the predicted operon (Supplementary Figure S9), there are also
numerous cases where several types of DNA restriction systems are encoded between the CARF gene
and the CoCoNuT (Figure 9). In these cases, nearly all genes are in an apparent operonic organization
that can extend upwards of 40 kb, such that they could all be transcribed as a single, giant polycistronic
mRNA (Figure 9). Although internal RtcR-independent promoters likely exist in these large loci, the
consistent directionality and close spacing of the genes in these superoperons suggests coordination of
expression. The complex organization of the CARF-CoCoNuT genomic regions, and by implication, the
corresponding defense mechanisms, might accomplish the same effect as Type III CRISPR-Cas, contriving
a no-win situation for the target virus. Under this scenario, the virus is either destroyed by the activity of
the DNA restriction systems, which would inhibit signal production (likely RNA fragments with cyclic
phosphate ends rather than cOA) and drive down CoCoNuT transcription, thereby preventing PCD, or as
the virus replicates, signaling and CoCoNuT transcription would continue until the infection is aborted by
PCD or dormancy caused by the degradation of host mRNA via the HEPN RNase(s) of the CoCoNuT. The
apparent toxicity of the CoCoNuT systems we expressed in *E. coli* (Supplementary Figure S7 A, B) is
compatible with this hypothesis, but as these were expressed from a plasmid, it is unclear if the toxicity
is naturally part of the CoCoNuT function. Future experiments will titrate the expression of the
CoCoNuTs to investigate this possibility further.

In many species of *Pseudomonas*, where CoCoNuTs are almost always associated with RtcR and
various ancillary factors, Type I RM systems often contain an additional gene that encodes a
transmembrane helix and a long coiled-coil fused to an RmuC-like nuclease. These proteins were
recently described as ShdA, the core component of the *Pseudomonas*-specific defense system Shield
[136]. These can potentially interact with coiled-coils in the CoCoNuTs, perhaps, guiding them to the
DNA from which RNA targeted by the CoCoNuTs is being transcribed or vice versa (Figure 9).

**Figure 9: Complex operonic associations of Type II CoCoNuTs**

Type II CoCoNuTs are frequently associated with RtcR homologs, and in many cases, ancillary defense genes are located
between the RtcR gene and the CoCoNuT, almost always oriented in the same direction in an apparent superoperon.
Abbreviations of domains: YprA – YprA-like helicase domain; DUF1998 - DUF1998 is often found in or associated with helicases
and contains four conserved, putatively metal ion-binding cysteine residues; PLD – Phospholipase D family nuclease domain;
SWI2/SNF2 – SWI2/SNF2-family ATPase; HsdR/M/S – Type I RM system restriction, methylation, and specificity factors; ShdA –
Shield system core component ShdA; TPR – Tetratricopeptide repeat protein; SLATT – Possible SLATT domain; Mod/Res – Type
III RM modification and restriction factors.

A notable difference between the generally similar, dual DNA and RNA-targeting mechanism of
many Type III CRISPR-Cas systems and the proposed mechanism of the CoCoNuTs is that specific DNA
cleavage activity of the Cas10-Csm complex, which in turn triggers non-specific RNA cleavage, is not
activated by cOA binding to a CARF domain, but rather by viral RNA binding. In contrast, in the CARF-
containing CoCoNuTs, both the DNA and RNA restriction factors appear to be arranged so that CARF
binding of a 2' 3' cyclic phosphate RNA fragment would initiate the expression of the entire cluster
[105]. In the case of the CoCoNuTs, signals of infection could activate transcription, first of the DNA
restriction systems, and then, the CoCoNuT itself, a predicted RNA restriction system. In these complex

configurations of the CoCoNuT genome neighborhoods (Figure 9), the gene order is likely to be
important, with Type I RM almost always directly following CARF genes and CoCoNuTs generally coming
last, although Druantia Type III sometimes follows the CoCoNuT. As translation in bacteria is co-
transcriptional, the products of genes transcribed first would accumulate before those of the genes
transcribed last, so that the full, potentially suicidal impact of the CoCoNuT RNA nucleolytic engine
would only be felt after the associated DNA restriction systems had ample time to act – and possibly, fail
[137].

Considering that the CoCoNuTs are potentially involved in degrading non-self and defective
mRNA and are abundant in Alphaproteobacteria, with a specialized variant, Type III-B, found mainly in
this class, some extant CoCoNuT CnuH homologs and the UPF1-like helicases required for NMD might
share a CoCoNuT-like common ancestor that was captured by emerging eukaryotes from the
protomitochondrial endosymbionts. In the case of NMD, recognition of a premature termination codon
(PTC), the mark of a nonsense message, is not completely understood, but a broadly conserved indicator
of a PTC seems to be inefficient translation termination [138-140]. Therefore, it is conceivable, and
perhaps not surprising, that the core regulator of NMD, UPF1, has prokaryotic roots in a family of
defensive systems that might exploit the bacterial tmRNA mechanism to degrade nonstop messages,
which stall ribosomes and prevent translation termination [78, 141]. The sophistication of these systems
in bacteria presages that of the NMD pathway, reinforcing the hypothesis on the origin of NMD from the
CoCoNuTs.

## Concluding remarks

In recent years, systematic searches for defense systems in prokaryotes, primarily by analysis of
defense islands, revealed enormous, previously unsuspected diversity of such systems that function
through a remarkable variety of molecular mechanisms. In this work, we uncovered the hidden diversity
and striking hierarchical complexity of a distinct class of defense mechanisms, the Type IV (McrBC)
restriction systems. We then zeroed in on a single major but previously overlooked branch of the McrBC
family, which we denoted CoCoNuTs for their salient features, namely, the presence of extensive coiled-
coil structures and tandem nucleases. Astounding complexity was also discovered at this level, with 3
distinct types and multiple subtypes of CoCoNuTs that differ by their domain compositions and genomic
associations. All CoCoNuTs contain domains interacting with translation systems components, such as
SmpB-like OBD, HSP70 homologs, or YTH domains, along with RNases, such as HEPN, suggesting that at
least one of the activities of these systems targets RNA. Most of the CoCoNuTs could be endowed with
DNA-targeting activity as well, either integral to the system, such as the McrC nuclease, or more loosely
associated, such as Type I RM and Druantia Type III systems that are encoded in the same predicted
superoperon with many CoCoNuTs. Numerous CoCoNuTs are associated with proteins containing CARF
domains, suggesting that cyclic (oligo)nucleotides regulate the CoCoNuT activity. Given the presence of
the RtcR-like CARF domains, it appears likely that the specific second messengers involved are RNA
fragments with cyclic phosphate termini. We hypothesize that the CoCoNuTs, in conjunction with
ancillary restriction factors, implement an echeloned defense strategy analogous to that of Type III
CRISPR-Cas systems, whereby an immune response eliminating virus DNA and/or RNA is launched first,
but then, if it fails, an abortive infection response leading to PCD/dormancy via host RNA cleavage takes
over. In a series of pilot experiments, we demonstrated that at least Type II CoCoNuTs can protect *E. coli*

against specific viruses and plasmids. The detailed exploration of the CoCoNuT functions and
mechanisms requires further, extensive experimentation that has the potential to uncover a variety of
currently unknown defense circuits.

## 985 Author contributions

## 989 Acknowledgments

The authors thank Dr. Becky Xu Hua Fu (University of California, San Francisco) for correspondence that
led to her contribution of the name CoCoNuT, Dr. Andrew Z. Fire and Usman Enam (Stanford University)
for critical reading of the manuscript and insightful comments, Dr. Joseph Bondy-Denomy (University of
California, San Francisco) for bringing the Shield factor ShdA to our attention, and Koonin group
members for helpful discussions. R.T.B., H.S., K.S.M., Y.I.W and E.V.K. are supported by the Intramural
Research Program of the National Institutes of Health (National Library of Medicine). J.J and I.S are
supported by Vilnius University Science Promotion fund project MSFJM-12/2021, I.S is funded by the
European Regional Development Fund under grant agreement number 01.2.2-CPVA-V-716-01-0001 with
the Lithuanian Central Project Management Agency (CPVA) and fellowship from The Lithuanian
Academy of Sciences.

1000

## References

1.  Burroughs, A.M., et al., *Comparative genomic analyses reveal a vast, novel network of nucleotide-centric systems in biological conflicts, immunity and signaling.* Nucleic Acids Research, 2015. **43**(22): p. 10633-10654.
2.  Gao, L., et al., *Diverse enzymatic activities mediate antiviral immunity in prokaryotes.* Science, 2020. **369**(6507): p. 1077-1084.
3.  Goldfarb, T., et al., *BREX is a novel phage resistance system widespread in microbial genomes.* Embo j, 2015. **34**(2): p. 169-83.
4.  Bell, R.T., Y.I. Wolf, and E.V. Koonin, *Modified base-binding EVE and DCD domains: striking diversity of genomic contexts in prokaryotes and predicted involvement in a variety of cellular processes.* BMC Biol, 2020. **18**(1): p. 159.
5.  Koonin, E.V. and L. Aravind, *Origin and evolution of eukaryotic apoptosis: the bacterial connection.* Cell Death & Differentiation, 2002. **9**(4): p. 394-404.
6.  Swarts, D.C., et al., *The evolutionary journey of Argonaute proteins.* Nature structural & molecular biology, 2014. **21**(9): p. 743-753.
7.  Anantharaman, V., L.M. Iyer, and L. Aravind, *Ter-dependent stress response systems: novel pathways related to metal sensing, production of a nucleoside-like metabolite, and DNA-processing.* Molecular bioSystems, 2012. **8**(12): p. 3142-3165.
8.  Kaur, G., et al., *Highly regulated, diversifying NTP-dependent biological conflict systems with implications for the emergence of multicellularity.* eLife, 2020. **9**: p. e52696.
9.  Loenen, W.A., et al., *Highlights of the DNA cutters: a short history of the restriction enzymes.* Nucleic Acids Res, 2014. **42**(1): p. 3-19.
10. Fire, A., et al., *Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans.* Nature, 1998. **391**(6669): p. 806-11.
11. Agrawal, N., et al., *RNA interference: biology, mechanism, and applications.* Microbiol Mol Biol Rev, 2003. **67**(4): p. 657-85.
12. Makarova, K.S., et al., *A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action.* Biology Direct, 2006. **1**(1): p. 7.
13. Makarova, K.S., et al., *Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants.* Nature Reviews Microbiology, 2020. **18**(2): p. 67-83.
14. Gasiunas, G., et al., *Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria.* Proc Natl Acad Sci U S A, 2012. **109**(39): p. E2579-86.
15. Jinek, M., et al., *A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity.* Science, 2012. **337**(6096): p. 816-821.
16. Raleigh, E.A., R. Trimarchi, and H. Revel, *Genetic and physical mapping of the mcrA (rglA) and mcrB (rglB) loci of Escherichia coli K-12.* Genetics, 1989. **122**(2): p. 279-296.
17. Luria, S.E. and M.L. Human, *A nonhereditary, host-induced variation of bacterial viruses.* J Bacteriol, 1952. **64**(4): p. 557-69.
18. Fleischman, R.A., J.L. Cambell, and C.C. Richardson, *Modification and restriction of T-even bacteriophages. In vitro degradation of deoxyribonucleic acid containing 5-hydroxymethylctosine.* J Biol Chem, 1976. **251**(6): p. 1561-70.
19. Dila, D., et al., *Genetic and sequence organization of the mcrBC locus of Escherichia coli K-12.* Journal of Bacteriology, 1990. **172**(9): p. 4888.
20. Sutherland, E., L. Coe, and E.A. Raleigh, *McrBC: a multisubunit GTP-dependent restriction endonuclease.* J Mol Biol, 1992. **225**(2): p. 327-48.

21.  Sukackaite, R., et al., *The recognition domain of the methyl-specific endonuclease McrBC flips out*
*5-methylcytosine.* Nucleic acids research, 2012. **40**(15): p. 7552-7562.
22.  Weigele, P. and E. Raleigh, *Biosynthesis and Function of Modified Bases in Bacteria and Their*
*Viruses.* Chemical reviews, 2016. **116**.
23.  Iyer, L.M., et al., *Evolutionary history and higher order classification of AAA+ ATPases.* Journal of
Structural Biology, 2004. **146**(1): p. 11-31.
24.  Nirwan, N., et al., *Structure-based mechanism for activation of the AAA+ GTPase McrB by the*
*endonuclease McrC.* Nature Communications, 2019. **10**(1): p. 3058.
25.  Panne, D., E.A. Raleigh, and T.A. Bickle, *The McrBC endonuclease translocates DNA in a reaction*
*dependent on GTP hydrolysis.* J Mol Biol, 1999. **290**(1): p. 49-60.
26.  Niu, Y., et al., *Structural asymmetry governs the assembly and GTPase activity of McrBC*
*restriction complexes.* Nat Commun, 2020. **11**(1): p. 5907.
27.  Altschul, S.F., et al., *Gapped BLAST and PSI-BLAST: a new generation of protein database search*
*programs.* Nucleic acids research, 1997. **25**(17): p. 3389-3402.
28.  Gabler, F., et al., *Protein Sequence Analysis Using the MPI Bioinformatics Toolkit.* Curr Protoc
Bioinformatics, 2020. **72**(1): p. e108.
29.  Zimmermann, L., et al., *A Completely Reimplemented MPI Bioinformatics Toolkit with a New*
*HHpred Server at its Core.* Journal of Molecular Biology, 2018. **430**(15): p. 2237-2243.
30.  Hauser, M., M. Steinegger, and J. Söding, *MMseqs software suite for fast and deep clustering*
*and searching of large protein sequence sets.* Bioinformatics, 2016. **32**(9): p. 1323-1330.
31.  Edgar, R.C., *MUSCLE: a multiple sequence alignment method with reduced time and space*
*complexity.* BMC bioinformatics, 2004. **5**: p. 113-113.
32.  Söding, J., *Protein homology detection by HMM–HMM comparison.* Bioinformatics, 2004. **21**(7):
p. 951-960.
33.  Söding, J., et al., *HHsenser: exhaustive transitive profile search using HMM–HMM comparison.*
Nucleic Acids Research, 2006. **34**(suppl_2): p. W374-W378.
34.  Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2 – Approximately Maximum-Likelihood Trees*
*for Large Alignments.* PLOS ONE, 2010. **5**(3): p. e9490.
35.  Robert, C.E., *High-accuracy alignment ensembles enable unbiased assessments of sequence*
*homology and phylogeny.* bioRxiv, 2022: p. 2021.06.20.449169.
36.  Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature, 2021.
**596**(7873): p. 583-589.
37.  Mariani, V., et al., *lDDT: a local superposition-free score for comparing protein structures and*
*models using distance difference tests.* Bioinformatics, 2013. **29**(21): p. 2722-2728.
38.  Holm, L., *DALI and the persistence of protein shape.* 2020. **29**(1): p. 128-140.
39.  van Kempen, M., et al., *Fast and accurate protein structure search with Foldseek.* Nature
Biotechnology, 2023.
40.  Pettersen, E.F., et al., *UCSF ChimeraX: Structure visualization for researchers, educators, and*
*developers.* 2021. **30**(1): p. 70-82.
41.  Berman, H.M., et al., *The Protein Data Bank.* Nucleic Acids Res, 2000. **28**(1): p. 235-42.
42.  Neuwald, A.F., et al., *AAA+: A class of chaperone-like ATPases associated with the assembly,*
*operation, and disassembly of protein complexes.* Genome Res, 1999. **9**(1): p. 27-43.
43.  Erzberger, J.P. and J.M. Berger, *Evolutionary relationships and structural mechanisms of AAA+*
*proteins.* Annu Rev Biophys Biomol Struct, 2006. **35**: p. 93-114.
44.  Pieper, U., et al., *The GTP-binding domain of McrB: more than just a variation on a common*
*theme?* J Mol Biol, 1999. **292**(3): p. 547-56.

45.    Hosford, C.J., A.Q. Bui, and J.S. Chappie, *The structure of the Thermococcus gammatolerans*
*McrB N-terminal domain reveals a new mode of substrate recognition and specificity among*
*McrB homologs.* Journal of Biological Chemistry, 2020. **295**(3): p. 743-756.
46.    Zimmermann, L., et al., *A Completely Reimplemented MPI Bioinformatics Toolkit with a New*
*HHpred Server at its Core.* J Mol Biol, 2018. **430**(15): p. 2237-2243.
47.    Simm, D., K. Hatje, and M. Kollmar, *Waggawagga: comparative visualization of coiled-coil*
*predictions and detection of stable single α-helices (SAH domains).* Bioinformatics, 2015. **31**(5):
p. 767-9.
48.    Varadi, M., et al., *AlphaFold Protein Structure Database: massively expanding the structural*
*coverage of protein-sequence space with high-accuracy models.* Nucleic Acids Research, 2022.
**50**(D1): p. D439-D444.
49.    Drozdetskiy, A., et al., *JPred4: a protein secondary structure prediction server.* Nucleic Acids
Research, 2015. **43**(W1): p. W389-W394.
50.    Raleigh, E.A., *Organization and function of the mcrBC genes of Escherichia coli K-12.* Mol
Microbiol, 1992. **6**(9): p. 1079-86.
51.    Ofir, G., et al., *DISARM is a widespread bacterial defence system with broad anti-phage*
*activities.* Nature Microbiology, 2018. **3**(1): p. 90-98.
52.    Colicelli, J., *Human RAS superfamily proteins and related GTPases.* Sci STKE, 2004. **2004**(250): p.
Re13.
53.    Bourne, H.R., D.A. Sanders, and F. McCormick, *The GTPase superfamily: conserved structure and*
*molecular mechanism.* Nature, 1991. **349**(6305): p. 117-27.
54.    Pillon, M.C., et al., *HEPN RNases - an emerging class of functionally distinct RNA processing and*
*degradation enzymes.* Crit Rev Biochem Mol Biol, 2021. **56**(1): p. 88-108.
55.    Mayer, M.P., *The Hsp70-Chaperone Machines in Bacteria.* Front Mol Biosci, 2021. **8**: p. 694012.
56.    Kishor, A., et al., *Hsp70's RNA-binding and mRNA-stabilizing activities are independent of its*
*protein chaperone functions.* J Biol Chem, 2017. **292**(34): p. 14122-14133.
57.    Zimmer, C., A. von Gabain, and T. Henics, *Analysis of sequence-specific binding of RNA to Hsp70*
*and its various homologs indicates the involvement of N- and C-terminal interactions.* Rna, 2001.
**7**(11): p. 1628-37.
58.    Kishor, A., et al., *Hsp70 is a novel posttranscriptional regulator of gene expression that binds and*
*stabilizes selected mRNAs containing AU-rich elements.* Mol Cell Biol, 2013. **33**(1): p. 71-84.
59.    Khadria, A.S. and A. Senes, *The transmembrane domains of the bacterial cell division proteins*
*FtsB and FtsL form a stable high-order oligomer.* Biochemistry, 2013. **52**(43): p. 7542-50.
60.    Chou, S.H. and M.Y. Galperin, *Diversity of Cyclic Di-GMP-Binding Proteins and Mechanisms.* J
Bacteriol, 2016. **198**(1): p. 32-46.
61.    Schnappauf, O., et al., *The Pyrin Inflammasome in Health and Disease.* Frontiers in Immunology,
2019. **10**.
62.    Park, H.H., et al., *The death domain superfamily in intracellular signaling of apoptosis and*
*inflammation.* Annu Rev Immunol, 2007. **25**: p. 561-86.
63.    Kaur, G., et al., *Bacterial death and TRADD-N domains help define novel apoptosis and immunity*
*mechanisms shared by prokaryotes and metazoans.* Elife, 2021. **10**.
64.    Stock, A.M., V.L. Robinson, and P.N. Goudreau, *Two-component signal transduction.* Annu Rev
Biochem, 2000. **69**: p. 183-215.
65.    Amir, M., et al., *Sequence, structure and evolutionary analysis of cold shock domain proteins, a*
*member of OB fold family.* Journal of Evolutionary Biology, 2018. **31**(12): p. 1903-1917.
66.    Hazra, D., C. Chapat, and M. Graille, *m⁶A mRNA Destiny: Chained to the rhYTHm by the YTH-*
*Containing Proteins.* Genes, 2019. **10**(1): p. 49.

67. Liao, S., H. Sun, and C. Xu, *YTH Domain: A Family of N6-methyladenosine (m6A) Readers.*
Genomics, Proteomics & Bioinformatics, 2018. **16**(2): p. 99-107.
68. Natarajan, K., M.G. Mage, and D.H. Margulies, *Immunoglobulin Superfamily*, in *Encyclopedia of*
*Life Sciences.* 2015. p. 1-7.
69. Halaby, D.M., A. Poupon, and J. Mornon, *The immunoglobulin fold family: sequence analysis and*
*3D structure comparisons.* Protein Eng, 1999. **12**(7): p. 563-71.
70. Gorbalenya, A.E. and E.V. Koonin, *Helicases: amino acid sequence comparisons and structure-*
*function relationships.* Current Opinion in Structural Biology, 1993. **3**(3): p. 419-429.
71. Fairman-Williams, M.E., U.P. Guenther, and E. Jankowsky, *SF1 and SF2 helicases: family matters.*
Curr Opin Struct Biol, 2010. **20**(3): p. 313-24.
72. Cheng, Z., et al., *Structural and functional insights into the human Upf1 helicase core.* Embo j,
2007. **26**(1): p. 253-64.
73. Chakrabarti, S., et al., *Molecular mechanisms for the RNA-dependent ATPase activity of Upf1 and*
*its regulation by Upf2.* Mol Cell, 2011. **41**(6): p. 693-703.
74. Zhou, C., S. Pourmal, and N.P. Pavletich, *Dna2 nuclease-helicase structure, mechanism and*
*regulation by Rpa.* eLife, 2015. **4**: p. e09832.
75. Kalathiya, U., et al., *Insights into the Effects of Cancer Associated Mutations at the UPF2 and*
*ATP-Binding Sites of NMD Master Regulator: UPF1.* Int J Mol Sci, 2019. **20**(22).
76. Morse, J.C., et al., *Elongation factor-Tu can repetitively engage aminoacyl-tRNA within the*
*ribosome during the proofreading stage of tRNA selection.* Proceedings of the National Academy
of Sciences, 2020. **117**(7): p. 3610-3620.
77. Barends, S., et al., *Simultaneous and functional binding of SmpB and EF-Tu-TP to the alanyl*
*acceptor arm of tmRNA.* J Mol Biol, 2001. **314**(1): p. 9-21.
78. Himeno, H., D. Kurita, and A. Muto, *tmRNA-mediated trans-translation as the major ribosome*
*rescue system in a bacterial cell.* Front Genet, 2014. **5**: p. 66.
79. Guyomar, C., et al., *Structures of tmRNA and SmpB as they transit through the ribosome.* Nat
Commun, 2021. **12**(1): p. 4909.
80. Yang, H., et al., *PAM-Dependent Target DNA Recognition and Cleavage by C2c1 CRISPR-Cas*
*Endonuclease.* Cell, 2016. **167**(7): p. 1814-1828.e12.
81. Zetsche, B., et al., *Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System.*
Cell, 2015. **163**(3): p. 759-771.
82. Liu, J.-J., et al., *CasX enzymes comprise a distinct family of RNA-guided genome editors.* Nature,
2019. **566**(7743): p. 218-223.
83. Altae-Tran, H., et al., *The widespread IS200/IS605 transposon family encodes diverse*
*programmable RNA-guided endonucleases.* Science, 2021. **374**(6563): p. 57-65.
84. Sasnauskas, G., et al., *TnpB structure reveals minimal functional core of Cas12 nuclease family.*
Nature, 2023. **616**(7956): p. 384-389.
85. Karvelis, T., et al., *Transposon-associated TnpB is a programmable RNA-guided DNA*
*endonuclease.* Nature, 2021. **599**(7886): p. 692-696.
86. Chavez, D.A., B.H. Greer, and B.F. Eichman, *The HIRAN domain of helicase-like transcription*
*factor positions the DNA translocase motor to drive efficient DNA fork regression.* J Biol Chem,
2018. **293**(22): p. 8484-8494.
87. Bessho, Y., et al., *Structural basis for functional mimicry of long-variable-arm tRNA by transfer-*
*messenger RNA.* Proceedings of the National Academy of Sciences, 2007. **104**(20): p. 8293-8298.
88. Gutmann, S., et al., *Crystal structure of the transfer-RNA domain of transfer-messenger RNA in*
*complex with SmpB.* Nature, 2003. **424**(6949): p. 699-703.
89. Giudice, E., K. Mace, and R. Gillet, *Trans-translation exposed: understanding the structures and*
*functions of tmRNA-SmpB.* Frontiers in Microbiology, 2014. **5**.

| 1188 | 90. | Gucinski, G.C., et al., *Convergent Evolution of the Barnase/EndoU/Colicin/RelE (BECR) Fold in* |
| 1189 | | *Antibacterial tRNase Toxins.* Structure, 2019. **27**(11): p. 1660-1674.e5. |
| 1190 | 91. | Pedersen, K., et al., *The bacterial toxin RelE displays codon-specific cleavage of mRNAs in the* |
| 1191 | | *ribosomal A site.* Cell, 2003. **112**(1): p. 131-40. |
| 1192 | 92. | Guglielmini, J. and L. Van Melderen, *Bacterial toxin-antitoxin systems: Translation inhibitors* |
| 1193 | | *everywhere.* Mob Genet Elements, 2011. **1**(4): p. 283-290. |
| 1194 | 93. | Goeders, N., P.L. Drèze, and L. Van Melderen, *Relaxed cleavage specificity within the RelE toxin* |
| 1195 | | *family.* J Bacteriol, 2013. **195**(11): p. 2541-9. |
| 1196 | 94. | Willmund, F., et al., *The cotranslational function of ribosome-associated Hsp70 in eukaryotic* |
| 1197 | | *protein homeostasis.* Cell, 2013. **152**(1-2): p. 196-209. |
| 1198 | 95. | Tsutakawa, S.E., H. Jingami, and K. Morikawa, *Recognition of a TG Mismatch: The Crystal* |
| 1199 | | *Structure of Very Short Patch Repair Endonuclease in Complex with a DNA Duplex.* Cell, 1999. |
| 1200 | | **99**(6): p. 615-623. |
| 1201 | 96. | Ipsaro, J.J., et al., *The structural biochemistry of Zucchini implicates it as a nuclease in piRNA* |
| 1202 | | *biogenesis.* Nature, 2012. **491**(7423): p. 279-283. |
| 1203 | 97. | Mendez, A.S., et al., *Site specific target binding controls RNA cleavage efficiency by the Kaposi's* |
| 1204 | | *sarcoma-associated herpesvirus endonuclease SOX.* Nucleic Acids Research, 2018. **46**(22): p. |
| 1205 | | 11968-11979. |
| 1206 | 98. | Laganeckas, M., M. Margelevicius, and C. Venclovas, *Identification of new homologs of PD-* |
| 1207 | | *(D/E)XK nucleases by support vector machines trained on data derived from profile-profile* |
| 1208 | | *alignments.* Nucleic Acids Res, 2011. **39**(4): p. 1187-96. |
| 1209 | 99. | Songailiene, I., et al., *HEPN-MNT Toxin-Antitoxin System: The HEPN Ribonuclease Is Neutralized* |
| 1210 | | *by OligoAMPylation.* Molecular Cell, 2020. **80**(6): p. 955-970.e7. |
| 1211 | 100. | Yamamoto, S., et al., *ATPase Activity and ATP-dependent Conformational Change in the Co-* |
| 1212 | | *chaperone HSP70/HSP90-organizing Protein (HOP)*.* Journal of Biological Chemistry, 2014. |
| 1213 | | **289**(14): p. 9880-9886. |
| 1214 | 101. | Chang, Y.W., et al., *Crystal structures of the 70-kDa heat shock proteins in domain disjoining* |
| 1215 | | *conformation.* J Biol Chem, 2008. **283**(22): p. 15502-11. |
| 1216 | 102. | Heinemann, U. and Y. Roske, *Cold-Shock Domains-Abundance, Structure, Properties, and* |
| 1217 | | *Nucleic-Acid Binding.* Cancers (Basel), 2021. **13**(2). |
| 1218 | 103. | Patil, D.P., B.F. Pickering, and S.R. Jaffrey, *Reading m6A in the Transcriptome: m6A-Binding* |
| 1219 | | *Proteins.* Trends in Cell Biology, 2018. **28**(2): p. 113-127. |
| 1220 | 104. | Makarova, K.S., et al., *Evolutionary and functional classification of the CARF domain superfamily,* |
| 1221 | | *key sensors in prokaryotic antivirus defense.* Nucleic Acids Res, 2020. **48**(16): p. 8828-8847. |
| 1222 | 105. | McMahon, S.A., et al., *Structure and mechanism of a Type III CRISPR defence DNA nuclease* |
| 1223 | | *activated by cyclic oligoadenylate.* Nature Communications, 2020. **11**(1): p. 500. |
| 1224 | 106. | Zhu, W., et al., *The CRISPR ancillary effector Can2 is a dual-specificity nuclease potentiating type* |
| 1225 | | *III CRISPR defence.* Nucleic Acids Research, 2021. **49**(5): p. 2777-2789. |
| 1226 | 107. | Makarova, K.S., et al., *Live virus-free or die: coupling of antivirus immunity and programmed* |
| 1227 | | *suicide or dormancy in prokaryotes.* Biology direct, 2012. **7**: p. 40-40. |
| 1228 | 108. | Koonin, E.V. and F. Zhang, *Coupling immunity and programmed cell suicide in prokaryotes: Life-* |
| 1229 | | *or-death choices.* BioEssays, 2017. **39**(1): p. e201600186. |
| 1230 | 109. | Koonin, E.V. and M. Krupovic, *Origin of programmed cell death from antiviral defense?* |
| 1231 | | Proceedings of the National Academy of Sciences, 2019. **116**(33): p. 16167. |
| 1232 | 110. | Kotta-Loizou, I., et al., *The RNA repair proteins RtcAB regulate transcription activator RtcR via its* |
| 1233 | | *CRISPR-associated Rossmann fold domain.* iScience, 2022. **25**(11): p. 105425. |
| 1234 | 111. | Hughes, K.J., et al., *An RNA Repair Operon Regulated by Damaged tRNAs.* Cell Rep, 2020. **33**(12): |
| 1235 | | p. 108527. |

112. Bitton, L., D. Klaiman, and G. Kaufmann, *Phage T4-induced DNA breaks activate a tRNA repair-*
*defying anticodon nuclease.* Mol Microbiol, 2015. **97**(5): p. 898-910.
113. van den Berg, D.F., et al., *Phage tRNAs evade tRNA-targeting host defenses through anticodon*
*loop mutations.* eLife, 2023. **12**: p. e85183.
114. Kaufmann, G., *Anticodon nucleases.* Trends Biochem Sci, 2000. **25**(2): p. 70-4.
115. Ishita, J., et al., *tRNA anticodon cleavage by target-activated CRISPR-Cas13a effector.* bioRxiv,
2021: p. 2021.11.10.468108.
116. Genschik, P., K. Drabikowski, and W. Filipowicz, *Characterization of the Escherichia coli RNA 3'-*
*terminal phosphate cyclase and its sigma54-regulated operon.* J Biol Chem, 1998. **273**(39): p.
25516-26.
117. Das, U. and S. Shuman, *2'-Phosphate cyclase activity of RtcA: a potential rationale for the operon*
*organization of RtcA with an RNA repair ligase RtcB in Escherichia coli and other bacterial taxa.*
Rna, 2013. **19**(10): p. 1355-62.
118. Tanaka, N. and S. Shuman, *RtcB is the RNA ligase component of an Escherichia coli RNA repair*
*operon.* J Biol Chem, 2011. **286**(10): p. 7727-7731.
119. Kosmaczewski, S.G., et al., *The RtcB RNA ligase is an essential component of the metazoan*
*unfolded protein response.* EMBO Rep, 2014. **15**(12): p. 1278-85.
120. Popow, J., et al., *HSPC117 is the essential subunit of a human tRNA splicing ligase complex.*
Science, 2011. **331**(6018): p. 760-4.
121. Kirby, J.E., J.E. Trempy, and S. Gottesman, *Excision of a P4-like cryptic prophage leads to Alp*
*protease expression in Escherichia coli.* J Bacteriol, 1994. **176**(7): p. 2068-81.
122. Han, Y., et al., *Structure of human RNase L reveals the basis for regulated RNA decay in the IFN*
*response.* Science, 2014. **343**(6176): p. 1244-8.
123. Chakrabarti, A., B.K. Jha, and R.H. Silverman, *New insights into the role of RNase L in innate*
*immunity.* J Interferon Cytokine Res, 2011. **31**(1): p. 49-57.
124. Bösl, M. and H. Kersten, *A novel RNA product of the tyrT operon of Escherichia coli.* Nucleic Acids
Res, 1991. **19**(21): p. 5863-70.
125. Kohanski, M.A., et al., *A common mechanism of cellular death induced by bactericidal*
*antibiotics.* Cell, 2007. **130**(5): p. 797-810.
126. Rossi, C.C., et al., *A computational strategy for the search of regulatory small RNAs in*
*Actinobacillus pleuropneumoniae.* Rna, 2016. **22**(9): p. 1373-85.
127. Iyer, L.M., A.M. Burroughs, and L. Aravind, *The prokaryotic antecedents of the ubiquitin-*
*signaling system and the early evolution of ubiquitin-like beta-grasp domains.* Genome Biol,
2006. **7**(7): p. R60.
128. Anderson, B.W., et al., *Molecular Mechanism of Regulation of the Purine Salvage Enzyme XPRT*
*by the Alarmones pppGpp, ppGpp, and pGpp.* J Mol Biol, 2020. **432**(14): p. 4108-4126.
129. Henry, J.T. and S. Crosson, *Ligand-binding PAS domains in a genomic, cellular, and structural*
*context.* Annu Rev Microbiol, 2011. **65**: p. 261-86.
130. Anantharaman, V., et al., *Comprehensive analysis of the HEPN superfamily: identification of*
*novel roles in intra-genomic conflicts, defense, pathogenesis and RNA processing.* Biol Direct,
2013. **8**: p. 15.
131. Panne, D., et al., *The McrBC restriction endonuclease assembles into a ring structure in the*
*presence of G nucleotides.* Embo j, 2001. **20**(12): p. 3210-7.
132. Maracci, C. and M.V. Rodnina, *Review: Translational GTPases.* Biopolymers, 2016. **105**(8): p.
463-75.
133. Jewett, M.C., et al., *Continued protein synthesis at low [ATP] and [GTP] enables cell adaptation*
*during energy limitation.* J Bacteriol, 2009. **191**(3): p. 1083-91.

134.   Rodnina, M.V., et al., *GTPases mechanisms and functions of translation factors on the ribosome.*
Biol Chem, 2000. **381**(5-6): p. 377-87.
135.   Millman, A., et al., *Diversity and classification of cyclic-oligonucleotide-based anti-phage*
*signalling systems.* Nature Microbiology, 2020. **5**(12): p. 1608-1615.
136.   Macdonald, E., et al., *Shield co-opts an RmuC domain to mediate phage defence across*
*Pseudomonas species.* 2022: p. 2022.11.04.515146.
137.   Irastortza-Olaziregi, M. and O. Amster-Choder, *Coupled Transcription-Translation in Prokaryotes:*
*An Old Couple With New Surprises.* Front Microbiol, 2020. **11**: p. 624830.
138.   Karousis, E.D. and O. Mühlemann, *Nonsense-Mediated mRNA Decay Begins Where Translation*
*Ends.* Cold Spring Harb Perspect Biol, 2019. **11**(2).
139.   Serdar, L.D., et al., *Inhibition of post-termination ribosome recycling at premature termination*
*codons in UPF1 ATPase mutants.* Elife, 2020. **9**.
140.   Kervestin, S. and A. Jacobson, *NMD: a multifaceted response to premature translational*
*termination.* Nat Rev Mol Cell Biol, 2012. **13**(11): p. 700-12.
141.   Kim, Y.K. and L.E. Maquat, *UPFront and center in RNA decay: UPF1 in nonsense-mediated mRNA*
*decay and beyond.* RNA, 2019. **25**(4): p. 407-422.

1299