Title: A diversified, widespread microbial gene cluster encodes homologs of

methyltransferases involved in methanogenesis.

Duncan J. Kountz[1] and Emily P. Balskus[1,2] [*]

Author information

Duncan J. Kountz djkountz@g.harvard.edu

Emily P. Balskus balskus@chemistry.harvard.edu, orcid.org/0000-0001-5985-5714

[1]Department of Chemistry and Chemical Biology, Harvard University, Cambridge,

Massachusetts 02138, United States.

[2]Howard Hughes Medical Institute, Harvard University, Cambridge, Massachusetts

02138, United States.

* Corresponding author

Competing interests: None

Short Title: Methanogen methyltransferases homologs

Abstract: Analyses of microbial genomes have revealed unexpectedly wide distributions of enzymes from specialized metabolism, including methanogenesis, providing exciting opportunities for discovery. Here, we identify a family of gene clusters (the type 1 *mlp* gene clusters (MGCs)) that encodes homologs of the soluble coenzyme M methyltransferases (SCMTs) involved in methylotrophic methanogenesis and is widespread in bacteria and archaea. Type 1 MGCs are expressed and regulated in medically, environmentally, and industrially important organisms, making them likely to be physiologically relevant. Enzyme annotation, analysis of genomic context, and biochemical experiments suggests these gene clusters play a role in methyl-sulfur and/or methyl-selenide metabolism in numerous anoxic environments, including the human gut microbiome, potentially impacting sulfur and selenium cycling in diverse, anoxic environments.

Keywords: Corrinoid methyltransferases, methanogenesis, selenium metabolism, sulfur metabolism, bacteria

Introduction

Other the past 50 years, biochemists and microbiologists have made much progress in elucidating the enzymology of methanogenesis (1,2). This reservoir of biochemical knowledge presents the opportunity to identify homologs of methanogen specific enzymes in organisms that are categorically not methanogens. In doing so, we may discover previously unappreciated metabolic pathways with global or ecological

relevance. Moreover, studying homologs of methanogen-specific enzymes in non-

methanogenic organisms could shed light on the evolution of methanogenic pathways

and organisms as well as the potential adaptation of methanogen genes to perform novel

functions.

Methanogenesis can be regarded as a specialized metabolism—its pathways are

biochemically distinct from those of most organisms and not expected to interface with

other metabolic processes when horizontally transferred. However, methanogen

cofactors and homologs of methanogen enzymes are indeed found in non-methanogens.

Examples include the use of methanofuran- and methanopterin-dependent enzymes in

aerobic methylotrophy (3), deazaflavin redox cofactor F_{420} in bacterial metabolism (4),

coenzyme M in alkene catabolism (5), heterodisulfide reductase-like enzymes in various

metabolisms, including sulfate reduction (6), acetogenesis (7), and the degradation of

benzoate (8). In each case the "methanogen" enzymes are performing an interesting,

novel, and important function.

We wondered if this logic extends to the soluble coenzyme M methyltransferases

(SCMTs) from methylotrophic methanogenesis. SCMTs are corrinoid-dependent

methyltransferases that transfer a methyl group from a methylated corrinoid-binding

protein (corrinoid protein) to coenzyme M to form methyl-coenzyme M, the immediate

precursor of methane (Fig. 1) (9). SCMTs belong to the very large and sequence-diverse

uroporphyrinogen decarboxylase (UroD) superfamily, which includes uroporphyrinogen

decarboxylase, cobalamin-independent methionine synthase, epoxide coenzyme M

transferase, and a variety of corrinoid methyltransferases including *O*-, *S*-, and halogen-

demethylases (10–16). However, we focused on proteins having >25% amino acid

identity to a characterized SCMT. We call these enzymes <u>Mt</u>aA-<u>l</u>ike <u>p</u>roteins, or Mlps,

after MtaA, the methanol-specific SCMT in *Methanosarcina* species (16). To our

knowledge, these Mlps have not been assigned a role outside of methanogens.

Fig. 1   Searching genomes for the soluble coenzyme M methyltransferase (SCMT) MtaA involved

in methylotrophic methanogenesis identifies related enzymes in non-methanogens. Overview of

genome mining strategy used in this work. The methanol-specific methanol:corrinoid methyltransferase

MtaB transfers the methyl group of methanol to the cognate corrinoid protein MtaC (the corrinoid cofactor

is represented by a square containing a cobalt ion). The SCMT MtaA then transfers to the methyl group

from MtaC to coenzyme M (CoM). The methyl group incorporated onto CoM is then disproportionated into

methane and carbon dioxide or solely reduced to methane, depending on the organism and environmental

conditions.

Here, we propose a prospective role for Mlps in a wide range of microorganisms by

identifying a diverse, widespread, and homologous set of gene clusters that encode

multiple Mlps, a corrinoid protein and a corrinoid reductive activase (Ram/RACE enzyme).

Gene cluster annotation, bioinformatic analysis, and biochemical logic suggests that the

encoded enzymes have two functions—methyl-selenide recycling and/or methionine

salvage—depending on the organism. These results illustrate the utility of using known

enzymes involved in metabolisms of interest to discover uncharacterized microbial gene

clusters that likely have intriguing and ecologically important functions.

Materials and methods

<u>General methods and procedures</u>

Manipulations and visualizations of genomic data were performed using Geneious

Prime 2020.0.4 software (17). BLASTP searches were conducted through the Geneious

interface using either National Center for Biotechnology Information (NCBI) nr/rt database

or custom databases generated by the user. In all cases we used the BLOSUM62 matrix

and a gap opening penalty of 11 and a gap extension penalty of 1. Protein alignments we

performed using MUSCLE (18).

<u>Identification of *mlp* gene clusters</u>

To identify homologs of SCMTs in *C. ljungdahlii*, we generated a protein database of

all the predicted proteins encoded in the *C. ljungdahlii* DSM 13528 genome, and

performed at BLASTP search of this database using the *Methanosarcina barkeri* MtbA

protein sequence (Swiss-Prot ID: O30640) as a query. We then inspected the genomic

context of the all the hits with E-values less than 1.0e-10.

To identify additional MGCs, we performed BLASTP searches against NCBI's nr/nt

database using *C. ljungdahlii* Mlp1, Mlp2, and Mlp3 (locus tags in Table S1) as queries.

We collected the top 1000 hits from each BLAST search. For the Mlp1 search, this yielded

hits with as low as 19.8% amino acid identity to the query. For the Mlp2 and Mlp3

searches, the minimum percent identities of the hits were 28.0% and 24.9%, respectively.

We then manually inspected the genomic context of each hit to determine whether they

were found in true MGCs. A "true" MGC met the following criteria: 1) it encodes at least

two Mlps, a corrinoid protein, and a Ram protein, 2) each of these components are

transcribed in the same direction with no intervening divergently-transcribed genes, 3)

there is no more than 2 kb of DNA in between the genes noted in 1, and 4) the gene

cluster encodes no homologs of tetrahydrofolate methylases within 20 kb of any of the

genes noted in 1. Using these methods and definitions, we identified 367 MGCs.

<u>SSN analysis of MGC proteins</u>

We generated all SSNs using the Enzyme Function Initiative Enzyme Similarity Tool

(EFI-EST) webtool (19). For each SSN, we first compiled the sequences of characterized

superfamily members and superfamily members that are encoded in the 367 MGCs

identified above, generating a superfamily-specific "preliminary input list". We then pruned

each list to ensure that only quality sequences (those lacking frameshifts or internal stops)

within a certain length window were included. For the Mlp SSN, length window was 200

to 470 amino acids. The resulting sequence list was termed the "input list". We next input

both the appropriate superfamily and input list into the EFI-EST web server. For the Mlp

SSN, the superfamily was IPR027980, for the Ram/RACE SSN, it was IPR027980, and

for the corrinoid protein SSN it was IPR006158. We selected alignment score thresholds

such that most MGC proteins fell into common network clusters, but were still as

separated as possible from non-MGC proteins. We selected node representation percent

identities to the highest that consistent with efficient use of the Cytoscape GUI on an

ordinary laptop MacIntosh computer. For each SSN, we used the UniRef90 option and

otherwise default parameters. Once generated, the SSNs were visualized with Cytoscape

software (version 3.8.0) using the "preferred" layout (20).

Phylogenetic analysis of *mlp* gene cluster proteins

We compiled proteins encoded by type I MGCs and sequences of related

characterized proteins and then aligned the resulting protein sequences using MUSCLE

with default parameters, and trimmed the alignments manually. All trees were generated

using FastTree (approximately-maximum likelihood method) with default parameters

(21).

We generated the rooted phylogenetic tree of the Mlps as follows. We first compiled

the sequences of Mlps1, Mlps2, and Mlps3 from type I MGCs included in the above SSN.

We added the SCMTs identified in the unrooted tree to this list, as well as the known

phenolic *O*-demethylases OdmB and VdmB (UniProt IDs: O87604 and Q6W001,

respectively). The SCMTs were identified as the smallest set of branches containing the

characterized SCMTs MtaA, MtbA, and MtsA (all from *Methanosarcina barkeri* Fusaro),

but excluding the Mlps3. The SCMTs were identified as the smallest clade in the unrooted

tree which contained the known SCMTs MtaA, MtbA, and MtsA. The tree was rooted on

the branch connecting the *O*-demethylases to the rest of the sequences, based on the

hypothesis that they serve as an appropriate outgroup.

Results

While manually inspecting the genome of *Clostridium ljungdahlii* DSM 13528, a

potential producer of biofuels, we noticed that this organism encodes multiple members

of the UroD superfamily. To see if these genes are closely related to SCMTs, we

performed BLASTP searches of the SCMT MtbA (*Methanosarcina barkeri* Fusaro,

*Mb*MtbA) against the predicted *C. lungdahlii* DSM 13528 proteome. We identified seven

hits with an E value < 1e-09 (Table S1). The highest hit was CLJU_RS09375, which

showed 32.5% amino acid (aa) identity to *Mb*MtbA. It is worth noting that the SCMTs

MtbA, MtaA, and MtsA (all from *M. barkeri* Fusaro) show only about 30% aa identity to

each other, suggesting that CLJU_RS09375 could be a close relative of the SCMTs.

CLJU_RS09375 had only 17.6% and 17.0% aa identity to the UroD superfamily phenolic

O-demethylases OdmB and VdmB, respectively.

To explore the physiological role of CLJU_RS09375, we next examined its

genomic context. CLJU_RS09375 is encoded in a gene cluster that includes genes

encoding a corrinoid protein (CLJU_RS09370), a "bacterial-type" corrinoid protein

reductive activase (Ram/RACE protein) (CLJU_RS09380), and two additional Mlps

(CLJU_RS09360 and CLJU_RS09365) (Fig. 2a). This gene cluster has been

independently predicted to form an operon by OperonDB, an automatic operon prediction

database (26). Because this gene cluster encodes multiple Mlps, we named it the *mlp*

gene cluster (MGC). We named the three Mlps in the gene cluster Mlp1, Mlp2, and Mlp3,

(specifically, *Cl*Mlp1, 2, and 3) in order of their appearance in the gene cluster from the

presumed transcriptional start site (see Fig. 2a).

Fig. 2 Mlp gene clusters (MGCs) are similar in genetic composition and protein seqence. a The

MGC from *C. ljungdahlii*. **b** Representative type I MGC. Coloring of the core MGC genes is the same as in

a, but non-core genes are colored grey. **c** A protein SSN of the UroD superfamily (IPR000257) showing the

clustering of Mlps1, 2, and 3. Nodes representing MtsA, MtbA, and MtaA (all from *M. barkeri* Fusaro) are

shown in dark blue. The alignment score cutoff was 40 and the node representation percent cut was 50 %

aa identity. The SSN shown is a trimmed version of the full SSN, with all nodes removed that are neither

directly or indirectly connected to the Mlps1, 2, or 3.

Given the relatively high sequence similarity of the MGC Mlps to SCMTs, it seems

reasonable to hypothesize that these enzymes are corrinoid methyltransferases. This is

consistent with the presence of genes encoding a corrinoid protein and a Ram/RACE

protein in the MGC. The former is a methyl group carrier (27), while the latter is an ATP-

dependent corrinoid protein reductase that reductively activates the corrinoid protein to

its catalytically-active Co(I) state (28,29). Corrinoid methyltransferases, corrinoid

proteins, and Ram/RACE proteins are frequently colocalized (30–32). In bacteria, such

gene clusters are involved in methylotrophic metabolism and participate in transferring a

methyl group from a donor substrate to tetrahydrofolate (H$_4$folate) via a methyl-corrinoid

protein intermediate, a reaction catalyzed by a methyl-corrinoid protein:H$_4$folate

methyltransferase (15,30). In most of the bacterial methyltransferase pathways examined

to date, the H$_4$folate methylase is encoded in the same gene cluster as the corrinoid

protein and methyl donor methyltransferase (30,31). It is therefore surprising that H$_4$folate

methyltransferase is missing from the *C. ljungdahlii* MGC. While it is possible that a

H$_4$folate methyltransferase encoded elsewhere in the *C. ljungdahlii* genome could

cooperate with the MGC methyltransferase system, it is also possible that H$_4$folate is not

involved or that one of the MGC Mlps is an undiscovered H$_4$folate methyltransferase.

In parallel, we located similar gene clusters in *C. sporogenes* DSM 795 and

Hungateiclostridium (*Clostridium*) *thermocellum* DSM 1313. Like the MGC from *C.*

ljungdahlii, the *C. sporogenes* and *H. thermocellum* gene clusters encode multiple Mlps

(each with about 30% aa identity to the SCMTs), a corrinoid protein, and a Ram/RACE

protein (Fig. 2b). These gene clusters also lack a H$_4$folate methylase. However, unlike *C.*

ljungdahlii, which encodes H$_4$folate methylases elsewhere in its genome, homologs of

H$_4$folate methylases are absent from the *C. sporogenes* and *H. thermocellum* genomes,

suggesting that H$_4$folate is not involved in the pathways encoded by these gene clusters.

Because of the similarity of the *C. sporogenes* and *H. thermocellum* gene clusters to the

C. ljungdahlii MGC, we refer to then as the *Cs*MGC, *Ht*MGC, and *Cl*MGC, respectively.

We next wondered if related MGCs are found in additional organisms. To address this,

we BLASTed *Cl*Mlp1, 2, and 3 against the NCBI non-redundant protein database. Many

of the organisms that had high alignment score hits for *Cl*Mlp1 also encoded high

alignment score hits for *Cl*Mlp2 and 3. When we examined the genomic contexts of these

hits, we identified gene clusters that resemble the *C. ljungdahlii* MGC (Fig. 2b).

We next asked if these MGCs are homologous or analogous. Because components

of a corrinoid methyltransferase pathway are commonly encoded in gene clusters

(14,15,30–32), evolutionary pressure to colocalize corrinoid methyltransferase genes

could have resulted in superficially similar MGCs that share no evolutionary relationship.

Alternatively, these gene clusters could be descended from a common ancestor (i.e. are

homologous), which would also suggested a shared function in diverse organisms. To

address this question, we identified 367 organisms encoding likely MGCs (see materials

and methods). We next constructed several protein sequence similarity networks (SSNs)

(19), reasoning that if MGCs are homologous, each conserved component should form

its own cluster in an SSN of the protein family to which that component belongs. Because

there are multiple *mlp* genes in each MGC, there are two SSN analysis outcomes that

are consistent with the homology hypothesis. First, all Mlps from the MGCs could be

grouped within a single cluster in the SSN, which would be expected if the multiple *mlp*

genes in each MGC arose from gene duplication. Second, the MGC Mlps could each

represent different lineages and would therefore likely form distinct clusters within an

SSN.

We first constructed an SSN (Fig. 2c) of the IPR000257 superfamily, which includes

the Mlps. We found that 80.9% of the MGC Mlps localized into one of three sub-clusters

within a larger cluster (Fig. 2c). Interestingly, these three sub-clusters correspond to the

three Mlps from the *C. ljungdahlii* MGC, with *Cl*Mlp1, 2, and 3 being grouped into distinct

sub-clusters. Inspection of other MGCs reveals that almost all encode Mlps from clusters

2 and 3, while a gene encoding an Mlp from cluster 1 was present a majority of the time,

but is often absent. We denote the proteins in the sub-cluster containing *Cl*Mlp1 as Mlps1,

and those from sub-clusters containing *Cl*Mlp2 and 3 as Mlps2, and Mlps3, respectively.

Altogether, this analysis shows that a large number of MGCs encode at least an Mlp2

and an Mlp3. We designate such gene clusters as "type 1 MGCs".

Because >95% of MGCs identified appear to be type 1 MGCs, we selected these

MGCs for further study. The fact that the type 1 MGC Mlps fall into three separate clusters

in the SSN suggests they might represent three distinct Mlp lineages, rather than a single

lineage that arose from recent gene duplication events. To solidify this inference, we

constructed a phylogenetic tree of the Mlps and other members of the UroD superfamily

that are close relatives of known corrinoid methyltransferases (Figure 3a). The Mlps 1, 2,

and 3 are situated in a large branch that includes the SCMTs, to the exclusion of other

known UroD superfamily corrinoid methyltransferases such as the aromatic *O*-

demethylases (13). When rooted (Fig. 3b), we found that the Mlps 1, 2 and 3 formed

distinct, well-supported clades. Interestingly, the Mlps2 form a sister clade to the clade

containing the SCMTs and the Mlps1 and Mlps3. Within this clade, the Mlps1 and Mlps3

form sister clades emerging from the SCMTs, which therefore appear to be paraphyletic.

Fig. 3 Type I MGCs encode orthologous proteins. a An approximately-maximum likelihood (FastTree)

phylogenetic tree of members of the UroD superfamily methyltransferases. The FastTree support values

for the Mlps1, 2, and 3 clades are 0.95, 0.94, and 1.00, respectively. Scale bar: branch length corresponding

to an average of one change per position. **b** A rooted approximately-maximum likelihood (FastTree)

phylogenetic tree of selected protein sequences. Shown are the soluble coenzyme M methyltransferases

(SCMTs), the Mlps1, 2 and 3, and the phenolic *O*-demethylases OdmB and VdmB (outgroup). Each branch

is labeled with the branch's FastTree support value. The number of sequences in each clade are as follows.

O-demethylases: 2; Mlps2: 267; Misc. SCMT clade: 20; MtbA clade: 85; Mlps1: 147; Mlps3: 233; MtaA +

MtsA clade: 252. **c** An approximately-maximum likelihood (FastTree) phylogenetic tree of members of the

methyltransferase corrinoid protein family. The type I MGC proteins are shown in red. The FastTree support

value for the type I MGC proteins is 0.92. Selected characterized family members are shown: MtbC1,

MtmC1, and MttC1 (all from *M. acetivorans* C2A); *Dh*MtgC from *Desulfitobacterium hafniense* Y51; MtqC

from *Eubacterium limosum* ATCC 8486; MtsB from *M. barkeri* Fusaro; OdmA and VdmA from

Acetobacterium dehalogenans DSM 11527. **d** An approximately-maximum likelihood (FastTree)

phylogenetic tree of members of the Ram/RACE protein family. The type I MGC proteins are shown in

orange. The FastTree support value for the type I MGC proteins is 0.98. A clade containing the corrinoid

iron-sulfur protein reductive activases is labeled "AcsVs". The locations of the activases RamQ (*E. limosum*

ATCC 8486) and OdmC (*A. dehalogenans* DSM 11527) are shown. Scale bars for all tree represent

respective branch length corresponding to an average of one change per position.

277   To further test this hypothesis, we generated two additional phylogenetic trees:

one containing the type 1 MGC corrinoid proteins and other monomeric methyltransferase

corrinoid proteins (Fig. 3c and 3d, respectively), and the other containing the type 1 MGC

Ram/RACE proteins and other Ram/RACE proteins (Fig. S4). In each case, the type 1

MGC proteins were found in the same branch, with a handful of exceptions. These

exceptions could be the result of gene conversion or the replacement of a "canonical"

type 1 MGC gene with an isofunctional gene from outside of the MGC.

Together, data from the SSN and phylogenetic analyses suggests that type 1 MGCs

are homologous in that they encode mutually orthologous proteins. The simplest

explanation for the homology of the type 1 MGCs is that they are horizontally transferred

as a unit. However, we cannot rule out more complicated scenarios including the transfer

of type 1 MGC genes individually into different regions of the genome followed by

selective pressure for colocalization. This latter possibility gains some credence when we

note the lack of conserved synteny in type 1 MGCs.

We next analyzed the phylogenetic distribution of type 1 MGCs (Fig. 4). To date, we

have identified over one hundred prokaryotic genera containing at least one type 1 MGC-

encoding strain. Of these, only three (*Methanococcus*, *Methanothermococcus*, and

Methanotorris) are methanogenic archaea. This is noteworthy because members of the

Methanococcales are not known to be methylotrophic and therefore would not be

expected to encode SCMTs. All type I MGC-encoding genera only obligate anaerobes,

with the majority (73%) belonging to the Firmicutes phylum. Although type 1 MGCs are

found a limited number of phyla, they are widely distributed across at least 35 orders of

bacteria and archaea. From this analysis, we conclude that type 1 MGCs are

phylogenetically widespread and typically associated with microorganisms from anoxic

environments. Notable species encoding type 1 MGCs include toxigenic clostridia

(*Clostridium botulinum*, *Paeniclostridium sordelii*, certain strains of *Clostridioides difficile*)

and industrially-relevant clostridia (*Hungateiclostridium* (*Clostridium*) *thermocellum*,

Clostridium ljungdahlii).

Firmicutes (74)
Desulfobacterota (11)
Synergistota (4)
Methanobacteriota (4)
Spirochaetota (3)
Fusobacteriota (2)
Planctomycetota (2)
Proteobacteria (1)

Fig. 4 Type I MGCs are found in several prokaryotic Phyla. Phylum composition of genera encoding type I MGCs. 101 genera are represented. The number of type I MGC-encoding genera within each phylum is enclosed in parentheses.

We next analyzed additional components of the type 1 MGCs and proposed a biochemical function for these genes. The UroD superfamily includes corrinoid methyltransferases that (de)methylate heteroatom-containing substrates (10,12,14,15,33). Several lines of evidence suggest that the type 1 MGCs encode a corrinoid S- or Se-methyltransferase system. First, the closest relationship between known UroD superfamily proteins and the type 1 mlp cluster Mlps is with the SCMTs (Fig. a and b), which are S-methyltransferases. Additional evidence comes from examination of the enzyme's Zn^{2+}-binding site. This Zn^{2+} cofactor coordinates the heteroatom of a substrate, activating it for (de)methylation (24,33,34). The primary coordination sphere of the Zn^{2+} cofactor is His-Cys-Cys in S-methyltransferases from the UroD superfamily, while in O-methyltransferases, one or both of the Cys ligands are substituted by carboxylates (Glu or Asp) or are substituted for non-coordinating residues (15,24,33,34). The His-Cys-Cys coordination environment favors the binding of a soft ligand such as a selenium or sulfur atom to the Zn^{2+} center, while substitution of Cys for a carboxylate

favors binding of an oxygen atom. The coordination sphere of the type 1 MGC Mlps is

(with the exception of several Mlps1) invariably His-Cys-Cys, suggesting that these

proteins are *S-* or *Se*-methyltransferases. Homology models of the *C. ljungdahlii* Mlps in

further confirmed that these residues are likely appropriately positioned to coordinate Zn^{2+}

(Fig. S3). Finally, examination of the non-core components of type I MGCs (genes in type

I MGCs that do not encode Mlps, corrinoid proteins, or Ram/RACE proteins) consistently

reveals genes involved in sulfur metabolism, including methionine aminotransferases,

sulfurtransferases, methionine transporters, and close homologs of the recently-

described MarHDK enzyme that catalyzes C–S bond cleavage to produce ethylene from

methylthioethanol (35). These observations suggest a potential role for many type I MGCs

in methyl-sulfur metabolism.

However, additional evidence suggests that most type I MGCs function in methyl-

selenium metabolism. Many type 1 MGCs contain genes provisionally linked to selenium

metabolism, such as the recently characterized "dicubane cluster protein" (36–38).

Several type 1 MGCs (Fig. 5) contain genes encoding authentic machinery for

selenocysteine synthesis, encoding, and decoding (SelABCD), as well as the enzymes

of the glycine reductase system, two of which have selenocysteine residues (39,40).

Notably, 87% of sequenced species that encode type I MGCs also encode selenocysteine

biosynthesis machinery within the same genome. For comparison, only 21.5% of bacteria

contain the requisite genes for selenocysteine biosynthesis, and even in the

selenocysteine-rich Clostridia, only 42% of 184 genomes encoded selenocysteine

biosynthesis machinery (36). Not all organisms that make use of selenium encode

selenocysteine biosynthetic machinery—some organisms that use the selenium cofactor

only encode selenide-water dikinase (SelD) (36). We therefore manually inspected

genomes that contain a type I MGC, but lack SelAB or C, finding that several genomes

lacked *selABC*, but had *selD* as well as selenium cofactor-dependent enzymes. As noted

below, type 1 MGCs from *C. ljungdahlii* and *C. drakei* are upregulated during autotrophic

growth on H_2 + CO_2, a condition where there is increased demand for selenoprotein

biosynthesis (41). Most importantly, during the preparation of this manuscript, we became

aware of prior study that used proteomic and genetic methods to implicate three type I

MGC-encoded proteins in the utilization of dimethyl-selenide as a selenium source in the

hydrogenotrophic methanogen *Methanococcus voltae* (42). The authors suggested that

the Mlps might demethylate (di)methyl-selenide and transfer the methyl group onto the

MGC corrinoid protein, liberating selenide for incorporation into biological

macromolecules. While the authors did not biochemically characterize the Mlps, their data

strongly support a role for the type I MGCs in methyl-selenol utilization.

Fig. 5 Type I MGCs often encode genes involved in selenium metabolism. Selenium metabolism gene

show are either close homologs of genes with experimentally-confirmed roles in selenium metabolism

("Selenium metabolism (known)") or have been associated with selenium metabolism by bioinformatic

methods ("Selenium metabolism (putative)") (36,37). The positions of selenocysteine codons are make with

"TGA." "TGA*" indicates a selenocysteine codon that has been bioinformatically predicted, but not

experimentally confirmed (37). Sec lyase: selenocysteine lyase. DCCP: dicubane cluster protein. DCCPR:

dicubane cluster protein reductase. grdA: glycine reductase subunit A. grdH: glycine betaine reductase

subunit H. selA: *L*-seryl-tRNA(Sec) selenium transferase. selB: selenocysteine-specific translation

elongation factor. selD: selenide water dikinase. selR: putative selenium transcriptional regulator (36). Sao:

putative "selenocysteine-assisted organometallic operon genes.

372   Because type 1 MGCs are present in several model organisms, we examined the

373  expression of these genes in existing proteomic and transcriptomic datasets for *C.*

374  *botulinum*, *C. ljungdahlii*, and *H. thermocellum* to gain additional information regarding

375  their functions. In all three organisms, the MGCs are expressed at a detectable level

376  under all conditions examined. First, we found that the *C. botulinum* MGC is upregulated

377  during growth at the organism's optimal growth temperature of 37 °C compared to growth

378  at 15 °C (transcripts 5.27-68.6 fold upregulated at 37 °C) (43). In *C. ljungdahlii*, we found

379  that the type 1 MGC is transcriptionally upregulated roughly 3-5 fold during autotrophic

380  growth on $H_2 + CO_2$ or CO versus heterotrophic growth on fructose (44–46). The same

381  pattern is found in *C. drakei* (6 to 13 fold transcriptional upregulation on $H_2 + CO_2$

382  compared to fructose) (41). In *H. thermocellum*, the type 1 MGC is upregulated during

383  growth with supplemental acetate in a strain from which the FeFe hydrogenase

384  maturation enzyme *ΔhydG* had been deleted (47). These findings imply that type 1 MGCs

385  are expressed, indicating their physiological relevance, and respond to environmental and

386  genetic perturbations in industrially and medically significant organisms.

387   Finally, we sought characterize the activity of an Mlp to obtain further insight into the

388  function of type I MGCs. Niess et al.(42) observed that deletion of the gene encoding an

389  Mlp2 (*sdmC*) resulted in loss of the ability of *M. voltae* to use dimethyl-selenide as a

390  selenium source, suggesting the function of Mlps2 may be to demethylate dimethyl-

391  selenide and/or methyl-selenol. We therefore selected *Cl*Mlp2 for biochemical assays.

392  Recombinant ClMlp2 was assayed with sodium selenide and methylcob(III)lalamin under

393  a strictly anoxic nitrogen atmosphere as described in the *Materials and Methods*. In brief,

this UV-visible spectroscopy assay assesses methylation of a methyl acceptor by

measuring the disappearance or shift of a large methylcob(III)alamin absorbance band

($\lambda_{max} \sim 525$ nm) when the oxidation and/or ligation state of the cobalt ion is changed. For

convenience, we assayed for methylcob(III)alamin:selenide methyl transfer (Fig 6a), the

reverse of the hypothesized physiological reaction. Addition of sodium selenide (666 μM)

to the reaction mixture resulted in a slow but significant decrease in A_{525nm} (Fig. 6b).

Control reactions lacking either *Cl*Mlp2 or sodium selenide exhibited rates of A_{525nm}

decrease that were 13.7- and 8.3-fold slower, respectively, and are attributable to

artefactual photolysis of methylcob(III)alamin inherent to measurement of absorbance

(48). Addition of sodium sulfide (666 μM) instead of sodium selenide resulted in a 4.3-fold

slower rate, which could suggest the enzyme also possesses low sulfide methylating

activity. Starting and ending spectra of the reactions only showed significant spectral

changes in the selenide and sulfide reactions. In both cases, the spectra showed a shift

in λ_{max} from 520 to 473 nm indicative of formation of base-on cob(II)alamin and therefore

demethylation (Fig. 6c). The net steady state formation of cob(II)alamin with minimal

accumulation of other intermediates (indicated by a steady absorbance trace at the

methylcob(III)alamin/cob(II)alamin isosbestic point of 485 nm (Fig S4)) is likely the result

of rapid oxidation of cob(I)alamin, the expected reaction product. Steady-state kinetic

analysis revealed an apparent k_{cat} of 0.778 min^{-1} and an apparent K_M for selenide of 79.8

μM with a selenide substate inhibition constant (K_I) of 3830 μM (Fig. 6d). The low k_{cat} is

likely due the use of methylcobalamin (rather than the physiological methyl-corrinoid

protein) as a methyl donor. The low selenide K_M, which is comparable to the 46 μM

selenide K_M measured for *E. coli* selenophosphate synthase (49), suggests that methyl-

selenol and/or dimethylselenide are physiological substrates of *Cl*Mlp2. Together, these

biochemical studies of *Cl*Mlp2 support the hypothesis that type I MGCs are involved in

methyl-selenide metabolism.

420
421
Fig. 6 *C. ljungdahlii* (Cl)Mlp2 catalyzes methylcobalamin:selenide methyl transfer a Schematic of the

reaction. Cbl: cobalamin. "Ox." signifies the observed formation of cob(II)alamin, presumably from

nonenzymatic reaction of small amounts of diselenides with highly labile cob(I)alamin. **b** 525 nm

(methylcob(III)alamin peak) traces indicating disappearance of methylcob(III)alamin. **c** Starting (t = 0 min)

and ending (t = 60 min) spectra for the "+ Na₂Se" reaction shown in **b**. **d** Steady state kinetic analysis of

ClMlp2 showing apparent kinetic parameters. Points and error bars represent the mean and standard

deviations from three independent reactions.

Discussion

Methylotrophic methanogenesis is restricted to a subset of the archaea, so it is

surprising to find homologs of enzymes from this pathway in non-methanogens. Here, we

identify a diversified, widespread set of gene clusters—type 1 MGCs—that encode

homologs of the SCMTs involved in methylotrophic methanogenesis. Type I MGCs are

found exclusively in anaerobes and are most common in the Firmicutes phylum. There is

a strong tendency for type I MGCs to be found in genomes that also contain

selenocysteine biosynthesis and decoding machinery.

What reactions do the enzymes encoded by type 1 MGCs catalyze and what is

their physiological function? The presence of a corrinoid protein, a corrinoid reductive

activase, and homologs of corrinoid methyltransferases (Mlps) implies that the type 1

MGCs encode components of a corrinoid methyltransferase pathway. Protein sequence

comparisons indicate that the Mlps are likely *S*- or *Se*-methyltransferases that

(de)methylate thiol and selenol metabolites.

The potential physiological roles of such reactivity may be explored through

considering features of the organisms that possess type I MGCs. Earlier studies clearly

implicate type I MGCs in methyl-selenol valorization (39). However, those authors

examined only the corrinoid protein and two closely-related Mlps2 in dimethyl-selenide

utilization—the more distantly related *M. voltae* Mlp1 and Mlp3 appear to have gone

unnoticed. It seems likely that the full type I MGC contains methyltransferases for

demethylation of several methyl-selenium compounds. While the genetic data in Ref. 39

implicates dimethyl-selenide and/or methyl-selenol as substates of the type I MGC Mlps2

(an inference that our assay of *Cl*Mlp2 supports), we propose that the Mlps1 and Mlps3

may be responsible for demethylation of other methyl-selenium compounds such as

methyl-selenocysteine and selenomethionine and concomitant methylation of the gene

cluster's encoded corrinoid proteins. We propose that the methyl-selenide-derived methyl

group is subsequently transferred from the methylated corrinoid protein to a thiol-

containing compound such as sulfide. The resulting methyl-sulfur metabolite could then

be excreted or recycled into methionine. Forms of methyl-selenium are present in many

environments, including the human gut (50). It may be that most type I MGCs function to

valorize these methyl-selenium compounds. However, many organisms (such as *C.*

ljungdahlii and *M. voltae*) upregulate their type I MGCs even when methyl-selenium

compounds are not added to growth media. This observation suggests that there may be

an endogenous source of methyl-selenium compounds. Indeed, it has been suggested

and evidentially supported that the selenide generated as an intermediate in

selenoprotein biosynthesis can react non-enzymatically with *S*-adenosylmethionine

(SAM) (51). This presents a problem for a cell that has a high demand for selenium

metabolism: methyl selenide formation from SAM wastes valuable selenium and

generates a potentially toxic dead-end metabolite. The enzymes encoded by type I MGCs

may solve this quandary by recycling the methyl selenide into methionine.

Some organisms, such as *H. thermocellum*, encode and express type I MGCs but lack

genes encoding known selenoenzymes or other indicators of selenium metabolism. We

found that all the selenium-trait deficient organisms have Mlps3 that fall into a single clade

of the Mlp phylogenetic tree (Fig. 3b). We suggest that the corresponding type I MGCs

are involved in methyl-sulfur metabolism. While one group of type I MGCs (>90%) is likely

involved in methyl-selenium metabolism, the other, which encodes Mlps3 from the non-

selenium clade, probably has a selenium-independent function.

Evidence from genomic context suggests that the non-selenium MGCs could be

involved in methionine salvage, defined as a metabolic pathway that converts methylthio-

adenosine (MTA) into methionine (52). MTA is a toxic by-product of the biosynthesis of

key metabolites derived from SAM (53). Methionine salvage pathways therefore support

sulfur metabolism both by detoxifying a by-product and preserving the metabolically

expensive methylthiol group of MTA (54). The so-called "universal" methionine salvage

pathway is O_2-dependent, and methionine salvage pathways that operate under

anaerobic conditions have only recently been discovered (52). The anaerobic methionine

salvage pathway from *Rhodospirillum rubrum* was recently shown to use the methylthio-

alkane reductase MarHDK system, homologs of the nitrogenase subunits NifHDK,

respectively (35). Remarkably, *MarHDK*-like genes belonging to the nitrogenase

superfamily IV-A group are prevalent in or near type I MGCs from the non-selenium clade

and, when not present in the type I MGC itself, are invariably present elsewhere in the

genome (Fig. 7a). This suggests that the non-selenium type I MGCs encode enzymatic

machinery for either methionine salvage, sulfur valorization from methyl-sulfide

compounds, or both.

a

Dendrosporobacter quercicolus DSM 1736

Geobacter daltonii FRC-32

Heliobacterium chlorum DSM 3682

Hungateiclostridium mesophilum N2K1

Moorella thermoacetica DSM 21394

Treponema azotonutricium ZAS-9

b

Clostridium sp. CM027

Clostridium aciditolerans DSM 17425

Clostridium algoriphilum DSM 16153

Clostridium sp. CF012

Fig. 7 Type I MGCs from encoding an Mlp3 from the non-selenium clade also encode putative sulfur

and methionine metabolism genes. a Non-selenium type I MGCs associated with group IV-A nitrogenase

superfamily genes and other genes involved in sulfur metabolism. Methylthioribulose-1-P methylsulfurase

is a methionine salvage enzyme originally described in *Rhodospirillum rubrum* (55). **b** Other non-selenium

type I MGCs associated with methionine metabolism. MetY: *O*-acetyl-homoserine sulfhydrylase. CysK:

cysteine synthase. CysI: assimilatory sulfite reductase. CysC: adenylyl-sulfate kinase. AprAB:

phosphoadeninosine-phospho-sulfate reductase subunits A and B. CysND: sulfate adenylyltransferase

subunits 1 and 2. MetC cystathionine beta-lyase. PatB: alternate cystathionine beta-lyase. Ald2: methylthio-

ribulose-1-phosphate aldolase. MtnK: methylthio-ribose kinase. MtnA: methylthio-ribose-1-phosphate

isomerase. MetB: cystathionine gamma-synthase. LuxS: *S*-ribosyl-homocysteine lyase. MetA: homoserine-

O-succinyltransferase. MetF: methylene-tetrahydrofolate reductase. MccA: *O*-acetyl-serine dependent

cystathionine beta-synthase. MetH: cobalamin-dependent methionine synthase. Hom: homoserine

dehydrogenase. SAM: *S*-adenosyl-methionine.

Additionally, the sporadic distribution of type I MGCs across phylogenetic trees is

potentially consistent with a role in methionine salvage, as such pathways are

dispensable under many growth conditions (54) and are not strictly conserved even in

closely related strains (56).

We propose how a type I MGC-encoded methionine salvage pathway could operate

in Fig. 8, considering a type I MGC that encodes only two Mlps. The MarHDK-like

enzymes may have a role in such a pathway, generating methyl-sulfide that might serve

as the methyl donor for the MGC methyltransferase system (R^1-SCH$_3$ with R^1 = H).

Alternatively, the MarHDK-like proteins might serve as part of a second, parallel

methionine salvage pathway, or even as a branch of a topologically bifurcating pathway

with two routes to methionine.

Intriguingly, this proposed methionine salvage pathway uses a distinct biochemical

logic from related pathways. Characterized methionine salvage pathways all incorporate

the methylthiol group of MTA intact into methionine (35,52,53,56). In the proposed

methionine salvage pathway involving the type 1 MGCs, the methylthio-ether of MTA is

dismantled into a methyl group which is carried by the corrinoid protein, leaving the free

thiol behind. This pathway architecture theoretically avoids production of volatile methyl-

sulfur intermediates that could be more readily lost from the cell. The proposed role of a

corrinoid cofactor in methionine salvage is also novel. Ultimately, biochemical

characterization of the enzymes encoded by type I MGCs will be required to test this

proposal.

Fig. 8 Type I MGCs likely encode components of corrinoid-dependent selenium salvage or methionine salvage pathways. The vast majority of type I MGC encoders likely use these methyltransferase systems to recycle methylated selenides formed from the non-enzymatic reaction of SAM

with selenol compounds, potentially recycling the methyl group into methionine. A smaller subset of type I

MGC encoders in the ref seq database (about 10%) lack known selenium metabolism traits. We propose

that these MGCs have evolved to participate in a methionine salvage pathway. MTA: 5'-methylthio-

adenosine. His: histidine. A: adenine.

In summary, we have identified a class of microbial gene clusters encoding

components of probable corrinoid *S*- and *Se*-methyltransferase pathways. The type I

MGCs are found in phylogenetically and metabolically diverse organisms encompassing

>100 microbial genera. Bioinformatic analyses and preliminary biochemical studies

suggest type I MGCs participate in methyl-selenide metabolism and/or methionine

salvage. Ultimately, this previously unappreciated example of methanogen specific

enzymes encoded outside of the archaea could provide insights into the vitally important

fields of sulfur and selenium cycling.

Acknowledgements: D.J.K. acknowledges support from National Institutes of Health

Training Grant #5T32GM095450. This work was supported by a Howard Hughes Medical

Institute (HHMI)-Gates Faculty Scholar Award (OPP1158186) and a National Science

foundation (NSF) Alan T. Waterman Award (CHE-20380529) to E.P.B. E.P.B. is Howard

Hughes Medical Institute Investigator. We also note that this article is subject to HHMI's

Open Access to Publications policy. HHMI lab heads have previously granted a

nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their

research articles. Pursuant to those licenses, the author-accepted manuscript of this

article can be made freely available under a CC BY 4.0 license immediately upon

publication.

Data Availability Statement: All relevant data has been made available in the main text,

Supplementary Information, and associated data files.

References

1. Kurth JM, Op den Camp HJM, Welte CU. Several ways one goal—methanogenesis

from unconventional substrates. Appl Microbiol Biotechnol. 2020 Aug;104(16):6839–

54.

2. Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R. Methanogenic archaea:

ecologically relevant differences in energy conservation. Nat Rev Microbiol. 2008

Aug;6(8):579–91.

3. Chistoserdova L. Wide Distribution of Genes for

Tetrahydromethanopterin/Methanofuran-Linked C1 Transfer Reactions Argues for

Their Presence in the Common Ancestor of Bacteria and Archaea. Front Microbiol

[Internet]. 2016 Sep 13 [cited 2023 May 29];7. Available from:

http://journal.frontiersin.org/Article/10.3389/fmicb.2016.01425/abstract

4. Grinter R, Greening C. Cofactor F420: an expanded view of its distribution,

biosynthesis and roles in bacteria and archaea. FEMS Microbiology Reviews. 2021

Sep 8;45(5):fuab021.

5. Allen JR, Clark DD, Krum JG, Ensign SA. A role for coenzyme M (2-

mercaptoethanesulfonic acid) in a bacterial pathway of aliphatic epoxide carboxylation.

Proc Natl Acad Sci USA. 1999 Jul 20;96(15):8432–7.

6. Venceslau SS, Stockdreher Y, Dahl C, Pereira IAC. The "bacterial heterodisulfide"

DsrC is a key protein in dissimilatory sulfur metabolism. Biochimica et Biophysica Acta

(BBA) - Bioenergetics. 2014 Jul;1837(7):1148–64.

7. Mock J, Wang S, Huang H, Kahnt J, Thauer RK. Evidence for a Hexaheteromeric

Methylenetetrahydrofolate Reductase in Moorella thermoacetica. J Bacteriol. 2014 Sep

15;196(18):3303–14.

8. Huwiler SG, Löffler C, Anselmann SEL, Stärk HJ, von Bergen M, Flechsler J, et al.

One-megadalton metalloenzyme complex in *Geobacter metallireducens* involved in

benzene ring reduction beyond the biological redox window. Proc Natl Acad Sci USA.

2019 Feb 5;116(6):2259–64.

9. Wongnate T, Ragsdale SW. The Reaction Mechanism of Methyl-Coenzyme M

Reductase. Journal of Biological Chemistry. 2015 Apr;290(15):9322–34.

10.   Tallant TC, Paul L, Krzycki JA. The MtsA Subunit of the Methylthiol:Coenzyme M

Methyltransferase of Methanosarcina barkeri Catalyses Both Half-reactions of

Corrinoid-dependent Dimethylsulfide: Coenzyme M Methyl Transfer. Journal of

Biological Chemistry. 2001 Feb;276(6):4485–93.

11.   Oelgeschläger E, Rother M. *In vivo* role of three fused corrinoid/methyl transfer

proteins in *Methanosarcina acetivorans*. Molecular Microbiology. 2009

Jun;72(5):1260–72.

12.   Fu H, Metcalf WW. Genetic Basis for Metabolism of Methylated Sulfur Compounds

in Methanosarcina Species. O'Toole GA, editor. J Bacteriol. 2015 Apr 15;197(8):1515–

24.

13.   Kaufmann F, Wohlfarth G, Diekert G. O-Demethylase from Acetobacterium

dehalogenans . Substrate specificity and function of the participating proteins. Eur J

Biochem. 1998 May;253(3):706–11.

14.   Kurth JM, Nobu MK, Tamaki H, de Jonge N, Berger S, Jetten MSM, et al.

Methanogenic archaea use a bacteria-like methyltransferase system to demethoxylate

aromatic compounds. ISME J [Internet]. 2021 Jun 18 [cited 2021 Aug 8]; Available

from: http://www.nature.com/articles/s41396-021-01025-6

15.   Naidu D, Ragsdale SW. Characterization of a Three-Component Vanillate *O* -

Demethylase from *Moorella thermoacetica*. J Bacteriol. 2001 Jun;183(11):3276–81.

16.   Sauer K, Thauer RK. Methanol:coenzyme M methyltransferase from

Methanosarcina barkeri -- substitution of the corrinoid harbouring subunit MtaC by free

cob(I)alamin: Substitution of *M. barkeri* Mta MtaC by free cob(I)alamin. European

Journal of Biochemistry. 1999 May;261(3):674–81.

17.   Geneious [Internet]. Available from: https://www.geneious.com

18.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

617   throughput. Nucleic Acids Research. 2004 Mar 8;32(5):1792–7.

618   19.   Oberg N, Zallot R, Gerlt JA. EFI-EST, EFI-GNT, and EFI-CGFP: Enzyme Function

619   Initiative (EFI) Web Resource for Genomic Enzymology Tools. Journal of Molecular

620   Biology. 2023 Jul;435(14):168018.

621   20.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape:

622   A Software Environment for Integrated Models of Biomolecular Interaction Networks.

623   Genome Res. 2003 Nov;13(11):2498–504.

624   21.   Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood

625   Trees for Large Alignments. Poon AFY, editor. PLoS ONE. 2010 Mar 10;5(3):e9490.

626   22.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

627   genomes with the Genome Taxonomy Database. Hancock J, editor. Bioinformatics.

628   2020 Mar 1;36(6):1925–7.

629   23.   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al.

630   SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic

631   Acids Research. 2018 Jul 2;46(W1):W296–303.

632   24.   Hoeppner A, Thomas F, Rueppel A, Hensel R, Blankenfeldt W, Bayer P, et al.

633   Structure of the corrinoid:coenzyme M methyltransferase MtaA from *Methanosarcina*

634   *mazei*. Acta Crystallogr D Biol Crystallogr. 2012 Nov 1;68(11):1549–57.

25.   Roberts DL, Zhao S, Doukov T, Ragsdale SW. The reductive acetyl coenzyme A

pathway: sequence and heterologous expression of active

methyltetrahydrofolate:corrinoid/iron-sulfur protein methyltransferase from Clostridium

thermoaceticum. J Bacteriol. 1994 Oct;176(19):6127–30.

26.   Okuda S, Yoshizawa AC. ODB: a database for operon organizations, 2011 update.

Nucleic Acids Research. 2011 Jan 1;39(Database):D552–5.

27.   Matthews RG. Cobalamin-Dependent Methyltransferases. Acc Chem Res. 2001

Aug 1;34(8):681–9.

28.   Siebert A, Schubert T, Engelmann T, Studenik S, Diekert G. Veratrol-O-

demethylase of Acetobacterium dehalogenans: ATP-dependent reduction of the

corrinoid protein. Arch Microbiol. 2005 Sep;183(6):378–84.

29.   Ferguson T, Soares JA, Lienard T, Gottschalk G, Krzycki JA. RamA, a Protein

Required for Reductive Activation of Corrinoid-dependent Methylamine

Methyltransferase Reactions in Methanogenic Archaea. Journal of Biological

Chemistry. 2009 Jan;284(4):2285–95.

30.   Ticak T, Kountz DJ, Girosky KE, Krzycki JA, Ferguson DJ. A nonpyrrolysine

member of the widely distributed trimethylamine methyltransferase family is a glycine

betaine methyltransferase. Proceedings of the National Academy of Sciences. 2014

Oct 28;111(43):E4668–76.

31.   Schilhabel A, Studenik S, Vödisch M, Kreher S, Schlott B, Pierik AY, et al. The
Ether-Cleaving Methyltransferase System of the Strict Anaerobe *Acetobacterium*
dehalogenans: Analysis and Expression of the Encoding Genes. J Bacteriol. 2009 Jan
15;191(2):588–99.

32.   Burke SA, Lo SL, Krzycki JA. Clustered Genes Encoding the Methyltransferases
of Methanogenesis from Monomethylamine. J Bacteriol. 1998 Jul;180(13):3432–40.

33.   Sauer K, Thauer RK. Methyl-coenzyme M formation in methanogenic archaea:
Involvement of zinc in coenzyme M activation. European Journal of Biochemistry. 2000
May;267(9):2498–504.

34.   Krüer M, Haumann M, Meyer-Klaucke W, Thauer RK, Dau H. The role of zinc in
the methylation of the coenzyme M thiol group in methanol:coenzyme M
methyltransferase from *Methanosarcina barkeri*: New insights from X-ray absorption
spectroscopy. European Journal of Biochemistry. 2002 Apr;269(8):2117–23.

35.   North JA, Narrowe AB, Xiong W, Byerly KM, Zhao G, Young SJ, et al. A
nitrogenase-like enzyme system catalyzes methionine, ethylene, and methane
biogenesis. Science. 2020 Aug 28;369(6507):1094–8.

36.   Lin J, Peng T, Jiang L, Ni JZ, Liu Q, Chen L, et al. Comparative Genomics Reveals
New Candidate Genes Involved in Selenium Metabolism in Prokaryotes. Genome
Biology and Evolution. 2015 Mar;7(3):664–76.

37. Haft DH, Gwadz M. Eight Unexpected Selenoprotein Families in Organometallic
Biochemistry in Clostridium difficile, in ABC Transport, and in Methylmercury
Biosynthesis. O'Toole G, editor. J Bacteriol. 2023 Jan 26;205(1):e00259-22.

38. Jeoung JH, Dobbek H. ATP-dependent substrate reduction at an [Fe$_8$S$_9$] double-
cubane cluster. Proc Natl Acad Sci USA. 2018 Mar 20;115(12):2994–9.

39. Böck A, Forchhammer K, Heider J, Leinfelder W, Sawers G, Veprek B, et al.
Selenocysteine: the 21st amino acid. Mol Microbiol. 1991 Mar;5(3):515–20.

40. Wells M, Basu P, Stolz JF. The physiology and evolution of microbial selenium
metabolism. Metallomics. 2021 May 31;13(6):mfab024.

41. Song Y, Lee JS, Shin J, Lee GM, Jin S, Kang S, et al. Functional cooperation of
the glycine synthase-reductase and Wood–Ljungdahl pathways for autotrophic growth
of Clostridium drakei. SYSTEMS BIOLOGY.

42. Niess UM, Klein A. Dimethylselenide Demethylation Is an Adaptive Response to
Selenium Deprivation in the Archaeon *Methanococcus voltae*. J Bacteriol. 2004
Jun;186(11):3640–8.

43. Dahlsten E, Isokallio M, Somervuo P, Lindström M, Korkeala H. Transcriptomic
Analysis of (Group I) Clostridium botulinum ATCC 3502 Cold Shock Response. Popoff
MR, editor. PLoS ONE. 2014 Feb 24;9(2):e89958.

44.   Al-Bassam MM, Kim JN, Zaramela LS, Kellman BP, Zuniga C, Wozniak JM, et al.

692   Optimization of carbon and energy utilization through differential translational

693   efficiency. Nat Commun. 2018 Oct 26;9(1):4474.

694   45.   Tan Y, Liu J, Chen X, Zheng H, Li F. RNA-seq-based comparative transcriptome

695   analysis of the syngas-utilizing bacterium Clostridium ljungdahlii DSM 13528 grown

696   autotrophically and heterotrophically. Mol BioSyst. 2013;9(11):2775.

697   46.   Aklujkar M, Leang C, Shrestha PM, Shrestha M, Lovley DR. Transcriptomic

698   profiles of Clostridium ljungdahlii during lithotrophic growth with syngas or H2 and CO2

699   compared to organotrophic growth with fructose. Sci Rep. 2017 Oct 13;7(1):13135.

700   47.   Biswas R, Wilson CM, Giannone RJ, Klingeman DM, Rydzak T, Shah MB, et al.

701   Improved growth rate in Clostridium thermocellum hydrogenase mutant via perturbed

702   sulfur metabolism. Biotechnol Biofuels. 2017 Dec;10(1):6.

703   48.   Hogenkamp HPC. The Photolysis of Methylcobalamin *. Biochemistry. 1966 Feb

704   1;5(2):417–22.

705   49.   Veres Z, Tsai L, Scholz TD, Politino M, Balaban RS, Stadtman TC. Synthesis of 5-

706   methylaminomethyl-2-selenouridine in tRNAs: 31P NMR studies show the labile

707   selenium donor synthesized by the selD gene product contains selenium bonded to

708   phosphorus. Proc Natl Acad Sci USA. 1992 Apr;89(7):2975–9.

50.   Nogueira CW, Zeni G, Rocha JBT. Organoselenium and Organotellurium

Compounds: Toxicology and Pharmacology. Chem Rev. 2004 Dec 1;104(12):6255–

86.

51.   Fernandes AP, Wallenberg M, Gandin V, Misra S, Tisato F, Marzano C, et al.

Methylselenol Formed by Spontaneous Methylation of Selenide Is a Superior Selenium

Substrate to the Thioredoxin and Glutaredoxin Systems. Netto LES, editor. PLoS ONE.

2012 Nov 30;7(11):e50727.

52.   Albers E. Metabolic characteristics and importance of the universal methionine

salvage pathway recycling methionine from 5′-methylthioadenosine. :11.

53.   Sauter M, Moffatt B, Saechao MC, Hell R, Wirtz M. Methionine salvage and S-

adenosylmethionine: essential links between sulfur, ethylene and polyamine

biosynthesis. Biochemical Journal. 2013 Apr 15;451(2):145–54.

54.   Miller AR, North JA, Wildenthal JA, Tabita FR. Two Distinct Aerobic Methionine

Salvage Pathways Generate Volatile Methanethiol in Rhodopseudomonas palustris.

Cavanaugh CM, editor. mBio. 2018 May 2;9(2):e00407-18.

55.   Warlick BP, Evans BS, Erb TJ, Ramagopal UA, Sriram J, Imker HJ, et al. 1-

Methylthio- D -xylulose 5-Phosphate Methylsulfurylase: A Novel Route to 1-Deoxy- D -

xylulose 5-Phosphate in *Rhodospirillum rubrum*. Biochemistry. 2012 Oct

23;51(42):8324–6.

56.   North JA, Wildenthal JA, Erb TJ, Evans BS, Byerly KM, Gerlt JA, et al. A

bifunctional salvage pathway for two distinct S-adenosylmethionine by-products that is

widespread in bacteria, including pathogenic *Escherichia coli*. Mol Microbiol. 2020

May;113(5):923–37.

735