1 **A conserved odorant receptor underpins borneol-mediated repellency in culicine**
2 **mosquitoes**
3

4 Yuri Vainer[2], Yinliang Wang[2,7], Robert M. Huff[1], Dor Perets[2], Evyatar Sar-Shalom[2], Esther
5 Yakir[2], Majid Ghaninia[5], Iliano V. Coutinho-Abreu Gomes[3], Carlos Ruiz [6], Dhivya
6 Rajamanickam, A. Warburg[4], Omar S. Akbari[3], Philippos A. Papathanos[2], R. Ignell[5], Jeffrey
7 A. Riffell[6], R. Jason Pitts[1*], Jonathan D. Bohbot[2*]
8
[1]Department of Biology, Baylor University, USA
[2]Department of Entomology, The Hebrew University of Jerusalem, Israel
[3]Division of Biological Sciences, University of California, USA
[4]Department of Microbiology and Molecular Genetics, The Hebrew University of Jerusalem,
Israel
[5]Department of Plant Protection Biology, Swedish University of Agricultural Sciences,
Sweden
[6]Department of Biology, University of Washington, USA
[7]Northeast Normal University, China
18
19 *Corresponding authors: jonathan.bohbot@mail.huji.ac.il & Jason_Pitts@baylor.edu

20
21 **Keywords:** mosquitoes, monoterpenoids, borneol, camphor, repellent, odorant receptor
22
Abstract

The use of essential oils derived from the camphor tree to repel mosquitoes is an ancient
practice that originated in Southeast Asia and gradually spread to China and across Europe
via the Maritime Silk Road. The olfactory mechanisms by which these oils elicit avoidance
behavior are unclear. Here we show that plant bicyclic monoterpenoids and borneol
specifically activate a neural pathway that originates in the orphan olfactory receptor neuron
of the capitate peg sensillum in the maxillary palp, and projects to the mediodorsal
glomerulus 3 in the antennal lobe. This neuron co-locates with two olfactory receptor neurons
tuned to carbon dioxide and octenol that mediate human-host detection. We also confirm that
borneol elicits repellency against human-seeking female mosquitoes. Understanding the
functional role of the mosquito maxillary palp is essential to investigating olfactory signal
integration and host-selection behavior.

35
Main
The use of plants to ward off insects has been a human practice since prehistorical times [1] and
is still used in many parts of the world [2,3]. Plant-based essential oils (EOs), including lemon
eucalyptus leaf oil [4], citronella oil [5], and coconut oil [6] exhibit different degrees of mosquito
repellency due to the presence of pyrethrins [7], phenol derivatives, and terpenoids [8]. The latter
includes oxygen-containing compounds with open-chain hydrocarbons, such as linalool,
citronellol, geraniol, and bicyclic derivatives such as cineole, fenchone, camphor and borneol
(Fig. 1a). The EO of the camphor tree *Cinnamomum camphora* is composed of over 70
volatile organic compounds (VOCs), most of which are oxygenated monoterpenes dominated
by camphor and other related compounds [9]. Camphor and borneol extracts are believed to
have originated from the camphor tree in the island of Borneo (Fig. 1a), where they were
initially traded with China and then introduced to the West due to their therapeutic,
refreshing, and repellent effects against mosquitoes [10]. How these terpenoids molecules exert
repelling effects against mosquitoes is not well-understood but is likely mediated by their
olfactory system.

The capitate-peg (cp) sensillum located on the mosquito maxillary palp comprises three
olfactory neurons, each distinguishable by size, olfactory receptor gene expression profile,
and odor response characteristics (Fig. 1b). In both culicine and anopheline mosquitoes, the
largest olfactory sensory neuron (cpA) expresses three gustatory receptors (*Grs*) that
specifically detect CO_2 [11,12]. The medium-sized cpB neuron of *Anopheles gambiae* and *Aedes*
aegypti expresses the 1-octen-3-ol odorant receptor *Or8* and its co-receptor Orco [13–15]. Both
CO_2 and 1-octen-3-ol elicit attraction and signal the presence of animal hosts in anopheline [16]
and culicine [17] mosquitoes. This cellular and functional organization have remained
remarkably conserved over 180 million years of mosquito evolution (Fig. 1b).
60
In anophelines, the small cpC neuron expresses the *Or28* gene [13], which responds to plant
volatile organic compounds (pVOC) [18]. In culicines, the cpC neuron expresses the *Or49* gene
[19], which is unrelated to *AgamOr28* (Fig. 1b). Using a pharmacological approach, we
expressed *Or49* in a heterologous expression system and exposed it to odorant mixtures, EOs
and individual volatile odorant compounds. Our findings provide strong evidence that the
OR49 receptor and the cpC neuron respond to plant-derived bicyclic monoterpenoids with a
marked selectivity towards borneol, traditionally used as a mosquito repellent. We observed
that a loss-of-function mutation in the *Or49* gene leads to a lack of electrophysiological
responses when stimulated by borneol. Furthermore, we show that borneol elicits selective
activation of the MD3 glomerulus in the antennal lobe of *Ae. aegypti*, indicating that all 3 cp
neurons project to the same brain area. Confirming our pharmacological and
electrophysiological results, our behavioral study reveals that borneol induces repellency in
human-seeking female mosquitoes. our findings lay the groundwork to understand the
detailed molecular and neural basis shaping olfactory integration and governing host
selection in mosquitoes.
76
Odorant receptor 49 is an evolutionary conserved borneol receptor
The odorant receptor 49 gene[18,19] is conserved in *Aedes albopictus*, *Culex quinquefasciatus,*
Toxorhynchites amboinensis and *Anopheles gambiae* (Supplementary Fig. 1). Phylogenetic
analyses support previous studies [19,20] describing the OR49 family as a Culicinae-specific
group (Supplementary Fig. 1c–d) that includes the more distantly related anopheline
OR48/49 group. The *Or49* genes are located on chromosome 2 in both *Aedes* and *Anopheles*
mosquitoes (Supplementary Fig. 1e) and exhibit conserved syntenic relationships with
neighbouring genes (Supplementary Fig. 1f).
85
As a continuation of our work on the functional identity of the cp sensillum in culicine
mosquitoes [19,21], we investigated the receptive field of *Ae. albopictus* OR49 (AalbOR49)
using two-electrode voltage clamp recordings (Fig. 1c–d) and a panel of 81 odorants
representing a variety of chemical classes (Supplementary Table 1). Oocytes expressing
functional receptor complexes were strongly activated by blends containing ketonic
compounds (Fig. 1c). Individual testing of the single compounds from the ketone blend
revealed the efficacy of bicyclic monoterpenoids, with camphor emerging as the most
efficacious member followed by cineole (also called eucalyptol) and fenchones (Fig. 1d).
Subsequently, we screened OR49 from the nectar-feeding *Tx. amboinensis* (*TambOR49*) with
8 *Cannabis* EOs (Fig. 2a, Supplementary Table 1) containing 42 plant VOCs dominated by
terpenes (Supplementary Fig. 2). The three most active mixtures were Pineapple Haze, OG
Kush, Adom #9 followed by Master Kush with a much lower efficacy (Fig. 2a), all evoking
consistent current responses (Supplementary Fig. 3). Successive fractionations of these EOs
down to 6 sub-mixtures and 4 single compounds were then administered of which the
racemic borneol showed the highest potency (Fig. 2b), consistent with the borneol content in

the original *Cannabis* mixtures (Fig. 2a, Supplementary Fig. 4a) and sub-mixtures
(Supplementary Fig. 4b).
103
In *An. gambiae*, the cpC neuron expresses OR28 (Figure 1a), which is activated by plant
VOCs, with acetophenone and 2,4,5-trimethylthiazole being the most effective ligands in the
Xenopus oocyte expression system [13,18]. The EC_{50} values of acetophenone, α-pinene, and α-
terpineol were in the low millimolar range indicating that more potent ligands remain to be
identified (Supplementary Fig. 5). Borneol did not elicit any response at the tested
concentrations suggesting that AgamOR28 is tuned to VOCs belonging to a different
chemical class.
111
TambOR49 and AaegOR49 did not exhibit enantioselectivity towards the (+) and (–)-borneol
(Figure 2c) as indicated by their EC_{50} values in the one-digit micromolar range. We
established concentration-response relationships between TambOR49, *Cx. quinquefasciatus*
OR38 (CquiOR49), *Ae. aegypti* OR49 (AaegOR49), and *Ae. allbopictus* OR49 (AalbOR49),
for the two most potent ligands, (+)-borneol and (+)-camphor (Fig. 2d). (+)-Borneol was 19-
32 times more potent than (+)-camphor in all cases except for TambOR49 in which both
compounds were equally active in the low micromolar range. In all four examined culicine
species, including the nectar-feeding *Tx. amboinensis*, (+)-borneol elicited significantly
higher responses than (+)-camphor across all concentrations (Fig. 2e, Supplementary Fig. 3).
This data indicates that OR49 is a selective borneol receptor.
122
The *Or49* gene confers sensitivity to borneol
To investigate the olfactory effect of camphor and borneol *in vivo*, we conducted
electropalpogram (EPG) recordings on three culicine (*Ae. albopictus*, *Ae. aegypti*, *Cx.
pipiens*) and one anopheline (*An. gambiae*) species. 1-Octen-3-ol, which served as a positive
control, induced significant responses in all mosquitoes tested. Camphor and borneol elicited
consistent palp responses in Culicinae mosquitoes, including *Ae. albopictus*, *Ae. aegypti*, and
Cx. pipiens but not in *An. gambiae* (Fig. 3a). Camphor and borneol evoked lower responses
than 1-octen-3-ol in the culicine palps, which is consistent with the smaller size of the cpC
neuron in comparison to the cpB neuron. In *Ae. albopictus*, (–)-borneol was more active than
(+)-camphor, while in *Ae. aegypti* both borneol enantiomers elicited significantly greater
EPG responses than camphor enantiomers. In *Cx. pipiens*, borneol and camphor elicited
comparable EPG responses. We established dose-response relationships with (+)-camphor
and (+)-borneol in *Ae. albopictus*, *Ae. aegypti*, and *Cx. pipiens* (Fig. 3b). Among the three
culicine species, *Cx. pipiens* exhibited the largest responses in response to (+)-camphor.
However, we did not find any significant statistical differences in activity between camphor
and borneol in the tested species.
139
To test a potential causal gene-function relationship between *Or49* and the borneol response,
we used homology directed repair (HDR) to knock-in a transgenic construct containing the
QF2 transactivator and a ubiquitously expressing eCFP fluorescent marker driven by the
OpiE2 promoter at the start codon of the *Or49* gene in *Ae. aegypti* (Supplementary Fig. 6).
To investigate the genetic mechanism determining borneol sensitivity in the palp, we
conducted single-sensillum recordings (Fig. 4a) from the cp sensillum of *Ae. aegypti* (Fig.
4b), *An. gambiae* (Fig. 4c), *Ae. albopictus* (Fig. 4d), and *Cx. quinquefasciatus* (Fig. 4e). The
response of the cpA and cpB neurons to CO_2 [22] (data not shown) and *R*-(–)-1-octen-3-ol [23],
respectively, has been well characterized in previous studies [13–15]. (+)-Borneol elicited a dose-
dependent response in the cpC neuron of *Ae. aegypti*, *Ae. albopictus*, *Cx. quinquefasciatus*
but not in *An. gambiae s.s.* (Fig. 4f). Single sensillum recordings identified the spontaneous

activity of three sensory neurons, distinguished by differences in spike amplitudes, housed in
the capitate peg sensilla of wild type *Ae. aegypti* (Fig. 4c) and *An. gambiae* (Fig. 4d). No SSR
responses were recorded from the cpC neuron in the *Ae. aegypti Or49* null mutant line in
response to (+)-borneol (Fig. 4c) indicating that *Or49* is sufficient and necessary to confer the
cpC neuron sensitivity to borneol. The lack of response of the cpC neuron in *Anopheles* (Fig.
4d and f) is consistent with the expression of *Or28* instead of *Or49*, which is not activated by
borneol (Supplementary Fig. 5). The absence of OR28 activation and *An. gambiae* palp EPG
responses by borneol and camphor support this conclusion.
159
Borneol and camphor specifically activate the MD3 glomerulus in the antennal lobe
To examine how *Ae. aegypti* processes borneol in the antennal lobe (AL), two-photon
imaging experiments were conducted using pan-neuronal GCaMP-expressing *Ae. aegypti*
mosquitoes. We utilized existing mosquito lines that contained a *QUAS-GCaMP6s* transgene
crossed with the *brp-QF2* driver line [24], allowing the directed expression of the calcium
indicator GCaMP6s in all neurons of the AL. Mosquitoes were glued to holders that
permitted two-photon imaging of calcium responses in the AL (Fig. 5a), allowing repeatable
registration of the AL glomeruli between preparations (Fig. 5b) and recording of glomerular
responses (Fig. 5c) [25]. Glomeruli from our two-photon imaging results were mapped to AL
atlases [26,27], allowing accurate identification of the glomeruli of interest in their spatial
context (Extended Data Fig. 5). We focused on the mediodorsal (MD) glomeruli, as these
glomeruli receive input from the capitate peg sensilla of maxillary palps (Fig. 5b and c) and
are responsive to host odors, including 1-octen-3-ol (MD2) and CO_2 (MD1) [27].
173
To determine the odor coding of the MD glomeruli, and identify the cognate glomerulus
representing borneol, we first recorded from the MD1-3 glomeruli while stimulating with
CO_2 (5%), 1-octen-3-ol (10^{-4} dilution), borneol (10^{-4} dilution), and the solvent control. For
the MD3 glomerulus, (+)-borneol elicited strong, tonic responses that were significantly
greater than the solvent control (P<0.001) (Fig. 5d and g). By contrast, responses of the MD1
and MD2 glomeruli to borneol were not significantly different from the control (P=0.09 and
0.23, respectively). The MD2 glomerulus showed the greatest responses to 1-octen-3-ol (Fig.
5e) (P = 0.00003), and the MD1 glomerulus to CO_2 (P=0.01) (Fig. 5f). Given MD3's
response to (+)-borneol, and OR49/cpC's responses to other terpene compounds, we next
examined how this glomerulus responded to a limited panel of different odorants, including
enantiomers of borneol and camphor. From this panel, (–)-borneol elicited the greatest
response in the MD3 glomerulus (Fig. 5g), closely followed by (+)-borneol, enantiomers of
camphor, and geosmin. These findings are consistent with previous studies demonstrating
that the three MD glomeruli receive input from the maxillary palp cp neurons whereby the
MD1 glomerulus receiving CO_2 input from the cpA neurons [28] while the medium size MD2
glomerulus receives 1-octen-3-ol input from the cpB neurons [27].
190
Borneol repels human-host seeking female *Ae. aegypti*
To examine the impact of (±)-borneol on blood-seeking *Ae. aegypti* mosquitoes, we
conducted an arm-in-a-cage assay (Extemded Data Fig. 1a), exposing 15 females to human
skin odor for 10 minutes (Figure 6). A protective glove covered the hand, allowing
mosquitoes to detect the odor through a dorsal open area, which was protected with a screen
and equipped with a chemical holder for the deposition of (±)-borneol (1M) (Extended data
Fig. 1). To monitor host-seeking behavior—specifically, the presence of females, their
walking path (in cm), and visit duration in the region of interest (ROI)—we fine-tuned a
custom object detection YOLOv8 model [29]. Our results demonstrated a significant reduction
(54%) in the number of trajectories within the ROI when the hand was treated with (±)-

borneol compared to the vehicle-treated hand (Fig. 6a). All trajectories across treatments and repetitions are shown in Figure 6b, where different colors indicate different trajectories. Vehicle and borneol-induced trajectories were 951 and 428, respectively This trend was consistent across time (Fig. 6c). On average, the total time spent in the ROI was three times higher in the control than in the (±)-borneol treatment (Fig. 6d). Additionally, the distance walked within the ROI was significantly lower in the presence of (±)-borneol (Fig. 6e). Figure 6f visually represents the mosquito detections throughout the experiment, highlighting the differences between the control and (±)-borneol treatments. A video sample is available in Extended Data Fig. 6. When purifying the *Or49* homozygote mutant, we noticed severe reduction in fitness, which precluded further behavioral experiments.

Discussion

We have identified a conserved mechanism in culicine mosquitoes responsible for the selective detection of borneol, a bicyclic monoterpenoid that has been used as an insect repellent since ancient times. Recent studies have shown that borneol is a broad-acting insect repellent against fungus gnats [30], the booklouse *Liposcelis bostrychophila* [31] and fire ants [32] suggesting that more than one olfactory-mediated mechanisms in insects are at play since *Or49* homologs in these insects have not been identified. Interestingly, borneol elicits responses in the *Ae. aegypti* antenna as well [33]. However, antennal activation by borneol requires doses 10,000 times greater than in the maxillary palp (Supplementary Fig. 7). This difference in sensitivity in favor of the palp indicates that whatever mechanism is involved in the antenna is likely not an ecologically-relevant response but rather the product of chemical overstimulation. Moreover, this alternative mechanism is *Or49*-independent since its expression has not been reported in the antenna.

In *Anopheles*, the antennal-expressed *Or48* [34], one of the two closest homologs to the culicine *Or49* (Supplementary Fig. 1c), responds to straight chains alcohols, ketones, and acetates [18,35] suggesting that this receptor is tuned to a different class of compounds setting apart the olfactory coding logic between these two mosquito subfamilies. Contrary to OR8, which discriminates between the 1-octen-3-ol enantiomers [21], the two borneol enantiomers elicited comparable activations at the pharmacological, physiological and AL levels. These findings suggest that the receptor binding pocket accommodates both forms of borneol or that a closely related structural analog of borneol, with greater potency, remains to be identified[36].

Our results suggest that in *Ae. aegypti* and other culicine mosquitoes, the repellent properties of traditional Chinese medicinal plants such as the camphor tree EO are largely mediated by the activation of OR49 by bicyclic monoterpenoids and more specifically by borneol. Borneol repels human host-seeking *Ae. aegypti*, *Cx. quinquefasciatus* and *An. stepehensi* [33,37,38] while its presence in EOs attracts gravid *Ae. aegypti* [39], suggesting that this compound has opposite effects on animal-host-seeking behavior and oviposition preferences. The reason for borneol being a signal mediating repellency is puzzling. While several monoterpenoids such as 1-8-cineol, limonene, and fenchone exhibit anticholinesterase activity [40,41] causing paralysis and death in a variety of insects, there are conflicting reports on the larvicidal activity of borneol and camphor against mosquito larvae [39,42–45]. Future studies should focus on the ecological role of borneol in different contexts, including foraging and oviposition site selection.

We found that the co-location of the three cp neurons in the cp sensillum is reflected by the anatomical proximity of their projections in the MD glomeruli cluster in the AL. Whether the grouping of these three glomeruli has a functional significance and why the cp sensillum

houses ORNs tuned to animal and plant host olfactory cues should be the focus of future
studies. The two largest cp neurons are tuned to animal-host attractants whereas the smallest
and third cpC neuron responds to a plant-host odorant with repellent activity. This
antagonism in valence whereby an attractive signal is detected by the largest of two ORNs
within the same sensillum while a deterring signal is detected by the smallest ORN is
widespread in animals[46]. We surmise that the main function of the mosquito cp sensillum is
to detect animal host odorants, while the cpC neuron exerts a presynaptic inhibitory effect
mediated by lateral coupling when activated by borneol.
259
Our findings indicate that the cp sensillum in the palp of culicine mosquitoes detects signals
from animal and plant hosts. Why OR49 is selectively tuned to this particular compound and
not to a more ubiquitous volatile phytochemical remains a mystery. Further experiments will
reveal how these three signals are integrated at the pre-synaptic level within the sensillum and
in the AL. Understanding the ecological and neurological significance of the mosquito cp
sensillum, i.e., the packaging of animal and plant host detectors within the same sensillum,
may be exploited to design novel repellent formulations.
267
Methods
Insects
Our mosquito colonies were reared according to previously published protocols [47]. *Ae.*
aegypti originated from a colony established by Prof. Joel Margalit. The *Ae. aegypti*
Liverpool strain was provided by the Akbari lab. *Aedes albopictus* was the FPA Foshan strain
collected (Foshan, China) reared at the insectary of the University of Pavia since 2013
(Palatini et al., 2017). *Anopheles gambiae* was the G3 strain originally isolated from West
Africa (MacCarthy Island, The Gambia) in 1975 (Federica Bernardini et al., 2017). *Culex*
pipiens originated from a wild type population was provided by Dr. Laor Orshan (Ministry of
Health, Israel).
278
Phylogenetic and genomic analyses
The AaegOR49 protein sequence (AAEL001303-PA) was used as a query to identify
homologs in other mosquito species. Amino acid sequences of homologous odorant receptors
were obtained from the Vectorbase database (vectorbase.org) and the *Tx. Amboinensis*
genome assembly (Zhou et al., 2014). The multiple sequence alignment was conducted using
ClustralW (Chenna et al., 2003). The Maximum-likelihood phylogenetic tree was constructed
using MEGAX software (Model: G, bootstraps: 5000). Exon-intron structure analysis was
conducted manually by aligning predicted amino-acid sequences to genomic regions.
Approximate mapping of the genes in chromosome location for each mosquito species and
construction of synteny maps done using Geneious prime software (Kearse et al., 2012) using
latest genome assemblies available for each mosquito species (Matthews et al., 2018, Palatini
et al., 2020, Boyle et al., 2021, Arensburger et al., 2012, Sharakhova et al., 2007).
Genomic DNA (gDNA) was isolated from whole bodies of *Tx. amboinensis* adults. gDNA
was diluted to 25ng/µL in nuclease free water and used as a template in polymerase chain
reactions to amplify odorant receptors (TambOR) using Taq polymerase and the following
primer sets: TambOr6.F1 (ATGCGCTTCTACGAGAAATAC), TambOr6.R1
(TCAGAAATTATCCTTCAGGATC); TambOr12.F1 (ATGCCATCGGTTTTCTTGGTT),
TambOr12.R1 (CTAAAACACTCGCTTCAATATC); TambOr13.F1
(ATGTTCTGCTTCAGGAAGATC), TambOr13.R1
(CTAGAAGTGGTTTTTCAATATAA); TambOr49.F1
(ATGTTGTTCAAGAACTGTTCC), TambOr49.R1
(TTAATAATTGAATCTTTCCTTCAG); TambOr71.F1

(ATGGGCAGCAGTGATGGTGAC), TambOr71.R1 (CTACTGGTTGATTTTACTGAGG).
Cycling conditions were 94°C for 60s; 30 cycles of 94°C for 20s, 56°C for 20s, 72°C for 30s;
and a final extension of 72°C for 5 minutes. Amplicons were analyzed by electrophoresis on
a 1% agarose gel, cloned into the TOPO-TA pCR2 plasmid, heat-shock transformed into
TOP10 competent *E. coli* cells, and grown overnight at 37°C on LB+ampicillin+X-gal agar
plates. Ampicillin resistant, white colonies were isolated using a sterile pipette tip and grown
overnight 37°C in 3mL of LB+ampicillin. Plasmids were isolated by DNA miniprep and the
Sanger method was used to determine the DNA sequence in both directions using standard
T7 and M13.rev primers. TambOR nucleotide sequences were compiled into contigs and
intron/exon regions were inferred by comparing to coding sequences described in a previous
publication [48].
312
Chemical reagents
The chemicals used for the deorphanization of receptors were obtained from Acros Organics
(Morris, NJ, USA), Alfa Aesar (Ward Hill, MA, USA), ChemSpace (Monmouth Junction,
NJ, USA), Sigma Aldrich (St. Louis, MO, USA), TCI America (Portland, OR, USA), and
Thermo Fisher Scientific (Waltham, MA, USA) and Penta Manufacturing Corp. (Livingston,
NJ, USA) at the highest purity available. *Cannabis* essential oils and sub-mixtures were
formulated and supplied by Eybna Technologies (Givat Hen, Israel) (Table S1 and
Supplementary Fig. 2–3).
321
Two-electrode voltage clamp of *Xenopus laevis* oocytes
In vitro transcription and two-microelectrode voltage clamp electrophysiological recordings
were carried out as previously described [49]. Experimental procedures for the *Ae. albopictus*
receptor clone was performed as previously described [50]. *AalbOr49* and *AalbOrco* templates
were synthesized by Twist Biosciences (San Francisco, CA, USA) and cloned into the
pENTR^{TM} vector using the Gateway^R directional cloning system (Invitrogen Corp., Carlsbad,
CA, USA) and subcloned into the *X. laevis* expression destination vector pSP64t-RFA.
Several of these genes were codon-optimized (Supplementary Table 2). For pairwise current
comparisons of (+)-borneol and (+)-camphor, two separate perfusion systems were
assembled, one for each compound. The outlet from each system was connected to a 2 to 1
perfusion manifold. At each tested concentration, (+)-borneol and (+)-camphor were
administered consecutively. Representative current traces can be seen in Supplementary
Figure 4. *Odorant receptor 49* and *Orco* clones and recording measurements can be found in
Supplementary Table 2 and 3, respectively. The use of *Xenopus laevis* frog eggs was carried
out according to The Hebrew University of Jerusalem Ethics committee.
337
Electropalpogram recordings
Based on the voltage clamp data, responses from the maxillary palps of *Ae. albopictus*, *Ae.
aegypti*, *Cx. pipiens* and *An. gambiae* were recorded with camphor or borneol.
Electopalpogram (EPG) assays were performed according to published procedures [51] with a
few modifications. Within the Pasteur pipette, 20 µL of odorants diluted in hexane were
deposited at desired dose onto a 4×20 mm Whatman filter paper strip. The odorants were
delivered into a consistent humidified airstream (at a flow rate of 50 cm/s) at approximately 2
cm from the maxillary palps. The pulse duration was 0.1 s, and the recording time was set for
5 s. A 2-min gap was allowed between stimuli to recover the EPG sensitivity. At least 5
individual females were tested, for each individual, 3 technical replicates were performed
with doses ranging from 0.01 ug to 100 µg. Negative controls (hexane and air) and positive
controls (1-octen-3-ol) were performed at both the beginning and end for each recording

session to monitor the decline in sensitivity of the maxillary palps. EPG peak responses were
normalized to the response of hexane.
352
Generating *Or49* knockout line in *Ae. aegypti* and isolating an *OR49*[-/-] homozygous line
To knock out the gene encoding the *Ae. aegypti* OR49 receptor, we used homology directed
repair (HDR) to knock in a transgenic construct containing the QF2 transactivator and a
ubiquitously expressing eCFP fluorescent marker driven by the OpiE2 promoter at the start
codon of the *Or49* gene (Supplementary Fig. 6a–b). HDR was triggered by double strand
breaks at the *Or49* promoter and the 1[st] exon mediated by two guide RNAs, whose cleavage
activities had been tested *in vitro* (Supplementary Fig. 6c). Genomic DNA was extracted
from whole bodies of *Ae. aegypti* Liverpool individuals using the DNeasy Blood & Tissue
Kit (Qiagen, Redwood City, CA). Homology arms of the *Or49* gene flanking 1 kb upstream
and downstream of the desired insertion site (Supplementary Fig. 6a) were amplified with the
Q5 High Fidelity DNA polymerase (New England Biolabs, Ipswich, MA), using primer pairs
Left_Arm_Or49_FWD and Left_Arm_Or49_REV as well as Right_Arm_Or49_FWD and
Right_Arm_Or49_REV, respectively (Supplementary Table 4). The QF2-ECFP DNA
cassette was amplified using the V1117A plasmid as template and the primer pair
QF2_OpIE_ECFP FWD and QF2_OpIE_ECFP REV. The cassette contains the *QF2*
sequence and the 3' UTR of the *HSP70* gene along with the *ECFP* gene under the control of
the *Opie2* promoter and the 3' UTR of the *SV40* gene. PCR bands were cut out from agarose
gels and purified with Zymoclean Gel DNA Recovery Kit (Zymo Research, Irvine, CA).
Upstream and downstream homology arms as well as the QF2-ECFP cassette PCR products
were assembled into the backbone of the V1117A plasmid (Supplementary Fig. 6b) using
Gibson assembly reaction, following the manufacturer recommendations. Gibson's reaction-
derived plasmid product was used to transform JM109 cells (Zymo Research), and colonies
were individually grown overnight for minipreps with Zyppy Plasmid Miniprep Kit (Zymo
Research). The sequence of the plasmid V1117F-Or49 was confirmed by restriction enzyme
digestion using the enzymes ScaI-HF and AvrII as well as by whole plasmid sequencing
(Primordium Labs, Monrovia, CA). V1117F-Or49 plasmid was used to retransform JM109
cells, which were grown for maxiprep purification using the PureLink Expi Endotoxin-Free
Maxi Plasmid Purification Kit (Thermo Fisher Scientifics, Waltham, MA).
For guide RNA synthesis, non-template reactions were carried out with the
gRNA_Left_Or49_F and gRNA_Left_Or49_R forward primers and the universal guide RNA
reverse primer (Universal-sgRNA_R). PCR bands were isolated from agarose gel and
purified as above described. Guide RNAs were synthesized with the Ambion MEGAscript kit
(Thermo Fisher Scientifics) for 4 hours at 37°C, using 300 ng of purified PCR product. Guide
RNAs were further purified with the Megaclear Kit (Thermo Fisher Scientifics).
To assess the cleavage activity of the synthesized guide RNAs, *in vitro* Cas9 cleavage assays
were carried out. A DNA fragment spanning 1,009 bp overlapping the cleavage sites of the
guide RNAs was amplified with primers Or49_cleav_F and Or49_cleav_R, DNA bands were
isolated from agarose gel, purified, and 100 ng of which was used in cleavage assays with
300 ng of recombinant Cas9 (PNA BIO, Thousand Oaks, CA) and 100 ng of each guide RNA
upon incubation at 37°C for 1 hour (Supplementary Fig. 6c).
For embryo microinjection, an injection mix was prepared with the plasmid V1117F-Or49 at
500 ng/ul, gRNAs left and right at 100 ng/ul each, and recombinant Cas9 at 300 ng/ul. Mix
was filtered with the Ultrafree-MC Centrifugal Filter UFC30GUOS (Millipore, Burlington,
MA). Freshly harvested *Ae. aegypti* embryos (Liverpool strain) were injected with the
transformation mix using a custom-made insect embryo microinjector (Hive Technologies,
Cary, NC) at 100 psi pulse and 3 psi back pressures, using quartz needles pulled with a P-
2000 needle puller (Sutter Instrument Co., Novato, CA). Genomic DNA of the G1

fluorescent mosquitoes (Supplementary Fig. 6d) were individually extracted with the Qiagen
blood kit, and PCR amplified for Sanger Sequencing reactions (Retrogen, San Diego, CA)
using the primer pairs Or49_diag_up_F and Or49_diag_up_R targeting the upstream and
primer pair Or49_diag_down_F and Or49_diag_down_R targeting the downstream insertion
site (Supplementary Fig. 6e). For whole DNA cassette sequencing (Primordium), 2.5kb DNA
fragments were amplified from the DNA of G1 individuals using the primer pair
Or49_diag_up_F and Or49_diag_down_R (Supplementary Fig. 6f).
For isolation of homozygous knockout individuals, pupae were sex sorted, and single pairs
were transferred into Narrow *Drosophila* Vials (Genesee Scientific, El Cajon, CA), filled
with 10 mL of deionized water (DI) and closed with cotton stoppers. Upon emergence, water
was drained, and sugar cottons were placed into the vials. Mosquitoes were allowed to mate
for 5 days in the vials, and then all individuals were transferred to a single cage for blood
feeding. Three days after blood feeding, females were individually transferred to a fresh vial
containing a piece of brown paper towel wetted with 1 mL DI water. Females were allowed
to lay eggs, and each egg batch was individually hatched in 250 mL Clear Pet Cups (9oz).
Larvae were screened for fluorescence, and the batches that resulted in 100% fluorescent
individuals were grown to adulthood and intercrossed to confirm homozygosity in the
following generation. Two out of thirty egg batches resulted in homozygous offspring. To
further confirm the homozygous status of these individuals, genomic DNA of three pools of
10 males, 10 females, and a mix of males and females were PCR amplified with the primer
pair Or49_diag_up_F and Or49_diag_down_R, which resulted in a single band for
homozygous individuals and two bands for heterozygous individuals (Supplementary Fig.
6g). Out of 739 injected embryos, 5.5% survived to the pupal stage. These G_0 individuals
were sex sorted and outcrossed with wildtypes. Sixteen G_1 larvae displayed blue (cyan)
fluorescent bodies (Supplementary Fig. 6d). Insertion of the transgenes within *Or49* was
confirmed in multiple G_1 individuals (Supplementary Fig. 6e–f) by sequencing of the
insertion sites as well as the transgene (Supplementary Fig. 6e–f) resulting in a nonfunctional
receptor. An *Or49* $^{-/-}$ homozygous mosquito strain was isolated by single-pair mating,
screening the fluorescence marker in the offspring for 100% penetrance, and PCR
confirmation for the presence of homozygous mutant alleles (Supplementary Fig. 6g).
430
Single-sensillum electrophysiology
Single sensillum recordings from capitate peg sensilla on the maxillary palp of wild type *Ae.*
aegypti (Liverpool)*, Ae. albopictus* (FPA)*, Cx. quinquefasciatus* (Thai) and *An. gambiae*
sensu stricto (G3) and *Ae. aegypti Or49*$^{-/-}$ mutant mosquitoes were performed using an
established protocol [52]. Spikes were quantified offline using the established nomenclature for
the sensory neurons [53]. The number of spikes counted during a 0.5 s stimulus delivery
interval was subtracted from the number of spikes counted during a 0.5 s prestimulus period,
and the result was multiplied by 2 to obtain the activity of individual sensory neurons housed
in the capitate peg sensillum as a spikes/s measurement.
To investigate the physiological activity of the A, B and C neurons housed in single capitate
peg sensilla, CO_2, *R*-(–)-1-octen-3-ol and (+)-borneol were used: gas cylinders containing
metered amounts of CO_2 (300, 600, 1200, 2400, or 4800 ppm) and oxygen (20%), balanced
by nitrogen (Strandmöllen AB, Ljungby, Sweden) were used to assess the activity of the A
neuron; serial decadic dilutions of *R*-(-)-1-octen-3-ol (CAS: 3687-48-7, Penta Manufacturing,
Livingston, USA), diluted in paraffin oil, were used to assess the activity of the B neuron;
and serial decadic dilutions of (+)-borneol (CAS: 464-43-7, Sigma-Aldrich, St. Louis, MO,
USA), diluted in diethyl ether (SupraSolv, Billerica, MA, USA) were used to assess the
activity of the C neuron. Pasteur pipettes were filled with the metered CO_2 and used
immediately to stimulate the preparation. A 15 μL aliquot of each dilution of *R*-(–)-1-octen-3-

ol and (+)-borneol was pipetted onto a filter paper (5 mm × 15 mm) inserted inside a Pasteur
pipette, and the diethyl ether was allowed 15 min to evaporate, before being used for stimulus
delivery. All stimuli were delivered into the airstream passing over the maxillary palp
preparation.
454
Calcium imaging in the *Ae. aegypti* antennal lobe (AL)
Odor-evoked responses in the *Ae. aegypti* antennal lobe (AL) were imaged using the brp-
QF2>QUAS-GCaMP6s progeny from the brp-QF2 and QUAS-GCaMP6s parental lines [24]. A
total of eight 6-8 day-old female mosquitoes were used for all calcium experiments. Each
mosquito was cooled on ice and transferred to a Peltier-cooled holder that allowed the
mosquito head to be fixed to a custom stage using ultraviolet glue. The stage permits the
superfusion of saline to the head capsule and space for wing and proboscis movement [25,54].
Once the mosquito was fixed to the stage, a window in its head was cut to expose the brain,
and the brain was continuously superfused with physiological saline [55]. Calcium-evoked
responses in the AL were imaged using the Prairie Ultima IV two-photon excitation
microscope (Prairie Technologies) and Ti-Sapphire laser (Chameleon Ultra; Coherent; at
1910 mW power). Experiments were performed at 75 µm depth from the ventral surface of
the AL, allowing characterization of the mediodorsal glomerular responses to olfactory
stimuli and allowing these glomeruli to be repeatedly imaged across preparations. To record
odor-evoked responses, images were collected from a 110 µm × 83 µm plane at 2 Hz (line
period of 1 ms), and for each odor stimulus, images were acquired for 25 s, starting 10 s
before the stimulus onset. Image data were imported into Matlab (v2017; Mathworks, Natick,
Massachusetts) for Gaussian filtering (2×2 pixel; σ = 1.5-3) and alignment using a single
frame as the reference at a given imaging depth and subsequently registered to every frame to
within ¼ pixel. Odor stimuli were diluted to 10^{-3} and 10^{-4} concentration in hexane (>99.5%
purity; Sigma), with hexane used as the solvent control. During an experiment, odor stimuli
were separated by intervals of 120 s to avoid receptor adaptation, and odor syringes were
used once per preparation to prevent decreased concentration within the cartridge. Calcium
responses are calculated as the change in fluorescence and time-stamped and synced with the
stimulus pulses. After an experiment, the AL was sequentially scanned at 0.5 µm depths from
the ventral to the dorsal surface to provide glomerular assignment and registration between
preparations. Glomeruli (1 µm^3 voxel) were mapped and registered based on the positions
and odor-evoked responses of the putative AL3, MD1-3, and AM2 glomeruli, using available
AL atlases [26,27] and the Amira software (v. 6.5, FEI Houston Inc.).
484
Behavioral assay
The role of borneol in human host-seeking female mosquitoes was examined with an arm-in-
a-cage assay (Extended Data Fig. 1a) described previously [47] with minor modifications
(Extended Data Fig. 1b and c) in an air-conditioned room (26±1 °C, 60±5 % RH). Briefly, the
experimenter's hand was presented to fifteen 5–10 days post emergence adult females. A
three-dimensional-printed interlocking ring with a diameter of 55 mm was used in this
experiment (Extended Data Fig. 1b). The ring was placed over the dorsal side of a nitrile
glove (powder-free latex). To evoke human odor and prevent mosquito bites, we replaced the
nitrile glove between the two ring components with a plastic net (Extemded Data Fig. 1c).
The interlocking ring included a central odorant delivery platform comprising a 10-mm-
diameter cover glass and two 5-mm-diameter filter discs (WHA10016508; Merck) for the
evaluation of VOCs (stl files are provided in Extended Data Fig. 2 and 3). Plastic net, nitrile
glove and odorant delivery platform were replaced between repetitions to avoid
contaminations. Mosquitoes were placed in a 20.3-cm^3 metal cage located in an experimental
room with a vent, under a video camera (EOS 70D, lens: MACRO 0.25/0.8ft; Canon Inc.,

Tokyo, Japan) and a light ring. Mosquitoes were allowed to acclimate for ten minutes before
recording. Mosquito behavior was recorded for 10 min at 25 frames per second. The number
of mosquito detection on the screen and ring were automatically counted using a custom
YOLOv8 model. We used the solvent diethyl ether (DEE) as a vehicle. On a blank filter disc,
25 μL of DEE, and DEE with racemic borneol (1M, 3.58 mg) were deposited on the filter
paper and allowed to evaporate for 2 min prior to mosquito exposure outside the experimental
room to avoid contamination. To enhance attractiveness, at the beginning of each experiment,
the experimenter first rubbed the ring-mounted glove against the shirt and skin for 1 minute.
Additionally, the experimenter blew twice, once into the cage and once into the glove. All
experiments were conducted during the first 4 hours of the diurnal period and lasted 10 min.
This schedule was chosen for practical reasons and because mosquitoes consistently
exhibited attraction to the human hand. Mosquito detection was normalized to the mean
detection of the control $(\frac{Sum\ of\ trajectories\ per\ minute}{Mean\ of\ summed\ trajectories\ for\ all\ minutes\ (control)}) * 100$. Statistical
comparisons of mosquito detections per minute were carried out with Mann Whitney U test
(P-value < 0.001, $n = 3$). For data variation across time and repetitions, see Figure 6. Data
was analyzed using R Studio [56].
516
YOLOv8 model
In this study, we fine-tuned the pre-trained YOLOv8m model [57] using a customized dataset
comprising images from the bioassay, along with additional data from previously published
literature [58], to investigate the behavioral role of borneol on mosquitoes. The training process
involved 250 epochs with a batch size of 10, utilizing separate sets of images and
corresponding annotations for training, validation, and testing (5,334, 1,498, and 1,479
images, respectively). To assess the model's performance, we employed the mean Average
Precision (mAP) metric across a range of intersection-over-union (IoU) thresholds from 50%
to 95% (mAP50-95). The resulting mAP50-95 value of 0.755 indicates the average precision
considering different IoU thresholds. Furthermore, the recall value of 0.978 indicates a high
proportion of true positive detections in this setup, while the precision value of 0.978 reflects
the accurate identification of mosquitoes. It's worth noting that detections were collected
with a minimum confidence threshold and IoU of 0.5 to ensure reliable results. Video
samples with automated detection are available in Extended Data Fig. 6. The model output
was proccesed with Python, resulting dataframe with the following columns, including frame,
x, y, treatment and trajectory identification. Statistical comparisons of all parameters were
carried out with Mann Whitney U test with Benjamini-Hochberg (BH) adjustment method.
Data was analyzed using R Studio (Team, 2021). The behavioral data can be found in
(Extended Data Fig. 4). All codes can be obtained by request to the corresponding authors.
536
537   **Figure legends**
538
Figure 1. Odorant receptor 49 is activated by bicyclic monoterpenoids.
a, Borneol and camphor oils from the camphor tree *Cinnamomum camphora* originated from
the island of Borneo and were traded with China and the rest of the Western world through
the maritime silk road during the classic age due to their medicinal and repellent properties.
The world map is a modified illustration Vecteezy.com (free license) where blue dots indicate
major trading ports. **b**, The capitate peg sensillum (cp) on the 4th segment of the mosquito
maxillary palps houses three neurons, including the CO_2 capitate-peg neuron (cpA)
expressing three gustatory receptors, the 1-octen-3-ol-sensitive neuron (cpB), and the orphan
neuron (cpC). **c,** The response profile of *Aedes albopictus* OR49 (AalbOR49) to odorant
mixtures (100 μM) belonging to a variety of chemical classes highlights the activity of

ketones. **d**, Bicyclic monoterpenoids, including camphor and fenchone are the most
efficacious activators of AalbOR49 (labeled in green). Statistical differences were evaluated
by ANOVA followed by a Kruskal-Wallis multiple comparisons test. ns, non-significant, *p
< 0.0332, **p < 0.0021, ***p < 0.0002 and ****p < 0.0001. Data indicate the means $\pm$ SEM.
553
Figure 2. Odorant receptor 49 is a borneol receptor.
a, Pineapple Haze (PH), OG Kush (OGK), Adom9 (A9), and Master Kush (MK) elicited the
highest currents from *Toxorhynchites amboinensis* OR49 (TambOR49). Borneol content is
indicated on the right Y-axis. A correlation between the OR49 response and the borneol
content in each Cannabis EO was observed ($R^2 = 0.89$). **b**, Response of TambOR49 to 6
Cannabis sub-mixtures and the constituents from sub-mixture 2. **c**, Concentration-response
relationships of TambOR49 and AaegOR49 to the two borneol enantiomers. **d**,
Concentration-response relationships of TambOR49, *Culex quinquefasciatus* OR38
(CquiOR38), *Ae. aegypti* OR49 (AaegOR49) and AalbOR49 in response to (+)-borneol and
(+)-camphor. Effective concentrations at 50% of the maximal response (EC_{50}) are circled. **e**)
Pairwise comparisons of the current responses of all four mosquito species OR49s elicited by
increasing concentrations of (+)-borneol and (+)-camphor. Statistical differences were
evaluated by multiple t-tests. *p < 0.0332, **p < 0.0021, ***p < 0.0002 and ****p < 0.0001.
Data indicate the means $\pm$ SEM. Representative current traces can be found in Supplementary
Fig. 3.
569
Figure 3. The maxillary palp of culicine mosquitoes respond to borneol and camphor.
a, Electropalpogram (EPG) responses in *Aedes albopictus*, *Aedes aegypti*, *Culex pipiens*, and
Anopheles gambiae. EPG responses to 1-octen-3-ol (blue, 10 μg), camphor (purple, 10 μg),
and borneol (green, 10 μg) enantiomers relative to hexane. Representative traces for each
odorant are shown below the x-axis. Statistical differences were evaluated via one-way
anova. *p < 0.05, **p < 0.01, ***p < 0.005 and ****p < 0.001. Data indicate the means $\pm$
SEM, $n = 15$. **b**, Dose-response relationships of the maxillary palps of three culicine
mosquito species in response to increasing concentrations of (+)-borneol and (+)-camphor.
Representative traces are shown on the right. Statistical differences were evaluated by
multiple t-tests. Data indicate the means $\pm$ SEM, $n = 15$.
580
Figure 4. Species-dependent response to (+)-borneol of the C neuron of the capitate peg
sensilla and its reliance of *Or49* in *Ae. aegypti*.
a, A tungsten electrode was used to record action potentials (spikes) from the capitate peg
sensillum on the maxillary palp, which houses three neurons. The cpA neuron responds to
CO_2 (large spike in traces), the cpB neuron responds to 1-octen-3-ol (medium size spikes)
and the cpC neuron responds to borneol (small size spikes). **b-e**, The spontaneous activity of
the three sensory neurons in the capitate peg sensilla of *Ae. aegypti* and *Anopheles gambiae*
s.s.. In the top traces, note the differences in spike amplitude of the A, B (orange markers)
and C neurons (green markers). *R*-(−)-1-octen-3-ol and (+)-borneol elicit responses in the B
and C neurons of *Ae. aegypti*, respectively. The response to (+)-borneol is abolished in
AaegOr49[−/−] mutant mosquitoes. The C neuron of *An. gambiae s.s.* does not respond to (+)-
borneol. **b-e**, Representative traces of the cp sensilla of *Ae. albopictus* and *Cx.*
quinquefasciatus. **f**, Species- and dose-dependent response of the C neuron to (+)-borneol
with mean $\pm$ s.e.m. at each concentration. The arrow and dotted line indicate the borneol dose
to the shown representative traces. *Ae. aegypti* ($n = 10$), *Ae. albopictus* ($n = 7$), *Cx.*
quinquefasciatus ($n = 2$), *Ae. aegypti Or49-/-* ($n = 8$), *An. gambiae* ($n = 15$).
597
Figure 5. Borneol elicits robust responses in the *Ae. aegypti* AL.

a, Schematic of the two-photon setup used to record calcium dynamics in the mosquito
antennal lobe (AL). **b**, AL atlas, highlighting the MD1 (blue), MD2 (orange), and MD3
(green) glomeruli. Non-responsive AL glomeruli (grey) and the mediodorsal glomeruli were
registered and mapped to previously published atlases. **c**, Pseudo color plot from a single
preparation of $\Delta F/F_0$ calcium responses (0-0.8 scale) to (+)-borneol (10^{-4} dilution), at a depth
of 75 μm from the surface of the AL. Borneol evoked a strong response in the region of
interest mapped to the MD3 glomerulus (highlighted in white). **d**, Glomerular responses
($\Delta F/F_0$) to (+)-borneol for the MD1, MD2, and MD3 glomeruli. Lines are the mean of one
glomerulus ($n = 4$-8 preparations); shaded areas are the SEM. The grey bar denotes stimulus
duration (2 s). **e**, Same as in D, except the glomeruli were stimulated with 1-octen-3-ol (10^{-4}
dilution). **f**, Same as in D, except the glomeruli were stimulated with CO_2 (5%). **g**, Tuning
curve for the MD3 glomerulus to a limited panel of 8 odorants, each tested at 10^{-4}
concentration. The MD3 glomerulus (bars in green) showed significant calcium responses to
enantiomers of borneol and camphor compared to the solvent control (Kruskal-Wallis test: χ^2
= 37.1, P < 0.0001; posthoc multiple comparisons: p < 0.05). Bars represent the mean ±
SEM.
615
Figure 6. Borneol inhibits host-seeking female *Ae. aegypti* (Liverpool) behavior.
a, The average number of females trajectoreis detection per minute (normalized to the control
group) in the ROI in response to control (DEE) and treatment (DEE + racemic borneol). **b**,
Schematics of all trajectories across treatment and repetitions (n=3). **c**, Average trajectories
over time (min). **d**, Sum of trajectory durations (seconds) within the ROI over time (minutes).
e, Total distance traveled per minute in the ROI. All comparisons were conducted with 30
data points for each treatment and tested with a Mann–Whitney U test (p-value < 0.001, $n = $
3). **f**, Schematics of all detections as a function of time.
624
Supplementary information
Supplementary Figure 1. The phylogenetically-conserved gene encoding OR49.
a, Amino-acid sequence alignment of OR49 homologs in *Culex quinquefasciatus* (Cqui),
Aedes albopictus (Aalb), *Aedes aegypti* (Aaeg), *Toxorhynchites amboinensis* (Tamb), and
Anopheles gambiae (Agam). The top chart presents the amino-acid consensus. High
consensus marked in green to yellow, low consensus are in red. **b**, Amino-acid sequence
identity matrix. Intensity of shading indicates percentage of homologies. **c**, Phylogenetic tree
of mosquito OR49 proteins (green). Closest brachyceran homologs (red branches) include
Glossina fuscipes (Gfuc) and *Drosophila melanogaster* (Dmel). Bootstrap support above
50% are shown. Gene exon structure of OR49 genes and other phylogenetically-related
genes. **d,** Exon compoistion of the OR49 group and related homologs. Exons are labeled in
white and black. The nucleotide positions of exon-exon boundaries are shown and intron
phases are color-coded. **e**, Chromosomic locations of Or49 homolog genes (green) and
indolergic ORs (indolORs, in red). **f**, Syntenic relationships of OR49 genes in mosquito
species and gene structure of OR49 genes.
640
641 **Supplementary Figure 2. List of volatile organic compounds present in *Cannabis***
642 **essential oils.**
643 Each Cannabis blend represented by a distinct color is composed of VOCs belonging mainly
644 to different terpene categories but also to non-terpene chemical classes.
645
Supplementary Figure 3. Representative current traces of oocytes expressing mosquito
OR49 proteins.

a, Representative traces of TambOR49+TambOrco injected oocytes vs monoterpenoid blends
and compounds. Pineapple Haze (PH), Adom #9 (A9), O.G Kush (OGK), Master Kush (MK)
and Jack Herrer (JH) triggered a measurable response. (±)-camphor, (+)-camphor, (+)-
fenchone and eucalyptol were used as positive controls at concentration of 0.1 mM, all
Cannabis essential oils were diluted by a factor of 1.5×10^5. **b**, Representative trace of
TambOR49+TambOrco vs sub-mixtures No. 1 through 6, which were formulated based on
overlapping active compounds from *Cannabis* essential oils. All sub-mixtures were diluted
by a factor of 1.5×10^6. Only sub-mixture No.2 activated TambOR49+TambOrco. (+)-
Camphor was used as positive control at concentration of 0.1 mM. **c**, Representative trace of
TambOR49+TambOrco vs the components of sub-mixture No. 2. (±)-Borneol elicited
currents 8-fold larger than the other tested single monoterpenoids. All compounds were tested
at concentration of 0.1 mM. **d**, Representative traces of TambOR49+TambOrco
concentration response curves (CRCs). Top left TambOR49+TambOrco vs (-)-borneol, Top
right TambOR49+TambOrco vs (+)-borneol and bottom left TambOR49+TambOrco vs (+)-
camphor. **e**, Representative traces of AaegOR49 CRC's. Left AaegOR49 vs (–)-Borneol;
right, AaegOR49 vs (+)-Borneol. **f**, Representative traces of CquiOR38+CquiOrco CRCs in
response to increasing concentrations of (+)-camphor (left) and (+)-borneol (right). **g**,
Representative traces of AaegOR49+Orco CRCs in response to increasing concentrations of
(+)-camphor (left) and (+)-borneol (right). **h**, Representative traces of AalbOR49+AalbOrco
CRCs in response to increasing concentrations of (+)-camphor (left) and (+)-borneol (right).
i) Representative traces of pairwise current comparisons between (+)-camphor and (+)-
borneol. From top to bottom, TambOR49+TambOrco, CquiOR38+CquiOrco,
AaegOR49+AaegOrco and AalbOR49+AalbOrco. Down arrows indicate odorant
administrations and concentrations. All concentrations are in micromolar [µM].
672
673   **Supplementary Figure 4. VOC content in *Cannabis* essential oils.**
674
Supplementary Figure 5. *Anopheles gambiae* OR28 (AgamOR28) does not respond to
borneol.
a, Concentration-response of AgamOR28 in response to increasing concentrations of four
plant volatile organic compounds, including aromatic (acetophenone and α-terpineol) and
terpenoid compounds (α-pinene and (+)-borneol). EC_{50} values shown in the inset are in the
low millimolar range. **b**, Representative current traces of AgamOR28-Orco activation by 4
plant volatile organic compounds. Arrowheads above the traces indicate the onset of the
odorant stimulus.
683
Supplementary Figure 6. Construction of *Ae. aegypti Or49* knockout line.
a, Diagram depicting the *Ae. aegypti Or49* gene and the gRNA target sites on the top and the
insertion cassette on the bottom flanked by the upstream (Left) and downstream (Right)
homology arms. The insertion cassette encompasses the *QF2* sequence and the 3' UTR of the
HSP70 gene along with the *ECFP* gene under the control of the *Opie2* promoter and the 3'
UTR of the *SV40* gene. Arrows indicate primer binding sites. Primers P1-12 are listed
in Table S4. **b**, Plasmid V1117F-Or49 map. **c**, *In vitro* Cas9 cleavage assay. Guide RNAs left
and right targeting the *Or49* gene almost completely digested PCR fragment containing target
sequence. (–) negative controls without guide RNAs show no PCR fragment digestion. **d**,
Sanger sequencing of amplified PCR fragments of G1 transgenic male individuals unveiling
the sequences overlapping upstream (left) and downstream (right) of the DNA cassette
inserted by homology directed repair. **e**, Sequencing of G1 individuals showing the complete
insertion of the DNA cassette containing *QF2* and *ECFP*. The presence of a single point
mutation was confirmed as an artifact and excluded by Sanger sequencing. **f**, Diagnostic PCR

reactions showing the presence of double bands in single G1 heterozygous individuals and
single upper band in pools of 10 individuals of the *OR49*$^{-/-}$ homozygous line.
700
701 **Supplementary Figure 7. The lowest borneol detection threshold of the maxillary palp is**
702 **0.1 μg.**
703 Bar plot of the maxillary palp responses to vehicle and increasing doses of borneol. Raw data
704 are shown in the adjacent table. Statistical differences were evaluated via one-way anova.
705 ***p < 0.005 and ****p < 0.001. Data indicate the means ± SEM, *n* = 15-27.
706
707 **Table S1. List of blends and single compounds used in the pharmacological screen.**
708
709 **Table S2. List of *Or* genes.**
710
711 **Table S3. Raw pharmacological data.**
712
713 **Table S4. Primer list (*Or49* knockout).**
714
715 **Extended data figures and tables**
716 **Extended Data Fig. 1 Diagrams of the odor delivery system.**
717 The odor delivery system is composed of interlocking top and bottom rings attached to a
718 removable chemical holder (overview, top, bottom and side views are provided). Dimensions
719 are provided in millimeters.
720
721 **Extended Data Fig. 2 Hand rings.stl.**
722 File format for 3D-printing of the two complementary hand rings.
723
724 **Extended Data Fig. 3 Chemical holder.stl.**
725 File format for 3D-printing of the chemical holder component.
726
727 **Extended Data Fig. 4 Behavioral data.**
728 CSV file of collected mosquito visits on the ROI.
729
730 **Extended Data Fig. 5 Video recording example of the of the female *Ae. aegypti* antennal**
731 **lobe.**
732
733 **Extended Data Fig. 6 Video recording example of the arm-in-a-cage assay.**
734 Thirty-second-long recording example at the 4-minute mark of mosquito landing behavior in
735 the presence of the vehicle diethyl ether (DEE) and racemic borneol.
736

Acknowledgments
This study was supported by ISF 997/19 awarded to A.W., ISF 719/21 awarded to J.B., NIH
RO1AI175152 and NSF IOS-2242604 awarded to O.S.A., NIH R01AI148300 awarded to
J.A.R, O.S.A., and R.J.P. J.A.R was funded by the Bill and Melinda Gates Foundation (INV-
021766). Science and Technology Development Plan Project of Jilin Province, China
(20200402001NC) awarded to Y.W. This research was supported by the Ministry of Science
& Technology, Israel to P.A.P. (grant numbers 3-16795 and 3-17985). We thank Ziv Kassner
for consultation, processing, analysis guidance, and technical support in the creation of the
mosquito detection model. We are especially grateful to Nadav Eyal at Eybna Terpene Based
Technologies for their time and supply of *Cannabis* essential oils. We are grateful to Dr. Laor
Orshan (Ministry of Health, Israel) for providing a colony of *Culex pipiens*. Special thanks to

Dr. Osnat Malka for productive discussions. Drs. Pitts and Bohbot are co-corresponding
authors.
750
Contributions
RJP initiated the functional characterization of OR49. JB conceived the study. YV identified
borneol as a key OR49 ligand. YW conducted mosquito electropalpograms. RMH identified
camphor as an OR49 agonist. OSA and IVC-A engineered the *Ae. aegypti Or49* knockout
strain. MG and RI designed and conducted the single sensillum recordings. ESS conducted
the behavioral experiments, created the YOLOv8 model and analyzed the data. DP and PP
conducted gene annotations, phylogenetic analyses and described OR49 syntenic
relationships. DR completed the *Toxorhynchites amboinensis* OR genomic PCRs and
sequencing. EY provided genetic constructs and injectable mRNA. AW provided funding and
scientific oversight. JB wrote the manuscript with individual contributions from all the
authors.
762
763 **Corresponding authors**
764 Correspondence to Jonathan D. Bohbot or Ronald J. Pitts.
765
766 **Ethics declarations**
767 Competing interests
768 The authors declare no competing interests.
769

References

1. Wadley, L. *et al.* Middle Stone Age Bedding Construction and Settlement Patterns at
Sibudu, South Africa. *Science* **334**, 1388–1391 (2011).

2. Pålsson, K. & Jaenson, T. G. T. Plant products used as mosquito repellents in Guinea
Bissau, West Africa. *Acta Trop* **72**, 39–52 (1999).

3. Maia, M. F. & Moore, S. J. Plant-based insect repellents: a review of their efficacy,
development and testing. *Malaria journal* **10 Suppl 1**, S11 (2011).

4. Moore, S. J. & Debboun, M. History of Insect Repellents. in *Insect Repellents: Principles,*
Methods, and Uses (eds. Debboun, M., Frances, S. P. & Strickman, D.) vol. 1 (CRC Press,
2007).

5. Kongkaew, C., Sakunrag, I., Chaiyakunapruk, N. & Tawatsin, A. Effectiveness of
citronella preparations in preventing mosquito bites: systematic review of controlled
laboratory experimental studies. *Trop. Med. Int. Heal.* **16**, 802–810 (2011).

6. Zhu, J. J. *et al.* Better than DEET Repellent Compounds Derived from Coconut Oil. *Sci.*
Rep. **8**, 14053 (2018).

7. Liu, F. *et al.* A dual-target molecular mechanism of pyrethrum repellency against
mosquitoes. *Nat Commun* **12**, 2553 (2021).

8. Iovinella, I., Pelosi, P. & Conti, B. A rationale to design longer lasting mosquito repellents.
Parasitology research **113**, 1813–1820 (2014).

9. Poudel, D. K. *et al.* The Chemical Profiling of Essential Oils from Different Tissues of
Cinnamomum camphora L. and Their Antimicrobial Activities. *Molecules* **26**, 5132 (2021).

10. Mei, Y. *et al.* The history, stereochemistry, ethnopharmacology and quality assessment of
borneol. *J Ethnopharmacol* **300**, 115697 (2023).

11. Erdelyan, C., Mahood, T., Bader, T. & Whyard, S. Functional validation of the carbon
dioxide receptor genes in Aedes aegypti mosquitoes using RNA interference. *Insect*
molecular biology **21**, 119–127 (2012).

12. McMeniman, C. J., Corfas, R. A., Matthews, B. J., Ritchie, S. A. & Vosshall, L. B.
Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction
to humans. *Cell* **156**, 1060–1071 (2014).

13. Lu, T. *et al.* Odor coding in the maxillary palp of the malaria vector mosquito Anopheles
gambiae. *Current biology* **17**, 1533–1544 (2007).

14. Grant, A. J. & Dickens, J. C. Functional characterization of the octenol receptor neuron
on the maxillary palps of the yellow fever mosquito, Aedes aegypti. *PLoS ONE* **6**, e21785
(2011).

15. Grant, A. J. & O'Connell, R. J. Electrophysiological responses from receptor neurons in
mosquito maxillary palp sensilla. in (ed. Block, E.) vol. 200 233-48-discussion 248-53-281–4
(John Wiley & Sons, Ltd., 1996).

16. Takken, W. & Kline, D. L. Carbon dioxide and 1-octen-3-ol as mosquito attractants.
Journal of the American Mosquito Control Association **5**, 311–316 (1989).

17. Gillies, M. T. The role of carbon dioxide in host-finding by mosquitoes (Diptera:
Culicidae): a review. *Bull Entomol Res* **70**, 525–532 (1980).

18. Wang, G., Carey, A. F., Carlson, J. R. & Zwiebel, L. J. Molecular basis of odor coding in
the malaria vector mosquito Anopheles gambiae. *PNAS* **107**, (2010).

19. Bohbot, J. *et al.* Molecular characterization of the Aedes aegypti odorant receptor gene
family. *Insect molecular biology* **16**, 525–537 (2007).

20. Matthews, B. J. *et al.* Improved reference genome of Aedes aegypti informs arbovirus
vector control. *Nature* **563**, 501–507 (2018).

21. Bohbot, J. D. & Dickens, J. C. Characterization of an enantioselective odorant receptor in
the yellow fever mosquito Aedes aegypti. *PLoS ONE* **4**, e7032 (2009).

22. Majeed, S., Hill, S. R., Dekker, T. & Ignell, R. Detection and perception of generic host
volatiles by mosquitoes: responses to CO2 constrains host-seeking behaviour. *Royal Society*
open science **4**, 170189 (2017).

23. Majeed, S., Hill, S. R., Birgersson, G. & Ignell, R. Detection and perception of generic
host volatiles by mosquitoes modulate host preference: context dependence of (R)-1-octen-3-
ol. *Royal Society open science* **3**, 160467 (2016).

24. Zhao, Z. *et al.* Mosquito brains encode unique features of human odour to drive host
seeking. *Nature* 1–7 (2022) doi:10.1038/s41586-022-04675-4.

25. Lahondère, C. *et al.* The olfactory basis of orchid pollination by mosquitoes. *Proceedings*
of the National Academy of Sciences USA (2019) doi:10.1073/pnas.1910589117.

26. Shankar, S. & McMeniman, C. J. An updated antennal lobe atlas for the yellow fever
mosquito Aedes aegypti. *Plos Neglect Trop D* **14**, e0008729 (2020).

27. Wolff, G. H., Lahondère, C., Vinauger, C., Rylance, E. & Riffell, J. A. Neuromodulation
and differential learning across mosquito species. *Proc. R. Soc. B* **290**, 20222118 (2023).

28. Anton, S., Loon, J. van, Meijerink, J. & Smid, H. Central projections of olfactory receptor
neurons from single antennal and palpal sensilla in mosquitoes. *Arthropod Structure &*
Development **32**, 319–327 (2003).

29. Jocher, G. *et al.* ultralytics/yolov5: v7.0 - YOLOv5 SOTA Realtime Instance
Segmentation. (2022) doi:10.5281/zenodo.7347926.

30. Cloyd, R. A., Marley, K. A., Larson, R. A., Dickinson, A. & Arieli, B. Repellency of
Naturally Occurring Volatile Alcohols to Fungus Gnat Bradysia sp. nr. coprophila (Diptera:
Sciaridae) Adults Under Laboratory Conditions. *J Econ Entomol* **104**, 1633–1639 (2011).

31. Liang, Y. *et al.* Evaluation of Repellency of Some Chinese Medicinal Herbs Essential
Oils Against Liposcelis bostrychophila (Psocoptera: Liposcelidae) and Tribolium castaneum
(Coleoptera: Tenebrionidae). *J Econ Entomol* **106**, 513–519 (2013).

32. Zhang, N. *et al.* Insecticidal, Fumigant, and Repellent Activities of Sweet Wormwood Oil
and Its Individual Components Against Red Imported Fire Ant Workers (Hymenoptera:
Formicidae). *J Insect Sci* **14**, 241 (2014).

33. Wood, M. J., Bull, J. C., Kanagachandran, K. & Butt, T. M. Development and laboratory
validation of a plant-derived repellent blend, effective against Aedes aegypti [Diptera:
Culicidae], Anopheles gambiae [Diptera: Culicidae] and Culex quinquefasciatus [Diptera:
Culicidae]. *PLOS ONE* **19**, e0299144 (2024).

34. Hill, C. A. *et al.* G protein-coupled receptors in Anopheles gambiae. *Science (New York,*
NY) **298**, 176–178 (2002).

35. Carey, A. F., Wang, G., Su, C.-Y., Zwiebel, L. J. & Carlson, J. R. Odorant reception in
the malaria mosquito Anopheles gambiae. *Nature* **464**, 66–71 (2010).

36. Bohbot, J. D. & Pitts, R. J. The narrowing olfactory landscape of insect odorant receptors.
Frontiers in Ecology and Evolution **3**, 39 (2015).

37. Hwang, Y.-S., Wu, K.-H., Kumamoto, J., Axelrod, H. & Mulla, M. S. Isolation and
identification of mosquito repellents in Artemisia vulgaris. *J Chem Ecol* **11**, 1297–1306
(1985).

38. Pitarokili, D., Michaelakis, A., Koliopoulos, G., Giatropoulos, A. & Tzakou, O. Chemical
composition, larvicidal evaluation, and adult repellency of endemic Greek Thymus essential
oils against the mosquito vector of West Nile virus. *Parasitol Res* **109**, 425–430 (2011).

39. Waliwitiya, R., Kennedy, C. J. & Lowenberger, C. A. Larvicidal and
oviposition☐altering activity of monoterpenoids, trans☐anithole and rosemary oil to the
yellow fever mosquito Aedes aegypti (Diptera: Culicidae). *Pest Manag Sci* **65**, 241–248
(2009).

40. Picollo, M. I., Toloza, A. C., Cueto, G. M., Zygadlo, J. & Zerba, E. Anticholinesterase
and pediculicidal activities of monoterpenoids. *Fitoterapia* **79**, 271–278 (2008).

41. Abdelgaleil, S. A. M., Mohamed, M. I. E., Badawy, M. E. I. & El-arami, S. A. A.
Fumigant and Contact Toxicities of Monoterpenes to Sitophilus oryzae (L.) and Tribolium
castaneum (Herbst) and their Inhibitory Effects on Acetylcholinesterase Activity. *J. Chem.*
Ecol. **35**, 518–525 (2009).

42. Rajkumar, S. & Jebanesan, A. Chemical composition and larvicidal activity of leaf
essential oil from Clausena dentata (Willd) M. Roam. (Rutaceae) against the chikungunya
vector, Aedes aegypti Linn. (Diptera: Culicidae). *J Asia-pac Entomol* **13**, 107–109 (2010).

43. Afify, A. & Galizia, C. G. Gravid females of the mosquito Aedes aegypti avoid
oviposition on m-cresol in the presence of the deterrent isomer p-cresol. *Parasite Vector* **7**,
315 (2014).

44. Raj, G. A., Chandrasekaran, M., Krishnamoorthy, S., Jayaraman, M. & Venkatesalu, V.
Phytochemical profile and larvicidal properties of seed essential oil from Nigella sativa L.
(Ranunculaceae), against Aedes aegypti, Anopheles stephensi, and Culex quinquefasciatus
(Diptera: Culicidae). *Parasitol Res* **114**, 3385–3391 (2015).

45. Nunes, R. K. V. *et al.* Evaluation of (–)-borneol derivatives against the Zika vector,
Aedes aegypti and a non-target species, Artemia sp. *Environ Sci Pollut R* **25**, 31165–31174
(2018).

46. Ng, R., Wu, S. & Su, C. Neuronal Compartmentalization: A Means to Integrate Sensory
Input at the Earliest Stage of Information Processing? *Bioessays* **42**, e2000026 (2020).

47. Dekel, A., Sar-Shalom, E., Vainer, Y., Yakir, E. & Bohbot, J. D. The oviposition cue
indole inhibits animal host attraction in Aedes aegypti (Diptera: Culicidae) mosquitoes.
Parasite Vector **15**, 422 (2022).

48. Zhou, X., Rinker, D. C., Pitts, R. J., Rokas, A. & Zwiebel, L. J. Divergent and conserved
elements comprise the chemoreceptive repertoire of the non-blood feeding mosquito
Toxorhynchites amboinensis. *Genome biology and evolution* **6**, 2883–2896 (2014).

49. Dekel, A., Pitts, R. J., Yakir, E. & Bohbot, J. D. Evolutionarily conserved odorant
receptor function questions ecological context of octenol role in mosquitoes. *Scientific*
reports **6**, 37330 (2016).

50. Huff, R. M. & Pitts, R. J. Carboxylic acid responses by a conserved odorant receptor in
culicine vector mosquitoes. *Insect Mol Biol* **29**, 523–530 (2020).

51. Wang, Y. *et al.* Molecular basis of peripheral olfactory sensing during oviposition in the
behavior of the parasitic wasp Anastatus japonicus. *Insect Biochem. Mol. Biol.* **89**, 58–70
(2017).

52. Herre, M. *et al.* Non-canonical odor coding in the mosquito. *Cell* **185**, 3104-3123.e28
(2022).

53. Ghaninia, M., Majeed, S., Dekker, T., Hill, S. R. & Ignell, R. Hold your breath –
Differential behavioral and sensory acuity of mosquitoes to acetone and carbon dioxide. *Plos*
One **14**, e0226815 (2019).

54. Melo, N. *et al.* Geosmin attracts Aedes aegypti mosquitoes to oviposition sites. *Current*
Biology **30**, 1–8 (2020).

55. Vinauger, C. *et al.* Visual-Olfactory Integration in the Human Disease Vector Mosquito
Aedes aegypti. *Current biology* **29**, 2509-2516.e5 (2019).

56. Team, R. C. *R: A Language and Environment for Statistical Computing.* (R Foundation
for Statistical Computing, Vienna, Austria, 2021).

57. Jocher, G., Chaurasia, A. & Qiu, J. YOLO by Ultralytics (Version 8.0. 0)[Computer
software]. (2023).

58. Janson, K. D. *et al.* Development of an automated biomaterial platform to study mosquito
feeding behavior. *Frontiers Bioeng Biotechnology* **11**, 1103748 (2023).

Figure 1

Figure 2

Figure 3

Figure 4

a

cp
sensillum

Tungsten electrode

Cuticle

Gr1,2,3-CO$_2$

Or8-1-octen-3-ol

Or49-borneol

b WT · *Ae. aegypti*

WT · Octenol

WT · Borneol

OR49-/- · Borneol

c *An. gambiae* (WT)

Borneol

d *Ae. albopictus* (WT)

Borneol

e *Cx. quinquefasciatus* (WT)

Borneol

f

- ● *Ae. aegypti* (WT)
- ■ *Ae. albopictus* (WT)
- ◆ *Cx. quinquefasciatus* (WT)
- ○ *Ae. aegypti (49-/-)*
- ■ *An. gambiae* (WT)

cpC, Spikes/sec

(+)-Borneol (%, W/W)

Figure 5

a 900 nm laser

Odor delivery

b MD3 MD2 MD1 20 μm

c $\Delta F/F_0$ 0.6 / 0.2 / 0.0

brp-QF2>QUAS-GCaMP6s

d Borneol MD3 MD2 MD1 5 sec

e 1-octen-3-ol

f CO_2

g CO_2 (+)-Borneol (+)-Camphor (−)-Borneol (−)-Camphor Geosmin 1-Octen-3-ol Control

Figure 6