*Title:* *The evolution of borneol repellency in culicine mosquitoes*

**Authors:** Yuri Vainer[1†], Yinliang Wang[1,5†], Robert M. Huff[2†], Majid Ghaninia[3], Iliano V.

Coutinho-Abreu[4], Evyatar Sar-Shalom[1], Carlos Ruiz[7], Dor Perets[1], Esther Yakir[1], Dhivya

Rajamanickam[2], Alon Warburg[6], Philippos Papathanos[1], Omar S. Akbari[4], Rickard Ignell[3], Jeff

A. Riffell[7], R. Jason Pitts[2§], Jonathan D. Bohbot[1§]

**Authors affiliations:**

1 Department of Entomology, The Hebrew University of Jerusalem, The Robert H. Smith

Faculty of Agriculture, Food and Environment, Rehovot, Israel

2 Department of Biology, Baylor University, Waco, Texas, USA

3 Unit of Chemical Ecology, Department of Plant Protection Biology, Swedish University of

Agricultural Sciences, Alnarp, Sweden

4 School of Biological Sciences, Department of Cell and Developmental Biology, University of

California, San Diego, La Jolla, CA, USA

5 Northeast Normal University, China

6 Department of Microbiology and Molecular Genetics, Institute for Medical Research Israel-

Canada, The Kuvin Centre for the Study of Infectious and Tropical Diseases, Faculty of

Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

7 Department of Biology, University of Washington, Seattle, WA, USA 98195

† Yuri Vainer, Yinliang Wang and Robert M. Huff contributed equally to this work.

*§ Co-corresponding authors*

**Funding:** ISF 997/19 awarded to A.W., ISF 719/21 awarded to J.B., NIH RO1AI175152 and

NSF IOS-2242604 awarded to O.S.A., NIH R01AI148300 awarded to J.A.R, O.S.A., and R.J.P.

J.A.R was funded by the Bill and Melinda Gates Foundation (INV-021766). Science and

Technology Development Plan Project of Jilin Province, China (20200402001NC) awarded to

Y.W. This research was supported by the Ministry of Science & Technology, Israel to P.A.P.

(grant numbers 3-16795 and 3-17985).

**Abstract**

Insects have developed remarkable adaptations to effectively interact with plant secondary metabolites and utilize them as cues to identify suitable hosts. Consequently, humans have used aromatic plants for centuries to repel mosquitoes. The repellent effects of plant volatile compounds are mediated through olfactory structures present in the antennae, and maxillary palps of mosquitoes. Mosquito maxillary palps contain capitate-peg sensilla, which house three olfactory sensory neurons, of which two are mainly tuned to either carbon dioxide or octenol - two animal host odorants. However, the third neuron, which expresses the OR49 receptor, has remained without a known ecologically-relevant odorant since its initial discovery. In this study, we used odorant mixtures and terpenoid-rich *Cannabis* essential oils to investigate the activation of OR49. Our results demonstrate that two monoterpenoids, borneol and camphor, selectively activate OR49, and OR9-expressing neurons, as well as the MD3 glomerulus in the antennal lobe. We confirm that borneol repels female mosquitoes, and knocking out the gene encoding the OR49 receptor suppresses the response of the corresponding olfactory sensory neuron. Importantly, this molecular mechanism of action is conserved across culicine mosquito species, underscoring its significance in their olfactory systems.

**Key words:** mosquitoes, borneol, monoterpenoid, repellent, odorant receptor, maxillary palp

## 1. Introduction

The use of plants to keep away insects has been a human practice since prehistorical times (Wadley et al., 2011) and is still a current practice in many parts of the world (Maia and Moore, 2011; Pålsson and Jaenson, 1999). This practice is shared with nonhuman primates that use

plants topically for their insect repelling properties (Baker, 1996). Plant-based essential oils

(EOs), including lemon eucalyptus leaf oil (Moore and Debboun, 2007), citronella oil

(Kongkaew et al., 2011), and coconut oil (Zhu et al., 2018) exhibit different degrees of repellent

activity due to the presence of pyrethrins (Liu et al., 2021), phenol derivatives, and terpenoids

(Iovinella et al., 2014). The latter chemical class includes oxygen-containing monoterpenes with

open-chain hydrocarbons, such as linalool, citronellol, geraniol, and cyclic derivatives such as

cineole, fenchone, camphor and borneol.

Borneol is believed to have derived its name from the island of Borneo, where it was initially

traded with China and then introduced to the West due to its therapeutic, refreshing, and

repellent effects against mosquitoes (Mei et al., 2023). Essential oils from various aromatic and

medicinal plants such as *Salvia* spp. (Amer and Mehlhorn, 2006; Conti et al., 2012; Dayaram

and Khan, 2016), *Valeriana officinalis* (Gordon et al., 2018), *Rosmarinus officinalis* (Alavez-

Rosas et al., 2022; Dayaram and Khan, 2016; Drapeau et al., 2009; Pratiwi and Purwati, 2021),

*Plectranthus* spp. (Kulkarni et al., 2013; Lukhoba et al., 2006), *Thymus* spp. (Pitarokili et al.,

2011) and *Coriandrum sativus* (Benelli et al., 2013; Dayaram and Khan, 2016) contain

significant levels of borneol that exhibit repellent and toxic activities against mosquito larvae

(Santos et al., 2010).

Our understanding of the molecular modes of repellent action remains fragmented. Catnip oil,

aside from its medical properties, exhibits repellency mediated by iridoids like nepetalactone,

which target the transient receptor potential cation channel TRPA1 (Melo et al., 2021) and

odorant receptors (Dickens and Bohbot, 2013) in insects. Similarly, the monoterpenoid

citronella, the primary compound found in citronella oil, has been shown to target both the

odorant receptor co-receptor (Orco) and the TRPA1 channel (Kwon et al., 2010).

Minimal information is available in the scientific literature regarding the molecular mode of

action underlying borneol detection by insects. However, previous research has demonstrated

that borneol, along with other monoterpenoids, induces Orco-mediated repellency in the

antennae of *Ae. aegypti* (Andreazza et al., 2023; Liu et al., 2021). Interestingly, an additional

mechanism for borneol-mediated repellency operates through a separate Orco-mediated pathway

in the antennae of *Ae. aegypti* (Andreazza et al., 2023; Liu et al., 2021).

The capitate-peg (cp) sensillum located on the maxillary palp comprises three olfactory

neurons, each distinguishable by size, olfactory receptor gene expression profile, and odor

response characteristics. In both culicine and anopheline mosquitoes, the largest olfactory

sensory neuron (cpA) expresses three gustatory receptors (*Grs*) that specifically detect $CO_2$

(Erdelyan and Mahood, 2011; McMeniman et al., 2014). The medium-sized cpB neuron of

*Anopheles gambiae* and *Aedes aegypti* expresses the 1-octen-3-ol odorant receptor *Or8* and its

co-receptor Orco (Grant and Dickens, 2011; Grant and O'Connell, 1996; Lu et al., 2007). Both

$CO_2$ and 1-octen-3-ol elicit attraction and signal the presence of animal hosts in anopheline

(Takken and Kline, 1989) and culicine (Gillies, 1980) mosquitoes. This functional organization

has remained remarkably conserved throughout over 180 million years of mosquito evolution

across these clades.

The third neuron (cpC) expresses a distinct *Or* gene in anopheline and culicine mosquitoes. In

anophelines, the cpC neuron expresses the *Or28* gene (Lu et al., 2007), which responds to

acetophenone and 2,4,5-trimethylthiazole, two plant volatile organic compounds (pVOC) (Wang

et al., 2010). Conversely, in culicines, the cpC neuron expresses the *Or49* gene (Bohbot et al.,

2007), which is unrelated to *AgamOr28*. Elucidating the function of OR49 within the capitate-

peg sensillum in mosquito palps will provide crucial insights into understanding the fundamental

aspects of cp neuron compartmentalization (Su et al., 2012).

Using a pharmacological approach, we expressed *Or49* in a heterologous expression system

and subjected it to odorant mixtures, plant EOs, and individual volatile odorant compounds. Our

findings provide strong evidence that the OR49 receptor is responsive to plant-derived bicyclic

monoterpenoids particularly showing a marked selectivity towards borneol—a well-known

repellent compound that also exhibits toxicity to *Aedes* mosquito larvae. Furthermore, we

observed that a loss-of-function mutation in the *Or49* gene leads to a lack of electrophysiological

responses when stimulated by borneol in the cpC neuron. Confirming our pharmacological and

electrophysiological results, behavioral studies reveal that borneol induces repellency in female

mosquitoes seeking animal hosts. It is likely that mosquitoes employ borneol as a signal to

distinguish between human and plant hosts. These findings highlight the pivotal role of OR49 in

shedding light on the function of cp neuron compartmentalization.

**Results**

*The conservation of the OR49 protein family spans across the evolutionary history of mosquitoes*

Using the AaegOR49 protein sequence as a query, we identified orthologs in *Ae. albopictus*, *Cx*

*quinquefasciatus, Tx. amboinensis* and *An. gambiae* (Figure 1A). Protein sequence alignment

shows high level of sequence coverage and a broad range of sequence identities that correlates

with mosquito phylogenetic relationships (Figure 1B). Our phylogenetic analysis supports

previous studies (Bohbot et al., 2007; Matthews et al., 2018) describing the OR49 family as a

Culicinae-specific group (Figure 1C). The culicine OR49, OR29, OR71 groups and the more

distantly related anopheline OR48/49 group form a monophyletic clade. These clades are further

supported by their exon structure and conserved intron phases (Figure 1D). The *Or49* genes are

located on chromosome 2 in both *Aedes* and *Anopheles* mosquitoes (Figure 1E). In *Aedes*

mosquitoes, *Or49* is located with *OR29* and *OR71* within several kb of each other and have

conserved syntenic relationships with neighbouring genes (Figure 1F). Except for the close

genomic proximity of OR49-related genes in *Cx. quinquefasciatus*, syntenic relationships in this

region are not shared with *Aedes* genomes. By contrast, *Anopheles* homologs (*Or48/49*) are

sitting next to each other, within a 5 kb distance of each other and without close neighboring *Or*

genes.

*OR49 is a selective borneol receptor*

To identify candidate OR49 ligands, we used the two-electrode voltage clamp of oocytes

expressing OR49 (Figure 2A, Table S1). We investigated the response of AalbOR49 to a panel

of odorants, which included 81 organic compounds representing a range of chemical classes

(Table S1). Oocytes expressing functional receptor complexes were strongly activated by blends

containing ketonic compounds (see in inset in Figure 2B). Individual testing of the single

compounds from the ketone blend revealed the efficacy of bicyclic monoterpenoids, with

camphor emerging as the most efficacious member followed by cineole (also called eucalyptol)

and fenchones (Figure 2B).

Subsequently, we screened OR49 from the nectar-feeding *Tx. amboinensis* (*TambOR49*) with

8 *Cannabis*-derived terpene mixtures (Figure 2C, Table S1, Figure S1). The three most active

mixtures were Pineapple Haze, OG Kush, Adom #9 followed by Master Kush with a much lower

efficacy. Six terpene sub-mixtures, each containing 3-4 terpenes present in all four active

*Cannabis* essential oils, were further tested using a single-blind approach. Sub-mixture No. 2

was the only pharmacologically-active treatment (Figure 2D). Single compounds from mixture

No. 2 were then prepared and individually tested in a single-blind approach as well. The racemic

borneol was significantly more active than fenchol, alpha-terpineol, geraniol and camphor

(Figure 2D). Subsequently, the borneol content in the original *Cannabis* mixtures was revealed,

establishing a correlation between the borneol content and the activity of the blend (Figure 2C,

Figure S2).

We established concentration-response curves (CRCs) to examine the relative sensitivity of

mosquito OR49s for the two enantiomers of borneol. TambOR49 and AaegOR49 did not exhibit

enantioselectivity towards the (+) and (–)-borneol (Figure 2E) as indicated by their $EC_{50}$ values

in the one-digit micromolar range. We established additional CRCs between TambOR49, *Cx.*

*quinquefasciatus* OR38 (CquiOR49), *Ae. aegypti* OR49 (AaegOR49), and *Ae. allbopictus* OR49

(AalbOR49), for the two most potent ligands, (+)-borneol and (+)-camphor (Figure 2F). (+)-

Borneol was 19-32 times more potent than (+)-camphor in all cases except for TambOR49 in

which both compounds were equally active in the low micromolar range. In all four examined

culicine species, including the nectar-feeding *Tx. amboinensis*, (+)-borneol consistently elicited

significantly higher responses than (+)-camphor (Figure 2G).

In *An. gambiae*, the cpC neuron expresses OR28 (Figure 2A), which is activated by panel of

diverse plant VOCs with acetophenone and 2,4,5-trimethylthiazole being the most effective

ligands in the *Xenopus* oocyte expression system (Lu et al., 2007; Wang et al., 2010). The $EC_{50}$

values of AgamOR28 for acetophenone, α-pinene, α-terpineol and (+)-borneol were in the low

millimolar range indicating that more potent ligands remain to be identified (Figure S4). Borneol

did not elicit any response at the tested concentrations suggesting that AgamOR28 is tuned to

VOCs belonging to different chemical classes.

*The maxillary palp of culicine mosquitoes responds to borneol and camphor*

To investigate the olfactory effect of camphor and borneol *in vivo*, we conducted electropalpogram (EPG) recordings on three culicine (*Ae. albopictus*, *Ae. aegypti*, *Cx. pipiens*) and one anopheline (*An. gambiae*) species. 1-Octen-3-ol, which served as a positive control, induced significant responses in all mosquitoes tested. Camphor and borneol elicited consistent palp responses in Culicinae mosquitoes, including *Ae. albopictus*, *Ae. aegypti*, and *Cx. pipiens* but not in *An. gambiae* (Figure 3). In *Ae. albopictus*, (–)-borneol was more active than (+)-camphor, while in *Ae. aegypti* both borneol enantiomers elicited significantly greater EPG responses than camphor enantiomers. In *Cx. pipiens*, borneol and camphor elicited comparable EPG responses.

We also established dose-response relationships with (+)-camphor and (+)-borneol in *Ae. albopictus*, *Ae. aegypti*, and *Cx. pipiens* (Figure 3B). Among the three culicine species, *Cx. pipiens* exhibited the largest responses in response to (+)-camphor. However, we did not find any significant statistical differences in activity between camphor and borneol in the tested species.

*The Or49 gene confers sensitivity to borneol*

To knock out the gene encoding the *Ae. aegypti* OR49 receptor, we used homology directed repair (HDR) to knock in a transgenic construct containing the QF2 transactivator and a ubiquitously expressing eCFP fluorescent marker driven by the OpiE2 promoter at the start codon of the *Or49* gene (Figure S5A,B). HDR was triggered by double strand breaks at the *Or49* promoter and the 1st exon mediated by two guide RNAs, whose cleavage activities had been tested *in vitro* (Figure S5C). Out of 739 injected embryos, 5.5% survived to the pupal stage.

These $G_0$ individuals were sex sorted and outcrossed with wildtypes. Sixteen $G_1$ larvae displayed

blue (cyan) fluorescent bodies (Figure S5D). Insertion of the transgenes within *Or49* was

confirmed in multiple $G_1$ individuals (Figure S5E,F) by sequencing of the insertion sites as well

as the transgene (Figure S5E,F) resulting in a nonfunctional receptor. An *Or49* $^{-/-}$ homozygous

mosquito strain was isolated by single-pair mating, screening the fluorescence marker in the

offspring for 100% penetrance, and PCR confirmation for the presence of homozygous mutant

alleles (Figure S5G).

To investigate the genetic mechanism determining borneol sensitivity in the palp, we

conducted single-sensillum recordings from the capitate-peg sensillum of *Ae. aegypti*, *Ae.*

*albopictus*, *Cx quinquefasciatus* and *An. gambiae* (Figure 4A, Figure S6). Single sensillum

recordings identified the spontaneous activity of three sensory neurons, distinguished by

differences in spike amplitudes, housed in the capitate peg sensilla of wild type *Ae. aegypti*

(Figure 4C) and *An. gambiae* (Figure 4D). The response of the cpA and cpB neurons to $CO_2$

(Majeed et al., 2017) (data not shown) and *R*-(–)-1-octen-3-ol (Figure 4C; Majeed et al., 2016),

respectively, has been well characterized in previous studies (Grant and Dickens, 2011; Grant

and O'Connell, 1996; Lu et al., 2007). (+)-Borneol elicited a dose-dependent response in the cpC

neuron of *Ae. aegypti*, *Ae. albopictus*, *Cx. quinquefasciatus* but not in *An. gambiae s.s.* (Figure

4B). No SSR responses were recorded from the cpC neuron in the *Ae. aegypti Or49* null mutant

line in response to (+)-borneol (Figure 4C).

*Two-photon imaging and borneol responses in the MD3 glomerulus*

To examine how *Ae. aegypti* processes borneol in the antennal lobe (AL), two-photon imaging

experiments were conducted using pan-neuronal GCaMP-expressing *Ae. aegypti* mosquitoes. We

utilized existing mosquito lines that contained a *QUAS-GCaMP6s* transgene crossed with the

*brp-QF2* driver line (Zhao et al., 2022), allowing the directed expression of the calcium indicator

GCaMP6s in all neurons of the AL. Mosquitoes were glued to holders that permitted two-photon

imaging of calcium responses in the AL (Figure 5A), allowing repeatable registration of the AL

glomeruli between preparations (Figure 5B) and recording of glomerular responses (Figure 5C)

(Lahondere et al., 2020). Glomeruli from our two-photon imaging results were mapped to AL

atlases (Shankar and McMeniman, 2021; Wolff et al., 2022), allowing accurate identification of

the glomeruli of interest in their spatial context (Additional file 7). We focused on the

mediodorsal glomeruli, as these glomeruli receive input from the capitate peg sensilla of

maxillary palps (Figure 5B,C) and are responsive to host odors, including 1-octen-3-ol (MD2)

and $CO_2$ (MD1) (Wolff et al., 2022).

To determine the odor coding of the mediodorsal glomeruli, and identify the cognate

glomerulus representing borneol, we first recorded from the MD1-3 glomeruli while stimulating

with $CO_2$ (5%), 1-octen-3-ol ($10^{-4}$ dilution), borneol ($10^{-4}$ dilution), and the solvent control. For

the MD3 glomerulus, (+)-borneol elicited strong, tonic responses that were significantly greater

than the solvent control (P<0.001) (Figure 5D). By contrast, responses of the MD1 and MD2

glomeruli to borneol were not significantly different from the control (P=0.09 and 0.23,

respectively). The MD2 glomerulus showed the greatest responses to 1-octen-3-ol (Figure 5E) (P

= 0.00003), and the MD1 glomerulus to $CO_2$ (P=0.01) (Figure 5F).  Given MD3's response to

(+)-borneol, and OR49 / cpC's responses to other terpene compounds, we next examined how

this glomerulus responded to a limited panel of different odorants, including enantiomers of

borneol and camphor. From this panel, (–)-borneol elicited the greatest response in the MD3

glomerulus (Figure 5G), closely followed by (+)-borneol and enantiomers of camphor.

*Borneol repels human-host seeking female Ae. aegypti mosquitoes*

To examine the impact of (+)-borneol on blood-seeking *Aedes aegypti* mosquitoes, we conducted

an arm-in-a-cage assay, exposing 15 females to human skin odor for 10 minutes (Figure 6). A

protective glove covered the hand, allowing mosquitoes to detect the odor through a dorsal open

area. This open area was protected with a screen and equipped with a chemical holder for the

deposition of (+)-borneol (1M). To monitor the number of females in the region of interest

(ROI), we trained a custom object detection YOLOv5 model (Jocher et al., 2022). Our results

showed a significant reduction (71.2%) in female landing on the ROI in the (±)-borneol-treated

hand compared to the vehicle-treated hand when presented separately (Figure 6A). The (+)-

borneol treatment consistently demonstrated lower detections, with less statistical variation

between repetitions than the vehicle treatment (Figure S7A). In both the control and borneol

treatments, we observed three waves of visits (Figure S7B). We used the trained YOLOv5 model

to monitor mosquitoes present in the ROI over time (Figure 6B) for all repetitions see Figure

S7C, and a video sample is available in Additional file 8. Figure S7B provides a visual

representation of the variation in total detections per minute, highlighting the detection

differences between the control and (+)-borneol treatments (total detections of 64,549 and

18,584, respectively) throughout the duration of the experiment. In the control treatment, the

highest number of detections was observed in the one-two minutes interval, accounting for

16.2% of the total detections. In the (+)-borneol treatment, the peak detections occurred during at

the 8-9 minute interval, comprising 14.4% of the total detections.

**4. Discussion**

Our results show that the culicine OR49 proteins are selective and sensitive bicyclic

monoterpenoid receptors with a marked selectivity for borneol. Furthermore, we show that the

deletion of the *Ae. aegypti Or49* gene suppresses cpC electrophysiological responses to borneol.

In all three blood-feeding culicine species, OR49s were more sensitive to borneol than to

camphor. Interestingly, the *Tx. amboinensis* OR49 exhibited overlapping sensitivity towards

borneol and camphor suggesting that camphor plays an increased role in this species. However,

camphor elicited significantly lower currents than borneol.

The OR49 clade is conserved in culicine mosquitoes. A complex process of gene duplication

events appears to have generated several related ORs, including OR37, OR38, OR71 and others.

Whether this OR subfamily expansion in culicines have specialized in detecting cyclic

monoterpenes will have to be examined. Antennal-expressed *AgamOr48* (Hill et al., 2002), one

of the two closest homologs to culicine *Or49*, responds to straight chains alcohols, ketones, and

acetates (Carey et al., 2010; Wang et al., 2010) suggesting that anopheline are tuned to a

different class of compounds of unknown ecological function.

In pharmacological assays, natural insect repellents interact with TRP channels (Melo et al.,

2021) and ORs (Bohbot et al., 2011; Bohbot and Dickens, 2010; Dekel et al., 2019; Xu et al.,

2019; Zeng et al., 2018) at the micromolar to low millimolar concentration range. By

comparison, our results show that OR49s exhibit sensitivities three log units below these

examples and is matched only by the *Drosophila melanogaster* OR56a-geosmin interaction

(Stensmyr et al., 2012).

Contrary to the well-characterized mosquito OR8, which discriminates between the two 1-

octen-3-ol enantiomers, the (+) and (–)-borneol elicited comparable activations at the

pharmacological, physiological and AL levels. These findings suggest that the receptor binding

pocket accommodates both forms of borneol eliciting the activation of the MD3 glomerulus in the brain. These findings align with previous studies demonstrating that the three MD glomeruli receive input from the maxillary palp cp neurons whereby the MD1 glomerulus receiving $CO_2$ input from the cpA neurons (Anton et al., 2003) while the medium size MD2 glomerulus receives 1-octen-3-ol input from the cpB neurons (Wolff et al., 2023).

Borneol and camphor evoked lower responses than 1-octen-3-ol in the culicine palps, which is consistent with the smaller size of the cpC neuron in comparison to the cpB neuron. Based on single sensillum recordings, these responses of the palp can be directly attributed to the *Or49* gene. The absence of OR28 activation and *An. gambiae* palp EPG responses by borneol and camphor support this conclusion.

Borneol and borneol-rich plants have long been known not only for their medicinal use (Mei et al., 2023) but also for their insecticidal properties against the granary weevil (Kordali et al., 2006) and fire ants (Zhang et al., 2014). This monoterpenoid has also larvicidal properties against a variety of mosquito species (Afify and Galizia, 2014; Nunes et al., 2018; Pitarokili et al., 2011; Raj et al., 2015; Rajkumar and Jebanesan, 2010; Waliwitiya et al., 2009). Borneol repels fungus gnats (Cloyd et al., 2011), the booklouse *Liposcelis bostrychophila* (Liang et al., 2013) and fire ants (Zhang et al., 2014). It also influences adult mosquito behavior, repelling human host-seeking *Ae. aegypti*, *Cx. quinquefasciatus* and *An. stepehensi* (Hwang et al., 1985; Pitarokili et al., 2011) while its presence in EOs stimulates oviposition in *Ae. aegypti* (Waliwitiya et al., 2009).

Is OR49 a molecular target of toxic plant-produced monoterpenoids, such as camphor and borneol, or do mosquito detect the volatile compounds in order to avoid their sources? Plants synthesize a variety of secondary metabolites to defend against herbivores and pathogens.

Bicyclic monoterpenoids, such as 1-8-cineole, fenchone, camphor and borneol, may act as

acetylcholinesterase inhibitors (Abdelgaleil et al., 2009; Picollo et al., 2008). Other classes of

terpenes have been proposed to target octopamine receptors (Enan, 2001) and the γ-aminobutyric

acid (GABA) type A receptor (Höld et al., 2000). Their overall effects include developmental

delays, oviposition and egg hatching reduction (Abdelgaleil et al., 2022). Intriguingly, gravid

females exhibit a contrasting behavior, laying more eggs in solutions containing borneol and

camphor (Waliwitiya et al., 2009), suggesting that these compounds have opposite effects on

host-seeking behavior and the oviposition preferences of gravid female mosquitoes.

Mosquitoes seem to avoid plants emitting toxic levels of bicyclic monoterpenoids, with

camphor and borneol ranking among the most potent compounds. This avoidance involves a

dedicated olfactory circuit, beginning with OR49, a receptor expressed by the palp cpC neuron

that projects to the MD3 glomerulus in the antennal lobe of *Ae. aegypti*. Whether borneol and

other bicyclic monoterpenoids serve as danger or attractive signals, the cpC neuron may

contribute to identify non-animal resources. Indeed, the significance of the stereotyped neuron

combination in the cp sensillum in the maxillary palp of culicine is remarkable from an

evolutionary and functional standpoint. The two largest neurons are tuned to animal host

attractants whereas the smallest and third neuron responds to a plant host odorant with repellent

activity. This antagonism in valence whereby attractive odorants is detected by the largest OSNs

while repellent odorants are detected by small OSNs is widespread in animals (Ng et al., 2020).

In this model, animal host odors are dominated by animal host odorants but can be modulated by

the presence of an odor that signals alternative resources such as a plant host or a source of

danger.

Innate repellency such as the one mediated by the OR49-borneol-MD3 labelled line is

probably widespread in the olfactory system of mosquitoes. However, the major challenge to

understand such mechanisms in mosquitoes and in insects is our limited odorant screening

capabilities. A better understanding of such processes will significantly increase our ability to

design potent repellent formulations targeting innate mosquito aversion.

**Material and Methods**

*Insects*

Our mosquito colonies were reared in air-controlled insect chambers at 26 ± 2 °C, under a 12

hr:12 hr photoperiod cycle. Larvae were kept in 1 L water-filled plastic containers and fed with a

ground mixture of Novocrabs feed (JBL GmbH & amp; Co, Germany). Pupae were transferred

into plastic cages until adults emerged. The adults were given 10% sucrose solution *ad libitum*.

For oviposition mosquitoes were provided water cup lines with Whatman filter paper. Females

were fed with cow blood using a membrane feeding system on days 4 and 5 following adult

emergence. *Ae. aegypti* originated from a colony established by Prof. Joel Margalit. The *Ae.*

*aegypti* Liverpool strain was provided by the Akbari lab. *Aedes albopictus* was the FPA Foshan

strain collected (Foshan, China) reared at the insectary of the University of Pavia since 2013

(Palatini et al., 2017). *Anopheles gambiae* was the G3 strain originally isolated from West Africa

(MacCarthy Island, The Gambia) in 1975 (Federica Bernardini et al., 2017). *Culex pipiens*

originated from a wild type population was provided by Dr. Laor Orshan (Ministry of Health,

Israel).

*Phylogenetic and genomic analyses*

The AaegOR49 protein sequence (AAEL001303-PA) was used as a query to identify

homologs in other mosquito species. Amino acid sequences of homologous odorant receptors

were obtained from the Vectorbase database ([vectorbase.org](vectorbase.org)) and the *Tx. amboinensis* genome

assembly (Zhou et al., 2014). The multiple sequence alignment was conducted using ClustralW

(Chenna et al., 2003). The Maximum-likelihood phylogenetic tree was constructed using

MEGAX software (Model: G, bootstraps: 5000). Exon-intron structure analysis was conducted

manually by aligning predicted amino-acid sequences to genomic regions. Approximate mapping

of the genes in chromosome location for each mosquito species and construction of synteny

maps done using Geneious prime software (Kearse et al., 2012) using latest genome assemblies

available for each mosquito species (Matthews et al., 2018, Palatini et al., 2020, Boyle et al.,

2021, Arensburger et al., 2012, Sharakhova et al., 2007).

Genomic DNA (gDNA) was isolated from whole bodies of *Tx. amboinensis* adults. gDNA

was diluted to 25ng/µL in nuclease free water and used as a template in polymerase chain

reactions to amplify odorant receptors (TambOR) using Taq polymerase and the following

primer sets: TambOr6.F1 (ATGCGCTTCTACGAGAAATAC), TambOr6.R1

(TCAGAAATTATCCTTCAGGATC); TambOr12.F1 (ATGCCATCGGTTTTCTTGGTT),

TambOr12.R1 (CTAAAACACTCGCTTCAATATC); TambOr13.F1

(ATGTTCTGCTTCAGGAAGATC), TambOr13.R1 (CTAGAAGTGGTTTTTCAATATAA);

TambOr49.F1 (ATGTTGTTCAAGAACTGTTTCC), TambOr49.R1

(TTAATAATTGAATCTTTCCTTCAG); TambOr71.F1 (ATGGGCAGCAGTGATGGTGAC),

TambOr71.R1 (CTACTGGTTGATTTTACTGAGG). Cycling conditions were 94°C for 60s; 30

cycles of 94°C for 20s, 56°C for 20s, 72°C for 30s; and a final extension of 72°C for 5 minutes.

Amplicons were analyzed by electrophoresis on a 1% agarose gel, cloned into the TOPO-TA

pCR2 plasmid, heat-shock transformed into TOP10 competent *E. coli* cells, and grown overnight

at 37°C on LB+ampicillin+X-gal agar plates. Ampicillin resistant, white colonies were isolated

using a sterile pipette tip and grown overnight 37°C in 3mL of LB+ampicillin. Plasmids were

isolated by DNA miniprep and the Sanger method was used to determine the DNA sequence in

both directions using standard T7 and M13.rev primers. TambOR nucleotide sequences were

compiled into contigs and intron/exon regions were inferred by comparing to coding sequences

described in a previous publication (Zhou et al., 2014).

*Chemical reagents*

The chemicals used for the deorphanization of receptors were obtained from Acros Organics

(Morris, NJ, USA), Alfa Aesar (Ward Hill, MA, USA), ChemSpace (Monmouth Junction, NJ,

USA), Sigma Aldrich (St. Louis, MO, USA), TCI America (Portland, OR, USA), and Thermo

Fisher Scientific (Waltham, MA, USA) and Penta Manufacturing Corp. (Livingston, NJ, USA) at

the highest purity available. *Cannabis* essential oils and sub-mixtures were formulated and

supplied by Eybna Technologies (Givat Hen, Israel) (Table S1 and Figure S1).

*Two-electrode voltage clamp of Xenopus laevis oocytes*

Experimental procedures for *Tx. amboinensis* receptor clones (Table S2) were performed as

followed: Stage V-VI *Xenopus laevis* oocytes were harvested, whole ovaries were treated with a

0.8 mg/ml collagenase solution (Sigma Aldrich, 9001-12-1) at 18°C with agitation for 3.5 hours

at 60 RPM. Following collagenase treatment, oocytes were washed 5 times in a $Ca^{+2}$ free Ringer

solution (NaCl 5.61 gr, KCl 0.14 gr, $MgCl_2$-6 $H_2O$ 0.4 gr, HEPES 1.19 gr, 1L double distilled

water [pH 7.6]). Subsequently oocytes were rinsed in a washing solution (96 mM NaCl, 2 mM

KCl, 5 mM MgCl2 and 5 mM HEPES, 100 µg/mL gentamycin pH 7.6). Finally, oocytes were

washed 5 times with incubation medium (96 mM NaCl, 2 mM KCl, 5 mM $MgCl_2$, 0.8 mM

$CaCl_2$ and 5 mM HEPES, pH 7.6), supplemented with 5% dialyzed horse serum, 50 µg/mL

tetracycline, 100 µg/mL streptomycin, and 550 µg/mL sodium pyruvate. Oocytes were allowed

to recover overnight prior to injection. Following the night, oocytes were microinjected with 27.6

ng cRNAs, TambOR49 was co-injected with its obligatory co-receptor TambORCO. These

genes were not cloned from *Tx. amboinensis* but were custom-synthesized (IDT Coralville,

USA) and supplied in a circularized plasmid. The genes were then cloned into *Xenopus* oocytes

expression vector pSP64T (DM#111) Addgene (CAT#15030) (Watertown, MA, USA), using

Takara-Clontech infusion kit (CAT# 638909). In vitro transcription was done using

mMESSAGE Mmachine sp6 Transcription kit (CAT# AM1340) Thermo Fisher (Waltham, MA,

USA).

Experimental procedures for *Ae. aegypti*, *Ae. albopictus*, *Cx. pipiens* receptor clones were

performed as follows: Stage V-VI *X. laevis* oocytes were ordered from Xenopus1 (Dexter, MI,

USA) and incubated in ND96 incubation media (96 mM NaCl, 2mM KCl, 5mM HEPES, 1.8mM

$CaCl_2$, 1mM $MgCl_2$, pH 7.6) supplemented with 5% dialyzed horse serum, 50 µg/mL

tetracycline, 100µg/mL streptomycin, 100µg/mL penicillin, and 550 µg/mL sodium pyruvate.

*AaegOr49*, *AaegOrco*, *AalbOr49*, *AalbOrco*, *CquiOr38*, and *CquiOrco* templates were

synthesized by Twist Biosciences (San Francisco, CA, USA) and cloned into the pENTR$^{TM}$

vector using the Gateway$^{R}$ directional cloning system (Invitrogen Corp., Carlsbad, CA, USA)

and subcloned into the *X. laevis* expression destination vector pSP64t-RFA. Several of these

genes were codon-optimized (Table S2). Plasmids were purified using GeneJET Plasmid

Miniprep Kit (ThermoFisher Scientific, Waltham, MA, USA) and sequenced in both directions

to confirm complete coding regions. cRNAs were synthesized from linearized pSP64t expression

vectors using the mMESSAGE mMACHINE® SP6 kit (Life Technologies). Oocytes were

injected with 27.6 nL (27.6 ng of each cRNA) of RNA using the Nanoliter 2010 injector (World

Precision Instruments, Inc., Sarasota, FL, USA).

Injected oocytes were then incubated at $18^{o}$C in 24 well-plates containing 1 ml incubation

medium as described above for 3 days prior to recording. OR response was recorded by

measuring whole cell currents using the two-electrode voltage clamp (TEVC) method. Holding

potential was maintained at −80 mV using an OC-725C oocyte clamp (Warner Instruments,

LLC, Hamden, CT, USA). Oocytes were placed in a RC-3Z oocyte recording chamber (Warner

Instruments, LLC, Hamden, CT, USA) and exposed for 8-10 s long stimuli. Currents were

allowed to return to baseline between odorant applications. Data acquisition was carried out with

a Digidata 1550A and pCLAMP10 (Molecular Devices, Sunny- vale, CA, USA). To assess OR

activation by ecologically relevant terpenes, selected terpenes were solubilized in 200 µl DMSO

prior to dilutions in ND96 buffer supplemented with 0.8 mM $CaCl_2$. For the initial screen,

selected terpenes were used at a concentration of 98 µM and all *Cannabis* essential oils

were diluted by a factor of $1.5*10^{5}$. For CRCs a stock solution of the tested compound was made

at concentration of 1M and then further diluted to the desired concentrations.

For pairwise current comparisons of (+)-borneol and (+)-camphor, two separate perfusion

systems were assembled, one for each compound. The outlet from each system was connected to

a 2 to 1 perfusion manifold. At each tested concentration, (+)-borneol and (+)-camphor were

administered consecutively. Oocyte baseline currents were allowed to fully recover between each

odorant administration. Each odorant concentration was administered for 8 seconds and the order

of odorant administration was reversed between each tested oocyte to control for any positional

effect. Representative current traces can be seen in Figure S3.

*Electropalpogram recordings*

Based on the voltage clamp data, responses from the maxillary palps of *Ae. albopictus*, *Ae.*

*aegypti*, *Tx. amboinensis*, *Cx. pipiens* and *An. gambiae* were recorded with camphor or borneol.

Electopalpogram (EPG) assays were performed according to published procedures (Wang et al.,

2017) with a few modifications. Briefly, electrically conductive gel (Parker Laboratories INC,

New Jersey, USA) was applied on the metal electrode surface, the head of a 3-6 days old virgin

female (not blood-fed) mosquito was cut and mounted on the reference electrode. The distal ends

of the maxillary palps were placed on the recording electrode, and the preparation was held using

an MP-15 micromanipulator (Syntech GmbH, Germany). EPG signals were acquired by a high-

impedance AC/DC IDAC-4 pre-amplifier, and the resulting data were analyzed with EAG Pro

version 2.1 software (Syntech GmbH, Germany). A Pasteur pipette was used as a cartridge to

deliver the odorant. Within this cartridge, 20 μL of odorants diluted in hexane were deposited at

desired dose onto a 4×20 mm Whatman filter paper strip. The odorants were delivered into a

consistent humidified airstream (at a flow rate of 50 cm/s) at approximately 2 cm from the

maxillary palps. The pulse duration was 0.1 s, and the recording time was set for 5 s. A 2-min

gap was allowed between stimuli to recover the EPG sensitivity. At least 5 individual females

were tested, for each individual, 3 technical replicates were performed with doses ranging from

0.01 ug to 100 μg. Negative controls (hexane and air) and positive controls (1-octen-3-ol) were

performed at both the beginning and end for each recording session to monitor the decline in

sensitivity of the maxillary palps. EPG peak responses were normalized to the response of

hexane.

*Generating Or49 knockout line in Ae. aegypti and isolating an OR49$^{-/-}$ homozygous line*

Genomic DNA was extracted from whole bodies of *Ae. aegypti* Liverpool individuals using the

DNeasy Blood & Tissue Kit (Qiagen, Redwood City, CA). Homology arms of the *Or49* gene

flanking 1 kb upstream and downstream of the desired insertion site (Figure S5A) were amplified

with the Q5 High Fidelity DNA polymerase (New England Biolabs, Ipswich, MA), using primer

pairs Left_Arm_Or49_FWD and Left_Arm_Or49_REV as well as Right_Arm_Or49_FWD and

Right_Arm_Or49_REV, respectively (Table S4). The QF2-ECFP DNA cassette was amplified

using the V1117A plasmid as template and the primer pair QF2_OpIE_ECFP FWD and

QF2_OpIE_ECFP REV. The cassette contains the *QF2* sequence and the 3' UTR of the *HSP70*

gene along with the *ECFP* gene under the control of the *Opie2* promoter and the 3' UTR of the

*SV40* gene. PCR bands were cut out from agarose gels and purified with Zymoclean Gel DNA

Recovery Kit (Zymo Research, Irvine, CA). Upstream and downstream homology arms as well

as the QF2-ECFP cassette PCR products were assembled into the backbone of the V1117A

plasmid (Figure S5B) using Gibson assembly reaction, following the manufacturer

recommendations. Gibson's reaction-derived plasmid product was used to transform JM109 cells

(Zymo Research), and colonies were individually grown overnight for minipreps with Zyppy

Plasmid Miniprep Kit (Zymo Research). The sequence of the plasmid V1117F-Or49 was

confirmed by restriction enzyme digestion using the enzymes ScaI-HF and AvrII as well as by

whole plasmid sequencing (Primordium Labs, Monrovia, CA). V1117F-Or49 plasmid was used

to retransform JM109 cells, which were grown for maxiprep purification using the PureLink

Expi Endotoxin-Free Maxi Plasmid Purification Kit (Thermo Fisher Scientifics, Waltham, MA).

For guide RNA synthesis, non-template reactions were carried out with the

gRNA_Left_Or49_F and gRNA_Left_Or49_R forward primers and the universal guide RNA

reverse primer (Universal-sgRNA_R). PCR bands were isolated from agarose gel and purified as

above described. Guide RNAs were synthesized with the Ambion MEGAscript kit (Thermo

Fisher Scientifics) for 4 hours at 37°C, using 300 ng of purified PCR product. Guide RNAs were

further purified with the Megaclear Kit (Thermo Fisher Scientifics).

To assess the cleavage activity of the synthesized guide RNAs, *in vitro* Cas9 cleavage assays

were carried out. A DNA fragment spanning 1,009 bp overlapping the cleavage sites of the guide

RNAs was amplified with primers Or49_cleav_F and Or49_cleav_R, DNA bands were isolated

from agarose gel, purified, and 100 ng of which was used in cleavage assays with 300 ng of

recombinant Cas9 (PNA BIO, Thousand Oaks, CA) and 100 ng of each guide RNA upon

incubation at 37°C for 1 hour (Figure S5C).

For embryo microinjection, an injection mix was prepared with the plasmid V1117F-Or49 at

500 ng/ul, gRNAs left and right at 100 ng/ul each, and recombinant Cas9 at 300 ng/ul. Mix was

filtered with the Ultrafree-MC Centrifugal Filter UFC30GUOS (Millipore, Burlington, MA).

Freshly harvested *Ae. aegypti* embryos (Liverpool strain) were injected with the transformation

mix using a custom-made insect embryo microinjector (Hive Technologies, Cary, NC) at 100 psi

pulse and 3 psi back pressures, using quartz needles pulled with a P-2000 needle puller (Sutter

Instrument Co., Novato, CA). Genomic DNA of the G1 fluorescent mosquitoes (Figure S5D)

were individually extracted with the Qiagen blood kit, and PCR amplified for Sanger Sequencing

reactions (Retrogen, San Diego, CA) using the primer pairs Or49_diag_up_F and

Or49_diag_up_R targeting the upstream and primer pair Or49_diag_down_F and

Or49_diag_down_R targeting the downstream insertion site (Figure S5E). For whole DNA

cassette sequencing (Primordium), 2.5kb DNA fragments were amplified from the DNA of G1

individuals using the primer pair Or49_diag_up_F and Or49_diag_down_R (Figure S5F).

For isolation of homozygous knockout individuals, pupae were sex sorted, and single pairs

were transferred into Narrow *Drosophila* Vials (Genesee Scientific, El Cajon, CA), filled with 10

mL of deionized water (DI) and closed with cotton stoppers. Upon emergence, water was

drained, and sugar cottons were placed into the vials. Mosquitoes were allowed to mate for 5

days in the vials, and then all individuals were transferred to a single cage for blood feeding.

Three days after blood feeding, females were individually transferred to a fresh vial containing a

piece of brown paper towel wetted with 1 mL DI water. Females were allowed to lay eggs, and

each egg batch was individually hatched in 250 mL Clear Pet Cups (9oz). Larvae were screened

for fluorescence, and the batches that resulted in 100% fluorescent individuals were grown to

adulthood and intercrossed to confirm homozygosity in the following generation. Two out of

thirty egg batches resulted in homozygous offspring. To further confirm the homozygous status

of these individuals, genomic DNA of three pools of 10 males, 10 females, and a mix of males

and females were PCR amplified with the primer pair Or49_diag_up_F and Or49_diag_down_R,

which resulted in a single band for homozygous individuals and two bands for heterozygous

individuals (Figure S5G).

*Single-sensillum electrophysiology*

Single sensillum recordings from capitate peg sensilla on the maxillary palp of wild type *Ae.*

*aegypti* (Liverpool)*, Ae. albopictus* (FPA)*, Cx. quinquefasciatus* (Thai) and *An. gambiae sensu*

*stricto* (G3) and *Ae. aegypti Or49*[-/-] mutant mosquitoes were performed using an established protocol (Herre et al., 2022). Briefly, a single cold-anesthetized female mosquito was placed ventrally on a microscope slide, which was covered with a double-sided sticky tape. Another piece of tape was placed on top of the dorsal side of the body and then pushed against the underlying tape to keep the insect from moving. The maxillary palps were then gently mounted on a piece of double-sided tape. Thereafter, the specimen was placed under a light microscope (700× magnification) to visualize the capitate peg sensilla for extracellular recordings.

Two tungsten microelectrodes were electrolytically sharpened and mounted in holders (Ockenfels Syntech GmbH, Buchenbach, Germany). The ground electrode was placed in the eye of the mosquito, while the recording electrode, which was attached to a high-impedance universal single ended probe (Ockenfels Syntech GmbH), was introduced into a sensillum with a piezo-controlled micro manipulator until electrical contact was achieved. Signals were routed to an Intelligent Data Acquisition Controller (Ockenfels Syntech GmbH), and were visualized on a computer screen. Spikes were quantified offline using the established nomenclature for the sensory neurons (Ghaninia et al., 2019). The number of spikes counted during a 0.5 s stimulus delivery interval was subtracted from the number of spikes counted during a 0.5 s prestimulus period, and the result was multiplied by 2 to obtain the activity of individual sensory neurons housed in the capitate peg sensillum as a spikes/s measurement.

To investigate the physiological activity of the A, B and C neurons housed in single capitate peg sensilla, $CO_2$, *R*-(–)-1-octen-3-ol and (+)-borneol were used: gas cylinders containing metered amounts of $CO_2$ (300, 600, 1200, 2400, or 4800 ppm) and oxygen (20%), balanced by nitrogen (Strandmöllen AB, Ljungby, Sweden) were used to assess the activity of the A neuron; serial decadic dilutions of *R*-(-)-1-octen-3-ol (CAS: 3687-48-7, Penta Manufacturing, Livingston,

USA), diluted in paraffin oil, were used to assess the activity of the B neuron; and serial decadic

dilutions of (+)-borneol (CAS: 464-43-7, Sigma-Aldrich, St. Louis, MO, USA), diluted in

diethyl ether (SupraSolv, Billerica, MA, USA) were used to assess the activity of the C neuron.

Pasteur pipettes were filled with the metered $CO_2$ and used immediately to stimulate the

preparation. A 15 μL aliquot of each dilution of $R$-(–)-1-octen-3-ol and (+)-borneol was pipetted

onto a filter paper (5 mm × 15 mm) inserted inside a Pasteur pipette, and the diethyl ether was

allowed 15 min to evaporate, before being used for stimulus delivery. All stimuli were delivered

into the airstream passing over the maxillary palp preparation.

*Calcium imaging in the Ae. aegypti antennal lobe (AL)*

Odor-evoked responses in the *Ae. aegypti* antennal lobe (AL) were imaged using the brp-

QF2>QUAS-GCaMP6s progeny from the brp-QF2 and QUAS-GCaMP6s parental lines (Zhao et

al., 2022). A total of eight 6-8 day-old female mosquitoes were used for all calcium experiments.

Each mosquito was cooled on ice and transferred to a Peltier-cooled holder that allowed the

mosquito head to be fixed to a custom stage using ultraviolet glue. The stage permits the

superfusion of saline to the head capsule and space for wing and proboscis movement

(Lahondère et al., 2019; Melo et al., 2020). Once the mosquito was fixed to the stage, a window

in its head was cut to expose the brain, and the brain was continuously superfused with

physiological saline (Vinauger et al., 2019). Calcium-evoked responses in the AL were imaged

using the Prairie Ultima IV two-photon excitation microscope (Prairie Technologies) and Ti-

Sapphire laser (Chameleon Ultra; Coherent; at 1910 mW power). Experiments were performed

at 75 µm depth from the ventral surface of the AL, allowing characterization of the mediodorsal

glomerular responses to olfactory stimuli and allowing these glomeruli to be repeatedly imaged

across preparations. To record odor-evoked responses, images were collected from a 110 µm ×

83 µm plane at 2 Hz (line period of 1 ms), and for each odor stimulus, images were acquired for

25 s, starting 10 s before the stimulus onset. Image data were imported into Matlab (v2017;

Mathworks, Natick, Massachusetts) for Gaussian filtering (2×2 pixel; $\sigma = 1.5$-3) and alignment

using a single frame as the reference at a given imaging depth and subsequently registered to

every frame to within ¼ pixel. Odor stimuli were diluted to $10^{-3}$ and $10^{-4}$ concentration in hexane

(>99.5% purity; Sigma), with hexane used as the solvent control. During an experiment, odor

stimuli were separated by intervals of 120 s to avoid receptor adaptation, and odor syringes were

used once per preparation to prevent decreased concentration within the cartridge. Calcium

responses are calculated as the change in fluorescence and time-stamped and synced with the

stimulus pulses. After an experiment, the AL was sequentially scanned at 0.5 µm depths from the

ventral to the dorsal surface to provide glomerular assignment and registration between

preparations. Glomeruli (1 $\mu m^3$ voxel) were mapped and registered based on the positions and

odor-evoked responses of the putative AL3, MD1-3, and AM2 glomeruli, using available AL

atlases (Shankar and McMeniman, 2020; Wolff et al., 2023) and the Amira software (v. 6.5, FEI

Houston Inc.).

*Behavioral assay*

The role of borneol in human host-seeking female mosquitoes was examined with an arm-in-a-

cage assay described previously (Dekel et al., 2022) with minor modifications (Additional File

1A) in an air-conditioned room (26±1 °C, 60±5 % RH). Briefly, the experimenter's hand was

presented to fifteen 5–10 days post emergence adult females. A three-dimensional-printed

interlocking ring with a diameter of 55 mm was used in this experiment (Additional File 1B).

The ring was placed over the dorsal side of a nitrile glove (powder-free latex). To evoke human

odor and prevent mosquito bites, we replaced the nitrile between the two ring components with a

plastic screen. The interlocking ring included a central odorant delivery platform (Additional File

1C) comprising a 10-mm-diameter cover glass and two 5-mm-diameter filter discs

(WHA10016508; Merck) for the evaluation of VOCs that interact with or blend with human odor

(stl files are provided in additional files 2 and 3). Plastic screen, nitrile glove and odorant

delivery platform were replaced between repetitions to avoid contaminations. Mosquitoes were

placed in a 20.3-$cm^3$ metal cage located in an experimental room with a vent, under a video

camera (EOS 70D, lance: MACRO 0.25/0.8ft; Canon Inc., Tokyo, Japan) and a light ring.

Mosquitoes were allowed to acclimate for ten minutes before recording. Mosquito behavior was

recorded for 10 min at 25 frames per second. The number of mosquito detection on the screen

and ring were automatically counted using a custom YOLOv5 model (see object detection

section below). The possible number of detections per experiment was calculated as 225,000,

obtained by multiplying the number of mosquitoes in the cage (15) by the frame rate (25 frames

per second) and the duration of each experiment (600 seconds). We used the solvent diethyl ether

(DEE) as a vehicle. On a blank filter disc, 25 μL of DEE, and DEE with racemic borneol (1M,

3.58 mg) were deposited on the filter paper and allowed to evaporate for 2 min prior to mosquito

exposure outside the experimental room to avoid contamination. To enhance attractiveness, at

the beginning of each experiment, the experimenter first rubbed the ring-mounted glove against

the shirt and skin for 1 minute. Additionally, the experimenter blew twice, once into the cage and

once into the glove. All experiments were conducted during the first 4 hours of the diurnal period

and lasted 10 min. This schedule was chosen for practical reasons and because mosquitoes

consistently exhibited attraction to the human hand. Mosquito detection was normalized to the

mean detection of the control $(\frac{detections\ for\ minute\ i}{mean\ detection\ for\ all\ minutes\ (control)} * 100$ . Statistical

comparisons of mosquito detections per minute were carried out with Mann Whitney U test (*P*-

value < 0.001, *n* = 3). For data variation across time and repetitions, see Figure S7. Data was

analyzed using R Studio (Team, 2021) The R-code and behavioral data can be found in

additional files 4-6.

**YOLOv5 Model Training and Custom Dataset**

In this study, we trained the YOLOv5l model using a customized dataset comprising images

from the bioassay, along with additional data from previously published literature (Janson et al.,

2023), to investigate the ecological role of borneol on mosquitoes. The training process involved

100 epochs with a batch size of 32, utilizing separate sets of images and corresponding

annotations for training, validation, and testing (4,759, 1,584, and 1,580 images, respectively).

To assess the model's performance, we employed the mean Average Precision (mAP) metric

across a range of intersection-over-union (IoU) thresholds from 50% to 95% (mAP50-95). The

resulting mAP50-95 value of 0.662 indicates the average precision considering different IoU

thresholds. At a 50% IoU threshold, the mAP50 was measured as 0.982, demonstrating high

accuracy in detection. Furthermore, the recall value of 0.969 indicates a high proportion of true

positive detections, while the precision value of 0.974 reflects accurate identification of

mosquitoes. It's worth noting that detections were collected with a minimum confidence

threshold of 0.5 to ensure reliable results. Video samples with automated detection are available

in additional file 8.

**Figure legends**

**Figure 1. The phylogenetically-conserved gene encoding OR49. A**) Amino-acid sequence

alignment of OR49 homologs in *Culex quinquefasciatus* (Cqui), *Aedes albopictus* (Aalb), *Aedes*

*aegypti* (Aaeg), *Toxorhynchites amboinensis* (Tamb), and *Anopheles gambiae* (Agam). The top

chart presents the amino-acid consensus. High consensus marked in green to yellow, low

consensus are in red. **B**) Amino-acid sequence identity matrix. Intensity of shading indicates

percentage of homologies. **C**) Phylogenetic tree of mosquito OR49 proteins (green). Closest

brachyceran homologs (red branches) include *Glossina fuscipes* (Gfuc) and *Drosophila*

*melanogaster* (Dmel). Bootstrap support above 50% are shown. Gene exon structure of OR49

genes and other phylogenetically-related genes. **D**) Exon compoistion of the OR49 group and

related homologs. Exons are labeled in white and black. The nucleotide positions of exon-exon

boundaries are shown and intron phases are color-coded. **E**) Chromosomic locations of Or49

homolog genes (green) and indolergic ORs (indolORs, in red). **F**) Syntenic relationships of

OR49 genes in mosquito species and gene structure of OR49 genes.

**Figure 2. Odorant receptor 49 is a borneol receptor. A**) The 4th segment of the mosquito

maxillary palps houses three neurons, including the $CO_2$ capitate-peg neuron (cpA) expressing

three gustatory receptors, the 1-octen-3-ol-sensitive neuron (cpB), and the orphan neuron (cpC).

**B**) The response profile of *Aedes albopictus* OR49 (AalbOR49) in response to odorant mixtures

(100 μM) belonging to a variety of chemical classes highlights the activity of ketones.

Terpenoids, including camphor and fenchone are the most efficacious activators of AalbOR49.

Bicyclic monoterpenoids are labeled in green. Statistical differences were evaluated by ANOVA

followed by a Kruskal-Wallis multiple comparisons test. ns, non-significant, *$p < 0.0332$, **$p <$

$0.0021$, ***$p < 0.0002$ and ****$p < 0.0001$. Data indicate the means ± SEM. **C**) Pineapple Haze

(PH), OG Kush (OGK), Adom9 (A9), and Master Kush (MK) elicited the highest currents from

*Toxorhynchites amboinensis* OR49 (TambOR49). Borneol content is indicated on the right Y-

axis. **D**) Response of TambOR49 to 6 *Cannabis* sub-mixtures and the constituents from sub-

mixture 2. **E**) Concentration-response relationships of TambOR49 to the two borneol

enantiomers. **F**) Concentration-response relationships of TambOR49, *Culex quinquefasciatus*

OR38 (CquiOR38), *Ae. aegypti* OR49 (AaegOR49) and AalbOR49 in response to (+)-borneol

and (+)-camphor. Effective concentrations at 50% of the maximal response ($EC_{50}$) are circled.

**G**) Pairwise comparisons of the current responses of all four mosquito species OR49s elicited by

increasing concentrations of (+)-borneol and (+)-camphor. Statistical differences were evaluated

by multiple t-tests. *$p < 0.0332$, **$p < 0.0021$, ***$p < 0.0002$ and ****$p < 0.0001$. Data indicate

the means ± SEM. Representative current traces can be found in Figure S3.

**Figure 3. The maxillary palp of culicine mosquitoes respond to borneol and camphor. A**)

Electropalpogram (EPG) responses in *Aedes albopictus*, *Aedes aegypti*, *Culex pipiens*, and

*Anopheles gambiae*. EPG responses to 1-octen-3-ol (blue, 10 μg), camphor (purple, 10 μg), and

borneol (green, 10 μg) enantiomers relative to hexane. Representative traces for each odorant are

shown below the x-axis. Statistical differences were evaluated via one-way anova. *$p < 0.05$, **p

$< 0.01$, ***$p < 0.005$ and ****$p < 0.001$. Data indicate the means ± SEM, $n = 15$. **B**) Dose-

response relationships of the maxillary palps of three culicine mosquito species in response to

increasing concentrations of (+)-borneol and (+)-camphor. Representative traces are shown on

the right. Statistical differences were evaluated by multiple t-tests. Data indicate the means ±

SEM, $n = 15$.

**Figure 4. Species-dependent response to (+)-borneol of the C neuron of the capitate peg**

**sensilla and its reliance of *Or49* in *Ae. aegypti*. A**) A tungsten electrode was used to record

action potentials (spikes) from the capitate peg sensillum on the maxillary palp, which houses

three neurons. The cpA neuron responds to $CO_2$ (large spike in traces), the cpB neuron responds

to 1-octen-3-ol (medium size spikes) and the cpC neuron responds to borneol (small size spikes).

**B**) Species- and dose-dependent response of the C neuron to (+)-borneol with mean $\pm$ s.e.m. at

each concentration. The arrow and dotted line indicate the borneol dose to the shown

representative traces. *Ae. aegypti* ($n = 10$), *Ae. albopictus* ($n = 7$), *Cx. quinquefasciatus* ($n = 2$),

*Ae. aegypti Or49-/-* ($n = 8$), *An. gambiae* ($n = 15$). Representative traces of the cp sensilla of *Ae.*

*albopictus* and *Cx. quinquefasciatus* are available in Figure S6. **C-D**) The spontaneous activity of

the three sensory neurons in the capitate peg sensilla of *Ae. aegypti* and *Anopheles gambiae s.s.*.

In the top traces, note the differences in spike amplitude of the A, B (orange markers) and C

neurons (green markers). **C**) *R*-(–)-1-octen-3-ol and (+)-borneol elicit responses in the B and C

neurons of *Ae. aegypti*, respectively. The response to (+)-borneol is abolished in *Or49$^{-/-}$* mutant

mosquitoes. **D**) The C neuron of *An. gambiae s.s.* does not respond to (+)-borneol.

**Figure 5. Borneol elicits robust responses in the *Ae. aegypti* AL. A**) Schematic of the two-

photon setup used to record calcium dynamics in the mosquito antennal lobe (AL). **B**) AL atlas,

highlighting the MD1 (blue), MD2 (orange), and MD3 (green) glomeruli. Non-responsive AL

glomeruli (grey) and the mediodorsal glomeruli were registered and mapped to previously

published atlases. **C**) Pseudo color plot from a single preparation of $\Delta F/F_0$ calcium responses (0-

0.8 scale) to (+)-borneol ($10^{-4}$ dilution), at a depth of 75 µm from the surface of the AL. Borneol

evoked a strong response in the region of interest mapped to the MD3 glomerulus (highlighted in

white). **D**) Glomerular responses ($\Delta F/F_0$) to (+)-borneol for the MD1, MD2, and MD3 glomeruli.

Lines are the mean of one glomerulus ($n$ = 4-8 preparations); shaded areas are the SEM. The

grey bar denotes stimulus duration (2 s). **E**) Same as in D, except the glomeruli were stimulated

with 1-octen-3-ol ($10^{-4}$ dilution). **F**) Same as in D, except the glomeruli were stimulated with

$CO_2$ (5%). (**G**) Tuning curve for the MD3 glomerulus to a limited panel of 8 odorants, each

tested at $10^{-4}$ concentration. The MD3 glomerulus (bars in green) showed significant calcium

responses to enantiomers of borneol and camphor compared to the solvent control (Kruskal-

Wallis test: $\chi^2$ = 37.1, P < 0.0001; posthoc multiple comparisons: p < 0.05). Bars represent the

mean ± SEM.

**Figure 6. Borneol inhibits host-seeking female *Ae. aegypti* (Liverpool) behavior.** Average

number of mosquito detections per minute (normalized to the control group) in the region of

interest (ROI) in response to control (DEE) and treatment (DEE + racemic borneol). Mosquito

detection on the ROI was normalized to the control. The statistical comparison involved

analyzing 30 data points of accumulative detections per minute for each treatment. The data was

analyzed with Pairwise comparisons using Wilcoxon rank sum test (p-value < 0.001, $n$ = 3). **B**)

A representative of the number of mosquitoes detected in the ROI as a function of time is shown

on the right side of each treatment (right y-axis).

**Supplementary information**

**Figure S1. List of volatile organic compounds present in *Cannabis* essential oils.** Each

Cannabis blend represented by a distinct color is composed of VOCs belonging mainly to

different terpene categories but also to non-terpene chemical classes.

**Figure S2. VOC content in *Cannabis* essential oils.**

**Figure S3. Representative current traces of mosquito OR49 proteins. A**) Representative traces of TambOR49+TambOrco injected oocytes vs monoterpenoid blends and compounds. Pineapple Haze (PH), Adom #9 (A9), O.G Kush (OGK), Master Kush (MK) and Jack Herrer (JH) triggered a measurable response. (±)-camphor, (+)-camphor, (+)-fenchone and eucalyptol were used as positive controls at concentration of 0.1 mM, all *Cannabis* essential oils were diluted by a factor of $1.5 \times 10^5$. **B**) Representative trace of TambOR49+TambOrco vs sub-mixtures No. 1 through 6, which were formulated based on overlapping active compounds from *Cannabis* essential oils. All sub-mixtures were diluted by a factor of $1.5 \times 10^6$. Only sub-mixture No.2 activated TambOR49+TambOrco. (+)-Camphor was used as positive control at concentration of 0.1 mM. **C**) Representative trace of TambOR49+TambOrco vs the components of sub-mixture No. 2. (±)-Borneol elicited currents 8-fold larger than the other tested single monoterpenoids. All compounds were tested at concentration of 0.1 mM. **D**) Representative traces of TambOR49+TambOrco concentration response curves (CRCs). Top left TambOR49+TambOrco vs (-)-borneol, Top right TambOR49+TambOrco vs (+)-borneol and bottom left TambOR49+TambOrco vs (+)-camphor. **E**) Representative traces of AaegOR49 CRC's. Left AaegOR49 vs (–)-Borneol; right, AaegOR49 vs (+)-Borneol. **F**) Representative traces of CquiOR38+CquiOrco CRCs in response to increasing concentrations of (+)-camphor (left) and (+)-borneol (right). **G**) Representative traces of AaegOR49+Orco CRCs in response to increasing concentrations of (+)-camphor (left) and (+)-borneol (right). **H**) Representative traces of AalbOR49+AalbOrco CRCs in response to increasing concentrations of (+)-camphor (left)

and (+)-borneol (right). **I**) Representative traces of pairwise current comparisons between (+)-

camphor and (+)-borneol. From top to bottom, TambOR49+TambOrco, CquiOR38+CquiOrco,

AaegOR49+AaegOrco and AalbOR49+AalbOrco. Down arrows indicate odorant

administrations and concentrations. All concentrations are in micromolar [µM].

**Figure S4.** *Anopheles gambiae* **OR28 (AgamOR28) does not respond to borneol. A**)

Concentration-response of AgamOR28 in response to increasing concentrations of four plant

volatile organic compounds, including aromatic (acetophenone and α-terpineol) and terpenoid

compounds (α-pinene and (+)-borneol). EC$_{50}$ values shown in the inset are in the low millimolar

range. **B**) Representative current traces of AgamOR28-Orco activation by 4 plant volatile

organic compounds. Arrowheads above the traces indicate the onset of the odorant stimulus.

**Figure S5. Construction of** *Ae. aegypti Or49* **knockout line. A**) Diagram depicting the *Ae.*

*aegypti Or49* gene and the gRNA target sites on the top and the insertion cassette on the bottom

flanked by the upstream (Left) and downstream (Right) homology arms. The insertion cassette

encompasses the *QF2* sequence and the 3' UTR of the *HSP70* gene along with the *ECFP* gene

under the control of the *Opie2* promoter and the 3' UTR of the *SV40* gene. Arrows indicate

primer binding sites. Primers P1-12 are listed in Table S4. **B**) Plasmid V1117F-Or49 map. **C**) *In*

*vitro* Cas9 cleavage assay. Guide RNAs left and right targeting the *Or49* gene almost completely

digested PCR fragment containing target sequence. (–) negative controls without guide RNAs

show no PCR fragment digestion. **D**) Sanger sequencing of amplified PCR fragments of G1

transgenic male individuals unveiling the sequences overlapping upstream (left) and downstream

(right) of the DNA cassette inserted by homology directed repair. **E**). Sequencing of G1

individuals showing the complete insertion of the DNA cassette containing *QF2* and *ECFP*. The

presence of a single point mutation was confirmed as an artifact and excluded by Sanger

sequencing. **F**) Diagnostic PCR reactions showing the presence of double bands in single G1

heterozygous individuals and single upper band in pools of 10 individuals of the *OR49*[-/-]

homozygous line.

**Figure S6. Response to (+)-borneol of the cpC neuron of the capitate peg sensilla of *Ae.***

***aegypti* and *Cx. quinquefasciatus*.** Representative SSR traces from the *Ae. albopictus* and *Cx.*

*quinquefasciatus* cp sensilla stimulated with 0.01% (+)-borneol. Note the differences in spike

amplitude of the A, B (orange markers) and C neurons (green markers).

**Figure S7. Behavior replicates. Detection variations in control and borneol treatments. A**)

Replicate behavioral experiments with the arm-in-a-cage assay. **B**) Box plot showing the

detection variation of each experimental replicate in the control and borneol treatments. The y-

axis represents the number of mosquito detections per minute over a 10-minute period. The x-

axis indicates the replicate. Number. **C**) Accumulative detections over time. The y-axis

represents the accumulative number of mosquito detection per minute.

**Table S1. List of blends and single compounds used in the pharmacological screen.**

**Table S2. List of *Or* genes.**

**Table S3. Raw pharmacological data.**

**Table S4. Primer list (*Or49* knockout).**

**Additional File 1: Diagrams of the odor delivery system.** The odor delivery system is

composed of interlocking top and bottom rings attached to a removable chemical holder

(overview, top, bottom and side views are provided). Dimensions are provided in millimeters.

**Additional file 2: Hand rings.stl.** File format for 3D-printing of the two complementary hand

rings.

**Additional file 3: Chemical holder.stl.** File format for 3D-printing of the chemical holder

component.

**Additional file 4: Behavioral data.** CSV file of collected mosquito visits on the ROI.

**Additional file 5 and 6: Codes for analyzing behavioral data.** R file to analyze the data and

generate figure 6 (arm-in-a-cage_with_yolov5.R) and S5 (Supplementary_plot.R). Post-

processing was carried out using Adobe Illustrator.

**Additional file 7. Video recording example of the of the female *Ae. aegypti* antennal lobe.**

**Additional file 8. Video recording example of the arm-in-a-cage assay.** Thirty-second-long

recording example at the 4-minute mark of mosquito landing behavior in the presence of the

vehicle diethyl ether (DEE) and racemic borneol.

**Authors contributions**

RJP initiated the functional characterization of OR49. JB conceived the study. YV identified

borneol as a key OR49 ligand. YW conducted mosquito electropalpograms. RMH identified

camphor as an OR49 agonist. OSA and IVC-A engineered the *Ae. aegypti Or49* knockout strain.

MG and RI designed and conducted the single sensillum recordings. ESS conducted the

behavioral experiments. DP and PP conducted gene annotations, phylogenetic analyses and

described OR49 syntenic relationships. DR completed the *Toxorhynchites amboinensis* OR

genomic PCRs and sequencing. EY provided genetic constructs and injectable mRNA. AW

provided funding and scientific oversight. JB wrote the manuscript with individual contributions

from all the authors.

**Acknowledgements**

We thank Ziv Kassner for consultation, processing and analysis guidance, and technical support

in the creation of the mosquito detection model. We are especially grateful to Nadav Eyal at

Eybna Terpene Based Technologies for their time and supply of *Cannabis* essential oils. We are

grateful to Dr. Laor Orshan (Ministry of Health, Israel) for providing a colony of *Culex pipiens*.

Dr. Pitts and Bohbot are co-corresponding authors.

**References**

Abdelgaleil, S.A.M., Al-Nagar, N.M.A., Abou-Taleb, H.K., Shawir, M.S., 2022. Effect of
   monoterpenes, phenylpropenes and sesquiterpenes on development, fecundity and fertility of
   Spodoptera littoralis (Boisduval). Int. J. Trop. Insect Sci. 42, 245–253.
   https://doi.org/10.1007/s42690-021-00539-y

Abdelgaleil, S.A.M., Mohamed, M.I.E., Badawy, M.E.I., El-arami, S.A.A., 2009. Fumigant and
   contact toxicities of monoterpenes to *Sitophilus oryzae* (L.) and *Tribolium castaneum*
   (Herbst) and their inhibitory effects on acetylcholinesterase activity. J. Chem. Ecol. 35, 518–
   525. https://doi.org/10.1007/s10886-009-9635-3

Afify, A., Galizia, C.G., 2014. Gravid females of the mosquito *Aedes aegypti* avoid oviposition
   on m-cresol in the presence of the deterrent isomer p-cresol. Parasite Vector 7, 315.
   https://doi.org/10.1186/1756-3305-7-315

Alavez-Rosas, D., Socorro-Benitez, C., Cruz-Esteban, S., 2022. Repellent and adulticidal effect
   of essential oils mixtures on *Aedes aegypti* females. Int J Trop Insect Sc 42, 1885–1892.
   https://doi.org/10.1007/s42690-021-00716-z

Amer, A., Mehlhorn, H., 2006. Repellency effect of forty-one essential oils against *Aedes*,
   *Anopheles*, and *Culex* mosquitoes. Parasitol Res 99, 478–490. https://doi.org/10.1007/s00436-
   006-0184-1

Andreazza, F., Valbon, W., Dong, K., 2023. Transfluthrin enhances odorant receptor-mediated
spatial repellency in *Aedes aegypti*. Pestic Biochem Phys 192, 105387.
https://doi.org/10.1016/j.pestbp.2023.105387

Anton, S., Loon, J. van, Meijerink, J., Smid, H., 2003. Central projections of olfactory receptor
neurons from single antennal and palpal sensilla in mosquitoes. Arthropod Structure &
Development 32, 319-327. https://doi.org/10.1016/j.asd.2003.09.002

Baker, M., 1996. Fur rubbing: Use of medicinal plants by capuchin monkeys (*Cebus capucinus*).
Am. J. Primatol. 38, 263–270. https://doi.org/10.1002/(sici)1098-2345(1996)38:3<;263::aid-
ajp5>3.0.co;2-x

Benelli, G., Flamini, G., Fiore, G., Cioni, P.L., Conti, B., 2013. Larvicidal and repellent activity
of the essential oil of *Coriandrum sativum* L. (Apiaceae) fruits against the filariasis vector
*Aedes albopictus* Skuse (Diptera: Culicidae). Parasitol Res 112, 1155–1161.
https://doi.org/10.1007/s00436-012-3246-6

Bohbot, J., Pitts, R.J., Kwon, H.W., Rutzler, M., Robertson, H.M., Zwiebel, L.J., 2007.
Molecular characterization of the Aedes aegypti odorant receptor gene family. Insect
molecular biology 16, 525–537. https://doi.org/10.1111/j.1365-2583.2007.00748.x

Bohbot, J.D., Dickens, J.C., 2010. Insect repellents: modulators of mosquito odorant receptor
activity. PLoS ONE 5, e12138. https://doi.org/10.1371/journal.pone.0012138

Bohbot, J.D., Fu, L., LE, T.C., Chauhan, K.R., Cantrell, C.L., Dickens, J.C., 2011. Multiple
activities of insect repellents on odorant receptors in mosquitoes. Medical and veterinary
entomology 25, 436–444. https://doi.org/10.1111/j.1365-2915.2011.00949.x

Carey, A.F., Wang, G., Su, C.-Y., Zwiebel, L.J., Carlson, J.R., 2010. Odorant reception in the
malaria mosquito *Anopheles gambiae*. Nature 464, 66–71.
https://doi.org/10.1038/nature08834

Cloyd, R.A., Marley, K.A., Larson, R.A., Dickinson, A., Arieli, B., 2011. Repellency of
naturally occurring volatile alcohols to fungus gnat *Bradysia* sp. nr. *coprophila* (Diptera:
Sciaridae) adults under laboratory conditions. J Econ Entomol 104, 1633–1639.
https://doi.org/10.1603/ec11066

Conti, B., Benelli, G., Leonardi, M., Afifi, F.U., Cervelli, C., Profeti, R., Pistelli, L., Canale, A.,
2012. Repellent effect of *Salvia dorisiana*, *S. longifolia*, and *S. sclarea* (Lamiaceae) essential
oils against the mosquito *Aedes albopictus* Skuse (Diptera: Culicidae). Parasitol Res 111,
291–299. https://doi.org/10.1007/s00436-012-2837-6

Dayaram, L., Khan, A., 2016. Repellent, fumigant and contact toxicity of *Salvia officinalis*,
*Rosmarinus officinalis* and *Coriandrum sativum* against *Callosobruchus maculatus* (Fab.)
(Coleoptera:Bruchidae). NAAS 34, 893–902.

Dekel, A., Sar-Shalom, E., Vainer, Y., Yakir, E., Bohbot, J.D., 2022. The oviposition cue indole inhibits animal host attraction in *Aedes aegypti* (Diptera: Culicidae) mosquitoes. Parasite Vector 15, 422. https://doi.org/10.1186/s13071-022-05545-8

Dekel, A., Yakir, E., Bohbot, J.D., 2019. The sulcatone receptor of the strict nectar-feeding mosquito *Toxorhynchites amboinensis*. Insect Biochemistry and Molecular Biology 111, 103174. https://doi.org/10.1016/j.ibmb.2019.05.009

Dickens, J.C., Bohbot, J.D., 2013. Mini Review: Mode of action of mosquito repellents. Pesticide Biochemistry and Physiology 106. 149–155.

Drapeau, J., Fröhler, C., Touraud, D., 2009. Repellent studies with *Aedes aegypti* mosquitoes and human olfactory tests on 19 essential oils from Corsica, France. Flavour and Fragrance Journal 24, 160-169. https://doi.org/10.1002/ffj.1928

Enan, E., 2001. Insecticidal activity of essential oils: octopaminergic sites of action. Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol. 130, 325–337. https://doi.org/10.1016/s1532-0456(01)00255-1

Erdelyan, C.N.G., Mahood, T.H., Bader T.S.Y., Whyard, S. 2011. Functional validation of the carbon dioxide receptor genes in *Aedes aegypti* mosquitoes using RNA interference. Insect molecular biology 21, 119-127. https://doi.org/10.1111/j.1365-2583.2011.01120.x

Ghaninia, M., Majeed, S., Dekker, T., Hill, S.R., Ignell, R., 2019. Hold your breath – Differential behavioral and sensory acuity of mosquitoes to acetone and carbon dioxide. Plos One 14, e0226815. https://doi.org/10.1371/journal.pone.0226815

Gillies, M.T., 1980. The role of carbon dioxide in host-finding by mosquitoes (Diptera: Culicidae): a review. Bull Entomol Res 70, 525–532.

Gordon, U., Ruther, J., BErnier, U.R., Rose, A., Geier, M., 2018. Development and evaluation of push-pull control strategies against *Aedes aegypti* (Diptera: Culicidae), in: Norris, E.J., Coats, J.R., Gross, A.D., Clark, J.M. (Eds.), Advances in the Biorational Control of Medical and Veterinary Pests. ACS Publications, pp. 187–204. https://doi.org/10.1021/bk-2018-1289.ch010

Grant, A.J., Dickens, J.C., 2011. Functional characterization of the octenol receptor neuron on the maxillary palps of the yellow fever mosquito, *Aedes aegypti*. PLoS ONE 6, e21785. https://doi.org/10.1371/journal.pone.0021785

Grant, A.J., O'Connell, R.J., 1996. Electrophysiological responses from receptor neurons in mosquito maxillary palp sensilla., in: Block, E. (Ed.), Ciba Found Symp. John Wiley & Sons, Ltd., pp. 233-48-discussion 248-53-281–4.

Herre, M., Goldman, O.V., Lu, T.-C., Caballero-Vidal, G., Qi, Y., Gilbert, Z.N., Gong, Z., Morita, T., Rahiel, S., Ghaninia, M., Ignell, R., Matthews, B.J., Li, H., Vosshall, L.B.,

Younger, M.A., 2022. Non-canonical odor coding in the mosquito. Cell 185, 3104-3123.e28. https://doi.org/10.1016/j.cell.2022.07.024

Hill, C.A., Fox, A.N., Pitts, R.J., Kent, L.B., Tan, P.L., Chrystal, M.A., Cravchik, A., Collins, F.H., Robertson, H.M., Zwiebel, L.J., 2002. G protein-coupled receptors in *Anopheles gambiae*. Science (New York, NY) 298, 176–178. https://doi.org/10.1126/science.1076196

Höld, K.M., Sirisoma, N.S., Ikeda, T., Narahashi, T., Casida, J.E., 2000. α-Thujone (the active component of absinthe): γ-Aminobutyric acid type A receptor modulation and metabolic detoxification. Proc. Natl. Acad. Sci. 97, 3826–3831. https://doi.org/10.1073/pnas.070042397

Hwang, Y.-S., Wu, K.-H., Kumamoto, J., Axelrod, H., Mulla, M.S., 1985. Isolation and identification of mosquito repellents in *Artemisia vulgaris*. J Chem Ecol 11, 1297–1306. https://doi.org/10.1007/bf01024117

Iovinella, I., Pelosi, P., Conti, B., 2014. A rationale to design longer lasting mosquito repellents. Parasitology research 113, 1813–1820. https://doi.org/10.1007/s00436-014-3827-7

Janson, K.D., Carter, B.H., Jameson, S.B., Verges, J.E. de, Dalliance, E.S., Royse, M.K., Kim, P., Wesson, D.M., Veiseh, O., 2023. Development of an automated biomaterial platform to study mosquito feeding behavior. Frontiers Bioeng Biotechnology 11, 1103748. https://doi.org/10.3389/fbioe.2023.1103748

Jocher, G., Chaurasia, A., Stoken, A., Borovec, J., NanoCode012, Kwon, Y., Michael, K., TaoXie, Fang, J., imyhxy, Lorna, Yifu, zeng, Wong, C., V, A., Montes, D., Wang, Z., Fati, C., Nadar, J., Laughing, UnglvKitDe, Sonck, V., tkianai, yxNONG, Skalski, P., Hogan, A., Nair, D., strobel, M., Jain, M., 2022. ultralytics/yolov5: v7.0 - YOLOv5 SOTA Realtime Instance Segmentation. https://doi.org/10.5281/zenodo.7347926

Kongkaew, C., Sakunrag, I., Chaiyakunapruk, N., Tawatsin, A., 2011. Effectiveness of citronella preparations in preventing mosquito bites: systematic review of controlled laboratory experimental studies. Trop. Med. Int. Heal. 16, 802–810. https://doi.org/10.1111/j.1365-3156.2011.02781.x

Kordali, S., Aslan, I., Çalmaşur, O., Cakir, A., 2006. Toxicity of essential oils isolated from three *Artemisia* species and some of their major components to granary weevil, *Sitophilus granarius* (L.) (Coleoptera: Curculionidae). Ind Crop Prod 23, 162–170. https://doi.org/10.1016/j.indcrop.2005.05.005

Kulkarni, R.R., Pawar, P.V., Joseph, M.P., Akulwad, A.K., Sen, A., Joshi, S.P., 2013. *Lavandula gibsoni* and *Plectranthus mollis* essential oils: chemical analysis and insect control activities against *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus*. J. Pest Sci. 86, 713–718. https://doi.org/10.1007/s10340-013-0502-1

Kwon, Y., Kim, S.H., Ronderos, D.S., Lee, Y., Akitake, B., Woodward, O.M., Guggino, W.B., Smith, D.P., Montell, C., 2010. *Drosophila* TRPA1 channel is required to avoid the naturally

occurring insect repellent citronellal. Current biology 20, 1672–1678.
https://doi.org/10.1016/j.cub.2010.08.016

Lahondère, C., Vinauger, C., Okubo, R.P., Wolff, G.H., Chan, J.K., Akbari, O.S., Riffell, J.A.,
2019. The olfactory basis of orchid pollination by mosquitoes. Proceedings of the National
Academy of Sciences USA. https://doi.org/10.1073/pnas.1910589117

Liang, Y., Li, J.L., Xu, S., Zhao, N.N., Zhou, L., Cheng, J., Liu, Z.L., 2013. Evaluation of
repellency of some chinese medicinal herbs essential oils against *Liposcelis bostrychophila*
(Psocoptera: Liposcelidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae). J Econ
Entomol 106, 513–519. https://doi.org/10.1603/ec12247

Liu, F., Wang, Q., Xu, P., Andreazza, F., Valbon, W.R., Bandason, E., Chen, M., Yan, R., Feng,
B., Smith, L.B., Scott, J.G., Takamatsu, G., Ihara, M., Matsuda, K., Klimavicz, J., Coats, J.,
Oliveira, E.E., Du, Y., Dong, K., 2021. A dual-target molecular mechanism of pyrethrum
repellency against mosquitoes. Nat Commun 12, 2553. https://doi.org/10.1038/s41467-021-
22847-0

Lu, T., Qiu, Y.T., Wang, G., Kwon, J.Y., Rutzler, M., Kwon, H.-W., Pitts, R.J., Loon, J.J.A.V.,
Takken, W., Carlson, J.R., Zwiebel, L.J., 2007. Odor coding in the maxillary palp of the
malaria vector mosquito *Anopheles gambiae*. Current biology 17, 1533–1544.
https://doi.org/10.1016/j.cub.2007.07.062

Lukhoba, C.W., Simmonds, M.S.J., Paton, A.J., 2006. *Plectranthus*: A review of ethnobotanical
uses. J. Ethnopharmacol. 103, 1–24. https://doi.org/10.1016/j.jep.2005.09.011

Maia, M.F., Moore, S.J., 2011. Plant-based insect repellents: a review of their efficacy,
development and testing. Malaria journal 10 Suppl 1, S11. https://doi.org/10.1186/1475-2875-
10-s1-s11

Majeed, S., Hill, S.R., Birgersson, G., Ignell, R., 2016. Detection and perception of generic host
volatiles by mosquitoes modulate host preference: context dependence of (*R*)-1-octen-3-ol.
Royal Society open science 3, 160467. https://doi.org/10.1098/rsos.160467

Majeed, S., Hill, S.R., Dekker, T., Ignell, R., 2017. Detection and perception of generic host
volatiles by mosquitoes: responses to CO2 constrains host-seeking behaviour. Royal Society
open science 4, 170189. https://doi.org/10.1098/rsos.170189

Matthews, B.J., Dudchenko, O., Kingan, S.B., Koren, S., Antoshechkin, I., Crawford, J.E.,
Glassford, W.J., Herre, M., Redmond, S.N., Rose, N.H., Weedall, G.D., Wu, Y., Batra, S.S.,
Brito-Sierra, C.A., Buckingham, S.D., Campbell, C.L., Chan, S., Cox, E., Evans, B.R.,
Fansiri, T., Filipović, I., Fontaine, A., Gloria-Soria, A., Hall, R., Joardar, V.S., Jones, A.K.,
Kay, R.G.G., Kodali, V.K., Lee, J., Lycett, G.J., Mitchell, S.N., Muehling, J., Murphy, M.R.,
Omer, A.D., Partridge, F.A., Peluso, P., Aiden, A.P., Ramasamy, V., Rašić, G., Roy, S.,
Saavedra-Rodriguez, K., Sharan, S., Sharma, A., Smith, M.L., Turner, J., Weakley, A.M.,
Zhao, Z., Akbari, O.S., Black, W.C., Cao, H., Darby, A.C., Hill, C.A., Johnston, J.S.,

Murphy, T.D., Raikhel, A.S., Sattelle, D.B., Sharakhov, I.V., White, B.J., Zhao, L., Aiden, E.L., Mann, R.S., Lambrechts, L., Powell, J.R., Sharakhova, M.V., Tu, Z., Robertson, H.M., McBride, C.S., Hastie, A.R., Korlach, J., Neafsey, D.E., Phillippy, A.M., Vosshall, L.B., 2018. Improved reference genome of *Aedes aegypti* informs arbovirus vector control. Nature 563, 501–507. https://doi.org/10.1038/s41586-018-0692-z

McMeniman, C.J., Corfas, R.A., Matthews, B.J., Ritchie, S.A., Vosshall, L.B., 2014. Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. Cell 156, 1060–1071. https://doi.org/10.1016/j.cell.2013.12.044

Mei, Y., Li, L., Fan, L., Fan, W., Liu, L., Zhang, F., Hu, Z., Wang, K., Yang, L., Wang, Z., 2023. The history, stereochemistry, ethnopharmacology and quality assessment of borneol. J Ethnopharmacol 300, 115697. https://doi.org/10.1016/j.jep.2022.115697

Melo, N., Capek, M., Arenas, O.M., Afify, A., Yilmaz, A., Potter, C.J., Laminette, P.J., Para, A., Gallio, M., Stensmyr, M.C., 2021. The irritant receptor TRPA1 mediates the mosquito repellent effect of catnip. Curr Biol. https://doi.org/10.1016/j.cub.2021.02.010

Melo, N., Wolff, G.H., Costa-da-Silva, A.L., Arribas, R., Triana, M.F., Gugger, M., Riffell, J.A., Degennaro, M., Stensmyr, M.C., 2020. Geosmin attracts *Aedes aegypti* mosquitoes to oviposition sites. Current Biology 30, 1–8. https://doi.org/10.1101/598698

Moore, S.J., Debboun, M., 2007. History of Insect Repellents. Chapter 1. In Insect Repellents: Principles, Methods, and Uses. Eds: Debboun, M., Frances, S.P., Strickman, D. CRC Press.

Ng, R., Wu, S., Su, C., 2020. Neuronal compartmentalization: a means to integrate sensory input at the earliest stage of information processing? Bioessays 42, e2000026. https://doi.org/10.1002/bies.202000026

Nunes, R.K.V., Martins, U.N., Brito, T.B., Nepel, A., Costa, E.V., Barison, A., Santos, R.L.C., Cavalcanti, S.C.H., 2018. Evaluation of (–)-borneol derivatives against the Zika vector, *Aedes aegypti* and a non-target species, *Artemia* sp. Environ Sci Pollut R 25, 31165–31174. https://doi.org/10.1007/s11356-018-2809-1

Pålsson, K., Jaenson, T.G.T., 1999. Plant products used as mosquito repellents in Guinea Bissau, West Africa. Acta Trop 72, 39–52. https://doi.org/10.1016/s0001-706x(98)00083-7

Picollo, M.I., Toloza, A.C., Cueto, G.M., Zygadlo, J., Zerba, E., 2008. Anticholinesterase and pediculicidal activities of monoterpenoids. Fitoterapia 79, 271–278. https://doi.org/10.1016/j.fitote.2008.01.005

Pitarokili, D., Michaelakis, A., Koliopoulos, G., Giatropoulos, A., Tzakou, O., 2011. Chemical composition, larvicidal evaluation, and adult repellency of endemic Greek *Thymus* essential oils against the mosquito vector of West Nile virus. Parasitol Res 109, 425–430. https://doi.org/10.1007/s00436-011-2271-1

Pratiwi, M.A.M., Purwati, 2021. The repellent activity test of rosemary leaf (*Rosmarinus*
*officinalis*) essential oil gel preparations influence on *Aedes aegypti* mosquito. J Phys Conf
Ser 1788, 012016. https://doi.org/10.1088/1742-6596/1788/1/012016

Raj, G.A., Chandrasekaran, M., Krishnamoorthy, S., Jayaraman, M., Venkatesalu, V., 2015.
Phytochemical profile and larvicidal properties of seed essential oil from *Nigella sativa* L.
(Ranunculaceae), against *Aedes aegypti*, *Anopheles stephensi*, and *Culex quinquefasciatus*
(Diptera: Culicidae). Parasitol Res 114, 3385–3391. https://doi.org/10.1007/s00436-015-
4563-3

Rajkumar, S., Jebanesan, A., 2010. Chemical composition and larvicidal activity of leaf essential
oil from *Clausena dentata* (Willd) M. Roam. (Rutaceae) against the chikungunya vector,
*Aedes aegypti* Linn. (Diptera: Culicidae). J Asia-pac Entomol 13, 107–109.
https://doi.org/10.1016/j.aspen.2010.02.001

Santos, S.R.L., Silva, V.B., Melo, M.A., Barbosa, J.D.F., Santos, R.L.C., Sousa, D.P. de,
Cavalcanti, S.C.H., 2010. Toxic effects on and structure-toxicity relationships of
phenylpropanoids, terpenes, and related compounds in *Aedes aegypti* larvae. Vector-Borne
Zoonotic Dis. 10, 1049–1054. https://doi.org/10.1089/vbz.2009.0158

Shankar, S., McMeniman, C.J., 2020. An updated antennal lobe atlas for the yellow fever
mosquito *Aedes aegypti*. Plos Neglect Trop D 14, e0008729.
https://doi.org/10.1371/journal.pntd.0008729

Stensmyr, M.C., Dweck, H.K.M., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe,
V., Steck, K., Lavista-Llanos, S., Wicher, D., Sachse, S., Knaden, M., Becher, P.G., Seki, Y.,
Hansson, B.S., 2012. A conserved dedicated olfactory circuit for detecting harmful microbes
in *Drosophila*. Cell 151, 1345–1357. https://doi.org/10.1016/j.cell.2012.09.046

Su, C.-Y., Menuz, K., Reisert, J., Carlson, J.R., 2012. Non-synaptic inhibition between grouped
neurons in an olfactory circuit. Nature 492, 66–71. https://doi.org/10.1038/nature11712

Takken, W., Kline, D.L., 1989. Carbon dioxide and 1-octen-3-ol as mosquito attractants. Journal
of the American Mosquito Control Association 5, 311–316.

Team, R.C., 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Vinauger, C., Breugel, F. van, Locke, L.T., Tobin, K.K.S., Dickinson, M.H., Fairhall, A.L.,
Akbari, O.S., Riffell, J.A., 2019. Visual-olfactory integration in the human disease vector
mosquito *Aedes aegypti*. Current biology 29, 2509-2516.e5.
https://doi.org/10.1016/j.cub.2019.06.043

Wadley, L., Sievers, C., Bamford, M., Goldberg, P., Berna, F., Miller, C., 2011. Middle stone
age bedding construction and settlement patterns at Sibudu, South Africa. Science 334, 1388–
1391. https://doi.org/10.1126/science.1213317

Waliwitiya, R., Kennedy, C.J., Lowenberger, C.A., 2009. Larvicidal and oviposition⬜altering
activity of monoterpenoids, trans⬜anithole and rosemary oil to the yellow fever mosquito
*Aedes aegypti* (Diptera: Culicidae). Pest Manag Sci 65, 241–248.
https://doi.org/10.1002/ps.1675

Wang, G., Carey, A.F., Carlson, J.R., Zwiebel, L.J., 2010. Molecular basis of odor coding in the
malaria vector mosquito *Anopheles gambiae*. PNAS 107.
https://doi.org/10.1073/pnas.0913392107

Wang, Y., Chen, Q., Guo, J., Li, J., Wang, J., Wen, M., Zhao, H., Ren, B., 2017. Molecular basis
of peripheral olfactory sensing during oviposition in the behavior of the parasitic wasp
*Anastatus japonicus*. Insect Biochem. Mol. Biol. 89, 58–70.
https://doi.org/10.1016/j.ibmb.2017.09.001

Wolff, G.H., Lahondère, C., Vinauger, C., Rylance, E., Riffell, J.A., 2023. Neuromodulation and
differential learning across mosquito species. Proc. R. Soc. B 290, 20222118.
https://doi.org/10.1098/rspb.2022.2118

Xu, P., Zeng, F., Bedoukian, R.H., Leal, W.S., 2019. DEET and other repellents are inhibitors of
mosquito odorant receptors for oviposition attractants. Insect Biochemistry and Molecular
Biology 113, 103224. https://doi.org/10.1016/j.ibmb.2019.103224

Zeng, F., Xu, P., Tan, K., Zarbin, P.H.G., Leal, W.S., 2018. Methyl dihydrojasmonate and lilial
are the constituents with an "off-label" insect repellence in perfumes. Plos One 13, e0199386.
https://doi.org/10.1371/journal.pone.0199386

Zhang, N., Tang, L., Hu, W., Wang, K., Zhou, Y., Li, H., Huang, C., Chun, J., Zhang, Z., 2014.
Insecticidal, fumigant, and repellent activities of sweet wormwood oil and its individual
components against red imported fire ant workers (Hymenoptera: Formicidae). J Insect Sci
14, 241. https://doi.org/10.1093/jisesa/ieu103

Zhao, Z., Zung, J.L., Hinze, A., Kriete, A.L., Iqbal, A., Younger, M.A., Matthews, B.J., Merhof,
D., Thiberge, S., Ignell, R., Strauch, M., McBride, C.S., 2022. Mosquito brains encode unique
features of human odour to drive host seeking. Nature 1–7. https://doi.org/10.1038/s41586-
022-04675-4

Zhou, X., Rinker, D.C., Pitts, R.J., Rokas, A., Zwiebel, L.J., 2014. Divergent and conserved
elements comprise the chemoreceptive repertoire of the non-blood feeding mosquito
*Toxorhynchites amboinensis*. Genome biology and evolution 6, 2883–2896.
https://doi.org/10.1093/gbe/evu231

Zhu, J.J., Cermak, S.C., Kenar, J.A., Brewer, G., Haynes, K.F., Boxler, D., Baker, P.D., Wang,
D., Wang, C., Li, A.Y., Xue, R., Shen, Y., Wang, F., Agramonte, N.M., Bernier, U.R., Filho,
J.G. de O., Borges, L.M.F., Friesen, K., Taylor, D.B., 2018. Better than DEET repellent
compounds derived from coconut oil. Sci. Rep. 8, 14053. https://doi.org/10.1038/s41598-018-
32373-7

1110

A. 900 nm laser / Odor delivery

B. MD3 (green), MD2 (orange), MD1 (blue), AD1. Scale bar: 20 µm

C. ΔF/F₀ color scale: 0.0, 0.2, 0.6. brp-QF2>QUAS-GCaMP6s

D. Borneol — MD3, MD2, MD1. 5 sec. ΔF/F₀

E. 1-octen-3-ol. ΔF/F₀

F. CO₂. ΔF/F₀

G. ΔF/F₀ — CO₂, (+)-Borneol *, (+)-Camphor *, (−)-Borneol **, (−)-Camphor *, Geosmin, 1-Octen-3-ol, Control

A

a

b

Detection number (%)

Control (DEE)  (±)-Borneol

B

Control (DEE)

0 1 2 3 4 5 6 7 8 9 10

No. of mosquitoes

(±)-Borneol

0 1 2 3 4 5 6 7 8 9 10
Time (min)

ROI