How do plant RNA viruses overcome the negative effect of Muller's ratchet despite strong transmission bottlenecks?

Guillaume Lafforgue[1], Marie Lefebvre[1], Thierry Michon[1] and Santiago F. Elena[2,3]

[1]*UMR Biologie du Fruit et Pathologie, INRAE, Université de Bordeaux, Villenave d'Ornon, France.*
[2]*Instituto de Biología Integrativa de Sistemas (I²SysBio), Consejo Superior de Investigaciones Científicas – Universitat de València, Paterna, 46980 València, Spain.* [3]*The Santa Fe Institute, Santa Fe, NM 87501, USA.*

ABSTRACT

Muller's ratchet refers to the irreversible accumulation of deleterious mutations in small populations, resulting in a decline in overall fitness. This phenomenon has been extensively observed in experiments involving microorganisms, including bacteriophages and yeast. While the impact of Muller's ratchet on viruses has been largely studied in bacteriophages and animal RNA viruses, its effects on plant RNA viruses remain poorly documented. Plant RNA viruses give rise to large and diverse populations that undergo significant bottlenecks during the colonization of distant tissues or through vector-mediated horizontal transmission. In this study, we aim to investigate the role of bottleneck size, the maximum population size between consecutive bottlenecks, and the generation of genetic diversity in countering the effects of Muller's ratchet. We observed three distinct evolutionary outcomes for tobacco etch virus under three different demographic conditions: (*i*) a decline in fitness following periodic severe bottlenecks in *Chenopodium quinoa*, (*ii*) a consistent fitness level with moderate bottlenecks in *C. quinoa*, and (*iii*) a net increase in fitness when severe bottlenecks in *C. quinoa* were alternated with large population expansions in *Nicotiana tabacum*. By fitting empirical data to an *in silico* simulation model, we found that initiating a lesion in *C. quinoa* required only 1-5 virions, and approximately 40 new virions were produced per lesion. These findings demonstrate that Muller's ratchet can be halted not only by increasing the number of founder viruses but also by incorporating phases of exponential growth in large populations between bottlenecks. Such population expansions generate genetic diversity, serving as a buffer against, and potentially even leveraging, the effects of genetic drift.

Key words: experimental evolution, Muller's ratchet, plant virus, potyvirus, RNA virus, simulations, transmission bottleneck, virus evolution.

INTRODUCTION

Viruses have very heterogeneous genomic structures and replication strategies. Among viruses, a defining feature of RNA viruses is their high mutation rate, with estimates ranging from 10^{-4} to 10^{-6} mutations per base per replication round (1 - 4). During the process of evolution, natural selection can only operate on the diversity present within a population. However, beside selection, random sampling of individuals may occur during transmission events, leading to a reduction in genetic variability, which influences the evolution of a viral population (5). The strength of the genetic drift depends on the effective population size (N_e, *i.e.*, size and diversity). Small N_e boosts the strength of genetic drift, magnifying stochastic effects throughout evolution (6). If virus infections are frequently initiated by a small number of virions (*i.e.*, small N_e), the resulting genetic bottlenecks may have a major impact on disease progression in individual hosts as well as in the evolution of virus population after multiple rounds of host infection. Furthermore, the frequency of mixed-genotype coinfections, including those by multipartite viruses, is determined by the number of infecting virions. Mixed-genotype infections can diminish pathogenesis-inducing interactions between viral genotypes while simultaneously decreasing evolutionary-relevant interactions such as recombination between distinct virus genotypes or segment segregation in segmented viruses. Although these effects are straightforward in and of themselves, their evolutionary consequences may not always be obvious (7).

The characteristics of N_e are likely to affect the genetic consequences of transmissions associated with repeated bottlenecks (7). Hence, starting from a parental population in which substantial genetic diversity may exist, the sampling of a small number of individuals can lead to a profoundly different genetic composition in the resulting population (8, 9). If this founder effect is strong and repeated, populations can diverge rapidly and bottlenecks tend to facilitate the random fixation of variants in populations regardless their selective value (10, 11). Repeated sampling of a small number of founders, combined with high mutation rates, can cause the serial fixation of deleterious mutations and a concomitant decrease of the mean fitness, the so-called Muller's ratchet effect (12). In nature, N_e widely varies for viruses as a function of inoculum dose and host susceptibility (13, 14). In the case of plant viruses, infection often begins with a few individuals on which the bottleneck effect is exerted during horizontal (15 - 19) or vertical transmission (19). The mean number of founders of infection can under some circumstances approach one for plant viruses (16, 20 - 23). High mutation rate may constitute a disadvantage for RNA virus survival, as most mutations shall be deleterious (24). Muller's ratchet has been demonstrated with bacteriophage Φ6 (25), vesicular stomatitis virus (26), foot-and-mouth disease virus (27), human immunodeficiency virus type 1 (28), and tobacco etch virus (TEV) (29), as well as for bacteria (30) and yeast (31). In all of these studies, in the absence of sexual reproduction, genetic mechanisms such as recombination cannot compensate for the accumulation of deleterious mutations (32, 33). In very large populations, fitness recovery can also occur through

fixation of compensatory or reversion mutations (34, 35). However, when N_e is small, the fitness loss due to Muller's ratchet is directly correlated to the fitness of the ancestral population (36).

So far, all experimental studies of the Muller's ratchet effect used protocols involving passages of viruses into cultured cells or unicellular organisms. These experimental conditions were therefore a poor representation of the complexity of natural hosts and possible vectors (37, 38): multicellular organisms display structures made of various differentiated tissues unevenly distributed in space and submitted to physical stress and possibly to immune responses, all these factors contributing to shape the genetic structure of infecting viral populations (39). Among the organisms used for experimental verification of Muller's ratchet effect, plant RNA viruses have been poorly represented.

TEV (species *Tobacco etch virus*, genus *Potyvirus*, family *Potyviridae*) possesses a positive-sense single-stranded RNA genome. The virus induces local lesions on *Chenopodium quinoa* Willd and a systemic infection in its natural host *Nicotiana tabacum* L. It has been reported that repeated transfers of local lesions generated by TEV in *C. quinoa* leads to a significant reduction of the Malthusian growth rate (29), a measure of absolute fitness. Because the virus is not moving systemically within the plant, local lesions can be used to separate a small subpopulation of viruses from the original pool (40); another characteristic of this model is a putative moderate virus multiplication rate due to the plant hypersensitive response that results in the necrotic lesions (41). Factors governing evolution in complex multicellular hosts systems are multiple. However, our experimental system allows dissecting the influence of two demographic parameters: the strength of individual bottlenecks and the amount of replication events between two consecutive bottlenecks, in a complex multi-cellular host plant. In this study, capitalizing on a previous study using the TEV/*C. quinoa* pathosystem (29), we investigated the impact of these two demographic parameters on this potyvirus infectivity in its natural host (*N. tabacum*), and how these factors impact the recovery or decrease of infectivity in the viral progeny.

MATERIALS AND METHODS

In vitro **RNA transcription and plant inoculation**

The pTEV-7DA infectious clone, provided by Prof. James C. Carrington (Donald Danforth Plant Sciences Center, St. Louis MO, USA), was the source for the ancestral virus for all experiments (37). Infectious plasmids were linearized with *Bgl*II (Fermentas, Vilnius, Lithuania) and transcribed into 5'-capped RNAs using SP6 mMessage mMachine® Kit (Ambion Inc., Austin TX, USA). Transcripts were precipitated (1.5 volumes of DEPC-treated water, 1.5 volumes of 7.5 M LiCl, 50 mM EDTA) and resuspended in DEPC-treated water. RNA integrity was assessed by gel electrophoresis and its concentration quantified spectrophotometrically with a NanoDrop (Thermo Fisher Scientific,

Waltham MA, USA). The obtained RNA was used to inoculate groups of 4-week-old *N. tabacum* cv. *Xanthi* plants by means of abrasion of the third true leaf with 4 μg of RNA (Fig. 1). Plants were maintained in the green house at 25 °C and 16 h light for one week. Symptoms appeared 5 - 6 days post-inoculation (dpi).

Fig. 1. Schematic representation of the serial transfer of a TEV population derived from an initial clone. At each transfer, either a random lesion (a, strong bottleneck, 'smallest N_e' treatment) or twenty lesions (b, weak bottleneck, 'intermediate N_e' treatment) were used to continue serial passages. In the third 'large N_e' treatment (c) one lesion was used to initiate infection in *N. tabacum* and then transferred on *C. quinoa*. After each transfer, the population fitness was measured on *N. tabacum*. Each evolutionary treatment was replicated 10 times.

Inoculation of *C. quinoa* and generation of mutation-accumulation lineages

An initial sap was prepared with 1 g of infected tissue (from *N. tabacum* plants 7 dpi) ground in 1.5 mL of phosphate buffer (50 mM phosphate, 3% polyethylene glycol 8000). Four different leaves from five different 4-week-old *C. quinoa* plants were rub-inoculated with 10 μL of sap. Lesions on inoculated leaves were harvested at 10 dpi. A well-isolated lesion was perforated from the leaf using the wide end of a 5 μL micropipette tip (approximately 4 mm diameter disc) and ground in a mortar with 24 μL of phosphate buffer. The extract was carefully collected, centrifuged for 1 min at 12,000 g and the supernatant was used for inoculation.

The experimental manipulation of the bottleneck size was done using two different modalities (Fig. 1). The 'smallest N_e' treatment was used to apply a strong bottleneck. Briefly, 10 lesions were

processed separately to create ten independent single lesion lineages: 5 µL of each lesion sap used to

inoculate four different leaves of *C. quinoa* to generate the next generation of lesions. After 10 dpi,

one lesion was used to generate the next generation. To generate a weaker bottleneck effect, the

'intermediate N_e' treatment was used. Ten independent pools of 20 lesions were grounded in 240 µL

of extraction buffer producing ten independent group lesion lineages. The same procedure was used

as for single lesion lineages (*i.e.*, 10 µL of lesion sap were used to inoculate ten *C. quinoa* leaves).

To introduce a larger number of generations between consecutive bottlenecks, the 'large N_e'

treatment was used. Briefly, in between passages in *C quinoa*, the progeny was sequentially inoculated

in TEV natural host, *N. tabacum*. After 10 dpi, a *N. tabacum* infected plant was grounded in phosphate

buffer (0.5 g in 500 µL) and used to inoculate ten *C. quinoa* leaves (10 µL).

It has been previously reported that single lesion lineages tend to have a significant loss of

absolute fitness until, in some cases, the lineages extinct (29). After five passages, the experimental

study ended with a loss of four lineages and a production of less than five lesions.

For all experimental treatments, lesions sampled at each passage were stored at −80 ºC to

perform all infectivity test later into a single block. No effect of storage at −80 °C on single lesion

infectivity was observed (data not shown).

Infectivity measurement

The fitness of each evolved lineage within each history (small, intermediate or large N_e treatments),

was assessed by infectivity measurement at each time point (*i.e.*, the end of a given passage). The

analysis of infectivity was performed on *N. tabacum*. From a given lesion on *C. quinoa*, only two

tobacco plants can be inoculated. Consequently, to be statistically relevant, for each passage, up to 10

single lesions were randomly selected for each lineage. The lesions were grounded in 24 µL of

phosphate buffer, and 10 µL were used to inoculate the third true leaf of two *N. tabacum* plants, for

each lesion. Presence of symptoms was recorded at 9 dpi. Hence, per history (small, intermediate or

large N_e treatments), a total of 200 tobacco plants were inoculated per passage. Infectivity was

expressed as the frequency of infected symptomatic plants out of the number of inoculated ones.

Statistical analyses

The linear regression of infectivity in *N. tabacum versus* passage number, fixing the origin at

the 0.261 ±0.035 (±1 SEM) infectivity value measured at the first passage (see below), returns a slope

that represents an estimate of the rate of infectivity evolution. These rates of evolution were determined

for each lineage of the small, intermediate and large N_e. A one-way robust Welch ANOVA was used

to evaluate the effect of demographic conditions in the rates of infectivity evolution. The magnitude

of the effect was evaluated using the η^2 statistic; usually $\eta^2 > 0.15$ are considered as large effects.

Variance components were estimated by maximum-likelihood. These statistical tests were done with
SPSS version 28.0.1.1 (IBM Corp., Armonk NY, USA).

An *in silico* model of the growth processes taking place within single lesions

To evaluate the demographic processes occurring within a local lesion during virus
multiplication, and to mimic the mechanisms acting for the small or intermediate N_e treatments, we
implemented a simulation model in R version 3.6.1 (https://doi.org/10.5281/zenodo.8199726).
Briefly, the experiment was simulated in three steps: (*i*) generation of lesions in *C. quinoa* from
infected *N tabacum*, (*ii*) multiplication within a random lesion and (*iii*) generation of lesions either
from a single (small N_e treatment) or from 20 digital lesions (intermediate N_e treatment). Steps (*ii*) and
(*iii*) were repeated four times with 200 replicates for each model.

These models take into account the following postulates: (*i*) a few number of particles shall be
enough to initiate a lesion (20), (*ii*) two types of viral particles are considered: type A virions which
can reproduce either in *C. quinoa* and *N. tabacum* and type D virions which can reproduce only in *C.*
quinoa but had lost their capacity to infect the natural host, and (*iii*) the percentage of virions A in a
lesion determines the probability of infection in *N. tabacum* defined as the infectivity rate:

$$I := \frac{Number\ of\ type\ A\ virions\ in\ a\ lesion}{Total\ number\ of\ virions\ in\ the\ lesion}. \tag{1}$$

Result for each passage is given by the mean of *I* for 12 random lesions.

The parameters of the model are the number of virions able to initiate a lesion $V \in (V_{min}, V_{max}\ |$
$V_{min} < V_{max})$ that take integer values in the range [0, 8] and the rates of replication in *C. quinoa* of type
A virons $r^A \in (r^A_{min}, r^A_{max}\ |\ r^A_{min} < r^A_{max})$ and D $r^D \in (r^D_{min}, r^D_{max}\ |\ r^D_{min} < r^D_{max})$ that take integer values
in the interval [1, 50].

The small N_e model was implemented as follows. First, a random number of founders *V* is
determined. Then, among these founders, the number of virions of types A and D are derived from
equation (1), *I* being given as the infectivity measured experimentally at the first passage (*i.e.*, 0.261
±0.035; see below). Replication rates in *C. quinoa* r^A and r^D are used to calculate the next pool of
virions (*G* + 1) able to generate the next lesion. Twelve simulated lesions are generated. A random
number in the interval (V_{min}, V_{max}) of founder particles is attributed to each lesion. From the *G* + 1
pool, the subpopulations of particles types A and D within each lesion was randomly assigned. For
this model, a single lesion was randomly picked to repeat the process until the fourth passage. An
exhaustive search in the parameter space values (V_{min}, V_{max}, r^A_{min}, r^A_{max}, r^D_{min}, r^D_{max}) was done, resulting
in $5.8 \cdot 10^7$ different combinations in the case of the small N_e model. For each of these simulations, 200
replicate runs were performed. Combinations of parameters resulting in estimates of infectivity and
intra-lineages genetic component of variance within the observed experimental 95% confidence

intervals were retained for further analyses. Furthermore, simulated rates of infectivity evolution for

these parameter combinations were obtained and contrasted against the 95% confidence intervals

empirically estimated. This step results in $1.8 \cdot 10^7$ combinations corresponding to measured values in

the small N_e treatment.

The intermediate Ne model uses the same canvas as for the small Ne model described above and

incorporates the $1.8 \cdot 107$ combinations previously retained. The difference resides in the generation of

20 lesions subsequently pooled together, the resulting mix being used to generate the $G + 1$.

Due to the objective of the simulations, which is to monitor the parameters associated with the

virus' life cycle within a lesion, and considering the significant increase in the number of additional

parameters involved in the large N_e treatment (such as mutation rate and multiplication within

tobacco), it was not considered in the analysis.

RESULTS

Critical infectivity can be restored

In all experiments, fitness was considered to be directly related to the capacity of the virus to

initiate disease in *N. tabacum*. Inoculation of *N. tabacum* with sap from infected tissue of *N. tabacum*

results in nearly 100% infectivity (data not shown). After a single passage of TEV in *C. quinoa*,

infectivity in *N. tabacum* was reduced to 0.261 ±0.035. This resulting virus population was then used

for subsequent evolution experiments.

Serial passages of single lesions (small N_e treatment) produced in *C. quinoa* resulted in a further

decrease in infectivity in *N. tabacum* (Fig. 2A). Indeed, the average infectivity among lineages after

five passages was 0.174 ±0.078, representing an average net reduction of 33.5% relative to the first

passage. After removing four outlier values, the average rate of infectivity evolution among these

lineages was -0.033 ± 0.005 per passage, which was found to be significantly negative ($t_{17} = -7.390$,

$P < 0.001$). Without removing outliers, the rate of infectivity evolution was -0.018 ± 0.009 per passage,

indicated that it was significantly negative ($t_{21} = -2.034$, one-tailed $P = 0.027$), showing a large

variation between lineages especially for this treatment.

When passaging was performed with pools of 20 local lesions (intermediate N_e treatment; Fig.

2B), the final average infectivity among the 20 lineages was 0.200 ±0.076, which represents a further

reduction of 21.6% in infectivity relative to the first passage. For this treatment, the rate of infectivity

evolution was -0.017 ± 0.006 per passage, a value that is also significantly negative ($t_{18} = -2.654$, $P =$

0.008). Finally, when passages included a population expansion in *N. tabacum* and the bottleneck in

C. quinoa (large N_e treatment), a net increase of infectivity over passages was observed (Fig. 2C), with

an average final infectivity of 0.447 ±0.038 and an average rate of infectivity evolution estimated to

be 0.037 ±0.005 per passage, a value significantly positive ($t_9 = 7.406$, $P < 0.001$).

Significant differences among the three treatments exist in the rates of infectivity evolution (Fig.

3D: $F_{2,31.281} = 28.122$, $P < 0.001$), being them of very large magnitude ($\eta^2 = 0.321$). However, these

differences were mostly driven by the differences between the large N_e treatment and the other two

(Bonferroni *post hoc* test: $P < 0.001$ in both comparison), but not among the small and intermediate

N_e treatments (Bonferroni *post hoc* test: $P = 1.000$).

Fig. 2. (A) Evolution of average infectivity among independent lineages at each bottleneck event (passage) for small $N_e = 1$ (lesion-to-lesion), (B) intermediate $N_e = 20$ (20 lesions-to-20 lesions) and (C) large N_e (population expansion in *N. tabacum*) experimental treatments. (D) Box plot for the estimated rates of infectivity evolution. Error bars represent ±1 SEM.

Variance intra- and inter-lineages

For a given demographic setup, the lineage fitness can vary in *N. tabacum* due to several factors

(adaptation, chance, contingency…). To evaluate the contribution of true genetic differences to the

evolution of infectivity, we decomposed the total observed variance for lineages evolved in the same

conditions into their additive (among lineages) and random (within lineages) components. Fig. 3A

shows the evolution of the additive (i.e., genetic) standard deviation for the three experimental

treatments. In all three treatments, genetic differences among lineages increased after the first passage,
inversely proportional to Ne: random fixation of alleles associated with strong bottlenecks contributes
to inflating the genetic component of variance. However, throughout the experiment, subsequent
strong bottlenecks resulted in a loss of genetic differences, stabilizing the variance.

Interestingly, at the beginning of the experiment, the intra-lineage variation among the lineages
with the smallest Ne appeared to be higher compared to the lineages derived from the other two
modalities. However, after three passages, the inter-lineage variance of the smallest Ne lineages
decreased significantly and reached a similar level as the inter-lineage variance observed in the
intermediate and large Ne lineages. This suggests a convergence in terms of genetic diversity among
the different lineages, regardless of their initial Ne values.

Fig. 3. Evolution of the genetic standard deviation among (A) or within (B) lineages along the passages.

The analysis of lineages with varying N_e revealed interesting trends in intra-lineage variation
over the course of passages. Specifically, the lineages with the smallest N_e exhibited a notable
decreasing trend in intra-lineage variation across passages (Fig. 3B). This finding suggests that there
was a reduction in genetic diversity within those lineages over time as they were serially passaged. On
the other hand, the intermediate and large N_e lineages did not display any significant changes in intra-
lineage variation throughout the passages. The lack of significant variation in these lineages suggests
that their genetic diversity remained relatively stable over time. This could be attributed to their larger
N_e, which may have provided a buffer against genetic drift and enabled the preservation of diverse

genetic traits within these lineages.

In the small N_e treatment, infectivity is independent from multiplication in *C. quinoa*

The observed decrease in infectivity in *N. tabacum* after serial passages of single lesions from

C. quinoa (small N_e treatment) may be attributed to factors related to the quantity and infectivity of

virions present in the inoculum derived from *C. quinoa* lesions.

It has been established that the number of lesions in C. quinoa is closely associated with the viral

load (29), which refers to the quantity of viral particles present within the infected host. Therefore, in

the context of the small N_e treatment, the decrease in infectivity observed in *N. tabacum* could be

linked to a lower quantity or even an absence of virions in the inoculum derived from *C. quinoa*

lesions, which are capable of infecting *N. tabacum*. However, upon further investigation, no significant

correlation was found between the number of lesions and the infectivity in *N. tabacum* (Spearman's

rank correlation coefficient $r_S = -0.017$, 118 d.f., $P = 0.854$). This finding indicates that the infectivity

rate in *N. tabacum* is influenced more by the genetic characteristics of the virions rather than the

concentration of the viral load (Supplementary Fig. S1). In other words, the genetic properties of the

viral particles present in the inoculum may play a more critical role in determining the infectivity in

N. tabacum than the sheer quantity of the viral particles.

Evaluation of growth parameters for individual lesions

From the $4.2 \cdot 10^7$ different combinations of parameters tested, $1.74 \cdot 10^3$ (*i.e.*, 0.04%) generated

mean infectivity values and non-additive standard deviations within lineages (*i.e.*, true noise) lying

within the observed 95% confidence intervals, thus being compatible with the experimental

observations for the small and intermediate N_e treatments. Fig. 4 summarizes the results of the

exhaustive search of space parameters compatible with the observed results. The modes of these six

distributions can be taken as the most representative values for the model parameters: $V_{min} = 1$, $V_{max} =$

4, $r_{min}^A = 11$, $r_{max}^A = 37$, $r_{min}^D = 21$, and $r_{max}^D >= 50$. These parameters show that very few particles are

enough to generate a necrotic lesion in *C. quinoa* and that the D mutant types show higher replication

rates in *C. quinoa* than the A types, with a replicative advantage that varies between 1.35- and 1.91-

fold. To qualitatively confirm the agreement between observed data and the estimated parameter

values, we run 20 independent simulations with the parameters above (Fig. 5).

The simulation results shown in Fig. 5A are in good agreement with the observed dynamics

shown in Fig. 2A and Fig. 3A: an overall trend of infectivity to decline along passages and a strong

decrease in intra-lineage variance, respectively. However, some differences exist between

experimental and simulation results. Firstly, most of the simulated lineages lost their infectivity after

the third passage while the majority of experimental lineages still were still infectious in *N. tabacum*

after the fifth passage. Secondly, some simulated lineages reached 100% infectivity in *N. tabacum*, something that was not observed in our experiments. Regarding the simulations for the intermediate N_e treatment (Fig. 5B), the 20 simulated lineages show a minor reduction in infectivity, also in good agreement with the data shown in Fig. 2B. In this case, a qualitative difference exists between observed and simulated data. In this case, some experimental lineages reached 0% infectivity in *N. tabacum*, while the simulations never show this lack of infectivity.

Fig. 4. Distribution of parameters (V_{min}, V_{max}, r_{min}^{A}, r_{max}^{A}, r_{min}^{D}, r_{max}^{D}) compatible with the experimental observations made for small and intermediate N_e demographic regimes.

Fig. 5. Twenty simulated lineages with the most common set of parameters. The infectivity and within-lineage error are plotted for (A) small N_e and (B) intermediate N_e treatments. Red lines represent the mean of the 20 replicates.

DISCUSSION

Although Muller's ratchet has been studied for several microorganisms, including viruses, so far, it

has been experimentally addressed only for one plant virus (29). Bottlenecks such as vector-mediated

transmission (16, 38), cell to cell movement (21, 39) and new leaves systemic colonization (15, 23,46)

are common during the multiple stages of the infectious cycle of plant viruses. There is experimental

evidence that the number of founders (*i.e.*, viral genomes triggering infection) depends on the stage of

the virus cycle affected by the bottlenecks. For instance, it was reported that from one to six viral

particles infect a cell (the multiplicity of infection, MOI) (44, 45, 437). Considering within-plant

bottlenecks, the lowest N_e reported was ~ 4 founders for wheat streak mosaic virus and the largest

~133 for pea seed-borne mosaic virus (43). It has also been shown that within-host bottleneck size can
largely vary in the range 6 - 107 among sink leaves (39). These different bottlenecks size shape
drastically the virus population. This means that the whole diversity present in a population can be lost
by a single infection event, or be highly reduced for cell-cell or systemic movement. Studying how
Muller's ratchet operates on a plant virus population during its complete cycle (inoculation, movement
and transmission) is therefore highly relevant to understanding its evolutionary dynamics.

In our study, each single lesion in *C. quinoa* represents a sample from a genetically complex viral
population. This feature allowed us to investigate the contribution of different factors influencing the
viral population. Using this approach, the role of bottleneck strength and multiplication rate on
evolution could be investigated demonstrating hereby that different combinations of these two factors
lead to global extinction (small bottleneck, low demographic dynamics), stabilization (intermediate
bottleneck, low demographic dynamics) or rescue of a population (small bottleneck, high demographic
dynamics). The generation of diversity is dependent on the occurrence of random mutations, and the
total number of mutations expected to occur will depend on the number of generations (*i.e.*, rounds of
replication). Our experimental results implicate three general phenomena, genetic drift, generation of
diversity and population size. Firstly, in the small N_e situation, only a few genomes are randomly
sampled every passage and then multiply at a low rate. This low multiplication in a local lesion
generates low diversity. The hypothesis is thus that the principal force acting in this case is genetic
drift through the Muller's ratchet process whereby in a defined population only some individuals
participate to the next generation.

Secondly, passages following the intermediate N_e conditions (weaker bottleneck) maintained the
original diversity generated in *N. tabacum*, but without any gain in fitness. In this modality, the missing
force is the generation of diversity through a high multiplication rate and selection for efficient
replication in *N. tabacum*. Regarding the results from the *in silico* experiments, we can extract a
different general range of maximum multiplication rates of virions inside a lesion between the assumed
virus particles A and D. This result could be explained by a better adaptation to *C. quinoa* of the
defective particle (virion D). Finally, a large range of values for the replication rate in *C. quinoa* either
for A or D virions can be explained by the variation due to the host's response and its ability to generate
the necrotic local lesions (41). Thirdly, these elements can be introduced by alternating passages in *C.*
quinoa and *N. tabacum* (large N_e treatment)*: despite the narrow bottleneck imposed by passaging only
a single *C. quinoa* local lesion, diversity was generated rapidly and selection acted efficiently in *N.*
tabacum. Such a net effect not only attenuated Muller's ratchet effect but also induced an increase in
fitness.

Modeling the experiment allows us to explore some features of the initiation and replication inside
a local lesion and even if results of simulated and experimental data are close, discrepancies could be

explained by different factors. First, we bounded the maximum replication rates for the viruses inside the lesion to 50. This is an arbitrary choice and this value can be larger, in particular for the D virions, where the maximum seems not be attained yet (Fig. 4). Anyway, the curves of occurrence of the values of r_{max}^D show a similar shape than observed for r_{max}^A, suggesting r_{max}^D shall be close to its maximum value. In the simulation of the small N_e model, two replicates show an intriguing pattern of increase in infectivity in *N. tabacum* (Fig. 5A). This suggests that several non-mutually exclusive explanations: (*i*) our plant experiments involved a limited number of lineages and thence we missed these exceptional cases. (*ii*) The assumption of only two types of virions (A and D) is far too simplistic and a gradient of fitness effects could exist. And (*iii*) some factors possibly implicated in the virus life cycle inside a lesion have not been well incorporated in the model, as for example the phenomenon of cooperation. Despite these differences, our model sets specific demographic parameters inside a lesion within a certain order of magnitude, which allows us to understand the virus cycle in such a particular environment. One first result of the model indicates that a single virion is enough to generate a lesion as previously described to initiate an infection (20). The maximum number of particles initiating a lesion (*Vmax*) is in the order of the single digit. These result echoes the number of virions transmitted by aphids (16, 38) and confirms the consequences of limited number of virions starting a new generation. In the case of the multiplication rate inside a lesion, the range is wider compared to the initiation of a lesion. This finding suggests that the host's response, which includes various defense mechanisms, can vary in effectiveness and speed of activation.

Fig. 6 depicts model of the effect of bottlenecks in plant virus transmission. After a first bottleneck in *C. quinoa* (Fig. 6A) the inter-lineage genetic variability was different for each demographic condition. For the case of small N_e, the founder effect resulted in a maximal inter-lineage variation (Fig. 6B), a moderate inter-lineage variation of infectivity for large N_e dynamic (Fig. 6C) and a reduce inter-lineage variation for large N_e dynamics together with a generation of diversity (Fig. 6D). The intra-lineage diversity produced is small, moderate and larger for small, intermediate and large N_e demographics respectively.

In the light of these results, combination of a founder effect with a better multiplication in *C. quinoa* of one variant can explain the rate of infectivity evolution of the small and the intermediate N_e modality.

Because of Muller's ratchet effect, it can be anticipated that an asexual organism with high rate of mutation and small population size will not recover its initial fitness (12). Muller's ratchet effects can be slowed or stopped by various factors as (*i*) increasing population size, (*ii*) selection and purge of deleterious genomes, (*iii*) reducing the mutation rate, (*iv*) appearance of compensatory or back-mutation, and (*v*) recombination. In our experimental model, passaging of local lesions in *C. quinoa* results in a drastic reduction of the probability that such events will occur. Indeed, the virus population

384 is contained in few cells and cannot expand. Given the small population size low generation number,
385 the distribution of mutational (44, 45) and recombination effects (46), the probability that any
386 fortuitous genetic changes be beneficial and not neutral or deleterious is very low.
387

Fig. 5. (A) Hypothetical effect of demographic bottlenecks of different size in the genetic diversity intra- and inter-linages after passage in *C. quinoa* after on passage. (B) For one lesion, (C) a group of lesions or (d) including a large population expansion in a systemic host such as *N. tabacum*.

RNA viruses encounter different bottlenecks during infection and expansion in plants from
inoculation by vectors to cell-cell and systemic movement (15). In this work, by modulating the
bottleneck size through single/grouped lesions and multiplication in *N. tabacum*, we have been able
to mimic the variety of bottlenecks encounter by the virus during its natural cycle. These bottlenecks
may allow the virus to explore the fitness landscape, by sampling viral population in the plant (7).
Moreover, in case of cellular co-infection, fitness loss during repeated narrow genetic bottlenecks is
an essential element of rapid selection. Plant RNA viruses rather than suffering from fitness losses use
this mechanism to favorized their adaptative variants (7, 8, 21). Moreover, when populations regularly
experience bottlenecks, as viruses, increasing the fitness cost of mutations so that unfit mutants are
less likely to fix at each passage is an adaptative solution (47). This approach to quickly remove non-
viable genomes is affected when serial bottlenecks are experimented and moreover in a new
environment leading to a weak selection. In this study, we show that viral populations draw their

strength from their ability to generate large populations. This ability coupled with high fitness costs during mutations allows facing the various bottlenecks encountered in a host.

It has been demonstrated that the average fitness of subclones does not predict the fitness of the whole clonal population at early time point (40, 48). This result underlines the advantage to group several lesions to avoid fitness losses. Moreover, this study has shown that differences in fitness between subclones and population disappear after a long period of replication; in our case evolution of populations' fitness in *C. quinoa* is stopped, although in *N. tabacum* the fitness can evolve.

These results highlight the combined effects of transmission bottlenecks, multiplication rate and selection in virus evolution in a plant system, as well as the life history of the founders. This mutation-selection balance can maximize adaptability while maintaining genetic structure of the virus population (48). Our data reinforce theoretical studies concluding that the mutation rate evolved in concert with the number of replication rounds coupled to bottlenecks events, thus a single founder particle is able to generate an expanding virus population.

ACKNOWLEDGEMENTS

We thank Francisca de la Iglesia for excellent technical assistance. SFE was supported by grants PID2022-136912NB-I00 funded by MCIN/AEI/10.13039/501100011033 and European Union NextGenerationEU/PRTR and CIPROM/2022/59 funded by Generalitat Valenciana. We are grateful to the genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, https://doi.org/10.15454/1.5572369328961167E12) and the computing facilities MCIA (Mésocentre de Calcul Intensif Aquitain) of the Université de Bordeaux and of the Université de Pau et des Pays de l'Adour for providing help and computing resources.

REFERENCES

1. Drake JW, Holland JJ. 1999. Mutation rates among RNA viruses. Proc Natl Acad Sci USA 96:13910–13913.

2. Sanjuán R, Nebot MR, Chirico N, Mansky LM, Belshaw R. 2010. Viral mutation rates. J Virol 84:9733–9748.

3. Malpica J, Fraile A, Moreno I, Obies CI, Drake JW, García-Arenal F. 2002. The rate and character of spontaneous mutation in an RNA virus. Genetics 162:1505–1511.

4. Tromas N, Elena SF. 2010. The rate and spectrum of spontaneous mutations in a plant RNA virus. Genetics 185:983–989.

5. Elena SF, Bedhomme S, Carrasco P, Cuevas JM, de la Iglesia F, Lafforgue G, Lalić J, Pròsper A,

Tromas N, Zwart MP. 2011. The evolutionary genetics of emerging plant RNA viruses. Mol Plant
Microbe Interact 24:287–293.

6. Wright S. 1931. Evolution in mendelian populations. Genetics 16:97–159.

7. Zwart MP, Elena SF. 2015. Matters of size: genetic bottlenecks in virus infection and their
potential impact on evolution. Annu Rev Virol 2:161–179.

8. Cases-González C, Arribas M, Domingo E, Lázaro E. 2008. Beneficial effects of population
bottlenecks in an RNA virus evolving at increased error rate. J Mol Biol 384:1120–1129.

9. Bull RA, Luciani F, McElroy K, Gaudieri S, Pham ST, Chopra A, Cameron B, Maher L, Dore
GJ, White PA, Lloyd AR. 2011. Sequential bottlenecks drive viral evolution in early acute
hepatitis C virus infection. PLoS Pathog 7:e1002243.

10. Kimura M, Ota T. 1969. The average number of generations until extinction of an individual
mutant gene in a finite population. Genetics 63:701–709.

11. Kimura M, Ohta T. 1969. The average number of generations until fixation of a mutant gene in a
finite population. Genetics 61:763–771.

12. Muller HJ. 1964. The relation of recombination to mutational advance. Mutant Res 1:2–9.

13. Zwart MP, Hemerik L, Cory JS, de Visser JAGM, Bianchi FJJA, Van Oers MM, Vlak JM,
Hoekstra RF, Van der Werf W. 2009. An experimental test of the independent action hypothesis
in virus-insect pathosystems. Proc R Soc B 276:2233–2242.

14. Van der Werf W, Hemerik L, Vlak JM, Zwart MP. 2011. Heterogeneous host susceptibility
enhances prevalence of mixed-genotype micro-parasite infections. PLoS Comput Biol
7:e1005097.

15. Ali A, Li H, Schneider WL, Sherman DJ, Gray S, Smith D, Roossinck MJ. 2006. Analysis of
genetic bottlenecks during horizontal transmission of cucumber mosaic virus. J Virol 80:8345–
50.

16. Betancourt M, Fereres A, Fraile A, Garcia-Arenal F. 2008. Estimation of the effective number of
founders that initiate an infection after aphid transmission of a multipartite plant virus. J Virol
82:12416–12421.

17. Couch RB, Cate TR, Douglas RG, Gerone PJ, Knight V. 1966. Effect of route of inoculation on
experimental respiratory viral disease in volunteers and evidence for airborne transmission.
Bacteriol Rev 30:517–529.

18. Moury B, Fabre F, Senoussi R. 2007. Estimation of the number of virus particles transmitted by
an insect vector. Proc Natl Acad Sci USA 104:17891–17896.

19. Da Silva W, Kutnjak D, Xu Y, Xu Y, Giovannoni J, Elena SF, Gray S. 2020. Transmission modes
affect the population structure of potato virus Y in potato. PLoS Pathog 16:e1008608.

20. Zwart MP, Daròs JA, Elena SF. 2011. One is enough: *in vivo* effective population size is dose-

dependent for a plant RNA virus. PLoS Pathog 7:e1002122.

21. Miyashita S, Kishino H. 2010. Estimation of the size of genetic bottlenecks in cell-to-cell movement of soil-borne wheat mosaic virus and the possible role of the bottlenecks in speeding up selection of variations in trans-acting genes or elements. J Virol 84:1828–1837.

22. Monsion B, Froissart R, Michalakis Y, Blanc S. 2008. Large bottleneck size in cauliflower mosaic virus populations during host plant colonization. PLoS Pathog 4:e1000174.

23. Sacristán S, Malpica JM, Fraile A, García-Arenal F. 2003. Estimation of population bottlenecks during systemic movement of tobacco mosaic virus in tobacco plants. J Virol 77:9906–9911.

24. Elena SF, Sanjuán R. 2005. Adaptive value of high mutation rates of RNA viruses: separating causes from consequences. J Virol 79:11555–11558.

25. Chao L. 1990. Fitness of RNA virus decreased by Muller's ratchet. Nature 348:454–455.

26. Duarte E, Clarke D, Moya A, Domingo E, Holland J. 1992. Rapid fitness losses in mammalian RNA virus clones due to Muller's ratchet. Proc Natl Acad Sci USA 89:6015–6019.

27. Escarmis C, Davila M, Charpentier N, Bracho A, Moya A, Domingo E. 1996. Genetic lesions associated with Muller's ratchet in an RNA virus. J Mol Biol 264:255–267.

28. Yuste E, Sánchez-Palomino S, Casado C, Domingo E, López-Galíndez C. 1999. Drastic fitness loss in human immunodeficiency virus type 1 upon serial bottleneck events. J Virol 73:2745–2751.

29. De la Iglesia F, Elena SF. 2007. Fitness declines in tobacco etch virus upon serial bottleneck transfers. J Virol 81:4941–4947.

30. Andersson DI, Hughes D. 1996. Muller's ratchet decreases fitness of a DNA-based microbe. Proc Natl Acad Sci USA 93:906–907.

31. Zeyl C, Mizesko M, de Visser JAGM. 2001. Mutational meltdown in laboratory yeast populations. Evolution 55:909–917.

32. Felsenstein J. 1974. The evolution advantage of recombination. Genetics.

33. Haigh J. 1978. The accumulation of deleterious genes in a population-Muller's Ratchet. Theor Popul Biol 14:251–267.

34. Novella IS, Duarte EA, Elena SF, Moya A, Domingo E, Holland JJ. 1995. Exponential increases of RNA virus fitness during large population transmissions. Proc Natl Acad Sci USA 92:5841–4.

35. Elena SF, Dávila M, Novella IS, Holland JJ, Domingo E, Moya A. 1998. Evolutionary dynamics of fitness recovery from the debilitating effects of Muller's ratchet. Evolution 52:309–314.

36. Elena SF, González-Candelas F, Novella IS, Duarte EA, Clarke DK, Domingo E, Holland JJ, Moya A. Evolution of fitness in experimental populations of vesicular stomatitis virus. Genetics 142:673–679.

37. Dolja VV, McBride HJ, Carrington JC. 1992. Tagging of plant potyvirus replication and

movement by insertion of beta-glucuronidase into the viral polyprotein. Proc Natl Acad Sci USA 89:10208–10212.

38. Pirone TP, Blanc S. 1996. Helper-dependent vector transmission of plant viruses. Annu Rev Phytopathol 34:227–247.

39. Tromas N, Zwart MP, Lafforgue G, Elena SF. 2014. Within-host spatiotemporal dynamics of plant virus infection at the cellular level. PLoS Genet 10:e1004186.

40. Cervera H, Elena SF. 2016. Genetic variation in fitness within a clonal population of a plant RNA virus. Virus Evol 2:vew006.

41. Loebenstein G. Local lesions and induced resistance. Adv Virus Res 75:73-117.

42. Zwart MP, Tromas N, Elena SF. 2013. Model-selection-based approach for calculating cellular multiplicity of infection during virus colonization of multi-cellular hosts. PLoS ONE 8:e64657.

43. Fabre F, Moury B, Johansen EI, Simon V, Jacquemond M, Senoussi R. 2014. Narrow bottlenecks affect pea seedborne mosaic virus populations during vertical seed transmission but not during leaf colonization. PLoS Pathog 10:e4003833.

44. Carrasco P, de la Iglesia F, Elena SF. 2007. Distribution of fitness and virulence effects caused by single-nucleotide substitutions in tobacco etch virus. J Virol 81:12979-12984.

45. Lalić J, Cuevas JM, Elena SF. 2011. Effect of host species on the distribution of mutational fitness effects for an RNA virus. PLoS Genet 7:e1002378.

46. Vuillaume F, Thébaud G, Urbino C, Forfert N, Granier M, Froissart R, Blanc S, Peterschmitt M. 2011. Distribution of the phenotypic effects of random homologous recombination between two virus species. PLoS Pathog 7:e1002028.

47. Merleau NSC, Pénisson S, Gerrish PJ, Elena SF, Smerlak M. 2021. Why are viral genomes so fragile? The bottleneck hypothesis. PLoS Comput Biol 17:e1009128.

48. Manrubia SC, Escarmís C, Domingo E, Lázaro E. 2005. High mutation rates, bottlenecks, and robustness of RNA viral quasispecies. Gene 347:273–282.

Supplementary Fig. S1. Relationship between infectivity on *N. tabacum* and number of lesions on *C. quinoa* before measure of infectivity.