The potential of young vegetative quinoa (*Chenopodium quinoa*) as a new

sustainable protein-rich winter leafy crop under Mediterranean climate

Aviv Asher[1], Reut Dagan[1], Yaron Lugasi[1], Shmuel Galili[2] and Lior Rubinovich[1]*

[1] Northern Agriculture R&D, MIGAL – Galilee Research Institute, Kiryat Shmona, Israel

[2] Institute of Plant Sciences, Agricultural Research Organization, The Volcani Center, Rishon LeZion, Israel

*Corresponding author

E-mail: liorr@migal.org.il

Abstract

The demand for protein products has significantly risen in the last few years. In western countries, animals are the primary source of protein; however, plants could take a share of this market due to lower production costs, among other advantages. Quinoa (*Chenopodium quinoa* Willd.) is a well-known but under-utilized protein-rich crop, commonly cultivated for grain production. These plants were recently evaluated for their use as a non-traditional, green leafy crop. Here we assessed the potential of young vegetative quinoa as a new sustainable winter leafy crop in Israel—serving as a model for Mediterranean semi-arid regions, by evaluating yield, protein content and quality. Five quinoa accessions were sown on three winter sowing dates over two consecutive years. Plants were harvested when they reached ~10–13% dry matter (DM). DM yield ranged between 574 and 1,982 kg ha^{-1} and was generally higher in the second year. Protein content ranged from 14.4–34% and was generally higher in the first year. Protein yield ranged from 111–471 kg ha^{-1} and was greatest on the December sowing date. DM and protein yields were positively correlated with plant density. Protein content was negatively correlated with plant density and DM yield. Our findings show that 200 g DM of young vegetative quinoa can meet the protein and most essential amino acid requirements for a 70 kg human adult. Prospects for cultivating young vegetative quinoa in Mediterranean countries as a new sustainable, protein-rich winter leafy crop are therefore high, as supported by its high protein yields and quality, and its requirement for only scant irrigation. Further studies should examine economic and other agrotechnical parameters toward the geographical distribution and expansion of young vegetative quinoa cultivation.

Keywords: amino acid; leaf; semi-arid; superfood; sustainable

1. Introduction

The demand for protein products has significantly risen in the last few years. The global protein ingredients market was valued at USD 38.5 billion in 2020 and is expected to expand at an annual growth rate of 10.5% from 2021 to 2028 [1]. A dietary shift in Western countries in the last century resulted in animals becoming the primary source of protein [2]. However, plant proteins could take a share of the animal protein market (meat, egg and dairy) due to lower production costs, and their being perceived by consumers as being more ethical, healthy and environmentally friendly [3,4]. The quinoa plant (*Chenopodium quinoa* Willd., Amaranthaceae) is a well-known but under-utilized protein-rich crop [5]. This pseudo-cereal originates from the South American Andes and is commonly grown for grain production for human consumption [6,7]. Quinoa is considered a climate-resilient crop and has a remarkable ability to grow in marginal environments [8,9]. Recently, the number of quinoa-producing countries has shown a rapid increase, and the crop is now grown commercially outside South America [10,11]. The quinoa grain's protein content (PC) is relatively high compared to the major cereal crops cultivated worldwide, such as wheat, corn and rice [12]; its high nutritional value is due to high levels of protein (11–19%) that contain all of the essential amino acids (EAA) [13,14]. It is also gluten-free, and may therefore be suitable for celiac patients [15,16]. Thus, quinoa is commonly considered a "superfood" or "functional food", enhancing its economic attractiveness [17].

Many studies have shown the possibility of cultivating quinoa outside its area of origin in diverse agroecological regions, including semi-arid and arid climates [6,11]. Other studies have examined the effects of agrotechnical parameters such as row spacing, plant density, irrigation and fertilization on quinoa grain production [18–21]. Numerous recent studies have shown that quinoa

can be used for purposes other than grain production. For example, several studies showed that the

residual straw following grain harvest might be used as livestock feed, as it contains relatively

high protein and mineral contents and is highly digestible [22–24]. Others showed that high quinoa

hay biomass and high hay quality suggest excellent prospects for its use as a high-quality feed for

ruminants [22,25,26]. Interestingly, the Incas and other earlier cultures included quinoa leaves in

their diet to balance the lack of animal protein [27]. Indeed, in recent years, quinoa plants have

also been evaluated for their use as a non-traditional, green leafy crop. The green leaves and tender

stalks have high nutritional value and can be consumed raw or cooked, similar to, for instance,

spinach [28,29]. Several studies have reported that the PC in quinoa leaves is greater than that in

quinoa grains, reaching over 25% in dry matter (DM) [30]. Quinoa leaves also contain higher PC

than broccoli, spinach or mangold [31]. Moreover, quinoa protein contains all of the EAA, and the

leaves have relatively low quantities of carbohydrates. Quinoa leaves also contain moderate levels

of calcium, phosphorus, sodium and zinc, and high levels of copper, manganese and potassium

[32]. Gawlik-Dziki et al. (2013) assessed the nutraceutical potential of quinoa leaves, and found

that their extract contains significant amounts of various bioactive polyphenols linked with an

inhibitory effect on prostate cancer cell proliferation [33]. Moreover, the content of saponins—

antinutritive compounds found in quinoa—is lower in the leaves than in the grains [34,35].

Previous field trials have evaluated the yield of green leafy quinoa. A study conducted in

Poland showed that quinoa should be harvested as a green leafy vegetable when it reaches a height

of 20–30 cm. In spring and summer sowing, young quinoa's average fresh weight biomass reached

317–1389 and 245–777 g m^{-2}, respectively [29]. In Egypt, young quinoa plants harvested 45 days

after sowing had greater fresh weight yield, 2.05–4.14 kg m^{-2} [36]. These findings indicate that

growing quinoa for use as a green leafy vegetable has economic potential.

As there is only scarce data regarding the cultivation of young vegetative quinoa in Mediterranean countries, we aimed to evaluate this crop's potential as a new protein-rich leafy crop in Israel. To that end, we sowed, grew, harvested and evaluated several quinoa accessions for two consecutive years. Winter is the rainy season in Israel, and we sowed the quinoa in November, December and January to examine its potential cultivation under low amounts of irrigation. We hypothesized that young quinoa's aerial vegetative parts can be used as a new, sustainable, protein-rich leafy crop for Israel and other Mediterranean and semi-arid regions. Our findings show high prospects for cultivating young vegetative quinoa in Mediterranean countries as a new sustainable, protein-rich winter leafy crop.

2. Materials and Methods

2.1. Experimental site

The experiments were conducted in northern Israel (The Avnei-Eitan research farm, altitude 375 masl, 32°81′64″N 35°76′28″E) from November 2020 to April 2022. The soil at the experimental site is of basaltic origin, composed of 60% clay, 35% silt and 5% sand with pH 7.6. The climate is Mediterranean, with a long-term annual average rainfall of 500–550 mm, mainly distributed between October and June. 92% of the total precipitation during the experimental season of 2020 to 2021 occurred between November 2020 and February 2021. Total precipitation during the growing season of 2021 to 2022 was slightly lower, with 95% of the total rainfall occurring from December 2021 to March 2022. Air temperature ranges were similar for the two seasons (Table 1).

Table 1

Monthly precipitation and temperatures at the Avnei-Eitan research farm during 2020–2021 and 2021–2022 winter growing seasons (Israeli Meteorological Services).

Month	2020 to 2021			2021 to 2022		
	Rainfall (mm)	Min. temp (°C)	Max. temp (°C)	Rainfall (mm)	Min. temp (°C)	Max. temp (°C)
October	0	13.5	38.1	4.6	11.2	35.2
November	119.1	8.8	26.6	19.7	7.8	29.6
December	100.3	6.2	22.2	96.5	3.7	24.1
January	124.6	0.9	23.3	167.4	-2	21.9
February	109.7	2	23.9	82.7	3.6	25.7
March	24.8	1.4	28.3	83.9	-0.9	24.7
April	12.3	2.7	37.7	0	6.2	35.1
Total	490.8			454.8		

2.2. Plant material and experimental design

Five quinoa accessions were used in the experiments: `Red Head`, `Mint Vanilla`, `Ivory`, `Oro de Valle` and `Peppermint`. All accessions were obtained from Wild Garden Seed (Philomath, OR, USA). In all experiments, flowerbeds were prepared after cultivation and rolling. Sowing density was selected to obtain low stem diameter [19]. Seeds were sown into six rows (16 cm between rows) at an intra-row density of 40–50 seeds m^{-1} using a manual seeder (a planned density of 240–300 seeds m^{-2}). Quinoa accessions were sown into four 8-m^2 (5 m long x 1.6 m wide) plots (repeats) in mid-November and mid-December 2020 and 2021, and mid-January 2021 and 2022. Plots were randomized on each sowing date. A single irrigation using sprinklers at a rate of 100 m^3 ha^{-1} was applied only in plots that did not receive rain within 10 days after sowing. The plots were

fertilized by ~100 kg N ha^{-1} as urea (46% N). All plots were manually weeded once during

cultivation.

2.3. Plant density and yield measurements

To determine plant density at harvest (plants m^{-2}), plants were counted in a representative

1-m^2 area in each plot. When the plants reached ~10–13% DM, the aerial parts of the plants in each

plot were hand-harvested ~5 cm above the ground and weighed for fresh matter yield. To determine

plant %DM, representative samples of each plot were collected and weighed before and after drying

at 60 °C for 48 h. Plot DM yield was calculated as fresh matter yield x plant %DM.

2.4. Plant protein content, protein yield and amino acid composition

Using the Kjeldahl method [32], each quinoa sample's PC (% of DM) was determined at the

Milouda and Migal laboratories in Kiryat Shmona, Israel. Plot protein yield was calculated as DM

yield x PC. The amino acid composition was determined at the Merieux NutriScience laboratories

in Resana, Italy. Briefly, the samples were subjected to hot enzymatic hydrolysis of the proteins.

The solution obtained from hydrolysis was diluted with deionized water and methanol and analyzed

by HPLC.

2.5. Statistical analysis

The data are presented as mean value ± standard error (SE). Results were subjected to one-

way analysis of variance (ANOVA) followed by Tukey's HSD test using JMP software version

11.0.0 (SAS Institute, Cary, NC, USA). Results were also subjected to a two-tailed Pearson

correlation matrix using Prism software version 9.0.0 (GraphPad Software, San Diego, CA, USA).

3. Results

Since Israel and other Mediterranean and semi-arid regions have an expensive and limited
water supply, we investigated the potential of young vegetative quinoa cultivation as a winter crop
(Fig 1). Plots were sown in November, December and January over two consecutive years.

Fig. 1. Young vegetative quinoa. A representative photograph of young vegetative quinoa grown at the Avnei-Eitan
research farm before harvest.

3.1. Plant density

In the first year, plants from the plots sown in November 2020, December 2020 and January
2021 were harvested 70, 73 and 69 days after sowing, respectively. Actual plant densities at harvest
are shown in Fig 2. In plots sown in November 2020, plant density ranged between 90 (accession
Oro de Valle) and 192 ('Peppermint') plants m^{-2}. In plots sown in December 2020, plant density
ranged between 179 and 264 plants m^{-2} (accessions Mint Vanilla and Red Head, respectively).

There were no differences ($P > 0.05$) between plant densities of the different accessions from these

sowing dates. In plots sown in January 2021, plant density ranged between 163 and 302 plants m^{-}

^{2}. Accession Red Head had the highest plant density, significantly ($P < 0.05$) higher than `Oro de

Valle`. In the second year, plants from the plots sown in November 2021, December 2021 and

January 2022 were harvested 99, 107 and 88 days after sowing, respectively (Fig 2a). In plots sown

in November 2021, plant density ranged between 150 and 333 plants m^{-2}. Accession Oro de Valle

had the highest plant density, significantly ($P < 0.05$) higher than `Mint Vanilla` and `Ivory`. In

plots sown in December 2021, plant density ranged between 330 and 521 plants m^{-2}. Accession

Mint Vanilla had the highest plant density, significantly ($P < 0.05$) higher than `Red Head`, `Oro

de Valle` and `Peppermint`. In plots sown in January 2022, plant density ranged between 142 and

567 plants m^{-2}. Accession Mint Vanilla had the highest plant density, significantly ($P < 0.05$)

higher than all other accessions (Fig 2b).

Fig. 2. Young vegetative quinoa plant density. Quinoa was sown in the Avnei-Eitan research farm during the winters

of 2020–2021 and 2021–2022. Results show plant density at harvest in plots sown in November 2020, December

2020, and January 2021 (a); and in November 2021, December 2021, and January 2022 (b). Results are presented as

mean value ± standard error of four 5-m^2 sample plots per accession on each sowing date; columns marked with

different letters differ significantly by Tukey-HSD ($P < 0.05$).

3.2. Dry matter yield

Quinoa DM yield is shown in Fig 3. In plots sown in November 2020, DM yield of young

vegetative quinoa ranged between 572 and 1200 kg DM ha^{-1} (accessions Oro de Valle and

Peppermint, respectively). In plots sown in December 2020, DM yield ranged between 693 and

1069 kg DM ha^{-1} (accessions Oro de Valle and Mint Vanilla), respectively. There were no

differences ($P > 0.05$) between the DM yields of the different accessions on November and

Decenmber 2020 sowing dates. In plots sown in January 2021, DM yield ranged between 701 and

1174 kg DM ha^{-1}. Accession Peppermint had the highest DM yield, significantly higher ($P < 0.05$)

than 'Oro de Valle' (Fig 3a). In the second year (Fig 3b), in plots sown in November and December

2021 and January 2022, DM yield ranged between 1088 and 1434 kg DM ha^{-1} ('Red Head' and

'Oro de Valle'), 1477 and 1982 kg DM ha^{-1} ('Red Head' and 'Ivory'), and 618 and 1384 kg DM

ha^{-1} ('Red Head' and 'Ivory'), respectively. There were no significant differences ($P > 0.05$)

between the DM yields of the different accessions for any of these sowing dates.

Fig. 3. Young vegetative quinoa yield. Quinoa was sown in the Avnei-Eitan research farm during the winters of 2020–2021 and 2021–2022. Results show DM yield in plots sown in November 2020, December 2020 and January 2021 (a); and in November 2021, December 2021, and January 2022 (b). Results are presented as mean value ± standard error of four 5-m^2 sample plots per accession on each sowing date; columns marked with different letters differ significantly by Tukey-HSD ($P < 0.05$).

3.3. Protein content and protein yield

Quinoa PC is shown in Fig 4. In plots sown in November 2020, the PC of young vegetative quinoa ranged between 30.6 and 32.4% (accessions Peppermint and Red Head, respectively). In plots sown in December 2020, quinoa PC ranged between 28.2 and 34% ('Red Head' and 'Mint

Vanilla', respectively). In plots sown in January 2021, quinoa PC ranged between 24.8 and 29.2%

('Ivory' and 'Mint Vanilla', respectively) (Fig 4a). In the second year (Fig 4b), in plots sown in

November 2021, quinoa PC ranged between 25.9 and 28.3% ('Oro de Valle' and 'Red Head',

respectively). In plots sown in December 2021, quinoa PC ranged between 19.4 and 23.7% ('Mint

Vanilla' and 'Ivory', respectively). In plots sown in January 2022, quinoa PC ranged between 14.4

and 19.4% ('Ivory' and 'Peppermint', respectively). There were no significant differences ($P >$

0.05) between the PC of the different accessions for any of these sowing dates.

Fig. 4. Young vegetative quinoa protein content. Quinoa was sown in the Avnei-Eitan research farm during the winters

of 2020–2021 and 2021–2022. Results show protein content (% DM) in quinoa plants sown in November 2020,

December 2020, and January 2021 (a); and in November 2021, December 2021, and January 2022 (b). Results are

presented as mean value ± standard error of four 5-m² sample plots per accession on each sowing date.

Quinoa protein yield is shown in Fig 5. In plots sown in November 2020, protein yield

ranged between 182 and 366 kg ha^{-1} (accessions Oro de Valle and Peppermint, respectively). In

plots sown in December 2020, protein yield ranged between 230 and 359 kg ha^{-1} ('Oro de Valle'

and 'Mint Vanilla', respectively). In plots sown in January 2021, protein yield ranged between 201

and 303 kg ha^{-1} ('Oro de Valle' and 'Peppermint', respectively) (Fig 5a). In the second year (Fig

5b), in plots sown in November 2021, protein yield ranged between 303 and 367 kg ha^{-1} ('Ivory'

and 'Oro de Valle', respectively). In plots sown in December 2021, protein yield ranged between

330 and 471 kg ha^{-1} ('Red Head' and 'Ivory', respectively). In plots sown in January 2022, protein

yield ranged between 111 and 240 kg ha^{-1} ('Red Head' and 'Mint Vanilla', respectively). There

were no significant differences ($P > 0.05$) between the protein yields of the different accessions

for each of these sowing dates.

Fig. 5. Young vegetative quinoa protein yield. Quinoa was sown in the Avnei-Eitan research farm during the winters

of 2020–2021 and 2021–2022. Results show protein yield in quinoa plants sown in November 2020, December 2020

and January 2021 (a); and in November 2021, December 2021, and January 2022 (b). Results are presented as mean

value ± standard error of four 5-m^2 sample plots per accession on each sowing date.

3.4. Correlation between traits

Young vegetative quinoa DM yield was positively correlated (r = 0.616, $P < 0.0001$) with

plant density (Fig 6a), but the opposite trend was found for PC (r = -0.376, $P < 0.0001$) (Fig 6b).

A positive correlation (r = 0.43, $P < 0.0001$) was found between protein yield and plant density

(Fig. 6c). The correlation between PC and DM yield was negative (r = -0.311, $P < 0.001$) (Fig 6d).

Fig. 6. Correlations of young vegetative quinoa yield parameters. Pearson's correlation coefficients between yield and plant density at harvest (a), protein content and plant density at harvest (b), protein yield and plant density at harvest (c), and protein content and yield (d). Values represent measurements from all plots from the winters of 2020–2021 and 2021–2022; each value represents measurements from one plot. The correlation coefficient value (r) is shown for each graph along with the significance of the Pearson correlation values ($***P < 0.001$, $****P < 0.0001$).

3.5. Amino acid composition

As accession Peppermint had relatively high protein yield values during the first year of the experiment, it was chosen for amino acid composition analysis. During the first year of the

investigation, the young vegetative quinoa contained all of the EAA (Table 2). Among them, the

quinoa was characterized by a high content of leucine and lysine. Their levels were significantly

($P < 0.05$) higher in plots sown in December 2020 compared to November 2020. Tryptophan levels

were significantly ($P < 0.05$) higher in plots sown in December 2020 and January 2021 compared

to November 2020. There were no significant differences ($P > 0.05$) in the other EAA levels

between the different sowing dates. For comparison, the recommended daily EAA intake for

humans (g per 70 kg body weight) is shown in Table 2. On the different sowing dates, histidine,

isoleucine, leucine, lysine, methionine, phenylalanine (with tyrosine), threonine, tryptophan and

valine levels in the young vegetative quinoa ranged from 50–63%, 41–57%, 53–68%, 60–112%,

37–50%, 79–110%, 84–119%, 70–164% and 53–63% of the recommended daily intake,

respectively. Among the non-essential amino acids (NEAA), young vegetative quinoa accession

Peppermint was characterized by high glutamic and aspartic acid contents. Aspartic acid and serine

levels were significantly ($P < 0.05$) higher in plots sown in December 2020 and January 2021

compared to November 2020. Glutamic acid and glycine levels were significantly ($P < 0.05$)

higher in plots sown in December 2020 compared to November 2020 and January 2021. Alanine

levels were significantly ($P < 0.05$) higher in plots sown in December 2020 compared to November

2020. In the experiment's first year, there were no significant differences ($P > 0.05$) in the other

NEAA levels between the different sowing dates (Table 2).

Table 2

Amino acid composition (g per100 g DM) of accession Peppermint from the 2020–2021 winter

sowing dates: November 2020, December 2020, and January 2021.

Amino acids (g per 100 g DM)	Nov '20	Dec '20	Jan '21	Recommended daily intake (g per 70 kg body weight) *
Essential				
Histidine	0.35 ± 0.02	0.39 ± 0.08	0.44 ± 0.02	0.7
Isoleucine	0.8 ± 0.07	0.78 ± 0.06	0.58 ± 0.04	1.4
Leucine	1.46 ± 0.07 b	1.86 ± 0.1 a	1.58 ± 0.1 ab	2.73
Lysine	1.27 ± 0.09 b	2.35 ± 0.33 a	1.80 ± 0.12 ab	2.1
Methionine	0.29 ± 0.02	0.35 ± 0.06	0.26 ± 0.04	0.7
Phenylalanine	0.88 ± 0.04	1.19 ± 0.12	0.99 ± 0.08	1.75**
Threonine	0.88 ± 0.04	1.25 ± 0.13	1.21 ± 0.09	1.05
Tryptophan	0.197 ± 0.08 b	0.46 ± 0.02 a	0.43 ± 0.15 a	0.28
Valine	0.96 ± 0.05	1.14 ± 0.03	1.00 ± 0.15	1.82
Non-essential				
Alanine	1.05 ±0.12 b	1.63 ± 0.05 a	1.33 ± 0.08 ab	
Arginine	0.97 ±0.06	1.30 ± 0.12	0.94 ± 0.22	
Aspartic acid	1.85 ± 0.09 b	2.86 ± 0.11 a	2.53 ± 0.09 a	
Cystine + Cysteine	0.26 ± 0.01	0.27 ± 0.03	0.23 ± 0.01	
Glutamic acid	2.54 ± 0.18 b	3.64 ± 0.41 a	2.45 ± 0.18 b	
Glycine	0.97 ± 0.04 b	1.47 ± 0.09 a	1.14 ± 0.03 b	

Proline	0.88 ± 0.08	0.58 ± 0.09	0.55 ± 0.22
Serine	0.89 ± 0.3 b	1.34 ± 0.09 a	1.22 ± 0.03 a
Tyrosine	0.51 ± 0.1	0.68 ± 0.17	0.95 ± 0.11

Results are expressed as mean value ± standard error of four samples per sowing date; different lowercase letters in a

row indicate a significant difference (Tukey HSD, $P < 0.05$).

*Data from [37].

**In total with tyrosine.

During the second year of the experiment, the young vegetative quinoa also contained all EAA and was characterized by a high content of leucine and lysine (Table 3). However, there were no significant differences ($P > 0.05$) in EAA levels between the different sowing dates. On the different sowing dates, histidine, isoleucine, leucine, lysine, methionine, phenylalanine (with tyrosine), threonine, tryptophan and valine levels in the young vegetative quinoa, ranged from 53–66%, 49–72%, 52–65%, 50–60%, 46–54%, 87–103%, 77–98%, 54–66% and 52–66% of the recommended daily EAA intake (Table 3), respectively. Among the NEAA, young vegetative quinoa was characterized by relatively high glutamic acid, aspartic acid and arginine contents. Alanine, arginine and serine levels were significantly ($P < 0.05$) higher in plots sown in November 2021 compared to January 2022. Glutamic acid levels were significantly ($P < 0.05$) higher in plots sown in November 2021 compared to December 2021 and January 2022. There were no significant differences ($P > 0.05$) in the other NEAA levels between the different sowing dates of the second year of the experiment (Table 3).

Table 3

311   Amino acid composition (g per100 g DM) of accession Peppermint from the 2021–2022 winter

312   sowing dates: November 2021, December 2021, and January 2022.

Amino acids (g per 100 g DM)				Recommended daily intake (g per 70 kg body weight) *
	Nov '21	Dec '21	Jan '22	
Essential				
Histidine	0.46 ± 0.04	0.41 ± 0.02	0.37 ± 0.05	0.7
Isoleucine	1.01 ± 0.06	0.98 ± 0.05	0.82 ± 0.1	1.4
Leucine	1.78 ± 0.11	1.63 ± 0.08	1.43 ± 0.18	2.73
Lysine	1.26 ± 0.1	1.17 ± 0.07	1.05 ± 0.13	2.1
Methionine	0.38 ± 0.03	0.37 ± 0.03	0.32 ± 0.04	0.7
Phenylalanine	1.09 ± 0.07	1.03 ± 0.05	0.89 ± 0.12	1.75**
Threonine	1.03 ± 0.07	0.89 ± 0.04	0.81 ± 0.1	1.05
Tryptophan	0.19 ± 0.016	0.15 ± 0.009	0.15 ± 0.02	0.28
Valine	1.20 ± 0.08	1.09 ± 0.06	0.94 ± 0.12	1.82
Non-essential				
Alanine	1.58 ±0.14 a	1.15 ± 0.05 ab	1.04 ± 0.13 b	
Arginine	2.04 ±0.31 a	1.29 ± 0.11 ab	1.09 ± 0.14 b	
Aspartic acid	2.02 ± 0.14	1.79 ± 0.08	1.6 ± 0.2	
Cystine + Cysteine	0.31 ± 0.02	0.29 ± 0.02	0.25 ± 0.03	

Glutamic acid	3.47 ± 0.48 a	2.17 ± 0.1 b	1.95 ± 0.26 b
Glycine	1.19 ± 0.09	1.03 ± 0.05	0.89 ± 0.11
Proline	0.98 ± 0.06	0.88 ± 0.05	0.8 ± 0.1
Serine	1.11 ± 0.1 a	0.86 ± 0.03 ab	0.79 ± 0.09 b
Tyrosine	0.72 ± 0.04	0.67 ± 0.03	0.63 ± 0.08

Results are expressed as mean value ± standard error of four samples per sowing date; different lowercase letters in a

row indicate a significant difference (Tukey HSD, $P < 0.05$).

*Data from [37].

**In total with tyrosine.

4. Discussion

We assessed the potential of young vegetative quinoa as a new winter leafy crop in Israel

and as a model for Mediterranean semi-arid regions by evaluating yield, PC and quality. Five

quinoa accessions were sown on three winter sowing dates over two consecutive years. The

greatest yield (1982 kg DM ha^{-1}) was achieved with accession Ivory sown in December 2021 (Fig

3). The cultivation of quinoa as a leafy crop is still rare worldwide, and there are therefore no

comparable data on its production parameters from the Food and Agriculture Organization of the

United Nations (FAO). However, the fresh yield of quinoa was relatively high compared to the

commercial yields of spinach, a more common leafy crop. According to the FAO, the fresh yield

of commercial spinach in Italy and Egypt from 2018 to 2021 ranged between 15,558 and 16,574

kg ha^{-1} and 14,952 and 16,411 kg ha^{-1}, respectively ([www.fao.org,](www.fao.org) accessed 2.2023). However, it

is essential to note that because crops are given more care and attention in small-scale field studies,

yields are often higher than in commercial-scale production. Even so, the quinoa yield obtained in

this study was higher than that reported in earlier small-scale field experiments. For example, a

recent study showed that under saline and non-saline conditions in Egypt, quinoa fresh vegetative

yield reached 4470 and 5985 kg ha^{-1}, respectively [30]. In another study conducted in India, the

fresh yield of the first harvest of quinoa foliage reached between 6240 and 4100 kg ha^{-1} in two

consecutive seasons [38]. Interestingly, the reported cumulative fresh yield of three consecutive

harvests of the same plots reached 18,999 and 13,900 kg ha^{-1} in the two respective seasons of the

experiment. Thus, quinoa yield may be significantly increased by continuing to grow the plants

after the first harvest and harvesting them at intervals, similar to spinach, alfalfa, elephant grass,

and other leafy field crops [39–41]. However, multiple harvests may eventually reduce quinoa

plant yield and quality [38]. Another suggested possibility for maximizing quinoa yield while

retaining quality is thus to establish a new cultivation on the same plot following harvest, similar

to spinach and rocket [41].

In this study, PC in young vegetative quinoa ranged between 14.4% and 34%, and in

general, it was higher in the first year of the experiment than in the second year (Fig 4). This can

be explained by the fact that actual plant density was generally lower in the first vs. second year

(Fig 2) and by the negative correlation between actual plant density and PC (Fig 6b). Still, PC was

relatively higher than in quinoa grains (Pereira et al., 2019): in a previous study conducted in Israel

under similar conditions, quinoa grain PC ranged between 5.3% and 14.2% (Asher et al., 2022);

in another study conducted in Vietnam, quinoa grain PC ranged between 15.01% and 18.49% [42].

In addition, the PC of young vegetative quinoa was relatively higher than in other vegetative crops,

such as spinach, with a reported PC of between 19.9% and 22.8% [36]. Hence, from these data,

young green quinoa may be cultivated in Israel and other Mediterranean countries as a potential

source of plant-based protein. Supporting this notion is the amino acid composition of the young

green quinoa protein (Tables 2 and 3). In most cases, the EAA composition in 100 g DM of young

green quinoa fulfilled over 50% of the recommended daily intake of EAA for a 70 kg human adult

[37]. In some cases, the EAA composition was above 100% of the recommended daily intake.

These data suggest the high quality of young green quinoa protein, because the nutritional quality

of a protein is determined by its concentration of EAA, which humans cannot synthesize [32].

Considering that the protein requirement for human adults stands at 0.66 g kg^{-1} per day [37], 200

g DM of young green quinoa can meet the protein and most of the EAA requirements for a 70 kg

adult. This is of clear economic and social significance, because the consumption of sufficient

high-quality protein in the diet poses a challenge for some populations, particularly those who are

limiting, or have eliminated their animal protein intake and must rely on cereals, grains, and

legumes for their protein consumption [12].

Plant density is a critical factor affecting quinoa and other commercial field crops' yield

and quality traits [43]. In this study, it was interesting to find that plant density differed in the plots

(Fig 2), possibly because of variation in seed quality, inaccuracies in the sowing operation or failed

seedling development due to biotic or abiotic stress conditions [44,45]. However, these

assumptions cannot be established in this present study, as germination studies were not conducted.

Young vegetative quinoa DM and protein yields were both positively correlated to plant density

at harvest (Fig 6a, c). In contrast, a study conducted in India showed that quinoa foliage yield

generally increases with increasing row spacing [38]. Thus, the optimal plant density should be

carefully considered, because low plant density may result in thicker quinoa stems [19], reducing

their palatability for humans. However, high plant density may reduce plant quality, as in this

study, young vegetative quinoa PC was negatively correlated to plant density at harvest (Fig 6b).

Future research should address this issue.

In both years of the experiment, young green quinoa protein yield was greatest on the December sowing date (Fig 5). In addition to this advantage, December is better for sowing crops because weed control is simpler than in November. As the weeds have already germinated in December, they can be effectively eliminated by soil cultivation or a suitable non-residual herbicide. With the help of this weed-control strategy, new quinoa shoots can grow with little or no requirement for herbicides [22]. Thus, the results of this present study suggest that the December sowing date is more appropriate for cultivation in the experimental setting. Additional in-depth research is necessary to further investigate the effect of climate and seasonal parameters on young vegetative quinoa production and quality. This is important, particularly in the Mediterranean region where precipitation is expected to decrease and temperature to warm significantly beyond the global average [46]. Moreover, protein yield was among the highest in most cases, albeit not significantly so, for the accession Peppermint (Fig 5). Therefore, this accession seems more suitable for cultivation in the experimental location when considering young vegetative quinoa for protein yield. Additional studies are required to further investigate how different genotypes impact the quality and production of young vegetative quinoa.

In previous studies conducted in Israel, quinoa grain yield reached 4266 kg DM ha^{-1}, whereas PC stood at 14.2% [19]. From these data, protein yield reached 605 kg ha^{-1}. In the present study, conducted under similar conditions, protein yield of young vegetative quinoa was about 50% lower (Fig 5). However, a significant advantage of cultivating young leafy quinoa is the short growing cycle from sowing to harvest, usually 60–80 days during the winter. In comparison, winter quinoa cultivation for grain production is much longer, 150–170 days. Therefore, multiple harvests of young vegetative quinoa or establishing a new cultivation on the same plot following harvest may produce a similar or greater protein yield than quinoa grown for grains. In addition, a shorter

growing cycle means that the plants may avoid various stresses, such as extreme weather events

(Lakshmanan and Robinson, 2013; Wahbi and Sinclair, 2005), and the young vegetative quinoa can be

cultivated between winter and summer yearly crop cycles, thus maximizing land and other

agricultural resource usages. Finally, the young vegetative quinoa requires scant irrigation, making

it highly suitable as a protein-rich crop for semi-arid and possibly even arid regions, where

irrigation infrastructure is a resource that is in short supply [47].

5. Conclusions

In conclusion, it seems that according to our research hypothesis, there are high prospects

for cultivating young vegetative quinoa in Israel and other Mediterranean countries as a new

sustainable protein-rich winter leafy crop. The fact that quinoa had high protein yield and quality,

even though the plants received only a small amount of irrigation ($100 \text{ m}^3 \text{ ha}^{-1}$), supports this

notion. This is crucial for agricultural production in arid and semi-arid regions where access to

irrigation water is challenging and expensive. Nevertheless, further studies are required to examine

other agrotechnical parameters, such as row spacing, plant density, irrigation and fertilization

regimes, and their effects on yield and quality parameters. Studies should also evaluate the

possibility of several growing cycles of young vegetative quinoa on the same plots with different

sowing dates to maximize land use and yearly yield potential. Finally, evaluating its potential as a

short-cycle high-quality forage crop and various economic aspects such as production cost will

enable young vegetative quinoa's geographical distribution and expansion.

Acknowledgements

The authors thank Shaul Graph, Itzik Abarbanel and the Avnei-Eitan research farm team

for their efforts in performing this study. The ICA foundation in Israel funded several aspects of

this study.

References

1.   Grandview research. Global Protein Ingredients Market Size Report, 2021-2028. 2020

[cited 29 Jan 2023]. Available: https://www.grandviewresearch.com/industry-

analysis/protein-ingredients-market

2.   de Boer J, Aiking H. On the merits of plant-based proteins for global food security:

Marrying macro and micro perspectives. Ecological Economics. 2011. pp. 1259–1265.

doi:10.1016/j.ecolecon.2011.03.001

3.   Pam Ismail B, Senaratne-Lenagala L, Stube A, Brackenridge A. Protein demand: Review

of plant and animal proteins used in alternative protein product development and

production. Animal Frontiers. 2020;10: 53–63. doi:10.1093/af/vfaa040

4.   Aschemann-Witzel J, Gantriis RF, Fraga P, Perez-Cueto FJA. Plant-based food and

protein trend from a business perspective: markets, consumers, and the challenges and

opportunities in the future. Critical Reviews in Food Science and Nutrition. Taylor and

Francis Inc.; 2020. pp. 1–10. doi:10.1080/10408398.2020.1793730

5.   Andreotti F, Bazile D, Biaggi C, Callo-concha D, Jacquet J, Jemal OM, et al. When

neglected species gain global interest: Lessons learned from quinoa's boom and bust for

teff and minor millet. Glob Food Sec. 2022;32: 100613. doi:10.1016/j.gfs.2022.100613

6.   Bazile D, Jacobsen S-E, Verniau A. The Global Expansion of Quinoa: Trends and Limits.

Front Plant Sci. 2016;7: 1–6. doi:10.3389/fpls.2016.00622

7.   Bazile D, Bertero HD, Nieto C. State of the Art Report on Quinoa around the World -

International Year of Quinoa 2013. Rome, Italy; 2015. Available:

http://www.fao.org/3/i4042e/I4042E.pdf

8.   Choukr-Allah R, Rao NK, Hirich A, Shahid M, Alshankiti A, Toderich K, et al. Quinoa

for marginal environments: Toward future food and nutritional security in MENA and

central Asia regions. Frontiers in Plant Science. Frontiers Research Foundation; 2016.

doi:10.3389/fpls.2016.00346

9.   Pulvento C, Bazile D. Worldwide Evaluations of Quinoa—Biodiversity and Food Security

under Climate Change Pressures: Advances and Perspectives. Plants. 2023;12: 868.

doi:10.3390/plants12040868

10.   Jacobsen SE. The scope for adaptation of quinoa in Northern Latitudes of Europe. J Agron

Crop Sci. 2017;203: 603–613. doi:10.1111/jac.12228

11.   Alandia G, Rodriguez JP, Jacobsen SE, Bazile D, Condori B. Global expansion of quinoa

and challenges for the Andean region. Glob Food Sec. 2020;26.

doi:10.1016/j.gfs.2020.100429

12.   Scanlin L, Lewis KA. Quinoa as a Sustainable Protein Source: Production, Nutrition, and

Processing. Sustainable Protein Sources. Elsevier Inc.; 2017. pp. 223–238.

doi:10.1016/B978-0-12-802778-3.00014-7

13.   Pereira E, Encina-Zelada C, Barros L, Gonzales-Barron U, Cadavez V, C.F.R. Ferreira I.

Chemical and nutritional characterization of *Chenopodium quinoa* Willd (quinoa) grains:

A good alternative to nutritious food. Food Chem. 2019;280: 110–114.

doi:10.1016/J.FOODCHEM.2018.12.068

14.   Angeli V, Silva PM, Massuela DC, Khan MW, Hamar A, Khajehei F, et al. Quinoa

(*Chenopodium quinoa* Willd.): An overview of the potentials of the "golden grain" and

socio-economic and environmental aspects of its cultivation and marketization. Foods.

2020;9. doi:10.3390/foods9020216

15.   Ceyhun Sezgin A, Sanlier N. A new generation plant for the conventional cuisine: Quinoa

(*Chenopodium quinoa* Willd.). Trends Food Sci Technol. 2019;86: 51–58.

doi:10.1016/J.TIFS.2019.02.039

16.   Dakhili S, Abdolalizadeh L, Hosseini SM, Shojaee-Aliabadi S, Mirmoghtadaie L. Quinoa

protein: Composition, structure and functional properties. Food Chemistry. Elsevier Ltd;

2019. doi:10.1016/j.foodchem.2019.125161

17.   Noulas C, Tziouvalekas M, Vlachostergios D, Baxevanos D, Karyotis T, Iliadis C.

Adaptation, Agronomic Potential, and Current Perspectives of Quinoa Under

Mediterranean Conditions: Case Studies from the Lowlands of Central Greece. Commun

Soil Sci Plant Anal. 2017;48: 2612–2629. doi:10.1080/00103624.2017.1416129

18.   Sellami MH, Pulvento C, Lavini A. Agronomic practices and performances of quinoa

under field conditions: A systematic review. Plants. 2021;10: 1–20.

doi:10.3390/plants10010072

19.   Asher A, Dagan R, Galili S, Rubinovich L. Effect of Row Spacing on Quinoa

(*Chenopodium quinoa*) Growth, Yield, and Grain Quality under a Mediterranean Climate.

Agriculture. 2022;12: 1298. doi:10.3390/agriculture12091298

20.   Walters H, Carpenter-Boggs L, Desta K, Yan L, Matanguihan J, Murphy K. Effect of

irrigation, intercrop, and cultivar on agronomic and nutritional characteristics of quinoa.

Agroecology and Sustainable Food Systems. 2016;40: 783–803.

doi:10.1080/21683565.2016.1177805

21.   Wang N, Wang F, Shock CC, Meng C, Qiao L. Effects of management practices on

quinoa growth, seed yield, and quality. Agronomy. 2020;10.

doi:10.3390/agronomy10030445

22.   Asher A, Galili S, Whitney T, Rubinovich L. The potential of quinoa (*Chenopodium*

quinoa) cultivation in Israel as a dual-purpose crop for grain production and livestock

feed. Sci Hortic. 2020;272: 109534. doi:10.1016/j.scienta.2020.109534

23.   Filik G. Biodegradability of quinoa stalks: The potential of quinoa stalks as a forage

source or as biomass for energy production. Fuel. 2020;266: 117064.

doi:10.1016/j.fuel.2020.117064

24.   Matías J, Cruz V, Reguera M. Heat Stress Impact on Yield and Composition of Quinoa

Straw under Mediterranean Field Conditions. Plants. 2021;10: 955.

doi:10.3390/PLANTS10050955

25.   Ebeid HM, Kholif AE, El-Bordeny N, Chrenkova M, Mlynekova Z, Hansen HH. Nutritive

value of quinoa (*Chenopodium quinoa*) as a feed for ruminants: in sacco degradability and

in vitro gas production. Environmental Science and Pollution Research. 2022;29: 35241–

35252. doi:10.1007/s11356-022-18698-x

26.   Ramos N, Cruz AM. Evaluation of seven seasonal crops for forage production during the

dry season in Cuba. Cuban Journal of Agricultural Science. 2002;36: 271–276.

27.   Camaggio G, Amicarelli V. The ancient crop of quinoa for world food security. In:

Mis`niakiewicz M, Popek S, editors. Future Trends and Challenges in the Food Sector.

Cracow, Poland: Polish Society of Commodity Science; 2014.

28.   Pathan S, Siddiqui RA. Nutritional Composition and Bioactive Components in Quinoa

(*Chenopodium quinoa* Willd.) Greens: A Review. Nutrients. 2022;14: 1–12.

doi:10.3390/nu14030558

29.   Adamczewska-Sowińska K, Sowiński J, Jama-Rodzeńska A. The effect of sowing date

and harvest time on leafy greens of quinoa (*Chenopodium quinoa*) willd.) yield and

selected nutritional parameters. Agriculture (Switzerland). 2021;11.

doi:10.3390/agriculture11050405

30.   El-Naggar A, Hussin S, Abd El-Samad E, Eisa S. Quinoa As a New Leafy Vegetable Crop

in Egypt. Arab Universities Journal of Agricultural Sciences. 2018;26: 745–753.

doi:10.21608/ajs.2018.16007

31.   Vazquez-Luna A, Cortés VP, Carmona FF, Díaz-Sobac R. Quinoa leaf as a nutritional

alternative. Cienc Investig Agrar. 2019;46: 137–143. doi:10.7764/rcia.v46i2.2098

32. Pathan S, Eivazi F, Valliyodan B, Paul K, Ndunguru G, Clark K. Nutritional Composition

of the Green Leaves of Quinoa (*Chenopodium quinoa* Willd.). J Food Res. 2019;8: 55.

doi:10.5539/jfr.v8n6p55

33. Gawlik-Dziki U, Swieca M, Sulkowski M, Dziki D, Baraniak B, Czyz J. Antioxidant and

anticancer activities of *Chenopodium quinoa* leaves extracts - In vitro study. Food and

Chemical Toxicology. 2013;57: 154–160. doi:10.1016/j.fct.2013.03.023

34. Stoleru V, Jacobsen SE, Vitanescu M, Jitareanu G, Butnariu M, Munteanu N, et al.

Nutritional and antinutritional compounds in leaves of quinoa. Food Biosci. 2022;45.

doi:10.1016/j.fbio.2021.101494

35. Lim JG, Park HM, Yoon KS. Analysis of saponin composition and comparison of the

antioxidant activity of various parts of the quinoa plant (*Chenopodium quinoa* Willd.).

Food Sci Nutr. 2020;8: 694–702. doi:10.1002/fsn3.1358

36. Abd El-Samad EH, Hussin SA, El-Naggar AM, El-Bordeny NE, Eisa SS. The potential

use of quinoa as a new non-traditional leafy vegetable crop. Bioscience Research.

2018;15: 3387–3403. Available: www.isisn.org

37. WHO. Protein and amino acid requirements in human nutrition: report of a joint

WHO/FAO/UNU expert consultation. 2007.

38. Bhargava A, Sudhir S, Deepak O. Effect of sowing dates and row spacings on yield and

quality components of quinoa (*Chenopodium quinoa*) leaves. Indian Journal of

Agricultural Sciences. 2007;77: 748–751. Available:

https://www.researchgate.net/publication/215713006

39. Woodard KR, Prine GM. Forage Yield and Nutritive Value of Elephantgrass as Affected

by Harvest Frequency and Genotype (AJ). Agron J. 1991;83: 541–546.

40. Robins JG, Bauchan GR, Brummer EC. Genetic mapping forage yield, plant height, and

regrowth at multiple harvests in tetraploid Alfalfa (Medicago sativa L.). Crop Sci.

2007;47: 11–18. doi:10.2135/cropsci2006.07.0447

41. Bantis F, Kaponas C, Charalambous C, Koukounaras A. Strategic successive harvesting of

rocket and spinach baby leaves enhanced their quality and production efficiency.

Agriculture (Switzerland). 2021;11. doi:10.3390/agriculture11050465

42. van Minh N, Hoang DT, van Loc N, Long NV. Effects of plant density on growth, yield

and seed quality of quinoa genotypes under rain-fed conditions on red basalt soil regions.

Aust J Crop Sci. 2020;14: 1977–1982. doi:10.21475/ajcs.20.14.12.2849

43. Spehar CR, Rocha JE da S. Effect of sowing density on plant growth and development of

quinoa, genotype 4.5, in the Brazilian Savannah Highlands. Bioscience Journal. 2009;25:

53–58.

44. Bellalou A, Daklo-Keren M, Abu Aklin W, Sokolskaya R, Rubinovich L, Asher A, et al.

Germination of *Chenopodium quinoa* cv. 'Mint Vanilla' seeds under different abiotic

stress conditions. Seed Science and Technology. 2022;50: 41–45.

doi:10.15258/sst.2022.50.1.05

45. Jacobsen S-E. The Worldwide Potential for Quinoa (*Chenopodium quinoa* Willd.). Food

Reviews International. 2003;19: 167–177. doi:10.1081/FRI-120018883

46.   Lionello P, Scarascia L. The relation between climate change in the Mediterranean region

568   and global warming. Reg Environ Change. 2018;18: 1481–1493. doi:10.1007/s10113-018-

569   1290-1

570   47.   Fuentes F, Bhargava A. Morphological Analysis of Quinoa Germplasm Grown Under

571   Lowland Desert Conditions. J Agron Crop Sci. 2011;197: 124–134. doi:10.1111/j.1439-

572   037X.2010.00445.x