Island demographics and trait associations in white-tailed deer

Brooklyn S. Cars[1,2], Camille Kessler[1], Eric A. Hoffman[3], Steeve D. Côté[4], Daniel Koelsch[5,6],

Aaron B.A Shafer[1,2]

[1]Environmental and Life Sciences Graduate Program, Trent University, 2140 East Bank Drive,

Peterborough, ON K9J 7B8, Canada

[2]Department of Forensics, Trent University, 2140 East Bank Drive, Peterborough, ON K9J 7B8,

Canada

[3]Department of Biology, University of Central Florida, 4000 Central Florida Blvd, Orlando, FL,

USA

[4]Département de Biologie and Centre d'Études Nordiques, Université Laval,

G1V 0A6, Québec, QC, Canada

[5]Fédération des chasseurs de Saint-Pierre et Miquelon. Saint-Pierre et Miquelon, France
16
17   *[6]Direction des Territoires de l'Alimentation et de la Mer, service Biodiversité. Saint-Pierre et*
18   *Miquelon, France.*
19
20

25   Corresponding author: Aaron Shafer (aaronshafer@trentu.ca)

26   **Word count:** 5,246

Abstract

When a population is isolated and composed of few individuals, genetic drift is the paramount evolutionary force that results in the loss of genetic diversity. Inbreeding might also occur, resulting in genomic regions that are identical by descent, manifesting as runs of homozygosity (ROHs) and the expression of recessive traits. Likewise, the genes underlying traits of interest can be revealed by comparing fixed SNPs and divergent haplotypes between affected and unaffected individuals. Populations of white-tailed deer (*Odocoileus virginianus*) on islands of Saint Pierre and Miquelon (SPM, France) have high incidences of leucism and malocclusions, both considered genetic defects; on the Florida Keys islands (USA) deer exhibit smaller body sizes, a polygenic trait. Here we aimed to reconstruct island demography and identify the genes associated with these traits in a pseudo case-control design. The two island populations showed reduced levels of genomic diversity and a build-up of deleterious mutations compared to mainland deer; there was also significant genome-wide divergence in Key deer. Key deer showed higher inbreeding levels, but not longer ROHs, consistent with long-term isolation. We identified multiple trait-related genes in ROHs including *LAMTOR2* which has links to pigmentation changes, and *NPVF* which is linked to craniofacial abnormalities. Our mixed approach of linking ROHs, fixed SNPs and haplotypes matched a high number (~50) of a-priori body size candidate genes in Key deer. This suite of biomarkers and candidate genes should prove useful for population monitoring, noting all three phenotypes show patterns consistent with a complex trait and non-Mendelian inheritance.

Introduction

Genomic architecture is the organized arrangement of genes and other functional elements underlying trait variation (Koonin 2009). Targeted population sampling combined with high-throughput sequencing can characterize the architecture and effect size of genomic variation underlying traits (Ceballos *et al.* 2018). Specifically, the genomes of individuals with and without traits of interest can be compared for the identification of divergent single nucleotide polymorphisms (SNPs) (Yang *et al.* 2008; Russell *et al.* 2012; Getmantseva *et al.* 2020). Reconstructing haplotypes is also useful for developing diagnostic tools and identifying candidate genes, with clear regions underlying body size variation in dogs detected for example (Parker *et al.* 2009). Large effect SNP and haplotypes, especially for polygenic traits, are typically found in low frequency (e.g. Yengo *et al.* 2022), and often require targeted phenotypic sampling to have enough power to detect in natural populations (e.g. Anderson *et al.* 2022).

The distribution of SNPs present in the genome is also useful for the identification of regions with reduced heterozygosity and segments of autozygosity (Kim *et al.* 2013); long rows of sequential homozygous sites known as runs of homozygosity (ROHs) can be used to map the genetic regions of recessive traits stemming from deleterious mutations (Hildebrandt *et al.* 2009). Runs of homozygosity result from the passing of identical haplotypes from one generation to the next as a result of mating between relatives, with the length of ROH corresponding to recency of the common ancestor (McQuillan *et al.* 2008; Kirin *et al.* 2010). Expression of recessive, often deleterious, traits due to homozygosity require ROHs to be present (Robinson *et al.* 2023), though purging might remove deleterious alleles (Wootton *et al.* 2023). A populations' demographic history, notably recent trends in effective size (N_e), also can correspond to ROH dynamics (Kardos *et al.* 2023) and estimates of ROH are now commonly included in

assessments of endangered species, with for example unmanaged scimitar-horned oryx (*Oryx*

dammah) populations manifesting significantly higher inbreeding (F_{ROH}) than those in captivity

(Humble *et al.* 2023). Therefore, detecting and mapping ROHs in a population can identify the

genomic impact of demographic change and map genes that are associated with recessive and

deleterious traits.

White-tailed deer (*Odocoileus virginianus*) are herbivorous, hooved ungulates that are

native to the western hemisphere, primarily North America, but their range has expanded

through human assistance to parts of Europe and various remote islands (Hewitt 2011; Villanova

et al. 2017; Fuller *et al.* 2019; Entfellner *et al.* 2021). Anthropogenic movement appears to have

impacted genetic load in white-tailed deer (Wootton *et al.* 2023), meaning the accumulation of

deleterious alleles, but overall, population differentiation (e.g. Cullingham *et al.* 2011) and

inbreeding across the species is low (F_{IS} <0.10; Hamlin *et al.* 2021). In some cases, introduced

island populations show high diversity (Fuller *et al.* 2019; Nelson *et al.* 2021), but ensuing island

dynamics have resulted in various phenotypic variants or anomalies in some deer populations.

For example, the native population of white-tailed deer that inhabit the Florida Keys - referred to

as Key deer - are significantly smaller compared to the white-tailed deer populations on the

mainland (Maffei *et al.* 1988; Folk and Klimstra 1991; Karns 2010). Island populations of deer

often experience reduced size and body mass from nutrition limitations (e.g. Anouk Simard *et al.*

2008; Bartareau 2019), but Key deer weigh less than the Florida mainland (Bartareau 2019) and

are considered the smallest subpspecies (Hewitt 2011). These patterns are consistent with the

Island rule of larger animals becoming smaller (Foster 1964; Benítez-López *et al.* 2021), but also

follow a general clinal decline in deer body sizes at southern latitudes (Wolverton *et al.* 2009).

Stature is generally a complex trait driven by many large and small effect genes (Kemper *et al.*

2012; Sidorenko *et al.* 2022) and Anderson *et al.*(2022) identified a suite of candidate genes

impacting body size in deer (See Supplemental File S1). Key deer are genetically differentiated

(Villanova *et al.* 2017), making disentangling drift from selection, and the Island rule,

particularly challenging without first reconstructing their demographic history. Mark-recapture

estimates place Key deer at 1,000 individuals (Villanova *et al.* 2017) and they are listed as

Endangered under the Endangered Species Act.

In contrast, the introduced population of white-tailed deer on the islands of Saint Pierre

and Miquelon (SPM) have relatively high rates of leucism and malocclusions, meaning incorrect

alignment of the maxillary bone and the mandible. This population stems from 14 animals being

introduced in the 1953; it is likely that the genes underlying these traits exhibit an increased

frequency due to their presence in a small founding population, though inbreeding has not been

assayed. While white-tailed deer are normally a uniform reddish-brown-grey with white fur

scattered on appendages, face, and rear, deer with leucism present a partial to entire lack of

pigmentation. Coat colour composition in ungulates is mainly determined by loci that control

melanocyte activity and distribution (Fontanesi *et al.* 2010). Mutations in the melanocortin

receptor locus (*MC1R*) in particular have been found to be major contributors to leucism (Haase

et al. 2013; Reiner *et al.* 2020). Brachygnathism and prognathism are both forms of

malocclusion (Faragalla 2000; Guan *et al.* 2015) and have been observed in both mule deer (*O.*

hemionus, Robinette and Aldous 1955) and white-tailed deer (Smits and Bubenik 1990).

Candidate genes have been identified for both leucism (Supplementary Table S1) and jaw

malocclusions (Supplementary Table S2); leucism is considered a recessive trait (Pereira *et al.*

2023) but malocclusion inheritance patterns are less clear (Cakan *et al.* 2012). Expression of both

traits in deer likely comes with a fitness cost (Wolff *et al.* 1993; Faragalla 2000; Reissmann and

Ludwig 2013).

The emergence of divergent phenotypes (Key deer) and traits with a fitness cost (SPM)

on islands present a pseudo case-control design allowing for characterizing the genetic

architecture of phenotypes. By leveraging these phenotypically distinct populations and

comparing to a large mainland white-tailed deer genomic data set (Kessler *et al.* 2023), we aim

to reconstruct each population's recent demographic history and genetic load, and test the

prediction that ROH and candidate genes are associated with expression of recessive traits in

SPM, while unique haplotypes with candidate body size genes underly the divergence of Key

deer.

Materials and Methods

Sampling and Sequencing

Phenotype information for leucism or malocclusions from SPM was obtained from local

managers and photos were examined to select individuals with one or more of the visible traits of

interest (see Supplemental Fig. S1). Twenty-six white-tailed deer tissue samples were then

selected from the two island populations of interest: SPM (France, but coastal Canada) and the

Florida Keys (United States) in addition to four samples from Anticosti Island, Quebec (Canada).

We leveraged available genome sequence data from four additional Key deer, one SPM sample

(normal phenotype), and 21 deer from the North American mainland (normal phenotypes) that

are part of a larger project (Fig. 1; PRJNA830519). Samples were provided by hunters

voluntarily, with some collected from road mortality. DNA extraction from the tissue samples

was carried out using the QIAGEN DNeasy Blood & Tissue Kit following the manufacturer's

instructions (*DNeasy® Blood & Tissue Handbook* 2020). We quantified the DNA concentrations

using the Invitrogen Qubit assay. Samples were sent to The Centre for Applied Genomics in

Toronto, Canada for low coverage (~4x) whole genome sequencing (WGS) on an Illumina

NovaSeq.

Data Preparation

The quality of the raw data libraries was examined using FastQC (v0.11.9; Andrews

2010) and regions of poor quality were removed using Trimmomatic (v0.39; Bolger, Lohse and

Usadel 2014). The trimmed sequence data was then re-examined using FastQC, mapped to the

annotated *O. virginianus* reference genome (GCA_023699985.1; Anderson *et al.* 2022) using

BWA-MEM (v0.7.17; Li 2013), and sorted using SAMtools sort (v1.15.1; Li *et al.* 2009). We

detected duplicated reads using Picard MarkDuplicates (v2.23.2; Broad Institute, GitHub

Repository 2019) and removed the duplicated reads using Sambamba (v0.8.2; Tarasov *et al.*

2015). Some samples had multiple sequence files and were merged before removing the

duplicated reads using SAMtools merge. To ensure that no errors had been made throughout the

process and that we had the desired coverage and depth to continue analysis we quantified

metrics with SAMtools coverage and depth, and sambamba flagstat.

SNPs were called using ANGSD (v0.939; Korneliussen, Albrechtsen and Nielsen 2014)

following the GATK model (-gl 2) to estimate genotype likelihoods (GLs) in a beagle genotype

likelihood format (-doGlf 2). The genotypes were printed (-doGeno 4) and required SNPs with a

minimum p-value of 1e−6 (-SNP_pval 1e-6), a minimum mapQ quality of 20 (-minMapQ 20),

and a minimum base quality of 20 (-minQ 20). Further, we assumed that the major and minor

alleles were known (-domaf 1) and that they can be inferred directly from likelihoods (-

doMajorMinor 1).

Estimating Subpopulation Diversity and Demography

We estimated the nucleotide diversity (π) among each subpopulation of our dataset using

VCFtools (v0.1.16; Danecek *et al.* 2011). We calculated π using a 10 kb step size in between

windows (-window-pi 10000). Tajima's *D* for each subpopulation was in windows of 10 kb (-

TajimaD 10000) using only bi-allelic sites. We estimated fixation indices (F_{ST}) between

populations using window sizes of 10 kb (-fst-window-size 10000). Using plink (v1.9) we

generated a PCA based on the variance-standardized relationship matrix for the visualization of

population stratification of the subpopulations. All plots were done in R version 4.2.2 (R Core

Team 2022). To estimate the historical N_e within the last 200 generations, we used the program

GONE that uses the linkage disequilibrium information for markers across the genome (Santiago

et al. 2020). We selected a maximum of 30,000 SNPs per chromosome, 0.5 cM/Mb (Kessler *et*

al. 2023), an *hc* of 0.04 (maximum recombination rate), and applied the Haldane correction for

genetic distance; we repeated the analysis 500 times. For a contemporary point estimate of N_e we

ran *currentNe* (Santiago *et al. 2*023) that extends the linkage disequilibrium approach but

includes aspects of the species' mating system. We ran *currentNe* with a random sample of

1,000,000 SNPs per island population with the mating structure parameter (*k*) inferred

empirically from the data.

Calling Runs of Homozygosity (ROHs)

Scaffolds making up the N90 were extracted using tabix (v.0.2.6.; Li 2011). We created

tped and tfam files using plink (v2.0) and identified ROH genotypes using the plink window-

based approach (v1.9). Data were lightly pruned for linkage disequilibrium ($r^2 > 0.90$; Danecek

et al. 2021). We used a sliding window size of 100 kb (-homozyg-kb 100), a minimum density of

1 SNP per 50 kb (-homozyg-density 50) and required a minimum of 50 SNPs to be considered a

ROH (-homozyg-snp 50). We used the default scanning window size of 50 SNPs (-homozyg-

window-snp 50), the default SNP hit rate of 0.05 (-homozyg-window-threshold 0.05) and

allowed up to 3 heterozygote sites (-homozyg-window-het 3) as recommended for low coverage

data (Ceballos *et al.* 2018) and 10 missing calls (-homozyg-window-missing 10) for each 100 kb

window within called ROHs. No minor allele frequency cut-off was applied (Meyermans et al.

2020). A 1,000 bp gap between variant sites (-homozyg-gap 1000) was required for two ROHs to

be considered separate and we grouped the samples in phenotypic pools to identify overlapping,

and possibly matching ROH segments (-homozyg-group). The percentage of the genome covered

by ROHs (F_{ROH}) per individual was estimated by summing the total distance spanned by the

ROH segments and dividing this by the total length of the N90 scaffolds.

We identified ROHs unique to each phenotypic group and found in >50% individuals,

meaning none of the SPM normal, mainland or Anticosti Island comparators presented this

ROH. Key deer we increased the required to >80% of samples have the ROH given the overall

increased divergence of the population. The genomic coordinates of the ROHs were identified

and matched to the *O. virginianus* gene annotation file (Anderson *et al.* 2022) and the biological

function information of the discovered genes within ROH were identified through Uniprot

(Pundir *et al.* 2016) and our candidate gene literature review (Supplementary Tables S1, S2, File

S1). Any gene without an entry ID was screened on NCBI blast to identify a putative function.

We inferred the approximate age of the identity by descent (IBD) event (no. of generations or *g*)

based on ROH length, where $g = 100/(2r\text{L})$, with *r* being the recombination rate of 0.50 cM/Mbp

(Kessler *et al.* 2023) length (L) of ROH in Mbp (Browning and Browning 2015; Kardos *et al.*

2017). To quantify whether our ROH-phenotype groupings could have occurred by chance, we

permutated mainland, Anticosti and SPM individuals IDs and re-ran the analysis for leucism

groupings 100 times.

Fixed SNPs and haplotypes

We identified SNPs that were fixed in the phenotypic group, with the alternate allele

found in >50% of wild-type samples (i.e. south mainland vs Key deer, SPM normal vs. leucism

and malocclusions; Fig. 1). Using VCFtools (v0.1.16) we extracted lists of SNPs fixed in the

local phenotypic group, requiring the alternate allele to be fixed in the mainland comparator. For

SPM and the Key deer, we extracted and assayed all fixed SNPs. We complemented this with a

formal haplotype analysis; here samples were first phased using SHAPEIT2 (Delaneau *et al.*

2013) then analyzed through a likelihood ratio test HapLRT (Garrison *et al.* 2022) to model the

haplotype length distributions of the target (island) and background (comparator) populations;

note the entire island sample of SPM was used to increase power. Here a positive sign from

HapLRT reflect longer haplotypes in the target population; haplotype blocks were then defined

as any region having a positive sign with successive p values <0.01 spanning 100 kb or longer

and separated by at least 5 kb from other blocks. All fixed SNPs and haplotype blocks were

matched to the *O. virginianus* gene annotation file (Anderson *et al.* 2022) and we identified

genes within haplotype blocks or within 50 kb of the fixed SNP grouping. Lastly, for all

phenotypic and mainland groupings we used SNPeff (Cingolani *et al.* 2012) to estimate and

summarize the putative impact of all present SNPs; we focused on genetic load by measuring

homozygote missense and loss of function (LOF) mutation, the latter being SNPs that resulted in

a premature stop codon.

Results

Sampling, Sequencing and Data Preparation

All deer samples were sequenced to ~4x coverage; this included 16 individuals from

SPM of which four presented with leucism and seven with malocclusions, as well as 10 Key deer

samples (smaller stature size). As comparators, we used 21 samples from the North American

mainland split into North and South groupings and four from Anticosti Island, all of which were

normal phenotypes (Fig. 1). We identified a total of 100,544,312 SNPs in all individuals and the

analyses were performed exclusively on the N90 scaffolds that harboured 90,993,235 SNPs.

Subpopulation Diversity

Tajima's D for SPM was the highest followed by Key deer that had the lowest value of π

(Table 1). Mainland and Anticosti deer showed negative Tajima's D values. The fixation index

(F_{ST}) and PCA showed high levels of genetic differentiation between the Key deer and every

other subpopulation; SPM formed its own cluster, while Anticosti Island was embedded within

the North Mainland samples (Supplementary Table S3; Supplementary Fig. S2). Recent

demographic trends revealed a stable long-term low N_e of Key deer, with a large crash in SPM

that predates the translocation (Fig. 2). Point estimates of current N_e are low for both

populations: N_e of SPM (108; 90% confidence interval [CI] 64 –180) and Key deer (287; 90% CI

95–870).

Distribution of ROH among phenotypes

The populations with the highest F_{ROH} were the Key deer followed bhy SPM, while the

populations from the North American Mainland and Anticosti Island all had a relatively low

F_{ROH} (Fig. 3A). The range of average ROH lengths within the Key deer individuals was shorter

than the other populations with ROH lengths of 136.8 kb – 192.1 kb (Table 1); this range of

ROH likely corresponds to >500 generations to the most recent common ancestor. Additionally,

all subpopulations had majority of their ROHs in the 100-300 kb size range (Fig. 3B), which

corresponds to the greater total ROH length of this size range as well (Fig. 3C).

In the pools for all four SPM white-tailed deer with leucism, we found 39 shared ROHs.

Of those, one was unique to the individuals with leucism, and it contained 19 genes (Table 2); no

a-priori candidate genes for leucism were detected (Supplementary Table S1). ROHs detected in

animals with leucism and absent from the comparators contained multiple genes, but notably

LAMTOR2 (Endosomal adaptor protein p14) (Table 2). A total of 21 ROHs were found to be

shared between all SPM white-tailed deer with malocclusions, but none were unique to this

phenotype. One ROH was unique to 4/7 deer with this phenotype, and only harbored one gene

(*NPVF*; Table 2). Lastly, 18 ROHs were shared among the smaller-stature Key deer. Of those,

one was unique to the Key deer, and it contained 1 gene (*CHMP3*); which was also found within

the Key deer's fixed SNPs (Table 2; Supplementary File S2). Within a pool of all 10 Key deer

and one SPM deer, there were two genes that matched the Anderson *et al.* (2022; Supplementary

File S1) candidate gene table: *ART1* and *ART5* (Table 2). In addition, nine out of 10 of the Key

deer shared eight ROHs that were unique to them, they collectively harboured 27 genes and had

one gene that matched the Anderson *et al.*(2022) candidate gene table: *RPL17* (Table 2). Pools

were unlikely to have arisen by chance (Table S4).

Fixed SNPs and haplotypes

For each fixed SNP we screened 50 kb flanking regions on either end to identify genes

that were in close proximity; here we identified 72,291 fixed SNPs for Key deer population that

collectively encompassed 9,081 genes. Of these, 20 overlapped with the ROHs and 427 matched

the Anderson *et al.* (2022) candidate gene table. There were 159 fixed SNPs within the SPM

leucism group that contained 65 genes, five fixed SNPs within the SPM malocclusion group that

contained two genes and four fixed SNPs within the Anticosti population that contained one

gene. There was no overlap for fixed SNPs and ROHs or candidate genes in either SPM

phenotypic group. Fixed SNPs and ROH overlapped SNPs detected with a standard genome-

wide association approach (Figure S4).

There were 695 haplotype blocks detected in Key deer, which encompassed 455 genes.

One haplotype overlapped with the ROHs for Key deer, 384 overlapped with the fixed SNPs for

Key deer, and 50 genes matched the candidate gene table, and notably included the gene *IGF1R*

(Supplementary File S2). For the SPM population, we identified four haplotype blocks; these

blocks harboured two genes, neither of which overlapped with the ROHs in SPM deer. The

absolute distribution of variants, including deleterious, were similar amongst populations (Table

S5). Homozygote LOF and missense mutations were notably elevated in Key deer (Fig. 4); there

was a modest increase in SPM, specifically relative to Anticosti and mainland deer (Fig. 4).

Discussion

Island populations are prone to drift; further, isolated populations composed of a small

number of individuals are more likely to suffer from inbreeding, accumulate deleterious alleles,

and manifest undesirable traits (Soule 1973; Villanova *et al.* 2017). Small populations, however,

might not show inbreeding depression (e.g. Robinson *et al*. 2018), with purging occurring under

the right demographic scenarios (Wootton *et al. 2*023). The presence of presumed deleterious

phenotypes and the corresponding genetic data (e.g. Fig. 4), suggest that purging has not

occurred in our focal populations. SPM deer have a relatively high frequency of malocclusions

and leucism, and Key deer are significantly smaller than their mainland counterparts (Folk and

Klimstra 1991). SPM samples clustered adjacent to northern mainland deer due to their recent

translocation, while the Key deer are significantly diverged (see Supplementary Fig. S2). The

Key deer became isolated approximately 8,000 years ago (Lazell 1989) while SPM were

introduced in 1953; consequently, the confounding effects of drift are much less of a factor in the

gene-trait associations detected in SPM compared to the Key deer. Still, clear patterns emerged

that not only impact our understanding of the genetic architecture of traits in wild populations,

but also the build-up of ROH and genetic load in light of recent demographic change.

A tale of two island populations

Key deer are an endangered species under the US Federal Endangered Species Act.

Genetic mark recapture studies suggests a population of ~1,000 individuals, with an N_e of only

11 (Villanova *et al.* 2017). Our genome wide estimates are higher, with a contemporary point

estimate of ~300. Here we suspect the cross-species amplification and small number of

microsatellites resulted in a downward bias estimate in Villanova *et al.* (2017) compared to

genome wide estimates based on linkage disequilibrium. Values of π, which are reflective of

long-term or coalescent N_e (Peart *et al.* 2020) are lowest in Key deer compared to all

comparators; the coalescent N_e / N_C ratio does appear to be linked to conservation status and

extinction risk (Peart *et al.* 2020; Dedato *et al.* 2022; Wilder *et al.* 2023); solving for $\pi = 4N_e\mu$

and with N_C estimates of Villanova *et al.*(2017) and μ of Kessler *et al.*(2023), would generate a

ratio coalescent N_e / N_C ratio >1 in Key deer, thereby supporting the endangered species

designation.

Key deer were almost hunted to extinction in the 1950s (U.S. Fish and Wildlife Service

1999); despite this, the N_e has been stable for >200 generations (Fig. 2A) and no signal of

overharvest is detected, as seen in mainland deer (Kessler and Shafer 2023). Collectively, this

suggests the moderate buildup of ROH (Fig. 3A), and genetic load (Fig. 4) is due to the historical

demographic bottleneck, and not to contemporary inbreeding or overharvest. Indeed, while Key

deer have the highest mean F_{ROH} of white-tailed deer examined (Table 1), the average length of

ROH is not longer, with the number of generations to the common ancestor of the ROH being

>500. And while there is an elevated genetic load, with values higher than relevant comparators

(Humble *et al.* 2023), there does not appear to be a link to population viability given the general

patterns seen in deer (Kessler and Shafer 2023) and robust population growth of white-tailed

deer in general (Webb 2018). The relatively high N_e in Key deer might also have limited

purging, as seen in ungulates with much lower N_e (e.g. mountain goats; Wootton *et al. 2023;*

Martchenko and Shafer 2023). Scandinavian wolves, for example, show most ROH due to recent

ancestry but also high variance in estimates with F_{ROH} ranging from 0.01 to 0.54 (Kardos *et al.*

*2*018). Accordingly, deleterious alleles have also become fixed (Smeds and Ellegren 2023). Here

the demographic collapse in wolves was considerably worse than Key deer, and recovery (or

growth) much slower which would explain these differences. Collectively, these data support

Villanova *et al.* (2017) who predicted long-term persistence of the Key deer population.

The SPM Island population originated from 14 white-tailed deer; in contrast 220 white-

tailed deer were introduced to the much larger Anticosti Island (7,953 km²) off the coast of

Quebec in 1896 (Fuller *et al.* 2019). Accordingly, diversity values (Table 1) reflect this

difference in founding number and island size, with Anticosti Island largely indistinguishable

from the mainland. Tajima's *D* values were also highest on SPM, suggesting a recent loss of rare

alleles due to this founding effect. The crash in recent N_e timepoints (Fig. 2B) predates the

translocation event; as genomes carry a historical record of the species, this signal is a carry-over

from the source population and colonial overharvest on the mainland (Kessler and Shafer 2023).

There does, however, appear to be a slight build-up of genetic load from the translocation event

(Fig. 4). In island simulations, Robinson *et al.* (2018) showed that strongly deleterious alleles

were purged, while moderate deleterious alleles accumulated. It is therefore conceivable that the

genetic load seen in both Key deer and SPM consists primarily of moderate to low effect

deleterious mutations allowing their persistence; temporal monitoring of SPM will help shed

light on this important evolutionary process given the recency of the founding event.

Inbreeding and isolation lead to detecting candidate genes

The Islands of SPM have a relatively high number of individuals with leucism and

malocclusions. None of the candidate genes for leucism (Hauswirth *et al.* 2012; Haase *et al.*

2013; Reiner *et al.* 2020) were found within our ROH segments. There were 19 new candidate

genes harboured by the ROHs unique to all leucism individuals, though none had a clear

biological link. Screening ROHs shared by 3/4 animals in the leucism pool identified the gene

LAMTOR2 (Endosomal adaptor protein p14) which has links to hypopigmentation (Bohn *et al.*

2007). Hypopigmentation is caused by the hindrance of melanocytes which are responsible for

the pigmentation of hair and skin (Dessinioti *et al.* 2009). *LAMTOR2* interacts with the *MITF*

transcription factor which regulates melanocyte activity and development; as leucism is known

to be caused by mutations in genes that are associated with melanocytes *MITF* could be the

leucism pathway in white-tailed deer. Our results also hint at non-Mendelian patterns of

inheritance as suggested in other vertebrates (Edelaar *et al.* 2011; Jablonski *et al.* 2017). No

ROHs were found to be unique to all of the white-tailed deer with malocclusions; however, a

unique ROH segment that was shared by 4/7 white-tailed deer with malocclusions contained the

gene *NPVF* . In humans and mice enhancers adjacent to the *NPVF* are implicated in craniofacial

abnormalities (Wildermann *et al.* 2022; Yu *et al.* 2023). While this region certainly appears

functionally relevant, not all animals with maloclussions have this ROH, suggesting again a

polygenic trait or incomplete penetrance (Stiles and Luke 1953). Kardos *et al.* (2018) suggested

that areas with low ROH abundance are likely to contain loci with deleterious recessive alleles

that contribute disproportionately to inbreeding depression. Thus, we cannot rule out,

homozygous deleterious variants that are not captured by our ROH quantification.

Stature size is a complex trait influenced by many potential genes that have both large

and small effects. Anderson *et al.* (2022) sequenced the genomes of extreme phenotypes in deer,

specifically the tail ends of the distribution for body size controlling for age and created an

extensive list of candidate genes (Supplemental File S1). Our study found thousands of fixed

SNPs, clearly due to drift and long-term isolation. Still, close to 50 genes that overlapped

previous genome-association work (Anderson *et al. 2*022) fell on both haplotypes and fixed

SNPs suggesting these are indeed likely linked to body size. Body size is clearly polygenic and it

would be tempting to create a story of how these genes influence body size (Pavlidis *et al.* 2012),

so we simply note that one haplotype block contained the gene *IGF1R* which has been seen to

cause a reduced size in dogs (*Canis familiaris*, Hoopes *et al.* 2012), which in our system is

consistent with the island rule.

Conclusion

Island white-tailed deer populations have arisen from a mix of human-assisted and

natural colonization or isolation. Demographic assessment shows a stable, but low *Ne* in both

SPM and Key deer, an accumulation of genetic load, but generally low levels of ROH. Our

results could be useful for wildlife management concerned with small and isolated populations

that may be experiencing a loss of genetic diversity due to inbreeding. Notably, combining

Tajima's *D* with demographic models, and ROH with genetic load metrics, helps elucidate the

role of demographic change in shaping population genomic health. Interestingly, the island

populations do not have longer ROH; thus, expression of putative recessive traits on SPM is not

due to recent inbreeding, rather chance autozygosity due to a high frequency of the gene

variant(s) in the founding population. Collectively, this supports more detailed demographic

models of Key suggesting they have been isolated for a considerable length of time (Kessler *et*

al. 2023), with this study showing no evidence of recent inbreeding, despite an increased genetic

load.

Our mixed approach of ROH, fixed SNPs and haplotypes did pull out a-priori candidate

genes and discovered others. Storytelling of gene ontologies is common and leads to false

positives (Pavlidis *et al.* 2012), but our findings support all three traits not having a simple

architecture, and identified promising genes involved with leucism (*LAMTOR2*), malocclusions

(*NPVF*), and stature size (*IGF1R*) in white-tailed deer. Despite the divergence in Key deer

limiting some approaches (i.e. fixed SNPs were associated with almost half the total genes

suggesting a large impact of genetic drift), both haplotype and ROH inferences overlapped with

a large number of candidate genes allowing us to weed out background noise. Given the

relatively small sample sizes, the validation of autozygosity segments and quantification of effect

sizes in independent samples and mouse models will be required to validate these genes.

Acknowledgements

Participation from hunters, managers and community members is essential to this research. We

want to thank the community of St-Pierre and Miquelon that provided samples. Additional

Florida Key samples were provided by George Zaragoza; the remaining providers are found

listed in (Kessler *et al.* 2023). BC was supported by a Natural Sciences and Engineering

Research Council (NSERC) Undergraduate Student Research Award and the Trent Summer

Work Experience Program. This work was supported by NSERC Discovery Grant (ABAS;

SDC); ComputeCanada Resources for Research Groups (ABAS); Canadian Foundation for

Innovation (ABAS): John R. Evans Leaders Fund (ABAS) and the Ontario Early Researcher

Award (ABAS). We thank four anonymous referees and the Editor for valuable comments. Trent

University is located on the traditional territory of the Michi Saagiig Anishnaabeg.

Author contribution statement

BC and ABAS conceived the study, BC, CK, EH, SDC, DK, ABAS collected the samples, BC

and CK performed the molecular laboratory work and the bioinformatic analyses with

contribution from ABAS. BC and ABAS wrote the initial draft of the manuscript, all authors

reviewed and commented on the manuscript.

Conflict of Interest

Authors declare no competing interests.
424

Data archiving

Raw reads for the 25 deer additional individuals were deposited on the NCBI Sequence Read

Archive (Accession number PRJNA830519). All scripts are available on GitLab

(https://gitlab.com/WiDGeT_TrentU/undergrad-theses/-/tree/master/%20Cars_2023)

Ethics approval

All experiments have been conducted as per the guidelines of the Trent University Research

Ethics Board. No approval of research ethics committees was required to accomplish the goals of

this study because sequencing work was conducted on samples donated by hunters or collected

as roadkill.

435
References

Anderson SJ, Côté SD, Richard JH, Shafer ABA (2022). Genomic architecture of phenotypic

extremes in a wild cervid. *BMC Genomics* **23**: 126.

Andrews S (2010). Babraham Bioinformatics - FastQC A Quality Control tool for High

Throughput Sequence Data.

Anouk Simard M, Côté SD, Weladji RB, Huot J (2008). Feedback effects of chronic browsing

on life-history traits of a large herbivore. Journal of Animal Ecology 77: 678–686.

Bartareau TM (2019). Estimating body mass of Florida white-tailed deer from standard age and

morphometric measurements. Wildl Res 46: 334–342.

Benítez-López A, Santini L, Gallego-Zamorano J, Milá B, Walkden P, Huijbregts MAJ, *et*

al.(2021). The island rule explains consistent patterns of body size evolution in terrestrial

vertebrates. *Nat Ecol Evol* **5**: 768–786.

Bohn G, Allroth A, Brandes G, Thiel J, Glocker E, Schaffer AA, *et al.*(2007). A novel human

primary immunodeficiency syndrome caused by deficiency of the endosomal adaptor

protein p14. *Nat Med* **13**: 38–46.

Bolger AM, Lohse M, Usadel B (2014). USADELLAB.org - Trimmomatic: A flexible read

trimming tool for Illumina NGS data.

Broad Institute, GitHub Repository (2019). Picard Tools.

Browning SR, Browning BL (2015). Accurate Non-parametric Estimation of Recent Effective

Population Size from Segments of Identity by Descent. *Am J Hum Genet* **97**: 404–418.

Cakan DG, Ulkur F, Taner T (Uğur) (2012). The genetic basis of facial skeletal characteristics

and its relation with orthodontics. *Eur J Dent* **6**: 340–345.

Ceballos FC, Hazelhurst S, Ramsay M (2018). Assessing runs of Homozygosity: a comparison

of SNP Array and whole genome sequence low coverage data. BMC Genomics 19: 106.

Ceballos FC, Joshi PK, Clark DW, Ramsay M, Wilson JF (2018). Runs of homozygosity:

windows into population history and trait architecture. *Nat Rev Genet* **19**: 220–235.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, *et al.*(2012). A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly*

(Austin) **6**: 80–92.

Cullingham CI, Merrill EH, Pybus MJ, Bollinger TK, Wilson GA, Coltman DW (2011). Broad

and fine-scale genetic analysis of white-tailed deer populations: estimating the relative

risk of chronic wasting disease spread. Evolutionary Applications 4: 116–131.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, *et al.*(2011). The variant

call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Dedato MN, Robert C, Taillon J, Shafer ABA, Côté SD (2022). Demographic history and

conservation genomics of caribou (Rangifer tarandus) in Québec. *Evol Appl* **15**: 2043–

2053.

Delaneau O, Howie B, Cox AJ, Zagury J-F, Marchini J (2013). Haplotype Estimation Using

Sequencing Reads. *Am J Hum Genet* **93**: 687–696.

Dessinioti C, Stratigos AJ, Rigopoulos D, Katsambas AD (2009). A review of genetic disorders

of hypopigmentation: lessons learned from the biology of melanocytes. *Exp Dermatol* **18**:

741–749.

DNeasy® Blood & Tissue Handbook (2020). Qiagen.

Edelaar P, Donazar J, Soriano M, Santillan MA, Gonzalez Zevallos DR, Garcia Borboroglu JP,

et al.(2011). Apparent selective advantage of leucism in a coastal population of Southern

caracaras (Falconidae).

Entfellner J-BD, Muchadeyi FC, Mpolya EA, Chebii VJ (2021). Genomics of Adaptations in

Ungulates. *Animals* **11**.

Faragalla (2000). Maxillary Brachygnathism in an Arab Mare. *J Vet Dent* **17**: 148–149.

Folk MJ, Klimstra WD (1991). Antlers of White-tailed Deer (Odocoieus virginianus) from

Insular and Mainland Florida. **19**: 97–105.

Fontanesi L, Martelli PL, Beretti F, Riggio V, Dall'Olio S, Colombo M, *et al.*(2010). An initial

comparative map of copy number variations in the goat (Capra hircus) genome. *BMC*

Genomics **11**: 639.

Foster JB (1964). Evolution of Mammals on Islands. *Nature* **202**: 234–235.

Fuller J, Ferchaud A, Laporte M, Le Luyer J, Davis TB, Côté SD, *et al.*(2019). Absence of

founder effect and evidence for adaptive divergence in a recently introduced insular

population of white-tailed deer (Odocoileus virginianus). *Mol Ecol* **29**: 86–104.

Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P (2022). A spectrum of free

software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim

and slivar. *PLOS Comput Biol* **18**: e1009123.

Getmantseva LV, Bakoev SY, Shevtsova VS, Kolosov AY, Bakoev NF, Kolosova MA (2020).

Assessing the Effect of SNPs on Litter Traits in Pigs. *Scientifica* **2020**: e5243689.

Guan X, Song Y, Ott J, Zhang Y, Li C, Xin T, *et al.*(2015). The ADAMTS1 Gene Is Associated

with Familial Mandibular Prognathism. *J Dent Res* **94**: 1196–1201.

Haase B, Signer-Hasler H, Binns MM, Obexer-Ruff G, Hauswirth R, Bellone RR, *et al.*(2013).

Accumulating Mutations in Series of Haplotypes at the KIT and MITF Loci Are Major

Determinants of White Markings in Franches-Montagnes Horses. *Plos One* **8**: e75071-

e75071.

Hamlin BC, Meredith EP, Rodzen J, Strand JM (2021). OdoPlex: An STR multiplex panel

optimized and validated for forensic identification and sex determination of North

American mule deer (Odocoileus hemionus) and white-tailed deer (Odocoileus

virginianus). Forensic Science International: Animals and Environments 1: 100026.

Hauswirth R, Haase B, Blatter M, Brooks SA, Burger D, Drögemüller C, *et al.*(2012). Mutations

in MITF and PAX3 Cause "Splashed White" and Other White Spotting Phenotypes in

Horses. *PLOS Genet* **8**: e1002653.

Hewitt D (2011). *Biology and management of white-tailed deer*. CRC Press: Boca Raton, FL.

Hildebrandt F, Heeringa SF, Ruschendorf F, Attanasio M, Nurnberg G, Becker C, *et al.*(2009). A

systematic approach to mapping recessive disease genes in individuals from outbred

populations. *PLoS Genet* **5**.

Hoopes BC, Rimbault M, Liebers D, Ostrander EA, Sutter NB (2012). The insulin-like growth

factor 1 receptor (IGF1R) contributes to reduced size in dogs. *Mamm Genome Off J Int*

Mamm Genome Soc **23**: 780–790.

Humble E, Stoffel MA, Dicks K, Ball AD, Gooley RM, Chuven J, *et al.*(2023). Conservation

management strategy impacts inbreeding and mutation load in scimitar-horned oryx.

Proc Natl Acad Sci **120**: e2210756120.

Jablonski D, Grula D, Christophoryová J (2017). Unusual colour and pattern variation of Lacerta

agilis (Squamata: Lacertidae) recorded from Central Europe. *Biharean Biol* **11**: 126–128.

Kardos M, Qvarnström A, Ellegren H (2017). Inferring Individual Inbreeding and Demographic

History from Segments of Identity by Descent in Ficedula Flycatcher Genome

Sequences. *Genetics* **205**: 1319–1334.

Kardos M, Åkesson M, Fountain T, Flagstad Ø, Liberg O, Olason P, *et al.*(2018). Genomic

consequences of intensive inbreeding in an isolated wolf population. Nat Ecol Evol 2:

124–131.

Kardos M, Zhang Y, Parsons KM, A Y, Kang H, Xu X, *et al.*(2023). Inbreeding depression

explains killer whale population dynamics. *Nat Ecol Evol* **7**: 675–686.

Karns G (2010). Understanding White-Tailed Deer: Florida and the Southeast by William M.

Giuliano; Elina Garrison; Brandon J. Schad. *J Wildl Manag* **74**: 1935–1936.

Kemper KE, Visscher PM, Goddard ME (2012). Genetic architecture of body size in mammals.

Genome Biol **13**: 244.

Kessler C, Shafer ABA (2023). Genomic analyses capture the human-induced demographic

collapse and recovery in a wide-ranging cervid. bioRxiv: 2023.07.19.549627.

Kessler C, Wootton E, Shafer ABA (2023). Speciation without gene-flow in hybridizing deer.

Mol Ecol **32**: 1117–1132.

Kim E-S, Cole JB, Huson H, Wiggans GR, Van Tassell CP, Crooker BA, *et al.*(2013). Effect of

Artificial Selection on Runs of Homozygosity in U.S. Holstein Cattle. *PLoS ONE* **8**:

e80813–e80813.

Kirin M, McQuillan R, Franklin CS, Campbell H, McKeigue PM, Wilson JF (2010). Genomic

runs of homozygosity record population history and consanguinity. *PloS One* **5**: e13996.

Koonin EV (2009). Evolution of genome architecture. *Int J Biochem Cell Biol* **41**: 298–306.

Korneliussen TS, Albrechtsen A, Nielsen R (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* **15**: 356.

Lazell JD (1989). Wildlife of the Florida Keys: a natural history.

Li H (2011). Tabix: fast retrieval of sequence features from generic TAB-delimited files.
Bioinformatics **27**: 718–719.

Li H (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, *et al.*(2009). The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Maffei MD, Klimstra WD, Thomas J. W (1988). Cranial and Mandibular Characteristics of the
Key Deer (Odocoileus virginianus clavium). *J Mammal* **69**: 403–407.

Martchenko D, Shafer ABA (2023). Contrasting whole-genome and reduced representation
sequencing for population demographic and adaptive inference: an alpine mammal case
study. Heredity 131: 273–281.

McQuillan R, Leutenegger A-L, Abdel-Rahman R, Franklin CS, Pericic M, Barac-Lauc L, *et
al.*(2008). Runs of homozygosity in European populations. *Am J Hum Genet* **83**: 359–
372.

Meyermans R, Gorssen W, Buys N, Janssens S (2020). How to study runs of homozygosity
using PLINK? A guide for analyzing medium density SNP data in livestock and pet
species. BMC Genomics 21: 94.

Nelson SL, Taylor SA, Reuter JD (2021). An isolated white-tailed deer (Odocoileus virginianus)
population on St. John, US Virgin Islands shows low inbreeding and comparable
heterozygosity to other larger populations. Ecology and Evolution 11: 2775–2781.

Parker HG, VonHoldt BM, Quignon P, Margulies EH, Shao S, Mosher DS, *et al.*(2009). An
Expressed Fgf4 Retrogene Is Associated with Breed-Defining Chondrodysplasia in
Domestic Dogs. *Science* **325**: 995–998.

Pavlidis P, Jensen JD, Stephan W, Stamatakis A (2012). A Critical Assessment of Storytelling:

Gene Ontology Categories and the Importance of Validating Genomic Scans. *Mol Biol*

Evol **29**: 3237–3248.

Peart CR, Tusso S, Pophaly SD, Botero-Castro F, Wu C-C, Aurioles-Gamboa D, *et al.*(2020).

Determinants of genetic variation across eco-evolutionary scales in pinnipeds. *Nat Ecol*

Evol **4**: 1095–1104.

Pereira JA, Wolfenson LI, Artero DF, Argerich EC, Varela D, Fracassi NG (2023). First records

of pigmentation anomalies in the marsh deer at its southernmost stronghold: a warning

message? *Eur J Wildl Res* **69**: 59.

Pundir S, Martin MJ, O'Donovan C, Consortium TU (2016). UniProt Tools. *Curr Protoc*

Bioinforma **53**: 1.29.1-1.29.15.

R Core Team (2022). R: The R Project for Statistical Computing.

Reiner G, Weber T, Nietfeld F, Fischer D, Wurmser C, Fries R, *et al.*(2020). A genome-wide

scan study identifies a single nucleotide substitution in MC1R gene associated with white

coat colour in fallow deer (Dama dama). *BMC Genet* **21**.

Reissmann M, Ludwig A (2013). Pleiotropic effects of coat colour-associated mutations in

humans, mice and other mammals. *Semin Cell Dev Biol* **24**: 576–586.

Robinette W, Aldous C (1955). 'Parrot Mouth' in Rocky Mountain Mule Deer. *J Mammal* **36**:

308–310.

Robinson JA, Brown C, Kim BY, Lohmueller KE, Wayne RK (2018). Purging of Strongly

Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding

Depression in Island Foxes. Current Biology 28: 3487-3494.e4.

Robinson J, Kyriazis CC, Yuan SC, Lohmueller KE (2023). Deleterious Variation in Natural

Populations and Implications for Conservation Genetics. *Annu Rev Anim Biosci* **11**: 93–

114.

Russell CD, Widdison S, Leigh JA, Coffey TJ (2012). Identification of single nucleotide

polymorphisms in the bovine Toll-like receptor 1 gene and association with health traits

in cattle. *Vet Res* **43**: 17.

Santiago E, Novo I, Pardiñas AF, Saura M, Wang J, Caballero A (2020). Recent Demographic

History Inferred by High-Resolution Analysis of Linkage Disequilibrium. *Mol Biol Evol*

37: 3642–3653.

Santiago E, Caballero A, Köpke C, Novo I (2023). Estimation of the contemporary effective

population size from SNP data while accounting for mating structure. Molecular Ecology

Resources

Sidorenko J, Arias JD, Graham SE, Highland HH, Lüll K, Bielak LF, *et al.*(2022). A saturated

map of common genetic variants associated with human height. *Nat Lond* **610**: 704–712.

Smeds L, Ellegren H (2023). From high masked to high realized genetic load in inbred

Scandinavian wolves. Mol Ecol 32: 1567–1580.

Smits B, Bubenik GA (1990). Congenital osteopetrosis in white-tailed deer (Odocoileus

virginianus). *J Wildl Dis* **26**: 567–571.

Soule M (1973). The Epistasis Cycle: A Theory of Marginal Populations.

Stiles KA, Luke JE (1953). The inheritance of malocclusion due to mandibular prognathism. *J*

Hered **44**: 241–245.

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P (2015). Sambamba: fast processing of NGS

alignment formats. *Bioinformatics* **31**: 2032–2034.

U.S. Fish and Wildlife Service (1999). South Florida multi-species recovery plan.

Villanova VL, Hughes PT, Hoffman EA (2017). Combining genetic structure and demographic

analyses to estimate persistence in endangered Key deer (Odocoileus virginianus

clavium). **18**: 1061–1076.

Webb G (2018). Searching the internet to estimate deer population trends in the U.S., California,

and Connecticut. 19: 163–173.

Wilder AP, Supple MA, Subramanian A, Mudide A, Swofford R, Serres-Armero A, *et al.*(2023).

The contribution of historical processes to contemporary extinction risk in placental

mammals. *Science* **380**: eabn5856.

Wilderman A, D'haene E, Baetens M, Yankee TN, Winchester EW, Glidden N, et al. (2022). A

distant global control region is essential for normal expression of anterior HOXA genes

during mouse and human craniofacial development. bioRxiv: 2022.03.10.483852.

628
Wolff G, Wienker TF, Sander H (1993). On the genetics of mandibular prognathism: analysis of

large European noble families. *J Med Genet* **30**: 112–116.

Wolverton S, Huston MA, Kennedy JH, Cagle K, Cornelius JD (2009). Conformation to

Bergmann's rule in white-tailed deer can be explained by food availability. *Am Midl Nat*

162: 403–418.

Wootton E, Robert C, Taillon J, Côté S, Shafer ABA (2023). Genomic health is dependent on

long-term population demographic history. : 2022.08.16.503900.

Yang Y, Li SS, Chien JW, Andriesen J, Zhao LP (2008). A systematic search for

SNPs/haplotypes associated with disease phenotypes using a haplotype-based stepwise

procedure. *BMC Genet* **9**: 90.

Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S, *et al.*(2022). A saturated

map of common genetic variants associated with human height. Nature 610: 704–712.

Yu H, Wang Y, Gao J, Gao Y, Zhong C, Chen Y (2023). Application of the neuropeptide NPVF

to enhance angiogenesis and osteogenesis in bone regeneration. Commun Biol 6: 1–12.

Tables

Table 1: Summary statistics of subpopulations analyzed.

Subpopulation	Tajima's *D*	Nucleotide Diversity (π)	Mean F_{ROH}	Range of average ROH length (kb)
Anticosti	-0.09	0.007	0.01	153.41 – 172.89
SPM	0.54	0.005	0.02	150.92 – 202.66
North Mainland	-0.39	0.006	0.01	151.85 – 233.50
South Mainland	-0.51	0.007	0.01	158.69 – 191.03
Florida Keys	0.22	0.003	0.05	136.80 – 192.08

Table 2: List of ROHs shared by subsets of each group and the genes found within them (Matching candidate gene table = red, fixed SNPs = blue, haplotypes = orange, and haplotypes and fixed SNPs = green)

Group	Scaffold	Position	Genes Found
Leucism (4/4)	ref0002511	920,074 – 1,368473	GPATCH4, NAXE, TTC24, MEF2D, IQGAP3, RBMS2, RHBG, TSACC, GLMP, CCT3, PAQR6, SMG5, SLC25A44, PMF1, TMEM79, BGLAP, MEX3A, SEMA4A, LMNA
Leucism (3/4)	ref0002511	1,368,616 – 1,525,114	SSR2, UBQLN4, **LAMTOR2**, RAB25, ARHGEF2, RXFP4
Leucism (3/4)	ref0002511	2,174,336 – 2,491,037	DCST2, EFNA4, ZBTB7B, ADAM15, DCST1, FLAD1, EFNA3, SHC1, PBXIP1, PYGO2, PMVK, KCNN3
Leucism (3/4)	ref0002253	43,510,764 – 43,742,984	GCSH
Malocclusions (4/7)	ref0001199	600,106 – 794,137	**NPVF**
Small Stature (10/10)	ref0000154	7,305,609 – 7,635,609	**CHMP3**
Small Stature (10/10) (+1)	ref0000849	25104544 - 25401608	**LAMTOR1, TOMT, FOLR1, NUMA1, IL18BP, RNF121**, LOC110131318, LOC110131306, **ART1, ART5**, CHRNA10
Small Stature (9/10)	ref0001279	18,430,311 – 19,119,250	**RPL17**, OSR2
Small Stature (9/10)	ref0001398	1,372,352 - 1,653,529	**SOWAHA, LEAP2**, UQCRQ, GDF9, SHROOM1, AFF4, HSPA4
Small Stature (9/10)	ref0001398	1,1643,985 – 1,1796,333	PYURF, PTPRQ
Small Stature (9/10)	ref0001825	20,565,681 – 20,840,670	BMPR2, NOP58
Small Stature (9/10)	ref0001994	144,267 – 547,283	HERC1, FBXL22, USP3
Small Stature (9/10)	ref0002478	1,856,109 – 2,093,885	MAB21L2
Small Stature (9/10)	ref0002507	4,010,013 – 4,212,606	GRPEL1, CCDC96, **TADA2B**, TBC1D14, KIAA0232
Small Stature (9/10)	ref0002754	8,036,746 – 8,316,086	**MARCHF5, IDE, HHEX, KIF11, LOC110122451**

Figure legends

Fig. 1: Sampling locations with phenotype represented by colours: Normal (black), Leucism

(yellow), Malocclusion (blue), and Small Stature (red).

Fig. 2: Effective population size changes for the last 400 years, x-axis as generations before

present with a generation time of two years A) Key Deer B) Saint Pierre et Miquelon

Fig. 3: A) Percentage of the genome covered by ROHs (F_{ROH}) for each population. B) Bar plot

showing the total number of ROHs in each population stratified by ROH size category in kb. C)

Bar plot showing the total length (kb) of all ROHs in each size category (kb).

Fig 4: A) Number of homozygous missense mutations based on SNPeff annotations. B) Number

of homozygous loss of function mutations based on SNPeff annotations. Populations are colour

coded by phenotype: Normal (black), Leucism (yellow), Malocclusion (blue), and Small Stature

(red).

Leucism   Malocclusions   Small Stature   Normal