Photosynthetic demands on translational machinery drive retention of redundant tRNA metabolism in plant organelles

Rachael Ann DeTar[1],*, Joanna Chustecki[2], Anna Martinez-Hottovy[2], Luis Federico Ceriotti[3,4], Amanda K. Broz[1], M. Virginia Sanchez-Puerta[3,4], Christian Elowsky[5], Alan C. Christensen[2], Daniel B. Sloan[1]

[1] Department of Biology, Colorado State University, Fort Collins, CO, USA
[2] School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA
[3] IBAM, Universidad Nacional de Cuyo, CONICET, Facultad de Ciencias Agrarias, Almirante Brown 500, M5528AHB Chacras de Coria, Argentina
[4] Facultad de Ciencias Exactas y Naturales, Padre Jorge Contreras 1300, Universidad Nacional de Cuyo, M5502JMA Mendoza, Argentina
[5] Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, NE, USA
* Corresponding Author

Keywords: organelle gene expression, plastome, mitogenome, tRNA, photosynthesis, aminoacyl-tRNA synthetase

Abstract

Eukaryotic nuclear genomes often encode distinct sets of translation machinery for function in the cytosol vs. organelles (mitochondria and plastids). This raises questions about why multiple translation systems are maintained even though they are capable of comparable functions and whether they evolve differently depending on the compartment where they operate. These questions are particularly interesting in plants because translation machinery, including aminoacyl-tRNA synthetases (aaRS), is often dual-targeted to the plastids and mitochondria. These organelles have different functions, with much higher rates of translation in plastids to supply the abundant, rapid-turnover proteins required for photosynthesis. Previous studies have indicated that plant organellar aaRS evolve more slowly compared to mitochondrial aaRS in eukaryotes that lack plastids. Thus, we investigated the evolution of nuclear-encoded organellar and cytosolic aaRS and tRNA maturation enzymes across a broad sampling of angiosperms, including non-photosynthetic (heterotrophic) plant species with reduced plastid gene expression, to test the hypothesis that translational demands associated with photosynthesis constrain the evolution of enzymes involved in organellar tRNA metabolism. Remarkably, heterotrophic plants exhibited wholesale loss of many organelle-targeted aaRS and other enzymes, even though translation still occurs in their mitochondria and plastids. These losses were often accompanied by apparent retargeting of cytosolic enzymes and tRNAs to the organelles, sometimes preserving aaRS-tRNA charging relationships but other times creating surprising mismatches between cytosolic aaRS and mitochondrial tRNA substrates. Our findings indicate that the presence of a photosynthetic plastid drives the retention of specialized systems for organellar tRNA metabolism.

Significance

The process by which endosymbionts are integrated into a host and become organelles results in a combination of gene loss, transfer to the nucleus, and retention in the organellar genome. It is not clear why some endosymbiont-derived genes may be retained when a functional host counterpart exists whose gene product could simply be retargeted to the organelles. This study revealed that the photosynthetic activity in plant plastids may be responsible for retention of functionally redundant tRNA processing machinery, while mitochondria are more flexible regarding substitution with cytosolic-type enzymes. Therefore, functional constraint in the plastid is likely more important than in the mitochondria for shaping the evolution and retention of functionally redundant proteins that are dual targeted to both organelles.

Introduction

The evolutionary transition from a free-living bacterium to an endosymbiont and eventually to an

organelle requires the physical, biochemical, and genomic integration of two organisms. One common

consequence of this process is a reduction in the genome of the endosymbiont, driven by relocation of

essential genes to the nucleus via horizontal transfer or outright loss of some genes that are no longer

useful after transition to an intracellular lifestyle. This trajectory has been observed repeatedly for

various endosymbionts and organelles, including mitochondria, plastids, and more recently acquired

intracellular bacteria such as the cyanobacterial-derived chromatophores of *Paulinella* amoeba or the

many heritable endosymbionts in insects (1–6). However, there are some endosymbiotically derived

genes that are recalcitrant to loss despite the presence of host counterparts with comparable functions.

Redundant genes may be maintained in the organelle genomes or in the nuclear genome alongside

genes encoding proteins for the same role in other cellular compartments. These observations elicit

some interesting questions; 1) Why are some functionally redundant genes retained? 2) Do these

redundant genes evolve differently depending on the cellular compartment where they operate?

The genes that encode the machinery required for organelle translation are widely retained, as

eukaryotes maintain distinct translation systems for nuclear and mitochondrial transcripts. Translation

becomes even more complicated in eukaryotes bearing three genomes. For example, plant cells contain

nuclear, mitochondrial, and plastid genomes. The translation machinery associated with each of these

cellular compartments includes ribosomal RNAs (rRNAs), ribosomal proteins, transfer RNAs (tRNAs), and

the accompanying enzymes involved in tRNA processing and aminoacylation. The aminoacyl-tRNA

synthetases (aaRS) that catalyze the "charging" of a tRNA with its cognate amino acid tend to be highly

conserved, with distinct versions for function in the organelles and cytosol (7). The degree of

redundancy of tRNAs and their interactors between cellular compartments also varies significantly

within eukaryotes. These characteristics make tRNAs and their interactors, particularly aaRS, an ideal

case study for understanding cytonuclear gene redundancy.

The concerted activity of the aaRS along with other enzymes involved in post-transcriptional

modification such as tRNases P and Z (end processing in tRNA maturation) and CCAses (addition of CCA

tail) ensures the tRNA has the proper structure to fulfill its role in translation. Generally, organisms

would need a complement of at least 20 aaRS, one for each amino acid. However, eukaryotes often have

more than this minimum requirement, with some overlapping sets of enzymes targeted to different

compartments (8, 9). It is not fully understood why some eukaryotes retain distinct, redundant sets of

tRNA processing machinery, while others do not. However, retention of organelle-targeted aaRS

correlates with the retention of tRNA genes within organelle genomes. For example, *Homo sapiens*

retains 38 aaRS with nearly distinct sets for the cytosol and mitochondria (9–11). Correspondingly, all

necessary tRNA genes are present in the mitochondrial genome (mitogenome) (8). In contrast,

trypanosomatids do not encode any tRNAs in their mitogenomes but rather import nuclear-encoded

tRNAs into the mitochondria and have a reduced set of aaRS that colocalize to the cytosol and the

mitochondria (9, 12, 13). In plants, closely related species may have very different numbers of tRNAs

encoded in their mitogenomes, with import of cytosolic tRNAs to compensate for losses (8, 14, 15).

Notably, plants also dual-target and import an organellar set of enzymes into both plastids and

mitochondria (9, 16). The co-occurrence of redundant tRNAs and aaRS could be explained by the

observation that organelle-encoded tRNAs are often structurally divergent from cytosolic tRNAs and

require specific enzymatic partners for recognition (8, 17).

For most eukaryotes, organellar and cytosolic aaRS are exclusively nuclear-encoded. The shared

genomic location of these genes facilitates direct comparison of sequence conservation because any

differences in evolutionary rate more likely reflect differences in selection, rather than other factors like

compartment-specific mutation rates or effective population size (18–21). Previous studies have

quantified rates of evolution for these enzymes in various animal lineages and a small number of plants.

Mitochondria-specific aaRS in animals exhibit lower sequence conservation than their cytosolic

counterparts (22–24). In contrast, plant organellar aaRS exhibit equivalent or greater sequence

conservation compared to their cytosolic counterparts (25). Furthermore, plastid genomes (plastomes)

tend to retain a set of 30 tRNA genes that is fully sufficient for translation (26, 27). Plastomes also

generally have high rates of transcription (28) and translation (29) due to the abundance and rapid

turnover of proteins required for photosynthesis. Thus, the disparity in evolutionary rates between

animal mitochondrial aaRS and plant dual-organelle targeted aaRS may be due to the specific

translational needs of the plastid. We hypothesize that aaRS and other translation-associated enzymes

operating in the plastid face great functional constraints, thus promoting unusually high sequence

conservation of organellar translation machinery compared to other eukaryotes. A prediction stemming

from this hypothesis is that plant lineages with reduced or nonexistent plastid gene expression would

exhibit more rapid evolution of organellar aaRS due to relaxed selection.

In this work, we test whether photosynthetic activity contributes to sequence conservation of

organellar tRNA processing and charging enzymes by analyzing evolution of this machinery across

diverse lineages of autotrophic and heterotrophic plants. It should be noted there is a gradient of

heterotrophy amongst plants – for the purposes of this study, we use "heterotrophic" to describe plants

that do not photosynthesize at all and exclusively rely on parasitism for acquisition of carbon. They do

this by tapping directly into host plants or by exploiting mycorrhizal fungi (mycoheterotrophy) (30, 31).

We refer to facultative autotrophs that supplement photosynthetic carbon with parasitism as

"hemiparasitic." The hallmark of many heterotrophic/hemiparasitic plants is a reduced or even non-

existent plastome (32). Highly expressed genes related to photosynthesis such as the Rubisco large

subunit and the Photosystem II D1 reaction center tend to be absent from plastomes in heterotrophs,

thereby drastically decreasing demands on plastid gene expression. In addition, many tRNA genes have

been lost from the plastomes of these plant species (32), providing a good system to examine the

relationship between evolutionary redundancy of aaRS, tRNA content of the organelle genomes, and

trophic lifestyle.

Results

Heterotrophic plants exhibit extensive loss of organellar aaRS and tRNA processing enzymes but not

ribosomal protein subunits

We searched for organellar and cytosolic aaRS orthologs in a broad sampling of angiosperm species with

a range of trophic lifestyles and plastome reduction (**Supp. Dataset 1**). For each

hemiparasitic/heterotrophic species in the analysis, we included the closest fully autotrophic relative

with a nuclear genome or transcriptome available, excepting the Balanophoraceae where we used

hemiparasite *Santalum album* as the reference autotroph. In total, we sampled 7 heterotrophs, 8

hemiparasites, and 10 autotrophs, including reference species *Arabidopsis thaliana.* Among

heterotrophs, we included the endoparasites *Rafflesia cantleyi* and *Sapria himalayana* from the family

Rafflesiaceae, likely the most extreme examples of plant heterotrophy, as several studies have

suggested wholesale loss of the plastome in these species (33–36).

We found orthologs across these diverse taxa by running Orthofinder (37, 38) on protein models

derived from nuclear genome or transcriptome data to find groups of likely orthologous proteins based

on similarity and phylogeny, i.e., "orthogroups". Then we extracted sequences that fall into an

orthogroup with known *A. thaliana* aaRS (**Supp. Dataset 2**). It is important to note that the number of

homologous sequences found for each species in this analysis may not directly reflect the absolute

number of genes or proteins due to the fragmented nature of gene models made from transcriptome

data and the inclusion of two independent sets of protein models for *Balanophora fungosa* and *S.*

himalayana. Unexpectedly, many lineages of heterotrophs did not have any orthologs for many of the

organellar aaRS. This loss of organelle-specific orthologs in heterotrophs was surprising as even species

entirely lacking the plastome must still conduct tRNA aminoacylation within mitochondria. Further

investigation using reciprocal-best-hit (RBH) searches with BLAST between *A. thaliana* and each species

supported that these were true losses. These losses primarily occurred in purely heterotrophic plants,

while hemiparasites retained most of their organellar aaRS. (**Fig. 1**). Yet, within the heterotrophs, there

was a gradient of loss. The two endophytic species from Rafflesiaceae (*R. cantleyi* and *S. himalayana*)

exhibited the most striking loss of organellar aaRS, retaining only PheRS. In contrast, obligate

mycoheterotroph *Monotropa hypopitys* retained more aaRS than any other heterotroph, missing only

AlaRS, CysRS, ThrRS, and TrpRS (**Fig. 1**).

This striking pattern led us to investigate the retention of other enzymes involved in tRNA

maturation and processing; see (15) and **Supplemental Dataset 2** for a comprehensive list.

Unsurprisingly, the CCAse enzyme responsible for adding CCA tails to tRNAs in all compartments was

retained across all species. Two enzymes that maintain distinct cytosolic and organellar versions, tRNase

Z (involved in 3' end processing) and Protein Only RNase P (PRORP; required for 5' end processing),

exhibited two different trends. We found that plastid tRNase Z (tRNase Z2) (39) was missing from all

heterotrophs except *M. hypopitys* and hemiparasites of the *Cuscuta* genus yet was retained in the

autotrophic relatives. Conversely, the organellar PRORP1 (40, 41) appeared to be universally retained

while cytosolic PRORP2/3 orthologs were missing from the Rafflesiaceae. We also investigated the

retention of bacterial-like organelle-specific machinery with no cytosolic counterpart, including the

methionyl-tRNA formyltransferase (MTF), tRNA-Ile lysidine synthetase (TilS), and the bacterial-type

glutamine amidotransferase (GatCAB) complex. We found orthologs for the putative *A. thaliana* MTF in

all species except the heterotrophs *Epipogium roseum* and *M. hypopitys,* and putative TilS orthologs

were absent only from the heterotrophs Rafflesiaceae and *B. fungosa*. GatCAB is the key complex used

by organelles in land plants to accurately charge tRNA^{Gln} in the absence of a dedicated organellar GlnRS

(42–44). This involves a two-step process, whereby tRNA^{Gln} is initially mis-acylated with Glu by GluRS.

GatCAB then catalyzes the transamidation of Glu to Gln. Most of our test species, even heterotrophs,

retained genes for at least some of the GatCAB subunits. However, Rafflesiaceae species appear to have

lost all subunits in this complex.

To assess whether other functionally redundant translation machinery was lost in heterotrophic

lineages, we analyzed presence/absence of nuclear-encoded riboproteins with exclusive targeting to a

given compartment (Table S1, Supp. Dataset 2, "Riboproteins"). While there was minor loss of cytosolic,

mitochondrial, and plastid subunits in all heterotrophic lines, the Rafflesiaceae lineages exhibited a

dramatic reduction in plastid subunits, which is expected because these lineages appear to have

dispensed with the plastid genome entirely and consequently no longer need plastid ribosomes. This

finding reveals a key contrast between the fate of organellar aaRS and riboproteins in response to loss of

photosynthesis and may reflect the fact that aaRS are shared between the two organelles, whereas the

riboproteins are compartment-specific.

Generally, we observed reductive evolution in all heterotrophs via loss of organellar aaRS and

bacterial-type enzymes for tRNA modification. This raises the question as to how gene expression is

accomplished in the organelles of these species. One possibility is that cytosolic enzymes are retargeted

to the organelles to functionally replace organellar enzymes.

Heterotrophic plants may compensate for loss of organellar aaRS by retargeting cytosolic

counterparts

To test for cytosolic aaRS retargeting, we applied two *in silico* targeting prediction programs to protein

models for all cytosolic aaRS in our heterotrophs and their close autotrophic relatives (**Fig. 2, Fig. S1)**.

Our analysis revealed that canonically cytosolic aaRS are disproportionately retargeted to mitochondria

in obligate heterotrophs compared to autotrophs ($p = 0.018$). There was also a non-significant trend

towards retargeting to the plastid ($p = 0.11$). Targeting of mitochondrial OXPHOS components and

plastid housekeeping proteins including subunits of the caseinolytic protease (Clp) complex and the

acetyl-CoA carboxylase (ACC) complex involved in producing malonyl-CoA for fatty acid biosynthesis

were included as known organellar positive controls (Supp. Dataset 2, "targeting controls"). Protein

degradation and fatty acid biosynthesis are essential plastid functions that are still necessary in the

absence of photosynthesis. Cell wall synthesis enzymes were included as negative controls. Of the

controls, the only group that showed a different degree of targeting than expected was the

mitochondrial OXPHOS components, where recovery of proteins with known mitochondrial targeting

was low (~60%) for both autotrophs and heterotrophs (Supp. Table 2). Thus, these *in silico* approaches

appear to successfully (albeit imperfectly) predict subcellular localization in both heterotrophic and

autotrophic plants.

We would specifically expect that a given cytosolic aaRS is more likely to be retargeted if its

organellar counterpart is lost. We examined this possibility and found a clear and statistically significant

correlation between loss of an organellar aaRS and retargeting of the corresponding cytosolic enzyme to

one or both compartments (p = 0.005, **Fig. 4, Fig. S2**).

The variable quality of gene models for each species and high false negative rate for retargeting

prediction confound interpretation of retargeting for individual enzymes in each species (**Fig. 4**, **Fig. S3**).

However, when looking only at *S. himalayana*, a species with relatively high-quality genome-based

models, we observed some interesting phenomena. Cytosolic AlaRS, GlnRS, MetRS, ProRS, and TrpRS

were predicted to be retargeted to mitochondria. Retargeting of cytosolic GlnRS could explain how this

species copes with loss of the GatCAB complex (**Fig. 1**). The cytosolic AspRS, GluRS, and GlyRS were

predicted to be dual targeted to both organelles, while cytosolic TyrRS and the PheRSα subunit were

predicted to be targeted to the plastid despite the likely absence of a plastid genome. All other cytosolic

enzymes were either not predicted to be retargeted or were missing a large portion of the N-terminus.

The fact that some cytosolic aaRS corresponding to lost organellar aaRS were not predicted to be

retargeted to the mitochondria and that some cytosolic aaRS were targeted to a plastid presumed to

lack gene expression may reflect an underestimation of mitochondrial retargeting due to the constraints

of *in silico* predictions and the potential divergence of TP sequences in Rafflesiaceae. Indeed, an

investigation of subunits for the protein import apparatus of plastids and mitochondria revealed that

the Rafflesiaceae are missing some key components of the plastid import apparatus (Fig. S6), some of

which are responsible for TP binding and recognition.

To assess whether our *in silico* predictions hold true *in vivo,* we conducted targeting assays on a

subset of aaRS (see **Supplemental Dataset 3** for sequence information) by inducing transient expression

of N-terminal fusions of *S. himalayana* cytosolic aaRS transit peptides (TPs) to green fluorescent protein

(GFP) in *Nicotiana benthamiana* (**Fig. 3, Fig. S4-S5**). We used eqFP611 tagged with the TP from

mitochondrial isovaleryl-CoA dehydrogenase (IVD) as a positive control for mitochondrial targeting.

Infiltration with single expression vector with just IVD_FP611 was used as a negative control (**Fig. S4**).

We found that proteins predicted to be retargeted *in silico* were almost universally retargeted to one or

both organelles *in vivo*. In some cases, putative TPs retargeted GFP to a different organelle than

predicted *in silico.* AsnRS displayed the plastid targeting that was predicted *in silico* and additional

mitochondrial targeting *in vivo* (**Fig. 3**)*.* GluRS and GlyRS were predicted to be dual-targeted but

exhibited targeting to only one of the two organelles. The only two TPs for which organellar GFP

localization completely mismatched the *in silico* prediction (but still exhibited organelle targeting) were

ProRS and TrpRS, which both showed strong plastid GFP fluorescence *in vivo* despite mitochondrial

predictions. These results were consistently observed with two different cloning and expression

strategies (Fig. 3, Fig. S5)**.** The cytosolic AspRS TP was the only one we cloned that did not exhibit

targeting to either organelle *in vivo* despite a clear biological expectation that it would have evolved

retargeting (**Fig. S4**). GFP tagged with this TP formed punctate structures in the cytosol, which may be

artefactual aggregates resulting from overexpression. Generally, *in vivo* assays provided evidence for

retargeting of cytosolic aaRS to at least one organelle and suggested that *in silico* prediction tools may

not be adept at distinguishing plastid from mitochondrial targeting.

Generally, organellar tRNAs differ from their cytosolic counterparts in evolutionary history,

sequence, and posttranscriptional modification (8). Identity elements that govern the specificity of

interaction between a tRNA and its conjugate aaRS may also differ between cytosolic and organellar

systems (45). However, in many systems (including plants), cytosolic tRNAs can be imported into the

organelles (8, 9, 14). Thus, our observation of retargeting opens some interesting questions – are the

retargeted aaRS charging organellar tRNAs? Or is there replacement of the entire organellar system of

tRNA aminoacylation, including both tRNAs and aaRS? We investigated the tRNA gene content of the

organelles of these species to gain insight into which tRNA substrates may be in use.

Organelle-encoded tRNA retention often, but not always, mirrors retention of organelle-type aaRS

To investigate the possibility of enzyme-substrate mismatch, we cataloged the presence/absence of

organellar tRNA-encoding genes (*trn*) for all sampled heterotrophic species (**Fig. 4, Supplemental**

Dataset 4)**.** Generally, there was strong correspondence between retention of organellar aaRS and

plastome/mitogenome tRNA genes. As the Rafflesiaceae species are thought to lack plastomes entirely,

they likely have no plastome-encoded tRNAs (**Fig. 4**). The Balanophoraceae species used in this study

also have been reported to exhibit loss of all plastid tRNA genes, except for an extremely divergent *trnE*

in *B. fungosa* (46–49). The Rafflesiaceae and Balanophoraceae have additionally lost most of their

mitogenome-encoded tRNAs. Correspondingly, these two families also have the most reduced sets of

organellar aaRS. At the opposite extreme, *M. hypopitys* retains most of its mitochondrial and plastid

tRNA genes and corresponding organellar aaRS.

However, there are some incidences where a cytosolic enzyme looks to be charging an

organellar tRNA due to loss of the organellar aaRS (**Fig. 4**). For example, the few remaining mitogenome-

encoded tRNAs in the Rafflesiaceae are missing their organellar aaRS partner (except *S. himalayana*

tRNA^{Phe}) and are likely charged by a cytosolic enzyme. Alternatively, they may be pseudogenes that have

been functionally replaced by an imported tRNA. For both compartments, there are numerous instances

where there is no encoded tRNA despite retention of the organellar aaRS (**Fig. 4**). For example, *R.*

phalloides retains its organellar SerRS, PheRS, and GlnRS but has lost the corresponding mitochondrial

tRNAs. No retargeting of cytosolic enzymes is predicted in these cases, suggesting the organellar enzyme

may be acting on an imported cytosolic tRNA.

Testing for accelerations in organellar aaRS sequence evolution in heterotrophs

Although wholesale loss of many organellar aaRS in heterotrophs precludes analysis of sequence

conservation, *M. hypopitys* retained the majority of organellar aaRS despite exhibiting an obligate

heterotrophic lifestyle. Thus, we quantified sequence conservation of organellar and cytosolic aaRS in

M. hypopitys and its autotrophic relative *Rhododendron williamsianum.* We built gene trees and ran

relative rate analysis to test for accelerated evolution on the *M. hypopitys* branch (**Supplemental**

Dataset 5). We also directly compared branch length ratios of *M. hypopitys* to *R. williamsianum* in

cytosolic and organellar aaRS (**Fig. S7, Supplemental Dataset 5).** Neither analysis showed a statistically

significant decrease in sequence conservation in organellar enzymes of *M. hypopitys* (**Fig. S7**). However,

the deep rooting of our tree on *A. thaliana* may result in poor resolution of the sequence evolution

between *M. hypopitys* and *R. williamsianum.* Additionally, the transition to heterotrophy on the *M.*

hypopitys branch may have occurred relatively recently compared to the split from *R. williamsianum*,

thus diluting any signals of accelerated evolution.

Discussion

Loss of photosynthesis relaxes constraints on both plastid and mitochondrial translation machinery

Our primary objective in this study was to test whether the high translational demands of the

photosynthetic chloroplast create functional constraint on the evolution of shared translation systems in

plastids and mitochondria. To do so, we investigated relative rates of protein evolution in cytosolic and

organellar aaRS in heterotrophic plants compared to their photosynthetic relatives. To our surprise,

heterotrophic plants underwent widespread loss of organellar aaRS and many bacterial-like enzymes

involved in tRNA processing. These losses suggest an extreme relaxation of selection pressures in the

absence of a photosynthesizing plastid and support our hypothesis that the translational demands in

chloroplasts are the linchpin responsible for the unusually strong functional constraint on the redundant

organelle-specific tRNA metabolism in plants. These correlated effects on mitochondrial translation

appear to be stronger when the same proteins are shared by mitochondria and plastids (e.g., aaRS)

because we found that the most reduced heterotrophs from Rafflesiaceae lost many plastid-specific

riboproteins but not their mitochondrial-specific counterparts.

There was variation in the degree of loss across the heterotrophs, often correlated with the

number of genes retained in the plastome, particularly tRNAs (see **Supp. Dataset 1** for

plastome/mitogenome content). For example, the most aaRS losses occurred in the Rafflesiaceae, all of

which are obligate endoparasites and have likely lost their plastomes entirely (33–36, 50). The fewest

aaRS losses occurred in *M. hypopitys*, which has the highest number of plastome-encoded genes of all

the heterotrophic species we sampled – and notably the greatest number of retained plastid tRNAs (51).

Thus, *M. hypopitys* may not be as far along the path of reductive evolution as the other species, which

might provide an additional explanation as to why our comparative rate analyses on the remaining

organellar aaRS in *M. hypopitys* did not support accelerated evolution compared to cytosolic enzymes

(**Fig. S7**). The Balanophoraceae and *Epipogium* species exhibited intermediate degrees of aaRS and

plastome gene loss, although the Balanophoraceae species used in this study have lost all plastid tRNAs

save for a truncated *trnE* in *B. fungosa* that is likely retained for use in the first step of heme

biosynthesis *(46).* An increased plastome mutation rate caused by loss of DNA repair machinery (52, 53)

in the Balanophoraceae may explain accelerated plastome (tRNA) gene losses relative to nuclear (aaRS)

gene loss. Generally, loss of organellar aaRS correlates well with reduction of the plastome across

heterotrophs.

Loss of organellar aaRS is associated with retargeting of cytosolic aaRS

The wholesale loss of organellar aaRS in heterotrophs was unexpected, as most non-photosynthetic

plants still contain a minimal plastome to support heme and fatty acid biosynthesis (32, 54) and fully

intact mitogenomes (see **Supp. Dataset 1**). This conundrum prompted us to investigate if cytosolic aaRS

are retargeted to organelles in heterotrophs to compensate for gene loss, as has been shown for

aaRS/tRNAs in mitochondria of other plant species (55). Retargeting a protein to the organelles

generally involves obtaining an N-terminal TP sequence that can be recognized by organellar protein

import complexes (56–58). We discovered that many cytosolic aaRS in the heterotrophs had N-terminal

sequences predicted to target to one or both organelles (**Fig. 2-3, S2-S5**). Therefore, the import of

cytosolic aaRS likely compensates for the loss of organellar aaRS and even the loss of the GatCAB

complex in Rafflesiaceae via retargeting of cytosolic GlnRS.

There was not perfect correspondence between loss of organellar aaRS and predicted

retargeting of cytosolic aaRS. However, the extent of retargeting may have been underestimated for

two main reasons. Firstly, TP sequence composition and protein import machinery vary considerably by

plant lineage (59), making _in silico_ predictions more challenging in uncharacterized species. Thus, some

organelle-imported cytosolic aaRS may eschew the use of typical TPs that are key for targeting

prediction _in silico_ (60). Indeed, some organellar proteins are known to not rely at all on N-terminal TPs

for import (61, 62). TP sequence divergence could also be accelerated by alteration of the sequence or

subunit composition of protein import machinery. Notably, our presence/absence analysis of plastid and

mitochondrial protein import machinery revealed some critical subunits for plastid protein import are

lost in the heterotrophs, especially the Rafflesiaceae, which could influence which sequences are

recognized as transit peptides (**Fig. S6**). For example, Toc159, one of the key outer envelope import

receptors involved in TP binding and recognition, is missing from the Rafflesiaceae (63). This subunit

specializes in binding and import of proteins involved in photosynthesis, whereas Toc132, which is

retained in these species, specializes in "non-green" protein import (64, 65). The Rafflesiaceae are also

missing Tic236, the protein that spans the space between the envelope membranes to physically

connect the inner and outer envelope (66). This raises interesting questions about the nature of plastid

protein import and the ultrastructure of envelope membranes in Rafflesiaceae. Other heterotrophs in

addition to the Rafflesiaceae are also missing some subunits present in *A. thaliana*; *Epipogium* species,

M. hypopitys and the Rafflesiaceae lack one or more Tic20 and Tic22 paralogs. The Tic22 protein family

is highly conserved and expressed, with a hypothesized role in outer membrane maturation in addition

to its role in protein import (67, 68). Study of Tic-22 loss-of-function mutants revealed that these

proteins are not essential, at least in *A. thaliana* (67, 69). Tic20 forms the structural basis of the inner

pore of the translocon, with Tic20-I/Tic20-IV being the essential isoforms for plant survival (63, 70–72).

Similar trends have also recently been found by Guo et al. 2024 (73). These alterations to the

composition of the TIC-TOC complex may reduce the specificity of plastid protein targeting predictions.

In contrast, the heterotrophic lines retained almost all TIM-TOM subunits forming the mitochondrial

translocon. The divergence of the TIC-TOC complex in some heterotrophs may result in accelerated

evolution of TPs, thus, targeting predictions made based on characteristics of photosynthetic plants may

not be fully accurate.

Secondly, uncertain protein models for some of our species may have led to true N-terminal

sequences being absent or incomplete. Retargeting of cytosolic aaRS may occur through two different

mechanisms; 1) alternative splicing or alternative transcription/translation start sites within one gene

resulting in cytosolic and organellar isoforms and 2) duplicated genes within the genome encoding

cytosolic and organellar paralogs. Our ability to investigate these two mechanisms was limited as we

generally only had transcriptome or genome information but not both for each species. However,

observations of transcriptomes assembled with consideration for alternative splice isoforms indicate

that both mechanisms are potentially at play (**Supplemental Dataset 6**). For example, *Rhopalocnemis*

phalloides has two separate paralogs for cytosolic ThrRS, one of which is predicted to be mitochondrial-

targeted (Rphallo_DN1669_c0_g1) while the other is not (Rphallo_DN1669_c0_g2). Conversely, *R.*

phalloides TyrRS is encoded by a single locus (Rphallo_DN204_c0_g1) with a multitude of splice

isoforms, some of which are putatively mitochondrial-targeted. Genome sequencing and long-read

mRNA sequencing would allow for more thorough analysis of which mechanism is most common in each

species.

While almost all the TPs that we investigated with *in vivo* microscopy assays exhibited targeting

to mitochondria and/or plastids, the specific organelle sometimes differed from the *in silico* prediction.

The ProRS and TrpRS results in particular were unexpected as the *in silico* predictions were strongly

mitochondrial (**Supp. Dataset 3**), but we observed clear plastid localization *in vivo* in two separate

rounds of imaging with different vectors (**Fig 3, Fig. S4, S5)**. There were other TPs where targeting

predictions were only partly supported by GFP localization experiments, including GluRS and GlyRS

which exhibited targeting to only one of the two predicted organelles, and AsnRS which was only

predicted to have plastid localization but additionally exhibited mitochondrial localization *in vivo.*

These results may reflect the fact that target peptide prediction models like those used in this

study do not have complete precision or sensitivity for a given subcellular compartment even when

applied to benchmark datasets. This is not surprising, as changing just a few amino acids in a transit

peptide can alter protein targeting from singular to dual targeting *in vivo* (74). Furthermore, predictions

made by a classifier trained on a broad sampling of TPs from diverse plant species cannot accurately

predict every TP for an individual species, since species-specific features may not be represented

enough to be detected by the model. Even if we assume our microscopy data are a perfect reflection of

"true" targeting in the native system, we might expect some inaccuracies in our targeting predictions

just based on the sensitivity and precision of the prediction algorithms (TargetP-2.0 and LOCALIZER). Of

the two models, TargetP-2.0 specifically identifies mitochondrial and plastid transit peptides with 85-

86% sensitivity and 87-90% precision and differentiates proteins lacking TPs with over 98% sensitivity

and precision (75, 76). The publication describing LOCALIZER does not directly quantify the tool's ability

to distinguish proteins with no transit peptide but has 60-73% sensitivity and 60-80% precision for

mitochondrial and plastid predictions respectively, counting proteins with predicted dual-targeting as

positive hits for both plastid and mitochondria.

The artificial nature of heterologous expression in *N. benthamian* can also create artefacts. We

can see that in some cases predictions for a given TP can change when it is fused to GFP as opposed to

the native aaRS sequence (Supp. Dataset 3). This implies that more than just the N-terminal extension of

a protein determines targeting and may create some artefacts in assessing targeting based on TP-GFP

fusions. Furthermore, alternative splicing or translation initiation sites can be a mechanism of

alternative targeting (77–79). Alternative translation initiation may require RNA cis regulatory factors

i.e., 5' UTR sequences not captured in our heterologous expression system (80). Additionally, the import

of a heterologously expressed TP-tagged GFP can change based on the age of leaves during infiltration,

the efficiency of protein folding, protein expression levels, and truncation of an internal targeting

sequence (81–83). As comparison of the *in silico* and heterologous expression assays is effectively

combining the errors created by both methods, we would not expect perfect concordance between the

methods regarding the specific identity of the retargeted organelle. Rather, these methods are

complementary tools to assess general retargeting to the endosymbiotic organelles rather than

differentiating between TPs specifically targeting mitochondria or plastids from those that are dual

targeted. An additional method for future investigation of targeting to plastids or mitochondria would

be to subfractionate tissue from one or more of these heterotrophic plants species and conduct

comparative high throughput proteomics on each fraction.

Potential rewiring of aaRS-tRNA interactions

Retargeting of cytosolic aaRS led us to investigate organelle genome tRNA content and whether

there was substrate-enzyme mismatch in the organelles, i.e. instances where the cytosolic enzyme was

interacting with an organellar tRNA or vice-versa. Our interpretations presume that loss of organellar

tRNA genes is compensated by import of the corresponding cytosolic tRNAs (9, 14, 84). We found that

the retention of organellar aaRS in heterotrophs generally matched the retention of plastid tRNAs but

not necessarily mitochondrial tRNAs (**Fig. 4**). This is further supported by observations in hemiparasite

Viscum album, which has lost most of its mitochondrial tRNAs, yet retains most organellar aaRS in

conjunction with a nearly full set of plastid tRNAs (32, 85, 86). Substrate-enzyme mismatch was rare

within the plastid and usually involved loss of a plastid tRNA yet continued presence of the

corresponding organellar aaRS (e.g., see *R. phalloides* where all plastid tRNAs are lost but a handful of

organellar aaRS remain). There were also instances of organellar aaRS apparently charging imported

cytosolic tRNAs in the mitochondria (e.g., *trnF* in *R. cantleyi and R. phalloides*). More surprisingly,

however, there were cases where a mitochondrial tRNA was retained despite loss of the corresponding

organellar aaRS and apparent retargeting of the cytosolic enzyme, particularly in the Rafflesiaceae. This

type of mismatch is unexpected because cytosolic aaRS are thought to generally be more discriminating

enzymes than their mitochondrial counterparts (8, 55, 87–89), making it more difficult to transition to a

new tRNA substrate. However, our results suggest some cytosolic enzymes may be interacting with

bacterial-type tRNAs in these heterotrophic species, although we cannot rule out the possibility that

cytosolic tRNAs are being redundantly imported. Indeed, our findings raise fascinating questions about

the process of functionally replacing tRNAs and interacting enzymes in these dynamic systems (55, 90),

which could be explored with comparative and biochemical approaches in closely related species still

undergoing these transitions. Given that parasitic plant lineages tend to be fast evolving across all three

genomes (91), the cytosolic aaRS in these species may have evolved lower specificity and the capability

to charge organellar tRNAs. Overall, our findings suggest that the plastid tRNA content, and not the

mitochondrial tRNA content, drives retention of organellar aaRS, again placing the photosynthetic

plastid as a central factor in the retention of functionally redundant translation machinery.

The identity of retained organellar aaRS versus those substituted by cytosolic counterparts was

seemingly non-random. GluRS and PheRS were the two mostly highly retained organellar aaRS across

these diverse lineages of plants (**Fig. 1**). Organellar PheRS is likely retained because the cytosolic PheRS

is unusual in that it is a heteromeric complex with independently encoded α and β subunits (92).

Therefore, functional replacement of the organellar PheRS would require two independent acquisitions

of TPs, as well as stoichiometric import and assembly within the organelles. Such a coordinated set of

changes may be unlikely to evolve (24, 55). The one exception was *E. aphyllum,* for which we could not

find an organellar PheRS. However, this may be an artefact of low expression, as protein models for this

species were curated from RNA-seq data. As tRNA^{Glu} is the most frequently retained tRNA in

heterotrophic plant plastomes due to its additional role in the first step of heme biosynthesis (54, 93,

94), it is not surprising that GluRS was among the most retained organellar aaRS in this study.

Intriguingly, the *Rafflesiaceae* are the only species that lost the organellar GluRS, as well as plastid *trnE*.

Yet, genes encoding heme synthesis enzymes are retained in these species (92), opening interesting

questions into how they carry out the first step in the pathway. In *S. himalayana*, we found retargeting

of the cytosolic GluRS to the plastid, which along with presumed import of the cytosolic tRNA^{Glu}, could

functionally replace the first step in the heme biosynthesis pathway. This has the interesting implication

that the glutamyl-tRNA reductase catalyzing the next step in heme biosynthesis is non-discriminating

towards cytosolic tRNAs. The organellar GluRS is also necessary for mis-aminoacylation of tRNA^{Gln} prior

to transamidation of Glu to Gln by the GatCAB complex(96). Cytosolic GlnRS is retargeted to the

mitochondria in *S. himalayana*, thus allowing for direct aminoacylation of tRNA^{Gln} assuming there is also

import of cytosolic tRNA[Gln]. This likely precipitated the loss of the GatCAB complex and negated the

need for a non-discriminating organellar GluRS.

At the other extreme, there was repeated loss of organellar AlaRS, which was absent from all

heterotrophs and the hemiparasitic *Cuscuta* species, possibly because plastomes of heterotrophs are

often AT-rich and thus have underrepresentation of amino acids with GC-rich codons like Ala, Gly and

Pro (47, 48). Thus, the corresponding organellar aaRS and tRNAs may be subject to decreased functional

constraint and be less likely to be retained. This hypothesis is supported by the fact that we also see

frequent loss of organellar GlyRS and ProRS in heterotrophs (**Fig. 1**).

Our results indicate that imported cytosolic aaRS and tRNAs are sufficient to sustain

mitochondrial translation in plants, supporting the hypothesis that the selective pressure to keep

organellar aaRS/tRNAs is maintained by translational demands associated with photosynthesis. Yet

other eukaryote lineages that never bore plastids often still retain organelle-specific tRNAs and aaRS for

their mitochondria (8, 9, 22, 24). This is even more puzzling when considering that various eukaryotic

lineages have jettisoned nearly all their mitochondrial tRNAs/aaRS, including cnidarians, ctenophores,

some fungi, and trypanosomes (12, 24, 97). One possible explanation is the logical parallel to our finding

that photosynthesis governs plastid translation rates and maintains gene redundancy; eukaryotic

lineages with high respiratory rates may require high mitochondrial gene expression and, therefore,

retain mitochondria-specific translation machinery. Indeed, mitochondrial gene expression tracks well

with respiratory rate and cellular activity across different tissues and metabolic states (98–100). This

connection to cell energetics would also explain why mitochondrial tRNAs/aaRS are dispensable in

parasitic eukaryotes like trypanosomes and heterotrophic plants, which presumably have lower energy

demands. By studying non-model plant species with unusual physiology and life histories, we were able

to link plastid bioenergetic processes to the evolution of both organellar and nuclear genomes. The

sequencing of mitogenomes and characterization of metabolism in non-model eukaryotes will provide

exciting opportunities to conduct similar studies to better understand how mitochondrial metabolism

effects gene redundancy and elucidate the principles governing endosymbiotic integration.

Methods

Data acquisition and ortholog search

Gene models for species used in the study were collected from publicly available data sources

(**Supplemental Dataset 1**). When protein sequences were not available, we generated sequences from

transcriptome data using Transdecoder v5.5.0 (101), applying a minimum protein length of 50 amino

acids and retaining only protein sequences with blast hits to any sequence in the Uniprot Taxonomic

Divisions Plant database (https://ftp.uniprot.org/pub/databases/uniprot/knowledgebase

/taxonomic_divisions/). Then, the Transdecoder primary_transcript.fa script was used to extract only

the longest translated protein sequence from the primary transcript for each gene to reduce the search

space. SeqKit 2.3.0 (102) was used to create uniform gene handles for all files and remove stop codons

from sequences. Orthofinder v2.5.5 (37) was run on all sets of protein models with a user-defined

species tree (see **Fig. 1** for tree topology) to find groups of likely orthologous proteins, i.e. "orthogroups"

in two steps as follows:

444   orthofinder.py -f <folder_with_protein_models> -t 24 -a 6

445   orthofinder.py -ft <folder_with_results_> -s userdefined_speciestree.txt -t 24 -a 6

Orthogroups containing proteins of interest including cytosolic/organellar aaRS and tRNA-processing

enzymes characterized in *A. thaliana* (**Supplemental Dataset 2**) were extracted and checked for

completeness via RBH against all protein models for a given species using BLASTP 2.15.0+ (103) and

custom Python/Bash scripts. After ortholog discovery via Orthofinder, full sets of protein models (not

just the longest) were searched for *A. thaliana* protein sets using BLASTP to ensure that narrowing of the

search space for Orthofinder had not resulted in loss of matching gene models. In cases where available

curated protein models excluded alternative transcripts, transcriptome data were obtained and

Transdecoder was run to predict protein-coding sequences (see **Supplemental Dataset 1** for details).

Note that ArgRS was not included in the ortholog search as there are no distinct organellar versions of

this enzyme in *A. thaliana.* One ArgRS gene product is cytosol-targeted, and the other is triple-targeted

to cytosol, mitochondria, and plastids (9, 16). In the case of GlnRS, there is no organellar ortholog as

plants typically use the bacterial-type GatCAB amidotransferase system to charge Gln-tRNA indirectly in

mitochondria and plastids (42). For CysRS, tRNase Z and PRORP, sequences for both cytosolic and

organellar proteins were clustered together in one orthogroup likely due to recent shared ancestry. For

these genes, cytosolic vs. organellar versions were differentiated by aligning sequences in each

orthogroup using MAFFT v7.525 (104) with the --auto setting, followed by trimming of the alignment

using trimAl v1.5.rev0 on the -automated1 setting (105). Alignments were then viewed using MEGA 11

(Tamura et al., 2021). Maximum-likelihood tree building was conducted using RAxML v 8.2.12 (107) as

follows:

raxmlHPC-PTHREADS-SSE3 -f a -# 100 -p 12345 -x 12345 -s <fasta file with orthogroup sequences> -n

<orthogroup> -m PROTGAMMALG -T 12

Inclusion in cytosolic versus organellar groups was then determined by relationship to known *A. thaliana*

cytosolic/nuclear/organellar proteins in maximum-likelihood trees (**Fig. S8-10**).

469   Once we had a curated set of protein sequences for all species, they were checked for

completeness of length by alignment with *A. thaliana* orthologs by MAFFT and subsequent visualization

with a custom Python script. A "full-length" ortholog was defined as having at least 50% alignment with

any *A. thaliana* protein in the orthogroup, excluding pseudogenes. Custom Python and Bash scripts are

available via Dryad (https://doi.org/10.5061/dryad.0cfxpnw7p).

In silico targeting predictions

For TP analysis, all protein models available for each gene with similarity to the original Orthofinder hit

(BLASTP e-value < 1e-6) were used, not just the longest. All transcripts for cytosolic aaRS, a "positive

control" set of genes involved in mitochondrial oxidative phosphorylation/ plastid housekeeping and a

"negative control" set of genes involved in cell-wall synthesis were run through targeting prediction

programs LOCALIZER-1.0.5 (76) and TargetP-2.0 (75), with commands as follows:

LOCALIZER.py -p -i <input file>

targetp -fasta <input file> -mature -org pl

Protein models for some species were often incomplete due to the fragmented nature of some of the

transcriptome assemblies. To ensure models missing the N-terminus were not used in retargeting

analysis, predictions for models that did not align without gaps for 15 residues within 100 amino acids of

the start codon for at least one *A. thaliana* ortholog were discarded. For transcript-by-transcript

heatmaps (**Supplemental Dataset 6**), a custom R script was used to plot all transcripts for all species. For

Fig. S3 (which shows predicted targeting for a multitude of different proteins) only the most retargeted

ortholog for each species was used. Orthologs were then binned based on probability of targeting to the

mitochondria or plastids. We chose a probability of 0.5 as a threshold for LOCALIZER to match the

original publication describing the tool(76). TargetP-2.0 uses a "winner take all" strategy to declare one

subcellular localization based on the location with the highest likelihood score, which is not compatible

with assessing dual targeting (75, 108). Thus, we used the lowest likelihood score for known organelle-

targeted *A. thaliana* proteins from our control dataset identified by TargetP-2.0 to be plastid or

mitochondria targeted (~0.5 for AT1G02410 cytochrome c oxidase 11). High confidence retargeting was

defined as a > 0.5 probability/likelihood of targeting for both LOCALIZER and TargetP-2.0. Moderate

confidence was defined as a > 0.5 probability/likelihood for one of the algorithms. Low confidence

retargeting was defined as neither algorithm providing a probability/likelihood > 0.5. For subsequent

analysis of ratios (**Fig. 2, Fig. S1-S2**), any ortholog with a "moderate" or better confidence was counted

as retargeted. Note that *R. cantleyi* and *S. himalayana* proteins were excluded from predictions of

targeting to plastids as these species likely lack plastomes. All analyses including plotting and statistics

were carried out using custom Bash, Python, and R scripts available via Dryad

(https://doi.org/10.5061/dryad.6hdr7sr5x).

In vivo localization of S. himalayana aaRS transit peptides via tobacco infiltration and confocal

microscopy

TPs were chosen based on having an at least moderate probability (see above) of targeting a protein to

either the mitochondria or plastid and/or having an N-terminal extension preceding conserved sequence

shared with *A. thaliana.* We selected either the sequence predicted to be a TP by TargetP-2.0 plus 10

residues or the entire N-terminal overhang plus the first 10 residues aligning with the functional portion

of the *A. thaliana* protein. These amino acid sequences were then converted into DNA sequences using

EMBOSS Backtranseq (109) with the *Nicotiana tabacum* codon usage table (see **Supplemental Dataset**

3). These sequences were then used to generate constructs for *Agrobacterium* transformation using a

newly designed dual-expression vector containing both the mitochondrial control IVD-FP611 and GFP

fused to the C-terminus of our TP of interest. Some TPs were also originally cloned into the single-gene

expression vector described in (55). Transformation into *N. benthamiana* and confocal microscopy was

constructed as described in (55). With our dual expression vectors, there was occasionally low GFP

and/or FP611 signal relative to chlorophyll autofluorescence (see Fig. 3 GlyRS and MetRS). Thus, we

scaled pixel intensity using look-up tables (LUTs) for all images to better see localization for low

expression constructs. It is important to note that in cases where GFP fluorescence is low and LUTs are

scaled, overlap of the chlorophyll florescence and GFP spectra can create the false impression of plastid

GFP targeting (see negative control in Fig. S4 for an example). Thus, we only determined a TP to be

plastid targeted if we either observed 1) GFP-fluorescing stromules in transformed plastids, an

indication of true GFP signal as these structures occlude thylakoids or 2) Increased GFP florescence in

plastids of transformed cells compared to untransformed cells in the same frame. Often guard cells

would not be transformed, so we also used these as an in-frame negative control. Note that *in silico*

predictions shown are for the TP attached to GFP, not the original protein model. Relative to the whole

protein models, the GFP predictions yielded higher targeting probabilities and affected classification for

MetRS (LOCALIZER likelihood score > 0.5 for mitochondria) and AsnRS (TargetP-2.0 likelihood score > 0.5

for plastid), but there were no substantive changes for any of the other transit peptides. For more

information see supplemental dataset 3. Constructs are available from Addgene; see supplemental

dataset 3 for ID numbers.

Mitogenome assembly

Prior to tRNA analysis, the mitogenomes of *S. himalayana* (SRR629601) and *R. cantleyi* (SRR629613)

were assembled *de novo* from raw reads available from the NCBI SRA database. To maximize the

likelihood of finding all mitogenome tRNAs, genomes were assembled in several ways. Reads were first

trimmed using FASTP (110), with either default parameters or "harsh" trimming where the --

qualified_quality_phred and --unqualified_percent_limit parameters were set to 20 and 10, respectively.

SPAdes v.3.15.4 (111) was then used to assemble trimmed reads *de novo,* with kmer sizes of 21, 33, 55,

and 77. To avoid potential assembly fragmentation due to excessive levels of sequence coverage, a

subset of reads (1,000,000 from S. *himalayana* and 500,000 from *R. cantleyi*) were sampled and

assembled from each species as well.

tRNA analysis

Excepting the Rafflesiaceae, for which we assembled mitogenomes *de novo,* the GenBank (.gbk) files for

mitogenomes and plastomes from species of interest were downloaded from NCBI (**Supplemental**

Dataset 1). A custom Perl script was used to extract annotated tRNAs from the .gbk files. tRNAscan-SE-

2.0.9 (112, 113) was run on downloaded genomes (parameters -BQH), and results were cross-checked

with preexisting GenBank annotations (**Supplemental Dataset 4**). For *S. himalayana* and *R. cantleyi*, the

same tRNAscan-SE-2.0.9 pipeline was used on all generated assemblies, and the consensus set of tRNAs

was checked for plant origin using BLASTN against all NCBI reference sequences and the *A. thaliana*

mitogenome (NC_037304.1) and plastome (NC_000932.1). tRNA presence/absence patterns were

plotted along with the corresponding aaRS presence/absence/retargeting data using a custom Python

script. Scripts are available via Dryad (https://doi.org/10.5061/dryad.np5hqc009). The highly divergent

and truncated trnE for *B. fungosa* was added based on prior publication(46).

Rate Analysis

Primary protein sequences for cytosolic and organellar aaRS from heterotroph *M. hypopitys* and related

autotroph *R. williamsianum* from the original Orthofinder search were used for comparison of rates,

with *A. thaliana* orthologs forming the outgroup. CysRS, LeuRS, ThrRS, and TrpRS were excluded from

the analysis because the organellar ortholog was missing or too short to create informative gene trees in

one or both test species. Predicted TPs were trimmed from sequences using output of TargetP-2.0.

Sequences were then aligned using MAFFT v7.490 (104) and trimmed manually in MEGA 11 (106).

Maximum likelihood trees were built using RAxML v 8.2.12 (107) as follows:

raxmlHPC-PTHREADS-SSE3 -f a -# 100 -p 12345 -x 12345 -s <orthogroup_aligned.fa> -n <orthogroup> -m

PROTGAMMALG -T 12

For orthogroups containing only three ortholog sequences, a "dummy" outgroup sequence represented

by the *A. thaliana* ortholog with one branch-specific substitution was created to overcome RAxML's

input requirement of four or more taxa. The resulting trees were then evaluated for topology. In

instances where species-specific paralogs were not sister to one another, separate multiple sequence

alignments and trees were re-made for each paralog as described above. In instances where species-

specific paralogs were sister, both sequences were retained in the same tree. Trees were rooted to the

A. thaliana ortholog(s), then branch lengths to the last shared node from each *H. monotropa* and *R.*

williamsianum ortholog were extracted from the trees. For instances where two sister paralogs from the

same species were retained, the branch lengths were averaged. Then, the $\log_{10}$-transformed ratio of the

H. monotropa/R williamsianum branches was plotted, and mean branch lengths for cytosolic and

organellar enzymes were calculated. Statistical difference in mean branch ratio in cytosolic vs. organellar

enzymes was tested using a *t*-test. Tajima's relative rate tests were run in MEGA 11 using *A. thaliana* as

the outgroup. Data is available in **Supplemental Dataset 5.** Scripts, alignments, and maximum likelihood

trees are available via Dryad (https://doi.org/10.5061/dryad.tb2rbp05r).

Acknowledgements

We thank Jessica Warren for helpful discussion and feedback on this project and Anna Pratt for helpful

advice on the cloning approach. We thank undergraduate researchers Xiaorui Lou and Janna Novak for

their assistance. This work was supported by grants MCB-2322154, MCB-2048407, and MCB-1933590

from the National Science Foundation (NSF) and utilized the RMACC Summit supercomputer (NSF ACI-

1532235 and ACI-1532236). RAD was supported by an NSF Postdoctoral Research Fellowship in Biology

(IOS-2208908). JC was supported by the University of Nebraska Foundation.

References

1.　A. N. Khachane, K. N. Timmis, V. A. P. Martins Dos Santos, Dynamics of reductive genome
evolution in mitochondria and obligate intracellular microbes. *Mol Biol Evol* **24**, 449–456 (2007).

2.　P. J. Keeling, J. D. Palmer, Horizontal gene transfer in eukaryotic evolution. *Nat Rev Genet* **9**, 605–
618 (2008).

3.　W. Martin, *et al.*, Evolutionary analysis of Arabidopsis, cyanobacterial, and chloroplast genomes
reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. *PNAS* **99**,
12246–12251 (2002).

4.　E. C. M. Nowack, *et al.*, Endosymbiotic gene transfer and transcriptional regulation of transferred
genes in Paulinella chromatophora. *Mol Biol Evol* **28**, 407–422 (2011).

5.　N. A. Moran, J. P. McCutcheon, A. Nakabachi, Genomics and evolution of heritable bacterial
symbionts. *Annu Rev Genet* **42**, 165–190 (2008).

6.　J. P. McCutcheon, N. A. Moran, Extreme genome reduction in symbiotic bacteria. *Nat Rev*
Microbiol **10**, 13–26 (2012).

7.　V. Rajendran, P. Kalita, H. Shukla, A. Kumar, T. Tripathi, Aminoacyl-tRNA synthetases: Structure,
function, and drug discovery. *Int J Biol Macromol* **111**, 400–414 (2018).

8.   T. Salinas-Giegé, R. Giegé, P. Giegé, TRNA biology in mitochondria. *Int J Mol Sci* **16**, 4518–4559
(2015).

9.   A. M. Duchêne, C. Pujol, L. Maréchal-Drouard, Import of tRNAs and aminoacyl-tRNA synthetases
into mitochondria. *Curr Genet* **55**, 1–18 (2009).

10.   L. Bonnefond, *et al.*, Toward the full set of human mitochondrial aminoacyl-tRNA synthetases:
Characterization of AspRS and TyrRS. *Biochemistry* **44**, 4805–4816 (2005).

11.   E. Tolkunova, H. Park, J. Xia, M. P. King, E. Davidson, The human lysyl-tRNA synthetase gene
encodes both the cytoplasmic and mitochondrial enzymes by means of an unusual: Alternative
splicing of the primary transcript. *Journal of Biological Chemistry* **275**, 35063–35069 (2000).

12.   S. Shikha, R. Brogli, A. Schneider, N. Polacek, TRNA biology in trypanosomes. *Chimia (Aarau)* **73**,
395–405 (2019).

13.   T. H. P. Tan, R. Pach, A. Crausaz, A. Ivens, A. Schneider, tRNAs in Trypanosoma brucei : Genomic
Organization, Expression, and Mitochondrial Import . *Mol Cell Biol* **22**, 3707–3717 (2002).

14.   J. M. Warren, *et al.*, Rapid Shifts in Mitochondrial tRNA Import in a Plant Lineage with Extensive
Mitochondrial tRNA Gene Loss. *Mol Biol Evol* **38**, 5735–5751 (2021).

15.   J. M. Warren, D. B. Sloan, Interchangeable parts: The evolutionarily dynamic tRNA population in
plant mitochondria. *Mitochondrion* **52**, 144–156 (2020).

16.   A.-M. Duchêne, *et al.*, Dual targeting is the rule for organellar aminoacyl-tRNA synthetases in
Arabidopsis thaliana. *PNAS* **102**, 16484–16489 (2005).

17.   K. Fujishima, A. Kanai, tRNA gene diversity in the three domains of life. *Front Genet* **5** (2014).

18.   M. Lynch, B. Koskella, S. Schaack, Mutation Pressure and the Evolution of Organelle Genomic
Architecture. *Science (1979)* **311**, 1727–1730 (2006).

19.   S. Schaack, E. K. H. Ho, F. MacRae, Disentangling the intertwined roles of mutation, selection and
drift in the mitochondrial genome. *Philosophical Transactions of the Royal Society B: Biological
Sciences* **375** (2020).

20.   K. Crosby, D. R. Smith, Does the Mode of Plastid Inheritance Influence Plastid Genome
Architecture? *PLoS One* **7** (2012).

21.   K. H. Wolfe, W.-H. Li, P. M. Sharp, Rates of nucleotide substitution vary greatly among plant
mitochondrial, chloroplast, and nuclear DNAs. *Proc. Nati. Acad. Sci. USA* **84**, 9054–9058 (1987).

22.   J. R. Adrion, P. S. White, K. L. Montooth, The roles of compensatory evolution and constraint in
aminoacyl trna synthetase evolution. *Mol Biol Evol* **33**, 152–161 (2016).

23.   F. S. Barreto, *et al.*, Genomic signatures of mitonuclear coevolution across populations of
Tigriopus californicus. *Nat Ecol Evol* **2**, 1250–1257 (2018).

24.   W. Pett, D. V Lavrov, Cytonuclear Interactions in the Evolution of Animal Mitochondrial tRNA
Metabolism. *Genome Biol Evol* **7**, 2089–2101 (2015).

25. D. B. Sloan, R. A. DeTar, J. M. Warren, Aminoacyl-tRNA synthetase evolution within the dynamic
tripartite translation system of plant cells. *Genome Biol Evol* **15** (2023).

26. S. Wicke, G. M. Schneeweiss, C. W. dePamphilis, K. F. Müller, D. Quandt, The evolution of the
plastid chromosome in land plants: Gene content, gene order, gene function. *Plant Mol Biol* **76**,
273–297 (2011).

27. M. Michaud, V. Cognat, A. M. Duchêne, L. Maréchal-Drouard, A global picture of tRNA genes in
plant genomes. *Plant Journal* **66**, 80–93 (2011).

28. E. S. Forsythe, *et al.*, Organellar transcripts dominate the cellular mRNA pool across plants of
varying ploidy levels. *PNAS* **119** (2022).

29. J. Mergner, *et al.*, Mass-spectrometry-based draft of the Arabidopsis proteome. *Nature* **579**, 409–
414 (2020).

30. J. Těšitel, Functional biology of parasitic plants: A review. *Plant Ecol Evol* **149**, 5–20 (2016).

31. L. Teixeira-Costa, C. C. Davis, Life history, diversity, and distribution in parasitic flowering plants.
Plant Physiol **187**, 32–51 (2021).

32. S. Wicke, J. Naumann, "Molecular Evolution of Plastid Genomes in Parasitic Flowering Plants" in
Advances in Botanical Research, (Academic Press Inc., 2018), pp. 315–347.

33. D. R. Smith, S. R. Asmail, Next-generation sequencing data suggest that certain
nonphotosynthetic green plants have lost their plastid genomes. *New Phytologist* **204**, 7–11
(2014).

34. H. T. Li, *et al.*, Plastid phylogenomic insights into relationships of all flowering plant families. *BMC
Biol* **19** (2021).

35. J. Molina, *et al.*, Possible loss of the chloroplast genome in the parasitic flowering plant Rafflesia
lagascae (Rafflesiaceae). *Mol Biol Evol* **31**, 793–803 (2014).

36. D. L. Nickrent, Y. Ouyang, R. J. Duff, C. W. Depamphilis, Do nonasterid holoparasitic flowering
plants have plastid genomes? *Plant Mol Biol* **34**, 717–729 (1997).

37. D. M. Emms, S. Kelly, OrthoFinder: Phylogenetic orthology inference for comparative genomics.
Genome Biol **20** (2019).

38. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol* **16** (2015).

39. G. Canino, *et al.*, Arabidopsis encodes four tRNase Z enzymes. *Plant Physiol* **150**, 1494–1502
(2009).

40. B. Gutmann, A. Gobert, P. Giegé, PRORP proteins support RNase P activity in both organelles and
the nucleus in Arabidopsis. *Genes Dev* **26**, 1022–1027 (2012).

41. A. Gobert, *et al.*, A single Arabidopsis organellar protein has RNase P activity. *Nat Struct Mol Biol*
17, 740–744 (2010).

42. C. Pujol, *et al.*, Dual-targeted tRNA-dependent amidotransferase ensures both mitochondrial and
chloroplastic Gln-tRNAGln synthesis in plants. *Proc Natl Acad Sci U S A* **105**, 6481–6485 (2008).

43. A. Nagao, T. Suzuki, T. Katoh, Y. Sakaguchi, T. Suzuki, Biogenesis of glutaminyl-mt tRNA Gln in
human mitochondria. *PNAS* **106**, 16209–16214 (2009).

44. L. Echevarría, *et al.*, Glutamyl-tRNAGln amidotransferase is essential for mammalian
mitochondrial translation in vivo. *Biochemical Journal* **460**, 91–101 (2014).

45. R. Giegé, G. Eriani, The tRNA identity landscape for aminoacylation and beyond. *Nucleic Acids Res*
51, 1528–1570 (2023).

46. L. F. Ceriotti, M. E. Roulet, M. V. Sanchez-Puerta, Plastomes in the holoparasitic family
Balanophoraceae: Extremely high AT content, severe gene content reduction, and two
independent genetic code changes. *Mol Phylogenet Evol* **162** (2021).

47. H. J. Su, *et al.*, Novel genetic code and record-setting AT-richness in the highly reduced plastid
genome of the holoparasitic plant Balanophora. *Proc Natl Acad Sci U S A* **116**, 934–943 (2019).

48. M. I. Schelkunov, M. S. Nuraliev, M. D. Logacheva, Rhopalocnemis phalloides has one of the most
reduced and mutated plastid genomes known. *PeerJ* **2019** (2019).

49. X. Chen, *et al.*, Comparative Plastome Analysis of Root-and Stem-Feeding Parasites of Santalales
Untangle the Footprints of Feeding Mode and Lifestyle Transitions. *Genome Biol Evol* **12**, 3663–
3676 (2019).

50. L. Cai, *et al.*, Deeply Altered Genome Architecture in the Endoparasitic Flowering Plant Sapria
himalayana Griff. (Rafflesiaceae). *Current Biology* **31**, 1002-1011.e9 (2021).

51. M. D. Logacheva, M. I. Schelkunov, V. Y. Shtratnikova, M. V. Matveeva, A. A. Penin, Comparative
analysis of plastid genomes of non-photosynthetic Ericaceae and their photosynthetic relatives.
Sci Rep **6** (2016).

52. M. I. Schelkunov, M. S. Nuraliev, M. D. Logacheva, Genomic comparison of non-photosynthetic
plants from the family Balanophoraceae with their photosynthetic relatives. *PeerJ* **9** (2021).

53. L. F. Ceriotti, L. Gatica-Soria, M. V. Sanchez-Puerta, Cytonuclear coevolution in a holoparasitic
plant with highly disparate organellar genomes. *Plant Mol Biol* **109**, 673–688 (2022).

54. A. C. Barbrook, C. J. Howe, S. Purton, Why are plastid genomes retained in non-photosynthetic
organisms? *Trends Plant Sci* **11**, 101–108 (2006).

55. J. M. Warren, *et al.*, Rewiring of aminoacyl-tRNA synthetase localization and interactions in plants
with extensive mitochondrial tRNA gene loss. *Mol Biol Evol* **40** (2023).

56. B. Bölter, En route into chloroplasts: preproteins' way home. *Photosynth Res* **138**, 263–275
(2018).

57. P. Chotewutmontri, K. Holbrook, B. D. Bruce, "Plastid Protein Targeting: Preprotein Recognition
and Translocation" in *International Review of Cell and Molecular Biology*, (Elsevier Inc., 2017), pp.
227–294.

58.   M. W. Murcha, *et al.*, Protein import into plant mitochondria: Signals, machinery, processing, and
709   regulation. *J Exp Bot* **65**, 6301–6335 (2014).

710   59.   R. W. Christian, S. L. Hewitt, G. Nelson, E. H. Roalson, A. Dhingra, Plastid transit peptides-where
711   do they come from and where do they all belong? Multi-genome and pan-genomic assessment of
712   chloroplast transit peptide evolution. *PeerJ* **8** (2020).

713   60.   S. Reinbothe, *et al.*, Trna-dependent import of a transit sequence-less aminoacyl-trna synthetase
714   (Leurs2) into the mitochondria of arabidopsis. *Int J Mol Sci* **22** (2021).

715   61.   W. Neupert, J. M. Herrmann, Translocation of proteins into mitochondria. *Annu Rev Biochem* **76**,
716   723–749 (2007).

717   62.   M. H. Chen, L. F. Huang, H. M. Li, Y. R. Chen, S. M. Yu, Signal peptide-dependent targeting of a
718   rice α-amylase and cargo proteins to plastids and extracellular compartments of plant cells. *Plant
719   Physiol* **135**, 1367–1377 (2004).

720   63.   L. G. L. Richardson, D. J. Schnell, Origins, function, and regulation of the TOC-TIC general protein
721   import machinery of plastids. *J Exp Bot* **71**, 1226–1238 (2020).

722   64.   H. Inoue, C. Rounds, D. J. Schnell, The molecular basis for distinct pathways for protein import
723   into Arabidopsis chloroplasts. *Plant Cell* **22**, 1947–1960 (2010).

724   65.   S. Bischof, *et al.*, Plastid proteome assembly without Toc159: Photosynthetic protein import and
725   accumulation of N-Acetylated plastid precursor proteins. *Plant Cell* **23**, 3911–3928 (2011).

726   66.   Y. L. Chen, *et al.*, TIC236 links the outer and inner membrane translocons of the chloroplast.
727   *Nature* **564**, 125–129 (2018).

728   67.   A. R. Kasmati, *et al.*, Evolutionary, Molecular and Genetic Analyses of Tic22 Homologues in
729   Arabidopsis thaliana Chloroplasts. *PLoS One* **8** (2013).

730   68.   J. Tripp, *et al.*, Structure and conservation of the periplasmic targeting factor Tic22 protein from
731   plants and cyanobacteria. *Journal of Biological Chemistry* **287**, 24164–24173 (2012).

732   69.   M. Rudolf, *et al.*, In vivo function of Tic22, a protein import component of the intermembrane
733   space of chloroplasts. *Mol Plant* **6**, 817–829 (2013).

734   70.   A. R. Kasmati, M. Töpel, R. Patel, G. Murtaza, P. Jarvis, Molecular and genetic analyses of Tic20
735   homologues in Arabidopsis thaliana chloroplasts. *Plant Journal* **66**, 877–889 (2011).

736   71.   E. Kovács-Bogdán, J. P. Benz, J. Soll, B. Bölter, Tic20 forms a channel independent of Tic110 in
737   chloroplasts. *BMC Plant Biol* **11** (2011).

738   72.   H. Liu, A. Li, J. D. Rochaix, Z. Liu, Architecture of chloroplast TOC–TIC translocon supercomplex.
739   *Nature* **615**, 349–357 (2023).

740   73.   X. Guo, H. Wang, D. Lin, Y. Wang, X. Jin, Cytonuclear evolution in fully heterotrophic plants:
741   lifestyles and gene function determine scenarios. *BMC Plant Biol* **24**, 989 (2024).

74. C. Carrie, J. Whelan, Widespread dual targeting of proteins in land plants: When, where, how and
why. *Plant Signal Behav* **8** (2013).

75. J. J. A. Armenteros, *et al.*, Detecting sequence signals in targeting peptides using deep learning.
Life Sci Alliance **2** (2019).

76. J. Sperschneider, *et al.*, LOCALIZER: Subcellular localization prediction of both plant and effector
proteins in the plant cell. *Sci Rep* **7** (2017).

77. M. Sharma, B. Bennewitz, R. B. Klösgen, Rather rule than exception? How to evaluate the
relevance of dual protein targeting to mitochondria and chloroplasts. *Photosynth Res* **138**, 335–
343 (2018).

78. L. Xu, C. Carrie, S. R. Law, M. W. Murcha, J. Whelan, Acquisition, conservation, and loss of dual-
targeted proteins in land plants. *Plant Physiol* **161**, 644–662 (2013).

79. D. B. Sloan, J. M. Warren, A. M. Williams, S. A. Kuster, E. S. Forsythe, Incompatibility and
Interchangeability in Molecular Evolution. *Genome Biol Evol* **15** (2023).

80. Y. R. Li, M. J. Liu, Prevalence of alternative AUG and non-AUG translation initiators and their
regulatory effects across plants. *Genome Res* **30**, 1418–1433 (2020).

81. S. K. Tanz, I. Castleden, I. D. Small, A. Harvey Millar, Fluorescent protein tagging as a tool to
define the subcellular distribution of proteins in plants. *Front Plant Sci* **4** (2013).

82. J. Jeong, B. Moon, I. Hwang, D. W. Lee, GREEN FLUORESCENT PROTEIN variants with enhanced
folding are more efficiently imported into chloroplasts. *Plant Physiol* **190**, 238–249 (2022).

83. M. Sharma, B. Bennewitz, R. B. Klösgen, Dual or not dual?—comparative analysis of fluorescence
microscopy-based approaches to study organelle targeting specificity of nuclear-encoded plant
proteins. *Front Plant Sci* **9** (2018).

84. M. W. Murcha, *et al.*, Plant-specific preprotein and amino acid transporter proteins are required
for tRNA import into mitochondria. *Plant Physiol* **172**, 2471–2490 (2016).

85. E. Skippington, T. J. Barkman, D. W. Rice, J. D. Palmer, Miniaturized mitogenome of the parasitic
plant viscum scurruloideum is extremely divergent and dynamic and has lost all nad genes. *PNAS*
112, E3515–E3524 (2015).

86. G. Petersen, A. Cuenca, I. M. Møller, O. Seberg, Massive gene loss in mistletoe (Viscum,
Viscaceae) mitochondria. *Sci Rep* **5** (2015).

87. A. Fender, *et al.*, Loss of a primordial identity element for a mammalian mitochondrial
aminoacylation system. *Journal of Biological Chemistry* **281**, 15980–15986 (2006).

88. Y. Kumazawa, H. Himeno, K.-I. Miura, K. Watanabe, Unilateral Aminoacylation Specificity
between Bovine Mitochondria and Eubacteria. *The Journal of Biochemistry* **109**, 421–427 (1991).

89. L. Bonnefond, M. Frugier, R. Giegé, J. Rudinger-Thirion, Human mitochondrial TyrRS disobeys the
tyrosine identity rules. *RNA* **11**, 558–562 (2005).

90.   I. Small, *et al.*, The Strange Evolutionary History of Plant Mitochondrial tRNAs and Their
Aminoacyl-tRNA Synthetases. *J Hered* **90:3**, 333–337 (1999).

91.   L. Bromham, P. F. Cowman, R. Lanfear, Parasitic plants have increased rates of molecular
evolution across all three genomes. *BMC Evol Biol* **13** (2013).

92.   M. Safro, N. Moor, O. Lavrik, Phenylalanyl-tRNA Synthetases. *Madame Curie Bioscience Database*
(2013).

93.   R. Tanaka, K. Kobayashi, T. Masuda, Tetrapyrrole Metabolism in Arabidopsis thaliana. *Arabidopsis*
Book (2011). https://doi.org/10.1199/tab.0145.

94.   A. Schon, *et al.*, The RNA required in the first step of chlorophyll biosynthesis is a chloroplast
glutamate transfer-RNA. *Nature* **322**, 281–284 (1986).

95.   X. Chen, *et al.*, Balanophora genomes display massively convergent evolution with other extreme
holoparasites and provide novel insights into parasite–host interactions. *Nat Plants* (2023).
https://doi.org/10.1038/s41477-023-01517-7.

96.   A. Schon, C. G. Kannangara, S. Gough, D. Sollt, Protein biosynthesis in organelles requires
misaminoacylation of tRNA. (1988).

97.   K. M. Haen, W. Pett, D. V. Lavrov, Parallel loss of nuclear-encoded mitochondrial aminoacyl-tRNA
synthetases and mtDNA-encoded tRNAs in cnidaria. *Mol Biol Evol* **27**, 2216–2219 (2010).

98.   I. Martin, O. Vinas, T. Mampel, F. Villarroya, Effects of cold environment on mitochondrial
genome expression in the rat: evidence for a tissue-specific increase in the liver, independent of
changes in mitochondrial gene abundance. *Biochemical Journal* **296**, 231–234 (1993).

99.   R. F. Hevner, M. T. T. Wong-Riley, Mitochondrial and Nuclear Gene Expression for Cytochrome
Oxidase Subunits Are Disproportionately Regulated by Functional Activity in Neurons. *The Journal*
of Neuroscience **13**, 1805–1819 (1993).

100. E. A. Amiott, J. A. Jaehning, Mitochondrial Transcription Is Regulated via an ATP "Sensing"
Mechanism that Couples RNA Abundance to Respiration. *Mol Cell* **22**, 329–338 (2006).

101. B. J. Haas, Transdecoder v5.5.0. (2018). Available at:
https://github.com/TransDecoder/TransDecoder [Accessed 14 May 2023].

102. W. Shen, S. Le, Y. Li, F. Hu, SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
manipulation. *PLoS One* **11** (2016).

103. C. Camacho, *et al.*, BLAST+: Architecture and applications. *BMC Bioinformatics* **10** (2009).

104. K. Katoh, K. Misawa, K.-I. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059–3066 (2002).

105. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: A tool for automated alignment
trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

106. K. Tamura, G. Stecher, S. Kumar, MEGA11: Molecular Evolutionary Genetics Analysis Version 11.
Mol Biol Evol **38**, 3022–3027 (2021).

107. A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

108. O. Emanuelsson, H. Nielsen, S. Brunak, G. Von Heijne, Predicting subcellular localization of
proteins based on their N-terminal amino acid sequence. *J Mol Biol* **300**, 1005–1016 (2000).

109. F. Madeira, *et al.*, Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic
Acids Res* **50**, W276–W279 (2022).

110. S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor in
Bioinformatics, (Oxford University Press, 2018), pp. i884–i890.

111. A. Bankevich, *et al.*, SPAdes: A new genome assembly algorithm and its applications to single-cell
sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

112. P. P. Chan, T. M. Lowe, "tRNAscan-SE: Searching for tRNA genes in genomic sequences" in
Methods in Molecular Biology, (Humana Press Inc., 2019), pp. 1–14.

113. P. P. Chan, B. Y. Lin, A. J. Mak, T. M. Lowe, TRNAscan-SE 2.0: Improved detection and functional
classification of transfer RNA genes. *Nucleic Acids Res* **49**, 9077–9096 (2021).

Supplemental Dataset 1: Species and datasets used for gene presence/absence, retargeting analysis, and tRNA content analysis.

Supplemental Dataset 2. Genes of interest and counts of gene copies for each species.

Supplemental Dataset 3: *S. himalayana* transit peptide sequences for transient expression and colocalization in *N. benthamiana*

Supplemental Dataset 4: tRNA content of organelle genomes.

Supplemental Dataset 5: Relative branch lengths and Tajima's relative rate test for aaRS in Ericales spp.

Supplemental Dataset 6: Heatmaps for probability of subcellular targeting to plastids or mitochondria on a protein-by-protein basis.

Figure 1: Presence/absence matrix of selected organellar and cytosolic aaRS and tRNA-processing enzymes shows major losses of organellar enzymes in heterotrophic lineages. Hue of boxes indicates metabolism (gray = autotroph, orange = hemiparasite, purple = heterotroph), shade indicates degree of presence/absence (white = no sequence found, light colors = partial sequence found, dark colors = complete or near complete sequence found). Note that there is no organellar GlnRS as plants typically use the bacterial-type GatCAB amidotransferase system to charge organellar Gln-tRNA. ArgRS is not presented as it has no distinct organellar enzyme; gene products are either cytosol-targeted, or triple-targeted to the cytosol, mitochondria, and plastids.

Figure 2: Proportion of retargeted cytosolic aaRS is higher in heterotrophs than autotrophs based on *in-silico* predictions. Proportion of enzymes with predicted retargeting (probability ≥ 0.5) to either mitochondria (top row) or plastids (bottom row) based on the outputs of LOCALIZER and/or TargetP. Autotrophs and heterotrophs are plotted in green and purple, respectively. Targeting predictions for proteins involved in mitochondrial oxidative phosphorylation (OXPHOS) are included as a positive control for mitochondrial targeting, and proteins involved in non-photosynthesis related housekeeping roles (ACC and Clp subunits) are included as a positive control for plastid targeting. Predictions for cell-wall synthesis associated enzymes are included as a negative control for organelle targeting. Reported *p*-values from *t*-tests or Mann-Whitney U tests depending on distribution of the data. Note that *R. cantleyi* and *S. himalayana* are not included in plastid retargeting analysis due to suspected loss of the plastome, but they were predicted to have retargeted 7.7% and 30.0% of their cytosolic aaRS to the plastid, respectively. These species have lost too many plastid housekeeping genes to assess proportion targeted. For both species, no cell wall synthesis enzymes exhibited plastid retargeting.

Figure 3: Cytosolic-type aaRS transit peptides from heterotroph *S. himalayana* can retarget enzymes to organelles in *Nicotiana benthamiana*. Transit peptides (TPs) from *S. himalayana* fused to GFP (middle right column) were transiently expressed in *N. benthamiana.* Chlorophyll autofluorescence (far left column) was used for colocalization with plastids. As a positive control for mitochondrial targeting, eqFP611 was tagged with the TP from mitochondrial isovaleryl-CoA dehydrogenase (middle left). The overlay of all three channels is shown on far right. Scale indicated in top right panel is the same for all images. As a means of differentiating true GFP florescence from chlorophyll autofluorescence, yellow ovals are shown in panels to indicate plastids in transformed cells, whereas white ovals indicate plastids in non-

transformed cells. Note the presence of GFP-fluorescing stromules in transformed plastids, an indication of true GFP signal. For more information, see methods and supplemental dataset 3.

Figure 4: Instances of both concordance and mismatch between organellar tRNA gene loss and aaRS loss/retargeting. Heterotrophic species are written in purple. *Arabidopsis thaliana* is included for reference. Left: mitogenome tRNAs. Right: plastome tRNAs. Color of boxes indicates presence/absence of tRNA. The – and + annotation denotes loss of corresponding organellar aaRS and retargeting of cytosolic tRNA respectively as described for Fig. 2 and Fig. S2. Retargeting predictions are specific for each organelle. Note that loss and retargeting of the corresponding aaRS was not assessed for *trnR* as there are no organelle-specific aaRS orthologs responsible for charging this tRNA. Loss of GatCAB and associated cytosolic GlnRS retargeting are presented for *trnQ*.

Supplemental Table 1: Count of Orthogroups Containing Riboprotein Ortholog(s)

Species	Plastid Specific	Mito. Specific	Cytosol Specific
Sapria himalayana	1	64	77
Rafflesia cantleyi	0	56	75
Arabidopsis thaliana	32	66	82

Supplemental Table 2: aaRS targeting and recovery of control groups

Protein Class	Mean % Mito-targeted		Mean % Plastid-targeted	
	Autotrophs	Heterotrophs	Autotrophs	Heterotrophs
Cytosolic aaRS	8.9%	20.3%	9%*	15.2%*
Mitochondrial OXPHOS	59.9%	58.5%	8.1%	8.3%
Plastid Housekeeping	20.5%	19.9%	92.6%*	86.6%*
Cell Wall Synth.	6.4%	11.2%	15.5%*	9%*

*Rafflesiaceae not included in calculation

Supplemental Figure 1: Proportion of Plastid Housekeeping Genes with Predicted Mitochondrial Retargeting (Left), and Proportion of Mitochondrial OXPHOS proteins with Predicted Plastid Retargeting (Right). Proportion of enzymes with predicted retargeting (probability ≥ 0.5) based on the outputs of LOCALIZER and/or TargetP. Autotrophs and heterotrophs are plotted in green and purple, respectively. Reported *p*-values from *t*-tests or Mann-Whitney U tests depending on distribution of the data. Note that *R. cantleyi* and *S. himalayana* are not included in plastid retargeting analysis due to suspected loss of the plastome. It should be noted that proportions of plastid housekeeping genes in the Rafflesiaceae represent predictions for only two proteins; all other proteins are missing from these lineages, see Fig. S3.

Supplemental Figure 2: Significant overlap between loss of organellar aaRS and retargeting of cytosolic counterpart in heterotrophs. Reported counts refer to number of aaRS "families" defined by amino acids (e.g., GluRS, PheRS, TrpRS, etc.). Organellar gene presence/absence determined as for Fig. 1. Partial sequences for organellar enzymes are counted as present. Predicted retargeting (probability ≥ 0.5) to either mitochondria or plastids was based on the outputs of LOCALIZER and/or TargetP. Note that aaRS where cytosolic targeting could not be tested (i.e., enzyme was not present, or N-terminus was incomplete) were excluded. Organellar enzyme present = shades of gray. Organellar enzyme absent = shades of orange. Cytosolic enzyme retargeting = darker shading. No retargeting = lighter shading. Reported *p*-value is from a Fisher's exact test for correlation between loss of organellar aaRS and retargeting of the corresponding cytosolic aaRS. Note that *R. cantleyi* and *S. himalayana* proteins are excluded from predictions of targeting to plastids as these species lack plastid genomes. Loss of GatCAB and associated cytosolic GlnRS retargeting are included in the counts.

Supplemental Figure 3: Targeting predictions for cytosolic-type aaRS. Deeper shades of green (plastid, right) and orange (mitochondrion, left) indicate probability of retargeting to that organelle is ≥ 0.5 based on both LOCALIZER and TargetP. Lighter shades of orange/green indicate one of the two algorithms yielded a probability ≥ 0.5. White indicates probability for both algorithms was < 0.5. Gray indicates that the gene/transcript was not present or that the N-terminus was incomplete, so no targeting prediction could be made. Heterotrophs are written in purple, closely related autotrophs in black. *Arabidopsis thaliana* is included as a reference.

| | Plastid Marker:
Chl Autoflouresence | Mito Marker:
FP611 | TP Marker:
GFP | Overlay |

FP611 only
in silico:
NA

AspRS TP
in silico:
mito

Supplemental Figure 4: Cytosolic-type aaRS transit peptides from heterotroph *S. himalayana* transiently expressed in *Nicotiana benthamiana*. Transit peptides (TPs) from *S. himalayana* fused to GFP (middle right column) were transiently expressed in *N. benthamiana.* Chlorophyll autofluorescence (far left column) was used for colocalization with plastids. As a positive control for mitochondrial targeting, eqFP611 was tagged with the TP from mitochondrial isovaleryl-CoA dehydrogenase (IVD) (middle left). Overlay of all three channels on far right. Top row: IVD_FP611 infiltration only- negative control. White ovals indicate plastids in untransformed cells, allowing assessment of bleed-over chlorophyll autofluorescence in the GFP channel. See main text for how true plastid localization was distinguished from bleed-over signal. Bottom row: Expression of AspRS TP led to apparent aggregates of GFP in the cytosol, potentially resulting from an artefact due to overexpression. Scale indicated in top right panel is the same for all images.

Supplemental Figure 5: Initial round of microscopy data using single expression vector PK7FWG2.
Transit peptides (TPs) from *S. himalayana* fused to GFP (middle right column) were transiently expressed in *N. benthamiana.* Chlorophyll autofluorescence (far left column) was used for colocalization with plastids. As a positive control for mitochondrial targeting, eqFP611 was tagged with the TP from mitochondrial isovaleryl-CoA dehydrogenase (middle left). The overlay of all three channels on far right. Arrows indicate

examples of plastid (yellow), mitochondrial (white) and potential nuclear (red) localization. Scale indicated in top right panel is the same for all images.

Supplemental Figure 6: Loss of key subunits for plastid import apparatus in *Rafflesiaceae*. For complete names of proteins and corresponding *A. thaliana* gene ID, see supplemental dataset 2. Hue of boxes indicates metabolism (gray = autotroph, orange = hemiparasite, purple = heterotroph), shade indicates degree of presence/absence (white = no sequence found, light colors = partial sequence found, dark colors = complete or near complete sequence found).

Supplemental Figure 7: Relative rates of sequence evolution of cytosolic and organellar aaRS are similar in heterotroph compared to autotrophic relative. Each point represents an aaRS branch length of *M. hypopitys* (heterotrophic) normalized to branch length of *R. williamsianum* (autotrophic) for canonically cytosolic and organellar aaRSs (log$_{10}$ scale).

Supplemental Figure 8: Maximum likelihood tree for CysRS. Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for organellar and cytosolic CysRS. Orthologs for *A. thaliana* organellar CysRS (AT2G31170) outlined in green, orthologs for *A. thaliana* cytosolic CysRS (AT3G56300, AT5G38830) outlined in red.

0.50

Supplemental Figure 9: Maximum likelihood tree for Protein Only RNase Ps (PRORPs). Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for nuclear and organellar PRORPs. Orthologs for *A. thaliana* organellar PRORP1 (AT2G32230) outlined in green, Orthologs for *A. thaliana* cytosolic PRORP2/3 (AT2G16650, AT4G21900) outlined in red.

Supplemental Figure 10: Maximum likelihood tree for tRNase Z 1/2. Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for tRNAse Z type proteins. Orthologs for *A. thaliana* cytosolic tRNase Z1 (AT1G74700) are outlined in red, and orthologs for *A. thaliana* plastid tRNase Z2 (AT2G04530) are outlined in green.