Photosynthesis drives retention of bacterial-like tRNA metabolism in plant organelles

Rachael Ann DeTar[1,]*, Joanna Chustecki[2], Anna Martinez-Hottovy[2], Luis Federico Cerrioti[3,4], Amanda K. Broz[1], M. Virginia Sanchez-Puerta[3,4], Christian Elowsky[5], Alan C. Christensen[2], Daniel B. Sloan[1]

[1] Department of Biology, Colorado State University, Fort Collins, CO, USA
[2] School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA
[3] IBAM, Universidad Nacional de Cuyo, CONICET, Facultad de Ciencias Agrarias, Almirante Brown 500, M5528AHB Chacras de Coria, Argentina
[4] Facultad de Ciencias Exactas y Naturales, Padre Jorge Contreras 1300, Universidad Nacional de Cuyo, M5502JMA Mendoza, Argentina
[5] Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, NE, USA
* Corresponding Author

Keywords: organelle gene expression, plastome, mitogenome, heme biosynthesis, tRNA, photosynthesis, aminoacyl-tRNA synthetase

Abstract

Eukaryotic nuclear genomes often encode distinct sets of protein translation machinery for function in the cytosol vs. organelles (mitochondria and plastids). This phenomenon raises questions about why multiple translation systems are maintained even though they are capable of comparable functions, and whether they evolve differently depending on the compartment where they operate. These questions are particularly interesting in land plants because translation machinery, including aminoacyl-tRNA synthetases (aaRS), is often dual-targeted to both the plastids and mitochondria. These two organelles have quite different metabolisms, with much higher rates of translation in plastids to supply the abundant, rapid-turnover proteins required for photosynthesis. Previous studies have indicated that plant organellar aaRS evolve more slowly compared to mitochondrial aaRS in other eukaryotes that lack plastids. Thus, we investigated the evolution of nuclear-encoded organellar and cytosolic translation machinery across a broad sampling of angiosperms, including non-photosynthetic (heterotrophic) plant species with reduced rates of plastid gene expression to test the hypothesis that translational demands associated with photosynthesis constrain the evolution of bacterial-like enzymes involved in organellar tRNA metabolism. Remarkably, heterotrophic plants exhibited wholesale loss of many organelle-targeted aaRS and other enzymes, even though translation still occurs in their mitochondria and plastids. These losses were often accompanied by apparent retargeting of cytosolic enzymes and tRNAs to the organelles, sometimes preserving aaRS-tRNA charging relationships but other times creating surprising mismatches between cytosolic aaRS and mitochondrial tRNA substrates. Our findings indicate that the presence of a photosynthetic plastid drives the retention of specialized systems for organellar tRNA metabolism.

Significance

The process by which endosymbionts are integrated into a host and become organelles results in a combination of gene loss, transfer to the nucleus, and retention in the organellar genome. It is not well understood why some endosymbiont-derived genes may be retained when a functional host counterpart exists whose gene product could simply be retargeted to the organelles. This study revealed that the photosynthetic activity in plant plastids may be responsible for retention of functionally redundant tRNA processing machinery, while mitochondria are more flexible regarding substitution with cytosolic type enzymes. Therefore, functional constraint in the plastid is likely more important than in the mitochondria for shaping the evolution and retention of translation machinery that is dual-targeted to both organelles.

Introduction

The evolutionary transition from a free-living bacterium to an endosymbiont and eventually to an

organelle requires the physical, biochemical, and genomic integration of two organisms. One common

consequence of this process is a reduction in the genome of the endosymbiont, driven by relocation of

essential genes to the nucleus via horizontal transfer or outright loss of some genes that are no longer

useful after transition to an intracellular lifestyle. This trajectory has been observed repeatedly for

various endosymbionts and organelles, including mitochondria, plastids, and more recently acquired

intracellular bacteria such as the cyanobacterial-derived chromatophores of *Paulinella* amoeba or the

many heritable endosymbionts in insects (1–6). However, there are some endosymbiotically derived

genes that are recalcitrant to loss despite the presence of host counterparts with comparable functions.

Redundant genes may be maintained in the organelle genomes or in the nuclear genome alongside

genes encoding proteins for the same role in other cellular compartments. These observations elicit

some interesting questions; 1) Why are some functionally redundant genes retained? 2) Do these

redundant genes evolve differently depending on the cellular compartment where they operate?

The genes that encode the machinery required for organelle translation are widely retained, as

eukaryotes maintain distinct translation systems for nuclear and mitochondrial transcripts. Translation

becomes even more complicated in eukaryotes bearing three genomes. For example, plant cells contain

nuclear, mitochondrial and plastid genomes. The translation machinery associated with each of these

cellular compartments includes ribosomal RNAs (rRNAs), ribosomal proteins, transfer RNAs (tRNAs), and

the accompanying enzymes involved in tRNA processing and aminoacylation. The aminoacyl-tRNA

synthetases (aaRS) that catalyze the "charging" of a tRNA with its conjugate amino acid tend to be highly

conserved, with distinct versions for function in the organelles and cytosol (7). The degree of

redundancy of tRNAs and their interactors between cellular compartments also varies significantly

within eukaryotes. These characteristics make tRNAs and their interactors, particularly aaRS, an ideal

case study for understanding cytonuclear gene redundancy.

The concerted activity of the aaRS along with other enzymes involved in post-transcriptional

modification such as tRNases P and Z (end processing in tRNA maturation) and CCAases (addition of CCA

tail) ensures the tRNA has the proper structure to fulfill its role in translation. Generally, organisms

would need a complement of at least 20 aaRS, one for each amino acid. However, eukaryotes often have

more than this minimum requirement, with some overlapping sets of enzymes targeted to different

compartments (8, 9). It is not fully understood why some eukaryotes retain distinct, redundant sets of

tRNA processing machinery, while others do not. However, retention of organelle-targeted aaRS

correlates with the retention of tRNA genes within organelle genomes. For example, *Homo sapiens*

retains 38 aaRS with nearly distinct sets for the cytosol and mitochondria (9–11). Correspondingly, all

necessary tRNA genes are present in the mitochondrial genome (mitogenome) (8). In contrast,

trypanosomatids do not encode any tRNAs in their mitogenomes but rather import nuclear-encoded

tRNAs into the mitochondria and have a reduced set of aaRS which colocalize to the cytosol and the

mitochondria (9, 12, 13). In plants, closely related species may have very different numbers of tRNAs

encoded in their mitogenomes, with import of cytosolic tRNAs to compensate for losses (8, 14, 15).

Notably, plants also dual-target and import an organellar set of enzymes into both plastids and

mitochondria (9, 16). The co-occurrence of redundant tRNAs and aaRS could be explained by the

observation that organelle-encoded tRNAs are often structurally divergent from cytosolic tRNAs and

require specific enzymatic partners for recognition (8, 17).

For most eukaryotes, organellar and cytosolic aaRS are exclusively nuclear-encoded. The shared

genomic location of these genes facilitates direct comparison of sequence conservation because any

differences in evolutionary rate should be more likely to reflect differences in selection, rather than

other factors like compartment-specific mutation rates or effective population size (18–21). Previous

studies have quantified rates of evolution for these enzymes in various animal lineages and a small

number of plants. Mitochondria-specific aaRS in animals exhibit lower sequence conservation than their

cytosolic counterparts (22–24). In contrast, plant organellar aaRS exhibit equivalent or greater sequence

conservation compared to their cytosolic counterparts (25). Furthermore, plastid genomes (i.e.,

plastomes) tend to retain a set of 30 tRNA genes that is fully sufficient for translation (26, 27). Plastomes

also generally have high rates of transcription (28) and translation (29) due to the abundance and rapid

turnover of proteins required for photosynthesis. Thus, the disparity in evolutionary rates between

animal mitochondrial aaRS and plant dual-organelle targeted aaRS may be due to the specific

translational needs of the plastid. We hypothesize that aaRS and other translation-associated enzymes

operating in the plastid face great functional constraints, thus promoting unusually high sequence

conservation of organellar translation machinery compared to other eukaryotes. A prediction stemming

from this hypothesis is that plant lineages with reduced or nonexistent plastid gene expression would

exhibit more rapid evolution of organellar aaRS due to relaxed selection.

In this work, we test whether photosynthetic activity contributes to sequence conservation of

organellar translation machinery by analyzing evolution of this machinery across diverse lineages of

autotrophic and heterotrophic plants. It should be noted there is a gradient of heterotrophy amongst

plants – for the purposes of this study, we use "heterotrophic" to describe plants that do not

photosynthesize at all and exclusively rely on parasitism for acquisition of carbon. They do this by

tapping directly into host plants or by exploiting mycorrhizal fungi (mycoheterotrophy) (30, 31). We

refer to facultative autotrophs that supplement photosynthetic carbon with parasitism as

"hemiparasitic." The hallmark of many heterotrophic/hemiparasitic plants is a reduced or even non-

existent plastome (32). Highly expressed genes related to photosynthesis such as the Rubisco large

subunit and the Photosystem II D1 reaction center tend to be absent from plastomes in heterotrophs,

thereby drastically decreasing demands on plastid gene expression. In addition, many tRNA genes have

been lost from the plastomes of these plant species (32), providing a good system to examine the

relationship between evolutionary redundancy of aaRS, tRNA content of the organelle genomes, and

trophic lifestyle.

Results

Heterotrophic plants exhibit extensive loss of organellar aaRS and tRNA processing enzymes

We searched for organellar and cytosolic aaRS orthologs in a broad sampling of angiosperm species with

a range of trophic lifestyles and plastome reduction (**Supp. Dataset 1**). For each

hemiparasitic/heterotrophic species in the analysis, we included the closest fully autotrophic relative

with a nuclear genome or transcriptome available, excepting the Balanophoraceae where we used

hemiparasite *Santalum album* as the reference autotroph. In total, we sampled 7 heterotrophs, 8

hemiparasites, and 10 autotrophs, including reference species *Arabidopsis thaliana.* Among

heterotrophs, we included the endoparasites *Rafflesia cantleyi* and *Sapria himalayana* from the family

Rafflesiaceae, likely the most extreme examples of plant heterotrophy, as several studies have

suggested wholesale loss of the plastome in these species (33–36).

We found orthologs across these diverse taxa by running Orthofinder (37, 38) on protein models

derived from nuclear genome or transcriptome data to find groups of likely orthologous proteins based

on similarity and phylogeny, i.e. "orthogroups". Then we extracted sequences that fall into an

orthogroup with known *A. thaliana* aaRS (**Supplemental Dataset 1**). Unexpectedly, many lineages of

heterotrophs did not have any orthologs for many of the organellar aaRS. This loss of organelle-specific

orthologs in heterotrophs was surprising as even species entirely lacking the plastome must still conduct

tRNA aminoacylation within mitochondria. Further investigation using reciprocal-best-hit (RBH) searches

with BLAST between *A. thaliana* and each species supported that these were true losses. These losses

primarily occurred in purely heterotrophic plants, while hemiparasites retained most of their organellar

aaRS. (**Fig. 1**). Yet, within the heterotrophs, there was a gradient of loss. The two endophytic species

from Rafflesiaceae (*R. cantleyi* and *S. himalayana*) exhibited the most striking loss of organellar aaRS,

retaining only PheRS. In contrast, obligate mycoheterotroph *Monotropa hypopitys* retained more aaRS

than any other heterotroph, missing only AlaRS, CysRS, ThrRS, and TrpRS (**Fig. 1**).

This striking pattern led us to investigate the retention of other enzymes involved in tRNA

maturation and processing; see (15) and **Supplemental Dataset 2** for a comprehensive list.

Unsurprisingly, the CCAase enzyme responsible for adding CCA tails to tRNAs in all compartments was

retained across all species. Two enzymes that maintain distinct cytosolic and organellar versions, tRNase

Z (involved in 3' end processing) and Protein Only RNase P (PRORP; required for 5' end processing),

exhibited two different trends. We found that plastid tRNase Z (39) was missing from all heterotrophs

except *M. hypopitys* and hemiparasites of the *Cuscuta* genus yet was retained in the autotrophic

relatives. Conversely, the organellar PRORP1 (40, 41) appeared to be universally retained while cytosolic

PRORP2/3 orthologs were missing from the Rafflesiaceae. We also investigated the retention of

bacterial-like organelle-specific machinery with no cytosolic counterpart, including the methionyl-tRNA

formyltransferase (MTF), tRNA-Ile lysidine synthetase (TilS) and the bacterial-type glutamine

amidotransferase (GatCAB) complex. We found orthologs for the putative *A. thaliana* MTF in all species

except the heterotrophs *Epigogium roseum* and *M. hypopitys,* and putative TilS orthologs were absent

only from the heterotrophs Rafflesiaceae and *Balanophora fungosa*. GatCAB is the key complex used by

organelles in land plants to accurately charge tRNA^{Gln} in the absence of a dedicated organellar GlnRS

(42–44). This involves a two-step process, whereby tRNA^{Gln} is initially mis-acylated with Glu by GluRS.

GatCAB then catalyzes the transamidation of Glu to Gln. Most of our test species, even heterotrophs,

retained genes for at least some of the GatCAB subunits. However, Rafflesiaceae species appear to have

lost all subunits in this complex.

Generally, we observed reductive evolution in heterotrophs via loss of organellar aaRS and

bacterial-type enzymes for tRNA modification. This begs the question as to how gene expression is

accomplished in the organelles of these species, particularly within mitochondria. One possibility is that

cytosolic enzymes are retargeted to the organelles to functionally replace organellar enzymes.

Heterotrophic plants may compensate for loss of organellar aaRS by retargeting cytosolic

counterparts

To test for cytosolic aaRS retargeting, we applied two *in silico* targeting prediction programs to protein

models for all cytosolic aaRS in our heterotrophs and their close autotrophic relatives. Our analysis

revealed that canonically cytosolic aaRS are disproportionately retargeted to mitochondria in obligate

heterotrophs compared to autotrophs (**Fig. 2,** p = 0.0341). There was also a non-significant trend

towards retargeting to the plastid (p = 0.190). Targeting of mitochondrial OXPHOS components and cell

wall synthesis enzymes were included as positive and negative controls, respectively. No plastid positive

control is provided as the reduced genome and altered metabolism of plastids in heterotrophic plants

precludes straightforward selection of a control gene set. For both heterotrophs and autotrophs, an

average of about 60% of OXPHOS proteins were predicted to be targeted to the mitochondria, and

about 10% to the plastid. About 10% of cell wall synthesis enzymes were predicted to be targeted to the

organelles for both groups (**Fig. 2**). Thus, our analysis appears to be detecting a legitimate biological

trend toward retargeting of cytosolic aaRS in heterotrophs, albeit with a generally high false negative

rate for mitochondrial retargeting. Furthermore, within the heterotrophic species there was a significant

correlation between loss of an organellar aaRS and retargeting of the corresponding cytosolic enzyme to

one or both compartments (p = 0.005, **Fig. S1**).

139   The variable quality of gene models for each species and high false negative rate for retargeting

prediction confound interpretation of retargeting for individual enzymes for each species (**Fig. S2**).

However, when looking only at *S. himalayana*, a species with relatively high-quality genome-based

models, we observed some interesting phenomena. Cytosolic GlnRS, MetRS, ProRS, and TrpRS were

predicted to be retargeted to mitochondria. Retargeting of cytosolic GlnRS could explains how this

species copes with loss of the organellar GatCAB complex (**Fig. 1**). The cytosolic AspRS and GlyRS were

predicted to be dual-targeted to both organelles, while AsnRS and TyrRS were predicted to be targeted

to the plastid despite the likely non-existence of a plastid genome. All other cytosolic enzymes were

either not predicted to be retargeted or were missing a large portion of the N-terminus.

To assess whether our *in silico* predictions hold true *in vivo,* we conducted targeting assays on a

subset of aaRS (see **Supplemental Dataset 3** for sequence information) by inducing transient expression

of N-terminal fusions of *S. himalayana* cytosolic aaRS transit peptides (TPs) to green fluorescent protein

(GFP) in *Nicotiana benthamiana* (**Fig. 3, Fig. S3**). We used GFP linked to the mitochondrial cytochrome C

TP from *S. himalayana* as an additional mitochondrial-targeted positive control (**Fig. S3**). We found that

in silico predictions were generally reflected *in vivo,* but some putative TPs targeted GFP to a different

compartment *in vivo*. For example, the AsnRS TP was predicted to be plastid-targeted *in silico*, yet

showed mitochondrial targeting during transient expression assays. Conversely, ProRS and TrpRS were

predicted to be mitochondrial *in silico*, yet exhibited plastid targeting *in vivo.* Additionally, we gained

further evidence that our *in silico* analysis produced false negatives for organellar targeting of some

aaRS TPs, as the N-terminal extension of ThrRS was not predicted to be an organellar transit peptide *in*

silico, yet retargeted GFP to the plastids in *N. benthamiana.* The only TP we cloned expecting organellar

retargeting which did not exhibit targeting to either organelle *in vivo* was AspRS (**Fig. S3**). GFP tagged

with this TP did not colocalize with mitochondrial FP611 florescence or plastid autofluorescence, but

rather formed punctate structures in the cytosol which are likely artefactual aggregates resulting from

overexpression. Generally, *in vivo* assays provided evidence for retargeting of cytosolic aaRS to at least

one organelle and suggested that *in silico* prediction tools may even be overly conservative.

Generally, organellar tRNAs differ from their cytosolic counterparts in evolutionary history,

sequence, and posttranscriptional modification (8). Identity elements which govern the specificity of

interaction between a tRNA and its conjugate aaRS may also differ between cytosolic and organellar

systems (45). However, in many systems (including plants), cytosolic tRNAs can be imported into the

organelles (8, 9, 14). Thus, our observation of retargeting opens some interesting questions – are the

retargeted aaRS charging organellar tRNAs? Or is there replacement of the entire organellar system of

tRNA aminoacylation, including both tRNAs and aaRS? We investigated the tRNA gene content of the

organelles of these species to get insight into which tRNA substrates may be in use.

Organelle-encoded tRNA retention often, but not always, mirrors retention of organelle-type aaRS

To investigate the possibility of enzyme-substrate mismatch, we cataloged the presence/absence of

organellar tRNA-encoding genes (*trn*) for all sampled heterotrophic species (**Fig. 4, Supplemental**

Dataset 4)**.** Generally, there was strong correspondence between retention of organellar aaRS and

plastome/mitogenome tRNA genes. As the Rafflesiaceae species are thought to lack plastomes entirely,

they likely have no plastome-encoded tRNAs (**Fig. 4**). The Balanophoraceae species also have been

reported to exhibit loss of all plastid tRNA genes (46–49). The Rafflesiaceae and Balanophoraceae have

additionally lost most of their mitogenome-encoded tRNAs. Correspondingly, these two families also

have the most reduced sets of organellar aaRS. At the opposite extreme, *M. hypopitys* retains most of its

mitochondrial and plastid tRNA genes and corresponding organellar aaRS.

However, there are some incidences where a cytosolic enzyme looks to be charging an

organellar tRNA due to loss of the organellar aaRS (**Fig. 4**). For example, the few remaining mitogenome-

encoded tRNAs in the Rafflesiaceae and Balanophoraceae species (excepting *S. himalayana* tRNA[Phe]

encoded by *trnF*) are missing their organellar aaRS partner and are likely charged by a cytosolic enzyme.

For both compartments, there are numerous instances where there is no encoded tRNA despite

retention of the organellar aaRS (**Fig. 4**).

Testing for accelerations in organellar aaRS sequence evolution in heterotrophs

Although wholesale loss of many organellar aaRS in heterotrophs precludes sequence conservation

analysis, *M. hypopitys* retained the majority of aaRS sequences while still exhibiting an obligate

heterotrophic lifestyle. Thus, we quantified sequence conservation of organellar and cytosolic aaRS in

M. hypopitys and its autotrophic relative *Rhododendron williamsianum.* We built gene trees and ran

relative rate analysis to test for accelerated evolution on the *M. hypopitys* branch (**Supplemental**

Dataset 5). We also directly compared branch length ratios of *M. hypopitys* to *R. williamsianum* in

cytosolic and organellar aaRS (**Fig. S4, Supplemental Dataset 5).** In neither analysis was there a

statistically significant trend towards decreased sequence conservation in organellar enzymes of *M.*

hypopitys. However, the deep rooting of our tree on *A. thaliana* may result in poor resolution of the

sequence evolution between *M. hypopitys* and *R. williamsianum.* Additionally, the transition to

heterotrophy on the *M. hypopitys* branch may have occurred relatively recently compared to the split

from *R. williamsianum*, thus diluting any signals of accelerated evolution.

Loss of GluRS and trnE in Rafflesiaceae may be facilitated by relocation of tetrapyrrole pathway to the

cytosol

In addition to its role in translation, aminoacylated plastid tRNA^{Glu} (encoded by the *trnE* gene) is the first

precursor in the plastid tetrapyrrole pathway. This bifurcating pathway begins with a common "stem"

set of enzymes providing the 5-aminoolevulinic acid (ALA) precursor for both the heme and chlorophyll

biosynthesis branches (50, 51). While heterotrophic plants do not need chlorophyll for photosynthesis,

heme is still a necessary cofactor for many enzymes and protein complexes involved in redox chemistry.

Accordingly, *trnE* is one of the most-retained genes in the plastomes of heterotrophic plant species (32)

and may even drive the retention of plastomes in non-photosynthetic organisms (52, 53). Yet, the

heterotrophs from Rafflesiaceae and Balanophoraceae used in this study have lost plastid *trnE* (35, 46).

We found that the Rafflesiaceae also are lacking the organellar GluRS typically required to aminoacylate

plastid tRNA^{Glu} for use in the tetrapyrrole pathway. Although a previous study of *R. cantleyi* (54)

reported the presence of a GluRS gene, which could potentially catalyze the first step in the tetrapyrrole

pathway, they did not investigate if it was an organellar or cytosolic enzyme. Intriguingly, the cytosolic

GluRS is not predicted to be retargeted to the plastid in the Rafflesiaceae, which prompted us to

investigate the presence/absence of tetrapyrrole pathway enzymes in our study species.

For both analyses, genes are grouped based on their placement in the tetrapyrrole pathway –

the shared "common pathway" or the heme and chlorophyll branches. Unsurprisingly, we found that

the heterotrophs (with exception of *Epipogium*) were lacking most genes in the chlorophyll biosynthesis

branch of the tetrapyrrole pathway (**Fig. S5**). The Rafflesiaceae also exhibited loss of the heme

oxygenase (HO) and phytochromobilin synthase (HY2) enzymes from the tetrapyrrole pathway, which

are necessary to produce phytochromobilin for light sensing. This may be a response to their

endoparasitic lifestyle where they only emerge from the host to flower and therefore can dispense with

light signaling. All heterotrophs retained most of the enzymes in the common pathway and the

ferrachelatase (FC), which catalyzes the final step in the production of heme. Therefore, it does not

appear that the tetrapyrrole pathway has been lost from any of the species, including those that lack

plastid *trnE* and/or organellar GluRS. A previous study on *Lophophytum* suggests that Balanophoraceae

species lacking plastid *trnE* likely import the nuclear-encoded tRNA^{Glu} but otherwise utilize the typical

plastid heme biosynthesis pathway (46). However, we have found the Rafflesiaceae lack both plastid

trnE and GluRS, and do not have predicted retargeting of the cytosolic GluRS. Thus, we hypothesized

that the Rafflesiaceae may adopt a different strategy to continue heme production by retargeting the

entire pathway to the cytosol and using nuclear-encoded tRNA^{Glu} and cytosolic GluRS. We investigated

this hypothesis using *in silico* targeting predictions and found that common pathway enzymes are not

predicted to be plastid-localized in the Rafflesiaceae, in contrast to autotrophic relative *Manihot*

esculenta, which retains plastid targeting (**Fig. S6**). Furthermore, plastid targeting was still predicted for

most tetrapyrrole enzymes in the other heterotrophic species that retain organellar GluRS, including the

Balanophoraceae species. Thus, it is possible that *S. himalayana* and *R. cantleyi* produce heme without a

functional plastid gene expression apparatus by relocating some steps in the tetrapyrrole pathway to

the cytosol.

Discussion

Loss of photosynthesis relaxes constraints on both plastid and mitochondrial translation machinery

Our primary objective in this study was to test if the high translational demands of the photosynthetic

chloroplast create functional constraint on the evolution of shared translation systems in plastids and

mitochondria. We accomplished this by investigating relative rates of protein evolution in

cytosolic/organellar aaRS in heterotrophic plants compared to their photosynthetic relatives. To our

surprise, heterotrophic plants underwent widespread loss of organellar aaRS and many bacterial-like

enzymes involved in tRNA processing. These losses suggest an extreme relaxation of selection pressures

in the absence of a photosynthesizing plastid and support our hypothesis that the translational demands

in chloroplasts are the linchpin responsible for the unusually strong functional constraint on the

redundant organelle-specific translation machinery in plants.

There was variation in the degree of loss across the heterotrophs, often correlated with the

number of genes retained in the plastome, particularly tRNAs (see **Supp. Dataset 1** for

plastome/mitogenome content). For example, the most losses occurred in the Rafflesiaceae, all of which

are obligate endoparasites and have likely lost their plastomes (33–36, 55). The fewest losses occurred

in *M. hypopitys*, which has the highest number of plastome-encoded genes of all the heterotrophic

species we sampled – and notably the greatest number of retained plastid tRNAs (56). Thus, *M.*

hypopitys may not be as far along the path of reductive evolution as the other species, which might

provide an additional explanation as to why our comparative rate analyses on the remaining organellar

aaRS in *M. hypopitys* did not support accelerated evolution compared to cytosolic enzymes (**Fig. S4**). The

Balanophoraceae and *Epipogium* species exhibited intermediate degrees of aaRS and plastome gene

loss, although the Balanophoraceae have lost all plastid tRNAs (Schelkunov *et al.*, 2015, 2019; Chen *et*

al., 2019; Serna-Sánchez *et al.*, 2021; Yu *et al.*, 2022). An increased plastome mutation rate caused by

loss of DNA repair machinery (57, 58) in the Balanophoraceae may explain accelerated plastome (tRNA)

gene losses relative to nuclear (aaRS) gene loss. Generally, loss of organellar aaRS correlates well with

reduction of the plastome across heterotrophs.

Loss of organellar aaRS is associated with retargeting of cytosolic aaRS and potential rewiring of

aaRS-tRNA interactions

Wholesale loss of organellar aaRS in heterotrophs was unexpected, as most non-photosynthetic plants

still contain a minimal plastome to support heme and fatty acid biosynthesis (32, 53) and fully intact

mitogenomes (see **Supp. Dataset 1**). This conundrum prompted us to investigate if cytosolic aaRS are

retargeted to organelles in heterotrophs to compensate for gene loss, as has been shown for

aaRS/tRNAs in mitochondria of other plant species (59). Retargeting a protein to the organelles

generally involves obtaining an N-terminal TP sequence that can be recognized by organellar protein

import complexes (60–62). We discovered that many cytosolic aaRS in the heterotrophs had N-terminal

sequences predicted to target to one or both organelles (**Fig. 2-3, S1-S3**). Therefore, the import of

cytosolic aaRS likely compensates for the loss of organellar aaRS and even the loss of the GatCAB

complex in Rafflesiaceae. There was not perfect correspondence between loss of organellar aaRS and

predicted retargeting of cytosolic aaRS. However, the extent of retargeting may have been

underestimated for several reasons: 1) TP sequence composition varies considerably by plant lineage

(63), making *in silico* predictions more challenging in uncharacterized species, 2) the low quality of

protein models for some of our species may have led to true N-terminal sequences being absent or

incomplete, and 3) some organelle-imported cytosolic aaRS may eschew the use of typical TPs that are

key for targeting prediction *in silico* (64). Retargeting of cytosolic aaRS may occur through two different

mechanisms; 1) alternative splicing or alternative transcription/translation start sites within one gene

resulting in cytosolic and organellar isoforms and 2) duplicated genes within the genome encoding

cytosolic and organellar paralogs. Our ability to investigate these two mechanisms was limited as we

generally only had transcriptome or genome information but not both for each species. However,

observations of transcriptomes assembled with consideration for alternative splice isoforms indicate

that both mechanisms are potentially at play (**Supplemental Dataset 6**). For example, *R. phalloides* has

two separate paralogs for cytosolic ThrRS, one of which is predicted to be mitochondrial-targeted

(Rphallo_DN1669_c0_g1) while the other is not (Rphallo_DN1669_c0_g2). Conversely, *R. phalloides*

TyrRS is encoded by a single locus (Rphallo_DN204_c0_g1) with a multitude of splice isoforms, some of

which are putatively mitochondrial-targeted. Genome sequencing and long-read mRNA sequencing

would allow for more thorough analysis of which mechanism is most common in each species.

Retargeting of cytosolic aaRS led us to investigate organelle genome tRNA content and whether

there was substrate-enzyme mismatch in the organelles, i.e. instances where the cytosolic enzyme was

interacting with an organellar tRNA or vice-versa. Our interpretations presume that loss of organellar

tRNA genes is compensated by import of the corresponding cytosolic tRNAs (9, 14, 65). We found that

the retention of organellar aaRS in heterotrophs generally matched the retention of plastid tRNAs but

not necessarily mitochondrial tRNAs (**Fig. 4**). This is further supported by observations in hemiparasite

Viscum album, which has lost most of its mitochondrial tRNAs, yet retains most organellar aaRS in

conjunction with a nearly full set of plastid tRNAs (32, 66, 67). Substrate-enzyme mismatch was rare

within the plastid and usually involved loss of a plastid tRNA yet continued presence of the

corresponding organellar aaRS (e.g., see *R. phalloides* where all plastid tRNAs are lost but a handful of

organellar aaRS remain). There were also instances of organellar aaRS apparently charging imported

cytosolic tRNAs in the mitochondria (e.g., *trnF* in *R. cantleyi and R. phalloides*). More surprisingly,

however, there were cases where a mitochondrial tRNA was retained despite loss of the corresponding

organellar aaRS and apparent retargeting of the cytosolic enzyme, particularly in the Rafflesiaceae. This

type of mismatch is unexpected because cytosolic aaRS are thought to generally be more discriminating

enzymes than their mitochondrial counterparts (8, 59, 68–70), making it more difficult to transition to a

new tRNA substrate. Our results suggest some cytosolic enzymes may be interacting with bacterial-type

tRNAs in these heterotrophic species, although we cannot rule out the possibility that cytosolic tRNAs

are being redundantly imported. Given that parasitic plant lineages tend to be fast evolving across all

three genomes (71), the cytosolic aaRS in these species may have evolved lower specificity and the

capability to charge organellar tRNAs. Overall, our findings suggest that the plastid tRNA content, and

not the mitochondrial tRNA content, drives retention of organellar aaRS, again placing the

photosynthetic plastid as a central factor in the retention of functionally redundant translation

machinery.

320   The identity of retained organellar aaRS versus those substituted by cytosolic counterparts was

seemingly non-random. GluRS and PheRS were the two mostly highly retained organellar aaRS across

these diverse lineages of plants (**Fig. 1**). As tRNA^{Glu} is the most frequently retained tRNA in heterotrophic

plant plastomes due to its additional role in heme biosynthesis (50, 53), it is not surprising that GluRS

was among the most retained organellar aaRS in this study. The Rafflesiaceae were the only exception,

as they might have evolved retargeting of the entire heme pathway to the cytosol to compensate for

lack of plastid *trnE*. Organellar PheRS is likely retained because the cytosolic PheRS is unusual in that it is

a heteromeric complex with independently encoded α and β subunits (72). Therefore, functional

replacement of the organellar PheRS would require two independent acquisitions of TPs, as well as

stoichiometric import and assembly within the organelles. Such a coordinated set of changes may be

unlikely to evolve (24, 59). The one exception was *E. aphyllum,* for which we could not find an organellar

PheRS. However, this may be an artefact of low expression, as protein models for this species were

curated from RNA-seq data. At the other extreme, there was repeated loss of organellar AlaRS, which

was absent from all heterotrophs and the hemiparasitic *Cuscuta* species, possibly because plastomes of

heterotrophs are often AT-rich and thus have underrepresentation of amino acids with GC-rich codons

like Ala, Gly and Pro (47, 48). Thus, the corresponding organellar aaRS and tRNAs may be subject to

decreased functional constraint and be less likely to be retained. This hypothesis is supported by the fact

that we also see frequent loss of organellar GlyRS and ProRS in heterotrophs (**Fig. 1**).

Our results indicate that imported cytosolic aaRS and tRNAs are sufficient to sustain

mitochondrial translation in plants, supporting the hypothesis that the selective pressure to keep

organellar aaRS/tRNAs is maintained by translational demands associated with photosynthesis. Yet

other eukaryote lineages that never bore plastids often still retain organelle-specific tRNAs and aaRS for

their mitochondria (8, 9, 22, 24). This is even more puzzling when considering that various eukaryotic

lineages have jettisoned nearly all their mitochondrial tRNAs/aaRS, including cnidarians, ctenophores,

some fungi, and trypanosomes (12, 24, 73). One possible explanation is the logical parallel to our finding

that photosynthesis governs plastid translation rates and maintains gene redundancy; eukaryotic

lineages with high respiratory rates may require high mitochondrial gene expression and, therefore,

retain mitochondria-specific translation machinery. Indeed, mitochondrial gene expression tracks well

with respiratory rate and cellular activity across different tissues and metabolic states (74–76). This

connection to cell energetics would also explain why mitochondrial tRNAs/aaRS are disposable in

parasitic eukaryotes which presumably have lower energy demands, like trypanosomes and

heterotrophic plants. By studying non-model plant species with unusual physiology and life histories, we

were able to link plastid bioenergetic processes to the evolution of both organellar and nuclear

genomes. The sequencing of mitogenomes and characterization of metabolism in non-model eukaryotes

will provide exciting opportunities to conduct similar studies to better understand how mitochondrial

metabolism effects gene redundancy and elucidate the principles governing endosymbiotic integration.

Methods

Data acquisition and ortholog search

Gene models for species used in the study were collected from publicly available data sources

(**Supplemental Dataset 1**). When protein sequences were not available, we generated sequences from

transcriptome data using Transdecoder v5.5.0 (77), applying a minimum protein length of 50 amino

acids and retaining only protein sequences with blast hits to any sequence in the Uniprot Taxonomic

Divisions Plant database (https://ftp.uniprot.org/pub/databases/uniprot/knowledgebase

/taxonomic_divisions/). Then, the Transdecoder primary_transcript.fa script was used to extract only

the longest translated protein sequence from the primary transcript for each gene to reduce the search

space. SeqKit 2.3.0 (78) was used to create uniform gene handles for all files and remove stop codons

from sequences. Orthofinder v2.5.4 (37) was run on all sets of protein models with a user-defined

species tree (see **Fig. 1** for tree topology) to find groups of likely orthologous proteins, i.e. "orthogroups"

in two steps as follows:

orthofinder.py -f <folder_with_protein_models> -t 24 -a 6

orthofinder.py -ft <folder_with_results_> -s userdefined_speciestree.txt -t 24 -a 6

Orthogroups containing proteins of interest including cytosolic/organellar aaRS and tRNA-processing

enzymes characterized in *A. thaliana* (**Supplemental Dataset 2**) were extracted and checked for

completeness via RBH, using blastp from the command line BLAST 2.13.0+ toolkit (79) and custom

Python/Bash scripts. After ortholog discovery via Orthofinder, full sets of protein models (not just the

longest) were searched for *A. thaliana* protein sets using blastp to ensure narrowing of search space for

Orthofinder did not result in loss of matching gene models. In cases where available curated protein

models excluded alternative transcripts, this required obtaining transcriptome data and running

Transdecoder (see **Supplemental Dataset 1** for details). Note that ArgRS was not included in the

ortholog search as there are no distinct organellar versions of this enzyme in *A. thaliana.* One ArgRS

gene product is cytosol-targeted, and the other is triple-targeted to cytosol, mitochondria, and plastids

(9, 16). In the case of GlnRS, there is no organellar ortholog as plants typically use the bacterial-type

GatCAB amidotransferase system to charge Gln-tRNA indirectly in mitochondria and plastids (42). For

CysRS, tRNase Z and PRORP, sequences for both cytosolic and organellar proteins were clustered

together in one orthogroup likely to due to recent shared ancestry. For these genes, cytosolic versus

organellar versions were differentiated by aligning sequences in each orthogroup using MAFFT v7.490

(80) with the --auto setting, followed by manual trimming of the alignment in MEGA 11 (Tamura et al.,

2021). Maximum-likelihood tree building was conducted using RaXML v 8.2.12 (82) as follows:

raxmlHPC-PTHREADS-SSE3 -f a -# 100 -p 12345 -x 12345 -s <fasta file with orthogroup sequences> -n

<orthogroup> -m PROTGAMMALG -T 12

Inclusion in cytosolic versus organellar groups was then determined by relationship to known *A. thaliana*

cytosolic/nuclear/organellar proteins in maximum likelihood trees (**Figs. S7-10**).

Once we had a curated set of sequences for aaRS and processing enzymes for all species,

orthologs were then checked for completeness of length by alignment with *A. thaliana* orthologs by

MAFFT and subsequent plotting with a custom Python script. A "full-length" ortholog was defined as

having at least 50% alignment with any *A. thaliana* protein in the orthogroup, excluding pseudogenes.

Custom Python and Bash scripts are available at https://doi.org/10.5061/dryad.0cfxpnw7p.

In silico targeting predictions

For TP analysis, all protein models available for each gene with similarity to the original Orthofinder hit

(blastp e-value < 1e-6) were used, not just the longest. All transcripts for cytosolic aaRS, a "positive

control" set of genes involved in mitochondrial oxidative phosphorylation and a "negative control" set of

genes involved in cell-wall synthesis were run through targeting prediction programs LOCALIZER-1.0.5

(83) and TargetP-2.0 (84), with commands as follows:

LOCALIZER.py -p -i <input file>

targetp -fasta <input file> -mature -org pl

Protein models for some species were often incomplete due to the fragmented nature of some of the

transcriptome assemblies. To ensure models missing the N-terminus were not used in retargeting

analysis, predictions for models that did not align within 100 amino acids of the start codon for at least

one *A. thaliana* ortholog were discarded. For transcript-by-transcript heatmaps (**Supplemental Dataset**

6), a custom R script was used to plot all transcripts for all species. For plotting in meta-heatmaps (**Fig.**

S2), only the most retargeted ortholog for each species was used. Orthologs were then binned based on

probability of targeting to the mitochondria or plastids. High confidence retargeting was defined as a >

0.25 probability of targeting for both LOCALIZER and TargetP-2.0. Moderate confidence was defined as a

> 0.25 probability for one of the algorithms. Low confidence retargeting was defined as neither

algorithm providing a probability > 0.25. For subsequent analysis of ratios (**Fig. 2, Fig. S1**), any ortholog

with a "moderate" or better confidence was counted as retargeted. Note that *R. cantleyi* and *S.*

himalayana proteins were excluded from predictions of targeting to plastids as these species likely lack

plastomes. All analyses including plotting and statistics were carried out using custom Bash, Python, and

R scripts available at https://doi.org/10.5061/dryad.6hdr7sr5x.

In-vivo localization of S. himalayana aaRS transit peptides via tobacco infiltration and confocal

microscopy

TPs were chosen based on having a moderate probability (see above) of targeting a protein to either the

mitochondria or plastid, and/or having an N-terminal extension preceding conserved sequence shared

with *A. thaliana.* We selected either the sequence predicted to be a TP by TargetP-2.0 plus 10 residues

or the entire N-terminal overhang plus the first 10 residues aligning with the functional portion of the *A.*

thaliana protein. These amino acid sequences were then converted into DNA sequence using EMBOSS

Backtranseq (85) with the *Nicotiana tabacum* codon usage table (see **Supplemental Dataset 3**). These

sequences were then used to generate constructs for *Agrobacterium* transformation of *N. benthamiana*

and confocal microscopy as described in (59). Constructs are available from Addgene (ID numbers

202646-202653).

Mitogenome assembly

Prior to tRNA analysis, the mitogenomes of *S. himalayana* (SRR629601) and *R. cantleyi* (SRR629613)

were assembled *de novo* from raw reads available from the NCBI SRA database*.* To maximize the

likelihood of finding all mitogenome tRNAs, genomes were assembled in several ways. Reads were first

trimmed using FASTP (86), with either default parameters or "harsh" trimming where the --

qualified_quality_phred and --unqualified_percent_limit parameters were set to 20 and 10, respectively.

SPAdes v.3.15.4 (87) was then used to assemble trimmed reads *de novo,* with kmer sizes of 21, 33, 55,

and 77*.* To avoid potential assembly fragmentation due to excessive levels of sequence coverage, a

subset of reads (1 million from S. *himalayana* and 500,000 from *R. cantleyi*) were sampled and

assembled from each species as well.

tRNA analysis

Excepting the Rafflesiaceae, for which we assembled mitogenomes *de novo,* the mitogenomes,

plastomes and accompanying GenBank (.gbk) annotation files for species of interest were downloaded

from NCBI (**Supplemental Dataset 1**). A custom Perl script was used to extract annotated tRNAs from

the .gbk files. Concurrently, tRNAscan-SE-2.0.9 (88, 89) was run on downloaded genomes (parameters -

BQH), and results were cross-checked with preexisting annotations from GenBank annotations

(**Supplemental Dataset 4**). For *S. himalayana* and *R. cantleyi*, the same tRNAscan-SE-2.0.9 pipeline was

used on all generated assemblies, and the consensus set of tRNAs was checked for plant origin using

blastn against all NCBI reference sequences and the *A. thaliana* mitogenome (NC_037304.1) and

plastome (NC_000932.1). tRNA presence/absence patterns were plotted along with the corresponding

aaRS presence/absence/retargeting data using a custom Python script. Scripts are available at

https://doi.org/10.5061/dryad.np5hqc009.

Rate Analysis

Primary protein sequences for cytosolic and organellar aaRS from heterotroph *M. hypopitys* and related

autotroph *R. williamsianum* from the original Orthofinder search were used for comparison of rates,

with *A. thaliana* orthologs forming the outgroup. CysRS, LeuRS, ThrRS, and TrpRS were excluded from

the analysis because the organellar ortholog was missing or too short to create informative gene trees in

one or both of the test species. Predicted TPs were trimmed from sequences using output of TargetP

2.0. Sequences were then aligned using MAFFT v7.490 (80) and trimmed manually in MEGA 11 (81).

Maximum likelihood trees were built using RaXML v 8.2.12 (82) as follows:

raxmlHPC-PTHREADS-SSE3 -f a -# 100 -p 12345 -x 12345 -s <orthogroup_aligned.fa> -n <orthogroup> -m

PROTGAMMALG -T 12

For orthogroups containing only three ortholog sequences, a "dummy" outgroup sequence represented

by the *A. thaliana* ortholog with one branch-specific substitution was created to overcome RaXML's

input requirement of four or more taxa. The resulting trees were then evaluated for topology. In

instances where species-specific paralogs were not sister to one another, separate multiple sequence

alignments and trees were re-made for each paralog as described above. In instances where species-

specific paralogs were sister, both sequences were retained in the same tree. Trees were rooted to the

A. thaliana ortholog(s), then branch lengths to the last shared node from each *H. monotropa* and *R.*

williamsianum ortholog were extracted from the trees. For instances where two sister paralogs from the

same species were retained, the branch lengths were averaged. Then, the $\log_{10}$-transformed ratio of the

H. monotropa/R williamsianum branches was plotted, and mean branch length for cytosolic and

organellar enzymes was calculated. Statistical difference in mean branch ratio in cytosolic vs. organellar

enzymes was tested using a *t*-test. Tajima's relative rate tests were run in MEGA 11 using *A. thaliana* as

the outgroup. Data are available in **Supplemental Dataset 5.** Scripts, alignments, and maximum

likelihood trees are available at https://doi.org/10.5061/dryad.tb2rbp05r.

Acknowledgements

We thank Jessica Warren for helpful discussion and feedback on this project. This work was supported

by grants MCB-2322154, MCB-2048407, and MCB-1933590 from the National Science Foundation (NSF)

and utilized the RMACC Summit supercomputer (NSF ACI-1532235 and ACI-1532236). RAD was

supported by an NSF Postdoctoral Research Fellowship in Biology (IOS-2208908).

References

1.   A. N. Khachane, K. N. Timmis, V. A. P. Martins Dos Santos, Dynamics of reductive genome
evolution in mitochondria and obligate intracellular microbes. *Mol Biol Evol* **24**, 449–456 (2007).

2.   P. J. Keeling, J. D. Palmer, Horizontal gene transfer in eukaryotic evolution. *Nat Rev Genet* **9**, 605–
618 (2008).

3.   W. Martin, *et al.*, Evolutionary analysis of Arabidopsis, cyanobacterial, and chloroplast genomes
reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. *PNAS* **99**,
12246–12251 (2002).

4.   E. C. M. Nowack, *et al.*, Endosymbiotic gene transfer and transcriptional regulation of transferred
genes in Paulinella chromatophora. *Mol Biol Evol* **28**, 407–422 (2011).

5.   N. A. Moran, J. P. McCutcheon, A. Nakabachi, Genomics and evolution of heritable bacterial
symbionts. *Annu Rev Genet* **42**, 165–190 (2008).

6.   J. P. McCutcheon, N. A. Moran, Extreme genome reduction in symbiotic bacteria. *Nat Rev*
Microbiol **10**, 13–26 (2012).

7.   V. Rajendran, P. Kalita, H. Shukla, A. Kumar, T. Tripathi, Aminoacyl-tRNA synthetases: Structure,
function, and drug discovery. *Int J Biol Macromol* **111**, 400–414 (2018).

8.   T. Salinas-Giegé, R. Giegé, P. Giegé, TRNA biology in mitochondria. *Int J Mol Sci* **16**, 4518–4559
(2015).

9.   A. M. Duchêne, C. Pujol, L. Maréchal-Drouard, Import of tRNAs and aminoacyl-tRNA synthetases
into mitochondria. *Curr Genet* **55**, 1–18 (2009).

10.   L. Bonnefond, *et al.*, Toward the full set of human mitochondrial aminoacyl-tRNA synthetases:
Characterization of AspRS and TyrRS. *Biochemistry* **44**, 4805–4816 (2005).

11.   E. Tolkunova, H. Park, J. Xia, M. P. King, E. Davidson, The human lysyl-tRNA synthetase gene
encodes both the cytoplasmic and mitochondrial enzymes by means of an unusual: Alternative
splicing of the primary transcript. *Journal of Biological Chemistry* **275**, 35063–35069 (2000).

12.   S. Shikha, R. Brogli, A. Schneider, N. Polacek, TRNA biology in trypanosomes. *Chimia (Aarau)* **73**,
395–405 (2019).

13.   T. H. P. Tan, R. Pach, A. Crausaz, A. Ivens, A. Schneider, tRNAs in Trypanosoma brucei : Genomic
Organization, Expression, and Mitochondrial Import . *Mol Cell Biol* **22**, 3707–3717 (2002).

14.   J. M. Warren, *et al.*, Rapid Shifts in Mitochondrial tRNA Import in a Plant Lineage with Extensive
Mitochondrial tRNA Gene Loss. *Mol Biol Evol* **38**, 5735–5751 (2021).

15.   J. M. Warren, D. B. Sloan, Interchangeable parts: The evolutionarily dynamic tRNA population in
plant mitochondria. *Mitochondrion* **52**, 144–156 (2020).

16.   A.-M. Duchêne, *et al.*, Dual targeting is the rule for organellar aminoacyl-tRNA synthetases in
Arabidopsis thaliana. *PNAS* **102**, 16484–16489 (2005).

17.   K. Fujishima, A. Kanai, tRNA gene diversity in the three domains of life. *Front Genet* **5** (2014).

18.   M. Lynch, B. Koskella, S. Schaack, Mutation Pressure and the Evolution of Organelle Genomic
Architecture. *Science (1979)* **311**, 1727–1730 (2006).

19.   S. Schaack, E. K. H. Ho, F. MacRae, Disentangling the intertwined roles of mutation, selection and
drift in the mitochondrial genome. *Philosophical Transactions of the Royal Society B: Biological
Sciences* **375** (2020).

20.   K. Crosby, D. R. Smith, Does the Mode of Plastid Inheritance Influence Plastid Genome
Architecture? *PLoS One* **7** (2012).

21.   K. H. Wolfe, W.-H. Li, P. M. Sharp, "Rates of nucleotide substitution vary greatly among plant
mitochondrial, chloroplast, and nuclear DNAs (plant molecular evolution/molecular
clock/mutation rate/organelle DNA/inverted repeat)" (1987).

22.   J. R. Adrion, P. S. White, K. L. Montooth, The roles of compensatory evolution and constraint in
aminoacyl trna synthetase evolution. *Mol Biol Evol* **33**, 152–161 (2016).

23.   F. S. Barreto, *et al.*, Genomic signatures of mitonuclear coevolution across populations of
Tigriopus californicus. *Nat Ecol Evol* **2**, 1250–1257 (2018).

24.   W. Pett, D. V Lavrov, Cytonuclear Interactions in the Evolution of Animal Mitochondrial tRNA
Metabolism. *Genome Biol Evol* **7**, 2089–2101 (2015).

25.   D. B. Sloan, R. A. DeTar, J. M. Warren, Aminoacyl-tRNA synthetase evolution within the dynamic
tripartite translation system of plant cells. *Genome Biol Evol* **15** (2023).

26.   S. Wicke, G. M. Schneeweiss, C. W. dePamphilis, K. F. Müller, D. Quandt, The evolution of the
plastid chromosome in land plants: Gene content, gene order, gene function. *Plant Mol Biol* **76**,
273–297 (2011).

27.   M. Michaud, V. Cognat, A. M. Duchêne, L. Maréchal-Drouard, A global picture of tRNA genes in
plant genomes. *Plant Journal* **66**, 80–93 (2011).

28.   E. S. Forsythe, *et al.*, Organellar transcripts dominate the cellular mRNA pool across plants of
varying ploidy levels. *PNAS* **119** (2022).

29.   J. Mergner, *et al.*, Mass-spectrometry-based draft of the Arabidopsis proteome. *Nature* **579**, 409–
414 (2020).

30.   J. Těšitel, Functional biology of parasitic plants: A review. *Plant Ecol Evol* **149**, 5–20 (2016).

31.   L. Teixeira-Costa, C. C. Davis, Life history, diversity, and distribution in parasitic flowering plants.
Plant Physiol **187**, 32–51 (2021).

32.   S. Wicke, J. Naumann, "Molecular Evolution of Plastid Genomes in Parasitic Flowering Plants" in
Advances in Botanical Research, (Academic Press Inc., 2018), pp. 315–347.

33.   D. R. Smith, S. R. Asmail, Next-generation sequencing data suggest that certain
nonphotosynthetic green plants have lost their plastid genomes. *New Phytologist* **204**, 7–11
(2014).

34.   H. T. Li, *et al.*, Plastid phylogenomic insights into relationships of all flowering plant families. *BMC
Biol* **19** (2021).

35.   J. Molina, *et al.*, Possible loss of the chloroplast genome in the parasitic flowering plant Rafflesia
lagascae (Rafflesiaceae). *Mol Biol Evol* **31**, 793–803 (2014).

36.   D. L. Nickrent, Y. Ouyang, R. J. Duff, C. W. Depamphilis, Do nonasterid holoparasitic flowering
plants have plastid genomes? *Plant Mol Biol* **34**, 717–729 (1997).

37.   D. M. Emms, S. Kelly, OrthoFinder: Phylogenetic orthology inference for comparative genomics.
Genome Biol **20** (2019).

38.   D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol* **16** (2015).

39.   G. Canino, *et al.*, Arabidopsis encodes four tRNase Z enzymes. *Plant Physiol* **150**, 1494–1502
(2009).

40.   B. Gutmann, A. Gobert, P. Giegé, PRORP proteins support RNase P activity in both organelles and
the nucleus in Arabidopsis. *Genes Dev* **26**, 1022–1027 (2012).

41.   A. Gobert, *et al.*, A single Arabidopsis organellar protein has RNase P activity. *Nat Struct Mol Biol*
17, 740–744 (2010).

42.   C. Pujol, *et al.*, Dual-targeted tRNA-dependent amidotransferase ensures both mitochondrial and
chloroplastic Gln-tRNAGln synthesis in plants. *Proc Natl Acad Sci U S A* **105**, 6481–6485 (2008).

43.   A. Nagao, T. Suzuki, T. Katoh, Y. Sakaguchi, T. Suzuki, Biogenesis of glutaminyl-mt tRNA Gln in
human mitochondria. *PNAS* **106**, 16209–16214 (2009).

44.   L. Echevarría, *et al.*, Glutamyl-tRNAGln amidotransferase is essential for mammalian
mitochondrial translation in vivo. *Biochemical Journal* **460**, 91–101 (2014).

45.   R. Giegé, G. Eriani, The tRNA identity landscape for aminoacylation and beyond. *Nucleic Acids Res*
51, 1528–1570 (2023).

46. L. F. Ceriotti, M. E. Roulet, M. V. Sanchez-Puerta, Plastomes in the holoparasitic family
Balanophoraceae: Extremely high AT content, severe gene content reduction, and two
independent genetic code changes. *Mol Phylogenet Evol* **162** (2021).

47. H. J. Su, *et al.*, Novel genetic code and record-setting AT-richness in the highly reduced plastid
genome of the holoparasitic plant Balanophora. *Proc Natl Acad Sci U S A* **116**, 934–943 (2019).

48. M. I. Schelkunov, M. S. Nuraliev, M. D. Logacheva, Rhopalocnemis phalloides has one of the most
reduced and mutated plastid genomes known. *PeerJ* **2019** (2019).

49. X. Chen, *et al.*, Comparative Plastome Analysis of Root-and Stem-Feeding Parasites of Santalales
Untangle the Footprints of Feeding Mode and Lifestyle Transitions. *Genome Biol Evol* **12**, 3663–
3676 (2019).

50. R. Tanaka, K. Kobayashi, T. Masuda, Tetrapyrrole Metabolism in Arabidopsis thaliana. *Arabidopsis*
Book (2011) https:/doi.org/10.1199/tab.0145.

51. K. Kobayashi, T. Masuda, Transcriptional regulation of tetrapyrrole biosynthesis in arabidopsis
thaliana. *Front Plant Sci* **7** (2016).

52. M. Kayama, *et al.*, Highly Reduced Plastid Genomes of the Non-photosynthetic
Dictyochophyceans Pteridomonas spp. (Ochrophyta, SAR) Are Retained for tRNA-Glu-Based
Organellar Heme Biosynthesis. *Front Plant Sci* **11** (2020).

53. A. C. Barbrook, C. J. Howe, S. Purton, Why are plastid genomes retained in non-photosynthetic
organisms? *Trends Plant Sci* **11**, 101–108 (2006).

54. S. M. Ng, *et al.*, Comparative analysis of nucleus-encoded plastid-targeting proteins in Rafflesia
cantleyi against photosynthetic and non-photosynthetic representatives reveals orthologous
systems with potentially divergent functions. *Sci Rep* **8** (2018).

55. L. Cai, *et al.*, Deeply Altered Genome Architecture in the Endoparasitic Flowering Plant Sapria
himalayana Griff. (Rafflesiaceae). *Current Biology* **31**, 1002-1011.e9 (2021).

56. M. D. Logacheva, M. I. Schelkunov, V. Y. Shtratnikova, M. V. Matveeva, A. A. Penin, Comparative
analysis of plastid genomes of non-photosynthetic Ericaceae and their photosynthetic relatives.
Sci Rep **6** (2016).

57. M. I. Schelkunov, M. S. Nuraliev, M. D. Logacheva, Genomic comparison of non-photosynthetic
plants from the family Balanophoraceae with their photosynthetic relatives. *PeerJ* **9** (2021).

58. L. F. Ceriotti, L. Gatica-Soria, M. V. Sanchez-Puerta, Cytonuclear coevolution in a holoparasitic
plant with highly disparate organellar genomes. *Plant Mol Biol* **109**, 673–688 (2022).

59. J. M. Warren, *et al.*, Rewiring of aminoacyl-tRNA synthetase localization and interactions in plants
with extensive mitochondrial tRNA gene loss. *Mol Biol Evol* **40** (2023).

60. M. W. Murcha, *et al.*, Protein import into plant mitochondria: Signals, machinery, processing, and
regulation. *J Exp Bot* **65**, 6301–6335 (2014).

61.   B. Bölter, En route into chloroplasts: preproteins' way home. *Photosynth Res* **138**, 263–275
614   (2018).

615   62.   P. Chotewutmontri, K. Holbrook, B. D. Bruce, "Plastid Protein Targeting: Preprotein Recognition
616   and Translocation" in *International Review of Cell and Molecular Biology*, (Elsevier Inc., 2017), pp.
617   227–294.

618   63.   R. W. Christian, S. L. Hewitt, G. Nelson, E. H. Roalson, A. Dhingra, Plastid transit peptides-where
619   do they come from and where do they all belong? Multi-genome and pan-genomic assessment of
620   chloroplast transit peptide evolution. *PeerJ* **8** (2020).

621   64.   S. Reinbothe, *et al.*, Trna-dependent import of a transit sequence-less aminoacyl-trna synthetase
622   (Leurs2) into the mitochondria of arabidopsis. *Int J Mol Sci* **22** (2021).

623   65.   M. W. Murcha, *et al.*, Plant-specific preprotein and amino acid transporter proteins are required
624   for tRNA import into mitochondria. *Plant Physiol* **172**, 2471–2490 (2016).

625   66.   E. Skippington, T. J. Barkman, D. W. Rice, J. D. Palmer, Miniaturized mitogenome of the parasitic
626   plant viscum scurruloideum is extremely divergent and dynamic and has lost all nad genes. *PNAS*
627   **112**, E3515–E3524 (2015).

628   67.   G. Petersen, A. Cuenca, I. M. Møller, O. Seberg, Massive gene loss in mistletoe (Viscum,
629   Viscaceae) mitochondria. *Sci Rep* **5** (2015).

630   68.   A. Fender, *et al.*, Loss of a primordial identity element for a mammalian mitochondrial
631   aminoacylation system. *Journal of Biological Chemistry* **281**, 15980–15986 (2006).

632   69.   Y. Kumazawa, H. Himeno, K.-I. Miura, K. Watanabe, Unilateral Aminoacylation Specificity
633   between Bovine Mitochondria and Eubacteria. *The Journal of Biochemistry* **109**, 421–427 (1991).

634   70.   L. Bonnefond, M. Frugier, R. Giegé, J. Rudinger-Thirion, Human mitochondrial TyrRS disobeys the
635   tyrosine identity rules. *RNA* **11**, 558–562 (2005).

636   71.   L. Bromham, P. F. Cowman, R. Lanfear, Parasitic plants have increased rates of molecular
637   evolution across all three genomes. *BMC Evol Biol* **13** (2013).

638   72.   M. Safro, N. Moor, O. Lavrik, Phenylalanyl-tRNA Synthetases. *Madame Curie Bioscience Database*
639   (2013) (May 24, 2023).

640   73.   K. M. Haen, W. Pett, D. V. Lavrov, Parallel loss of nuclear-encoded mitochondrial aminoacyl-tRNA
641   synthetases and mtDNA-encoded tRNAs in cnidaria. *Mol Biol Evol* **27**, 2216–2219 (2010).

642   74.   I. Martin, O. Vinas, T. Mampel, F. Villarroya, Effects of cold environment on mitochondrial
643   genome expression in the rat: evidence for a tissue-specific increase in the liver, independent of
644   changes in mitochondrial gene abundance. *Biochemical Journal* **296**, 231–234 (1993).

645   75.   R. F. Hevner, M. T. T. Wong-Riley, Mitochondrial and Nuclear Gene Expression for Cytochrome
646   Oxidase Subunits Are Disproportionately Regulated by Functional Activity in Neurons. *The Journal
647   of Neuroscience* **13**, 1805–1819 (1993).

648   76.   E. A. Amiott, J. A. Jaehning, Mitochondrial Transcription Is Regulated via an ATP "Sensing"
649   Mechanism that Couples RNA Abundance to Respiration. *Mol Cell* **22**, 329–338 (2006).

650   77.   B. J. Haas, Transdecoder v5.5.0 (2018) (May 14, 2023).

651   78.   W. Shen, S. Le, Y. Li, F. Hu, SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
652   manipulation. *PLoS One* **11** (2016).

653   79.   C. Camacho, *et al.*, BLAST+: Architecture and applications. *BMC Bioinformatics* **10** (2009).

654   80.   K. Katoh, K. Misawa, K.-I. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence
655   alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059–3066 (2002).

656   81.   K. Tamura, G. Stecher, S. Kumar, MEGA11: Molecular Evolutionary Genetics Analysis Version 11.
657   *Mol Biol Evol* **38**, 3022–3027 (2021).

658   82.   A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
659   phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

660   83.   J. Sperschneider, *et al.*, LOCALIZER: Subcellular localization prediction of both plant and effector
661   proteins in the plant cell. *Sci Rep* **7** (2017).

662   84.   J. J. A. Armenteros, *et al.*, Detecting sequence signals in targeting peptides using deep learning.
663   *Life Sci Alliance* **2** (2019).

664   85.   F. Madeira, *et al.*, Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic
665   Acids Res* **50**, W276–W279 (2022).

666   86.   S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor in
667   *Bioinformatics*, (Oxford University Press, 2018), pp. i884–i890.

668   87.   A. Bankevich, *et al.*, SPAdes: A new genome assembly algorithm and its applications to single-cell
669   sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

670   88.   P. P. Chan, T. M. Lowe, "tRNAscan-SE: Searching for tRNA genes in genomic sequences" in
671   *Methods in Molecular Biology*, (Humana Press Inc., 2019), pp. 1–14.

672   89.   P. P. Chan, B. Y. Lin, A. J. Mak, T. M. Lowe, TRNAscan-SE 2.0: Improved detection and functional
673   classification of transfer RNA genes. *Nucleic Acids Res* **49**, 9077–9096 (2021).

Supplemental Dataset 1: Species and datasets used for gene presence/absence, Species and datasets used for retargeting analysis, Species and datasets used for tRNA content analysis.

Supplemental Dataset 2. Genes of interest and counts of gene copies for each species.

Supplemental Dataset 3: *S. himalayana* transit peptide sequences for transient expression and colocalization in *N. benthamiana*

Supplemental Dataset 4: tRNA content of organelle genomes.

Supplemental Dataset 5: Relative branch lengths and Tajima's relative rate test for aaRS in Ericales spp.

Supplemental Dataset 6: Heatmaps for probability of subcellular targeting to plastids or mitochondria on a protein-by-protein basis.

Figure 1: Presence/absence matrix of selected organellar and cytosolic aaRS and tRNA-processing enzymes shows major losses of organellar enzymes in heterotrophic lineages. Hue of boxes indicates metabolism (gray = autotroph, orange = hemiparasite, purple = heterotroph), shade indicates degree of presence/absence (white = no sequence found, light colors = partial sequence found, dark colors = complete or near complete sequence found). Note that there is no organellar GlnRS as plants typically use the bacterial-type GatCAB amidotransferase system to charge organellar Gln-tRNA. ArgRS is not presented as it has no distinct organellar enzyme; gene products are either cytosol-targeted, or triple-targeted to the cytosol, mitochondria, and plastids.

Figure 2: Proportion of retargeted cytosolic aaRS is higher in heterotrophs than autotrophs based on *in-silico* predictions. Proportion of enzymes with predicted retargeting (probability ≥ 0.25) to either mitochondria (top row) or plastids (bottom row) based on the outputs of LOCALIZER and/or TargetP. Autotrophs and heterotrophs are plotted in green and purple, respectively. Targeting predictions for proteins involved in mitochondrial oxidative phosphorylation (OPXHOS) are included as a positive control for mitochondrial targeting. Predictions for cell-wall synthesis associated enzymes are included as a negative control for organelle targeting. Reported *p*-values from *t*-tests or Mann-Whitney U tests depending on distribution of the data. Note that *R. cantleyi* and *S. himalayana* are not included in plastid retargeting analysis due to suspected loss of the plastome.

Figure 3: Cytosolic-type aaRS transit peptides from heterotroph *S. himalayana* can retarget enzymes to organelles in *Nicotiana benthamiana*. Transit peptides (TPs) from *S. himalayana* fused to GFP (middle right column) were transiently expressed in *N. benthamiana.* Chlorophyll autoflorescence (far left column) was used for colocalization with plastids. As a positive control for mitochondrial targeting, eqFP611 was tagged with the TP from mitochondrial isovaleryl-CoA dehydrogenase (middle left). The overlay of all three channels on far right shows mitochondrial localization for GFP tagged with AsnRS, GlyRS, and MetRS TPs. Minor/weak mitochondrial localization for ProRS TP. MetRS and ProRS TPs additionally target GFP to plastids. TrpRS and ThrRS TPs target GFP to plastids, cytosol, and nucleus. Note presence of punctate structures for ThrRS TP, possibly inclusion bodies. Arrows indicate good examples of plastid (yellow) and mitochondrial (white) and potential nuclear (red) localization. Scale indicated in top right panel is the same for all images.

Figure 4: Instances of both concordance and mismatch between organellar tRNA gene loss and aaRS loss/retargeting. Heterotrophic species are written in purple. *Arabidopsis thaliana* is included for reference. Left: mitogenome tRNAs. Right: plastome tRNAs. Color of boxes indicates presence/absence of tRNA. The – and + annotation denotes loss of corresponding organellar aaRS and retargeting of cytosolic tRNA respectively as described for figure 4. Retargeting predictions are based on individual predictions for each organelle. Note that loss and retargeting of the corresponding aaRS was not assessed for *trnR* as there are no organelle-specific aaRS orthologs responsible for charging this tRNA. Loss of GatCAB and associated cytosolic GlnRS retargeting are presented for *trnQ*. Additionally, plastid targeting predictions for Rafflesiaceae are not included here because of the suspected loss of the plastome in this taxon.

Supplemental Figure 1: Significant overlap between loss of organellar aaRS and retargeting of cytosolic counterpart in heterotrophs. Reported counts refer to number of aaRS "families" defined by amino acids (e.g., GluRS, PheRS, TrpRS, etc.). Organellar gene presence/absence determined as for figure 1. Partial sequences for organellar enzymes are counted as present. Predicted retargeting (probability ≥ 0.25) to either mitochondria or plastids was based on the outputs of LOCALIZER and/or TargetP. Note that aaRS where cytosolic targeting could not be tested (i.e., enzyme was not present, or N-terminus was incomplete) were excluded. Organellar enzyme present = shades of gray. Organellar enzyme absent = shades of orange. Cytosolic enzyme retargeting = hashing. No retargeting = solid bars. Reported *p*-value is from a Fisher's exact test for correlation between loss of organellar aaRS and retargeting of the corresponding cytosolic aaRS. Note that *R. cantleyi* and *S. himalayana* proteins are excluded from predictions of targeting to plastids as these species lack plastid genomes. Loss of GatCAB and associated cytosolic GlnRS retargeting are included in the counts.

Supplemental Figure 2: Targeting predictions for cytosolic-type aaRS. Deeper shades of green (plastid, right) and orange (mitochondrion, left) indicate probability of retargeting to that organelle is ≥ 0.25 based on both LOCALIZER-1.0.5 and TargetP-2.0. Lighter shades of orange/green indicate one of the two algorithms yielded a probability ≥ 0.25. White indicates probability for both algorithms was < 0.25. Gray indicates that the gene/transcript was not present or that the N-terminus was incomplete, so no targeting prediction could be made. Heterotrophs are written in purple, closely related autotrophs in black. *A. thaliana* is included as a reference.

Supplemental Figure 3: Cytosolic-type aaRS transit peptides from heterotroph *S. himalayana* transiently expressed in *Nicotiana benthamiana*. Transit peptides (TPs) from *S. himalayana* fused to GFP (middle right column) were transiently expressed in *N. benthamiana.* Chlorophyll autofluorescence (far left column) was used for colocalization with plastids. As a positive control for mitochondrial targeting, eqFP611 was tagged with the TP from mitochondrial isovaleryl-CoA dehydrogenase (middle left). Overlay of all three channels on far right. Arrows indicate good examples mitochondrial (white) localization. From top down: Cytochrome C (CytC) TP was included as a positive control showing mitochondrial targeting of a *S. himalayana* TP. AspRS TP (bottom row) primarily shows GFP fluorescence from non-mitochondrial punctate structures in the cytosol, potentially inclusion bodies. Thus, fluorescence patterns in this line appear to be an expression artifact and are not informative for addressing the question of targeting. Scale indicated in top right panel is the same for all images.

Supplemental Figure 4: Relative rates of sequence evolution of cytosolic and organellar aaRS are similar in heterotroph compared to autotrophic relative. Each point represents an aaRS branch length of *M. hypopitys* (heterotrophic) normalized to branch length of *R. williamsianum* (autotrophic) for canonically cytosolic and organellar aaRSs ($\log_{10}$ scale).

Supplemental Figure 5: Heterotrophic plants often lose chlorophyll biosynthesis enzymes yet retain heme biosynthesis enzymes in the tetrapyrrole pathway. "Common Pathway" = Enzymes for the base pathway for both heme and chlorophyll. "Heme" = Enzymes for heme (FC) and phytochromobilin (HO and HY2) biosynthesis. "Chlorophyll" = enzymes for the biosynthesis of chlorophyll A and B. Numbering of protein names represents their order within the pathway. For complete names of enzymes and corresponding *A. thaliana* gene ID, see supplemental dataset 2. Hue of boxes indicates metabolism (gray = autotroph, orange = hemiparasite, purple = heterotroph), shade indicates degree of presence/absence (white = no sequence found, light colors = partial sequence found, dark colors = complete or near complete sequence found).

Supplemental Figure 6: Targeting predictions for tetrapyrrole pathway enzymes suggests potential relocation of pathway to cytosol in *Rafflesiaceae*. Deeper shades of green indicate probability of targeting to the plastid is ≥ 0.25 based on both LOCALIZER and TargetP. Lighter shades of green indicate one of the two algorithms yielded a probability ≥ 0.25. White indicates probability for both algorithms was < 0.25. Gray indicates that the gene/transcript was not present or that the N-terminus was incomplete, so no targeting prediction could be made. Enzymes are divided based on whether they are involved in the common (shared) pathway, the heme/phytochromobilin branch, or the chlorophyll branch. Numbering of protein names indicates their order within the pathway. Heterotrophs are written in purple, closely related autotrophs in black. *A. thaliana* is included as a reference.

Supplemental Figure 7: Maximum likelihood tree for CysRS. Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for organellar and cytosolic CysRS. Orthologs for *A. thaliana* organellar CysRS (AT2G31170) outlined in green, orthologs for *A. thaliana* cytosolic CysRS (AT3G56300, AT5G38830) outlined in orange.

Supplemental Figure 8: Maximum likelihood tree for Protein Only RNase Ps (PRORPs). Bootstrap values are indicated at nodes. Branch length labels represent average substitions per site. Red arrow denotes *A. thaliana* sequences for nuclear and organellar PRORPs. Orthologs for *A. thaliana* organellar PRORP1 (AT2G32230) outlined in green, Orthologs for *A. thaliana* cytosolic PRORP2/3 (AT2G16650, AT4G21900) outlined in orange.

Supplemental Figure 9: Maximum likelihood tree for tRNase Z 1/2. Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for tRNAse Z type proteins. Orthologs for *A. thaliana* cytosolic tRNAse Z1 (AT1G74700) are outlined in orange, and orthologs for *A. thaliana* plastid tRNAse Z2 (AT2G04530) are outlined in green.

Supplemental Figure 10: Maximum likelihood tree for tRNase Z 3/4. Bootstrap values are indicated at nodes. Branch length labels represent average substitutions per site. Red arrow denotes *A. thaliana* sequences for tRNAse Z type proteins.