Gorse (*Ulex europeaus*) wastes with 5,6-dimethylbenzimidazole supplementation can support growth of vitamin B_{12} producing commensal gut microbes.

Ajay Iyer[1,3], Eva C. Soto-Martín[1], Gary A. Cameron[1], Petra Louis[1], Sylvia H. Duncan[1], Charles S. Bestwick[1], and Wendy R. Russell[1]

[1]University of Aberdeen, Rowett Institute, Foresterhill, Scotland AB25 2ZD

[2]University of Aberdeen, Institue of Medical Sciences, Foresterhill, Scotland AB25 2ZD

[3]**Current:** Teagasc, Fermoy, Co Cork, Ireland, P61 C996

Abstract

Many commensal gut microbes are recognized for their potential to synthesize vitamin B_{12}, offering a promising avenue to address deficiencies through probiotic supplementation. While bioinformatics tools aid in predicting B_{12} biosynthetic potential, empirical validation remains crucial to confirm production, identify cobalamin vitamers, and establish biosynthetic yields.

This study investigates vitamin B_{12} production in three human colonic bacterial species: *Anaerobutyricum hallii* DSM 3353, *Roseburia faecis* DSM 16840, and *Anaerostipes caccae* DSM 14662, along with *Propionibacterium freudenreichii* DSM 4902 as a positive control. These strains were selected for their potential use as probiotics, based on speculated B_{12} production from prior bioinformatic analyses. Cultures were grown in M2GSC, chemically defined media (CDM), and Gorse extract medium (GEM). The composition of GEM was similar to CDM, expect that the carbon and nitrogen source was replaced with protein-depleted liquid wastes obtained after subjecting Gorse to a leaf protein extraction process. B_{12} yields were quantified using liquid chromatography with tandem mass spectrophotometry.

The results suggest that the three butyrate-producing strains could indeed produce B_{12}, although the yields were notably low, and with no B_{12} being detected in the extracellular media. Next, GEM outperformed the conventional M2GSC growth medium in enhancing B_{12} production. The positive control, *P. freudenreichii* DSM 4902 produced B_{12} at concentrations ranging from $7\,ng \cdot mL^{-1}$ to $12\,ng \cdot mL^{-1}$. Univariate-scaled Principal Component Analysis of data from previous publications

investigating B_{12} production in *P. freudenreichii* revealed that B_{12} yields diminished when the carbon source concentration was $\leq 30\,\text{g·L}^{-1}$.

In conclusion, the protein-depleted wastes from the leaf protein extraction process from Gorse can be valorised as a viable substrate for culturing B_{12}-producing colonic gut microbes. Furthermore, this is the first report attesting to the ability of *A. hallii, R. faecis*, and *A. caccae* to produce B_{12}. However, these microbes seem unsuitable for industrial applications owing to low B_{12} yields.

Keywords: B_{12} production, cobalamin, gut commensals, circular economy.

Abbreviations: CDM: chemically defined medium, GEM: Gorse extract medium, PDF: protein depleted fraction, PCA: principal component analysis

Introduction

The biological role and molecular structure of vitamin B$_{12}$: Vitamin B$_{12}$ is essential for various enzymatic reactions, including isomerases, methyltransferases, and dehalogenases (Banerjee & Ragsdale, 2003), with B$_{12}$-dependent dehalogenases being exclusive to bacteria (Payne et al., 2015). It is a critical cofactor in the type 1 reaction, catalysing the isomerization of L-methylmalonyl-CoA to succinyl-CoA, that subsequently enters the Krebs cycle (Takahashi-Iñiguez et al., 2012). Furthermore, B$_{12}$ is crucial for the type 2 reaction of methyl transfer from 5-methyltetrahydrofolate to homocysteine, to produce tetrahydrofolate and methionine (Froese et al., 2019). Hindrance in this type 2 reaction owing to B$_{12}$ deficiency results in the malformation of red blood corpuscles, and manifests as pernicious anaemia (World Health Organization & Food and Agriculture Organization of the United Nations, 2004).

The bioactivity of vitamin B$_{12}$ hinges on three critical ligands, namely, the central cobalt ion located in the upper corrin ring, the methyl side-group, and 5,6-dimethylbenzimidazole (DMB) (see Figure 1). The methyl side-group and DMB are attached to the cobalt ion with a coordinate bond. The cobalt ion switches from the oxidised intermediate (III) to the reduced (I) state during the transfer of the methyl-group to homocysteine to form methionine. The oxidised cobalamin is subsequently re-alkylated with n-methyl-THF. The exact mechanism has been described previously by Matthews et al. (2008).

Dietary vitamin B$_{12}$ however may have either the methyl, adenosyl, hydroxyl or cyano group associated with the cobalt atom, with the cyano vitamer being a semi-synthetic stable derivative of B$_{12}$ (Herbert, 1988). Once absorbed in the body, the molecule is converted to the hydroxyl-form, that is then re-alkylated to the methyl form; ultimately entering the methionine synthesis pathway as described previously (Froese et al., 2019). There is variation in the second ligand moiety, where either the biologically active DMB, or its structural homologue, adenyl-ribofuranose phosphate, may be present. Only the DMB-based vitamer is capable of participating in the catalytic reaction facilitated by methionine synthase.

Harnessing microbes for industrial B$_{12}$ production: The biosynthesis of B$_{12}$ requires several enzymes, that are only found in the bacteria and archaea domains of life (Raux et al., 2000; Stamford et al., 1997). The complete set of genes required for *de novo* production of cobalamins; particularly B$_{12}$, has been observed in numerous bacteria such as *Pseudomonas dentrificans*, *Bacillus megaterium*, *Rhodobacter capusulatus*, *Bacillus megaterium* and *Thermosipho melanesiensis*, to name a few (Fang et al., 2017). Many other bacteria such as *Escherichia coli*, *Kosmotoga olearia* and *Fervidobacterium nodosum* can only use salvage pathways to synthesise cobalamin from precursors. The population of these microbes capable of cobalamin production thought salvage pathways play an important role in

Figure 1: The variable ligands of B_{12} vitamers. Variable side-groups form coordinate bonds with the central cobalt ion. The cyano-group is marked in blue as it is a semi-synthetic derivate of B_{12}. The ligands at location L_l determine if the cobalamin vitamer is biologically active in humans with 5,6-dimethybenzimidazole (DMB) or inactive with adenine (marked in red). B_{12} vitamers can be found with combinations of S_v and L_l ligands.

the micro-ecology of the niche they inhabit firstly through commensalistic or even symbiotic association with primary B_{12} producers (Raux et al., 2000), and secondly, by facilitating tertiary associations with other microbes that are heterotrophic for the precursor. In industrial setups however, candidate selection of B_{12} producers is largely influenced by the genetic stability of the strain, bioconversion efficiency, B_{12} yields, extracellular production capability, and rapid turnover.

Many microbial species; either naturally, or through genetic engineering, are capable of over-synthesising vitamin B_{12}. The most commonly used species in industry are *Propionibacterium freudenreichii*, *Propionibacterium shermanii* and *Pseudomonas denitrificans*. Growth conditions are generally aerobic as *de novo* biosynthesis of vitamin B_{12} is an energetically demanding process, particularly the synthesis

of DMB (Raux et al., 2000).

Consequently, there is great variability in the growth substrates used, depending on the metabolic
capability of the microbe employed. For example, many media formulations use additional supple-
mentation of cobalt, DMB, and in rare cases, haeme (Chen & Wolin, 1981). This helps cope with the
increased demand for micronutrients, and alleviates the metabolic burden of *de novo* DMB or corrin
synthesis. Interestingly, this has lead to investigations on waste stream valorisation with cobalt and
DMB supplementation for the growth of B_{12} producing bacteria. For example, *P. freudenreichii* has
been shown to grow on apple pomace and potato wastes for large scale production of propionic acid,
with vitamin B_{12} obtained as a secondary product in significant quantities (Piwowarek et al., 2022;
Piwowarek et al., 2023). Overall, such endeavours help generate high-value products and circularise
waste streams otherwise fated for landfill or waste-water treatment plants.

B_{12} from Gorse wastes using gut microbes: The investigation reported herein attempts to
valorise the liquid wastes generated from a previously described leaf protein extraction process, that
was obtained from the biomass of Gorse (*Ulex europaeus*), which has become invasive in many parts of
the world (Iyer et al., 2022). The process generated a protein depleted fraction (PDF) that contained
residual proteins, plant bioactives, and carbohydrates. In an effort to valorise this waste stream, three
butyrate-producing microbial candidates, namely *Anaerobutyricum hallii* DSM 3353, *Roseburia faecis*
DSM 16840, and *Anaerostipes caccae* DSM 14662, were investigated for their ability to grow on media
comprising of PDF as the sole carbon and nitrogen source. These microbial candidates were chosen
due to their potential to produce vitamin B_{12} (Soto-Martín et al., 2020) and their purported beneficial
impact in the human gut (Arruda et al., 2022; Montalban-Arques et al., 2021). *Propionibacterium*
freudenreichii DSM 4902, was used as a positive control for vitamin B_{12} production.

Materials and methods

All chemicals were purchased from Merck, (Darmstadt, Germany) unless stated otherwise. All water used was from the in-house Milli-Q®purification system. All experiments were performed in triplicate.

Media

Media preparation was carried out in an anaerobic cabinet (Don Whitley Scientific, West Yorkshire, UK) set for 10% H_2, 10% CO_2, and 80% N_2, with Milli-Q water which was first flushed with N_2 and autoclaved. The water was then allowed to equilibrate in the cabinet prior to preparation for 48 h.

Media preparation

M2GSC: The recipe for M2GSC is provided in **S1 Text**, and was prepared as described previously by Cummings and Macfarlane (1991). The medium was allowed to equilibrate in the anaerobic cabinet for 48 h before use.

Chemically defined medium (CDM):   CDM was prepared as described previously by Soto-Martín et al. (2020), except with the inclusion of 5,6-dimethylbenzimidazole (DMB, 20 mg·L^{-1}). The medium was filter-sterilised using 0.2 μm filter (Millipore, Merck, US) and allowed to equilibrate in the anaerobic cabinet. The composition of the CDM is provided in **S2 Text**.

Gorse extract medium (GEM):   GEM was prepared by replacing the carbon and nitrogen source of CDM with the gorse protein-depleted fraction (PDF). The composition of PDF is provided in a previous publication by Iyer et al. (2022). It contained residual protein (53.9 mg·g^{-1} dry mass) and polysaccharides / sugars (glucose equivalent of 88.9 mg·g^{-1} dry mass). Given that the plant mass was previously treated with cellulase as part of the protein extraction process, the carbohydrate profile of the PDF was expected to comprise of low molecular weight polysaccharides and glucose. Other monomers such as xylose, fucose, and glucuronic acid were expected to remain largely absent as pectinase, xylanase and other cell-wall digesting enzymes were not employed in the process.

The PDF was used at a final concentration of 25 g·L^{-1}. This meant that GEM contained about 1.35 g·L^{-1} of proteins and 2.3 g·L^{-1} of carbohydrates; values comparable to CDM.

Bacterial strains

Four bacterial strains were chosen for B_{12} production, namely, *P. freudenreichii* DSM 4902, *A. caccae* DSM 14662, *A. hallii* DSM 3353, and *R. faecis* DSM 16840, based on the work previously described by Soto-Martín et al. (2020). *A. caccae* DSM 14662, *A. hallii* DSM 3353, and *R. faecis* DSM 16840 were available from the in-house culture stock. *P. freudenreichii* DSM 4902 was purchased from NCIMB (Aberdeen, Scotland, NCIMB 5959).

Culture revival and growth

Microbial species were revived from frozen stocks using M2GSC (5 mL) in Hungate tubes (sealable with rubber caps) in anaerobic conditions and incubated at 37 °C for five days. The strains were sub-cultured by inoculating 0.5 mL of the culture in 4 mL of fresh M2GSC and incubation for two days. Growth was measured with optical density at 650 nm using Novaspec-II visible spectrophotometer (Pharmacia LKB Biotechnology AB, Uppsala, Sweden).

Production of B_{12} samples

From the revived stock, 200 µL was inoculated in triplicate in M2GSC, CDM or GEM (10 mL) and gently mixed through repeated aspiration. The culture was statically incubated at 37 °C for 48 h. Sample (1 mL) was drawn at time points 0 h, 6 h, 12 h, 24 h, 30 h, 36 h and 48 h and centrifuged at 10 000 × g for 20 min at 4 °C (Biofuge (Fresco), Heraeus Instruments, Germany). The supernatant was recovered and analysed for B_{12} content. The pellet was resuspended in PBS buffer (1 mL) and slowly cooled to −80 °C by first placing them in an ice-bath during sample collection, followed by incubation in −20 °C freezer overnight. Samples were then placed in a −80 °C freezer for a couple of days. This was performed to facilitate gradual growth of ice crystals to pierce and rupture the bacterial cell wall. Samples were then thawed to 4 °C overnight and sonicated (MSE Soniprep 150, UK) in two 10 s bursts to rupture the cells. The cells were centrifuged at 10 000 × g for 20 min and the supernatant was recovered and analysed for B_{12} content.

B_{12} Quantification using LC-MS

B_{12} estimation was performed using mass spectrophotometry in tandem with liquid chromatography as described elsewhere (Jiang et al., 2003). Calibration standards (all from Sigma-Aldrich; Saint Louis, USA) used were adenosylcobalamin, hydroxycobalamin, and methylcobalamin at concentrations 0.3 ng·mL^{-1}, 1.0 ng·mL^{-1}, 3.0 ng·mL^{-1}, 10 ng·mL^{-1}, 30 ng·mL^{-1}, 100 ng·mL^{-1} and 300 ng·mL^{-1} in dimethyl sulphoxide (DMSO). Cyanocobalamin (internal standard) was used at 2 µg·mL^{-1}. Samples or

standards were injected at $100\,\mu L$ with $10\,\mu L$ internal standard. Details of the LC-MS setup is provided in **S3 Text**.

Literature review

To retrieve previous literature on B_{12} production in *P. freudenreichii*, the search structure:

```
(propionibacterium AND (cobalamin OR B12))
```

was used in PubMed and Scopus. The relevant titles were selected and analysed by eye to obtain values for B_{12} yield, the carbohydrate concentration used (Carbon $g\cdot L^{-1}$), fermentation time (Time h), cobalt salt used ($mg\cdot L^{-1}$) and added DMB ($mg\cdot L^{-1}$). These values were used for the construction of a univariate-scaled PCA model to help compare and contextualise the B_{12} yields obtained in the work described herein. The details of the study and B_{12} are provided in **S1 Table**.

Statistical Analysis

All statistical analyses were performed using R (4.2.2) (R Core Team, 2021) using the tidyverse (2.0.0) (Wickham et al., 2019) package for data processing and visualisation. Normality of the data distribution was checked using the Shapiro-Wilks test. The Carbon ($g\cdot L^{-1}$), Time (h), and B_{12} ($mg\cdot L^{-1}$) values were $\log_{10}$ transformed to achieve normality.

The k-means analysis followed by univariate-scaled principal component analysis was performed using the cluster (2.1.4) (Maechler et al., 2022), and factoextra (1.0.7) (Kassambara & Mundt, 2020) packages. A dimension reduction model was made using previously published B_{12} production with *P. freudenreichii*. A k-means analysis (MacQueen algorithm) was performed to observe clustering in the raw data. The raw data was then subjected to a univariate-scaled PCA analysis and the clusters established from k-means analysis was overlaid on the plot to observe their relative positions. This PCA model was then used to interpolate the relative position of the data points representing the experimental conditions described in our study. This was used to understand which cluster of experimental conditions our setup matched and compare the corresponding B_{12} yields. The raw data and codes can be found at https://doi.org/10.17605/osf.io/3yb2r to reproduce the analyses.

Results and discussion

Growth profiles

Four bacterial strains, namely, *A. caccae* DSM 14662, *A. hallii* DSM 3353, *R. faecis* DSM 16840 and *P. freudenreichii* DSM 4902 were incubated in CDM, GEM and M2GSC over a period of 48 hours. The growth profiles are provided in Figure 2.

Figure 2: Growth profile of *A. caccae* DSM 14662, *A. hallii* DSM 3353, *P. freudenreichii* DSM 4902, and *R. faecis* DSM 16840 in chemically defined medium (CDM), gorse extract medium (GEM) and M2GSC.

All tested strains demonstrated robust growth in M2GSC and GEM media, reaching ΔO.D values >1.0. In contrast, CDM medium only supported the growth of *A. hallii* DSM 3353 and *P. freudenreichii* DSM 4902, with ΔO.D values <0.51.

In previous experiments by Soto-Martín et al. (2020), *A. caccae* DSM 14662 and *R. faecis* DSM 16840 were grown in CDM variants to assess trophic capabilities. They were initially grown in a semi-defined, vitamin-controlled medium using vitamin-free casein acid hydrolysate fortified with L-tryptophan, L-serine, L-threonine, L-glutamine, and L-asparagine to check for auxotrophy for each individual vitamin. Subsequently, they were grown in an amino-acid controlled CDM supplemented with all vitamins to check for auxotrophy for each amino acid. All three strains used here were able to grow well in both versions of CDM used by Soto-Martín et al. (2020). The growth profile in the panel for CDM in Figure 2 differs from previous reports, and may be explained by the pre-culture conditions. Soto-Martín et al. (2020) subjected the microbes through two passages in CDM before measuring their growth, while in this study, the microbes were revived in M2GSC and directly inoculated in CDM.

The GEM and CDM used were identical in their base compositions of added vitamins (as well as

absence of natural cobalamins), minerals, micronutrients, and precursors. The key difference lay in their carbohydrate and protein sources. While GEM contained residual proteins and peptides, CDM contained purified amino acids. This difference may have contributed to the lack of growth by *A. caccae* DSM 14662 and *R. faecis* DSM 16840 in CDM. Certain bacterial species are known to prefer peptides as a nitrogen source rather than individual amino acids (Kampen, 2014) under anaerobic conditions (Cotta & Russell, 1982). The uptake of individual amino acids is energetically demanding and requires active Na^{2+} mediated transmembrane transport (Heyne et al., 1991) and is the primary rate-limiting step in nitrogen uptake (Cotta & Russell, 1982). *A. caccae* DSM 14662 and *R. faecis* DSM 16840 are typically cultured in YCFA medium (La Rosa et al., 2019; Marquet et al., 2009) and the cobalamin-devoid, free-amino-acid based CDM may have been energetically incompatible for their growth.

B_{12} PRODUCTION IN EXPERIMENTAL SPECIES

The extent of vitamin B_{12} production in the experimental species is shown in Table 1 below.

Table 1: Total B_{12} measured in experimental fractions at 48 h incubation

Species	Media	Lysate	Supernatant
A. caccae	GEM	16.10±7.59	N/D
	CDM	N/G	N/G
	M2GSC	N/D	N/D
A. hallii	GEM	7.40±5.90	N/D
	CDM	N/D	N/D
	M2GSC	N/D	N/D
P. freudenreichii	GEM	15.21±1.00	N/D
	CDM	N/D	N/D
	M2GSC	4.23±0.60[a]	7.95±1.40
R. faecis	GEM	16.34±0.92	N/D
	CDM	N/G	N/G
	M2GSC	0.77±0.12[b]	N/D

Values expressed in mean±standard deviation µg·L^{-1}.
Marked values were significantly different from other yields measured across the microbial species for a given medium.
N/D = below detection level, N/G = no growth
B_{12} in all media at time 0 was N/D.

Vitamin B_{12} was below detection levels at the 0 h time point across all tested media. *P. freudenreichii* DSM 4902 and *R. faecis* DSM 16840 were able to produce B_{12} at the 48 h end-point in M2GSC as shown in Table 1. *P. freudenreichii* DSM 4902 showed B_{12} in the supernatant as well as the cell lysate fraction.

Vitamin B$_{12}$ was detected in the lysates of *A. caccae* DSM 14662 and *A. hallii* DSM 3353 cultured in GEM. In contrast, no B$_{12}$ was observed in M2GSC, despite it being the ideal growth medium. DMB was exclusively added to GEM and was likely critical to B$_{12}$ production. The growth of *A. caccae* and *A. hallii* in M2GSC without detectable B$_{12}$ in the lysates could be attributed to the synthesis of pseudo-B$_{12}$ vitamers, that were not detectable by the LC/MS method employed in the study. This may also explain the lower B$_{12}$ levels in M2GSC compared to GEM for *P. freudenreichii* DSM 4902 and *R. faecis* DSM 16840. Although these microbes demonstrated the ability to synthesize biologically active vitamers in anaerobic conditions, the addition of DMB likely alleviated the metabolic burden required for *de novo* synthesis, resulting in an increased total B$_{12}$ production in GEM compared to M2GSC.

In this experiment, B$_{12}$ measurement was performed using LC/MS, capable of detection at ng quantities in the sample. Physiologically, such quantities are far higher than the metabolic requirements of the cell (Burgess et al., 2009). This suggests that the chosen bacterial species were able to overproduce B$_{12}$, but the experimental conditions, or perhaps the media design was unsuitable for any real commercial application. Moreover, this contributes towards confirmation of the speculation (Engels et al., 2016) that *A. hallii* (formerly *E. hallii*) could produce B$_{12}$ vitamers in the gut.

Concerning probiotic applications, effective vitamin production necessitates high production in the extracellular environment. Since the chosen microbes failed to meet both requirements, they may not be ideal candidates for probiotic applications.

Of particular concern was the observed low B$_{12}$ yields for *P. freudenreichii* DSM 4902, which served as the positive control for this experiment. Generally, *P. freudenreichii* DSM 4902 has yields $\geq 0.2\,\mu\text{g}\cdot\text{mL}^{-1}$, yet the corresponding values noted in Table 1 were about 13$\times$ lower. A comparative analysis with other literature investigating *P. freudenreichii* for B$_{12}$ production was conducted to explore potential causes.

Component analyses using previous literature

The systematic online searches yielded 1651 hits from which 230 research titles were retrieved post title screening. A final list of 36 titles were selected based on the relevance of the research described therein. The data points for the concentration of the carbon source (g·L^{-1}), cobalt (mg·L^{-1}), DMB (mg·L^{-1}) and total fermentation time (hours) were retrieved and is presented in **S1 Table**. This only included cases of batch culturing of *P. freudenreichii*.

The data in **S1 Table** was first subjected to k-means clustering to look for grouping patterns in the various experimental conditions that may correspond to B$_{12}$ yields. Based on the "elbow method", three groups were deemed ideal to explain clustering in the data (**S Codes**). Next, the data was subjected

to a univariate-scaled PCA analysis and the corresponding grouping schema was overlaid to visualise their clustering on the PCA plot, as shown in Figure 3.

Figure 3: Univariate-scale PCA plot of the experimental conditions employed for B_{12} production using *P. freudenreichii* in previous publications. The point marked 'E' refers to the condition employed in this experiment. The position of the 'E' point was predicted by the model, which appears to associate to cluster 2. Ellipses represent 95% confidence limits of each cluster.

In Figure 3, the PCA model could account for 80.6% of the total observed variance in the data across Dimensions 1 and 2. The factors in the model, namely Carbon, Cobalt, DMB and Time accounted for 15.7%, 37.9%, 36.8% and 9.6% of the variance accounted in Dimension 1. In Dimension 2, Carbon, Cobalt, DMB and Time accounted for 39.4%, >0.1%, 0.1% and 60.6% of the variance.

Cluster 1 represents conditions with media employing high concentrations of carbon source, as well as a longer fermentation time, Cluster 2 represents the absence of precursors and low carbon source concentrations, and Cluster 3 represents high use of the precursors.

The point labelled "E" represents the experimental conditions and plots the closest to the mid-point of Cluster 2 ($d = 0.718$) compared to Clusters 1 ($d = 2.57$) and 3 ($d = 5.86$), where d represents the Euclidean distance between the points. This implies that the experimental condition described in this work was similar to that represented by Cluster 2. The median value of the conditions represented in each cluster is presented in Table 2.

Table 2: Median values of the experimental parameters and B_{12} yield represented in the clusters identified using k-means and PCA analysis.

Groups	Carbon	Cobalt	DMB	Time	B_{12}
Cluster 1	30 (25-35)	5.5 (5-12.7)	10 (0.9-15.0)	120 (72-16.0)	5.7 (1.3-16.1)
Cluster 2	10 (8.4-10)[a]	0 (0-0)	0 (0-0)	72 (72-102)	0.22 (0.16-2.21)
Cluster 3	40 (31.3-42.0)	40 (29-45)	70 (70-70)	48[b] (47.5-48.0)	6.7 (4.2-7.4)
Experimental	6[a]	10	20	48[b]	0.01

Values expressed as median (25% quantile to 75% quantile) to indicate the data distribution.
Carbon represents carbohydrates in the media (g·L^{-1}).
Cobalt and DMB (5,6-dimethylbenzimidazole) are expressed in mg·L^{-1}.
Time represents the total fermentation time (h).
B_{12} was the final reported yield (mg·L^{-1}).
Values with the same superscript were statistically similar using one-sample t-test.

The yield in Cluster 2 (0.2 mg·L^{-1}) was significantly lower than in Clusters 1 (5.7 mg·L^{-1}) and 3 (6.7 mg·L^{-1}), respectively ($F_{(2, 44)} = 5.298$, p = 0.009). These findings suggest that the availability of carbon source (via carbohydrates), was lower in the experimental setup described here compared to previous publications. As presented in Table 1, B_{12} content was predominantly retained within the cell lysate, indicating that the microbes did not secrete B_{12} extracellularly. Consequently, to achieve desired B_{12} yields, it is imperative to supply essential nutrients and precursors at sufficiently high concentrations, as indicated by the values presented in Table 2. **S1 Fig** illustrates boxplots comparing the levels of Cobalt, Carbon, DMB, and Time for each cluster.

Thus the data from Table 2 suggest that the yields of B_{12} are impacted when the carbohydrate concentration in the media is >30 g·L^{-1}.

LIMITATIONS AND SCOPE FOR FURTHER WORK

This experiment was designed to investigate two primary objectives: first, to determine the growth potential of selected gut microbes on media composed of PDF generated from Gorse wastes, and second, to assess their ability to produce vitamin B_{12}. The focus of this study was primarily to test B_{12} production in the candidate microbes and their ability to grow on Gorse waste media, rather than delve into the intricate mechanisms behind vitamin B_{12} production or enhancing yields for industrial applications.

Consequently, there were some aspects of this experiment that may warrant additional investigation. For example, *A. hallii* has previously been reported to produce pseudo-vitamin B_{12} (Belzer et al., 2017). Thus, while this experiment could demonstrate that biologically active B_{12} was produced by *A. hallii*, it is difficult to conclude if the microbe can facultatively produce the bioactive vitamer without a control medium devoid of DMB.

Expanding on the investigation, it's worth noting that other colonic microbes, such as *Lactobacillus reuteri*, are known to produce pseudo-cobalamin under anaerobic conditions. These microbes present intriguing candidates for future studies exploring their potential to produce the bioactive vitamer when DMB is introduced into the media.

Lastly, the individual vitamers and pseudo-vitamers were not characterised. More work will be required to understand if there are unique stoichiometries in which these vitamers are produced.

Conclusion

The investigation could show for the first time that *A. hallii*, *R. faecis* and *A. caccae* can produce vitamin B$_{12}$, albeit in very small quantities. However, these candidates are not suitable candidates for commercial B$_{12}$ production. The Protein Depleted Fraction (PDF) stands out as a valuable carbon and nitrogen source, capable of supporting the growth of gut microbes. This sets the stage for developing circular processes to cultivate microbes with higher B$_{12}$ production potential and pave the way for sustainable probiotic products.

Supporting information

M2GSC recipe The recipe for the M2GSC medium used for the growth of the experimental cells is provided section 1.1 in the Supplementary Materials file.

Chemically defined medium recipe The recipe for the the chemically defined medium (CDM) used for the growth of the experimental cells is provided section 1.2 in the Supplementary Materials file.

LC/MS conditions Details of the LC/MS setup to measure B$_{12}$ is provided section 1.3 in the Supplementary Materials file.

S Codes The codes used to analyse the data is annotated and available the Supplementary Codes file.

S1 Fig. Measures of experimental parameters in k-means clusters Median levels of B$_{12}$ yield, carbon source, precursors, and fermentation time in the k-means groups, shown in Figure 3. This can be found in the Supplementary Materials file.

Supplementary text and details Details on media recipe and LC/MS conditions. This can be found in the Supplementary Materials file.

S1 Table. B$_{12}$ yields in previous publications This can be found in the Supplementary Materials file.

Acknowledgments

We thank Freda Farquharson and Jenny Martin for their help and support in the laboratory. We thank Wiley Barton for their comments on the manuscript. This work was funded by the Scottish Government through Rural and Environment Science and Analytical Services Division (RESAS) as part of its strategic research programme. This work was performed as part of A.I's PhD project (2017 to 2021).

References

Arruda, H. S., Geraldi, M. V., Cedran, M. F., Bicas, J. L., Marostica Junior, M. R., & Pastore, G. M. (2022). Prebiotics and probiotics. In *Bioactive Food Components Activity in Mechanistic Approach* (pp. 55–118). Elsevier. https://doi.org/10.1016/B978-0-12-823569-0.00006-0

Banerjee, R., & Ragsdale, S. W. (2003). The Many Faces of Vitamin B $_{12}$: Catalysis by Cobalamin-Dependent Enzymes. *Annual Review of Biochemistry*, *72*(1), 209–247. https://doi.org/10.1146/annurev.biochem.72.121801.161828

Belzer, C., Chia, L. W., Aalvink, S., Chamlagain, B., Piironen, V., Knol, J., & de Vos, W. M. (2017). Microbial Metabolic Networks at the Mucus Layer Lead to Diet-Independent Butyrate and Vitamin B $_{12}$ Production by Intestinal Symbionts (N. Dubilier, Ed.). *mBio*, *8*(5), e00770–17. https://doi.org/10.1128/mBio.00770-17

Burgess, C. M., Smid, E. J., & van Sinderen, D. (2009). Bacterial vitamin B2, B11 and B12 overproduction: An overview. *International Journal of Food Microbiology*, *133*(1-2), 1–7. https://doi.org/10.1016/j.ijfoodmicro.2009.04.012

Chen, M., & Wolin, M. J. (1981). Influence of heme and vitamin B12 on growth and fermentations of Bacteroides species. *Journal of Bacteriology*, *145*(1), 466–471. https://doi.org/10.1128/jb.145.1.466-471.1981

Cotta, M. A., & Russell, J. B. (1982). Effect of Peptides and Amino Acids on Efficiency of Rumen Bacterial Protein Synthesis in Continuous Culture. *Journal of Dairy Science*, *65*(2), 226–234. https://doi.org/10.3168/jds.S0022-0302(82)82181-4

Cummings, J., & Macfarlane, G. (1991). The control and consequences of bacterial fermentation in the human colon. *Journal of Applied Bacteriology*, *70*(6), 443–459. https://doi.org/10.1111/j.1365-2672.1991.tb02739.x

Engels, C., Ruscheweyh, H.-J., Beerenwinkel, N., Lacroix, C., & Schwab, C. (2016). The Common Gut Microbe Eubacterium hallii also Contributes to Intestinal Propionate Formation. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.00713

Fang, H., Kang, J., & Zhang, D. (2017). Microbial production of vitamin B12: A review and future perspectives. *Microbial Cell Factories*, *16*(1), 15. https://doi.org/10.1186/s12934-017-0631-y

Froese, D. S., Fowler, B., & Baumgartner, M. R. (2019). Vitamin B $_{12}$, folate, and the methionine remethylation cycle—biochemistry, pathways, and regulation. *Journal of Inherited Metabolic Disease*, *42*(4), 673–685. https://doi.org/10.1002/jimd.12009

Herbert, V. (1988). Vitamin B-12: Plant sources, requirements, and assay. *The American Journal of Clinical Nutrition*, *48*, 852–858. https://doi.org/10.1093/ajcn/48.3.852

Heyne, R. I., de Vrij, W., Crielaard, W., & Konings, W. N. (1991). Sodium ion-dependent amino acid transport in membrane vesicles of Bacillus stearothermophilus. *Journal of Bacteriology*, *173*(2), 791–800. https://doi.org/10.1128/jb.173.2.791-800.1991

Iyer, A., Guerrier, L., Leveque, S., Bestwick, C. S., Duncan, S. H., & Russell, W. R. (2022). High throughput method development and optimised production of leaf protein concentrates with potential to support the agri-industry. *Journal of Food Measurement and Characterization*, *16*(1), 49–65. https://doi.org/10.1007/s11694-021-01136-w

Jiang, H., Pin, Z. G., & Yu, K. (2003). *LC-MS Analysis of Vitamin B12*. Waters. Retrieved March 15, 2021, from https://www.waters.com/content/dam/waters/en/app-notes/2003/720000758/720000758-en.pdf

Kampen, W. H. (2014). Nutritional Requirements in Fermentation Processes. In *Fermentation and Biochemical Engineering Handbook* (pp. 37–57). Elsevier. https://doi.org/10.1016/B978-1-4557-2553-3.00004-0

Kassambara, A., & Mundt, F. (2020). *Factoextra: Extract and Visualize the Results of Multivariate Data Analyses* (Version 1.0.7). https://CRAN.R-project.org/package=factoextra
Programmers: _:n3203.

La Rosa, S. L., Leth, M. L., Michalak, L., Hansen, M. E., Pudlo, N. A., Glowacki, R., Pereira, G., Workman, C. T., Arntzen, M. Ø., Pope, P. B., Martens, E. C., Hachem, M. A., & Westereng, B. (2019). The human gut Firmicute Roseburia intestinalis is a primary degrader of dietary beta-mannans. *Nature Communications*, *10*(1), 905. https://doi.org/10.1038/s41467-019-08812-y

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2022). *Cluster: Cluster Analysis Basics and Extensions.* (Version 2.1.4). https://CRAN.R-project.org/package=cluster
Programmers: _:n3786.

Marquet, P., Duncan, S. H., Chassard, C., Bernalier-Donadille, A., & Flint, H. J. (2009). Lactate has the potential to promote hydrogen sulphide formation in the human colon. *FEMS Microbiology Letters*, *299*(2), 128–134. https://doi.org/10.1111/j.1574-6968.2009.01750.x

Matthews, R. G., Koutmos, M., & Datta, S. (2008). Cobalamin-dependent and cobamide-dependent methyltransferases. *Current Opinion in Structural Biology*, *18*(6), 658–666. https://doi.org/10.1016/j.sbi.2008.11.005

Montalban-Arques, A., Katkeviciute, E., Busenhart, P., Bircher, A., Wirbel, J., Zeller, G., Morsy, Y., Borsig, L., Glaus Garzon, J. F., Müller, A., Arnold, I. C., Artola-Boran, M., Krauthammer, M., Sintsova, A., Zamboni, N., Leventhal, G. E., Berchtold, L., De Wouters, T., Rogler, G., . . . Scharl, M. (2021). Commensal Clostridiales strains mediate effective anti-cancer immune

response against solid tumors. *Cell Host & Microbe*, *29*(10), 1573–1588.e7. https://doi.org/10.1016/j.chom.2021.08.001

Payne, K. A. P., Quezada, C. P., Fisher, K., Dunstan, M. S., Collins, F. A., Sjuts, H., Levy, C., Hay, S., Rigby, S. E. J., & Leys, D. (2015). Reductive dehalogenase structure suggests a mechanism for B12-dependent dehalogenation. *Nature*, *517*(7535), 513–516. https://doi.org/10.1038/nature13901

Piwowarek, K., Lipińska, E., Hać-Szymańczuk, E., Kolotylo, V., & Kieliszek, M. (2022). Use of apple pomace, glycerine, and potato wastewater for the production of propionic acid and vitamin B12. *Applied Microbiology and Biotechnology*, *106*(17), 5433–5448. https://doi.org/10.1007/s00253-022-12076-w

Piwowarek, K., Lipińska, E., & Kieliszek, M. (2023). Reprocessing of side-streams towards obtaining valuable bacterial metabolites. *Applied Microbiology and Biotechnology*, *107*(7-8), 2169–2208. https://doi.org/10.1007/s00253-023-12458-8

R Core Team. (2021). *R: A Language and Environment for Statistical Computing* (Version 4.1.2). Vienna, Austria. https://www.R-project.org/
Programmers: _:n2750.

Raux, E., Schubert, H. L., & Warren*, M. J. (2000). Biosynthesis of cobalamin (vitamin B12): A bacterial conundrum: *Cellular and Molecular Life Sciences*, *57*(13), 1880–1893. https://doi.org/10.1007/PL00000670

Soto-Martín, E. C., Warnke, I., Farquharson, F. M., Christodoulou, M., Horgan, G., Derrien, M., Faurie, J.-M., Flint, H. J., Duncan, S. H., & Louis, P. (2020). Vitamin Biosynthesis by Human Gut Butyrate-Producing Bacteria and Cross-Feeding in Synthetic Microbial Communities (D. A. Relman, Ed.). *mBio*, *11*(4), e00886–20. https://doi.org/10.1128/mBio.00886-20

Stamford, N. J., Duggan, S., Li, Y., Alanine, A. I., Crouzet, J., & Battersby, A. R. (1997). Biosynthesis of vitamin B12: The multi-enzyme synthesis of precorrin-4 and factor IV. *Chemistry & Biology*, *4*(6), 445–451. https://doi.org/10.1016/S1074-5521(97)90196-4

Takahashi-Iñiguez, T., García-Hernandez, E., Arreguín-Espinosa, R., & Flores, M. E. (2012). Role of vitamin B12 on methylmalonyl-CoA mutase activity. *Journal of Zhejiang University SCIENCE B*, *13*(6), 423–437. https://doi.org/10.1631/jzus.B1100329

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., . . . Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

World Health Organization & Food and Agriculture Organization of the United Nations (Eds.). (2004). *Vitamin and mineral requirements in human nutrition* (2nd ed). World Health Organization ; FAO.
OCLC: ocm61189687.