Title: The multi-Siglec inhibitor AL009 reprograms suppressive macrophages and activates innate and adaptive tumor immunity

Authors: Sam C. Nalle,* Helen Lam, Ling Leung, Spencer Liang, Daniel Maslyar, Arnon Rosenthal

Affiliation:

Alector, Inc., South San Francisco, CA, USA.

*Corresponding author. Email: Sam.Nalle@alector.com

Abstract:

Sialic acid–binding immunoglobulin-type lectins (Siglecs) are cell surface receptors that regulate innate and adaptive immunity, with inhibitory Siglecs promoting immune tolerance. In the tumor microenvironment, overexpression of sialic acid glycans exploits inhibitory Siglec signaling, leading to a cancer-permissive phenotype. AL009 is an engineered Siglec-9-Fc fusion molecule that functions as a sialic acid trap and reprograms suppressive macrophages to activate an anti-tumor immune response. AL009 treatment of human myeloid-derived suppressor cells (MDSCs), an in vitro model of tumor-associated macrophages (TAMs), resulted in an increase in proinflammatory cytokines and chemokines, changes in cell surface markers, and potent relief of T cell inhibition in a co-culture assay. In syngeneic mouse tumor models, AL009 engineered with a mouse Fc (AL009m) reduced tumor growth as a monotherapy and in combination with the checkpoint inhibitor anti–PD-L1. In addition, AL009m synergized with the tumor-targeting therapy anti-TRP1 to reduce lung nodules in the B16-F10 intravenous model. Pharmacodynamic

marker analysis in syngeneic and humanized mouse tumor models supported an AL009

mechanism of action based on reprogramming TAMs and enhanced T cell activation. Future

clinical studies are warranted to further elucidate the safety and efficacy of AL009.

One Sentence Summary: Preclinical data for AL009 support investigation of this therapeutic

for treatment of cancers rich in suppressive macrophages.

Main Text:

INTRODUCTION

Sialic acid–binding immunoglobulin-like lectins (Siglecs) are immune regulatory molecules that

interact with sialic acid residues in the glycocalyx to modulate innate and adaptive immune

response (*1-3*). Inhibitory Siglecs, such as Siglec-3, -5, -7, -9, and -10, contain an

immunoreceptor tyrosine-based inhibition motif (ITIM) that recruits tyrosine phosphatases (such

as SHP1 and SHP2) involved in dampening the immune response to maintain homeostasis (*1, 2,*

4, 5). This is particularly important for recognition of self and resolution of inflammation (*6*).

However, some tumors have exploited this process by overexpressing sialic acid ligands on the

tumor cell surface to engage inhibitory Siglecs and evade immune detection, thus promoting an

immunosuppressive, cancer-permissive phenotype (*1, 5, 7, 8*). Disrupting the signaling between

immune-inhibitory Siglecs and their sialic acid ligands on hypersialylated cancer cells could

confer a therapeutic benefit, making both Siglecs and their sialic acid ligands attractive targets

for cancer immunotherapies (*2, 8, 9*).

Siglecs are expressed on various immune cells but predominantly on myeloid cells, including

macrophages and myeloid-derived suppressor cells (MDSCs) (*2, 5*). The immunoregulatory

activity of macrophages, whether proinflammatory or immunosuppressive, can change

depending on the chemokines, cytokines, and enzymes in the surrounding environment, with

tumor-associated macrophages (TAMs) being primarily immunosuppressive due to the nature of

the tumor microenvironment (TME) (*5, 10-13*). TAMs are not only influenced by the TME but

also contribute to the immunosuppressive signaling therein by releasing anti-inflammatory

chemokines and cytokines, such as IL-10 and TGF-β, which ultimately lead to T cell inactivation

(*10, 13*). Cell surface markers can be used to monitor the current immunomodulatory status of

macrophages; MHC-II, CD80, and CD86 are increased on proinflammatory macrophages, while

suppressive macrophages display elevated CD163 and CD206 (*5, 13-15*). TAM infiltration into

tumors has been associated with worse prognosis, particularly when high levels of Siglecs are

displayed (*16, 17*).

Targeted therapeutics designed to reprogram TAMs from an immunosuppressive to a

proinflammatory phenotype are currently being investigated (*11*). Thus far, antibodies against

macrophage receptor with collagenous structure (MARCO), the leukocyte immunoglobulin-like

receptor B (LILRB) family, Clever-1, and P-selectin glycoprotein ligand-1 (PSGL-1), as well as

CD40 agonists, have all shown potential in converting TAMs into proinflammatory effectors

(*18-22*). Interferon gamma (IFNɣ) has also shown potential to skew macrophages towards a

proinflammatory phenotype (*15, 18, 23*). The immunosuppressive characteristic of inhibitory

Siglecs along with their expression on TAMs make them ideal targets to modulate macrophage

function. However, since multiple inhibitory Siglecs are expressed on TAMs, broad targeting is

required to effectively interrupt immunosuppressive Siglec signaling (*4, 5*).

Here, we present data on AL009, an engineered multi-Siglec inhibitor that functions as a sialic

acid trap and reprograms suppressive macrophages to activate an anti-tumor immune response.

In vitro, AL009 effectively reprograms suppressive macrophages, as demonstrated by changes in

cell surface markers as well as increased secretion of proinflammatory cytokines. In animal

tumor models, monotherapy with AL009 or combination therapy with AL009 and anti–PD-L1 or

anti-TRP1 reduced tumor burden. Taken together, these data support further investigation of

AL009 as a targeted immunotherapy across a broad range of tumors high in Siglec-expressing

TAMs.

RESULTS

AL009 displays specific blocking between inhibitory Siglecs and their ligands

AL009 is composed of a Siglec-9 extracellular domain (ECD) and an engineered Fc to localize

the therapeutic to immune cells and promote cooperative binding of the ECD. Given the

redundancy of inhibitory Siglecs, an effective therapeutic would require broad blocking between

multiple inhibitory Siglecs and their respective ligands. To assess the degree to which AL009

could disrupt Siglec-ligand binding on MDSCs, the binding between Siglec-3, -5, -7, -9, and -10

Fc fusion proteins and human MDSCs was quantified in the presence of increasing

concentrations of AL009. With all the Siglec Fcs tested, an AL009 dose-dependent decrease in

binding was observed, demonstrating a broadly effective blockade of ligand recognition on cells

(Fig. 1A). In order to assess specific ligand blockade, a biochemical blocking assay using AL009

was established with a glycan array containing over 100 sialic acids. Using this assay format,

AL009 blocked binding of known sialic acids recognized by inhibitory Siglecs -3, -5, and -7

(Fig. 1B-D). As the binding portion of AL009 is composed of a Siglec-9 ECD, Siglec-9 binding

relative to other inhibitory Siglecs was evaluated side by side using a glycan array. Siglec-9

displayed an extensive sialic acid binding profile compared to other inhibitory Siglecs (Fig. S1),

helping to explain the ability of AL009 to block ligands recognized by multiple Siglecs on

MDSCs. Overall, these results demonstrate that AL009 possesses the ability to block specific

ligand binding of at least 5 inhibitory Siglecs present on myeloid cells.

Fig. 1. AL009 blocks binding of inhibitory Siglec receptors and their ligands. (A) AL009

blocked binding of Siglec-3, -5, -7, -9, and -10 from binding to MDSCs in a dose-dependent

98   manner. (**B-D**) In a glycan array containing over 100 sialic acids, AL009 blocked Siglec-3, -5,

99   and -7 from binding their respective ligands. Fc, fragment crystallizable; IgG, immunoglobulin

100   G; MDSCs, myeloid-derived suppressor cells; Siglec, sialic acid–binding immunoglobulin-type

101   lectin.

AL009 reprograms MDSCs

The phenotypic effect of AL009 blocking recognition between inhibitory Siglecs and their

ligands was assessed by incubating MDSCs with AL009 and evaluating cell surface markers,

cytokines, and chemokines. AL009 treatment resulted in a dose-dependent increase in immune

activation marker CD86 and a dose-dependent decrease in the immunosuppressive macrophage

marker CD163, consistent with reprogramming of MDSCs (Fig. 2A, 2B). Further, tumor

necrosis factor (TNF) production was increased in a dose-dependent manner (Fig. 2C). In order

to gain a greater understanding of the reprogramming effect of AL009 on MDSCs, RNAseq

analysis was performed on MDSCs after AL009 treatment. AL009 induced robust expression of

cytokines, chemokines, and co-stimulatory molecules compared to isotype control in MDSCs

(Fig. 2D).

We considered whether the reprogramming effect of AL009 treatment was specific to the

Siglec-9 ECD portion of the molecule. To evaluate this, MDSCs were treated with Siglec-3, -5, -

7, -9, or -10 ECD-Fc fusions and the reprogramming effect was assessed. The Siglec-9–mIgG

fusion resulted in the largest increase in activation markers CD86 and CD40 and the largest

reduction in suppressive macrophage markers CD163 and CD206 (Fig. S2). The Siglec-9–mIgG

fusion was also the only Siglec-Fc fusion that induced a dose-dependent increase in CCL4

production by MDSCs, consistent with reprogramming of these cells (Fig. S3).

Next, we examined whether AL009 was able to rescue MDSC suppression of T cells in a co-

culture system. MDSCs were treated with ligand-blocking antibodies against Siglec-3, -7, or -9,

a combination of all 3 antibodies, or AL009 prior to co-culture with activated T cells. The

amount of IFNɣ produced by the T cells was reduced in the presence of MDSCs and was not

rescued by any treatment except AL009 (Fig. 2E). In addition to increasing IFNɣ production in T

cells, we tested AL009's impact on T cell proliferation. After treatment with increasing

concentrations of AL009, T cell proliferation was quantified, with a half maximal effective

concentration (EC_{50}) of 1 to 2 nM AL009 (Fig. S4). Overall, AL009 is effective at potently

repolarizing MDSCs, resulting in phenotypic changes consistent with a proinflammatory

response and relief of T cell inhibition.

Fig. 2. AL009 repolarizes suppressive MDSCs. AL009 (**A**) enhanced CD86, (**B**) repressed

CD163, and (**C**) induced TNF secretion after treatment on MDSCs. (**D**) AL009 treatment of

134   MDSCs led to robust changes in gene expression. (**E**) AL009 rescued T cells suppressed by

135   MDSCs in a co-culture system. Error bars represent standard error of the mean. IFNɣ, interferon

136   gamma; IgG, immunoglobulin G; mAbs, monoclonal antibodies; MDSCs, myeloid-derived

137   suppressor cells; MedFI, median fluorescence intensity; Siglec, sialic acid–binding

138   immunoglobulin-type lectin; TNF, tumor necrosis factor. **$p < 0.01$, 2-sided t-test.

AL009 activity depends on the ECD and Fc regions

We hypothesized that both the Siglec-9 ECD and engineered Fc regions of AL009 were

important for the function of the molecule. To determine the contribution of the Siglec-9 ECD

portion of AL009, we first created domain mutation constructs. Production of the chemokine

CCL5 was induced with AL009, but that effect was ablated when the Siglec-9 ECD portion of

AL009 was deleted (Fig. 3A). Next, the Siglec-9 ECD and Fc fusion portions of AL009 were

decoupled, and MDSCs were incubated with plate-bound IgG to engage Fcγ receptors (FcγRs),

with or without a soluble AL009 mutant consisting of only the Siglec-9 ECD, then analyzed for

TNF production as a marker of MDSC programming (Fig. 3B). FcγR engagement by increasing

concentrations of plate-bound IgG induced a low level of TNF that was enhanced by the addition

of Siglec-9 ECD (Fig. 3C). We next asked if the Siglec-9 ECD could augment a separate

activation signal beyond FcγRs. Soluble anti-TREM1 or anti-TREM2 agonist antibodies were

incubated with MDSCs at increasing concentrations in addition to Siglec-9 ECD. Anti-TREM1

and anti-TREM2 had very minimal effects on MDSCs alone, but the addition of a fixed amount

of Siglec-9 ECD resulted in increased secretion of TNF in combination with TREM1 or TREM2

antibodies (Fig. 3D). These results demonstrate that Siglec-9 ECD alone is active and synergizes

with an activation signal in order to reprogram MDSCs.

Fig. 3. The Siglec-9 ECD relieves immune suppression in the presence of an activating

signal. (**A**) Truncation of the ECD domain of AL009 ablated its ability to induce CCL5. (**B**)

Schematic of experimental setup. (**C**) Levels of TNF produced by MDSCs upon IgG cross-

linking were increased in the presence of Siglec-9 ECD. (**D**) Levels of TNF produced by MDSCs

were not induced by anti-TREM1 or anti-TREM2 alone but were induced when Siglec-9 ECD

was present. ECD, extracellular domain; Fc, fragment crystallizable; IgG, immunoglobulin G;

MDSCs, myeloid-derived suppressor cells; MedFI, median fluorescence intensity; Siglecs, sialic

acid–binding immunoglobulin-type lectins; TNF, tumor necrosis factor.

The contribution of the Fc portion of AL009 was also investigated. The ability of AL009 to

relieve T cell suppression in an MDSC–T cell co-culture assay was greatly reduced when the Fc

portion of AL009 was silenced via mutations that ablate binding to FcγRs (Fig. 4A). To further

examine the contribution of the Fc portion of AL009 in binding MDSCs, dissociation constants

on cells were determined. The dissociation constant for AL009 when incubated with MDSCs

was 1.3 nM. However, when non-Fc receptor–expressing cells were used in place of MDSCs,

binding was much weaker (Kd = 220 nM). Similarly, when the Fc portion of AL009 was

silenced, binding to MDSCs was greatly reduced (Kd = 97 nM). In order to determine which Fc

gamma receptors were important for AL009 function, the ability of AL009 to increase CD86

expression at the cell surface, as a marker of MDSC reprogramming, was assessed in the

presence of antibodies against FcγRIIa, FcγRIIb, or a combination of FcγRIIa and FcγRIIb. The

N325S/L328F (NSLF) mutation contained within the Fc portion of AL009 prevents binding to

FcγRIIIa and FcγRIIIb; therefore, as expected there was no effect when MDSCs were incubated

with an FcγRIII blocking antibody prior to AL009 treatment. FcγRIIa blockade reduced AL009

activity, while FcγRIIb had no effect. Consistent with these results, the combination of FcγRIIa-

and FcγRIIb-blocking antibodies was equivalent to FcγRIIa blockade alone (Fig. 4B). Taken

together, these results suggest FcγRIIa is necessary for AL009 activity on MDSCs.

Fig. 4. AL009 activity requires Fc receptor interactions. (A) While AL009 was able to

reverse MDSC suppression and lead to higher levels of IFNγ compared to control, AL009 Fc

188   silent was not able to elevate IFNɣ to the same extent. (**B**) FcɣRIIa was required for functional

189   activity of AL009 on MDSCs. Data are shown as mean ± SEM. Fc, fragment crystallizable;

190   IFNɣ, interferon gamma; IgG, immunoglobulin G; MDSC, myeloid-derived suppressor cell;

191   MedFI, median fluorescence intensity; SEM, standard error of the mean.

AL009 preferentially binds myeloid cells

In whole human blood, binding of AL009 to a variety of immune cell subsets, including

monocytes, granulocytes, B cells, natural killer (NK) cells, T cells, and platelets, was assessed.

AL009 preferentially bound monocytes compared to the other immune cell subsets tested (Fig.

5A). To further elucidate cell subsets that are preferentially bound by AL009, donor blood

samples were used to compare AL009 binding to monocytes, macrophages, and MDSCs. The

highest binding observed was between AL009 and MDSCs, followed by non-activated M0

macrophages, and then classically activated M1 macrophages (Fig. 5B).

AL009m, a mouse reactive surrogate for AL009 containing a mouse Fc, was created to assess the

activity of AL009 in vivo. A biodistribution study was performed with AL009m by fluorescently

labeling the molecule and then injecting it intravenously in mice. AL009m accumulation was

higher than the isotype control in macrophage-rich mouse tissue samples, such as liver and

spleen, while no difference between AL009m or isotype control was seen in other tissues, such

as heart and lung (Fig. 5C-5F). The results from both the whole human blood binding study and

biodistribution study provide evidence that AL009 preferentially binds myeloid cells both in

vitro and in vivo.

Fig. 5. **AL009 preferentially binds myeloid cells.** (**A**) In whole human blood, AL009

preferentially bound monocytes over granulocytes, B cells, NK cells, T cells, and platelets. (**B**)

AL009 preferentially bound MDSCs over other myeloid subsets, including M0 and M1

macrophages and monocytes. (**C-F**) In a biodistribution study in mice, AL009m accumulated in

macrophage-rich tissues, such as liver and spleen (n = 3 mice per timepoint). Error bars represent

216 standard error of the mean. FLI, fluorescence intensity; IgG, immunoglobulin G; MDSC,

217 myeloid-derived suppressor cell; MedFI, median fluorescence intensity; NK, natural killer.

219 **AL009m demonstrates anti-tumor activity in mouse syngeneic tumor models**

220 Several syngeneic mouse tumor models were used to assess the anti-tumor efficacy of AL009m

221 in vivo. As a monotherapy, AL009m was tested in the E0771 triple negative breast cancer model.

222 Tumor volume was reduced and survival improved with AL009m treatment compared to isotype

223 control (Fig. 6A, 6B). To assess the impact of AL009m in combination with other therapies, the

224 B16-F10 intravenous injection model in combination with anti-TRP1 was utilized. In this model,

225 anti-TRP1 induces macrophage-mediated antibody-dependent cellular phagocytosis (ADCP) and

226 antibody-dependent cellular cytotoxicity (ADCC) of B16-F10 melanoma cells (*24*). Combination

227 treatment with AL009m and anti-TRP1 resulted in a significant reduction of lung nodules

228 compared to anti-TRP1 and isotype control (Fig. 6C). These results demonstrate that AL009m

229 can enhance ADCP/ADCC in vivo. To test the combination of AL009 with checkpoint

230 inhibitors, anti–PD-L1 was utilized along with AL009m in the MC38 and E0771 models. In both

231 models, anti–PD-L1 alone was not significantly different than isotype control, whereas

232 combination treatment with AL009m led to a significant reduction in tumor volume (Fig. 6D,

233 6E). Taken together, these efficacy results demonstrate that AL009m is active as a monotherapy

234 and in combination with an agent that induces ADCP/ADCC and the representative checkpoint

235 inhibitor anti–PD-L1.

Fig. 6. **AL009m demonstrated anti-tumor activity in multiple mouse syngeneic tumor**

models. (**A**) In the E0771 breast cancer model, AL009m treatment resulted in a significant

reduction in tumor volume as well as (**B**) increased survival. (**C**) In the intravenous B16-F10

tumor model, combination treatment with anti-TRP1 and AL009 resulted in reduced lung

241 nodules. Line at mean. (**D**) In the MC38 tumor model, combination treatment with AL009m and

242 anti–PD-L1 resulted in reduced tumor volume. (**E**) In the E0771 breast cancer model,

243 combination treatment with AL009m and anti–PD-L1 showed a synergistic effect. (D-E) Data

244 are shown as mean ± SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, 2-sided t-test. IgG,

245 immunoglobulin G; ns, not significant.

Pharmacodynamic markers show enhanced immune activation in the tumor and spleen

following AL009 treatment

To evaluate the pharmacodynamic effect of AL009m in the context of a tumor, spleen and tumor

samples from the E0771 mouse tumor models were analyzed by flow cytometry following

treatment with AL009m alone or in combination with anti–PD-L1. As a monotherapy, AL009m

treatment resulted in a significant decrease in Ly6C$^+$ cells, as a percentage of CD11b$^+$ cells,

indicating a decrease in MDSCs in tumor tissue (Fig. 7A). In spleen samples, an increase in

HLA-DR with AL009m treatment signaled activation of splenic macrophages (Fig. 7B). Further,

following AL009m treatment, there was a concurrent increase in activated CD4$^+$ cells in the

spleen, as measured by an enhanced percentage of CD69$^+$ cells (Fig. 7C). There was also an

increase in activated CD8$^+$ cells induced by AL009m that was further enhanced by anti–PD-L1

treatment, indicating a synergistic effect (Fig. 7D). In addition to the effects observed in the

tumor and spleen, AL009m also induced a dose-dependent increase in surface CD86 on

circulating monocytes (Fig. S5), similar to what was observed on human MDSCs following

AL009 treatment (Fig. 2A). The pharmacodynamic effect of AL009 was further evaluated in a

humanized mouse tumor model. In this humanized model, immunodeficient mice were engrafted

with human CD45$^+$ stem cells, implanted with a human A375 tumor weeks later, and treated

with AL009 approximately 2 weeks post implantation. Flow cytometry was used to analyze

tumor and spleen samples. Treatment with AL009 significantly decreased surface CD163 on

TAMs and splenic macrophages (Fig. 7E, 7F), demonstrating an in vivo reprogramming effect

consistent with the in vitro reprogramming observed with human MDSCs.

Fig. 7. Pharmacodynamic biomarker data provides evidence that AL009m repolarizes

myeloid cells and activates T cells in vivo. In the E0771 mouse tumor model, AL009m

treatment (**A**) decreased MDSCs in tumor tissue as shown by a decrease in Ly6C$^+$ cells, (**B**)

activated macrophages in spleen tissue as shown by increased Class II high cells, and (**C**)

activated CD4[+] T cells, as shown by an increase in CD69[+] cells. (**D**) CD8[+] T cells were also

increased with anti–PD-L1 treatment alone or in combination with AL009m treatment. (**E-F**) In

a humanized mice tumor model, AL009 treatment decreased levels of the pharmacodynamic

biomarker CD163, a marker for suppressive M2 macrophages, in the tumor and spleen. IgG,

immunoglobulin G; MDSC, myeloid-derived suppressor cell; MedFI, median fluorescence

intensity. Mean is shown. *$p < 0.05$, **$p < 0.01$, 2-sided t-test.

DISCUSSION

The current state of immunotherapy offers significant promise to treat cancer; however, while up

to half of all patients may be eligible for treatment with immunotherapy, less than 15% derive

clear benefit (*25*). TAMs heavily influence the TME and may comprise up to 50% of tumor

mass, which creates barriers to anti-tumor immunity and may underlie the resistance of some

patients to checkpoint inhibitors such as anti-PD1, anti–PD-L1, and anti-CTLA4 (*26*).

Consequently, an aspect of cancer immunotherapy drug development has shifted focus to

targeting the suppressive myeloid cells associated with tumors. There are currently TAM-

targeting therapeutics under investigation which aim to either deplete TAMs, block recruitment

of macrophages to the tumor, or reprogram TAMs to a proinflammatory phenotype (*26, 27*).

Depletion and blocking recruitment of macrophages disregard the anti-tumor potential of

creating reprogrammed, proinflammatory macrophages in the TME (*27*). Reprogramming TAMs

to a proinflammatory phenotype results in stimulation of cytotoxic T cells and tumor regression

in preclinical studies (*26*). Therapeutics aiming to reprogram TAMs may be effective as a

monotherapy or in synergy with checkpoint inhibitors (*26, 28*). As a TAM-reprogramming

therapeutic, AL009 provides two potential avenues for enabling immunotherapeutic activity: as a

monotherapy to drive macrophage modulation of the TME towards a more immune-responsive

environment, and in combination with checkpoint inhibitors to disrupt inherent immune-

mediated inhibition of the adaptive immune response. AL009 therefore represents a novel, multi-

functional therapeutic that has capacity for interrupting multiple negative regulatory signals in

TAMs with significant potential for treating cancer.

With its Siglec-9 ECD and engineered Fc region, AL009 targets immune cells to widely block

inhibitory signaling from Siglec receptors, leading to reprogramming of suppressive

macrophages. Siglecs are responsible for regulating immune homeostasis, with inhibitory Siglecs

playing a critical role in dampening immune responses (*4, 8*). The data from our experiments

confirm the ability of the Siglec-9 ECD of AL009 to bind sialic acid ligands of not only Siglec-9

but also other inhibitory Siglecs, thus broadly blocking immune-inhibitory Siglec–sialic acid

interactions. The disruption of this interaction allows anti-tumor activation signaling to proceed

(Fig. 8). The engineered Fc portion of AL009 facilitates localization to immune cells and

cooperative binding with the Siglec-9 ECD results in increased activity on innate immune cells.

(Fig. 8).

Deactivated anti-tumor response
Siglecs bind to sialic acid, inhibiting TAM activation

Activated anti-tumor response
AL009 binds sialic acid, relieving cis and trans Siglec-induced inhibition and allowing activation signaling to proceed

Fig. 8. AL009 mechanism of action. AL009 consists of an extracellular Siglec domain and an engineered Fc portion for immune cell targeting. The Siglec portion of AL009 binds sialic acid ligands in a cis or trans fashion, relieving Siglec-mediated immune suppression.

Overall, AL009 is a differential approach to test the therapeutic hypothesis that TAM reprogramming is an effective method of generating an anti-tumor effect in humans. Thus far, pre-clinical studies have not identified any significant toxicology signals. In addition, serum and immune cell pharmacodynamic biomarkers have been identified, and immunohistochemistry companion diagnostic development is underway. Taken together, these preclinical results support clinical testing of the safety and efficacy of AL009.

MATERIALS AND METHODS

Study design

The objective of this study was to evaluate the activity of AL009 in vitro and in mouse tumor models. Experiments were designed to elucidate the role of AL009's Siglec-9 ECD and its engineered Fc region, as well as its interactions with and impact on human myeloid cells. Additionally, AL009 pharmacodynamics and efficacy were tested in mouse models.

Animals and reagents

C57BL/6 mice harboring a BAC-expressing human Siglec-3, -7, and -9 (hSiglec BAC Tg) were generated and maintained by Alector. Expression of human Siglecs was confirmed by flow cytometry. Mice were maintained by crossing Tg^{het} mice with wildtype (WT) C57BL/6 mice, and all mice used in studies were Tg^{het} mice. Anti–PD-L1 was produced at Alector based on a reported sequence. Siglec Fcs and isotype control antibodies were produced at Alector unless otherwise stated. In mouse syngeneic tumor studies, a chimeric form of AL009 was produced with a mouse IgG2a Fc (AL009m). huNOG-EXL humanized mice were purchased from Taconic Biosciences (Germantown, NY) and housed at Alector.

Cell binding competition assay with AL009

MDSCs were generated from healthy human donors by differentiation of CD14$^+$ monocytes isolated using a RosetteSep human monocyte enrichment kit (STEMCELL Technologies). In order to generate MDSCs, CD14$^+$ monocytes were cultured for 7 days in Roswell Park Memorial Institute (RPMI) media containing 10-ng/mL human granulocyte-macrophage colony-stimulating factor (GM-CSF) and 10-ng/mL human IL-6 (PeproTech). For cell binding competition with AL009, MDSCs were incubated with titrating amounts of AL009, followed by

incubation with Siglec-3, -5, -7, -9, or -10 mouse IgG1 Fc fusion molecules. Bound Siglec-3, -5,

-7, -9, and -10 Fcs were assessed by detecting with an anti-mouse IgG-specific secondary

(Thermo Fisher) and analyzed via flow cytometry (BD Biosciences).

Glycan array

We assessed binding to a library of 300 glycan structures for Fc fusion proteins sourced from a

commercial vendor (R&D Systems) composed of a fusion of Siglec-3, -5, -7, -9, -10, or -15

ECDs with a human IgG1 Fc. Siglec fusion proteins and control human IgG1 were assessed for

glycan binding on the Glycan-300 (RayBiotech). The staining and analysis were performed as a

service by RayBiotech using the manufacturer's protocol. Briefly, the recombinant proteins were

incubated with the glycan array for 2 to 3 hours, followed by detection with anti-hIgG biotin and

streptavidin-Cy3. Samples were read and fluorescence was normalized across samples. For

blocking experiments, AL009 and mIgG1 or mIgG1 alone were applied first before incubation

with Siglec Fcs and human IgG.

MDSC stimulation and RNAseq analysis of MDSCs treated with AL009

MDSCs were treated with AL009 for 48 hours at 37° C and cell surface markers and cytokine

production were analyzed via flow cytometry and LEGENDplex (Biolegend), respectively. For

RNAseq analysis, MDSCs were treated with 10 µg/mL of AL009 for 24 hours. Cells were

harvested and pelleted, followed by resuspension in RLT buffer (Qiagen). RNA was extracted

with the RNeasy Plus Microkit (Qiagen). Illumina compatible libraries were created using

Lexogen QuantSeq 3' mRNA-Seq Library Prep Kit FWD and sequenced on a Nextseq P2.

MDSC–T cell co-culture

The activity of AL009 was evaluated alongside several different functional Siglec antibodies in a

MDSC–T cell co-culture assay. MDSCs were harvested and co-cultured with autologous CD8[+] T

cells in the presence of Dynabeads Human T-activator CD3/CD28 (Thermo) and either a Siglec-

Fc fusion or antibodies. Proliferation via carboxyfluorescein succinimidyl ester (CFSE, Thermo

Fisher) dilution and cytokine production via LEGENDplex were assessed after 3 to 5 days.

MDSC stimulation with plate-bound IgG and anti-TREM1/anti-TREM2

MDSCs were stimulated with plate-bound human IgG1 with or without 10-µg/mL Siglec-9 ECD

co-incubation for 48 hours followed by analysis of TNF production (Meso Scale Discovery).

Separately, MDSCs were stimulated with anti-TREM1 or anti-TREM2 agonistic antibodies with

or without Siglec-9 ECD for 48 hours followed by analysis of TNF production.

Fcγ receptor blockade

MDSCs were incubated with the 10-µg/mL FcγRIIa (clone IV.3)- or FcγRIIb (clone 2B6)-

blocking antibodies alone or in combination for 30 minutes on ice, followed by treatment with

3.3-µg/mL AL009 for 24 hours at 37° C and analysis of cell surface markers by flow cytometry.

Specific binding of AL009

For peripheral immune cell binding specificity, healthy human blood was incubated with AF647-

labeled AL009 using SiteClick reagents (Thermo Fisher) followed by analysis via flow

cytometry. For myeloid binding, healthy human monocytes from peripheral blood, M0

macrophages differentiated from human monocytes with 50-ng/mL human M-CSF, M1

macrophages differentiated from human monocytes with 50-ng/mL M-CSF, 100-ng/mL LPS

(InvivoGen), and 50-ng/mL IFNγ (PeproTech), or MDSCs, were incubated with AF647-labeled

AL009 followed by analysis via flow cytometry.

Biodistribution of AL009

Female C57BL/6 mice bearing established subcutaneous MC38.NCI.TD1 murine colon carcinoma were injected intravenously with 10-mg/kg AF647-labeled AL009m. Ex vivo fluorescence imaging (FLI) using the IVIS Spectrum imaging system was performed on liver, spleen, heart, and lungs collected from each mouse at 1, 6, 24, 48, and 96 hours post injection. Three animals were imaged per time point.

Effect of AL009 in mouse models

The MC38 and B16-F10 cell lines were purchased from ATCC. The E0771 cell line was purchased from MI Bioresearch. Female S3/7/9 Tg mice that were 7 to 12 weeks old were implanted with MC38 or E0771 cells subcutaneously in the flank region at a density of 1.0×10^5 or 4.0×10^5 cells per mouse, depending on the model. When tumors reached an average volume of ~100 mm^3, mice were randomized into treatment groups and dosing was initiated. Mice received intraperitoneal injections of 10- to 20-mg/kg AL009m or isotype control with or without 3-mg/kg anti–PD-L1 twice per week for 3 weeks. Tumors were measured twice per week with calipers. For the B16-F10 intravenous model, 1.5×10^5 B16-F10 cells were administered via intravenous injection. Mice were treated with 27-μg/mouse anti-TRP1 (BioXCell) on day 1 in addition to 10-mg/kg AL009m or isotype control on days 1, 4, 7, 10, and 13. A satellite group of animals was dosed only with control antibodies, and several mice in this group were euthanized every 2 days starting on day 9 to assess lung nodules. When the satellite group reached 50 to 100 lung nodules, the remaining experimental groups were euthanized for analysis. Statistical testing was performed by 2-way Student's t-test.

Pharmacodynamic biomarkers

Siglec-3/7/9 BAC Tg mice were implanted with E0771 tumors and randomized for treatment as

described above. The mice received intraperitoneal 20-mg/kg AL009m or isotype control on

days 0, 4, and 7, and were sacrificed on day 8. Splenic and tumor tissues were mechanically

dissociated and passed through a 70-μm filter to obtain single-cell suspensions. Peripheral blood

was treated with ammonium-chloride-potassium (ACK) lysis buffer to lyse red blood cells and

washed with phosphate-buffered saline (PBS). Cell surface markers on immune cell subsets were

analyzed by flow cytometry.

For pharmacodynamic analysis of humanized mice, 14- to 16-week-old female huNOG-EXL

mice were implanted with A375 melanoma cells subcutaneously in the flank region at a density

of with 3.0×10^6 cells per mouse. When tumors reached an average volume of 200 to 400 mm^3,

mice received intraperitoneal injections of 10-mg/kg AL009 or isotype control on days 0 and 3

and were sacrificed 24 hours later. Cell surface markers on immune cells subsets were analyzed

by flow cytometry.

Statistical analyses

Statistical analysis was performed using GraphPad Prism v9. Data are displayed as mean ± SEM.

A 2-tailed t-test was used for comparisons between 2 groups. A 1-way ANOVA followed by the

Holm-Sidak method of multiple comparisons was used for comparisons of more than 2 groups.

A *p* value of < 0.05 was considered significant.

Study approval

All in-house animal studies were reviewed by, approved by, and performed in compliance with

Alector's Institutional Animal Care and Use Committee protocols. Outsourced studies were

conducted in accordance with the applicable Covance or Charles River Laboratories standard

operating procedures, which approved these studies.

Supplementary Materials

Supplementary Materials and Methods

Activation of myeloid cells by Siglecs

The ability of different Siglec family members to reprogram MDSCs was evaluated. The ECDs

of Siglec-3, -5, -7, -9, and -10 were expressed as mouse IgG1 fusion proteins and incubated with

human MDSCs for 48 hours. Flow cytometry to assess cell surface molecules and analysis of

conditioned media for chemokines were performed.

Supplementary Figures

Fig. S1. Siglec-9-Fc demonstrated binding of a variety of sialic acid ligands.

Fig. S2. Siglec-9-Fc has a unique impact on MDSC activation and suppression markers.

Fig. S3. Siglec-9-Fc induced CCL4 production by MDSCs.

Fig. S4. AL009 relieved MDSC-mediated suppression of T cells with an EC_{50} of 1-2 nM.

Fig S5. AL009 induced dose-dependent increases in CD86 on monocytes.

References

1. P. R. Crocker, J. C. Paulson, A. Varki, Siglecs and their roles in the immune system. *Nat. Rev. Immunol.* **7**, 255-266 (2007).

2. H. Laubli, A. Varki, Sialic acid-binding immunoglobulin-like lectins (Siglecs) detect self-associated molecular patterns to regulate immune responses. *Cell. Mol. Life Sci.* **77**, 593-605 (2020).

3. J. S. Dos Reis, I. Diniz-Lima, M. Santos, P. M. Barcelos, K. M. da Costa, R. D. C. Valente, L. S. Chaves, L. P. de Campos, A. C. Dos Santos, R. G. Correia de Lima, D. Decote-Ricardo, A. Morrot, J. O. Previato, L. Mendonca-Previato, C. G. Freire-de-Lima, L. M. D. Fonseca, L. Freire-de-Lima, The blessed union of glycobiology and immunology: A marriage that worked. *Medicines (Basel)* **10**, (2023).

4. M. S. Macauley, P. R. Crocker, J. C. Paulson, Siglec-mediated regulation of immune cell function in disease. *Nat. Rev. Immunol.* **14**, 653-666 (2014).

5. K. C. M. Santegoets, P. R. Gielen, C. Bull, B. M. Schulte, E. D. Kers-Rebel, B. Kusters, S. Bossman, M. Ter Laan, P. Wesseling, G. J. Adema, Expression profiling of immune inhibitory Siglecs and their ligands in patients with glioma. *Cancer Immunol. Immunother.* **68**, 937-949 (2019).

6. J. C. Paulson, M. S. Macauley, N. Kawasaki, Siglecs as sensors of self in innate and adaptive immune repsonses. *Ann. N. Y. Acad. Sci.* **1253**, 37-48 (2012).

7. E. Rodriguez, S. T. T. Schetters, Y. van Kooyk, The tumour glyco-code as a novel immune checkpoint for immunotherapy. *Nat. Rev. Immunol.* **18**, 204-211 (2018).

8. H. Laubli, S. C. Nalle, D. Maslyar, Targeting the Siglec-Sialic acid immune axis in cancer: Current and future approaches. *Cancer Immunol. Res.* **10**, 1423-1432 (2022).

9.   G. Murugesan, B. Weigle, P. R. Crocker, Siglec and anti-Siglec therapies. *Curr. Opin.*

Chem. Biol. **62**, 34-42 (2021).

10.   A. N. Orekhov, V. A. Orekhova, N. G. Nikiforov, V. A. Myasoedova, A. V. Grechko, E.

B. Romanenko, D. Zhang, D. A. Chistiakov, Monocyte differentiation and macrophage

polarization. *Vessel Plus* **3**, 10 (2019).

11.   J. Zhou, Z. Tang, S. Gao, C. Li, Y. Feng, X. Zhou, Tumor-associated macrophages:

recent insights and therapies. *Front. Oncol.* **10**, 188 (2020).

12.   V. Kumar, S. Patel, E. Tcyganov, D. I. Gabrilovich, The nature of myeloid-derived

suppressor cells in the tumor microenvironment. *Trends Immunol.* **37**, 208-220 (2016).

13.   Y. Chen, Y. Song, W. Du, L. Gong, H. Chang, Z. Zou, Tumor-associated macrophages:

An accomplice in solid tumor progression. *J. Biomed. Sci.* **26**, 78 (2019).

14.   M. Lv, X. Zhuang, S. Shao, X. Li, Y. Cheng, D. Wu, X. Wang, T. Qiao, Myeloid-derived

suppressor cells and CD68(+)CD163(+)M2-like macrophages as therapeutic response

biomarkers are associated with plasma inflammatory cytokines: A preliminary study for

non-small cell lung cancer patients in radiotherapy. *J. Immunol. Res.* **2022**, 3621496

(2022).

15.   H. Cai, Y. Zhang, J. Wang, J. Gu, Defects in macrophage reprogramming in cancer

therapy: The negative impact of PD-L1/PD-1. *Front. Immunol.* **12**, 690869 (2021).

16.   S. Q. Qiu, S. J. H. Waaijer, M. C. Zwager, E. G. E. de Vries, B. van der Vegt, C. P.

Schroder, Tumor-associated macrophages in breast cancer: Innocent bystander or

important player? *Cancer Treat. Rev.* **70**, 178-189 (2018).

494 17. N. Xiao, X. Zhu, K. Li, Y. Chen, X. Liu, B. Xu, M. Lei, J. Xu, H. C. Sun, Blocking

495   siglec-10(hi) tumor-associated macrophages improves anti-tumor immunity and enhances

496   immunotherapy for hepatocellular carcinoma. *Exp. Hematol. Oncol.* **10**, 36 (2021).

497 18. Z. Duan, Y. Luo, Targeting macrophages in cancer immunotherapy. *Signal Transduct.*

498   *Target Ther.* **6**, 127 (2021).

499 19. M. Viitala, R. Virtakoivu, S. Tadayon, J. Rannikko, S. Jalkanen, M. Hollmen,

500   Immunotherapeutic blockade of macrophage clever-1 reactivates the CD8(+) T-cell

501   response against immunosuppressive tumors. *Clin. Cancer Res.* **25**, 3289-3303 (2019).

502 20. A. M. Georgoudaki, K. E. Prokopec, V. F. Boura, E. Hellqvist, S. Sohn, J. Ostling, R.

503   Dahan, R. A. Harris, M. Rantalainen, D. Klevebring, M. Sund, S. E. Brage, J. Fuxe, C.

504   Rolny, F. Li, J. V. Ravetch, M. C. Karlsson, Reprogramming tumor-associated

505   macrophages by antibody targeting inhibits cancer progression and metastasis. *Cell Rep.*

506   **15**, 2000-2011 (2016).

507 21. L. La Fleur, J. Botling, F. He, C. Pelicano, C. Zhou, C. He, G. Palano, A. Mezheyeuski,

508   P. Micke, J. V. Ravetch, M. C. I. Karlsson, D. Sarhan, Targeting MARCO and IL37R on

509   immunosuppressive macrophages in lung cancer blocks regulatory T cells and supports

510   cytotoxic lymphocyte function. *Cancer Res.* **81**, 956-967 (2021).

511 22. A. Mantovani, P. Allavena, F. Marchesi, C. Garlanda, Macrophages as tools and targets

512   in cancer therapy. *Nat. Rev. Drug Discov.* **21**, 799-820 (2022).

513 23. G. Genard, S. Lucas, C. Michiels, Reprogramming of tumor-associated macrophages

514   with anticancer therapies: radiotherapy versus chemo- and immunotherapies. *Front.*

515   *Immunol.* **8**, 828 (2017).

24.   H. Benonisson, H. S. Sow, C. Breukel, J. Claassens, C. Brouwers, M. M. Linssen, M. F.

Fransen, M. Sluijter, F. Ossendorp, T. van Hall, J. S. Verbeek, High FcgammaR

expression on intratumoral macrophages enhances tumor-targeting antibody therapy. *J.*

Immunol. **201**, 3741-3749 (2018).

25.   A. Haslam, V. Prasad, Estimation of the percentage of US patients with cancer who are

eligible for and respond to checkpoint inhibitor immunotherapy drugs. *JAMA. Netw.*

Open. **2**, e192535 (2019).

26.   P. Pathria, T. L. Louis, J. A. Varner, targeting tumor-associated macrophages in cancer.

Trends Immunol. **40**, 310-327 (2019).

27.   N. Kumari, S. H. Choi, Tumor-associated macrophages in cancer: recent advancements in

cancer nanoimmunotherapies. *J. Exp. Clin. Cancer Res.* **41**, 68 (2022).

28.   X. Xiang, J. Wang, D. Lu, X. Xu, Targeting tumor-associated macrophages to synergize

tumor immunotherapy. *Signal Transduct. Target Ther.* **6**, 75 (2021).

Acknowledgments: Editorial support and publication assistance were provided by SCIENT

Healthcare Communications (Cedar Knolls, NJ, USA).

Funding: All studies were supported by Alector, Inc.

Author contributions:

Conceptualization: SCN, SL, AR

Investigation: SCN, HL, LL

Analysis: SCN, HL, LL

Writing – original draft: SCN, DM

Writing – review and editing: SCN, DM

Competing interests: All authors are shareholders of Alector, Inc., or were shareholders of

Alector, Inc., when the studies were being conducted. SCN and SL are inventors on a patent

application related to this work.

Data and materials availability:

All data are available in the main text or supplementary materials.