3   **Seasonally migratory songbirds have different historic population size characteristics than**
4   **resident relatives.**
5

Kevin Winker[a,1] and Kira Delmore[b]

[a] University of Alaska Museum and Department of Biology and Wildlife, Fairbanks, Alaska, USA
99775. kevin.winker@alaska.edu

[b] Department of Biology, Texas A&M University, College Station, Texas, USA 77843.
kdelmore@bio.tamu.edu

[1] To whom correspondence may be addressed.
Kevin Winker
University of Alaska Museum
907 Yukon Drive
Fairbanks, AK 99775
907-474-7027; fax 907-474-1987
Email: kevin.winker@alaska.edu

ORCID IDs:

Kevin Winker: https://orcid.org/0000-0002-8985-8104
Kira Delmore: https://orcid.org/0000-0003-4108-9729

Running title: Migrant vs resident bird population sizes

Keywords: *Catharus* thrushes, effective population size, North and South America, PSMC, taxon
cycles.

Numbers of words: Abstract (208), main text (4,406).
Number of references: 78
Numbers of: figures (2), tables (1). (Plus 4 supplementary figures and 2 supplementary tables.)
36
37

Abstract

Modern genomic methods enable estimation of a lineage's long-term effective population sizes back to its origins. This ability allows unprecedented opportunities to determine how adoption of a major life-history trait affects lineages' populations relative to those without the trait. We used this novel approach to study the population effects of the life-history trait of seasonal migration in evolutionary time. Seasonal migration is a common life-history strategy, but its effects on long-term population sizes relative to lineages that don't migrate are largely unknown. Using whole-genome data, we estimated effective population sizes over millions of years in closely related seasonally migratory and resident lineages in a group of songbirds. Our main predictions were borne out: Seasonal migration is associated with larger effective population sizes (N_e), greater long-term variation in N_e, and a greater degree of initial population growth than among resident lineages. Initial growth periods showed phylogenetic signal, and their length (0.75-4.3 Myr) parallels the expansion and adaptation phases of taxon cycles, a framework of lineage expansion and eventual contraction over time encompassing biogeography and evolutionary ecology. Heterogeneity among lineages is noteworthy, despite geographic proximity (including overlap) and close relatedness. Seasonal migration imbues these lineages with fundamentally different population size attributes through evolutionary time compared to closely related resident lineages.

KEYWORDS *Catharus*, effective population size, North America, PSMC, South America, taxon cycles, thrushes, Turdidae.

INTRODUCTION

Modern genomic methods enable estimation of a lineage's long-term effective population sizes back to its origins (Li & Durbin, 2011; Mazet et al., 2015; Nadachowska-Brzyska et al., 2015). This ability allows unprecedented opportunities to study the populational effects of past evolutionary and climatic phenomena on organismal lineages, and this new area of paleodemographics is likely to grow rapidly (Nadachowska-Brzyska et al. 2021, Germain et al. 2023a,b). We use this approach in a novel way to test hypotheses about how adoption of a major life-history trait affected lineages' populations relative to those without the trait.

Seasonal migration is a common life-history strategy among the world's birds, but its effects on long-term population sizes relative to lineages that don't have this trait are largely unknown (Newton, 2008; Rappole, 2013). Demographic differences between seasonally migratory and resident lineages have been studied for decades, but, thus far, ecological, behavioral, and genetic approaches have necessarily been focused on recent and microevolutionary phenomena (e.g., Greenberg, 1980; Pérez-Tris et al., 2004; Sandercock & Jaramillo, 2002). Here we look much deeper, contrasting changes in effective population sizes through millions of years between closely related migratory and resident lineages.

Thrushes in the genus *Catharus* and this group's sister *Hylocichla mustelina* (Aves: Turdidae) are a model songbird system for studying the evolutionary effects of seasonal migration. This group has both migratory and resident lineages, and it has been important in the study of seasonal migration, divergence, and speciation (e.g., Delmore et al,. 2015, 2016; Outlaw et al., 2003; Ruegg et al., 2014; Termignoni-Garcia et al., 2022; Voelker et al., 2013; Winker, 2000, 2010; Winker and Pruett, 2006). These animals are relatively small, omnivorous or insectivorous, forest-related birds with lineages that are long-distance seasonal migrants in North America and others that are resident in tropical

North and South America (Collar, 2005). The breeding grounds of the migratory lineages span

Siberia, Alaska, northern Canada, and the United States, and their wintering ranges include Mexico,

the Caribbean, Central America, and tropical South America. In contrast, the resident lineages

occupy Mexico, Central America, and northwestern South America (Collar, 2005; Fig. S1).

We made three predictions about effective population sizes (N_e) in seasonally migratory

versus resident lineages of these thrushes: 1) N_e would on average be higher among seasonal

migrants; 2) variation in population size would also be higher; and 3) early population growth would

be higher as seasonal migration opens novel ecological and geographic space, achieving the first

prediction.

These predictions stem from the following reasoning. We can think of North America

roughly as an inverted triangle, with substantially more geographic space in the north than in the

south. Among these thrushes, the migratory lineages breed in that northern, on average, larger space.

The resident lineages instead currently occupy, on average, smaller geographic ranges, even in South

America, where they are largely montane (Fig. S1). Thus, larger breeding ranges among migrants, on

average, should yield higher population sizes. But climatic variability is also higher with increased

latitude, often temporarily rendering vast swaths of northern North America unoccupiable to many

long-distance migrant species (Pielou, 1991). Thus, higher-latitude breeding occupancy should cause

larger fluctuations in population size. Finally, seasonal migration is likely to cause enhanced

ecological (and in this case also geographic) release (i.e., in the occupancy of new niche space) in

relation to resident relatives. Thus, as migratory lineages' mobility enabled them to take advantage of

seasonal resource blooms in ecosystems at higher latitudes and expand their breeding ranges, their

populations should have grown proportionally more and at a faster rate than sedentary lineages.

Here we test these predictions by reconstructing effective population sizes of these thrush lineages

through time using whole-genome data and the pairwise sequential Markovian coalescent (PSMC; Li

& Durbin, 2011).

RESULTS

The focal variables of our hypotheses all lacked phylogenetic signal. Mean effective

population sizes and their variation were both higher among migratory lineages, as predicted (Table

1; $U = 6$ and 9, $p < 0.01$ and 0.05, respectively; one-tailed Mann-Whitney U-test). The degree of

early population growth was also proportionally higher among migrants, as predicted (Table 1; $U =$

9, $p < 0.05$; one-tailed Mann-Whitney U-test), but the rate of this early growth was not different

between the two groups (Table 1; $U = 21.5$, $p > 0.05$).

To examine why the degree of initial migrant population growth was higher but the rate of

growth was not, we considered that this might be because that growth extended over a longer period

of time. To our surprise, we found that among migrants these growth periods were remarkably long,

ranging from 1.67-4.26 My and averaging 2.97 My (Table 1, Fig. 1). It appeared that migratory

lineages did have a longer initial period of growth (migrants mean $deltaT = 2.97$ My vs. resident

mean $deltaT = 1.49$ My; Table 1). Unlike our other variables, initial growth period ($deltaT$) showed

significant phylogenetic signal ($p = 0.02$; Fig. S3), requiring PGLS analyses, which supported this

post-hoc hypothesis, both with actual branch lengths and with branch lengths set to 1.0 (Table S2).

We also used this approach to examine the relationship between $deltaT$ and breeding latitude (also

significant) and to confirm the relationship between migration and breeding latitude (Table S2).

In addition to testing our hypotheses, our results reveal other important attributes of these

songbirds. First, they appear to show few of the general population declines at the beginning of the

last glacial period (LGP, ~115 Kya) that Nadachowska-Brzyska et al. (2016) found among a global

sampling of birds (Figs. 2, S4). Secondly, and more importantly, there is a general lack of temporal

synchrony in historic population size changes (Figs. 2, S4), despite considerable geographic

proximity and overlap of ranges (especially during the nonbreeding season). Finally, a pattern widely

shared in the group is that most of the recent populations are below each lineage's historic peak, and

this trend predates the arrival of humans in the New World (Figs. 2, S4). A striking exception is *C.*

ustulatus swainsoni, which has had remarkable long-term growth (Fig. S4).

DISCUSSION

To our knowledge, this is the first study to contrast long-term effective population size

changes among closely related species to test *a priori* hypotheses about the population effects of a

major life-history trait; here we focused on seasonal migration. This perspective offers several new

insights into migrant demographics and evolutionary ecology. Migratory thrush lineages showed

higher effective population sizes, greater population size variation, and proportionally larger initial

population growth than resident relatives. A migratory life-history strategy thus imbues these

lineages with population size characteristics that are fundamentally different, on average, from those

of resident relatives.

Although migratory thrush lineages showed higher proportional initial growth than resident

relatives, this was not achieved through rapid adaptation to new environments as we predicted.

Population growth in migrants and residents was positive early in the lineages' histories (*deltaT*) in all

cases but one (*C. aurantiirostris*), and it occurred over fairly long periods of time (Table 1, Fig. 1).

That this process initially (on average) involves millions of years of population growth in this group,

apparently extended by a migratory life-history strategy that provided access to higher latitudes

(Tables 1, S2), is an insight from our study that bears examination from a broader taxonomic

perspective. While actual, census population sizes would likely have shown considerable variation

across this time, especially after the glacial cycles of the Pleistocene began ~2.6 Mya, the long-term

effective population sizes of migrants tended to show steady increases, achieving peaks on average

almost 3 My after the lineages' PSMC inception (Figs. 1, S4; Table 1).

With *deltaT* being both long and showing phylogenetic signal (Table 1, Fig. S3), one can

envision in this group a sort of ur-thrush trajectory in which unique, lineage-specific processes occur

as each evolves, influenced by ancestry, geography, ecological factors, and life-history strategy. These

long periods of early growth are concordant with the initial expansion or adaptation phases of taxon

cycles, a conceptual framework of lineage expansion and contraction over time through interactions

between biogeography and evolutionary ecology (Pepke et al., 2019; Ricklefs & Bermingham, 2002;

Ricklefs & Cox, 1972; Wilson, 1961).

Another noteworthy pattern widely shared in this group is that most recent populations are

below each lineage's historic peak (Figs. 2, S4). This, too, is reminiscent of taxon cycles (smaller

populations in later stages), and this trend both predates the arrival of humans in the New World

(Fig. S4) and reflects a broader pattern of historic avian declines (Germain et al. 2023b). Yet another

pattern is a general lack of temporal synchrony in historic population size changes, despite

considerable geographic proximity and overlap of ranges (Figs. 2, S4). In their review of taxon

cycles, Ricklefs & Bermingham (2002) inferred from such a lack of similarity among lineages that

each is in a unique population-environment relationship, responding idiosyncratically in a

coevolutionary relationship with factors such as parasites, predators, pathogens, and competitors.

Population structure

PSMC estimates of population size can be affected by population structure: a change in

structure can produce the signal of a change in size (Mazet et al., 2015, 2016; Wakeley, 1999). Some

of the population size changes we observed might therefore reflect changes only in levels of

population connectivity. There are well-understood differences between seasonal migrants and

residents in population structure. Migrants generally have lower levels of geographic partitioning of

both phenotypic (subspecies) and genetic variation (Belliure et al., 2000; Delmore et al., 2020; Mayr,

1963; Montgomery, 1896). The resident lineages we studied are more prone to this structure than

the migrant lineages, with migrants having an average of 2.5 subspecies per lineage and residents

having an average of 5.9 (Collar, 2005). Migratory lineages therefore come closer to meeting the

assumption of PSMC analyses that populations are panmictic. Thus, in addition to our geographic

scenario differing between the migrant and resident lineages we studied, we also have a pervasive

effect of population structure affecting our results, likely increasingly complementing geographic

expansion among seasonal migrants as lineage ages approach the recent.

Accounting for population structure in these models to distinguish between the two

phenomena is a difficult computational problem, and the correlation is not simple, and it might not

be tractable with data from a single individual (Mazet et al., 2016; Nadachowska-Brzyska et al., 2016,

2022). However, each individual has a genome amalgamated from the entire lineage metapopulation,

including individuals and populations that have not been sampled (Mazet et al., 2015). When not

focusing on relatively recent or rapid changes (e.g., in human or other bottlenecked systems, where

this has been most studied), these issues are probably not as important (Mazet et al., 2016). For

example, sister lineages usually show identical effective population size attributes at their origins,

reflecting their shared histories (Delmore et al., 2020; Li & Durbin, 2011; Nadachowska-Brzyska et

al., 2016). Inasmuch as many of our most interesting results are in deep time, where we expect

current individuals of a lineage to have the most thorough mixing or amalgamation of lineage-

specific metapolulation history, the effects of population structure are likely to become less

important and the results reflect (as we have necessarily interpreted) more of a lineage-wide

phenomenon than a variably structured, demically biased one. However, it is clear that the latter bias

becomes stronger as lineages progress toward the present, and for more recent times population

structure will be an important aspect of comparing effective population size estimates and their

changes between migrant and resident lineages (in this respect we were conservative in ignoring the

most recent 50 Kyr).

At present, there is no way to fully disentangle the effects of population structure and

geographic space on our results, but we make two observations leading us to infer that general

differences in population structure are not the main factor driving our results. First, structure will

have smaller effects on the PSMC signal in the deep past than in more recent history. Second,

climatic changes are known to cause coordinated shifts among codistributed taxa, e.g., in promoting

the partitioning of genetic variation among glacial refugia (Hewitt, 2000). A lack of coordinated

shifts in our PSMC results suggests that climate-driven shifts in population structure are not a major

feature (Fig. S4).

Geography and evolutionary ecology

Early population growth among migrant lineages is likely affected by some degree of

ecological release as these birds engage with the novel environments their increased movements

expose them to and occupy new niche space. However, in this system we cannot readily decouple

the expanded geographic and ecological opportunities that seasonal migration provides. The

relationship between avian range size and migration is complex, but correlation with latitude is

notably strong in North America (Pegan & Winger, 2020). Two aspects of our results suggest that

geography operating alone is unlikely to explain the larger migrant population sizes observed (Table

1, Fig. S4). The first is that early population growth is positive in all but one of the lineages we

studied; despite a probable lack of marked geographic expansion among residents, they, too,

experienced long-term growth. The second is that these initial growth periods are quite long,

extending over much longer periods than we should expect populations to respond to opportunities

for geographic expansion.

PSMC analyses excel at long time frames, but initial growth in residents and migrants

suggests a strong role for ecological factors. Global climate reconstruction data do not suggest that

there were similar long, steady trends during these time periods to match these thrush population

size changes (Lisiecki & Raymo, 2005; Fig. S4). Perhaps geographic space does make a difference, on

average, between the two life-history strategies. That these growth periods are long also suggest the

importance of ecological factors, though. For example, the boreal and subarctic woodland breeding

habitats of most *C. ustulatus swainsoni* and *C. minimus* expanded dramatically across North America as

glaciers receded since the last glacial maximum (LGM), a period of < 20 Kyr (Pielou, 1991). A

spatial response of such rapidity, less than 1% of the time of the initial historic growth phases of

these two lineages (> 3 Myr; Table 1), leads us to infer that breeding range size alone is not a

satisfactory explanation for our results. This reflects census size, and not effective population size

(N_e), but it implies that a migratory *Catharus* lineage's matching (or mapping) of N_e to average long-

term available geographic space is not likely to take millions of years. Or, to put it another way, their

comparatively recent rapid expansion into new space suggests a sort of preadaptation that is not

apparent early in the lineages' histories. Together, these aspects of our results suggest that while

geographic factors are involved, ecological factors are also important. Teasing these factors apart

might also explain the recent finding of Germain et al. (2023a) that migrants in general have had

larger populations than residents during the last 0.67 my.

246     Identifying key ecological mechanisms affecting population sizes among migrants is difficult,

and they will likely vary among lineages. Biotic interactions in general show greater effects on range

limits at the warmer than the cooler edges (Paquette & Hargreaves, 2021). Long-distance migrants

especially, in their extensive latitudinal movements, would receive a range expansion advantage in

this respect regardless of continental landmass shape. Many ecological factors affect the population

sizes of migrants versus residents, but the relationships between seasonal migration and predators,

parasites, diseases, and competitors are extremely complex and key aspects remain poorly

understood (Altizer et al., 2011; Newton, 2008).

254     Seasonal migration can be integrally related to predator avoidance, but among birds its

effects are not understood in the full, circannual context relative to resident relatives and they are

not generalizable (Brönmark et al., 2008; McKinnon et al., 2010; Newton, 2008; Rappole, 2013).

Exposure to parasites and pathogens varies both spatially and seasonally. Although breeding range

diminishment of these exposures can be important in some migrants (probably not among all

lineages), full annual exposure to both disease and parasite diversity is likely higher, resulting in at

least some lineages showing larger immune defense organs (Bennett & Fallis, 1960; Figuerola &

Green, 2000; Jenkins et al., 2011; Møller & Erritzoe, 1998; Piersma, 1997; Ricklefs et al., 2017).

Exposure to higher circannual parasite and pathogen diversity might not be correlated with

susceptibility, however; migration creates a geographic break that can lower transmission, causing

migrants overall to be less vulnerable to population declines (Hall et al., 2014).

265     Presently *C. minimus* shows an increased breeding and migratory prevalence and diversity of

avian malaria, a blood parasite, but aspects of this relationship probably changed over the time scales

involved (Pulgarín-R et al., 2019; Ricklefs et al., 2014). Avian migrants are more likely to be affected

by blood parasites in ecological rather than evolutionary time, although the latter are also operating

(Hellgren et al., 2007; Ricklefs et al., 2014). At the deeper times of avian families, seasonal migration

is an important factor affecting blood parasite-host coevolution by breaking down the congruence

of phylogenetic relationships between hosts and parasites, and there are also important differences

among biting-fly vectors and their transmission of avian blood parasite hosts (Jenkins et al., 2011;

Ricklefs et al., 2014). These issues suggest that reconstructing associations and their effects in deep

time will be difficult.

Interspecific competition has long been studied between migrants and residents, but its

potential role in affecting relative population sizes between the two groups is largely unknown

(Rappole, 2013). As close relatives, these thrushes are likely to be each others' closest competitors,

so it is conceivable that the increased relative abundance of migrants might depress realized

population sizes among residents. In this group, geographic ranges overlap broadly during the

nonbreeding season, but the tropical residents tend to occupy higher elevations than nonbreeding

migrants (Collar, 2005). Both interspecific competition and inland, montane, and mature-forest

retreats among older taxa are major factors in taxon cycles (Ricklefs & Bermingham, 2002; Ricklefs

& Cox, 1972; Wilson, 1961) and might also be operating in this system.

Conclusions

High-quality, genome-scale data offer important new insights into the effects of the major

life-history trait of seasonal migration on long-term effective population sizes. Our results indicate

larger and more variable migrant populations relative to residents, and these populations grow in a

way suggesting an important role not only for seasonal migration and geography, but also for

ancestry and evolutionary ecology. The differences we found are between group averages, and

variation among lineages includes residents that are migrant-like (*C. occidentalis*; Table 1, Figs. 1, 2, S4)

and vice-versa (e.g., *C. u. ustulatus*; Table 1, Fig. S4). Understanding why residents like *C. occidentalis*

show migrant-like historic population characteristics will help us understand the mechanisms

affecting these lineages. Such reasons could be as simple as realized range expansion (i.e., geographic

and ecological opportunity), and this hypothesis appears to fit the deeply split eastern-western sister

lineages within species in which smaller western ranges have smaller effective population sizes (*C.*

guttatus and *C. ustulatus*; Fig. S4). Or the reasons might be as complex as a lineage dropping out of

migration to become sedentary (*C. occidentalis* is sister to the migrant *C. guttatus*; Fig. S3).

What other patterns will emerge across geographic, temporal, and phylogenetic space?

Further sampling among other closely related groups containing mixed lineages of migrants and

residents are needed, as are complementary approaches to reconstructing demographic history, using

more individuals per lineage and other population genomic methods applicable to that sampling

(e.g., Beichman et al., 2017; Delmore et al., 2020; Marchi et al., 2021). Such studies will also help

determine the relative contributions of seasonal migration and its range and ecological expansion

components in the context of what appears to be a normal and phylogenetically influenced tendency

for early lineage growth (Figs. 1, S4, Table 1).

MATERIALS AND METHODS

The thrush lineages in our study are from the genus *Catharus* and its sister, the single species

in the genus *Hylocichla*. These comprise Nearctic-Neotropic seasonal migrants and Neotropical

residents. Among the latter, some lineages (*C. aurantiirostris, occidentalis, frantzii,* and *mexicanus*) have

some populations that exhibit some short-distance migration—likely partial migration or hard-

weather movements, either in elevation or with extreme northern populations moving south (Collar,

2005). These lineages thus comprise two groups, which for simplicity we will refer to as 'seasonal

migrants' and 'residents': seasonal migrants with extensive latitudinal movements, and residents that

are either wholly sedentary or with some individuals moving rather limited distances within the

Neotropics. The eight migrant lineages are: *Hylocichla mustelina*, *Catharus fuscescens*, *C. guttatus* E

(eastern lineage), *C. guttatus* W (western lineage), *C. minimus*, *C. ustulatus swainsoni*, *C. u. ustulatus*, and

C. bicknelli. (Note that four of these lineages represent two deeply divergent lineages each within

what are currently considered two biological species, *C. guttatus* and *C. ustulatus*; Topp et al., 2013).

The six resident lineages are: *Catharus aurantiirostris*, *C. fuscater*, *C. frantzii*, *C. gracilirostris*, *C. mexicanus*,

and *C. occidentalis*. Over the full evolutionary history of this *Hylocichla-Catharus* clade, changes among

lineages occurred in the trait of long-distance migration (Outlaw et al., 2003; Voelker et al., 2013;

Winker and Pruett, 2006). Our study does not assume that this trait was constant within each lineage

after the speciation events when these switches occurred. Instead, we consider that for a major trait

like long-distance migration, losses and gains at the full lineage level are likely to be infrequent, and

that current evidence of trait distribution on a phylogeny provides a powerful comparative

framework in which to elucidate the effects of that trait on other lineage attributes.

DNA was extracted from high-quality voucher specimens or blood samples for 14 thrush

lineages (Table S1). Whole-genome sequencing libraries were constructed using Nextera (Illumina)

DNA Flex Library Prep kits, and libraries were sequenced on a NovaSeq S4 (Illumina, San Diego,

CA) using paired-end 150 bp reads. Reads were aligned to the *C. ustulatus* genome (inland subspecies,

Reference bCatUst1, NCBI **PRJNA613294**) using bwa (mem algorithm with default settings, Li and

Durbin 2009). Resulting .sam files were converted to .bam format with samtools (Li et al., 2009);

picardtools was used to clean, sort and add read groups to .bam files

(http://broadinstitute.github.io/picard/). Between 97 and 98% of reads mapped to the reference for

an average read depth of 23 bp (range 15.9-29.8).

We analyzed whole-genomic data from the .bam files using the pairwise sequential

Markovian coalescent (PSMC) approach in the software package psmc (Li & Durbin, 2011). This

analysis uses pairwise (allelic) estimates of coalescent times for each locus across the genome,

developing a distribution of times to most recent common ancestor (TMRCAs), and the frequency

of these coalescent events is inversely proportional to effective population size (Li & Durbin, 2011;

Mather et al., 2020). We used a generation time of 2.5 yr, taken as an average among many of these

species (Dellinger et al., 2020; Evans et al., 2020; Mack & Yong, 2020; Saether et al., 2005;

Townsend et al., 2020; Whitaker et al., 2020), and a mutation rate of 2.3×10^{-9} mutations per site

per year (Smeds et al., 2016). Single individuals represent lineage characteristics using this approach

(Li & Durbin, 2011; Mazet et al., 2015; Nadachowska-Brzyska et al., 2015). Importantly for our focal

questions, variations in these parameters do not affect the shape of the curve of population sizes

(N_e) through time, but rather the values of the time and N_e estimates (Nadachowska-Brzyska et al.

2015). Being closely related and of similar size, we expect these taxa to have very similar generation

times and mutation rates (e.g., Bird et al., 2020; Healy et al., 2014).

The PSMC method is not foolproof; for example, aspects of selection, inbreeding, and

isolation can affect population size estimates (Mather et al., 2020). No method of genomic

population demographic reconstruction is known to be consistently and reliably accurate (Marchi et

al., 2021), but uniform application of a powerful method like PSMC to genome-scale data in a

closely related group of lineages can provide new insights into the evolutionary effects of major life-

history strategies.

For all variables, PSMC output for times more recent than 50 Kya was ignored, because

recent estimates of effective population size are less reliable (Li & Durbin, 2011; Nadachowska-

Brzyska et al., 2016). We consider this to be a conservative cutoff date. Our two focal variables,

mean effective population size (N_e) and the coefficient of variation (SD/mean), were obtained from

standard PSMC output. Our examination of early population growth involved deriving two

additional variables from the PSMC output: degree of initial growth (1 - (N_{trough}/N_{peak})), and rate of

growth (degree/time period over which initial growth occurred; the latter we term *deltaT*; Fig. S2).

We note that estimates of the dates of divergence events in *Catharus* evolution vary by study, genetic

marker, methodology, and parameter estimates such as generation time and mutation rates (Outlaw

et al., 2003; Topp et al., 2013; Voelker et al., 2013; Winker and Pruett, 2006). Date estimates in our

study that differ from previous work reflect these methodological issues, and here it is the relative

values among lineages that are most important.

We tested the assumption of phylogenetic independence of continuous characters obtained

from the PSMC analyses using a phylogeny reconstructed using maximum likelihood from 1.5 Mb

of autosomal sequence from chromosome 22 in MEGA X (Kumar et al. 2018), which produced the

expected topology given prior work (e.g., Everson et al. 2019), and, for phylogenetic independence,

the Abouheif-Moran approach (Abouheif, 1999; Münkenmüller et al., 2012) as implemented in the R

package adephylo (Jombart et al., 2010). For those variables not showing phylogenetic signal, we

tested our hypotheses using a one-tailed Mann-Whitney *U*-test. This nonparametric test has the

advantage of not requiring that particular assumptions are met (e.g., homogeneity of variance), nor

precise measurements (thus accommodating uncertainty in our N_e estimates; Sokal & Rohlf, 1995;

Whitlock & Schluter, 2015).

Phylogenetic generalized least squares (PGLS) was used in a post-hoc analyses to test for

differences in the single variable (*deltaT*) that showed phylogenetic signal and in followup analyses

bringing in breeding latitude directly. PGLS can accommodate tests for relationships with discrete

traits (here migratory-sedentary) and is basically a linear regression weighted in this case by *detaT*'s

phylogenetic relationships (Freckleton, 2009; Symonds & Blomberg, 2014). Because this genus has

experienced considerable interspecific gene flow (Everson et al. 2019), the accuracy of branch

lengths is uncertain, so we ran analyses both using actual branch lengths and setting branch lengths

to 1.0. We also extended our analysis of *deltaT* to examine its relationship to breeding latitude

directly (using the same approach). Midpoints of breeding range latitudes were derived using

ArcMap (ESRI 2021, ver.10.8.2) and range maps from BirdLife International and the Handbook of

the Birds of the World (2021). PGLS analyses were done using the R packages phytools (ver. 0.7-80,

Revell 2012) and nmle (ver. 3.1-152, Pinheiro et al., 2013) and using the corPagel model approach.

REFERENCES

Abouheif E. 1999. A method for testing the assumption of phylogenetic independence in
comparative data. *Evolutionary Ecology Research* **1**:895-909.

Altizer S, Bartel R, Han BA. 2011. Animal migration and infectious disease risk. *Science* **331**:296-302.

Beichman AC, Phung TN, Lohmueller KE. 2017. Comparison of single genome and allele
frequency data reveals discordant demographic histories. *Genes Genomes Genetics* **7**:3605-3620.

Belliure J, Sorci G, Møller AP, Clobert J. 2000. Dispersal distances predict subspecies richness in
birds. *Journal of Evolutionary Biology* **13**:480–487.

Bennett GM, Fallis AM. 1960. Blood parasites of birds in Algonquin Park, Canada, and a discussion
of their transmission. *Can. J. Zool.* **38**:261-273.

Bird JP, Martin R, Akçakaya HR, Gilroy J, Burfield IJ, Garnett ST, Symes A, Taylor J, Şekercioğlu
ÇH, Butchart SHM. 2020. Generation lengths of the world's birds and their implications for
extinction risk. *Conservation Biology* **34**:1252-1261.

BirdLife International and the Handbook of the Birds of the World. 2021. Bird species distribution
maps of the world. Version 2021.1. Available at http://datazone.birdlife.org/species/requestdis.

Brönmark C, Skov C, Brodersen J, Nilsson PA, Hansson L-A. 2008. Seasonal migration determined
by a trade-off between predator avoidance and growth. *PLoS ONE* **3**:e1957.
doi:10.1371/journal.pone.0001957

Collar N J. 2005. Family Turdidae (thrushes). Pp. 514-807 *in Handbook of the Birds of the World*,
Volume 10 (J. del Hoyo et al., eds.). Lynx Edicions, Barcelona.

Dellinger R, Wood PB, Jones PW, Donovan TM. 2020. Hermit Thrush (*Catharus guttatus*), version
1.0. In Birds of the World (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
https://doi.org/10.2173/bow.herthr.01

Delmore K, Winker K. 2022. *Catharus* genomes project; NCBI-SRA; Version (pending); doi:
pending.

Delmore K, Illera JC, Pérez-Tris J, Segelbacher G, Ramos JSL, Durieux G, Ishigohoka J, Liedvogel
M. 2020. The evolutionary history and genomics of European blackcap migration. *eLife* **9**:e54462.

Evans M, Gow E, Roth RR, Johnson MS, Underwood TJ. 2020.. Wood Thrush (*Hylocichla mustelina*),
version 1.0. In Birds of the World (A. F. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY,
USA. https://doi.org/10.2173/bow.woothr.01

ESRI. 2021. ArcGIS Desktop: Release 10.8.2. Redlands, CA: Environmental Systems Research
Institute.

Figuerola J, Green AJ. 2000. Hematozoan parasites and migratory behavior in waterfowl. *Evolutionary*
Ecology **14**:143-153.

Freckleton RP. 2009. The seven deadly sins of comparative analysis. *Journal of Evolutionary Biology*
22:1367-1375.

Freckleton RP, Harvey PH, Pagel M. 2002. Phylogenetic analysis and comparative data: a test and
review of evidence. *American Naturalist* **160**:712-26. doi: 10.1086/343873

Germain RR, Shaohong F, Buffan L, Carmona CP, Chen G, Graves GR, Tobias JA, Rahbek C, et al.
2023a. Changes in the functional diversity of modern bird species over the last million years.
Proceedings of the National Academy of Sciences USA **120**:e2201945119.
https://doi.org/10.1073/pnas.2201945119

Germain RR, Shaohong F, Buffan L, Carmona CP, Chen G, Graves GR, Tobias JA, Rahbek C, et al.
2023b. Species-specific traits mediate avian demographic responses under past climate change.
Nature Ecology and Evolution **7**:862-872.

Greenberg R. 1980. Demographic aspects of long-distance migration. Pp. 493-503 *in* A. Keast and
E. S. Morton, eds. *Migrant Birds in the Neotropics*. Smithsonian Institution Press, Washington, D.C.

Hall RJ, Altizer S, Bartel RA. 2014. Greater migratory propensity in hosts lowers pathogen
transmission and impacts. *Journal of Animal Ecology* **83**:1068-1077. doi:10.1111/1365-2656.12204

Healy K, Guillerme T, Finlay S, Kane A, Kelly SBA, McClean D, Kelly DJ, Donohue I, Jackson A

L, Cooper N. 2014. Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proceedings of the Royal Society B* **281**:20140298.

Hellgren O, Waldenström J, Peréz-Tris J, Ösi ES, Hasselquist D, Krizanauskiene A, Ottosson U, Bensch S. 2007. Detecting shifts of transmission areas in avian blood parasites—a phylogenetic approach. *Molecular Ecology* **16**:1281-1290.

Hewitt GM. 2000. The genetic legacy of the Quaternary. *Nature* **405**:907-913.

Jenkins T, Thomas GH, Hellgren O, Owens IPF. 2011. Migratory behavior of birds affects their coevolutionary relationship with blood parasites. *Evolution* **66**:740-751.

Jombart T, Balloux F, Dray S. 2010. *adephylo*: new tools for investigating the phylogenetic signal in biological traits. *Bioinformatics* **26**:1907-1909.
 https://doi.org/10.1093/bioinformatics/btq292

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* **35**:1547-1549.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**:1754–1760.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature* **475**:493-496.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078-2079.

Lisiecki LE, Raymo ME. 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records. *Paleoceanography* **20**: PA1003. doi:10.1029/2004PA001071.

Marchi N, Schlichta F, Excoffier L. 2021. Demographic inference. *Current Biology* **31**:R267-R281.

Mack DE, Yong W. 2020. Swainson's Thrush (*Catharus ustulatus*), version 1.0. In Birds of the World (A. F. Poole and F. B. Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.swathr.01

Mather N, Traves SM, Ho SYW. 2020. A practical introduction to sequentially Markovian coalescent methods for estimating demographic history from genomic data. *Ecology and Evolution* **10**:579-589.

Mayr E. 1963. *Animal Species and Evolution.* Belknap Press, Harvard University, Cambridge, Massachusetts.

Mazet O, Rodriguez W, Chikhi L. 2015. Demographic inference using genetic data from a single individual: Separating population size variation from population structure. *Theoretical Population Biology*

104:46–58. http://dx.doi.org/10.1016/j.tpb.2015.06.003

Mazet O, Rodriguez W, Grusea S, Boitard S, Chikhi L. 2016. On the importance of being structured: instantaneous coalescence rates and human evolution—lessons for ancestral population size inference. *Heredity* **116**:362-371. doi:10.1038/hdy.2015.104

McKinnon L, Smith PA, Nol E, Martin JL, Doyle FI, Abraham KF, Gilchrist HG, Morrison RIG, Bêty J. 2010. Lower predation risk for migratory birds at high latitudes. *Science* **327**:326-327. https://doi.org/10.1126/science.1183010

Møller AP, Erritzoe J. 1998. Host immune defense and migration in birds. *Evolutionary Ecology* **12**: 945-953.

Montgomery TH Jr. 1896. Extensive migration in birds as a check upon the production of geographical varieties. *American Naturalist* **30**:458–464.

Münkenmüller T, Lavergne S, Bzeznik B, Dray S, Jombart T, Schiffers K, Thuiller W. 2012. How to measure and test phylogenetic signal. *Methods in Ecology and Evolution* **3**:743-756.

Nadachowska-Brzyska K, Li C, Smeds L, Zhang G, Ellegren H. 2015. Temporal dynamics of avian populations during Pleistocene revealed by whole-genome sequences. *Current Biology* **25**:1375-1380.

Nadachowska-Brzyska K, Burri R, Smeds L, Ellegren H. 2016. PSMC analysis of effective population sizes in molecular ecology and its application to black-and-white *Ficedula* flycatchers. *Molecular Ecology* **25**:1058-1072.

Nadachowska-Brzyska K, Konczal M, Babik W. 2022. Navigating the temporal continuum of effective population size. *Methods in Ecology and Evolution* **13**:22-41.

Newton I. 2008. *The Migration Ecology of Birds*. Elsevier, Amsterdam.

Outlaw DC, Voelker G, Mila B, Girman DJ. 2003. Evolution of long-distance migration in and historical biogeography of Catharus thrushes: A molecular phylogenetic approach. *Auk* **120**:299-310.

Paquette A, Hargreaves AL. 2021. Biotic interactions are more often important at species' warm versus cool range edges. *Ecology Letters* **24**:2427-2438.

Pegan TM, Winger BM. 2020. The influence of seasonal migration on range size in temperate North American passerines. *Ecography* **43**:1191-1202.

Pepke ML, Irstedt M, Fjeldså J, Rahbek C, Jønsson KA. 2019. Reconciling supertramps, great speciators and relict species within the taxon cycle stages of a large island radiation (Aves: Campephagidae). *Journal of Biogeography* **46**:1214-1225.

Pérez-Tris J, Bensch S, Carbonell R, Helbig AJ, Tellería JL. 2004. Historical diversification of migration patterns in a passerine bird. *Evolution* **58**:1819-1832.

Pielou EC. 1991. *After the Ice Age: The return of Life to Glaciated North America*. University of Chicago Press, Chicago.

Piersma T. 1997. Do global patterns of habitat use and migration strategies co-evolve with relative investments in immunocompetence due to spatial variation in parasite pressure? *Oikos* **80**:623–631. https://doi.org/10.2307/3546640

Pinheiro J, Bates D, DebRoy S, Sarkar D, the R Development Core Team. 2013. nlme: Linear and nonlinear mixed effects models. R package version 3.1-108.

Pulgarín-R PC, Gómez C, Bayly NJ, Bensch S, FitzGerald AM, Starkloff N, Kirchman JJ, González-Prieto AM, Hobson KA, Ungvari-Martin J, Skeen H, Castaño MI, Cadena CD. 2019. Migratory birds as vehicles for parasite dispersal? Infection by avian haemosporidians over the year and throughout the range of a long-distance migrant. *Journal of Biogeography* 46:83–96. https://doi.org/10.1111/jbi.13453

Rappole JH. 2013. *The Avian Migrant: The Biology of Bird Migration*. Columbia University Press, New York.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**:217-223.

Ricklefs RE, Cox GW. 1972. Taxon cycles in the West Indian avifauna. *American Naturalist* **106**:195-219.

Ricklefs RE, Bermingham E. 2002. The concept of the taxon cycle in biogeography. *Global Ecology and Biogeography* **11**:353-361.

Ricklefs RE, Outlaw DC, Svensson-Coelho M, Medeiros MC, Ellis VA, Latta S. 2014. Species formation by host shifting in avian malaria parasites. *Proceedings of the National Academy of Sciences USA* **111**:14816-14821.

Ricklefs RE, Medeiros M, Ellis VA, Svensson-Coelho M, Blake JG, Loiselle BA, Soares L, Fecchio A, Outlaw D, Marra PP, Latta SC, Valkiūnas G, Hellgren O, Bensch S. 2017. Avian migration and the distribution of malaria parasites in New World passerine birds. *Journal of Biogeography* **44**:1113-1123.

Ruegg K, Anderson EC, Boone J, Pouls J, Smith TB. 2014. A role for migration linked genes and genomic islands in divergence of a songbird. *Molecular Ecology* **23**:4757–4769. https://doi.org/10.1111/mec.12842.

Saether B-E, Lande R, Engen S, Weimerskirsch H, Lillegård M, Altwegg R, Becker PH, Bregnballe J, et al. 2005. Generation time and temporal scaling of bird population dynamics. *Nature* **436**:99-102.

Sandercock BK, Jaramillo A. 2002. Annual survival rates of wintering sparrows: Assessing

603   demographic consequences of migration. *Auk* **119**:149-165.
604
605   Smeds L, Qvarnström A, Ellegren H. 2016. Direct estimate of the rate of germline mutation in a
606   bird. *Genome Research* **26**:1211-1218.
607
608   Sokal RR, Rohlf FJ. 1995. *Biometry: The Principles and Practice of Statistics in Biological Research*, 3rd ed. W.
609   H. Freeman and Company, New York.
610
611   Symonds MRE, Blomberg SP. 2014. A primer on phylogenetic generalised least squares. Pp. 105-130
612   in *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology* (L. Z. Garamszegi,
613   ed.). Springer-Verlag, Berlin.
614
615   Termignoni-Garcia F, Kirchman JJ, Clark J, Edwards SV. 2022. Comparative population genomics
616   of cryptic speciation and adaptive divergence in Bicknell's and Gray-cheeked thrushes (Aves:
617   *Catharus bicknelli* and *Catharus minimus*). *Genome Biol Evol.* **14**:evab255. doi: 10.1093/gbe/evab255.
618
619   Topp CM, Pruett CL, McCracken KG, Winker K. 2013. How thrushes conquered North America:
620   A comparative phylogeography approach. *PeerJ* 1:e206. http://dx.doi.org/10.7717/peerj.206
621
622   Townsend JM, McFarland KP, Rimmer CC, Ellison WG, Goetz JE. 2020. Bicknell's Thrush
623   (*Catharus bicknelli*), version 1.0. In Birds of the World (P. G. Rodewald, Editor). Cornell Lab of
624   Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.bicthr.01
625
626   Voelker G, Bowie RC, Klicka J. 2013. Gene trees, species trees and Earth history combine to shed
627   light on the evolution of migration in a model avian system. *Molecular Ecology* **22**:3333–3344.
628   https://doi.org/10.1111/mec.12305.
629
630   Wakeley J. 1999. Nonequilibrium migration in human history. *Genetics* **153**:1863-1871.
631
632   Whitaker DM, Warkentin IG, McDermott JPB, Lowther PE, Rimmer CC, Kessel B, Johnson SL,
633   Ellison WG. 2020. Gray-cheeked Thrush (*Catharus minimus*), version 1.0. In Birds of the World (P.
634   G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
635   https://doi.org/10.2173/bow.gycthr.01
636
637   Whitlock MC, Schluter D. 2015. *The Analysis of Biological Data*, 2nd ed. Roberts and Company
638   Publishers, Greenwood Village, Colorado.
639
640   Wilson EO. 1961. The nature of the taxon cycle in the Melanesian any fauna. *American Naturalist*
641   **95**:169-193.
642
643   Winker K. 2000. Migration and speciation. *Nature* **404**:36.
644
645   Winker K. 2010. On the origin of species through heteropatric differentiation: A review and a model
646   of speciation in migratory animals. *Ornithological Monographs* **69**:1-30.
647
648   Winker K, Pruett CL. 2006. Seasonal migration, speciation, and morphological convergence in the

649 *Catharus* thrushes (Aves: Turdidae). *Auk* **123**:1052-1068.
650

Tables and Figures

Table 1. Data from pairwise sequential Markovian coalescent (PSMC) analyses reflecting effective population sizes (N_e) through history at depths > 50 Kyr and the five variables derived and analyzed from that output. Taxa shaded in gray are Neotropical residents.

Figure 1. Partial historic effective population size curves from all lineages in this study, based on pairwise sequential Markovian coalescent (PSMC) analyses. Each peak of initial growth is set to zero years to set a common framework in which to visualize the periods and magnitudes of initial growth among migrant (red) and resident (blue) lineages. See specific lineages with their bootstrapped results in Fig. S2.

Figure 2. Historic effective population size curves from all lineages, based on pairwise sequential Markovian coalescent (PSMC) analyses. Migrant lineages are in red and resident lineages in blue. See specific lineages with their bootstrapped results in Fig. S2.

Supplementary Information

Table S1. Specimen data and NCBI-SRA numbers. Vouchered specimens are housed in the following institutions: UAM (University of Alaska Museum), MSB (Museum of Southwestern Biology, University of New Mexico), LSUMNS (Louisiana State University Museum of Natural Science), and FMNH (Field Museum of Natural History).

Table S2. Results of phylogenetic generalized least squares (PGLS) regressions using corPagel and associated lambda values (see Revell 2012).

Figure S1. Distribution maps of the thrush taxa in this study. Among seasonal migrants, green indicates breeding range and yellow is wintering range. Among sedentary lineages (those shaded in gray), purple indicates year-round range. Data are from BirdLife International and the Handbook of the Birds of the World (2021), Bird species distribution maps of the world. Version 2021.1. Available at http://datazone.birdlife.org/species/requestdis.

Figure S2. Graphic presentation of the *Catharus minimus* PSMC dataset, showing effective population size (N_e) from 50 kyr back in time to the lineages' origins as estimated from genomic data. The variables in our analyses are the mean and SD of the effective population size (N_e) values, *deltaT* of initial growth (time$_{trough}$ – time$_{peak}$), the degree of that growth ($1 - [N_{trough}/N_{peak}]$), and the rate of that growth (degree/*deltaT*).

Figure S3. The phylogenetic tree of *Catharus* and outgroup *Hylocichla* with the population growth metric *deltaT* mapped onto it. Resident taxa are highlighted in gray. Positive centered and scaled *deltaT* values are in black and negative values are in white, and their size is proportional to the absolute value. This trait has significant phylogenetic signal (see Jombart et al. 2012) and therefore comprises attributes both of migratory/sedentary lineages and of their deeper evolutionary history.

Figure S4. Montage of the historic effective population size curves of all lineages analyzed in this

697   study, with each lineage in a separate panel (based on pairwise sequential Markovian coalescent, or
698   PSMC, analyses). Sedentary lineages are highlighted in gray. Note that scales on both axes vary
699   among panels, and that the X axis is on a log scale. Bold red lines are the main curves from the
700   original data, and pink lines reflect 100 replicates from bootstrapped sequences. Bold red curves are
701   all overlaid on common axes in the single panel of Fig. 1.
702

703 Table 1. Data from pairwise sequential Markovian coalescent (PSMC) analyses reflecting effective
704 population sizes (N_e) through history at depths > 50 Kyr and the five variables derived and analyzed
705 from that output. Taxa shaded in gray are Neotropical residents.
706

Taxon	Mean N_e (± SD)	SD/mean	Degree of early growth 1 - (N_{trough}/N_{peak})	Rate of early growth degree/deltaT	deltaT
Hylocichla mustelina	35.63 (± 20.11)	0.56	0.86	2.92E-07	2,946,862
Catharus fuscescens	96.22 (± 56.07)	0.58	0.83	1.94E-07	4,256,514
C. guttatus E	83.12 (± 84.11)	1.01	0.90	4.47E-07	2,023,341
C. guttatus W	39.72 (± 24.20)	0.61	0.81	3.28E-07	2,477,684
C. minimus	63.71 (± 28.83)	0.45	0.78	2.17E-07	3,594,914
C. ustulatus swainsoni	76.61 (± 66.61)	0.87	0.79	2.12E-07	3,740,764
C. ustulatus ustulatus	39.56 (± 12.15)	0.31	0.63	2.14E-07	2,943,763
C. bicknelli	46.90 (± 24.12)	0.51	0.78	2.79E-07	2,781,554
C. aurantiirostris	12.38 (± 2.83)	0.23	-0.75	-1.00E-06	747,451
C. fuscater	11.38 (± 1.66)	0.15	0.39	6.18E-07	627,449
C. frantzii	24.71 (± 8.41)	0.34	0.17	2.20E-07	774,024
C. gracilirostris	30.21 (± 10.05)	0.33	0.58	3.52E-07	1,639,266
C. mexicanus	37.10 (± 16.18)	0.44	0.75	3.34E-07	2,250,295
C. occidentalis	76.40 (± 53.14)	0.70	0.91	2.11E-07	4,292,405
Means (± SD)					
migrants	60.18 (± 39.52)**	0.61 (± 0.23)*	0.798 (± 0.08)*	2.73E-7 (± 0.85E-7)	3,095,674 (± 725,912)†
residents	32.03 (± 20.11)	0.36 (± 0.19)	0.340 (± 0.59)	1.23E-7 (± 5.69E-7)	1,721,815 (+ 1,409,920)

* $p < 0.05$; ** $p < 0.01$; † $p < 0.10$.

712
713

714
715
716 Figure 1. Partial historic effective population size curves from all lineages in this study, based on
717 pairwise sequential Markovian coalescent (PSMC) analyses. Each peak of initial growth is set to zero
718 years to set a common framework in which to visualize the periods and magnitudes of initial growth
719 among migrant (red) and resident (blue) lineages. See specific lineages with their bootstrapped
720 results in Fig. S2.
721

722
723
724 Figure 2. Historic effective population size curves from all lineages, based on pairwise sequential
725 Markovian coalescent (PSMC) analyses. Migrant lineages are in red and resident lineages in blue. See
726 specific lineages with their bootstrapped results in Fig. S2.

Table S1. Specimen data and NCBI-SRA numbers. Vouchered specimens are housed in the following institutions: UAM (University of Alaska Museum), MSB (Museum of Southwestern Biology, University of New Mexico), LSUMNS (Louisiana State University Museum of Natural Science), and FMNH (Field Museum of Natural History).

Institution	Catalog #	Species	Year	Field No.	Locality	NCBI-SRA
UAM	28620	*H. mustelina*	2010	KSW5403	Belize: Toledo District; Big Falls	pending
UAM	27774	*C. fuscescens*	2007	KSW5151	Belize: Toledo District; Big Falls	pending
UAM	15202	*C. guttatus* E	1992	KSW4013	USA: Vermont; Brandon	pending
UAM	26337	*C. guttatus* W	2008	UAMX5095	USA: Alaska; Kodiak	pending
UAM	22642	*C. minimus*	2003	KSW5000	USA: Alaska; Fairbanks	pending
n.a. (blood)	KF15K01	*C. ustulatus swainsoni*	2011	KF15K01	Canada: British Columbia, Kamloops	pending
n.a. (blood)	KF01K01	*C. u. ustulatus*	2011	KF01K01	Canada: British Columbia, Kamloops	pending
UAM	19996	*C. bicknelli*	2000	KSW3633	USA: Vermont; Mt Mansfield	pending
UAM	25341	*C. aurantiirostris*	2004	MJM1154	Panama: Chiriqui; El Salto	pending
MSB	31939	*C. fuscater*	2008	MSB31939	Peru: Amazonas; 4.5 km N Tullanya	pending
UAM	25098	*C. frantzii*	2004	KSW4485	Panama: Chiriqui; Volcan Baru	pending
LSUMNS	138784	*C. gracilirostris*	1990	JMB1065; B-16270	Costa Rica: San Jose; Cerro de la Muerte	pending
UAM	10352	*C. mexicanus*	1994	PEP2489	Mexico: Veracruz; Volcan San Martin	pending
FMNH	343305	*C. occidentalis*	1989	MEX408	Mexico: Oaxaca; Totontepec	pending

Table S2. Results of phylogenetic generalized least squares (PGLS) regressions using corPagel and associated lambda values (see Revell 2012).

variables	branch lengths	lambda	*p*-value
deltaT vs *migration*	actual	0.87	0.011
"	1.0	1.00	0.027
deltaT vs *latitude*	actual	1.00	0.049
"	1.0	1.00	0.019
latitude vs *migration*	actual	1.00	0.015
"	1.0	1.00	<0.001

Figure S1. Distribution maps of the thrush taxa in this study. Among seasonal migrants, green indicates breeding range and yellow is wintering range. Among sedentary lineages (those shaded in gray), purple indicates year-round range. Data are from BirdLife International and the Handbook of the Birds of the World (2021), Bird species distribution maps of the world. Version 2021.1. Available at http://datazone.birdlife.org/species/requestdis.

Figure S2. Graphic presentation of the *Catharus minimus* PSMC dataset, showing effective population size (N_e) from 50 kyr back in time to the lineages' origins as estimated from genomic data. The variables in our analyses are the mean and SD of the effective population size (N_e) values, *deltaT* of initial growth (time$_{trough}$ – time$_{peak}$), the degree of that growth (1 – [N$_{trough}$/N$_{peak}$]), and the rate of that growth (degree/*deltaT*).

Figure S3. The phylogenetic tree of *Catharus* and outgroup *Hylocichla* with the population growth metric *deltaT* mapped onto it. Resident taxa are highlighted in gray. Positive centered and scaled *deltaT* values are in black and negative values are in white, and their size is proportional to the absolute value. This trait has significant phylogenetic signal (see Jombart et al. 2012) and therefore comprises attributes both of migratory/sedentary lineages and of their deeper evolutionary history.

Figure S4. Montage of the historic effective population size curves of all lineages analyzed in this study, with each lineage in a separate panel (based on pairwise sequential Markovian coalescent, or PSMC, analyses). Sedentary lineages are highlighted in gray. Note that scales on both axes vary among panels, and that the X axis is on a log scale. Bold red lines are the main curves from the original data, and pink lines reflect 100 replicates from bootstrapped sequences. Bold red curves are all overlaid on common axes in the single panel of Fig. 1.

Hylocichla mustelina

Catharus ustulatus swainsoni

Catharus frantzii

Catharus fuscescens

Catharus ustulatus ustulatus

Catharus gracilirostris

Catharus guttatus W

Catharus bicknelli

Catharus mexicanus

Catharus guttatus E

Catharus aurantiirostris

Catharus occidentalis

Catharus minimus

Catharus fuscater

Figure S2. Graphic presentation of the *Catharus minimus* PSMC dataset, showing effective population size (N_e) from 50 kyr back in time to the lineages' origins as estimated from genomic data. The variables in our analyses are the mean and SD of the effective population size (N_e) values, *deltaT* of initial growth (time$_{trough}$ − time$_{peak}$), the degree of that growth ($1 − [N_{trough}/N_{peak}]$), and the rate of that growth (degree/*deltaT*).

C_mexicanus

C_aurantiirostris

C_fuscater

C_guttatus_W

C_guttatus_E

C_occidentalis

C_gracilirostris

C_frantzii

C_ustulatus_ustulatus

C_ustulatus_swainsoni

C_minimus

C_bicknelli

C_fuscescens

H_mustelina

-2   -1   1   2

deltaT