Multifunctionality of V-type ATPase during asexual growth and development of
Plasmodium falciparum

Neeta Shadija[1], Swati Dass[1], Wei Xu[1], Hangjun Ke[1]

[1] Center for Molecular Parasitology, Department of Microbiology and Immunology, Drexel
University College of Medicine, Philadelphia, Pennsylvania, USA, 19129

Correspondence, hk84@drexel.edu

Abstract

V-type ATPases are highly conserved hetero-multi-subunit proton pumping machineries found in
all eukaryotic organisms. They use ATP hydrolysis to pump protons, acidifying intracellular or
extracellular compartments, and are thus crucial for various biological processes. Despite being
evolutionarily conserved in malaria parasites, this proton pump remains understudied. To
understand the localization and biological function of V-type ATPase in the deadliest human
malaria parasite *Plasmodium falciparum*, we utilized CRISPR/Cas9 to endogenously tag the
subunit A of the V_1 domain at the C-terminus. V_1A (PF3D7_1311900) was tagged with a triple
hemagglutinin (3HA) epitope and TetR-DOZI-aptamers for conditional expression under the
regulation of anhydrotetracycline. Through immunofluorescence assays, we identified that V-
type ATPase was expressed throughout the intraerythrocytic developmental cycle and was
mainly localized on the digestive vacuole and plasma membrane. Immuno-electron microscopy
further revealed that V-type ATPase was also localized on secretory organelles, such as
rhoptries in merozoites. Knockdown of V_1A led to cytosolic pH imbalance and blockage of
hemoglobin digestion in the digestive vacuole, resulting in an arrest of parasite development in
the trophozoite stage and, ultimately, parasite demise. Using BN-PAGE/Western blot, we
detected a large molecular weight complex (~ 1.0 MDa) corresponding to the total molecular
weights of V_1 and V_o domains. The complex was readily disrupted by the V-type ATPase specific
inhibitor Bafilomycin A1, but not by low glucose conditions or treatment with chloroquine.
Together, our data suggest that V-type ATPase is localized on several subcellular compartments
in *P. falciparum* and plays critical roles to support malaria parasites to grow and replicate inside
red blood cells.

Introduction

Malaria remains one of the deadliest infectious diseases worldwide. In 2021, the WHO reported 247,000,000 malaria cases and 619, 000 deaths[1]. The causative agent of this disease is the protozoan parasite called *Plasmodium*, belonging to the phylum Apicomplexa. Among the five *Plasmodium* species infecting humans (*P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae*, *P. knowlesi*), *P. falciparum* is responsible for the majority of malaria cases and deaths worldwide. All clinical symptoms of malaria arise due to the repetitive growth of the parasites in red blood cells (RBCs); hence, the intraerythrocytic developmental cycle (IDC) or the asexual blood stage has long been recognized as a crucial target for developing antimalarial interventions.

Throughout the IDC, malaria parasites reside in a parasitophorous vacuole and contain many intracellular compartments, including the cytosol, the digestive vacuole (DV) for hemoglobin digestion, the nucleus/ER/Golgi apparatus, the mitochondrion, the apicoplast, and secretory organelles in merozoites. Some of these subcellular compartments are known to have distinct pH, which is key for sustaining the biological processes inside them. For instance, the parasite cytosol maintains a static pH near 7.3[2], whereas the DV requires a lower pH near 5.0 for hemoglobin digestion[3-6]. The mitochondrion has a slightly basic pH (7.37 ± 0.09) whereas the apicoplast is more neutral (7.12 ± 0.40)[7].

Regulation of pH in distinct subcellular compartments involves multiple players, including buffers, enzymes, transporters, and proton pumps[8]. Proton pumps are critical because they can directly transport protons against the concentration gradient, actively regulating subcellular pH[9]. In *P. falciparum*, two types of proton pumps are present[10], including the pyrophosphate (PPi) driven H^+-pumping pyrophosphatases (H^+-PPases) or vacuolar pyrophosphatases (V-PPases)[11-13], and the ATP driven vacuolar type ATPases (V-type ATPases)[14]. VP1 (vacuolar pyrophosphatase 1) is an example of PPi driven proton pumping pyrophosphatase. In *P. falciparum*, a recent study has shown that VP1 is mainly localized on the parasite plasma membrane, regulates cytosolic pH, and plays essential roles in the ring stage and the ring-to-trophozoite transition[15]. On the other hand, V-type ATPase is a large molecular weight ATP hydrolyzing proton pumping machinery[16], consisting of a V_1 domain containing subunits of A-H and a V_o domain containing subunits of a, c, c', c'', d, e. The cytosolic V_1 domain holds the ATP hydrolysis hexamer made of A and B subunits, whereas the membrane bound V_o domain contains the H^+ channel, which is involved in proton transport across the membrane. A unique feature of this machinery lies in the reversible assembly and disassembly of the V_1V_o domains[17], regulating the pump's function in response to various extracellular conditions[18]. The *P. falciparum* genome encodes all V_1 domain subunits and most V_o domain subunits that are present in other eukaryotes (www.PlasmoDB.org). The recent genome-wide mutagenesis study in *P. falciparum*[19] and the knockout survey in *P. berghei*[20] reveal that nearly all V-type ATPase subunits are essential in the asexual blood stage. Despite the significance, our understanding of the physiological role of V-type ATPase in malaria parasites remains rather limited.

In classical model organisms, including yeast, humans, and plants, V-Type ATPase is localized on the membranes of multiple subcellular compartments such as endosome, lysosomes, vacuoles, Golgi, and the plasma membrane[21-23]. In marine phytoplankton such as diatoms, dinoflagellates and coccolithophores, V-type ATPase localizes to the membranes of chloroplasts derived from endosymbiosis and significantly enhances carbon fixation and oxygen production[24]. In *P. falciparum*, previous studies have suggested that V-type ATPase is a major proton pump responsible for maintaining cytosolic pH[2] and DV pH[5] in trophozoite-stage

parasites. However, its localization has remained elusive or somewhat controversial. An early
study indicated that V-type ATPase has heterogenous but not well-defined localization[25]. In
another study, immunofluorescence assays (IFAs) and immuno-EM suggested that V-type
ATPase is localized on the parasite periphery, DV, and small clear vesicles[26]. However, the use
of antibodies raised against the subunits of the bovine V-type ATPase in the study[26] was likely
problematic due to limited specificity towards the *Plasmodium* counterparts. Furthermore,
another report revealed that V-type ATPase also localizes on the RBC membrane, apparently
being exported to the RBC cytosol[27]. Thus, studies of the *Plasmodium falciparum* V-type
ATPase have not provided a consensus on its localization. Therefore, further investigation is
needed to clarify the localization and function of this important proton pump in malaria parasites.

In this study, we employed genetic and biochemical approaches to better understand the
localization and functionality of V-type ATPase in *P. falciparum*. Our data suggest that V-type
ATPase is expressed throughout the IDC and exhibits dynamic localization patterns that vary
depending on the specific developmental stages of the parasite. We have also discovered the
intriguing localization of V-type ATPase on rhoptries and secretory vesicles in merozoites.
Moreover, our results underscore the essentiality of V-type ATPase throughout the IDC.
Depletion of the proton pump leads to the arrest of trophozoites, which contain undigested
hemoglobin in the DV. Using the V-type ATPase specific inhibitor Bafilomycin A1, we have
discovered that V-type ATPase potentially plays important roles during parasite egress and
invasion. Finally, our study demonstrates that the V-type ATPase is sensitive to disruption by
Bafilomycin A1, but the complex stability is less susceptible to certain treatments, such as low
glucose or chloroquine.

Results

Endogenous tagging of V_1A via CRISPR/Cas9 for localization and conditional knockdown

The highly conserved V-Type ATPase has 2 domains, the V_1 domain that hydrolyses ATP and
the V_o domain carrying out proton translocation using the free energy released from ATP
hydrolysis. *P. falciparum* encodes all subunits required to form the V_1 subunit, including subunits
of A (PF3D7_1311900), B (PF3D7_0406100), C (PF3D7_0106100), D (PF3D7_1341900), E
(PF3D7_0934500), F (PF3D7_1140100), G (PF3D7_1323200), and H (PF3D7_1306600). The
V_o includes subunits of a (PF3D7_0806800), c (PF3D7_0519200), c" (PF3D7_1354400) and e
(PF3D7_0721900). Subunits A and B form the ATP hydrolyzing hexamer to power the rotational
movement of the entire proton pump. To study V-type ATPase in *P. falciparum*, we successfully
tagged the endogenous locus of subunit A (V_1A) C-terminally with the triple HA epitope (3xHA)
and elements required for conditional knockdown using the TetR-DOZI-aptamer system[28,29].
This was achieved via CRISPR/Cas9[30,31] with two different gRNA plasmids (**Figure 1A**). We
termed the resulted transgenic parasite line NF54attB-V_1A-3HA[apt]. We also performed limiting
dilution to clone the transgenic parasite line to obtain pure parasites. The D11 clone was chosen
for all subsequent experiments. The genotype of the NF54attB-V_1A-3HA[apt] line (D11 clone) was
verified by diagnostic PCR (**Figure 1B**). The expression of V_1A-3HA proteins at the expected
molecular weight was confirmed by Western blot (**Figure 1C**).

To verify the localization of V-type ATPase in *P. falciparum*, we tightly synchronized the parasites
with alanine/HEPES and performed Immunofluorescence studies (IFAs) over the 48 h lifecycle.
We used Exp2[32], the maker for the Parasitophorous Vacuolar Membrane (PVM), as the indicator
of the parasite periphery. From the time of invasion (T0), we harvested samples every 8 hours in

the ring stage and every 4 hours in trophozoite or schizont stages. V_1A was seen to be
expressed in all asexual blood stages (**Figure 2A**). In the ring stage, V_1A localized to the
parasite periphery, showing good colocalization with Exp2. In the trophozoite stage, V_1A was
predominantly localized on both the Parasite Plasma Membrane (PPM) and the Digestive
Vacuole (DV). This localization pattern was also observed in the early schizont stage. However,
in the late schizont stage, some punctate signals appeared near the parasite periphery. To
further characterize this, we performed IFA on schizonts pre-treated with the egress inhibitor,
ML10[33], a reversible blocker of protein kinase G. Treatment of ML10 inhibits parasite egress,
leading to an increased number of mature late schizont stage parasites. In the treated mature
schizonts, V_1A's localization appeared more punctate, lacking confined signals showing circular
or membranous structures (**Figure 2B**). This prompted us to further verify V-type ATPase's
localization via Immuno-Electron Microscopy. Indeed, immuno-EM revealed a specific
localization of V_1A on rhoptries and some small vesicles near them (**Figure 3**), indicating that V-
type ATPase is also present on the secretory organelles in merozoites. Altogether, we have
used IFA and Immuno-EM to uncover the dynamic localization of V-type ATPase in different
asexual stages of *P. falciparum*. This dynamic localization of V-type ATPase may be associated
with its multifunctionality throughout the 48 h lifecycle.

V-type ATPase is essential for asexual growth and development in *P. falciparum*

To understand the essentiality of V-type ATPase in *P. falciparum*, we performed knockdown
studies using two different methods to remove aTc. In the first method, we synchronized the
parasites several times and started aTc removal in the early ring stage (**Figure 4A**). The
knockdown parasites appeared healthy 24 h post aTc removal, but they failed to produce any
new ring stage parasites in the next 24 h. At 48 h of knockdown, the parasites were arrested in
the trophozoite stage and lysed at 72 h. Thus, V_1A knockdown from the early ring stage resulted
in trophozoite arrestment and parasite demise within one cycle. In the second method, we
started aTc removal from the synchronized trophozoite stage parasites (**Figure 4D**). The
knockdown parasites were able to produce new ring stage parasites at 24 h post aTc removal,
but were arrested again in the trophozoite stage at 48 h. The arrested parasites did not initiate
another IDC, but remained arrested or lysed at 72 h. Thus, both knockdown methods revealed
the essentiality of V-type ATPase for asexual growth and development in *P. falciparum*. In each
experiment, quantification of parasite growth in the aTc (±) conditions was shown in **Figure 4B
and 4E**. The V_1A protein levels of the knockdown experiments were shown by Western blot in
Figure 4C and 4F.

To better understand parasite developmental blockage upon V_1A knockdown, we observed the
morphological changes of the parasites in a more detailed time course experiment (**Figure 4G**).
In tightly synchronized parasites, we collected thin blood smears every 8 hours during the ring
stage and every 4 hours during the trophozoite and schizont stages. Unlike the aTc (+) parasites
that completed their development normally, we observed an arrest in parasite growth in the aTc
(-) culture at 32 h post aTc removal. The knockdown parasites were arrested in the mid-
trophozoite stage and did not progress significantly in the remaining hours of the IDC.
Altogether, using different approaches, we have shown that knockdown of V-type ATPase leads
to the arrest of parasite development in the trophozoite stage within one IDC post aTc removal.
Overall, our results confirm that V-type ATPase is essential for *P. falciparum*, with the
trophozoite stage being the most susceptible to the loss of this proton pump.

V-type ATPase is essential for hemoglobin digestion

V-type ATPase has been shown to regulate the pH of DV in trophozoite stage parasites[3-5]. A low
pH (~ 5.0) is essential for hemoglobin digestion, which is extremely important for malaria
parasites to obtain nutrients and gain space to grow inside RBCs[34]. To get a deeper
understanding of the knockdown effects on the DV, we performed Transmission Electron
Microscopy (TEM) studies. The knockdown experiment was initiated from Percoll-enriched
schizont stage parasites, and we purified parasites by a magnetic column at 36 h post aTc
removal. The parasitemia of both aTc (±) cultures reached nearly 100% (**Figure 5A**). The aTc (-)
culture displayed more cell lysis compared to the aTc (+) culture after magnetic enrichment,
indicating that the integrity of some arrested trophozoites was compromised. Under TEM, the
knockdown parasites exhibited striking features (**Figure 5B**). The DV of knockdown parasites
showed significantly higher electron density than the control. Additionally, we observed
membrane-bound large vesicles inside the DV, indistinguishable from the RBC cytosol in terms
of electron density. These structures apparently contained large amounts of undigested
hemoglobin. Although not measured quantitatively, we also noticed a significant reduction in
hemozoin crystals in the knockdown parasites compared to the control. Consequently,
knockdown of V-type ATPase leads to defects in several steps of hemoglobin metabolism,
including the release of hemoglobin into the DV, breakdown of hemoglobin into small peptides,
and formation of hemozoin.

Hemoglobin uptake is initiated at a specialized structure called cytostome[35], and hemoglobin-
containing vesicles (HCvs) are pinched off from the membranes and fused with the DV to
release their contents for proper digestion and heme detoxification. The hemoglobin containing
vesicles are thus wrapped with two membranes, with the PPM on the outside and the PVM on
the inside[36]. Interestingly, knockdown of V-type ATPase blocked hemoglobin digestion in the DV,
but the upstream trafficking of HCvs enroute to the DV was seemingly not affected. This differed
from a previous study that showed accumulation of HCvs inside the parasite cytosol upon
conditional inactivation of PfVPS45[37], a protein involved in vesicular trafficking. Moreover, the
fusion of the outer membrane of HCVs to the DV membrane is potentially a rapid process as
this event has been rarely observed in wildtype parasites[38]. In the V_1A knockdown parasites, the
undigested hemoglobin was surrounded by a single membrane (**Figure 5B, red arrows**). This
information indicated that fusion of HCVs with the DV membrane was likely normal upon
depletion of V-type ATPase, but degradation of the inner membrane of HCvs was blocked.
Interestingly, we also observed that some vesicles in the DV of the knockdown parasites had
much reduced electron density, likely due to a more advanced level of hemoglobin digestion
(**Figure 5B, blue arrow**). Overall, our results show that V-type ATPase is critical for hemoglobin
release and digestion in the DV.

V-type ATPase is essential for regulating cytosolic pH

Apart from its role in the DV, V-type ATPase is known to regulate cytosolic pH by pumping
protons across the PPM in *P. falciparum*[2]. This understanding was based on the observation
that Bafilomycin A1, an inhibitor of V-type ATPase, caused a dramatic decline of cytosolic pH. A
recent study suggests that Bafilomycin A1 causes a stearic hinderance and blocks the c-ring
rotation by preventing the interaction between subunit a and the c-ring[39]. The effect of V-type
ATPase on cytosolic pH, however, has not been confirmed by genetic studies in *Plasmodium*. To
address this, we utilized the well-established pH measurement technique with BCECF-AM (2',7'-
Bis-(2-Carboxyethyl)-5-(and-6)-Carboxyfluorescein, Acetoxymethyl Ester) in saponin-treated
trophozoites[2]. We quantified cytosolic pH of aTc ± cultures at 36 h post knockdown from Percoll-

enriched schizont stage parasites. We included Bafilomycin A1 as a positive control. Our assays
revealed that cytosolic pH was 7.24±0.08 in the control parasites but reduced to 6.01±0.28 upon
knockdown of V-type ATPase (**Figure 6**). This result confirms that V-type ATPase is the major
proton pump for regulating cytosolic pH in trophozoite stage parasites. Consistent with the
published data[2], we also observed that Bafilomycin A1 caused a profound decrease in cytosolic
pH from 7.24 to 6.51±0.1 within 5 min of treatment. Interestingly, the combination of V_1A
knockdown and Bafilomycin A1 resulted in an even lower cytosolic pH (5.77±0.15), although this
effect was not statistically significant compared to knockdown alone ($p > 0.05$). Overall, our
findings confirm that V-type ATPase plays a major role in regulating cytosolic pH in the
trophozoite stage parasites.

Viability of the V-type ATPase knockdown parasites

Our data have shown that knockdown of V-type ATPase led to the arrest of trophozoite-like
parasites that contained large amounts of undigested hemoglobin in the DV and reduced
cytosolic pH. To understand the viability of these parasites, we performed an aTc addback
experiment post knockdown (**Figure 7A**). We initiated the knockdown from tightly synchronized
Percoll-enriched schizont stage parasites, and performed the addback at different time points
post knockdown, 28 h, 36 h, 40 h, and 48 h. Consistent with **Figure 4G**, the knockdown
parasites at 28 h did not display obvious morphological defects in Giemsa-stained smears
(**Figure 7B**). Upon aTc addback, these parasites were able to produce new ring stage parasites
in the next 24 h. At 36 h post knockdown, a delay in parasite development was readily
detectable. Upon addback, however, most of the arrested parasites managed to produce new
ring stage parasites in the next 24 h. These ring stage parasites continued to become
trophozoites at 48 h of addback. This result suggested that the arrested parasites at 36 h post
knockdown remained largely viable, despite having severe defects in hemoglobin digestion and
pH regulation. At 40 h post knockdown, a smaller percentage of the arrested parasites still
succeeded in producing new ring stage parasites at 24 h post addback. However, just a few
hours later at 48 h post knockdown, aTc addback failed to rescue any parasite growth. The
knockdown parasites either displayed abnormal morphologies or lysed. Altogether, these results
imply that *P. falciparum* parasites can sustain viability in a severely stressed condition for at
least several hours. Remarkably, aTc addback allowed the arrested parasites to recover from
various defects caused by V_1A knockdown, such as reduced cytosolic pH, blockage of
hemoglobin digestion in the DV, and other possible disruptions.

The effect of Bafilomycin A1 on parasite egress and invasion

The localization of V-type ATPase on rhoptries and other secretory organelles prompted us to
investigate the potential role of this proton pump in parasite egress and invasion. However, in
our knockdown studies initiated from the ring, trophozoite, or schizont stage (**Figures 4-7**), an
extended period of ~ 30 h was consistently required before the onset of detectable defects. In
addition, all knockdown approaches led to the arrest of trophozoites, making examinations of
schizonts infeasible. To address the delayed effects of genetic knockdown, we reasoned that a
pool of pre-existing V_1A proteins, present before aTc removal, might permit parasite survival and
maturation over an extended period until the proteins were fully exhausted. To overcome this
limitation, we used Bafilomycin A1 to rapidly block V-type ATPase in the late schizonts and
observed its effect on parasite egress and invasion. As shown previously, Bafilomycin A1
reduced cytosolic pH from ~ 7.3 to ~ 6.5 in about 5 min in *in vitro* assays (**Figure 6**).

In brief, we tightly synchronized the parasites with alanine/HEPES and used ML10 to block
parasite egress, thereby increasing the percentage of mature schizonts. At the mature schizont
stage (~ 0 h), we added Bafilomycin A1 (100 nM) for 2 h, washed it out, and allowed parasites
to invade new RBCs for 4 h. At 6 h post the initiation of the experiment, we counted the
parasitemia of newly formed ring stage parasites and used it as the readout of successful
egress and invasion (**Figure 8A**). Parasite growth and morphology were also examined at 24 h
and 48 h post the initiation of the experiment. At 6 h, while the new ring stage reached
4.77%±0.25% in the control culture, the Bafilomycin A1 treated culture had a significantly lower
parasitemia (0.24%±0.07%) (**Figure 8B**). In the treated culture, events of merozoite invasion
were commonly observed, however, formation of new rings was drastically reduced (**Figure
8C**). In the next 24 h and 48 h, the control culture followed the normal progression of the
lifecycle. In contrast, the treated culture barely increased parasitemia (**Figure 8B**). The
parasites displayed abnormal morphologies or failed to invade RBCs. It was interesting to note
that in the treated culture, some merozoite-like parasites appeared to be attached to the RBC
surface even at 48 h post the initiation of the experiment (**Figure 8C**). Collectively, treatment of
Bafilomycin A1 in the mature schizont stage leads to dramatic reduction in the formation of new
ring stage parasites.

Stability of the V-type ATPase complex under various conditions

In model eukaryotes, it has been well established that V-type ATPase undergoes a reversible
process of assembly and disassembly to rapidly regulate its function upon encountering
different conditions[17]. For example, the yeast V-type ATPase is very sensitive to the bioenergetic
status of the cell and V_1 and V_o disassociate in a low glucose condition[40]. Disassociation of the
two domains completely abolishes the proton pumping function as well as the ATP hydrolysis
activity of the V_1 domain[41]. To assess if the *Plasmodium falciparum* V-type ATPase can regulate
its function by changing complex stability, we treated synchronized trophozoite stage parasites
with Bafilomycin A1, chloroquine, and 1 mM glucose and verified V_1V_o complex formation
through BN/PAGE. In parallel, the expression level of the V_1A monomers was monitored by
SDS/PAGE. As shown in **Figure 9**, in the control protein lysate, we detected a high-molecular-
weight band near the marker of 1.026 MDa, which approximately corresponded to the total
molecular weights of the entire V-type ATPase[21]. In addition, in BN/PAGE, smaller bands near
the markers of 720 kDa, 240 kDa, and 66 kDa were also detected. The nature of these smaller
complexes is uncertain at present. They could be assembly intermediates of the V_1V_o complex
or electrophoretic degradation products, although enough protease inhibitor cocktails were
included throughout the experiment. Upon Bafilomycin A1 (10 nM) treatment, the top band near
1.026 MDa was detectable after 2 h, but gradually decreased in intensity after 4 h and 8 h. By
contrast, the monomer V_1A remained more stable throughout the time course. This result
indicated that the V-type ATPase dissembled upon Bafilomycin A1 treatment longer than 2 h.
Upon treatment with chloroquine (100 nM) or 1 mM glucose, however, the top band near 1.026
MDa remained largely constant even after 8 h. The V_1A monomer was stably expressed in
chloroquine treated samples and gradually decreased in the low glucose treated samples.
Together, these data implied that the V-type ATPase complex was readily disrupted by
Bafilomycin A1, but not by treatment of chloroquine or a low glucose condition.

Discussion

In this study, we have characterized the essentiality, localization, and biological function of V-
type ATPase in *Plasmodium falciparum*. By tagging the subunit A of V_1 via CRISPR/Cas9, we

generated a conditional knockdown parasite line under aTc regulation and obtained several
interesting results.

We have found that V-type ATPase is essential for *P. falciparum* during asexual growth and
development, with the trophozoite stage being particularly vulnerable to death when this proton
pump is depleted (**Figures 4-7**). The localization of *P. falciparum* V-type ATPase is highly
dynamic throughout the 48 h asexual lifecycle (**Figures 2-3**). Specifically, in trophozoite stage
parasites, V-type ATPase localizes on both the parasite plasma membrane (PPM) and digestive
vacuole (DV). This localization pattern is consistent with its role in regulating cytosolic pH[2] and
DV pH[5]. We also observed a drastic reduction in cytosolic pH in the V_1A knockdown parasites
during the trophozoite stage (**Figure 6**). Moreover, our data suggest that V-type ATPase is also
expressed and localized on the PPM in the ring stage. However, we need to be cautious when
defining the precise localization of V-type ATPase in the small ring stage parasites. A recent
study implied that V-type ATPase might not play a major role in proton pumping during the ring
stage, as the pyrophosphate driven proton pump, PfVP1, is essential for this stage[15].
Conditional depletion of PfVP1 blocks ring stage development and the ring-to-trophozoite
transition, which is apparently not rescued by the V-type ATPase. This indicates that V-type
ATPase might either not fulfill proton pumping across the PPM in the ring stage or have a
negligible role. In the late schizont stage, we have observed punctate signals of V_1A in IFA
without defined membranous staining (**Figure 2B**). Immuno-EM further revealed V_1A's
localization on rhoptries and secretory organelles near rhoptries (**Figure 3**). To our best
knowledge, this is the first report that observed such localization of V-type ATPase on mature
secretory organelles in protozoa. A previous study highlighted the function of V-type ATPase in
facilitating the maturation of secretory organelles in *Toxoplasma gondii*, an apicomplexan
parasite related to *Plasmodium*[42], but the presence of V-type ATPase on mature rhoptries or
micronemes was not detected[42]. Thus, the unique localization of V-type ATPase on rhoptries
and secretory organelles in *Plasmodium* has led us to investigate the role of this proton pump in
parasite egress and invasion. To alleviate the slow effect of genetic knockdown, we used
Bafilomycin A1 to chemically inhibit V-type ATPase in a short time (2 h) and observed serious
defects of the parasites in initiating the next lifecycle (**Figure 8**). Bafilomycin A1's specificity as a
V-type ATPase blocker has been well documented and it has been used as a standard reagent
in *P. falciparum* studies[2,5]. We quantified the number of new ring stage parasites post
Bafilomycin A1 treatment and used it as a readout to indicate the overall effects on parasite
egress and invasion. However, our data did not provide detailed mechanisms underlying the
mode of action of Bafilomycin A1 in mature schizont stage parasites. It is possible that
Bafilomycin A1 exerts its effects through multiple mechanisms, such as killing the schizonts,
blocking egress, and inhibiting invasion. Future investigations will be necessary to characterize
these processes in detail. Nevertheless, our data implied that V-type ATPase likely plays a role
in parasite egress and invasion.

In the trophozoite stage, we have discovered that V-type ATPase is extremely important for
hemoglobin digestion in the DV. This finding aligns with the previously reported role of V-type
ATPase in maintaining DV pH[5]. However, observation of undigested hemoglobin in the DV of
V_1A knockdown parasites was astonishing and stands out as a novel finding of this study
(**Figure 5**). To date, the processes of hemoglobin uptake from cytostomes and the delivery of
hemoglobin containing vesicles (HCvs) to the DV remain enigmatic in *P. falciparum,* as these
processes are likely dynamic, transient, and difficult to capture. A previous study reported the
rare fusion of one HCv with the DV membrane among > 600 examined thin sections under

transmission electron microscopy[38]. Since cytostomes are invaginated from the PPM and PVM,
the HCvs are enclosed by the PPM on the outside and the PVM on the inside. The fusion event
indicates that the PPM of HCvs fuses with the DV membrane[38]. Therefore, the membrane
surrounding the undigested hemoglobin in our data should be the PVM, not the PPM. We can
speculate that after HCvs are fused with the DV membrane, the leftover PVM needs to be
digested first by specific enzymes (likely lipases) before hemoglobin can be digested by aspartic
proteases and cysteine proteases[43]. The enzyme responsible for digesting the PVM of HCvs
remains unknown, but it is evidently pH dependent and not among the list of enzymes known to
digest hemoglobin[44].

Moreover, we have detected the native *P. falciparum* V-type ATPase complex by BN/PAGE. In
addition to the large molecular weight band near 1.0 MDa, many smaller subcomplex-like
moieties were also detected at various molecular weights (**Figure 9**). Though unknown, we
speculate that formation of the *P. falciparum* V-type ATPase complex is a highly regulated
process, and the intermediate complexes might be involved in the assembly and/or disassembly
processes. In mammalian cells, the V-type ATPase was also examined by BN/PAGE but few
intermediate complexes were detected[45]. In addition, we noted that the complex was readily
disrupted upon Bafilomycin A1 treatment but remained more stable upon challenges of
chloroquine and low glucose. The glucose effect on *P. falciparum* V-type ATPase was not
expected since low glucose levels are known to dissociate the complex in yeast[40]. This could
reflect the unique features of V-type ATPase in *P. falciparum* in a bioenergetically different
environment from that of yeast.

In summary, our data have provided valuable insights into the localization and functions of V-
type ATPase in *Plasmodium falciparum*, highlighting its critical roles in regulating pH in various
subcellular compartments. As we prepared this manuscript, a very recent report by Alder and
colleagues also unveiled the localization and function of the *P. falciparum* V-type ATPase, using
an inducible knockout approach[46]. Their study has characterized defects related to knockdown
of subunit B in the V_1 domain and subunits a and c in the V_o domain. Interestingly, our findings
from knockdown of V_1A align well with those obtained from inducible knockout of V_1B. Given
that subunits A and B work together in the ATP hydrolysis center, genetic knockdown (or
knockout) of either A or B leads to common downstream effects. Nonetheless, the combined
studies on the two major subunits of V-type ATPase, both from our work and others[46], have
collectively revealed the dynamic localization of V-type ATPase in the asexual blood stage of *P.*
falciparum. These studies shed light on the significant role of V-type ATPase in parasite growth
and replication, thereby enhancing our understanding of *Plasmodium* biology. Moreover, results
of our work and Alder *et al.*[46] could provide valuable guidance for drug development strategies
as V-type ATPase was implicated as a drug target[47,48].

Figure Legends

Figure 1. Endogenous tagging of V_1A via CRISPR/Cas9 for conditional expression

A, Schematic of modifying the V_1A genetic locus via CRISPR/Cas9 mediated double crossover
recombination. The primer positions were indicated by black arrows with sequences listed in
Table 1.
395

B, Diagnostic PCR/DNA electrophoresis showing the corrected modification of the V_1A genetic locus in the D11 clone of NF54attB-V_1A-3HA^{apt}. The band sizes of PCR products were 0.8 kb (lane 1), 1.2 kb (lane 4), and 1.9 kb (lane 8). One non-specific band was detected in lane 1 (star).

C, Western blot showing the expression of V_1A-3HA proteins in the transgenic parasite line, NF54attB-V_1A-3HA^{apt}.

Figure 2. Localization of V_1A throughout the asexual blood stages

A, Localization of V_1A throughout the lifecycle. In the NF54attB-V_1A-3HA^{apt} line, V_1A was detected by anti HA and FITC-labeled secondary antibodies. Exp2 was detected by anti-Exp2[49] and TRITC-labeled secondary antibodies.

B, Localization of V_1A in mature schizont stage parasites treated with ML10. The synchronized late-trophozoite stage parasites were treated with 25 nM of ML10 for 14 h to reach the mature schizont stage. V_1A was detected by anti HA and FITC-labeled secondary antibodies. DNA was stained with DAPI.

In each condition, representative images of at least 25 parasites were shown.

Figure 3. Localization of V_1A on secretory organelles in mature schizont stage parasites

A-F, Detection of V_1A signals on rhoptries and secretory organelles near rhoptries in the parasite line NF54attB-V_1A-3HA^{apt} by immuno-electron microscopy. Bars, 500 nm.

G-H, Negative control images with the primary antibody omitted throughout the experiment. Bars, 500 nm.

Figure 4. Essentiality of V-type ATPase in the asexual blood stages

Giemsa-stained images showing parasite morphologies resulted from knocking down of V_1A from the ring stage (**A**), or from the trophozoite stage (**D**), in the NF54attB-V_1A-3HA^{apt} line.

Quantification of parasitemia in the aTc +/- cultures (**B, E**). At each time point, parasitemia was determined from counting 1000 RBCs under a light microscope. Error bars indicated the s.d. of three technical measurements.

Western blot showing the expression level of V_1A in the knockdown experiments (**C, F**). The same blot was re-probed with anti-Exp2 to show loading controls.

G, Giemsa-stained images showing parasite morphological changes over the time course experiment.

Experiments of A-F were repeated three times. Panel G was repeated two times.

Figure 5. V-type ATPase is essential for hemoglobin digestion in the digestive vacuole

A, Giemsa-stained images of control and V_1A knockdown parasites enriched by a magnetic column.

B, Electron Transmission Microscopic images of control and V_1A knockdown parasites. In the aTc - parasites, many vesicles inside the digestive vacuole had electron dense signals and were

434 surrounded by membranes (red arrows). Some vesicles had translucent appearance but were
435 still surrounded by membranes (blue arrow).

Figure 6. V-type ATPase regulates cytosolic pH in trophozoite stage parasites

437 At 36 h post knockdown, aTc +/- cultures of the NF54attB-V$_1$A-3HA^{apt} line were subjected to pH
438 measurement using the BCECF-AM method in the absence of presence of 100 nM Bafilomycin
439 A1 (100 nM) for 5 min. Error bars indicated the s.d. of 3-4 measurements in each condition.
440 Statistical analysis was done by student t-test. This experiment was repeated two times.

Figure 7. Viability of the V-type ATPase knockdown parasites

442 **A**, Schematic diagram of the knockdown/addback experiment. At various time points post
443 knockdown, the cultures were fed with aTc + medium and further monitored for 48 h.

444 **B**, Giemsa-stained images showing morphological changes of the parasites upon aTc addback.

445 This experiment was repeated three times.

Figure 8. Effects of Bafilomycin A1 on parasite egress and invasion

A, Schematic illustration of the experimental design generated by BioRender. Mature schizont
stage parasites (NF54attB-V$_1$A-3HA^{apt} under 250 nM) were treated with Bafilomycin A1 (100
nM) for 2 h. The drug was washed out and parasites were allowed to invade new RBCs for 4 h.
At 6 h, the parasitemia of newly formed ring stage parasites was monitored. Parasite
morphology and number were further examined at 24 h and 4 h post the initiation of the
experiment. aTc (250nM) was maintained throughout the experiment.

B, Giemsa-stained imaging showing parasite morphologies in the control and treated cultures.

C, Quantification of parasite growth throughout the experiment. Parasitemia was determined by
counting 1000 RBCs in each smear under a light microscope. Error bars indicated the s.d. of 3
measurements at each time point.

This experiment was repeated five times.

Figure 9. The stability of V-type ATPase complex under various conditions

459 Synchronized trophozoite stage parasites (NF54attB-V$_1$A-3HA^{apt} under 250 nM) were treated
460 with Bafilomycin A1 (10 nM), Chloroquine (100 nM), or 1 mM glucose for 2 h, 4 h, and 8 h.
461 Samples were harvested and analyzed by BN-PAGE and SDS-PAGE. The arrow indicated the
462 large molecular weight band corresponding to the estimated size of the *P. falciparum* V-type
463 ATPase. aTc (250nM) was maintained throughout the experiment.

Materials and Methods

Table 1. Primers and oligoes used in this study

ID	PRIMER NAME	SEQUENCE
P1	A-VATPase gRNA1	TCATATTAAGTATATAATATTAGTAGGTTTAAGAAACTTGAGTTTCAGAGCTATGCTGGA
P2	A-VATPase gRNA2	CATATTAAGTATATAATATTGAAACTTGATGGAAAAATGATGTTTCAGAGCTATGCTGGA
P3	A-VATPase gRNA1-N20	AGTAGGTTTAAGAAACTTGA
P4	A-VATPase gRNA2-N21	GAAACTTGATGGAAAAATGAT
P5	N20CheckR	ATATGAATTACAAATATTGCATAAAGA
P6	RevOligo	TAGGAAATAATAAAAAAGCACCGACTCG
P7	Super CPF1	CTTTTATTTTTACTGTAATATAATTTTTTATAATGTAAAAATAAAGGG
P8	A-VATPase 5HRF	gatatctacgatatcccgcggtccggaGGTTCTCCATCTCGTATAGG
P9	A-VATPase 5HRR	ATGGTAACCTGcTTcTCCATaAgaTTaCgaAgtCCaACaGTTATTTCTTCTTCAAGTGCC
P10	A-VATPase 3UTRF	atgcgcgcGTGATTTATTTATTAGTTGAATAATTAAAGC
P11	A-VATPase 3UTRR	tccggaccgcgggatatcgtagatatcGAAAAAGTTTACCTTAGTATGATAAAAAAATG
P12	pMG75-BBHA REV	TGGGCCCGAATTCTCATCATTGTGC
P13	pMG75SeqF	CTTTAAATTCATGCAAAAATTTAC
P14	A-VATPase 5chk	GGTTATCCAGCTTATTTAGGTGC
P15	A-VATPase 3chk New	GATCTACTTACAATTTTTTTCCTTC

1, Plasmid Construction

Subunit A of *Plasmodium falciparum* V-Type ATPase (V$_1$A, PF3D7_1311900) was endogenously tagged with a 3xHA epitope and the TetR-DOZI-Aptamer system at the C-terminus via CRISPR/Cas9. Two NFCas9/gRNA constructs were made according to our cloning procedures published earlier[50]. The DNA oligoes used for cloning and sequencing were listed in **Table 1** (P1-P7). To make the pMG75 template plasmid for double crossover recombination, the 5' and 3' homologous regions (5'HR and 3'HR) of V$_1$A were PCR amplified from the wildtype genomic DNA using primers P8 and P9 for 5'HR and P10 and P11 for 3'HR. The PCR product was purified and then digested with SacII/BstEII for 5'HR and BssHII/SacII for 3'HR. The digested fragments were sequentially cloned into the pMG75 vector bearing Blasticidin deaminase by T4 DNA ligase. The clones were screened by proper restriction digestion. The positive clones were further confirmed by Sanger sequencing using the primers P12-P13. The final maxi-prep

plasmid of pMG75-BSD-V_1A-3HA was linearized with EcoRV before being transfected into
parasites.

2, Parasite Culture and Transfection

Plasmodium falciparum wildtype strain NF54attB was generously provided by Dr. Joshua Beck
(Iowa State University). This line is compatible with all three transfection markers that are
commonly used, including BSD (Blasticidin deaminase), hDHFR (human dihydrofolate
reductase) and yDHODH (yeast dihydroorotate dehydrogenase). *P. falciparum* was cultured in
human O^+ red blood cells (5% hematocrit, Interstate Blood Bank, Inc.) in the standard RPMI
1640 medium containing 0.5% AlbuMAX® II (Invitrogen), 15 mM HEPES, 10 mg/L
hypoxanthine, 25 mM $NaHCO_3$, and 50 mg/L gentamicin. Transfections were performed using
the standard methods in the ring stage with details shown previously[50,51]. The concentration of
aTc was 250 nM in all aTc (+) cultures.

3, Immunofluorescence assays (IFA)

Parasites were tightly synchronized with alanine (0.5 M)/HEPES (10 mM) several times. At each
time point of the experiment, ~ 50 µL parasitized RBCs were collected and fixed with 4%
paraformaldehyde, 0.0075% glutaraldehyde in 1X PBS. Blocking was done by 3% BSA/PBS for
overnight at 4°C. Fixed cells were permeabilized by Triton X-100 (0.25%)/PBS and incubated
with primary antibodies overnight at 4°C, including α-HA (1:300, mouse, sc-7392, Santa Cruz
Biotechnology) and α-Exp2 (1:500, rabbit)[49]. Following thorough washing, the samples were
incubated with FITC- or TRITC- labeled secondary antibodies for 3 h at room temperature.
Other procedures followed standard IFA protocols. Images were captured using the Nikon Ti
microscope.

4, Western Blot for BN-PAGE and SDS-PAGE

Parasite cultures were lysed by Saponin (0.05%)/PBS and host hemoglobin was removed by
several washes with 1X PBS. From one culture condition, two aliquots of samples were
obtained for BN-PAGE and SDS-PAGE, respectively. For BN-PAGE, the saponin lysed pellet
was solubilized with 1% glycodiosgenin (GDN) in MESH buffer (250 mM sucrose, 10 mM
HEPES, 1 mM EDTA, 1X protease inhibitor cocktail) for overnight at 4°C. The supernatant after
a high-speed centrifugation (13,000 rpm, 10 min) was used for Blue-Native gel electrophoresis
and Western blot. For SDS-PAGE, the saponin lysed pellet was solubilized by 2% SDS/Tris (65
mM, pH 6.8) for overnight at 4°C. The supernatant after a high-speed centrifugation (13,000
rpm, 10 min) was used for denatured electrophoresis. For all samples, the protein
concentrations were determined by the Pierce™ BCA Protein Assay Kit (ThermoFisher
Scientific) to ensure equal loading. Antibodies used for Western blot included α-HA (1:10,000,
mouse, sc-7392, Santa Cruz Biotechnology), α-Exp2 (1:10,000, rabbit)[49], HRP-conjugated goat
anti-mouse secondary (1:10,000, A16078, ThermoFisher Scientific), HRP-conjugated goat anti-
rabbit secondary (1:10,000, 31460, ThermoFisher Scientific). The membranes were incubated
with Pierce ECL Western Blot Substrates and developed using the Bio-Rad chemidoc imaging
system.

5, Transmission Electron Microscopy

Parasites were synchronized with alanine (0.5 M)/HEPES (10 mM) several times and the
schizont stage parasites were enriched by Percoll. After several washes with 1X PBS, the

schizonts were added with new RBCs and cultured in aTc +/- conditions for 36 h. Both cultures
were then passed through a MACS magnetic column (MiltenyiBiotec) to enrich trophozoite
stage parasites, which were fixed with freshly made 2% paraformaldehyde, 2.5% glutaraldehyde
in 100mM sodium cacodylate buffer for overnight at 4°C. The samples were shipped to Dr.
Wandy Beatty from the Molecular Microbiology Imaging Facility at Washington University, St.
Louis, MO for further processing and imaging. Other procedures were carried out as shown
previously[51].

6, Immuno-Electron Microscopy

Parasites were synchronized with alanine (0.5 M)/HEPES (10 mM) several times and the
trophozoite stage parasites were enriched using a MACS magnetic column (MiltenyiBiotec). The
enriched parasites were treated with 25 nM ML10[33] for 14 h and then fixed with freshly made
4% paraformaldehyde/0.05% glutaraldehyde/100mM PIPES buffer for overnight at 4°C. The
samples were shipped to Dr. Wandy Beatty from the Molecular Microbiology Imaging Facility at
Washington University, St. Louis, MO for further processing and imaging. Other procedures
were carried out as previously published[50]. Antibodies used in the study included α-HA (1:300,
mouse, sc-7392, Santa Cruz Biotechnology) and goat anti-mouse IgG 18 nm colloidal gold-
conjugated secondary antibody (1:100, Jackson ImmunoResearch Laboratories). A parallel
sample with the primary antibody omitted was used as the negative control, which did not react
to the secondary antibody.

7, pH measurement with BCECF-AM, a ratio-metric pH indicator

pH measurement was carried out according to the published protocols[2]. Post 36 h of aTc
removal, the control and knockdown cultures were incubated with 4 µM BCECF-AM (2',7'-Bis-
(2-Carboxyethyl)-5-(and-6)-Carboxyfluorescein, Acetoxymethyl Ester, Fisher Scientific) for 30
min at 37°C. Pluronic at 1:1000 dilution was added to increase cellular permeability to BCECF-
AM. The parasites were treated with rapid saponin lysis to remove hemoglobin. The pellet was
washed two times with saline/glucose buffer (NaCl 125 mM, KCl 5 mM, MgCl2 1 mM, glucose
20 mM, HEPES 25 mM, pH 7.4) and resuspended in 1 mL of warm saline/glucose buffer. The
sample's fluorescence was monitored using a fluorescence spectrometer (Hitachi F-7000) at
535 nm with excitation wavelengths set at 490/440 nm. When Bafilomycin A1 (100 nM) was
used in aTc +/- conditions, the drug was incubated with saponin-lysed and washed parasites for
5 min before they were further analyzed by Hitachi F-7000. The fluorescence ratios were back
calculated to pH values according to a linear regression curve generated from several known
pH standards (pH 6.8, 7.1, 7.4), as shown previously[2].

Acknowledgement

We are grateful for the technical assistance and intellectual input from members of the Center
for Parasitology at Drexel University College of Medicine. We thank Dr. Joshua Beck (Iowa
State University) for providing the parasite line, Drs. Jaquin Niles (MIT) and Sean Prigge (Johns
Hopkins University) for providing the TetR-DOZI-aptamer plasmids, Drs. Baker and Ooij for
providing the compound, ML10. We thank Dr. Wandy Beatty at Washington University in St.
Louis for performing transmission electron microscopy and immuno-electron microscopy
studies.

References

WHO. World Malaria Report. (2022).

Saliba, K. J. & Kirk, K. pH regulation in the intracellular malaria parasite, Plasmodium falciparum. H(+) extrusion via a V-type H(+)-ATPase. *J Biol Chem* **274**, 33213-33219 (1999). https://doi.org:10.1074/jbc.274.47.33213

Kuhn, Y., Rohrbach, P. & Lanzer, M. Quantitative pH measurements in Plasmodium falciparum-infected erythrocytes using pHluorin. *Cell Microbiol* **9**, 1004-1013 (2007). https://doi.org:10.1111/j.1462-5822.2006.00847.x

Bennett, T. N. *et al.* Drug resistance-associated pfCRT mutations confer decreased Plasmodium falciparum digestive vacuolar pH. *Mol Biochem Parasitol* **133**, 99-114 (2004). https://doi.org:10.1016/j.molbiopara.2003.09.008

Saliba, K. J. *et al.* Acidification of the malaria parasite's digestive vacuole by a H+-ATPase and a H+-pyrophosphatase. *J Biol Chem* **278**, 5605-5612 (2003). https://doi.org:10.1074/jbc.M208648200

Klonis, N. *et al.* Evaluation of pH during cytostomal endocytosis and vacuolar catabolism of haemoglobin in Plasmodium falciparum. *Biochem J* **407**, 343-354 (2007). https://doi.org:10.1042/BJ20070934

Mohring, F. *et al.* Determination of glutathione redox potential and pH value in subcellular compartments of malaria parasites. *Free Radic Biol Med* **104**, 104-117 (2017). https://doi.org:10.1016/j.freeradbiomed.2017.01.001

Casey, J. R., Grinstein, S. & Orlowski, J. Sensors and regulators of intracellular pH. *Nat Rev Mol Cell Biol* **11**, 50-61 (2010). https://doi.org:10.1038/nrm2820

Cosse, M. & Seidel, T. Plant Proton Pumps and Cytosolic pH-Homeostasis. *Front Plant Sci* **12**, 672873 (2021). https://doi.org:10.3389/fpls.2021.672873

Ginsburg, H. Abundant proton pumping in Plasmodium falciparum, but why? *Trends Parasitol* **18**, 483-486 (2002). https://doi.org:10.1016/s1471-4922(02)02350-4

McIntosh, M. T. & Vaidya, A. B. Vacuolar type H+ pumping pyrophosphatases of parasitic protozoa. *Int J Parasitol* **32**, 1-14 (2002). https://doi.org:10.1016/s0020-7519(01)00325-3

McIntosh, M. T., Drozdowicz, Y. M., Laroiya, K., Rea, P. A. & Vaidya, A. B. Two classes of plant-like vacuolar-type H(+)-pyrophosphatases in malaria parasites. *Mol Biochem Parasitol* **114**, 183-195 (2001). https://doi.org:10.1016/s0166-6851(01)00251-1

Luo, S., Marchesini, N., Moreno, S. N. & Docampo, R. A plant-like vacuolar H(+)-pyrophosphatase in Plasmodium falciparum. *FEBS Lett* **460**, 217-220 (1999). https://doi.org:10.1016/s0014-5793(99)01353-8

Karcz, S. R., Herrmann, V. R. & Cowman, A. F. Cloning and characterization of a vacuolar ATPase A subunit homologue from Plasmodium falciparum. *Mol Biochem Parasitol* **58**, 333-344 (1993). https://doi.org:10.1016/0166-6851(93)90056-4

Omobukola Solebo, L. L., Jing Zhou, Tian-min Fu, Hangjun Ke. Malaria parasites utilize pyrophosphate to fuel an essential proton pump in the ring stage and the transition to trophozoite stage. *BioRxiv* (2021). https://doi.org:doi: https://doi.org/10.1101/2021.10.25.465524

Vasanthakumar, T. & Rubinstein, J. L. Structure and Roles of V-type ATPases. *Trends Biochem Sci* **45**, 295-307 (2020). https://doi.org:10.1016/j.tibs.2019.12.007

Kane, P. M. Disassembly and reassembly of the yeast vacuolar H(+)-ATPase in vivo. *J Biol Chem* **270**, 17025-17032 (1995).

Wang, H. & Rubinstein, J. L. CryoEM of V-ATPases: Assembly, disassembly, and inhibition. *Curr Opin Struct Biol* **80**, 102592 (2023). https://doi.org:10.1016/j.sbi.2023.102592

19 Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite
Plasmodium falciparum by saturation mutagenesis. *Science* **360** (2018).
https://doi.org:10.1126/science.aap7847
20 Bushell, E. *et al.* Functional Profiling of a Plasmodium Genome Reveals an Abundance
of Essential Genes. *Cell* **170**, 260-272 e268 (2017).
https://doi.org:10.1016/j.cell.2017.06.030
21 Harrison, M. A. & Muench, S. P. The Vacuolar ATPase - A Nano-scale Motor That Drives
Cell Biology. *Subcell Biochem* **87**, 409-459 (2018). https://doi.org:10.1007/978-981-10-
7757-9_14
22 Graham, L. A., Flannery, A. R. & Stevens, T. H. Structure and assembly of the yeast V-
ATPase. *J Bioenerg Biomembr* **35**, 301-312 (2003).
https://doi.org:10.1023/a:1025772730586
23 Lupanga, U. *et al.* The Arabidopsis V-ATPase is localized to the TGN/EE via a seed
plant-specific motif. *Elife* **9** (2020). https://doi.org:10.7554/eLife.60568
24 Yee, D. P. *et al.* The V-type ATPase enhances photosynthesis in marine phytoplankton
and further links phagocytosis to symbiogenesis. *Curr Biol* **33**, 2541-2547 e2545 (2023).
https://doi.org:10.1016/j.cub.2023.05.020
25 Karcz, S. R., Herrmann, V. R., Trottein, F. & Cowman, A. F. Cloning and characterization
of the vacuolar ATPase B subunit from Plasmodium falciparum. *Mol Biochem Parasitol*
65, 123-133 (1994). https://doi.org:10.1016/0166-6851(94)90121-x
26 Hayashi, M. *et al.* Vacuolar H(+)-ATPase localized in plasma membranes of malaria
parasite cells, Plasmodium falciparum, is involved in regional acidification of parasitized
erythrocytes. *J Biol Chem* **275**, 34353-34358 (2000).
https://doi.org:10.1074/jbc.M003323200
27 Marchesini, N., Vieira, M., Luo, S., Moreno, S. N. & Docampo, R. A malaria parasite-
encoded vacuolar H(+)-ATPase is targeted to the host erythrocyte. *J Biol Chem* **280**,
36841-36847 (2005). https://doi.org:10.1074/jbc.M507727200
28 Rajaram, K., Liu, H. B. & Prigge, S. T. Redesigned TetR-Aptamer System To Control
Gene Expression in Plasmodium falciparum. *mSphere* **5** (2020).
https://doi.org:10.1128/mSphere.00457-20
29 Ganesan, S. M., Falla, A., Goldfless, S. J., Nasamu, A. S. & Niles, J. C. Synthetic RNA-
protein modules integrated with native translation mechanisms to control gene
expression in malaria parasites. *Nat Commun* **7**, 10727 (2016).
https://doi.org:10.1038/ncomms10727
30 Wagner, J. C., Platt, R. J., Goldfless, S. J., Zhang, F. & Niles, J. C. Efficient CRISPR-
Cas9-mediated genome editing in Plasmodium falciparum. *Nat Methods* **11**, 915-918
(2014). https://doi.org:10.1038/nmeth.3063
31 Ghorbal, M. *et al.* Genome editing in the human malaria parasite Plasmodium falciparum
using the CRISPR-Cas9 system. *Nat Biotechnol* **32**, 819-821 (2014).
https://doi.org:10.1038/nbt.2925
32 Ho, C. M. *et al.* Malaria parasite translocon structure and mechanism of effector export.
Nature **561**, 70-75 (2018). https://doi.org:10.1038/s41586-018-0469-4
33 Ressurreicao, M. *et al.* Use of a highly specific kinase inhibitor for rapid, simple and
precise synchronization of Plasmodium falciparum and Plasmodium knowlesi asexual
blood-stage parasites. *PLoS One* **15**, e0235798 (2020).
https://doi.org:10.1371/journal.pone.0235798
34 Krugliak, M., Zhang, J. & Ginsburg, H. Intraerythrocytic Plasmodium falciparum utilizes
only a fraction of the amino acids derived from the digestion of host cell cytosol for the
biosynthesis of its proteins. *Mol Biochem Parasitol* **119**, 249-256 (2002).
https://doi.org:10.1016/s0166-6851(01)00427-3

35   Goldberg, D. E. Hemoglobin degradation in Plasmodium-infected red blood cells. *Semin*
Cell Biol **4**, 355-361 (1993). https://doi.org:10.1006/scel.1993.1042
36   Spielmann, T., Gras, S., Sabitzki, R. & Meissner, M. Endocytosis in Plasmodium and
Toxoplasma Parasites. *Trends Parasitol* **36**, 520-532 (2020).
https://doi.org:10.1016/j.pt.2020.03.010
37   Jonscher, E. *et al.* PfVPS45 Is Required for Host Cell Cytosol Uptake by Malaria Blood
Stage Parasites. *Cell Host Microbe* **25**, 166-173 e165 (2019).
https://doi.org:10.1016/j.chom.2018.11.010
38   Milani, K. J., Schneider, T. G. & Taraschi, T. F. Defining the morphology and mechanism
of the hemoglobin transport pathway in Plasmodium falciparum-infected erythrocytes.
Eukaryot Cell **14**, 415-426 (2015). https://doi.org:10.1128/EC.00267-14
39   Wang, R. *et al.* Molecular basis of V-ATPase inhibition by bafilomycin A1. *Nature*
Communications **12**, 1782 (2021). https://doi.org:10.1038/s41467-021-22111-5
40   Parra, K. J. & Kane, P. M. Reversible association between the V1 and V0 domains of
yeast vacuolar H+-ATPase is an unconventional glucose-induced effect. *Mol Cell Biol*
18, 7064-7074 (1998). https://doi.org:10.1128/MCB.18.12.7064
41   Parra, K. J., Keenan, K. L. & Kane, P. M. The H subunit (Vma13p) of the yeast V-ATPase
inhibits the ATPase activity of cytosolic V1 complexes. *J Biol Chem* **275**, 21761-21767
(2000). https://doi.org:10.1074/jbc.M002305200
42   Stasic, A. J. *et al.* The Toxoplasma Vacuolar H(+)-ATPase Regulates Intracellular pH and
Impacts the Maturation of Essential Secretory Proteins. *Cell Rep* **27**, 2132-2146 e2137
(2019). https://doi.org:10.1016/j.celrep.2019.04.038
43   Francis, S. E., Sullivan, D. J., Jr. & Goldberg, D. E. Hemoglobin metabolism in the
malaria parasite Plasmodium falciparum. *Annu Rev Microbiol* **51**, 97-123 (1997).
https://doi.org:10.1146/annurev.micro.51.1.97
44   Liu, J., Istvan, E. S., Gluzman, I. Y., Gross, J. & Goldberg, D. E. Plasmodium falciparum
ensures its amino acid supply with multiple acquisition pathways and redundant
proteolytic enzyme systems. *Proc Natl Acad Sci U S A* **103**, 8840-8845 (2006).
https://doi.org:10.1073/pnas.0601876103
45   Ludwig, J. *et al.* Identification and characterization of a novel 9.2-kDa membrane sector-
associated protein of vacuolar proton-ATPase from chromaffin granules. *J Biol Chem*
273, 10939-10947 (1998). https://doi.org:10.1074/jbc.273.18.10939
46   Alder, A. *et al.* The role of Plasmodium V-ATPase in vacuolar physiology and antimalarial
drug uptake. *Proc Natl Acad Sci U S A* **120**, e2306420120 (2023).
https://doi.org:10.1073/pnas.2306420120
47   Hameed, P. S. *et al.* Triaminopyrimidine is a fast-killing and long-acting antimalarial
clinical candidate. *Nat Commun* **6**, 6715 (2015). https://doi.org:10.1038/ncomms7715
48   Reader, J. *et al.* Multistage and transmission-blocking targeted antimalarials discovered
from the open-source MMV Pandemic Response Box. *Nat Commun* **12**, 269 (2021).
https://doi.org:10.1038/s41467-020-20629-8
49   Meibalan, E. *et al.* Host erythrocyte environment influences the localization of exported
protein 2, an essential component of the Plasmodium translocon. *Eukaryot Cell* **14**, 371-
384 (2015). https://doi.org:10.1128/EC.00228-14
50   Ling, L. *et al.* Genetic ablation of the mitoribosome in the malaria parasite Plasmodium
falciparum sensitizes it to antimalarials that target mitochondrial functions. *J Biol Chem*
295, 7235-7248 (2020). https://doi.org:10.1074/jbc.RA120.012646
51   Ke, H., Dass, S., Morrisey, J. M., Mather, M. W. & Vaidya, A. B. The mitochondrial
ribosomal protein L13 is critical for the structural and functional integrity of the
mitochondrion in Plasmodium falciparum. *J Biol Chem* **293**, 8128-8137 (2018).
https://doi.org:10.1074/jbc.RA118.002552

A

B

Lanes 1, 4, 7: V$_1$A transgenic DNA
Lanes 2, 5, 8: WT DNA
Lanes 3, 6, 9: No DNA controls

C

Figure 1. Endogenous tagging of V$_1$A via CRISPR/Cas9 for conditional expression

Figure 2. Localization of V_1A throughout the asexual blood stages

Figure 3. Localization of V_1A on the secretory organelles in mature schizont stage parasites

Figure 4. Essentiality of V-type ATPase in the asexual blood stages

Figure 5. V-type ATPase is essential for hemoglobin digestion in the digestive vacuole

Figure 6. V-type ATPase regulates cytosolic pH in trophozoite stage parasites

A

B

Figure 7. Viability of the V-type ATPase knockdown parasites

Figure 8. Effects of Bafilomycin A1 on parasite egress and invasion

Figure 9. The stability of V-type ATPase complex under various conditions