Phylogeny and divergence time estimation of Io moths and relatives (Lepidoptera: Saturniidae: *Automeris*)

Authors

Chelsea Skojec[1,2], Chandra Earl[3], Christian D. Couch[1,2], Akito Y. Kawahara[1,2]

Affiliations

1. McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of Florida, Gainesville, FL 32611 USA
2. Department of Biology, University of Florida, Gainesville, FL, 32611 USA
3. Bernice Pauahi Bishop Museum, Honolulu, HI, 96817 USA

Keywords

anti-predation, dating, deimatism, evolution, eyespots, moth, museums

Abstract

The saturniid moth genus *Automeris* includes 145 described species. Their geographic distribution ranges from the eastern half of North America to as far south as Peru. *Automeris* moths are cryptically colored, with forewings that resemble dead leaves, and conspicuously colored, elaborate eyespots hidden on their hindwings. Despite their charismatic nature, the evolutionary history and relationships within *Automeris* and between closely related genera, remain poorly understood. In this study, we present the most comprehensive phylogeny of *Automeris* to date, including 80 of the 145 described species. We also incorporate two morphologically similar hemileucine genera, *Pseudautomeris* and *Leucanella*, as well as a morphologically distinct genus, *Molippa*. We obtained DNA data from both dry-pinned and ethanol-stored museum specimens and conducted Anchored Hybrid Enrichment (AHE) sequencing to reconstruct a high-quality dataset for phylogenetic analysis. The resulting phylogeny supports *Automeris* as a paraphyletic genus, with *Leucanella* and *Pseudautomeris* nested within, with the most recent common ancestor dating back to 26 mya. This study lays the foundation for future research on various aspects of *Automeris* biology, including anti-predator defense mechanisms, ecological adaptations, geographical distribution patterns, and potential drivers of speciation.

Introduction

Saturniidae, commonly known as wild silkmoths, are renowned for their remarkable morphological features, such as large body size, vibrant wing patterns, and distinctive hindwing markings. According to the checklist (Kitching et al. 2018), this family comprises a staggering 3,454 validated species, divided into eight subfamilies and 180 genera. The genus *Automeris*

(subfamily Hemileucinae) is the most diverse genus within the family. These species can be found across a vast geographic range, from North America to Peru. They thrive in diverse biotopes, spanning from tropical rainforests to arid habitats, in a wide range of altitudinal levels, from the sea level up to 4000 m (Deacaens and Herbin 2002).

One distinctive feature of almost all *Automeris* species, as well as closely related genera, is their recognizable wing ornamentation on both the fore- and hindwings. Forewings typically exhibit drab coloration and a cryptic leaf-like pattern. In stark contrast, hindwings have vibrant colors and feature an eyespot. Eyespots on the hindwings of *Automeris* moths make up the conspicuous coloration component of a deimatic display– thought to deter possible predators by startling, frightening, or confusing them, causing predators to pause or abandon their pursuit (Umbers and Mappes, 2016; Drinkwater et al. 2022). The presence of eyespots in most species of *Automeris* suggests they confer a selective benefit against predation (Olofsson et al. 2013), although eyespots of a few species are minimized or vestigial, which may be a secondary evolutionary loss. There has been extensive research on molecular mechanisms of eyespot development, primarily in butterflies (Monteiro 2013; Nijhout 2017; Matsuoka and Monteiro 2021) and some studies have examined the evolutionary origins of eyespots in a few representative species of Lepidoptera (Kodandaramaiah 2011; Monteiro 2015; Beldade 2021) but research into the evolution of eyespots in *Automeris* and relatives has yet to be conducted. The evolution of eyespot size and shape is believed to be driven by adaptive pressures instead of developmental constraints, thus they are not likely to constrain adaptive radiation of size, shapes, and patterns of eyespots (Beldade et al. 2002; Evans and Marcus 2006).

Historically, researchers have heavily relied on wing pattern variation to establish systematic hypotheses and determine relationships within *Automeris* (Tuskes and McElfresh, 1995; Tuskes et al. 1996), with the assumption that species most similar in appearance are most closely related. In his extensive study of the subfamily Hemileucinae, Lemaire (2002) identified 135 species within *Automeris*, classified into nine species groups based on their physical appearance and genitalia morphology. Despite being a charismatic and extremely diverse genus, knowledge regarding the evolutionary relationships between *Automeris* species, closely related genera, and the timing of the group's divergence from other hemileucine lineages remains limited. While some studies have included *Automeris* and other hemileucine species in large-scale phylogenetic analyses of Lepidoptera (Kitching et al. 2018, Kawahara et al. 2019), a comprehensive phylogenetic analysis focused on *Automeris* is yet to be conducted.

In this study we present the most comprehensive *Automeris* phylogeny that includes 80 of the 135 described species and 6 subspecies (Lemaire 2002). Additionally, we include two morphologically similar hemileucine genera - *Pseudautomeris* and *Leucanella*, and a morphologically distinct hemileucine genus, *Molippa,* to investigate relationships of *Automeris* with these genera. By constructing a well-represented phylogeny, we aim to uncover the diversification patterns within the genus and shed light on the timing of its evolutionary divergence from other lineages. Furthermore, a detailed phylogeny will provide a framework for future studies on various aspects of *Automeris* biology, including their anti-predatory defense, ecological adaptations, geographical distribution patterns, and potential drivers of speciation.

Materials and Methods

Taxon sampling and extraction

We sampled and extracted all previously unsequenced species and subspecies of *Automeris* (85), *Leucanella* (4), *Pseudautomeris* (6), and *Molippa* (2) available in the McGuire Center for Lepidoptera and Biodiversity (MGCL), Florida Museum of Natural History (FLMNH). Specimens were obtained from two collection types: dry, pinned specimens and those stored in ethanol at -80°C, which have been wing-vouchered following Cho et al. (2016) and are specifically stored for use in molecular studies.

One specimen per taxon was selected from the collections. Pinned specimens were carefully selected based on their condition, focusing on those with intact wings and body parts. Priority was given to more recently collected specimens during the selection process to enhance the likelihood of successful DNA extraction. Two legs were pulled from each sample and placed in a 1.5 mL microcentrifuge tube. Corresponding identification labels for samples were recorded on the respective tubes, linking them to those collection data present in the MGCL specimen database. Similarly, molecular collection samples were taken from their respective cold storage boxes, and partial thoracic tissue was dissected and placed in a labeled 1.5 mL microcentrifuge tube, and all relevant information was recorded in the collective database. The extracted thorax tissue was then placed in a labeled 1.5 mL microcentrifuge tube, and all relevant information was recorded in a collective database. Determination of DNA extraction from legs or thorax tissue was based on availability within the MGCL pinned and molecular collection. Due to the likelihood of higher DNA yield, specimens from the molecular collection were given priority. If specimens were available in both molecular and pinned collections, specimens were chosen from the molecular collection.

Extractions were completed following the specified protocol of the Qiagen DNeasy Blood and Tissue kit (Qiagen, Hilden, Germany). For quality control, extracted DNA was checked for concentration and fragmentation with a Qubit 2.0 fluorometer and electrophoresis gels (Fisherbrand Electrophoresis Power Supplies, FB200). After DNA extraction, if the desired concentration of DNA was not achieved, samples were re-extracted if tissue was still available. Otherwise, species with multiple samples were re-extracted with a different individual. Once the quantity of DNA was determined (8 ng/uL minimum and 100 ng/uL maximum), extracted samples were sent for Anchored Hybrid Enrichment (AHE) library preparation at RAPiD Genomics in Gainesville, Florida, USA. Higher concentration samples were prioritized. Out of 97 extracted samples, 95 showed promising DNA concentration of ≥ 8 ng/uL. Library preparation, hybridization enrichment, and Illumina HiSeq 2500 sequencing (PE100) was carried out at RAPiD Genomics.

DNA sequencing and assembly

We used the "BOM1" Anchored Hybrid Enrichment probe set (Hamilton et al. 2019), which was developed to target 921 loci across Bombycoidea. We used this probe set because it has been proven effective in capturing sequence data from both dry-pinned and ethanol-preserved

specimens (Hamilton et al. 2019; Dowdy et al. 2020; Li et al. 2022) This probe kit includes 58 loci across 24 "legacy" Sanger sequencing-based genes (Regier et al. 2008), eight bombycoid vision-related loci, and 855 loci designated as the Lepidoptera Agilent Custom SureSelect Target Enrichment "LEP1" probe kit from Breinholt et al. (2018). Anchored Hybrid Enrichment is a targeted-sequencing methodology designed to capture hundreds of unique orthologous loci (i.e., single copy, phylogenetically-informative markers) from across the genome, for resolving both shallow and deep-level evolutionary relationships. Probes are designed to anchor in conserved regions that are flanked by variable regions randomly spread throughout the genome. This approach creates a diverse set of informative loci that include exons, introns, intergenic, and conserved regions of the genome (Lemmon et al. 2012).

Raw AHE sequences were assembled following the methods of Breinholt et al. (2018), which implements an Iterative Baited Assembly (IBA) approach. This involves using the original sequencing probes to identify matching raw reads, which are then assembled into a novel probe. The newly assembled probe is subsequently used as a query to match against the remaining raw reads, and the process is repeated iteratively until confident assemblies can no longer be obtained. The pipeline also checks for quality and cross contamination due to barcode leakage and removes paralogs. Resultant assemblies extend beyond the boundaries of the initial sequencing probe, thereby leading to the production of two distinct datasets - one comprising sequences solely from the probe region of the assembly (Probe dataset), and the other comprising sequences from the complete assembly, encompassing both the probe and outer flanking regions (PF dataset).

Because the IBA approach often resulted in multiple assembled sequences for each locus per specimen, sequences were aligned using MAFFT v7.245 (Katoh and Stanley 2013) and a 50% consensus generated using FASconCAT-G v1.02 (Kück and Longo 2014) with the '-c -c -c' command to result in one sequence per locus per specimen. To minimize the extent of missing data in the final concatenated dataset, loci that were only obtained from three or fewer species were excluded from the datasets. Loci were concatenated across all species into one supermatrix using FASconCAT-G with the '-s' command.

In addition to specimens newly sequenced with AHE (95 taxa), we also included 18 previously sequenced hemileucine taxa in our lab that were sequenced and assembled via the same described methodologies. Four outgroup species were chosen to provide secondary calibrations for the divergence time analysis and to provide a root for the tree. All outgroups chosen were genera included in the (Hamilton et al. 2019) phylogeny and all had available raw transcriptomes or genomes. We chose two outgroup taxa in Saturniidae, *Attacus atlas* and *Therinia lactucina*, one from Sphingidae, *Manduca sexta*, and one from Bombycidae, *Bombyx mori* (The International Silkworm Genome Consortium 2008; Breinholt et al. 2013; Kawahara and Breinholt 2014; Kanost 2016). The two transcriptomes (Saturniidae) and two genomes (*Bombyx* and *Manduca*) that we used for the outgroups were assembled to the BOM1 AHE probe regions using the methods described above. This enabled us to combine the transcriptome and genome datasets with our newly sequenced data that were also assembled to the BOM1 AHE probe regions.

Alistat v1.6 (Wong et al. 2020) was used to evaluate the completeness of the concatenated alignment for Probe and PF datasets. We created Probe only and PF datasets because PF datasets can sometimes yield greater robustness to phylogenetic analyses of AHE data (e.g., (Kawahara et al. 2018; Hamilton et al. 2019; Homziak et al. 2019; St. Laurent et al. 2021). Completeness scores (C-scores) were computed for each taxon (Cr), and those taxa with a C-score < 0.15 were removed to avoid specimens with poor capture quality. We also removed loci that were captured across ≤ 3 taxa to improve dataset quality.

Phylogenetic analysis

Phylogenetic inference was conducted on concatenated supermatrices using a maximum likelihood analysis with IQTREE v. 1.5 (Nguyen et al. 2015). We constructed two datasets, one with just the probe region, and another with both probe and PF regions. Both datasets were analyzed as nucleotides, and we determined the best substitution model and partitioning scheme using the command '-m MFP+MERGE' using ModelFinder (Lanfear et al. 2012) as implemented in IQTREE. The command '-B 1000 -bnni' was used to perform 1000 ultrafast bootstrap (UFBS) replicates, while optimizing each bootstrap tree using a hill-climbing nearest neighbor interchange (NNI) search to reduce the risk of overestimating branch supports. All trees were rooted to *Bombyx mori*. We refer to high node support as those with UFBS ≥ 95.

Divergence time estimation

We performed divergence time estimation in BEAST v2.6.2 (Bouckaert et al. 2019) using a fixed topology via the MultiMonophyleticConstraint parameter to increase precision and accuracy. The topology generated from the PF dataset in IQTREE was used, given the higher bootstrap values when compared to the Probe dataset. However, when running BEAST using the PF dataset for dating, the resulting 95% confidence intervals were disproportionately small, indicating the high likelihood that the large dataset caused the underlying Bayesian analysis to become easily stuck at a local minima (Kawahara et al. 2019; Rougerie et al. 2022). Due to this, we ultimately chose to use the Probe dataset for dating.

Among insects, there are disproportionately few Lepidoptera fossils (Labandeira and Sepkoski 1993). Therefore, our tree was calibrated using fixed dates obtained from Kawahara et al. (2019). We used four secondary calibration points with uniform distributions to constrain the most recent common ancestors (MRCAs) of Hemileucinae and each chosen outgroup taxon. We used the Yule model of speciation (Yule 1924) as the tree prior, applied a strict clock rate, and ran the analyses for 10 million generations, sampling every 1,000 generations. Runs were assessed using TRACER v1.6.0 (Rambaut and Drummond 2013) to examine convergence, and a tree with mean heights was created from 9,000 trees after discarding a burn-in of 1,000 trees. All pipeline steps and phylogenomic analyses were conducted on the University of Florida HiPerGator high-performance computing cluster (http://www.hpc.ufl.edu/).

Results and Discussion

We sampled 85 species and subspecies of *Automeris*, two species of *Molippa*, four species of *Leucanella*, and six species of *Pseudautomeris*, resulting in a total of 97 sampled specimens. Of the 97, 74 were dry-pinned specimens and 23 were ethanol-stored specimens. The resulting DNA concentration range was 0.158-133 ng/uL. Samples that passed the DNA concentration cutoff (>8 ng/µL) were included in further analyses and sequencing.

In total, 95 newly sequenced samples had sufficient data for sequence assembly. We also supplemented these sequences with 22 sequences that were sequenced and assembled prior to this project. These included 18 ingroup sequences and 4 outgroups, resulting in a total of 117 taxa that were included in the beginning data matrix. After removing low quality sequences from each dataset, the Probe dataset contained 589 loci across 150,646 nucleotide sites, covering 105 taxa and the PF dataset contained 906 loci across 106 taxa. The average length of each locus, as well as the percentage of missing data, were observed to be 256 nucleotides and 25% for the Probe dataset, and 2,236 nucleotides and 84% for the PF dataset, respectively. The high missing data percentage of the PF dataset is due to the nature of the flanking regions. These regions differ vastly among taxa, since they extend as far as possible across unconserved regions, and the terminus varies depending on the coverage and quality of raw reads.

Both probe and PF topologies support the monophyly and general placement of genera *Molippa*, *Leucanella* and *Pseudautomeris* and the paraphyly of *Automeris*. All taxa represented by multiple subspecies were recovered as monophyletic with strong support. Considering the methodology employed and the higher support values, we favor the phylogeny derived from the PF dataset (Fig. 1). However, it is important to note that both phylogenies exhibit nearly identical topologies, differing only in a few relationships at the tips while maintaining the same backbone relationships.

Figure 1. Time calibrated tree showing relationships of *Automeris species and relatives, Leucanella, Pseudautomeris, Molippa* and outgroups. Blue circles represent nodes where UFBoot ≥ 90, though all are above 95. Red circles represent nodes where UFBoot ≤ 69. Divergence dates with 95% confidence intervals for each labeled node are in **Table 1**. Asterisks (*) notate pictured species.

Recently, Rougerie et al. (2022) reconstructed a phylogeny of Saturniidae, which was the only modern analysis providing a hypothesis of relationships of *Automeris*, *Leucanella, Pseudoautomeris* and *Molippa*. We find general congruence with the relationships uncovered by Rougerie et al. (2022), although *Automeris* is paraphyletic our study contained more species of *Automeris* included in the analysis. There was strong support in our study for a monophyletic *Leucanella + Pseudautomeris*, which splits *Automeris* into two distinct groups - a larger clade (*Automeris* Clade A) branching around 21 million years ago (mya) and a smaller clade, including *Automeris io* (*io* Clade B), branching earlier at approximately 23 mya (see Table 1). Notably,

none of the species within the smaller clade were included in the analysis by Rougerie et al. (2022). However, in both analyses, the relationships between the species included in both studies and *Leucanella*, *Molippa,* and *Pseudautomeris* were found to be consistent. We also found *Molippa* to be monophyletic, emerging as the sister group (UFBS = 100) to the remaining Hemileucines. This placement suggests that the most recent common ancestor of this clade (*Automeris + Leucanella + Molippa + Pseudautomeris*) dates to approximately 26 mya. This timeframe aligns closely with the findings of Rougerie et al. (2022), who estimated a divergence of approximately 22 mya. It should be noted that our analyses specifically targeted four genera within a subfamily characterized by significant diversity, species complexes and apparent paraphyly. However, to gain a comprehensive understanding of relationships of these genera within and among the rest of the subfamily, broader sampling would be beneficial.

Table 1: Major clade confidence intervals. Minimum 95% confidence interval (CI), median, and maximum 95% CI ages of major clades and outgroup taxa in Millions Years Ago (mya). Node numbers correspond to those in Figure 1.

	Node Number	Minimum - 95%	Median	Maximum - 95%
Automeris **Clade A**	1	0.202	0.2089	0.2163
Pseudautomeris	2	0.1627	0.1685	0.1747
Leucanella	2	0.1627	0.1685	0.1747
io **Clade B**	3	0.2167	0.2241	0.232
Molippa	4	0.2487	0.2576	0.2665
Attacus atlas	5	04759	0.4917	0.508
Therinia lactucina	6	0.5686	0.5864	0.6035
Manduca sexta	7	0.6606	0.6822	0.7053
Bombyx mori	8	0.6607	0.6825	0.7056

Eyespots of many *Leucanella* and *Pseudautomeris* are elaborate in their shape and color, and we hypothesized that these two genera are more closely related to *Automeris* than *Molippa*, as the latter have smaller, drab, potentially vestigial eyespots, or lack eyespots entirely. Our tree revealed that both *Leucanella* and *Pseudautomeris* are nested within *Automeris* with strong branch support. We postulate that predation pressure in the Neotropics drove the diversification

and complexity of eyespots in species in this clade. Like many effective anti-predatory traits, eyespots in this group are likely under positive selection due to a selective benefit for survival. Diversification and complexity of eyespots may have been driven by predation pressure, given the observed anti-predatory deimatic display of eyespots in many *Automeris* species. Previous research suggests that eyespots may be adaptive in some geographic regions, and maladaptive in others, which may explain secondary losses of eyespots in *Molippa* sp. (Kodandaramaiah 2011). There may be greater selective pressure on eyespots in the Neotropics than eyespots in North American species, driving the elaborate features and colors in these species (Janzen 1985).

Conclusion

Using Anchored Hybrid Enrichment techniques and developed analyses, we generated a robust phylogeny encompassing 106 taxa across *Automeris* and three closely related genera - *Leucanella*, *Pseudautomeris* and *Molippa*. This analysis reveals that *Leucanella* and *Pseudautomeris* are nested within *Automeris* with robust branch support, supporting the paraphyly of *Automeris* and suggesting close evolutionary relationships between these genera. Though this study helps clarify part of the complex Hemileucinae subfamily, a more complete sampling across species would provide greater understanding of the evolutionary patterns and processes that led to the larger diversification and evolutionary drivers of the subfamily. We hope this phylogeny will serve as a foundational framework for future investigations into the evolutionary dynamics and ecological adaptations of *Automeris* and its closely related genera. Future studies should focus on investigating eyespot trait morphology, to further clarify the diversification across species within the genus and sister groups

Data Accessibility

All assembled DNA sequences, alignments, phylogenies, and specimen metadata are available via a Dryad Deposition [here](). The final phylogeny is available on Open Tree of Life (https://tree.opentreeoflife.org/curator/study/view/ot_2240/).

Acknowledgements

We thank Nick Homziak for his comments and suggestions on the manuscript, Lawrence Reeves who took some of the images in Figure 1, and Amanda Markee for help with specimen extraction.

Literature Cited

1. Beldade P, Koops K, Brakefield PM. Developmental constraints versus flexibility in morphological evolution. Nature. 2002 Apr; 416(6883):844–7.   https://doi.org/10.1038/416844a.

 2. Beldade P, Monteiro A. Eco-evo-devo advances with butterfly eyespots. Current Opinion in Genetics & Development. 2021 Aug 1;69:6–13.   https://doi.org/10.1016/j.gde.2020.12.011.

3. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kuhnert D, De Maio N, Matschiner M, Mendes FK, Muller NF, Ogilvie HA, du Plessis L, Popinga A, Rambaut A, Rasmussen D, Siveroni I, Suchard MA, Wu C, Xie D, Zhang C, Stadler T, Drummond A. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLOS Computational Biology. 2019 Apr 8;15(4):e1006650. https://doi.org/10.1371/journal.pcbi.1006650.

4. Breinholt JW, Kawahara AY. Phylotranscriptomics: Saturated third codon positions radically influence the estimation of trees based on next-gen data. Genome Biology and Evolution. 2013 Nov 1;5(11):2082–92.   https://doi.org/10.1093/gbe/evt157.

5. Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. Resolving Relationships among the megadiverse butterflies and moths with a novel pipeline for anchored phylogenomics. Systematic Biology. 2018 Jan 1;67(1):78–93. https://doi.org/10.1093/sysbio/syx048.

6. Cho S, Epstein SW, Mitter K, Hamilton CA, Plotkin D, Mitter C, et al. Preserving and vouchering butterflies and moths for large-scale museum-based molecular research. PeerJ. 2016 Jun 22;4:e2160.   0.7717/peerj.2160

7. Decaens, T. and Herbin, D. Description of new *Automeris* species from Bolivia. *Zobodat - Literatur Artikel* 2002.   https://www.zobodat.at/publikation_articles.php?id=304493.

8. Dowdy NJ, Keating S, Lemmon AR, Lemmon EM, Conner WE, Scott Chialvo CH, et al. A deeper meaning for shallow-level phylogenomic studies: nested anchored hybrid enrichment offers great promise for resolving the tiger moth tree of life (Lepidoptera: Erebidae: Arctiinae). Systematic Entomology. 2020;45(4):874–93.   https://doi.org/10.1111/syen.12433

9. Drinkwater, E, Allen, WL, Endler JA, Hanlon RT, Holmes G, Homziak NT, Kang C, Leavell BC, Lehtonen J, Loeffler-Henry K, Ratcliffe JM, Rowe C, Ruxton GD, Sherratt TN, Skelhorn J,

Skojec C, Smart HR, White TE, Yack JE, Young CM, Umbers KDL. A synthesis of deimatic behaviour. Biological Reviews. 2022;97(6):2237–67. https://doi.org/10.1111/brv.12891.

10. Evans TM, Marcus JM. A simulation study of the genetic regulatory hierarchy for butterfly eyespot focus determination. Evolution & Development. 2006;8(3):273–83. https://doi.org/10.1111/j.1525-142X.2006.00098.x.

11. Hamilton CA, St Laurent RA, Dexter K, Kitching IJ, Breinholt JW, Zwick A, Timmermans MJTN, Barber JR, Kawahara AY. Phylogenomics resolves major relationships and reveals significant diversification rate shifts in the evolution of silk moths and relatives. BMC Evolutionary Biology. 2019 Sep 18;19(1):182. https://doi.org/10.1186/s12862-019-1505-1.

12. Homziak NT, Breinholt JW, Branham MA, Storer CG, Kawahara AY. Anchored hybrid enrichment phylogenomics resolves the backbone of erebine moths. Molecular Phylogenetics and Evolution. 2019 Feb 1;131:99–105. https://doi.org/10.1016/j.ympev.2018.10.038

13. Hossie TJ, Skelhorn J, Breinholt JW, Kawahara AY, Sherratt TN. Body size affects the evolution of eyespots in caterpillars. Proceedings of the National Academy of Sciences. 2015 May 26;112(21):6664–9. https://doi.org/10.1073/pnas.141512111

14. Janzen DH. On ecological fitting. Oikos. 1985 Dec;45(3):308. https://doi.org/10.2307/3565565.

15. Kanost MR, Arrese EL, Cao X, Chen YR, Chellapilla S, Goldsmith MR, Grosse-Wilde E, Heckel DG, Herndon N, Jiang H, Papanicolaou A. Multifaceted biological insights from a draft genome sequence of the tobacco hornworm moth, *Manduca sexta*. Insect biochemistry and molecular biology. 2016 Sep 1;76:118-47. https://doi.org/10.1016/j.ibmb.2016.07.005.

16. Katoh K, Standley DM. MAFFT Multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution. 2013 Apr 1;30(4):772–80. https://doi.org/10.1093/molbev/mst010.

17. Kawahara AY, Breinholt JW. Phylogenomics provides strong evidence for relationships of butterflies and moths. Proceedings of the Royal Society B: Biological Sciences. 2014 Aug 7;281(1788):20140970. https://doi.org/10.1098/rspb.2014.0970.

18. Kawahara AY, Breinholt JW, Espeland M, Storer C, Plotkin D, Dexter KM, et al. Phylogenetics of moth-like butterflies (Papilionoidea: Hedylidae) based on a new 13-locus

target capture probe set. Molecular Phylogenetics and Evolution. 2018 Oct 1;127:600–5. https://doi.org/10.1016/j.ympev.2018.06.002

19. Kawahara AY, Plotkin D, Espeland M, Meusemann K, Toussaint EF, Donath A, Gimnich F, Frandsen PB, Zwick A, Dos Reis M, Barber JR. Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths. Proceedings of the National Academy of Sciences. 2019 Nov 5;116(45):22657-63.   https://doi.org/10.1073/pnas.1907847116.

20. Kitching IJ, Rougerie R, Zwick A, Hamilton CA, St Laurent RA, Naumann S, Mejia LB, Kawahara AY. A global checklist of the Bombycoidea (Insecta: Lepidoptera). Biodiversity Data Journal. 2018(6).   https://doi.org/10.3897/BDJ.6.e22236.

21. Kodandaramaiah U. The evolutionary significance of butterfly eyespots. Behavioral Ecology. 2011 Nov 1;22(6):1264–71.   https://doi.org/10.1093/beheco/arr123.

22. Kück P, Longo GC. FASconCAT-G: Extensive functions for multiple sequence alignment preparations concerning phylogenetic studies. Frontiers in Zoology. 2014 Nov 18;11(1):81. https://doi.org/10.1186/s12983-014-0081-x.

23. Labandeira CC, Sepkoski JJ. Insect diversity in the fossil record. Science. 1993 Jul 16;261(5119):310–5.   https://doi.org/10.1126/science.11536548.

24. Lanfear R, Calcott B, Ho SYW, Guindon S. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. Molecular Biology and Evolution. 2012 Jun 1;29(6):1695–701.   https://doi.org/10.1093/molbev/mss020.

25. Lemaire, C. The Saturniidae of America - Hemileucinae. Goecke & Evers; 2002.

26. Lemmon AR, Emme SA, Lemmon EM. Anchored hybrid enrichment for massively high-throughput phylogenomics. Systematic Biology. 2012 Oct 1;61(5):727–44. https://doi.org/10.1093/sysbio/sys049.

27. Li, X., Hamilton, C.A., St Laurent, R., Ballesteros-Mejia, L., Markee, A., Haxaire, J., Rougerie, R., Kitching, I.J. and Kawahara, A.Y., 2022. A diversification relay race from Caribbean-Mesoamerica to the Andes: historical biogeography of Xylophanes hawkmoths. Proceedings of the Royal Society B, 289 (1968) p.20212435. https://doi.org/10.1098/rspb.2021.2435

28. Matsuoka Y, Monteiro A. Hox genes are essential for the development of eyespots in *Bicyclus anynana* butterflies. Genetics [Internet]. 2021. Jan 1. 217(1). https://doi.org/10.1093/genetics/iyaa005.

29. Monteiro A. Origin, development, and evolution of butterfly eyespots. Annual Review of Entomology. 2015 Jan; 60(1):253–71. https://doi.org/10.1146/annurev-ento-010814-020942

30. Monteiro A, Chen B, Ramos DM, Oliver JC, Tong X, Guo M, et al. Distal-Less Regulates Eyespot Patterns and Melanization in Bicyclus Butterflies. Journal of Experimental Zoology Part B: Molecular and Developmental Evolution. 2013;320(5):321–31. https://doi.org/10.1002/jez.b.22503.

31. Nijhout HF. The common developmental origin of eyespots and parafocal elements and a new model mechanism for color pattern formation. In: Diversity and Evolution of Butterfly Wing Patterns. Springer, Singapore; 2017. p. 3–19.

32. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution. 2015:32(1):268–74. https://doi.org/10.1093/molbev/msu300.

33. Olofsson M, Løvlie H, Tibblin J, Jakobsson S, Wiklund C. Eyespot display in the peacock butterfly triggers antipredator behaviors in naïve adult fowl. Behavioral Ecology. 2013 Jan 1;24(1):305–10. https://doi.org/10.1093/beheco/ars167.

34. Rambaut A, Drummond AJ. Tracer v1. 5 http://beast.bio.ed.ac.uk. Tracer; 2013.

35. Regier JC, Cook CP, Mitter C, Hussey A (2008) A phylogenetic study of the 'bombycoid complex' (Lepidoptera) using five protein-coding nuclear genes, with comments on the problem of macrolepidopteran phylogeny. Systematic Entomology 33: 175–189. https://doi.org/10.1111/j.1365-3113.2007.00409.x

36. Rougerie R, Cruaud A, Arnal P, Ballesteros-Mejia L, Condamine FL, Decaëns T, et al. Phylogenomics illuminates the evolutionary history of wild silkmoths in space and time (Lepidoptera: Saturniidae) [Internet]. bioRxiv; 2022 [cited 2023 Jul 30]. p. 2022.03.29.486224. https://www.biorxiv.org/content/10.1101/2022.03.29.486224v2

37. St Laurent RA, Carvalho APS, Earl C, Kawahara AY. Food plant shifts drive the diversification of sack-bearer moths. The American Naturalist. 2021 Nov;198(5):E170–84. https://doi.org/10.1086/716661

38. The International Silkworm Genome Consortium. The genome of a lepidopteran model insect, the silkworm *Bombyx mori*. Insect Biochemistry and Molecular Biology. 2008 Dec 1;38(12):1036–45. https://doi.org/10.1016/j.ibmb.2008.11.004.

39. Tuskes PM, McElfresh S. The biology and distribution of *Hemileuca electra* (Saturniidae) populations in the United States and Mexico, with descriptions of two new subspecies. Journal of the Lepidopterists Society. 1995;49(1):49–71.

40. Tuskes PM, Tuttle JP, Collins MM. The wild silk moths of North America: a natural history of the Saturniidae of the United States and Canada. Cornell University Press; 1996.

41. Umbers K, Mappes J. Towards a tractable working hypothesis for deimatic displays. Animal Behaviour. 2016 Feb 1;113. https://doi.org/10.1016/j.anbehav.2016.01.002.

42. Wong TKF, Kalyaanamoorthy S, Meusemann K, Yeates DK, Misof B, Jermiin LS. A minimum reporting standard for multiple sequence alignments. NAR Genomics and Bioinformatics. 2020 Jun 1;2(2):lqaa024. https://doi.org/10.1093/nargab/lqaa024.

43. Yule GU, A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis, F. R. S | Philosophical Transactions of the Royal Society of London. Series B, Containing Papers of a Biological Character [Internet]. [cited 2023 May 31]. https://royalsocietypublishing.org/doi/abs/10.1098/rstb.1925.0002