Unique T cell signatures are associated with reduced *Chlamydia trachomatis* reinfection

in a highly exposed cohort

Kacy S Yount[1], Avinash Kollipara[1], Chuwen Liu[2], Xiaojing Zheng[1,2], Catherine M O'Connell[1],

Bruce Bagwell[3], Harold C Wiesenfeld[4], Sharon L Hillier[4], Toni Darville[1,5]

[1]Department of Pediatrics, University of North Carolina at Chapel Hill, Chapel Hill, NC;

[2]Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC; [3]Verity

Software House, Topsham, Maine; [4]The University of Pittsburgh School of Medicine and the

Magee-Womens Research Institute, Pittsburgh, PA; [5]Department of Microbiology and

Immunology, University of North Carolina at Chapel Hill, Chapel Hill, NC

Corresponding Author:

Toni Darville

lad@email.unc.edu

501-944-3130

4008B Mary Ellen Jones, 116 Manning Drive, Chapel Hill, NC, 27599

The authors have declared that no conflict of interest exists.

ABSTRACT

Chlamydia trachomatis (CT) is the most common bacterial sexually transmitted infection (STI) in

the United States, despite effective antibiotics. Information regarding natural immunity to CT will

inform vaccine design. The objectives of this study were to determine immune cell populations

and functional features associated with reduced risk of CT reinfection or endometrial CT infection.

PBMCs were collected from a cohort of CT-exposed women who were tested for CT and other

STIs at the cervix and endometrium (to determine ascension) and were repeatedly tested over

the course of a year (to determine reinfection). Mass cytometry identified major immune

populations and T cell subsets. Women with CT had increased CD4+ effector memory T cells

(TEM) compared to uninfected women. Specifically, Th2, Th17, and Th17 DN CD4+ TEM were

increased. Th17 and Th17 DN CD4+ central memory T cells (TCM) were increased in women

who did not experience follow-up CT infection, suggesting that these cells may be important for

protection. These data indicate that peripheral T cells display distinct features that correlate with

natural immunity to CT and suggest that the highly plastic Th17 lineage plays a role in protection

against reinfection.

INTRODUCTION

Chlamydia trachomatis (CT) is the most common bacterial sexually transmitted infection (STI) globally (1). In women, CT infects the cervix and can ascend to the uterus and oviducts to cause pelvic inflammatory disease that can result in pelvic pain, infertility, and ectopic pregnancy (2). Although antibiotic treatment is effective, 80% of infections are asymptomatic and may go undetected (3), contributing to the risk for chronic sequelae (4).

Although repeated CT infections are common, they elicit partially protective immunity. Women who spontaneously cleared infection without treatment were less likely to become reinfected (5) and persons with confirmed prior infection have lower chlamydial loads (6). Determining the aspects of natural immunity that drive protection against CT will advance understanding of chlamydial immunity and aid design of an effective vaccine. Severe combined immune deficient (SCID), RAG-1-/-, TCRα-/-, CD4-/-, and MHCII-/- mice display delayed or failed resolution of *Chlamydia* infection, while MHCI-deficient (B2M-/-), B cell-deficient (µMT), and CD8-/- mice resolve infection with comparable kinetics to wild-type (7-11). These data emphasize the importance of CD4 T cell mediated immunity in a murine model of severe genital tract disease.

The T Cell Response Against Chlamydia (TRAC) cohort, a unique resource containing clinical samples and information from women highly exposed to CT and other STIs, was established to improve understanding of human natural immunity to CT (12). We observed that CD4 T cells producing CT-specific IFNγ were associated with protection from reinfection (13), consistent with findings of others (14), and we additionally determined that CD8 T cells producing CT-specific IFNγ were associated with protection from infections that ascended to the endometrium (13). Further, we identified a single nucleotide polymorphism in the gene for CD151, a tetrospanin molecule important for immune cell adhesion and migration and T cell proliferation, as a candidate genetic marker for reduced susceptibility for endometrial CT infection (15). Examining the chlamydial-specific responses of TRAC participants, we determined that anti-CT

antibodies are insufficient to protect women from ascending or recurrent infections (16, 17). TRAC

was specifically designed to characterize two important clinical outcomes of CT infection: (1)

distinguishing lower tract infection from ascended upper tract infection, which has direct

implications for disease severity, and (2) rates of reinfection, determined by diagnostic STI testing

and self-reports at 1-, 4-, 8-, and 12-months post-enrollment. In summary, these data indicate that

CD4 T cells, and to a lesser extent CD8 T cells, but not antibody, contribute to protection against

CT reinfection or ascension. The objective of this study was to further probe specific, cell-

mediated features of natural protective chlamydial immunity by using mass cytometry to

immunophenotype peripheral blood from TRAC participants for whom reinfection and ascension

status were defined.

Mass cytometry enabled inclusion of sufficient markers to determine broad populations of

immune cells while simultaneously focusing on CD4 and CD8 T cell subsets with distinct

functions, such as T follicular helper (Tfh) cells, regulatory T (Treg) cells, central memory T (TCM)

cells, effector memory T (TEM) cells, and stem-cell like memory T (TSCM) cells. TEM and TCM

T-helper cell subsets were defined by surface expression of chemokine receptors CCR6, CXCR3,

and CCR4 (18-20). Th1 cells (CCR6-CXCR3+CCR4-) produce mainly IFNγ and are important for

macrophage activation, phagocytosis, and bacterial defense, while Th2 cells (CCR6-CXCR3-

CCR4+) produce mainly IL-4, IL-5, and IL-13 and are important for supporting antibody production

(21). Th17 cells are a highly plastic family. Classical Th17 cells (CCR6+CXCR3-CCR4+) produce

primarily IL-17 and are important for neutrophil recruitment and defense at mucosal sites (22);

Th1/17 cells (CCR6+CXCR3+CCR4-) share Th1 and Th17 functionality, respond to IL-12 and IL-

23, and produce mostly IFNγ (23); Th17 double positive (DP) cells (CCR6+CXCR3+CCR4+)

produce IFNγ and low levels of IL-17 (19), and double negative (DN) cells (CCR6+CXCR3-CCR4-

) preferentially produce IL-17 and express transcriptional markers of early Th17 development (IL-

17F, STAT3), lymph-node homing (CCR7, CD62L), follicular help (CXCR5, BCL6, ASCL2), and

self-renewal (LEFI, MYC, TERC) (19, 24). We observed increased frequencies of CD4+ memory

Th17 and Th17 DN cells in participants with active CT infection and determined that these cells

were additionally enriched in participants protected from recurrent infection during the year they

were enrolled in the study.

RESULTS

Overall participant characteristics.

This study acquired CyTOF data from PBMCs of 82 of the 246 participants enrolled in

TRAC (12). The sociodemographic characteristics and self-reported STI diagnoses, and exposure

histories of this sub-cohort resemble those previously described for the full cohort (Supplemental

Table 1, Supplemental Table 2) (12).

CT genital tract infection elicits monocytes, plasma cells, and CD4 T effector memory cells

in the periphery.

Study participants were classified by their infection status at enrollment: Uninfected (CT-

;GC/MG-), CT monoinfection (CT+;GC/MG-), or CT coinfection with *Neisseria gonorrheoae* (GC)

and/or *Mycoplasma genitalium* (MG) (CT+;GC/MG+), and their PBMCs, collected at enrollment

and 1 month post treatment (1M), were immunophenotyped by CyTOF using a 33-marker panel

(Fig 1A, Supplemental Table 3). The manual gating hierarchy and definitions used to determine

frequencies of each cell type are outlined in Supplemental Table 4. Frequencies of total myeloid

cells were increased in CT+ women at enrollment and remained elevated at 1M, regardless of

coinfection (Fig 1B). Frequencies of total B cells, CD4 T cells, and CD8 T cells were similar across

all groups (Fig 1C-E). While dendritic cells (myeloid and plasmacytoid) and NK cell frequencies

were similar across patient subgroups (data not shown), classical, intermediate, and non-classical

monocytes were increased in CT+ women at enrollment and 1M, regardless of coinfection with

significant increases noted in non-classical monocytes (Fig 1F-H). Interestingly, frequencies of

plasma cells were increased in CT+;GC/MG- participants, but not CT+;GC/MG+ participants, at

enrollment and 1M (Fig 1I). Previous studies have shown that CT drives a strong and specific

antibody response (12, 25), which would be supported by a high frequency of plasma cells.

Frequencies of CD4 TEM cells were significantly increased in CT+ women at enrollment and 1M,

regardless of coinfection (Fig 1J). CD8 TEM were slightly increased in CT+ women (Fig 1K),

although to a lesser extent than CD4 TEM. In contrast, CD4 and CD8 TCM frequencies were

similar or slightly reduced in CT+ women compared to uninfected (Fig 1L, M). Frequencies for

populations defined in Fig 1A but not shown individually were not different between groups (data

not shown). Overall, these data demonstrate that CT infection drives increased frequencies of

peripheral monocytes, plasma cells, and CD4 effector memory cells, responses that remain

elevated for at least one month after treatment with antibiotics.

Participants with CT have increased peripheral frequencies of long-lived Th2 and Th17-

lineage CD4 TEM cells compared to uninfected participants.

It is broadly accepted that CD4 TEM, and specifically IFNγ-producing Th1 cells, are the

primary contributors to protection against CT in animals and humans (13, 26). Therefore, the

CyTOF antibody panel was designed with an investigational focus on T-cell subsets. Initial

investigation of major cell populations revealed a significant increase in peripheral CD4 TEM cells

in participants with CT (Fig 1J). Expression of the chemokine receptors CCR6, CXCR3, and CCR4

further characterized CD4 and CD8 T-helper subtypes of peripheral TEM (Fig 2A) (18-20).

Frequencies of CD4 TEM Th1 cells were not different between CT+ and uninfected individuals

(Fig 2B). Frequencies of CD4 TEM Th2 cells were markedly increased in CT+ participants at

enrollment and 1M, regardless of coinfection (Fig 2C), which is consistent with strong antibody

responses detected in response to CT (12, 25). Frequencies of CD4 TEM Th17 and Th17 DN

cells were increased in CT+;CT/MG- participants at enrollment and 1M, whereas there was no

difference in frequencies of Th1/17 or Th17 DP cells (Fig 2D-G). These differences were

maintained in CT+;GC/MG+ participants for CD4 TEM Th2 but not Th17 lineage cells (Fig 2C-F).

We similarly applied chemokine receptor definitions to investigate cells comprising the

CD8 memory compartment (Tc1, Tc2, Tc17, etc.). Frequencies of CD8 TEM Tc1 and Tc2 were

significantly increased in CT+;GC/MG- participants compared to uninfected (Fig 2H-I). There was

no change in CD8 TEM Tc1/17 (Fig 2J), but Tc17 DN cells trended toward an increase in

CT+;GC/MG- participants (Fig 2K), both of which parallel the results for CD4 TEM subsets

described above. CD8 TEM Tc17 and Tc17 DP were not detected. There were no differences in

either CD4 or CD8 TCM subtypes (data not shown), which is not surprising considering that

overall frequencies of CD4 and CD8 TCM were unchanged (Fig 1L-M). Frequencies of Treg and

Tfh cells were also determined but were not different between uninfected and CT+ participants

(data not shown). Overall, these results suggest that CT infection drives a peripheral adaptive

immune response dominated by CD4 and CD8 TEM Th2/Tc2 and CD4 TEM Th17-lineage cells.

Frequencies of CD4 TSCM are enriched in the peripheral blood of participants with

endometrial CT infection.

All participants were tested for CT, GC, and MG infection at cervical and endometrial sites

at enrollment, enabling bisection of the CT+;GC/MG- and CT+;GC/MG+ groups into CT infections

that had ascended to the upper genital tract (Endo+) and CT infections that were limited to the

cervix (Endo-). All major populations and T cell subsets were defined as above (Supplemental

Table 4), and comparisons of population frequencies performed between Endo+ and Endo-

participants. Women who were CT+;GC/MG- and Endo+ exhibited significantly increased

frequencies of CD4 TSCM, which was not observed in CT+;GC/MG+ women (Fig 3A). In

CT+;GC/MG- and Endo- participants, CD8 TEM and TCM Tc17 DN cells trended toward increase

(Fig 3C-D), suggesting expansion of these cells may contribute to protection from ascending CT

infection. Overall, however, there were comparatively minimal differences in peripheral blood T

cell populations between Endo+ and Endo- CT.

CD4 TCM Th1 and Th17 family subsets are enriched in peripheral blood of women without

documented CT reinfection.

Individuals treated for CT often experience repeat infections (6). Nevertheless, evidence

for acquired natural immunity includes documented increased resistance to CT reinfection in

women who previously spontaneously cleared an infection (14), and decreased CT infection rates

with increasing age after controlling for sexual activity (27). The TRAC study included a

longitudinal component designed to investigate attributes of the immune response contributing to

resistance to reinfection. Participants completed follow-up visits at 1, 4, 8, and 12 months where

they underwent diagnostic testing (cervix) and were asked to report any STI diagnoses that

occurred since their last visit. Using this information, we again bisected CT+;GC/MG- and

CT+;GC/MG+ groups into FU+ (positive for CT by diagnostic test or self-report at any follow-up

visit) and FU- (completed at least 3 out of 4 follow-up visits, and negative by diagnostic test and

self-report at all visits). Nine participants completed fewer than 3 follow-up visits and were

excluded from this analysis.

We compared frequencies of manually gated T cell populations between FU- and FU+

individuals. All cell populations described above (Supplemental Table 4) were defined but only

significant findings are presented in Figure 4. Frequencies of total CD4 and total CD8 T cells and

frequencies of CD4 and CD8 stem-cell like memory (TSCM) cells were enriched in FU-

participants without co-infection at enrollment and remained expanded 1 month post treatment

(Fig 4A-D). CD4 Treg cells were enriched in FU- participants at the enrollment visit (Fig 4E).

Interestingly, frequencies of CD4 TEM Th2 cells were increased in FU+ participants (Fig 4F),

suggesting that although these cells are induced by CT infection, they are not protective. In

contrast, frequencies of CD4 TCM Th1, Th17, Th1/17, and Th17 DN cells were increased in FU-

participants (Fig 4G-J). These data suggest that T cells, specifically CD4 and CD8 TSCM and

CD4 TCM Th1/Th17 lineage cells are important for developing natural immunity against incident

CT infection. Interestingly, differences in frequencies of T cell subsets between FU- and FU+

participants were only detected in CT+;GC/MG- and not CT+;GC/MG+ participants, which

suggests that immune responses specific to CT are diluted by coinfection.

DISCUSSION

Using samples from a uniquely characterized and clinically-relevant cohort, this study

immunophenotyped responses to CT infection with high dimensional analysis of surface marker

expression on PBMCs and found specific T cell populations correlated with reduced incidence of

CT reinfection. Overall, broad immune cell populations in peripheral blood were different between

uninfected and CT+ participants, but similar between CT monoinfection and coinfection. Notably,

CT+ participants had increased frequencies of mononuclear myeloid cells, consistent with an

inflammatory response to an acute infection. CT+ participants displayed increases in frequencies

of non-classical (CD14-CD16+) monocytes, important for antigen presentation, T cell proliferation,

patrolling tissues for signs of injury, and increased Fc receptor-mediated phagocytosis (28).

Frequencies of monocytes remained elevated in some samples of PBMCs collected one month

later from participants who were CT+ at enrollment, likely driven by ongoing inflammation induced

by danger associated molecular patterns released from bacteria and dying cells.

While frequencies of immune populations were largely similar in CT+ participants

regardless of coinfection, plasma cells were elevated in participants with CT monoinfection, but

their frequencies in CT+;GC/MG+ participants were similar to uninfected. This suggests that

plasma cells are driven by CT infection, consistent with well-documented robust but non-

protective antibody responses to CT infection (16, 29, 30), but that this response is suppressed

or impeded by coinfection with other STIs. *N. gonorrhoeae* opacity proteins bind

carcinoembryonic antigen-related cellular adhesion molecule 1 (CEACAM1 or CD66a) on B cells,

leading to inhibitory signals which promote B cell death (31). Plasma cells also express CD66a

(32). Thus, in women coinfected with GC, opacity protein interactions with CD66a could contribute

to reduced numbers of plasma cells, especially in women with chronic GC infection.

CD4 TEM frequencies were strikingly increased in CT+ participants compared to CD8

TEM frequencies. These data are consistent with our prior investigation of CT antigen specific

responses in a subset of TRAC participants where we observed higher levels of IFNγ production

and a greater breadth of recognition of CT antigens among CD4 T cells compared to CD8 T cells

(13). That CD8 TEM frequencies were not increased is also consistent with engaging the PD-1

receptor on CD8 T cells by antigen presenting cells inhibits generation of CT-specific memory

CD8 T cells during murine chlamydial infection (33). Frequencies of TEM but not TCM were

increased in CT+ compared to uninfected participants in our study. In contrast, an analysis of *C.*

pneumoniae-specific peripheral blood T cells in healthy donors found that CD4 TCM were

enriched compared to TEM (34). These contrasting data may be explained by the difference

between acute infection experienced by TRAC participants and relatively remote infection

experienced by individuals in the *C. pneumoniae* study.

Further investigation of CD4 TEM subsets revealed that CT+ participants displayed higher

frequencies of peripheral Th2, Th17, and Th17 DN cells, with unchanged frequencies of Th1 cells.

Th1, Th2, and Th17 cells support different aspects of the overall downstream immune response.

CD4 Th1 cells are classically defined by production of IFNγ and expression of CXCR3 and/or

CCR5 (18). They are well-documented contributors to clearance of *Chlamydia* and protection from

reinfection in mouse models (35-37) and associate with resistance to CT reinfection in humans

(13, 14). We did not find evidence for expansion of Th1 cells in the peripheral blood of CT+

participants compared to uninfected. The absence of such an important protective cell type in the

periphery may explain why CT reinfection is so common. However, Th1 cells express the

chemokine receptors CXCR3 and CCR5, which bind the chemokines CXCL10 and CCL5

(RANTES), respectively (38). We previously reported detecting both of these chemokines in

cervical secretions of TRAC participants (39), and they are highly expressed in genital tract

tissues of *C. muridarum* infected mice (37). Thus, it is possible that despite overall expansion of

Th1 cells, they were preferentially recruited to the site of infection and consequently not increased

in the periphery. In contrast, peripheral CD4 Th2 cells were significantly enriched in CT+ women,

consistent with the fact that Th2 responses are generally supportive of antibody production and

that CT infection is known to drive strong but non-protective antibody responses (6, 12, 16).

Th17 cells are a highly plastic family of T-helper cells. They are mainly defined by their

expression of CCR6 (40) but can gain or lose chemokine receptors and their associated effector

functions (e.g., cytokine production) (41). Th17 cells can be pathogenic, but they are important

contributors to protection against mucosal pathogens such as *Mycobacterium tuberculosis* and

Bordetella pertussis (42, 43). We used expression of CCR6, CXCR3, and CCR4 to define four

subsets of Th17 family cells: classical Th17, Th1/17, Th17 DP, and Th17 DN (19). Th17 DN cells

express transcriptional markers of early Th17 development (IL-17F, STAT3), lymph-node homing

(CCR7, CD62L), follicular help (CXCR5, BCL6, ASCL2), and self-renewal (LEFI, MYC, TERC)

(19), consistent with their highly plastic nature. We found increased frequencies of classical Th17

and Th17 DN cells in CT+ participants compared to uninfected. These findings, when considered

alongside classifications of whether participants experienced CT infection that ascended to the

endometrium (Endo+ vs Endo-) and whether CT reinfection was documented over the course of

one year of follow-up visits (FU+ vs FU-), provided important insights into specific responses that

may be protective. Future studies will examine Th17 family subsets to determine the functional

cytokine responses of Th17 and Th17 DN cells in cervical and endometrial tissues from CT-

infected women.

Peripheral immune cell populations were similar overall between participants with

endometrial CT infections and those that had CT infections limited to the cervix. Frequencies of

CD4 TSCM were significantly increased in Endo+ and FU- participants. This is consistent with our

prior finding that endometrial infection was associated with reduced risk of follow-up CT infection

(12), likely because the increased bacterial burden associated with endometrial infection drives a

more robust memory response. CD8 TEM Tc17 DN cells trended toward increased frequencies

in CT+;GC/MG- Endo- participants, consistent with our previous study showing that CT-specific

CD8 T cells weakly associated with reduced CT ascension (13). However, evaluation of immune

cell populations in the genital tract is necessary to define the specific features of immunity that

prevent or limit ascending infection. Future studies will investigate immune cells from cervical and

endometrial TRAC samples.

In contrast to the relatively minor differences identified between Endo+ and Endo-

participants, several T cell subtypes were found to be strongly associated with presence or

absence of recurrent CT infection. Our detection of increased frequencies of total CD4 T cells in

FU- participants at enrollment and maintenance of an increase one month post treatment is

indicative of developing natural immunity in some participants. We also detected increased

frequencies of CD4 and CD8 TSCM in FU- individuals. TSCM resemble naïve T cells in their

expression of CD45RA and CCR7, but they also express CD95, a marker of antigen experience

and memory (44). They are minimally differentiated memory cells that have a high propensity for

stemness and self-renewal as well as the multipotency to differentiate into a broad range of

memory T-helper phenotypes (44), which may explain why they are associated with FU-

outcomes, The importance of the quality of the T cell response is demonstrated by the contrasting

subsets that were expanded in FU- and FU+ participants. Expansion of TEM Th2 cells, which

were strikingly increased overall in CT+ participants, was associated with a FU+ outcome, while

frequencies of TCM Th1 cells, which were not different between CT+ and uninfected participants,

were associated with FU- status. An important role for Th17-lineage cells in protection from

reinfection is suggested by the striking increase in TCM Th17, along with increased frequencies

of TCM Th1/17 and TCM Th17 DN cells in FU- participants, again suggesting that Th17-family

cells may contribute to CT natural immunity.

The finding that Th1 and Th17 subsets were associated with reduced reinfection (and that

Th2 responses were not) provides the opportunity to design vaccines that induce more protective

responses than natural immunity. Different vaccine platforms and adjuvants can shape different

T-helper responses (45). CAF01, a Th1/Th17 adjuvant, combined with recombinant chlamydial

antigen elicited resident memory Th1/Th17 responses that increased protection of mice against

chlamydial challenge (46). Together, these findings support the design of vaccines that prioritize

CD4 Th1 and Th17-polarized responses to elicit the best protection against CT.

Our study represents a major advancement in the understanding of natural immunity to

CT in humans, but it is somewhat limited because we defined populations increased in CT+

participants compared to uninfected participants, but the cells were not restimulated with CT

antigens to confirm their CT-specificity. Immunophenotyping PBMCs directly without restimulation

allowed us to confidently define T cell subsets based on chemokine receptors without the risk of

downregulation upon restimulation. However, the important open question of whether Th17 and

Th17 DN cells produce CT-specific IL-17 and/or IFNγ remains to be addressed in future studies.

Another limitation of this study is that cell population frequencies could not be correlated with

endometrial inflammation and pelvic inflammatory disease. Many of the women participating in

TRAC selected contraceptive methods that promote thin uterine lining, making it difficult to acquire

sufficient tissue for evaluation of pathology. However, ongoing studies are investigating local T

cell responses in endometrial tissue sections from participants with sufficient tissue.

Overall, we found that frequencies of CD4 TEM, specifically Th2, Th17, and Th17 DN

subsets, were enriched in individuals with active CT infection. We also identified that increased

frequencies of CD4 TCM Th1, Th17, and Th17 were associated with individuals who had no

documented reinfection over the course of a year. Together, these findings increase our

understanding of natural immunity to CT and inform strategies for future vaccine design.

METHODS

Patient Populations.

A subset of a previously described T Cell Response Against Chlamydia (TRAC) cohort (12) was

used for this study.

Definition of Clinical Classifications

Uninfected, CT+;GC/MG-, and CT+;GC/MG+ Groups:

Nucleic acid amplification testing (NAAT) was used to test cervical swabs collected from study

particpants at enrollment for CT, *Neisseria gonorrhoeae* (gonococcus, GC), and *Mycoplasma*

genitalium (MG). Particpants whose swabs tested negatively for all 3 STIs at enrollment were

classified as "uninfected". Participants testing positively for CT but negative for GC and MG were

classified as "CT+;GC/MG-". Enrollees testing positively for CT and GC, MG, or both were

classified as "CT+;GC/MG+". Of 82 total TRAC participants studied, 12 were uninfected, 44 were

CT+;GC/MG-, and 26 were CT+;GC/MG+ (Supplemental Table 5).

Endometrial positive and endometrial negative CT:

Endometrial biopsies were also NAAT tested by to determine whether infection was limited to the

cervix or had ascended to the upper genital tract. Within CT+;GC/MG- and CT+;GC/MG+ groups,

enrollees testing positively for CT at the cervix but not the endometrial biopsy were classified as

Endo-, while those testing positively for CT at both sites were classified as Endo+. No participants

tested positive for CT solely in the endometrial sample. Of 44 CT+;GC/MG- participants, 19 were

Endo+ and 25 were Endo-. Of 26 CT+;GC/MG+ participants, 11 were Endo+ and 15 were Endo-

(Supplemental Table 5).

Follow-up CT positive and follow-up CT negative:

NAAT testing was performed on cervical brush samples at enrollment and at 1, 4, 8, and 12 month

follow-up visits. Participants with a positive cervical CT test at enrollment and a positive cervical

CT test or self-reported CT diagnosis from an outside clinic in the time since last visit at any follow-

up visit (regardless of how many visits they completed) were classified as follow-up positive (FU+).

Participants that tested positive for cervical CT at enrollment, completed at least 3 out of 4 follow-

up visits, tested negative for cervical CT at all follow-up visits, and self-reported no CT diagnosis

from an outside clinic in the time since last visit at any follow-up visit were classified as follow-up

negative (FU-). 9 participants attended less than 3 follow-up visits (FU N/A) and were excluded

from follow-up analysis. Of CT+;GC/MG- participants, 19 were FU-, 21 were FU+, and 4 were FU

N/A. Of CT+;GC/MG+ pariticpants, 9 were FU-, 12 were FU+, and 5 were FU N/A (Supplemental

Table 5). All uninfected samples were FU- and the group was therefore not bisected.

PBMC Processing

Whole blood samples from individual patients at each visit were diluted with equal volume of PBS

and peripheral blood mononuclear cells (PBMCs) were isolated via a density gradient using

lymphocyte separation media (Corning). Cells were frozen in 90% FBS with 10% DMSO and

stored in liquid nitrogen until thawed for CyTOF analysis.

Mass Cytometry

Mass cytometry (CyTOF) was performed on 156 total PBMC samples (Supplemental Table 5). A

mass cytometry panel comprising of 33 surface markers (Supplemental Table 3) was used to

profile immune cell populations in detail. Frozen PBMCs were thawed in a 37°C water bath, and

an equal amount of pre-warmed complete RPMI (cRPMI) with 50 units/mL benzonase (Sigma)

was added dropwise. Cells were then centrifuged at 300xg for 5 min at room temperature (RT).

Pellets were washed in cRPMI and enumerated using trypan blue and a hemocytometer. 3 x 10^6

viable participant cells and 1 x 10^6 PBMCs from a common negative control donor were

transferred to one FACS tube and washed with PBS (CyTOF grade, Rockland Immunochemicals

Inc.). Dead cells were stained with 1 µM Cell-ID Cisplatin 198-Pt (Fluidigm) for 5 min at RT and

washed twice with cell staining buffer (CSB, Fluidigm). An antibody cocktail comprising 33

antibodies (Supplemental Table 3) and human Fc block (eBioscience) in was added to cells in 50

µL staining volume and incubated for 30 min at RT. Cells were washed twice with CSB followed

by secondary antibody (Anti FITC-144Nd) incubation. Stained cells are fixed with 2% PFA in PBS

for 1 hr at RT followed by two washes in PBS. 1 mL of 1:2000 Cell-ID Intercalator-Ir-125 µM

(Fluidigm) solution in fix and perm buffer (Fluidigm) was added to the cells to stain the DNA and

incubated for 30 min at RT or 4°C overnight. Cells were further washed with CSB twice and

resuspended in CAS (Cell Acquisition Solution; Fluidigm). Cell concentrations were adjusted to

0.5 x 10^6 cells/mL in CAS with Four Element Calibration beads (Fluidigm). 200,000 cells per

sample were acquired on a Helios instrument (Fluidigm) at a flow rate of 250 cells/sec by the UNC

Mass Cytometry Core.

Data Analysis

Data Cleaning. Raw CyTOF files were cleaned using the Pathsetter algorithm (47). All samples

were of good quality and viability. FCS files containing cleaned viable events were created and

used for remaining analysis.

Manual Gating Phenotypic Analysis. All analysis was done in Cytobank (Beckman Coulter) and

FlowJo (BD) software. The manual gating strategy and surface marker designations selected for

immune cell phenotypes are summarized in Supplemental Table 4.

Statistics

Statistical calculations were all computed in R. For comparisons between uninfected,

CT+;GC/MG-, and CT+;GC/MG+, a non-parametric Kruskal-Wallis test followed by Dunn's test

and Bonferroni correction for multiple comparisons was used. For comparisons between Endo-

vs Endo+ and FU- vs FU+, a non-parametric Wilcoxon ranked sum test was used. $P < 0.05$ was

used to determine a significant result. Significance symbols used in figures are defined as:

*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Study Approval

The institutional review boards for human research of the University of Pittsburgh and the

University of North Carolina approved the study protocol. All participants provided written informed

consent at the time of enrollment and agreed to be contacted to return for follow-up visits 1, 4, 8,

and 12 months after enrollment.

AUTHOR CONTRIBUTIONS

KSY, AK, XZ, and TD designed research studies. KSY and AK conducted experiments. HCW and

SLH collected participant samples and performed diagnostic testing. KSY, CL, and BB analyzed

data. KSY, CMO and TD wrote the manuscript.

ACKNOWLEDGEMENTS

We thank the women who agreed to participate in this study; Ingrid Macio, Melinda Petrina, Carol

Priest, Abi Jett, and Lorna Rabe, for their efforts in the clinic and the microbiology laboratory; and

the staff at the Allegheny County Health Department STD Clinic. The UNC Mass Cytometry Core

is supported by the University Cancer Research Fund (UCRF) and UNC Cancer Center Core

Support Grant #P30CA016086. This work was supported by the National Institute of Allergy and

Infectious Diseases through U19AI144181 (TD). KSY is funded by UNC STI/HIV T32 postdoctoral

training grant AI007001.

BIBLIOGRAPHY

1. Newman L, Rowley J, Vander Hoorn S, Wijesooriya NS, Unemo M, Low N, et al. Global Estimates of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting. *PLoS One.* 2015;10(12):e0143304.
2. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, and Ness RB. Risk of sequelae after Chlamydia trachomatis genital infection in women. *J Infect Dis.* 2010;201 Suppl 2:S134-55.
3. Stamm WE. Chlamydia trachomatis infections: progress and problems. *JInfectDis.* 1999;179 Suppl 2:S380-S3.
4. Wiesenfeld HC, Hillier SL, Meyn LA, Amortegui AJ, and Sweet RL. Subclinical pelvic inflammatory disease and infertility. *Obstet Gynecol.* 2012;120(1):37-43.
5. Geisler WM, Lensing SY, Press CG, and Hook EW, 3rd. Spontaneous resolution of genital Chlamydia trachomatis infection in women and protection from reinfection. *The Journal of infectious diseases.* 2013;207(12):1850-6.
6. Batteiger BE, Tu W, Ofner S, Van Der Pol B, Stothard DR, Orr DP, et al. Repeated Chlamydia trachomatis genital infections in adolescent women. *J Infect Dis.* 2010;201(1):42-51.
7. Yang X, Hayglass KT, and Brunham RC. Different roles are played by alpha beta and gamma delta T cells in acquired immunity to Chlamydia trachomatis pulmonary infection. *Immunology.* 1998;94(4):469-75.
8. Morrison RP, Feilzer K, and Tumas DB. Gene knockout mice establish a primary protective role for major histocompatibility complex class II-restricted responses in Chlamydia trachomatis genital tract infection. *Infection and immunity.* 1995;63(12):4661-8.
9. Magee DM, Igietseme JU, Smith JG, Bleicker CA, Grubbs BG, Schachter J, et al. Chlamydia trachomatis pneumonia in the severe combined immunodeficiency (SCID) mouse. *Reg Immunol.* 1993;5(6):305-11.
10. Su H, Feilzer K, Caldwell HD, and Morrison RP. Chlamydia trachomatis genital tract infection of antibody-deficient gene knockout mice. *Infection and immunity.* 1997;65(6):1993-9.
11. Morrison SG, Su H, Caldwell HD, and Morrison RP. Immunity to murine Chlamydia trachomatis genital tract reinfection involves B cells and CD4(+) T cells but not CD8(+) T cells. *Infection and immunity.* 2000;68(12):6979-87.
12. Russell AN, Zheng X, O'Connell CM, Taylor BD, Wiesenfeld HC, Hillier SL, et al. Analysis of Factors Driving Incident and Ascending Infection and the Role of Serum Antibody in Chlamydia trachomatis Genital Tract Infection. *The Journal of infectious diseases.* 2016;213(4):523-31.
13. Russell AN, Zheng X, O'Connell CM, Wiesenfeld HC, Hillier SL, Taylor BD, et al. Identification of Chlamydia trachomatis Antigens Recognized by T Cells From Highly Exposed Women Who Limit or Resist Genital Tract Infection. *The Journal of infectious diseases.* 2016;214(12):1884-92.
14. Bakshi RK, Gupta K, Jordan SJ, Chi X, Lensing SY, Press CG, et al. An Adaptive Chlamydia trachomatis-Specific IFN-gamma-Producing CD4(+) T Cell Response Is Associated With Protection Against Chlamydia Reinfection in Women. *Front Immunol.* 2018;9:1981.
15. Zhong W, Kollipara A, Liu Y, Wang Y, O'Connell CM, Poston TB, et al. Genetic susceptibility loci for Chlamydia trachomatis endometrial infection influence expression of genes involved in T cell function, tryptophan metabolism and epithelial integrity. *Front Immunol.* 2022;13:1001255.

16. Darville T, Albritton HL, Zhong W, Dong L, O'Connell CM, Poston TB, et al. Anti-chlamydia
IgG and IgA are insufficient to prevent endometrial chlamydia infection in women, and
increased anti-chlamydia IgG is associated with enhanced risk for incident infection.
American journal of reproductive immunology (New York, NY : 1989). 2019;81(5):e13103.
17. Liu C, Hufnagel K, O'Connell CM, Goonetilleke N, Mokashi N, Waterboer T, et al. Reduced
Endometrial Ascension and Enhanced Reinfection Associated With Immunoglobulin G
Antibodies to Specific Chlamydia trachomatis Proteins in Women at Risk for Chlamydia.
The Journal of infectious diseases. 2022;225(5):846-55.
18. Bonecchi R, Bianchi G, Bordignon PP, D'Ambrosio D, Lang R, Borsatti A, et al. Differential
expression of chemokine receptors and chemotactic responsiveness of type 1 T helper
cells (Th1s) and Th2s. *J Exp Med.* 1998;187(1):129-34.
19. Wacleche VS, Goulet JP, Gosselin A, Monteiro P, Soudeyns H, Fromentin R, et al. New
insights into the heterogeneity of Th17 subsets contributing to HIV-1 persistence during
antiretroviral therapy. *Retrovirology.* 2016;13(1):59.
20. Mousset CM, Hobo W, Woestenenk R, Preijers F, Dolstra H, and van der Waart AB.
Comprehensive Phenotyping of T Cells Using Flow Cytometry. *Cytometry Part A.*
2019;95(6):647-54.
21. Spellberg B, and Edwards JE, Jr. Type 1/Type 2 Immunity in Infectious Diseases. *Clinical*
Infectious Diseases. 2001;32(1):76-102.
22. Khader SA, Gaffen SL, and Kolls JK. Th17 cells at the crossroads of innate and adaptive
immunity against infectious diseases at the mucosa. *Mucosal immunology.* 2009;2(5):403-
11.
23. Duhen T, and Campbell DJ. IL-1β promotes the differentiation of polyfunctional human
CCR6+CXCR3+ Th1/17 cells that are specific for pathogenic and commensal microbes.
Journal of immunology (Baltimore, Md : 1950). 2014;193(1):120-9.
24. Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia A, et
al. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper
memory cells. *Nature immunology.* 2007;8(6):639-46.
25. Rahman KS, Darville T, Russell AN, O'Connell CM, Wiesenfeld HC, Hillier SL, et al.
Comprehensive Molecular Serology of Human Chlamydia trachomatis Infections by
Peptide Enzyme-Linked Immunosorbent Assays. *mSphere.* 2018;3(4).
26. Johansson M, Schön K, Ward M, and Lycke N. Studies in knockout mice reveal that anti-
chlamydial protection requires TH1 cells producing IFN-gamma: is this true for humans?
Scand J Immunol. 1997;46(6):546-52.
27. Arno JN, Katz BP, McBride R, Carty GA, Batteiger BE, Caine VA, et al. Age and clinical
immunity to infections with Chlamydia trachomatis. *Sexually transmitted diseases.*
1994;21(1):47-52.
28. Sampath P, Moideen K, Ranganathan UD, and Bethunaickan R. Monocyte Subsets:
Phenotypes and Function in Tuberculosis Infection. *Front Immunol.* 2018;9:1726.
29. LaVerda D, Albanese LN, Ruther PE, Morrison SG, Morrison RP, Ault KA, et al.
Seroreactivity to Chlamydia trachomatis Hsp10 correlates with severity of human genital
tract disease. *Infection and immunity.* 2000;68(1):303-9.
30. El Hakim EA, Gordon UD, and Akande VA. The relationship between serum Chlamydia
antibody levels and severity of disease in infertile women with tubal damage. *Arch Gynecol*
Obstet. 2010;281(4):727-33.
31. Pantelic M, Kim YJ, Bolland S, Chen I, Shively J, and Chen T. Neisseria gonorrhoeae kills
carcinoembryonic antigen-related cellular adhesion molecule 1 (CD66a)-expressing
human B cells and inhibits antibody production. *Infection and immunity.* 2005;73(7):4171-
9.

32.   Lee C, Guinn BA, Brooks SE, Richardson D, and Orchard K. CD66a (CEACAM1) is the
521   only CD66 variant expressed on the surface of plasma cells in multiple myeloma: a refined
522   target for radiotherapy trials? *Br J Haematol.* 2010;149(5):795-6.
523   33.   Fankhauser SC, and Starnbach MN. PD-L1 limits the mucosal CD8+ T cell response to
524   Chlamydia trachomatis. *Journal of immunology (Baltimore, Md : 1950).* 2014;192(3):1079-
525   90.
526   34.   Bunk S, Schaffert H, Schmid B, Goletz C, Zeller S, Borisova M, et al. Chlamydia
527   pneumoniae-induced memory CD4+ T-cell activation in human peripheral blood correlates
528   with distinct antibody response patterns. *Clinical and vaccine immunology : CVI.*
529   2010;17(5):705-12.
530   35.   Olive AJ, Gondek DC, and Starnbach MN. CXCR3 and CCR5 are both required for T cell-
531   mediated protection against C. trachomatis infection in the murine genital mucosa.
532   *Mucosal immunology.* 2011;4(2):208-16.
533   36.   Poston TB, Qu Y, Girardi J, O'Connell CM, Frazer LC, Russell AN, et al. A Chlamydia-
534   Specific TCR-Transgenic Mouse Demonstrates Th1 Polyfunctionality with Enhanced
535   Effector Function. *Journal of immunology (Baltimore, Md : 1950).* 2017;199(8):2845-54.
536   37.   Belay T, Eko FO, Ananaba GA, Bowers S, Moore T, Lyn D, et al. Chemokine and
537   chemokine receptor dynamics during genital chlamydial infection. *Infection and immunity.*
538   2002;70(2):844-50.
539   38.   Cockwell P, Calderwood JW, Brooks CJ, Chakravorty SJ, and Savage CO.
540   Chemoattraction of T cells expressing CCR5, CXCR3 and CX3CR1 by proximal tubular
541   epithelial cell chemokines. *Nephrol Dial Transplant.* 2002;17(5):734-44.
542   39.   Poston TB, Lee DE, Darville T, Zhong W, Dong L, O'Connell CM, et al. Cervical Cytokines
543   Associated With Chlamydia trachomatis Susceptibility and Protection. *The Journal of
544   infectious diseases.* 2019;220(2):330-9.
545   40.   Liu H, and Rohowsky-Kochan C. Regulation of IL-17 in human CCR6+ effector memory T
546   cells. *Journal of immunology (Baltimore, Md : 1950).* 2008;180(12):7948-57.
547   41.   Muranski P, and Restifo NP. Essentials of Th17 cell commitment and plasticity. *Blood.*
548   2013;121(13):2402-14.
549   42.   Ogongo P, Tezera LB, Ardain A, Nhamoyebonde S, Ramsuran D, Singh A, et al. Tissue-
550   resident-like CD4+ T cells secreting IL-17 control Mycobacterium tuberculosis in the
551   human lung. *J Clin Invest.* 2021;131(10).
552   43.   Borkner L, Curham LM, Wilk MM, Moran B, and Mills KHG. IL-17 mediates protective
553   immunity against nasal infection with Bordetella pertussis by mobilizing neutrophils,
554   especially Siglec-F(+) neutrophils. *Mucosal immunology.* 2021;14(5):1183-202.
555   44.   Gattinoni L, Speiser DE, Lichterfeld M, and Bonini C. T memory stem cells in health and
556   disease. *Nat Med.* 2017;23(1):18-27.
557   45.   Safar HA, Mustafa AS, Amoudy HA, and El-Hashim A. The effect of adjuvants and delivery
558   systems on Th1, Th2, Th17 and Treg cytokine responses in mice immunized with
559   Mycobacterium tuberculosis-specific proteins. *PloS one.* 2020;15(2):e0228381.
560   46.   Nguyen N, Guleed S, Olsen AW, Follmann F, Christensen JP, and Dietrich J. Th1/Th17 T
561   cell Tissue-Resident Immunity Increases Protection, But Is Not Required in a Vaccine
562   Strategy Against Genital Infection With Chlamydia trachomatis. *Front Immunol.*
563   2021;12:790463.
564   47.   Bagwell CB, Hunsberger B, Hill B, Herbert D, Bray C, Selvanantham T, et al. Multi-site
565   reproducibility of a human immunophenotyping assay in whole blood and peripheral blood
566   mononuclear cells preparations using CyTOF technology coupled with Maxpar Pathsetter,
567   an automated data analysis system. *Cytometry B Clin Cytom.* 2020;98(2):146-60.
568

Figure 1. CT genital tract infection elicits monocytes, plasma cells, and CD4 T effector memory cells in the periphery. (A) Major immune populations defined by manual gating in peripheral blood of 82 participants at enrollment and 1 month post-treatment. (B-M) Population percentages of total PBMCs. Horizontal bar represents mean. Significance was determined by Kruskal-Wallis test followed by Dunn's test and Bonferroni correction for multiple comparisons. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 2. Participants with CT have increased peripheral frequencies of long-lived Th2 and Th17-lineage CD4 TEM cells compared to uninfected participants.
(A) Subsets of CD4 and CD8 TEM were identified by expression of chemokine receptors CCR6, CXCR3, and CCR4. (B-K) Population percentages of total T Cells. Horizontal bar represents mean. Significance was determined by Kruskal-Wallis test followed by Dunn's test and Bonferroni correction for multiple comparisons. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 3. Frequencies of CD4 TSCM are enriched in the periphery of participants with endometrial CT infection.

Frequencies of manually gated T cell subpopulations were stratified by cervix only CT (Endo-) or cervix and endometrial CT (Endo+) status. (A-D) T cell subpopulation percentages. Horizontal bar represents mean. Significance was determined by Wilcoxon rank-sum test.

Figure 4. CD4 TCM Th1 and Th17 family subsets are enriched in peripheral blood of women without documented CT reinfection.

Frequencies of manually gated T cell subpopulations were stratified by follow-up negative (FU-) and follow-up positive (FU+) status. (A-J) T cell subpopulation percentages. Horizontal bar represents mean. Significance was determined by Wilcoxon rank-sum test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.