Unique T cell signatures associated with reduced *Chlamydia trachomatis* reinfection in a

highly exposed cohort

Kacy S Yount[1], Chi-Jane Chen[2], Avinash Kollipara[1], Chuwen Liu[3], Neha V Mokashi[3], Xiaojing

Zheng[1,3], C Bruce Bagwell[4], Taylor B Poston[1], Harold C Wiesenfeld[5], Sharon L Hillier[5], Catherine

M O'Connell[1], Natalie Stanley[2,6], Toni Darville[1,7]

[1]Department of Pediatrics, University of North Carolina at Chapel Hill, Chapel Hill, NC;

[2]Department of Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC;

[3]Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC; [4]P.O.

Box 247, Topsham, ME; [5]The University of Pittsburgh School of Medicine and the Magee-

Womens Research Institute, Pittsburgh, PA; [6]Computational Medicine Program, University of

North Carolina at Chapel Hill, Chapel Hill, NC; [7]Department of Microbiology and Immunology,

University of North Carolina at Chapel Hill, Chapel Hill, NC

Corresponding Author:

Toni Darville

lad@email.unc.edu

501-944-3130

4008B Mary Ellen Jones, 116 Manning Drive, Chapel Hill, NC, 27599

The authors have declared that no conflict of interest exists.

ABSTRACT

Chlamydia trachomatis (CT) is the most common bacterial sexually transmitted infection

globally. Understanding natural immunity to CT will inform vaccine design. This study aimed to

profile immune cells and associated functional features in CT-infected women, and determine

immune profiles associated with reduced risk of ascended endometrial CT infection and CT

reinfection. PBMCs from CT-exposed women were profiled by mass cytometry and random forest

models identified key features that distinguish outcomes. CT+ participants exhibited higher

frequencies of CD4+ Th2, Th17, and Th17 DN CD4 T effector memory (TEM) cells than

uninfected participants with decreased expression of T cell activation and differentiation markers.

No significant differences were detected between women with or without endometrial CT infection.

Participants who remained follow-up negative (FU-) showed higher frequencies of CD4 T central

memory (TCM) Th1, Th17, Th1/17, and Th17 DN but reduced CD4 TEM Th2 cells than FU+

participants. Expression of markers associated with central memory and Th17 lineage were

increased on T cell subsets among FU- participants. These data indicate that peripheral T cells

exhibit distinct features associated with resistance to CT reinfection. The highly plastic Th17

lineage appears to contribute to protection. Addressing these immune nuances could

promote efficacy of CT vaccines.

GRAPHICAL ABSTRACT

INTRODUCTION

Chlamydia trachomatis (CT) is the most common bacterial sexually transmitted infection (STI) globally (1). In women, CT infects the cervix and can ascend to the uterus and oviducts to cause pelvic inflammatory disease (PID) that can result in pelvic pain, infertility, and ectopic pregnancy (2). Although antibiotic treatment is effective, 80% of infections are asymptomatic and may go undetected (3), contributing to the risk for chronic sequelae (4). Although repeated CT infections are common, they can elicit partially protective immunity. Women who spontaneously cleared infection without treatment were less likely to become reinfected (5) and persons with confirmed prior infection had lower chlamydial loads (6). Determining the aspects of natural immunity that drive protection against CT will advance understanding of chlamydial immunity and aid design of an effective vaccine.

Mouse models have elucidated protective mechanisms of different arms of the immune response. The importance of CD4 T cell mediated immunity is revealed from studies in severe combined immune deficient (SCID), RAG-1-/-, TCRα-/-, CD4-/-, and MHCII-/- mice, which display delayed or failed resolution of Chlamydia infection, while MHCI-deficient (B2M-/-), B cell-deficient (μMT), and CD8-/- mice resolve infection with comparable kinetics to wild-type (7-11). Th1 cells secrete IFNγ and TNFα that promote recruitment and activation of macrophages. IFNγ also induces expression of indoleamine 2,3-dioxygenase (IDO) in epithelial cells, which catabolizes tryptophan, an essential nutrient for chlamydia growth (12). IFNγ-knockout, antibody depletion, and adoptive transfer studies in mice confirmed that IFNγ-producing CD4 T cells (Th1) are key mediators of protection against Chlamydia (13-15), although this may be independent of T-bet, the defining transcription factor expressed by Th1 cells (16). Th2 cells produce IL-4, IL-5, and IL-13 that drive antibody responses, but a Ct-specific Th2 clone did not protect against murine genital infection and displayed reduced trafficking to the genital mucosa, whereas a polyfunctional Th1 clone mediated chlamydial clearance (17, 18). Classical Th17 cells produce IL-17, recruit

neutrophils, promote secretory IgA, and are instrumental in protection against many mucosal

pathogens, but can be pathogenic. Mice lacking T-bet had Th17-skewed responses and similar

clearance of *C. muridarum* infection to wild-type (16) while those lacking RORγt (Th17-deficient)

exhibited delayed clearance (19), highlighting a potential protective role for Th17. However, we

found that IFNγ-deficient mice had chronic pathology related to a compensatory increase in Th17

cells and downstream induction of neutrophils (20), highlighting the importance for a balance in

Th17 responses.

The T Cell Response Against Chlamydia (TRAC) cohort is a unique resource containing

clinical samples and information from women highly exposed to CT and other STIs that was

established to improve understanding of human natural immunity to CT (21). In this cohort, we

observed that peripheral blood CD4 T cells producing CT-specific IFNγ were associated with

protection from reinfection (22), consistent with findings of others (23), and we additionally

determined that CD8 T cells producing CT-specific IFNγ were associated with protection from

endometrial infection (22). Further, we identified a single nucleotide polymorphism in the gene for

CD151, a tetrospanin molecule important for immune cell adhesion and migration and T cell

proliferation, as a candidate genetic marker for reduced susceptibility to endometrial infection

(24). In contrast, we determined that anti-CT antibodies were insufficient to protect participants

from endometrial or recurrent infection (25, 26). These data indicate that CD4 T cells, and to a

lesser extent CD8 T cells, contribute to protection against CT reinfection or ascension.

The objective of this study was to immunophenotype peripheral blood from TRAC

participants to identify specific T cell subsets and surface markers associated with T cell function

that associated with resistance to CT reinfection and/or endometrial infection. Therefore, a CyTOF

panel containing 33 markers was designed to deeply profile T-helper cell subsets such as T

follicular helper (Tfh; CXCR5+) cells, regulatory T (Treg; CTLA4+ CD25+ CD127lo) cells, naïve T

cells (TN; CD45RA+ CCR7+), central memory T (TCM; CD45RA- CCR7+) cells, effector memory

T (TEM; CD45RA- CCR7-) cells, terminally differentiated T effector (TEMRA; CD45RA+ CCR7-)

cells, and stem-cell like memory T (TSCM; CD45RA+ CCR7+ CD28+ CD95+) cells. TEM and

TCM T-helper cell subsets were defined by surface expression of chemokine receptors CCR6,

CXCR3, and CCR4 (27) because multiple studies have shown that CCR6-CXCR3+CCR4- (Th1),

CCR6-CXCR3-CCR4+ (Th2), and CCR6+CXCR3-CCR4+ (Th17) cells polarized or sorted from

human peripheral blood express the canonical cytokines and transcription factors associated with

their corresponding T-helper phenotype (27-31). Th17-lineage cells are also highly plastic and

modify chemokine receptor and effector cytokine expression depending on their environment (32,

33). Classical Th17 cells (CCR6+CXCR3-CCR4+) produce primarily IL-17 and are important for

neutrophil recruitment and defense at mucosal sites (34); Th1/17 cells (CCR6+CXCR3+CCR4-)

share Th1 and Th17 functionality, respond to IL-12 and IL-23, and produce mostly IFNγ (35);

Th17 double positive (DP) cells (CCR6+CXCR3+CCR4+) produce IFNγ and low levels of IL-17

(29), and double negative (DN) cells (CCR6+CXCR3-CCR4-) preferentially produce IL-17 and

express transcriptional markers of early Th17 development (IL-17F, STAT3), lymph-node homing

(CCR7, CD62L), follicular help (CXCR5, BCL6, ASCL2), and self-renewal (LEFI, MYC, TERC)

(29, 36).

To enhance the depth of our analyses, we incorporated 13 surface markers that relate to

T cell function into a machine learning model for predicting clinical outcomes. The panel included

surface markers associated with: activation/proliferation (CD71, HLA-DR, CD38, CD25);

degranulation/cytotoxicity (CD94, CD107a); differentiation/memory (loss of CD27, CD28,

CD127); Treg/checkpoint inhibitor/exhaustion (CTLA4, PD-1); Tfh/germinal center homing

(CXCR5); and mucosal migration (CCR5). While frequencies of T cell subsets could be analyzed

individually, the incorporation of expression levels of T cell functional features resulted in over

400 subset-marker pairs and required use of computational methods to analyze this high-

dimensional dataset. Using a random forest machine learning model, we determined whether T

cell subset frequencies and marker expression levels could distinguish clinical outcomes. When

accurate prediction was achieved, feature importance analysis by Gini score (37) determined the

most discriminatory features.

Increased frequencies of CD4 TEM Th2 and Th17-lineage subsets and reduced

expression of T cell activation/differentiation markers among CT+ or CT+CoP participants

compared to uninfected were most informative to the model's correct classification of CT infection.

The model was unable to accurately predict endometrial infection but was successful in identifying

features associated with absence of reinfection. The most discriminating features were increased

frequencies of CD4 TCM Th1, Th17, Th1/17, and Th17 DN cells and increased expression of

markers associated with central memory and Th17-lineage. Although these data suggest a

protective role for Th17 cells, their contribution to human immunopathogenesis remains

undetermined.

RESULTS

CT genital tract infection elicits monocytes, plasma cells, and CD4 T effector memory cells

in the periphery.

This study acquired CyTOF data from PBMCs of 82 of the 246 participants enrolled in

TRAC (21). The sociodemographic characteristics, self-reported STI diagnoses, and exposure

histories of this sub-cohort resemble those previously described for the full cohort (Table S1, S2)

(21). Study participants were assigned to infection outcome groups according to the presence of

CT or copathogens (CoP) *Neisseria gonorrhoeae* (NG) and/or *Mycoplasma genitalium* (MG) at

enrollment: Uninfected (CT- NG- MG-), CT+ (CT+ NG- MG-), or CT+CoP (CT+; NG+ and/or

MG+). PBMCs, collected at enrollment and 1 month post treatment (1M), were

immunophenotyped by CyTOF using a 33-marker panel (Fig 1A, Table S3). The manual gating

hierarchy and definitions used to determine frequencies of each cell type are outlined in Table

S4. Frequencies of total myeloid cells, B cells, CD4 T cells, and CD8 T cells were similar across

all groups (Fig 1B-E), as were dendritic cells and NK cells (data not shown). However, classical,

intermediate, and non-classical monocytes were increased in CT+ and CT+CoP women at

enrollment and 1M, with significant increases noted in non-classical monocytes, which are

important for antigen presentation (Fig 1F-H). Macrophage frequencies were variable but slightly

increased in CT+ participants at enrollment and 1M (Fig S1A). Although frequencies of total B

cells were not different at enrollment and 1M, frequencies of plasma cells were increased in CT+

but not CT+CoP participants (Fig 1I). Previous studies have shown that CT drives a strong and

specific antibody response (21, 38), consistent with a high frequency of plasma cells. Although

frequencies of total T cells were not different, frequencies of CD4 TEM cells were significantly

increased in CT+ and CT+CoP participants at enrollment and 1M (Fig 1J), while CD8 TEM were

only slightly increased in CT+ and CT+CoP participants (Fig 1K). In contrast, CD4 and CD8 TCM

frequencies were similar or slightly reduced in CT+ and CT+CoP participants compared to

uninfected (Fig 1L, M). Frequencies of CD4 and CD8 stem cell-like memory T cells (TSCM) were

moderately decreased in CT+ and CT+CoP participants at enrollment and 1M, likely a reflection

of the proportional increase in TEM (Fig S1B, C). Overall, these data demonstrate that CT

infection drives increased frequencies of peripheral non-classical monocytes, plasma cells, and

CD4 TEM cells, and that responses remain elevated for at least one month after treatment.

CT+ participants have increased peripheral frequencies of Th2 and Th17-lineage CD4 TEM

cells.

We used differential expression of the chemokine receptors CCR6, CXCR3, and CCR4 to

define Th1, Th2, and Th17 subsets (Fig 2A, Table S4) (27-29). CD4 T cells, and specifically IFNγ-

producing Th1 cells, are broadly accepted as the primary contributors to protection against CT in

animals and humans (22, 39), and yet, frequencies of CD4 TEM Th1 cells were not different

between CT-infected and uninfected participants (Fig 2B). Instead, frequencies of CD4 TEM Th2

cells were significantly increased in CT+ and CT+CoP participants at enrollment and 1M (Fig 2C),

consistent with strong antibody responses detected in response to CT (21, 38). Frequencies of

CD4 TEM Th17 and Th17 DN cells were also increased in CT+ participants at enrollment and 1M,

whereas there was no difference in frequencies of Th1/17 or Th17 DP cells (Fig 2D-G). These

differences were maintained in CT+CoP participants for CD4 TEM Th2 but were dampened in the

case of Th17 lineage cells (Fig 2C-F), suggesting that coinfection dilutes Th17 responses elicited

by CT.

We similarly investigated cells comprising the CD8 memory compartment (Tc1, Tc2, Tc17,

etc.; Fig 2A, Table S4). Frequencies of CD8 TEM Tc1 and Tc2 were significantly increased in

CT+ participants compared to uninfected (Fig 2H-I). There was no change in CD8 TEM Tc1/17

(Fig 2J), but Tc17 DN cells trended toward an increase in CT+ participants (Fig 2K), both of which

parallel the results for CD4 TEM subsets. Again, these differences were reduced when comparing

uninfected to CT+CoP participants. CD8 TEM Tc17 and Tc17 DP were not detected because

there was very low co-expression of CCR6 and CCR4 on CD8 T cells. CD4 or CD8 TCM subset

frequencies were similar or slightly decreased in CT+ participants (data not shown), following the

trend previously observed for frequencies of total CD4 and CD8 TCM (Fig 1L-M). Overall, these

results suggest that CT infection drives a peripheral adaptive immune response dominated by

CD4 and CD8 TEM Th2/Tc2 and CD4 TEM Th17-lineage cells.

We then implemented an unsupervised clustering-based strategy to increase the

resolution of our findings obtained by supervised manual gating (Fig 3A-F). Clustering with the *k-*

means algorithm partitioned cells based on expression of six phenotypic markers (CD45RA,

CCR7, CD95, CCR6, CXCR3, CCR4). We previously defined TSCM cells as

CD45RA+CCR7+CD28+CD95+ by manual gating (Table S4), but since expression of CD28 was

similar across all clusters, this marker was excluded, and TSCM were defined as

CD45RA+CCR7+CD95+. Following clustering, memory and T-helper subtypes were annotated

based on marker expression as previously (Table S4, S5; Fig 3A-F). Annotation revealed similar

clusters to manual gating overall, but enhanced resolution allowed detection of heterogeneity in

TN subsets (Fig 3C, F). CD4 clusters 1, 20, and 22 and CD8 clusters 6, 15, and 22 expressed

high levels of both CD45RA and CCR7, representing the classical definition of TN. However, CD4

clusters 10, 14, 16 and 19 expressed lower levels of CD45RA and/or CCR7, suggesting that these

cells were differentiating from a naïve state to a memory state. Additionally, CD8 clusters 1, 19,

4, and 13 were CD45RA+CCR7+ like traditional TN cells but also expressed CXCR3, a marker

typically reserved for memory Th1 cells. Studies have shown that naïve human CD8 T cells

expressing CXCR3 have greater potential for effector differentiation (40).

Clusters with frequencies that were statistically significant (P<0.05 by Wilcoxon test)

between Uninfected and CT+ or CT+CoP participants at Enrollment or 1 Month are represented

in Fig 3G as a distribution of cluster frequencies per participant. Frequencies of CD4 clusters 10

(TN with low CCR7), 14 (Transitional TCM), 25 (TEM), and 3 and 9 (TEM Th2) were increased in

CT+ and CT+CoP participants compared to uninfected, while frequencies of clusters 1 and 8

(classical TN), 5 and 23 (TCM), and 17 (TCM Th2) were decreased, suggesting that CD4 T cells

respond to CT by evolving away from a naïve state. Similar immune signatures were observed at

enrollment and 1 month, suggesting that changes in immune responses in the periphery are

maintained after treatment. These findings were consistent with manual gating where overall CD4

TEM (Fig 1J), and especially TEM Th2 cells (Fig 2C), were increased in CT-infected participants

compared to uninfected while overall frequencies of CD4 TCM were trending towards decrease

(Fig 1L).

CT+ and CT+CoP participants express reduced T cell activation and differentiation

markers on several CD4 and CD8 subsets compared to uninfected participants.

While the unsupervised clustering approach lent confidence to our previous findings and

uncovered previously unappreciated nuance in the TN population, it demonstrated limited

resolution of low frequency T cell subsets defined by manual gating, such as Th17-lineage cells,

that were of particular interest to us. Therefore, for further analysis of marker expression on each

T cell subset, we used manually gated subsets. We implemented a random forest machine

learning model trained by T cell subset frequencies as well as expression levels of the phenotypic

markers used above to define T cell subsets (CD45RA, CCR7, CD95, CCR6, CXCR3, CCR4)

and 13 surface markers that relate to T cell function to predict clinical outcomes. The panel

included surface markers associated with the following T cell functions: activation/proliferation

(CD71,   HLA-DR,   CD38,   CD25);   degranulation/cytotoxicity   (CD94,   CD107a);

differentiation/memory (loss of CD27, CD28, CD127); Treg/checkpoint inhibitor/exhaustion

(CTLA4, PD-1); Tfh/germinal center homing (CXCR5); and mucosal migration (CCR5). The upper

quartile expression of each of these 19 markers was calculated for each CD4 or CD8 manually

gated subset (as previously defined, Fig 2, Table S4) and for each participant. These upper

quartile expression levels (functional features) were then used in addition to the manually gated

frequency features to train the random forest model. This machine learning approach allowed

interrogation of over 400 features defined by subset-marker pairs.

Classification of uninfected versus CT+ or CT+CoP participants subset frequency and

functional features was successful for CD4 (Fig 4A; AUC range 0.96-1.00) or CD8 T cells (Fig4H;

AUC range 0.93-0.98). Together, these results suggest that frequency and functional features of

CD4 and CD8 T cells contribute to distinction between CT-infected participants and uninfected

participants. The similarity between model success at enrollment and 1 month suggests that these

signatures were maintained after infection resolution.

To identify prominent cellular subsets that contributed to prediction, a measure of feature

importance, or Gini score, was assigned to each frequency feature or functional feature (37).

Briefly, the Gini score encodes the extent to which each frequency feature or functional feature

provides the model with useful information, such that subsets with high Gini scores are likely to

be the most useful in distinguishing between clinical outcomes. The Gini scores for predicting

classification of infection status are represented in Table S6 for CD4 features and Table S7 for

CD8 features. In general, features contributing to accurate classification of uninfected versus CT+

and uninfected versus CT+CoP were similar for the models informed by CD4 and CD8 features,

consistent with similar accuracy between the two comparisons and weak classification accuracy

for CT+ versus CT+CoP. A subset of upper quartile functional features with highest importance

by Gini score and significant difference in expression between uninfected and CT+ or CT+CoP

by Wilcoxon test (P<0.001) is shown in Fig 4B-G (CD4), I-M (CD8). Several subsets of both CD4

and CD8 T cells in CT infected participants expressed lower levels of CD25 (IL-2Rα), CD95 (Fas),

and CD38, markers associated with T cell activation and antigen experience, than the same

subsets in uninfected participants (Fig 4B-D,I-J). Loss of costimulatory molecules CD27 and

CD28 is associated with T cell differentiation, and CD28 expression was higher in some CD4 and

CD8 subsets from CT+ and CT+CoP participants than in uninfected participants (Fig 4E, K),

suggesting that peripheral blood from infected participants contained less differentiated T cells.

Activated, antigen experienced, differentiated T cells may have trafficked to the infected genital

tract, resulting in a reciprocal decrease in differentiation markers in the blood. Several subsets,

especially TSCM and TCM subsets, also expressed lower levels of CCR7 in CT+ or CT+CoP

participants than uninfected participants (Fig 4F, L). This aligns with increased frequencies of

effector memory (CCR7-) T cells in CT+ and CT+CoP participants and suggests that the central

memory cells of infected participants are shifted more toward an effector phenotype. Additionally,

expression of the mucosal homing marker CCR5, a marker known to be important for trafficking

of T cells to the genital tract during murine chlamydial infection (41, 42), was decreased in several

subsets in CT+ or CT+CoP participants compared to uninfected participants (Fig 4G, M). This

CCR5 expression pattern could suggest that CCR5+ T cells had migrated to the genital mucosa

at the time of sampling. Frequencies of T cell subsets, especially TEM Th2, TEM Th17DN, and

TEM Th17 among CD4 T cells (Table S6) and TEM Tc2 and generic TEM among CD8 T cells

(Table S7), were also important contributors to the differentiation of CT+ from uninfected

participants. Trends in feature importance analysis were similar for both CD4 and CD8 T cells at

1 month (Fig 4, Table S6,7), suggesting maintenance of the T cell signature following infection

resolution.

Discrimination of CT+ versus CT+CoP participants using CD4 or CD8 frequency

demonstrated only weak to moderate success, with greater success from CD4 features (Fig 4A;

AUC range: 0.69-0.74) than CD8 (Fig 4H; AUC range: 0.61-0.64). The top contributing feature by

Gini score was higher CTLA4 expression on CD4 T cell subsets among CT+CoP participants

compared to CT+ participants (Table S6). CTLA4 (cytotoxic T lymphocyte antigen 4) plays an

important role in CD4 Treg function by downregulating T cell activation. NG evades host immunity

by inducing TGFβ, which is important for Treg development. The increase in CTLA4 expression

on CD4 T cell subsets among CT+CoP participants compared to CT+ participants could be

reflective of NG-mediated immune evasion.

Neither T cell subset frequencies nor functional expression features predict endometrial

CT infection.

Endometrial biopsies from all participants were tested for CT, NG, and MG infection at

enrollment, enabling bisection of the CT+ and CT+CoP groups into CT infections that had

ascended to the upper genital tract (Endo+) and CT infections that were limited to the cervix

(Endo-). All major cell populations and T cell subsets were defined by manual gating (Table S4),

and comparisons of population frequencies performed between Endo+ and Endo- participants.

Endo+ participants exhibited significantly increased frequencies of CD4 TSCM, which were not

observed in Endo+ CT+CoP participants (Fig S2A). Frequencies of CD8 TEM were not different

between Endo+ and Endo- outcomes for CT+ or CT+CoP participants (Fig S2B). Among Endo-

participants, there was a trend toward increased frequencies of CD8 TEM Tc17 and TCM Tc17

DN cells (Fig S2C-D), suggesting expansion of these cells may contribute to protection from

ascending CT infection. However, the random forest model trained on manually gated subset

frequencies with the addition of upper quartile expression of functional markers (Fig S2E-F) failed

to accurately classify Endo+ from Endo- participants based on CD4 or CD8 features. Overall,

there were relatively minimal differences in the frequency and functional marker expression

patterns in peripheral blood T cell subsets between participants with or without ascended CT

infection.

Increased TSCM, CD4 TCM Th1, and CD4 TCM Th17-lineage subsets but decreased CD4

TEM Th2 cells in peripheral blood are associated with absence of CT reinfection.

Individuals treated for CT often experience repeat infections (6). Nevertheless, evidence

exists for partially protective immunity after infection (23). TRAC study participants were followed

for one year to determine if they became reinfected (21). We again bisected CT+ and CT+CoP

groups into FU+ (positive for CT by diagnostic test or self-report at any follow-up visit) and FU-

(completed at least 3 out of 4 follow-up visits, and CT negative by diagnostic test and self-report

at all visits). Five out of 20 participants in the CT+ FU+ group tested positive for CT at 1M, while

none of the participants in the CT+CoP FU+ group tested positive for CT at 1M. Removing these

5 participants in the CT+FU+ group from the analysis did not change the overall differences

observed between FU- and FU+ participants.

We compared frequencies of manually gated T cell subsets between FU- and FU+

individuals. Frequencies of CD4 and CD8 total T cells and CD4 and CD8 TSCM cells were

enriched in FU- participants without co-infection at enrollment and remained expanded 1 month

post treatment (Fig 5A-D). CD4 Treg cells were enriched in FU- participants at the enrollment visit

(Fig 5E). Interestingly, frequencies of CD4 TEM Th2 cells were increased in FU+ participants (Fig

5F), suggesting that although these cells are induced by CT infection (Fig 2C), they are not

protective. In contrast, frequencies of CD4 TCM Th1, Th17, Th1/17, and Th17 DN cells were

increased in FU- participants (Fig 5G-J). These data suggest CD4 and CD8 TSCM and CD4 TCM

Th1/Th17 lineage cells, are important for natural immunity against incident CT infection.

Differences in frequencies of Th17 family subsets between FU- and FU+ participants were only

detected among CT+ and not CT+CoP participants, which suggests that Th17 immune responses

specific to CT are diluted by coinfection.

We then repeated the comparison using frequencies of clusters generated by *k*-means

(Fig 3A-F) and observed trends toward higher frequencies of clusters 15 (TSCM) and 12 and 21

(Th17 family) and lower frequencies of clusters 3, 4, and 9 (Th2 TEM) in FU- participants

compared to FU+ (Fig S3; see Fig 3C for cluster annotations). These trends were consistent with

results obtained by traditional manual gating (Fig 5) and suggest a protective role for TSCM and

TCM Th17-lineage cells but not for Th2 TEM. However, there was limited resolution of Th17 family

subsets by clustering compared to manual gating.

Central memory and Th17-lineage markers distinguish participants who did not become

reinfected with CT.

We used a random forest machine learning model trained with manually gated T cell

subset frequencies and functional expression level features to classify FU+ versus FU- status (Fig

6). Classification of FU+ versus FU- was most accurate for CT+ participants at enrollment (CD4

ROC AUC = 0.72; CD8 ROC AUC = 0.70) (Fig 6A,B). Analysis of feature importances by Gini

score identified cell-types and subset-specific expression patterns that were most distinct (Table

S8). Among CT+ participants, the most prominent CD4 features contributing to successful FU-

versus FU+ classification were decreased frequencies of Th2 cells and increased frequencies of

Th17 family subsets in FU- participants (Table S8). The upper quartile functional features with

highest importance by Gini score and significant difference in expression between FU+ and FU-

by Wilcoxon test (P<0.05) are highlighted in Fig 6C-F. Important CD4 features included increased

expression of CCR7 on T cell subsets among FU- participants, consistent with increased central

memory, and increased CCR6 expression on Th17 subsets among FU- participants, consistent

with increased Th17 lineage (Fig 6C-D). Important CD8 features among FU- participants included

increased frequencies of CD8 TSCM, increased expression of CCR7 (similar to the CD4 model),

and reduced expression of CD127 (IL-7 receptor, loss of which is associated with differentiation

and memory) (Table S8, Fig 6E-F). Overall, the model was most successful in distinguishing FU+

from FU- participants among CT+ participants when informed by CD4 T cell frequencies at

enrollment. CD8 T cell features and T cell features at 1 month were less informative, and

coinfection with NG or MG diluted the success of the model. Overall, these results suggest a

protective role for CD4 TCM Th17-lineage cells against CT reinfection.

DISCUSSION

Using samples from a uniquely characterized and clinically relevant cohort, this study

immunophenotyped responses to CT infection with high dimensional analysis of surface marker

expression on PBMCs and found specific T cell subsets correlated with reduced incidence of CT

reinfection. Overall, broad immune cell populations in peripheral blood differed between

uninfected and CT+ participants, but were similar between CT monoinfection and coinfection with

NG and/or MG. CT+ participants displayed increased frequencies of non-classical (CD14-CD16+)

monocytes, important for antigen presentation, T cell proliferation, patrolling tissues for signs of

injury, and increased Fc receptor-mediated phagocytosis (43).

While frequencies of immune populations were largely similar in CT+ participants

regardless of coinfection, plasma cells were elevated in participants with CT monoinfection, but

their frequencies among CT+CoP participants were similar to uninfected. This is consistent with

robust antibody responses to CT infection (25, 44, 45), but this response appears suppressed by

coinfection with other STIs. NG opacity proteins bind carcinoembryonic antigen-related cellular

adhesion molecule 1 (CEACAM1 or CD66a) on B cells, leading to inhibitory signals which promote

B cell death (46). Plasma cells also express CD66a (47). Thus, opacity protein interactions with

CD66a could contribute to reduced numbers of plasma cells.

CD4 TEM frequencies were strikingly increased in CT+ and CT+CoP participants

compared to CD8 TEM frequencies. These data are consistent with our prior investigation of CT

antigen specific responses in a subset of TRAC participants where we observed higher levels of

IFNγ production and a greater breadth of recognition of CT antigens among CD4 T cells compared

to CD8 T cells (22). Lower CD8 TEM frequencies are also consistent with engagement of the PD-

1 receptor on CD8 T cells by antigen presenting cells inhibiting generation of CT-specific memory

CD8 T cells (48). Frequencies of TEM but not TCM were increased in CT+ compared to uninfected

participants in our study.

Investigation of CD4 TEM subsets revealed that CT+ participants had higher frequencies

of peripheral Th2, Th17, and Th17 DN cells compared to uninfected participants, with unchanged

frequencies of Th1 cells. CD4 Th1 cells are classically defined by production of IFNγ and

expression of CXCR3 and/or CCR5 (28). They are well-documented contributors to chlamydial

clearance and protection from reinfection in mouse models (18, 41, 42) and associate with

resistance to reinfection in humans (22, 23). We did not find that Th1 cells were expanded in the

peripheral blood of CT+ participants, potentially explaining why CT reinfection is common.

However, Th1 cells express the chemokine receptors CXCR3 and CCR5, which bind the

chemokines CXCL10 and CCL5 (RANTES), respectively (49). We previously reported detecting

these chemokines in cervical secretions of TRAC participants (50), and they are highly expressed

in genital tract tissues of *C. muridarum* infected mice (41). In this study, we found decreased

expression of CCR5 on T cell subsets among CT+ and CT+CoP participants. Thus, it is possible

that despite overall expansion of Th1 cells, they were preferentially recruited to the site of infection

and consequently not increased in the periphery. In contrast, peripheral CD4 Th2 cells were

significantly enriched in CT+ and CT+CoP participants, consistent with high antibody titers to CT

infection that fail to prevent reinfection (6, 21, 25).

Th17 cells are highly plastic. They are mainly defined by their expression of CCR6 (51)

but can gain or lose chemokine receptors and their associated effector functions (e.g., cytokine

production) (32). Th17 cells can be pathogenic, but they are also important contributors to

protection against mucosal pathogens (52, 53). We used expression of CCR6, CXCR3, and

CCR4 to define four subsets of Th17 family cells: classical Th17, Th1/17, Th17 DP, and Th17 DN

(29). We found increased frequencies of classical Th17 and Th17 DN cells in CT+ participants

compared to uninfected, which were reduced by coinfection. Frequencies of parallel populations

of CD8 TEM cells including Tc2 and Tc17 DN cells were also increased in CT+ and CT+CoP

participants, albeit to a lesser extent than CD4 counterparts. Co-expression of CCR6 and CCR4

was extremely rare on CD8 T cells, and therefore Tc17 and Tc17 DP populations were excluded

from analysis.

An important goal of the study was to analyze expression of additional surface markers

associated with activation/proliferation (CD71, HLA-DR, CD38, CD25); degranulation/cytotoxicity

(CD94, CD107a); differentiation/memory (loss of CD27, CD28, CD127); Treg/checkpoint

inhibitor/exhaustion (CTLA4, PD-1); Tfh/germinal center homing (CXCR5); and mucosal

migration (CCR5) on each T cell subset and to compare expression patterns between clinical

outcomes. While T cell subsets could be analyzed individually, investigating differences in

expression of phenotypic and functional markers on each subset resulted in over 400 subset-

marker pairs. Utilizing a random forest machine learning model allowed us to determine whether

T cell subset frequencies and functional expression features were distinct between observed

clinical outcomes of CT infection (Uninfected, CT+, CT+CoP), ascension (Endo+/-), or reinfection

(FU+/-). If the random forest model was successful using a combination of frequency and

functional features to classify clinical outcomes, subsequent feature importance analysis

identified which of the T cell subset frequencies and or subset-marker expression pairs were most

informative to the model. The model successfully classified CT+ or CT+CoP from uninfected

participants using CD4 or CD8 features. Increased frequencies of CD4 TEM Th2, Th17, and Th17

DN cells and distinct expression patterns of CD25, CD95, CD38, CCR7, CD28, and CCR5 were

defined as important features for the model. CD25 is the IL-2 receptor and is crucial for T cell

activation and proliferation. CD95, or Fas, is involved in apoptosis but is also a marker of antigen

experience and stem-cell memory. CD38 is a receptor involved in cell activation, adhesion, and

signaling, and is commonly associated with T cell activation. All three of these markers were

expressed at lower levels in CT+ or CT+CoP participants than uninfected participants. Lower

levels of CCR7 expression on T cells among CT+ and CT+ CoP participants was consistent with

higher frequencies of TEM. Additionally, higher levels of CD28 expression on T cell subsets

among CT+ or CT+CoP participants suggested less differentiation. Lower levels of CCR5, a

molecule important for trafficking of effector T cells to the female genital tract during murine

infection (42) and slightly associated with reduced chlamydia burden in humans (54), were also

observed on T cell subsets among CT+ or CT+CoP participants. Together, these data suggested

that even though CD4 TEM cells were increased in peripheral blood of participants with CT

infection, these T cells were not more activated or differentiated than the same subsets in

uninfected participants. The chronicity often associated with chlamydial infections could lead to a

reduced differentiation state associated with T cell suppression in the periphery, as well as

promote ongoing exit of activated, differentiated T cells from the peripheral blood to the infected

mucosa. Supporting this hypothesis of reciprocal decrease in activation/homing markers from

peripheral blood compared to the site of infection, other studies found that CD38 and/or CCR5

expression was higher among CD4 and CD8 T cells isolated from the genital tract than from

peripheral blood of CT+ women (55, 56).

Since CT ascension is a prerequisite for generation of upper tract disease, we sought to

identify T cell subsets and expression patterns associated with ascended infection. Peripheral

immune cell populations were similar overall between participants with endometrial CT infection

and those that had CT infection limited to the cervix. CD8 TEM Tc17 DN cells trended toward

increased frequencies in Endo- participants, consistent with our previous study showing that CT-

specific CD8 T cells weakly associated with reduced CT ascension (22). Frequencies of CD4

TSCM were significantly increased in Endo+ as well as FU- participants. This is consistent with

our prior finding that endometrial infection was associated with reduced risk of follow-up CT

infection (21), likely because the increased bacterial burden associated with endometrial infection

drives a more robust memory response. However, a random forest model trained with manually

gated T cell subset frequencies and functional expression features was unsuccessful in

classifying Endo+ from Endo- CT infections, suggesting that immune responses in the periphery

are too weak to distinguish whether a participant had ascended CT infection. Evaluation of

immune cell populations in the genital tract will likely be necessary to define the specific features

of immunity that prevent or limit ascending infection. Future studies will investigate immune cells

from cervical and endometrial TRAC samples.

Several T cell subsets were found to be strongly associated with the presence or absence

of recurrent CT infection. Sustained increased frequencies of total CD4 T cells in FU- participants

may indicate developing natural immunity in some participants. We also detected increased

frequencies of CD4 and CD8 TSCM in FU- individuals. TSCM resemble naïve T cells in their

expression of CD45RA and CCR7, but they also express CD95, a marker of antigen experience

and memory (57). They are minimally differentiated memory cells that have a high propensity for

stemness and self-renewal as well as the multipotency to differentiate into a broad range of

memory T-helper phenotypes (57). This may explain why they are associated with a FU- outcome.

The importance of the nature of the T cell response is demonstrated by the contrasting subsets

that were expanded in FU- and FU+ participants. Higher frequencies of TEM Th2 cells, which

were strikingly increased in CT+ participants, were associated with a FU+ outcome, while higher

frequencies of TCM Th1 cells, which were not different between CT+ and uninfected participants,

were associated with FU- status. Overall, an important role for Th1 and Th17-lineage cells in

protection from reinfection is suggested by the striking increase in frequencies of TCM Th1 and

Th17 cells among FU- participants, along with increased frequencies of TCM Th1/17 and TCM

Th17 DN cells. The increased stem-ness and differentiation potential of TSCM and Th17 lineage

cells (31, 58), especially Th17 DN (29, 36), is potentially impactful because their plasticity may

allow them to adapt quickly to fight infection. A recent study detected enrichment of

Mycobacterium tuberculosis (Mtb)-specific Th17-like cells in people who were resistant to Mtb

infection following exposure compared to those who were latently infected (59). Similar trends

were observed when comparing frequencies of T cell subsets as defined by *k*-means clusters.

Overall, clusters annotated as TSCM and Th17 lineage were increased in FU- participants and

clusters annotated as TEM Th2 were decreased in FU- participants.

Our finding that TCM subsets (TCM Th1, TCM Th17, TCM Th1/17 and TCM Th17 DN

cells) rather than TEM subsets were associated with FU- status, is consistent with our detection

of CT antigen-specific CD4 TCMs months post treatment among TRAC participants (60). In

addition, an analysis of *C. pneumoniae*-specific peripheral blood T cells in healthy seropositive

donors found that antigen-specific CD4 TCM were enriched compared to TEM (61). Thus, TCM

appear to make up a greater frequency of antigen-specific CD4 T cells important for protection

against subsequent infection, although we found that active CT infection drives expansion of TEM

cells that remain 1M post-treatment,.

A random forest machine learning model trained to classify FU- versus FU+ informed by

manually gated T cell subset frequencies and functional expression features confirmed that the

most distinct differences were increased frequencies of TCM Th17, Th17 DN, Th1/17, and Th1

cells and decreased frequencies of TEM Th2 cells in FU- participants. Additionally, we identified

distinct expression patterns of surface markers including CCR7 and CCR6. In both the CD4 and

CD8 models for CT+ participants, higher expression of CCR7 was observed on several subsets

among FU- participants. Since CCR7 is a defining marker of central memory T cells and is

important for trafficking to the lymph node, it suggests that central memory cells with the

propensity to be long-lived are associated with protection from reinfection. Higher expression of

CCR6 on subsets of CD4 T cells among FU- participants suggests an importance not only for

Th17 lineage cells, which were defined by the presence of CCR6, but also for its surface

expression levels. CCR6 binds CCL20 is highly expressed in mucosal tissues, driving migration

of Th17 cells to the site of infection. Additionally, high expression of CCR6 is associated with

increased secretion of Th17 effector cytokines and chemokines such as IL-17A, IL-17F, IL-22,

and CCL20, and increased expression of *IL17A*, *IL17F*, *IL22*, *IL23R*, *CCL20*, *RORA*, and *RORC*

(58).

The finding that CD4 Th1 and Th17 subsets were associated with reduced reinfection

informs vaccine development. Different vaccine platforms and adjuvants can shape T-helper

responses (62). Vaccination of mice with a recombinant chlamydial antigen with CAF01 (63), or

a STING agonist (64) elicited Th1 and Th17 responses that led to reduced chlamydial burden

upon challenge. In contrast, a viral vector vaccine failed to elicit antigen-specific CD4 T cell

responses or protection (65).

Our study is limited because although we defined immune cell populations increased in

CT+ participants compared to uninfected participants, the cells were not restimulated with CT

antigens to confirm their CT-specificity. However, immunophenotyping PBMCs directly without

restimulation allowed us to confidently define T cell subsets based on chemokine receptors

without the risk of downregulation upon restimulation. An important open question is whether

Th17 and Th17 DN cells produce CT-specific IL-17 and/or IFNγ, and this should be addressed in

future studies. Another limitation of this study is that cell population frequencies could not be

correlated with endometrial CT infection. Many of the women participating in TRAC selected

contraceptive methods that promote a thin uterine lining, making it difficult to acquire sufficient

tissue for evaluation of pathology. However, ongoing studies are investigating local T cell

responses in endometrial tissue sections from participants with sufficient tissue.

Overall, we found that frequencies of CD4 TEM, specifically Th2, Th17, and Th17 DN

subsets, were enriched in individuals with active CT infection. We also identified that increased

frequencies of CD4 TCM Th1, Th1/Th17, and Th17 were associated with individuals who had no

documented reinfection over the course of a year. Together, these findings increase our

understanding of natural immunity to CT and inform strategies for future vaccine design.

METHODS

Sex as a Biological Variable

The TRAC cohort was restricted to persons with a cervix and uterus because the objective of the study was to determine factors associated with ascension of CT from the cervix to the endometrium.

Study Population

This study involved a subset of 82 of the 246 participants enrolled in the T Cell Response Against Chlamydia (TRAC) cohort (21). TRAC participants were asymptomatic and presented to the clinic for STI screening due to sexual behavior that put them at risk for CT (16). None had signs or symptoms of PID. At enrollment, all participants received antibiotics recommended by the Centers for Disease Control and Prevention for treatment of CT and NG, and individual participants that tested positive for STI at any follow-up visit were retreated (16).

Definition of Clinical Classifications

Uninfected, CT+, and CT+CoP Groups:

Nucleic acid amplification testing (NAAT) was used to test cervical swabs collected from study participants at enrollment for CT, NG, and MG. Participants whose swabs tested negatively for all 3 STIs at enrollment were classified as "uninfected". Participants testing positively for CT but negative for NG and MG were classified as "CT+". Enrollees testing positively for CT and NG, MG, or both were classified as "CT+CoP". Of 82 total TRAC participants studied, 12 were uninfected, 44 were CT+, and 26 were CT+CoP (Table S9). Of 26 CT+CoP participants, 11 were CT+ NG+ MG-, 12 were CT+ MG+ NG-, and 3 were CT+ NG+ MG+.

Endometrial positive and endometrial negative CT:

Endometrial biopsies were also tested by NAAT to determine whether infection was limited to the cervix or had ascended to the upper genital tract. Within CT+ and CT+CoP groups, enrollees

testing positively for CT at the cervix but not the endometrial biopsy were classified as Endo-,

while those testing positively for CT at both sites were classified as Endo+. No participants tested

positive for CT solely in the endometrial sample. Of 44 CT+ participants, 19 were Endo+ and 25

were Endo-. Of 26 CT+CoP participants, 11 were Endo+ and 15 were Endo- (Table S10).

Follow-up CT positive and follow-up CT negative:

NAAT testing was performed on cervical brush samples at enrollment and at 1, 4, 8, and 12 month

follow-up visits. Participants with a positive cervical CT test at enrollment and a positive cervical

CT test or self-reported CT diagnosis from an outside clinic in the time since last visit at any follow-

up visit (regardless of how many visits they completed) were classified as follow-up positive (FU+).

Participants that tested positive for cervical CT at enrollment, completed at least 3 out of 4 follow-

up visits, tested negative for cervical CT at all follow-up visits, and self-reported no CT diagnosis

from an outside clinic in the time since last visit at any follow-up visit were classified as follow-up

negative (FU-). 9 participants attended less than 3 follow-up visits (FU N/A) and were excluded

from follow-up analysis. Of CT+ participants, 19 were FU-, 21 were FU+, and 4 were FU N/A. Of

CT+CoP pariticpants, 9 were FU-, 12 were FU+, and 5 were FU N/A (Table S10). All uninfected

samples were FU- and the group was therefore not bisected.

PBMC CyTOF

Peripheral blood mononuclear cells (PBMCs) were isolated from whole blood via a density

gradient using lymphocyte separation media (Corning). Cells were frozen in 90% FBS with 10%

DMSO and stored in liquid nitrogen until thawed for analysis. Dead cells were stained with 1 µM

Cell-ID Cisplatin 198-Pt (Fluidigm). Cells were stained with an antibody cocktail comprising 33

antibodies (Table S3) and human Fc block (eBioscience) in cell staining buffer (CSB, Fluidigm),

followed by staining with secondary antibody (Anti FITC-144Nd). Stained cells were fixed with 2%

PFA in PBS followed by 1:2000 Cell-ID Intercalator-Ir-125 µM (Fluidigm) in fix and perm buffer

(Fluidigm). Cell concentrations were adjusted to 0.5 x 10^6 cells/mL in CAS (Cell Acquisition

Solution; Fluidigm) with Four Element Calibration beads (Fluidigm). Samples were acquired on a

Helios instrument (Fluidigm) by the UNC Mass Cytometry Core.

Data Analysis

Defining T cell subsets by manual or automated gating.

Raw CyTOF files were cleaned using the Pathsetter algorithm (66).

Manual gating. Cleaned FCS files were analyzed in Cytobank (Beckman Coulter) and FlowJo

(BD) software. The manual gating strategy and surface marker designations selected for immune

cell phenotypes are summarized in Table S4. Some gated CD8 populations were undetectable

(<0.1% of T cells) and were excluded from further analysis (Table S4).

Automated gating. Batch effects were removed from the dataset using the combat algorithm (67).

A random subset of 15,000 cells was selected from each sample, resulting in 1,230,000 total cells

for further analysis. We then performed automated cell-population discovery followed by feature

engineering, as outlined in Stanley et al. (68). CD4 and CD8 T cell populations were each

partitioned into 25 clusters using *k*-means clustering. Briefly, each cell was represented by

expression of six phenotypic markers (CD45RA, CCR7, CD95, CCR6, CXCR3, CCR4), which

were input to the clustering algorithm. Moreover, each cell was labelled in terms of one of the 25

clusters, based on its expression across these six phenotypic markers (Table S5). Modeled after

canonical immune features that are often computed based on manual gates, we engineered per-

cluster frequency features, where for each sample, we computed the proportion of its cells

assigned across each of the 25 clusters.

Feature engineering

While previous studies have created functional features based on the mean or median expression

of functional markers in each cluster (68, 69), some functional markers in this dataset were

expressed by a small proportion of cells, such that there were some cases where median

expression of a marker was zero. Therefore, we computed the upper quartile expression of each

marker across each manually gated T cell subset for each sample. For samples with 0.1% events assigned to a particular subset, the upper quartile features were imputed with the mean of each subset-marker feature. Additionally, subset-marker features for which upper quartile expression was below 2 (range 0-500) were excluded from the training dataset. Upper quartile features were arcsinh transformed with co-factor 5. In all, 19 frequency features and 233 upper quartile expression features were included in the CD4 model training dataset, and 14 frequency features and 174 upper quartile expression features were included in the CD8 model training dataset.

Model training

We trained random forest models using engineered manually gated frequency features and upper quartile expression features to classify and predict clinical outcomes for each participant. For each clinical outcome comparison, we performed 30 trials of five-fold cross validation, whereby each fold used 80% of samples for training and 20% for testing. Reported area under the ROC curves (AUCs) represent the mean accuracy across 30 trials. For biological interpretability of each of the trained random forest models, we further computed per-feature importance measures with the Gini index (37). A higher per-feature Gini score implied that the feature was important in the clinical prediction task. We further imposed directionality to the computed per-feature Gini scores to readily identify condition-specific, predictive features by first choosing a particular clinical outcome to be the *positive (negative) direction*, such that we multiplied the Gini score by 1 (-1), if it was associated with the positive (negative) direction.

Statistics

Statistical calculations were computed in R and python. Statistical tests used are outlined in the appropriate figure legends. $P < 0.05$ was used to determine a significant result. Significance symbols used in figures are defined as: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Study Approval

The institutional review boards for human research of the University of Pittsburgh and the University of North Carolina approved the study protocol. All participants provided written informed consent at the time of enrollment and agreed to be contacted to return for follow-up visits 1, 4, 8, and 12 months after enrollment.

Data and Code Availability

CyTOF data are available for download via ImmPort at https://www.immport.org under study accession number SDY2772. Manually gated T cell subset frequencies reported in the Figures are available in Supporting Data Values. Code for (1) automated gating, (2) feature engineering, (3) random forest training and Gini feature importance analysis, and (4) upper quartile expression boxplots are available via GitHub at https://github.com/ksyount/TRAC_PBMC_CyTOF.

AUTHOR CONTRIBUTIONS

ACKNOWLEDGEMENTS

We thank the women who agreed to participate in this study; Ingrid Macio, Melinda Petrina, Carol Priest, Abi Jett, and Lorna Rabe, for their efforts in the clinic and the microbiology laboratory; and the staff at the Allegheny County Health Department STD Clinic. We thank Jason Whitmire for critical review of the manuscript. We also thank Marie Iannone and the UNC Mass Cytometry Core, which is supported by the University Cancer Research Fund (UCRF) and UNC Cancer Center Core Support Grant #P30CA016086. This work was supported by the National Institute of Allergy and Infectious Diseases through U19AI144181 (TD) and R01AI170959 (CMO). KSY

668   received funding from UNC STI/HIV postdoctoral training grant T32AI007001 and the American

669   Association of Immunologists Intersect Fellowship Program for Computational Scientists and

670   Immunologists.

672
673
674 **Figure 1. CT genital tract infection elicits monocytes, plasma cells, and CD4 T effector**
675 **memory cells.** (A) Major immune populations defined by manual gating in peripheral blood of 82
676 participants at enrollment and 1 month post-treatment. (Created with Biorender.com) (B-M)
677 Population percentages of total PBMCs. White triangle represents mean. Significance determined
678 by Dunn's test with Bonferroni correction for multiple comparisons.
679

680
681
682 **Figure 2. Increased frequencies of Th2 and Th17-lineage CD4 TEM cells among CT+**
683 **participants compared to uninfected participants.** (A) Subsets of CD4 and CD8 TEM were
684 identified by expression of chemokine receptors (Created with Biorender.com). (B-K) Population
685 percentages of total T Cells. Triangle represents mean. Significance determined by Dunn's test
686 with Bonferroni correction for multiple comparisons.
687

Figure 3. CD4 but not CD8 automated gating frequency features differentiate CT+ and CT+CoP participants from uninfected participants.

(A-F) Automated gating by *k*-means clustering and cluster annotation. (A-B) Expression of phenotypic markers (CD45RA, CCR7, CD95, CCR6, CXCR3, and CCR4) was used to partition CD4 T cells into 25 *k*-means clusters. (C) Resulting CD4 clusters were annotated based on marker expression (See Table S4, S11). (D-E) CD8 T cells were partitioned into 25 *k*-means clusters based on the expression of the same phenotypic markers as for CD4 T cells and (F) were annotated. (G) Boxplots representing frequencies of CD4 clusters with significant differences in frequency (Wilcoxon P<0.05). Statistical comparisons between CT+ or CT+CoP and Uninf by Wilcoxon test. White triangle represents mean.

700
Figure 4. CT+ and CT+CoP participants express reduced T cell activation and
differentiation markers on several CD4 and CD8 subsets compared to uninfected
participants. A random forest model trained on (A) CD4 or (H) CD8 manually gated subset
frequency features and upper quartile expression features was used to discriminate between
uninfected, CT+, and CT+CoP outcomes at enrollment (En) or 1 month (1M). Receiver operating
characteristic (ROC) curves describe performance of the model. Top markers of (B-G) CD4 or (I-

M) CD8 T cells were identified by feature importance analysis (see Suppl tables 6-7). Features
were included on the plot if they were statistically significant (as defined by $P<0.001$ by Wilcoxon
rank sum test between outcomes) for at least one comparison between uninfected versus CT+
and/or uninfected versus CT+CoP. Boxplots represent the overall distribution of marker
expression. Individual points represent the upper quartile expression for each participant in the
group. (AUC = area under the ROC curve)

Figure 5. CD4 TCM Th1 and Th17 family subsets are enriched in peripheral blood of participants without documented CT reinfection. Frequencies of manually gated T cell subsets were stratified by follow-up negative (FU-) and follow-up positive (FU+) status. (A-J) T cell subset percentages. White triangle represents mean. Statistical significance determined by Wilcoxon rank-sum test.

729
Figure 6. Increased central memory and Th17-lineage markers distinguish participants
without documented CT reinfection. A random forest model trained on (A) CD4 or (B) CD8
manually gated subset frequency features and upper quartile expression features was used to
discriminate between FU+ and FU- groups at enrollment (En) or 1 month (1M). Receiver operating
characteristic (ROC) curves visualize the performance of the model. Top markers of (C-D) CD4
or (E-F) CD8 T cells among CT+ participants were identified by feature importance analysis (see
Table S9). Features were included on the plot based on a significance threshold of $P<0.05$ by
Wilcoxon rank sum test for FU+ verus FU- at Enrollment. Boxplots represent the overall
distribution of marker expression. Individual points represent the upper quartile expression for
each participant in the group. (AUC = area under the ROC curve)

740
741

BIBLIOGRAPHY

1. Newman L, Rowley J, Vander Hoorn S, Wijesooriya NS, Unemo M, Low N, et al. Global Estimates of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting. *PLoS One.* 2015;10(12):e0143304.

2. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, and Ness RB. Risk of sequelae after Chlamydia trachomatis genital infection in women. *J Infect Dis.* 2010;201 Suppl 2:S134-55.

3. Stamm WE. Chlamydia trachomatis infections: progress and problems. *JInfectDis.* 1999;179 Suppl 2:S380-S3.

4. Wiesenfeld HC, Hillier SL, Meyn LA, Amortegui AJ, and Sweet RL. Subclinical pelvic inflammatory disease and infertility. *Obstetrics and gynecology.* 2012;120(1):37-43.

5. Geisler WM, Lensing SY, Press CG, and Hook EW, 3rd. Spontaneous resolution of genital Chlamydia trachomatis infection in women and protection from reinfection. *J Infect Dis.* 2013;207(12):1850-6.

6. Batteiger BE, Tu W, Ofner S, Van Der Pol B, Stothard DR, Orr DP, et al. Repeated Chlamydia trachomatis genital infections in adolescent women. *J Infect Dis.* 2010;201(1):42-51.

7. Yang X, Hayglass KT, and Brunham RC. Different roles are played by alpha beta and gamma delta T cells in acquired immunity to Chlamydia trachomatis pulmonary infection. *Immunology.* 1998;94(4):469-75.

8. Morrison RP, Feilzer K, and Tumas DB. Gene knockout mice establish a primary protective role for major histocompatibility complex class II-restricted responses in Chlamydia trachomatis genital tract infection. *Infect Immun.* 1995;63(12):4661-8.

9. Magee DM, Igietseme JU, Smith JG, Bleicker CA, Grubbs BG, Schachter J, et al. Chlamydia trachomatis pneumonia in the severe combined immunodeficiency (SCID) mouse. *Reg Immunol.* 1993;5(6):305-11.

10. Su H, Feilzer K, Caldwell HD, and Morrison RP. Chlamydia trachomatis genital tract infection of antibody-deficient gene knockout mice. *Infect Immun.* 1997;65(6):1993-9.

11. Morrison SG, Su H, Caldwell HD, and Morrison RP. Immunity to murine Chlamydia trachomatis genital tract reinfection involves B cells and CD4(+) T cells but not CD8(+) T cells. *Infect Immun.* 2000;68(12):6979-87.

12. Beatty WL, Belanger TA, Desai AA, Morrison RP, and Byrne GI. Tryptophan depletion as a mechanism of gamma interferon-mediated chlamydial persistence. *Infect Immun.* 1994;62(9):3705-11.

13. Rank RG, and Whittum-Hudson JA. Protective immunity to chlamydial genital infection: evidence from animal studies. *J Infect Dis.* 2010;201 Suppl 2:S168-77.

14. Perry LL, Feilzer K, and Caldwell HD. Immunity to Chlamydia trachomatis is mediated by T helper 1 cells through IFN-gamma-dependent and -independent pathways. *JImmunol.* 1997;158(7):3344-52.

15. Stary G, Olive A, Radovic-Moreno AF, Gondek D, Alvarez D, Basto PA, et al. VACCINES. A mucosal vaccine against Chlamydia trachomatis generates two waves of protective memory T cells. *Science.* 2015;348(6241):aaa8205.

788   16.   Rixon JA, Depew CE, and McSorley SJ. Th1 cells are dispensable for primary
789   clearance of Chlamydia from the female reproductive tract of mice. *PLoS Pathog.*
790   2022;18(2):e1010333.
791   17.   Lampe MF, Wilson CB, Bevan MJ, and Starnbach MN. Gamma interferon
792   production by cytotoxic T lymphocytes is required for resolution of Chlamydia
793   trachomatis infection. *Infect Immun.* 1998;66(11):5457-61.
794   18.   Poston TB, Qu Y, Girardi J, O'Connell CM, Frazer LC, Russell AN, et al. A
795   Chlamydia-Specific TCR-Transgenic Mouse Demonstrates Th1 Polyfunctionality
796   with Enhanced Effector Function. *J Immunol.* 2017;199(8):2845-54.
797   19.   Rixon JA, Fong KD, Morris C, Nguyen AT, Depew CE, and McSorley SJ.
798   Elimination of Chlamydia muridarum from the female reproductive tract is IL-12p40
799   dependent, but independent of Th1 and Th2 cells. *PLoS Pathog.*
800   2024;20(1):e1011914.
801   20.   Scurlock AM, Frazer LC, Andrews CW, Jr., O'Connell CM, Foote IP, Bailey SL, et
802   al. Interleukin-17 contributes to generation of Th1 immunity and neutrophil
803   recruitment during Chlamydia muridarum genital tract infection but is not required
804   for macrophage influx or normal resolution of infection. *Infect Immun.*
805   2011;79(3):1349-62.
806   21.   Russell AN, Zheng X, O'Connell CM, Taylor BD, Wiesenfeld HC, Hillier SL, et al.
807   Analysis of Factors Driving Incident and Ascending Infection and the Role of Serum
808   Antibody in Chlamydia trachomatis Genital Tract Infection. *J Infect Dis.*
809   2016;213(4):523-31.
810   22.   Russell AN, Zheng X, O'Connell CM, Wiesenfeld HC, Hillier SL, Taylor BD, et al.
811   Identification of Chlamydia trachomatis Antigens Recognized by T Cells From
812   Highly Exposed Women Who Limit or Resist Genital Tract Infection. *J Infect Dis.*
813   2016;214(12):1884-92.
814   23.   Bakshi RK, Gupta K, Jordan SJ, Chi X, Lensing SY, Press CG, et al. An Adaptive
815   Chlamydia trachomatis-Specific IFN-gamma-Producing CD4(+) T Cell Response
816   Is Associated With Protection Against Chlamydia Reinfection in Women. *Front*
817   *Immunol.* 2018;9:1981.
818   24.   Zhong W, Kollipara A, Liu Y, Wang Y, O'Connell CM, Poston TB, et al. Genetic
819   susceptibility loci for Chlamydia trachomatis endometrial infection influence
820   expression of genes involved in T cell function, tryptophan metabolism and
821   epithelial integrity. *Front Immunol.* 2022;13:1001255.
822   25.   Darville T, Albritton HL, Zhong W, Dong L, O'Connell CM, Poston TB, et al. Anti-
823   chlamydia IgG and IgA are insufficient to prevent endometrial chlamydia infection
824   in women, and increased anti-chlamydia IgG is associated with enhanced risk for
825   incident infection. *Am J Reprod Immunol.* 2019;81(5):e13103.
826   26.   Liu C, Hufnagel K, O'Connell CM, Goonetilleke N, Mokashi N, Waterboer T, et al.
827   Reduced Endometrial Ascension and Enhanced Reinfection Associated With
828   Immunoglobulin G Antibodies to Specific Chlamydia trachomatis Proteins in
829   Women at Risk for Chlamydia. *J Infect Dis.* 2022;225(5):846-55.
830   27.   Mousset CM, Hobo W, Woestenenk R, Preijers F, Dolstra H, and van der Waart
831   AB. Comprehensive Phenotyping of T Cells Using Flow Cytometry. *Cytometry Part*
832   *A.* 2019;95(6):647-54.

28. Bonecchi R, Bianchi G, Bordignon PP, D'Ambrosio D, Lang R, Borsatti A, et al.
Differential expression of chemokine receptors and chemotactic responsiveness
of type 1 T helper cells (Th1s) and Th2s. *J Exp Med.* 1998;187(1):129-34.

29. Wacleche VS, Goulet JP, Gosselin A, Monteiro P, Soudeyns H, Fromentin R, et al.
New insights into the heterogeneity of Th17 subsets contributing to HIV-1
persistence during antiretroviral therapy. *Retrovirology.* 2016;13(1):59.

30. Singh SP, Zhang HH, Foley JF, Hedrick MN, and Farber JM. Human T cells that
are able to produce IL-17 express the chemokine receptor CCR6. *J Immunol.*
2008;180(1):214-21.

31. Takeshita M, Suzuki K, Kassai Y, Takiguchi M, Nakayama Y, Otomo Y, et al.
Polarization diversity of human CD4+ stem cell memory T cells. *Clin Immunol.*
2015;159(1):107-17.

32. Muranski P, and Restifo NP. Essentials of Th17 cell commitment and plasticity.
Blood. 2013;121(13):2402-14.

33. Basdeo SA, Cluxton D, Sulaimani J, Moran B, Canavan M, Orr C, et al. Ex-Th17
(Nonclassical Th1) Cells Are Functionally Distinct from Classical Th1 and Th17
Cells and Are Not Constrained by Regulatory T Cells. *J Immunol.*
2017;198(6):2249-59.

34. Khader SA, Gaffen SL, and Kolls JK. Th17 cells at the crossroads of innate and
adaptive immunity against infectious diseases at the mucosa. *Mucosal Immunol.*
2009;2(5):403-11.

35. Duhen T, and Campbell DJ. IL-1β promotes the differentiation of polyfunctional
human CCR6+CXCR3+ Th1/17 cells that are specific for pathogenic and
commensal microbes. *J Immunol.* 2014;193(1):120-9.

36. Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia
A, et al. Surface phenotype and antigenic specificity of human interleukin 17-
producing T helper memory cells. *Nature immunology.* 2007;8(6):639-46.

37. Menze BH, Kelm BM, Masuch R, Himmelreich U, Bachert P, Petrich W, et al. A
comparison of random forest and its Gini importance with standard chemometric
methods for the feature selection and classification of spectral data. *BMC
Bioinformatics.* 2009;10(1):213.

38. Rahman KS, Darville T, Russell AN, O'Connell CM, Wiesenfeld HC, Hillier SL, et
al. Comprehensive Molecular Serology of Human Chlamydia trachomatis
Infections by Peptide Enzyme-Linked Immunosorbent Assays. *mSphere.*
2018;3(4).

39. Johansson M, Schön K, Ward M, and Lycke N. Studies in knockout mice reveal
that anti-chlamydial protection requires TH1 cells producing IFN-gamma: is this
true for humans? *Scand J Immunol.* 1997;46(6):546-52.

40. De Simone G, Mazza EMC, Cassotta A, Davydov AN, Kuka M, Zanon V, et al.
CXCR3 Identifies Human Naive CD8(+) T Cells with Enhanced Effector
Differentiation Potential. *J Immunol.* 2019;203(12):3179-89.

41. Belay T, Eko FO, Ananaba GA, Bowers S, Moore T, Lyn D, et al. Chemokine and
chemokine receptor dynamics during genital chlamydial infection. *Infect Immun.*
2002;70(2):844-50.

42. Olive AJ, Gondek DC, and Starnbach MN. CXCR3 and CCR5 are both required for T cell-mediated protection against C. trachomatis infection in the murine genital mucosa. *Mucosal Immunol.* 2011;4(2):208-16.

43. Sampath P, Moideen K, Ranganathan UD, and Bethunaickan R. Monocyte Subsets: Phenotypes and Function in Tuberculosis Infection. *Front Immunol.* 2018;9:1726.

44. LaVerda D, Albanese LN, Ruther PE, Morrison SG, Morrison RP, Ault KA, et al. Seroreactivity to Chlamydia trachomatis Hsp10 correlates with severity of human genital tract disease. *Infect Immun.* 2000;68(1):303-9.

45. El Hakim EA, Gordon UD, and Akande VA. The relationship between serum Chlamydia antibody levels and severity of disease in infertile women with tubal damage. *Arch Gynecol Obstet.* 2010;281(4):727-33.

46. Pantelic M, Kim YJ, Bolland S, Chen I, Shively J, and Chen T. Neisseria gonorrhoeae kills carcinoembryonic antigen-related cellular adhesion molecule 1 (CD66a)-expressing human B cells and inhibits antibody production. *Infect Immun.* 2005;73(7):4171-9.

47. Lee C, Guinn BA, Brooks SE, Richardson D, and Orchard K. CD66a (CEACAM1) is the only CD66 variant expressed on the surface of plasma cells in multiple myeloma: a refined target for radiotherapy trials? *Br J Haematol.* 2010;149(5):795-6.

48. Fankhauser SC, and Starnbach MN. PD-L1 limits the mucosal CD8+ T cell response to Chlamydia trachomatis. *J Immunol.* 2014;192(3):1079-90.

49. Cockwell P, Calderwood JW, Brooks CJ, Chakravorty SJ, and Savage CO. Chemoattraction of T cells expressing CCR5, CXCR3 and CX3CR1 by proximal tubular epithelial cell chemokines. *Nephrol Dial Transplant.* 2002;17(5):734-44.

50. Poston TB, Lee DE, Darville T, Zhong W, Dong L, O'Connell CM, et al. Cervical Cytokines Associated With Chlamydia trachomatis Susceptibility and Protection. *J Infect Dis.* 2019;220(2):330-9.

51. Liu H, and Rohowsky-Kochan C. Regulation of IL-17 in human CCR6+ effector memory T cells. *J Immunol.* 2008;180(12):7948-57.

52. Ogongo P, Tezera LB, Ardain A, Nhamoyebonde S, Ramsuran D, Singh A, et al. Tissue-resident-like CD4+ T cells secreting IL-17 control Mycobacterium tuberculosis in the human lung. *J Clin Invest.* 2021;131(10).

53. Borkner L, Curham LM, Wilk MM, Moran B, and Mills KHG. IL-17 mediates protective immunity against nasal infection with Bordetella pertussis by mobilizing neutrophils, especially Siglec-F(+) neutrophils. *Mucosal Immunol.* 2021;14(5):1183-202.

54. Gérard HC, Stanich JA, Oszust CE, Whittum-Hudson JA, Carter JD, Schumacher HR, et al. Functional CCR5 receptor protects patients with arthritis from high synovial burden of infecting Chlamydia trachomatis. *Am J Med Sci.* 2010;340(6):448-51.

55. Ogendi BMO, Bakshi RK, Sabbaj S, Brown L, Lee JY, Kapil R, et al. Distinct peripheral vs mucosal T-cell phenotypes in chlamydia-infected women. *Am J Reprod Immunol.* 2017;78(6).

56. Ficarra M, Ibana JS, Poretta C, Ma L, Myers L, Taylor SN, et al. A distinct cellular profile is seen in the human endocervix during Chlamydia trachomatis infection. *Am J Reprod Immunol.* 2008;60(5):415-25.

57. Gattinoni L, Speiser DE, Lichterfeld M, and Bonini C. T memory stem cells in health and disease. *Nat Med.* 2017;23(1):18-27.

58. Singh SP, Parween F, Edara N, Zhang HH, Chen J, Otaizo-Carrasquero F, et al. Human CCR6+ Th Cells Show Both an Extended Stable Gradient of Th17 Activity and Imprinted Plasticity. *J Immunol.* 2023;210(11):1700-16.

59. Sun M, Phan JM, Kieswetter NS, Huang H, Yu KKQ, Smith MT, et al. Specific CD4(+) T cell phenotypes associate with bacterial control in people who 'resist' infection with Mycobacterium tuberculosis. *Nature immunology.* 2024;25(8):1411-21.

60. Li Y, Warren JA, Poston TB, Clutton G, Shaw FR, Conrad SZ, et al. Chlamydia trachomatis induces low-frequency, sustained CD4 T cell responses in most women, predominantly targeting chlamydial protease-like activity factor, CPAF. *J Infect Dis.* 2024.

61. Bunk S, Schaffert H, Schmid B, Goletz C, Zeller S, Borisova M, et al. Chlamydia pneumoniae-induced memory CD4+ T-cell activation in human peripheral blood correlates with distinct antibody response patterns. *Clin Vaccine Immunol.* 2010;17(5):705-12.

62. Safar HA, Mustafa AS, Amoudy HA, and El-Hashim A. The effect of adjuvants and delivery systems on Th1, Th2, Th17 and Treg cytokine responses in mice immunized with Mycobacterium tuberculosis-specific proteins. *PLoS One.* 2020;15(2):e0228381.

63. Nguyen N, Guleed S, Olsen AW, Follmann F, Christensen JP, and Dietrich J. Th1/Th17 T cell Tissue-Resident Immunity Increases Protection, But Is Not Required in a Vaccine Strategy Against Genital Infection With Chlamydia trachomatis. *Front Immunol.* 2021;12:790463.

64. Poston TB, Girardi J, Kim M, Zwarycz P, Polson AG, Yount K, et al. Intranasal immunization with CPAF combined with cyclic-di-AMP induces a memory CD4 T cell response and reduces bacterial burden following intravaginal infection with Chlamydia muridarum. *bioRxiv.* 2024.

65. Poston TB, Girardi J, Polson AG, Bhardwaj A, Yount KS, Jaras Salas I, et al. Viral-vectored boosting of OmcB- or CPAF-specific T-cell responses fail to enhance protection from Chlamydia muridarum in infection-immune mice and elicits a non-protective CD8-dominant response in naïve mice. *Mucosal Immunol.* 2024.

66. Bagwell CB, Hunsberger B, Hill B, Herbert D, Bray C, Selvanantham T, et al. Multi-site reproducibility of a human immunophenotyping assay in whole blood and peripheral blood mononuclear cells preparations using CyTOF technology coupled with Maxpar Pathsetter, an automated data analysis system. *Cytometry B Clin Cytom.* 2020;98(2):146-60.

67. Johnson WE, Li C, and Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics.* 2006;8(1):118-27.

68. Stanley N, Stelzer IA, Tsai AS, Fallahzadeh R, Ganio E, Becker M, et al. VoPo leverages cellular heterogeneity for predictive modeling of single-cell data. *Nature Communications.* 2020;11(1):3738.

967   69.   Bruggner RV, Bodenmiller B, Dill DL, Tibshirani RJ, and Nolan GP. Automated
968   identification of stratifying signatures in cellular subpopulations. *Proceedings of the*
969   *National Academy of Sciences.* 2014;111(26):E2770-E7.

970