Body weight, gonadectomy, and other risk factors for diagnosis of osteoarthritis in companion dogs.

Jessica L Graves, MS[1†].; Brennen A McKenzie, VMD[1†*]; Zane Koch, BS[1]; Alexander Naka, PhD[1]; Nathaniel Spofford, MPH[2]; JoAnn Morrison, DVM, DACVIM[2]

[1]Loyal, San Francisco, CA, Unites States
[2]Banfield Pet Hospital, Portland, OR, United States

† These authors contributed equally to this work.

*Correspondence:
Brennen McKenzie
brennen@loyalfordogs.com

Keyword: dogs, osteoarthritis, degenerative joint disease, risk factors, incidence

Abstract

OBJECTIVE
To evaluate age, sex, body weight, breed, neuter status, and age at neutering as risk factors for diagnosis of osteoarthritis in companion dogs

ANIMALS
Dogs seen as patients at Banfield Pet Hospital in the United States from 1998-2019 with a date of death in 2019. The final cohort consisted of 131,140 dogs.

METHODS
In this retrospective cohort study, Cox proportional hazard models were used to test for associations between osteoarthritis incidence and age at baseline, sex, maximum body weight, maximum body condition score, neuter status, and age at neutering. The same model was used to test these associations in 12 representative breeds, chosen based on breed weight and sample size.

RESULTS
Older age, higher adult body weight, gonadectomy, and younger age at gonadectomy were significantly associated with higher risks of osteoarthritis in the total cohort and in all 12 breeds evaluated. Higher body condition score and sex were also significantly associated with osteoarthritis but with minimal effect sizes in the overall cohort, and these risk factors were not consistently significant in all breeds tested.

CLINICAL RELEVANCE
These results will assist veterinarians in identifying dogs at higher risk for osteoarthritis and applying appropriate diagnostic, preventative, and treatment interventions. An understanding of

45 potentially modifiable risk factors, such as body condition, and neutering, will support evidence-
46 based discussions with dog owners about risk management in individual patients.

Introduction

Musculoskeletal pain and lameness are among the most common clinical problems seen
in companion dogs.(1–3) A large proportion of these cases involve osteoarthritis (OA), the most
common joint disorder in dogs. There is inconsistency in literature regarding the terminology for
this condition. Terms such as "osteoarthritis," osteoarthrosis, and "degenerative joint disease"
may be used interchangeably, or some authors may use different terms for conditions with
specific etiology or pathogenesis. In this report, we use the term "osteoarthritis" to refer to a
diagnosis of one of the following structured diagnostic codes: osteoarthritis, arthritis or
degenerative joint disease – excluding rheumatoid, septic or immune mediated arthritic
conditions.

Prevalence of OA ranges from 2.5% to over 80% depending on study methods and
population characteristics.(1,4,5) This variation likely reflects both true differences in prevalence
between populations, related to differences in breed, age, husbandry, and other causal factors, as
well as differences in diagnostic methods and case definitions between studies.(6)

As a progressive, incurable condition, OA can affect comfort and quality of life for a
substantial proportion of a patient's life.(4) This condition compromises the welfare of affected
dogs and places a significant burden on human caregivers.(7) Musculoskeletal pain and
locomotor dysfunction, often due to OA, are also among the most frequent reasons for euthanasia
in dogs.(8–11)

Currently, the primary approach to mitigating the negative impact of canine OA is
treatment once clinical signs have manifested. Earlier detection and interventions to prevent or
delay the development of OA could have significant benefits. Effective approaches for
preventing or delaying the development of OA depend on a clear understanding of relevant risk
factors and identifying individuals at increased risk.

Many putative risk factors have been associated with development and progression of
canine OA.(6) There is strong evidence for the role of genetic factors influencing both individual
risk and breed differences in susceptibility to OA. These factors are often associated indirectly
with OA, causing predisposition conditions such as hip or elbow dysplasia or a propensity for
cranial cruciate ligament rupture, which then lead to development of arthritis.(6,12)

Body weight is another factor associated with OA risk. However, studies often do not
clearly distinguish between body size and body condition. Larger breeds appear to be at greater
risk, as do individuals who are overweight or obese, but the relationship between these different
body-size variables is not always clear.(6,13–15)

Osteoarthritis is considered a disease of aging, and increased age is often associated with
increased OA prevalence. This association is potentially complicated, however, by the lack of
surveillance and diagnostic markers of early, pre-clinical joint disease. Predisposing conditions
and early OA may be present undetected in young dogs while older individuals may be more

likely to be diagnosed with OA because of greater diagnostic attention or because the condition
has progressed to more apparent clinical signs.(6)
92
93   The evidence is limited and conflicting for many potential OA risk actors. Sex, for
94 example, is often associated with OA prevalence, but both males and females have been reported
95 to be at increased risk and the potential for confounding by body size, activity, and neuter status
96 is high.(6)
97
One of the most debated risk factors for OA is neuter status.(6,16–19) While most reports
indicate neutered dogs are at higher risk than intact dogs, the details of the relationship between
neuter status and OA are unclear. For example, this association seems to be consistently true for
large-breed dogs and less often found in smaller breeds.(18) Neutered dogs are also at greater
risk for obesity, so the degree to which the relationship between neutering and OA is mediated or
confounded by body condition is uncertain.(6,20–22)
104
Some studies find that age at neutering influences the impact of gonadectomy on the risk
of OA and predisposing conditions, such as cruciate ligament disease.(19,23) This suggests
gonadal hormones are protective primarily through effects of skeletal development, and
neutering after puberty or skeletal maturity may be less likely to promote OA. However, other
studies report residual increased risk in dogs neutered after skeletal maturity and suggest that
gonadal hormones may have an ongoing protective effect.(24) There is also significant variation
in the existence and strength of associations between neutering and orthopedic disease found in
different breeds and different research studies.(18) Many other factors, including diet, activity
patterns, and even birth month have been associated with the risk of OA, but detailed causal links
have not been clearly identified.(6)
115
From a preventative-medicine perspective, risk factors for OA can be considered
modifiable or non-modifiable. Most genetic factors are not directly modifiable in individuals,
and the risk presented by specific genotypes, breed, and conformation are fixed at birth or during
development. Recent advances in the study of epigenetics suggest it may be possible to mitigate
the impact of some genes through environmental modification of the regulation of gene activity,
but clinical interventions for doing so have not yet been validated.(25,26) On a population level,
genetic factors which influence the occurrence of OA can potentially be modified by selective
breeding targeting both individual genes associated with increased OA risk as well as
conformations that predispose to the disease.(25,26)
125
126   Other factors associated with OA are clearly modifiable in individuals, including body
127 weight and body condition, diet, activity patterns, and neutering practices.
128
Common dietary recommendations to delay or prevent OA include reduced feeding to
prevent obesity and modulate skeletal development in growing puppies. Lifelong caloric
restriction in a cohort of Labrador retrievers delayed the onset of hip OA and reduced the
severity of the disorder.(27,28) Less definitive effects were reported for elbow and shoulder
OA.(29,30) Whether these effects were due solely to differences in body condition or other
influences of caloric restriction is uncertain.
135

136   Experimental studies have also shown that reduction in caloric and calcium content of
137 diets can reduce the risk of developmental abnormalities in giant-breed dogs, such as hip
138 dysplasia, that frequently lead to OA later in life.(31,32) Other nutritional interventions may also
139 influence development of OA and predisposing conditions,(33) but there is still significant
140 uncertainty about the efficacy of most dietary approaches.
141
Greater clarity about the role of key risk factors in development of canine OA would be
useful for informing preventative strategies. The purpose of this study was to examine selected
risk factors for development of OA in a large, retrospective cohort study of companion dogs
using medical record data from primary care veterinary practices. We examined previously
reported risk factors including age, sex, breed, body weight, and body condition. We also sought
to further investigate the relationships between OA, neuter status and age at neutering as well as
the variability of these relationships among breeds.

Materials and Methods

Data summary and processing

By the end of 2019 the Banfield Pet Hospital network included 1,084 primary care small
animal hospitals in 42 U.S. states, the District of Columbia and Puerto Rico. All hospitals used
the same proprietary practice information management system (PIMS; PetWare®) to enter
patient and visit information and the resulting electronic medical records were uploaded nightly
to a central data warehouse. The PIMS contained both structured and unstructured fields for data
entry. Structured data included diagnoses and clinical signs, physical examination findings, and
invoiced services and medications; unstructured data included narrative text related to subjective
and objective observations, patient assessment and treatment plan.
159
All in-network visits from dogs with a death date in 2019 were extracted from the data
warehouse, with visit information ranging from 1998-2019. Patient information for each visit
included breed, sex, neuter status, visit age, and visit weight. Visit information included the
location and reason for the visit, all structured clinical signs and diagnoses, examination findings
related to musculoskeletal issues and body condition, and any invoiced items or services related
to the diagnosis or treatment of osteoarthritis. Invoiced items or services related to in-hospital
euthanasia were also extracted in order to more accurately identify patient death date.
167
To improve data quality the visit-level data was cleaned and evaluated for data entry
errors. First, only visits in which the dog received a physical examination by a veterinarian were
included. Some data were removed due to likely data entry errors. These include visits in which
there were discrepant entries, visit dates occurring after 2019 or weights recorded over 300 lbs,
or dogs who had implausible data values (e.g., negative age). To minimize cohort effects, only
dogs born after 1997 were included for analysis. Lastly, the body condition score (BCS)
assessment has changed over time (e.g., the use of 3-point (clinician-based assessment), 5-point
and 9-point BCS scales), scores were harmonized and converted into a single 9-point scale
(**Table 1**).
177
Osteoarthritis was defined as the earliest diagnosis of the following structured diagnostic
codes: osteoarthritis, arthritis or degenerative joint disease – excluding rheumatoid, septic or

immune mediated arthritic conditions. Follow-up time was calculated as the number of years
between a dog's age at their first visit and their age at OA diagnosis or death, whichever
occurred first – dogs with follow-up times greater than 20 years were removed. In order to
capture incidence of aging-related OA, dogs who were not observed into maturity (2 years and
up) or who already had OA at their first visit were not included for analysis.
185
Desexing status was defined based on veterinarian assessment and if the dog was
gonadectomized at a Banfield clinic. Age at desexing was defined as age during a visit with a
desexing procedure. Age at desexing was only ascertained for dogs desexed at a Banfield clinic.
To preserve temporality between desexing as a risk factor and OA incidence, dogs desexed after
being diagnosed with OA were treated as intact. Dogs desexed after 5 years of age were not
included in age at desexing analyses, as desexing later has a higher likelihood of being a
therapeutic rather than an elective procedure.
193
Multiple weight and BCS metrics were created. To estimate the effects of maximum body
size or condition, the 75th-percentile of adult (2+ years of age) body weights and BCS were used
to approximate a 'maximum' weight or BCS. 75th percentiles were used to minimize the
influence of outliers. Median weight and BCS between ages 1.5 and 2.5 years of age were used
to approximate the weight and BCS at developmental maturity. Not all dogs were seen between
1.5 and 2.5 years of age. In these cases, weight or BCS at desexing (so long as desexing occurred
after full development, e.g., 2+ years of age) was used.
201
202   To estimate effects of weight gain after desexing, weight and BCS after desexing was
203 defined as the 75th-percentile of weight and BCS after desexing and 2+ years of age. Percent
204 change in weight and BCS after desexing was calculated as 100*(observation after desexing -
205 observation at maturity)/observation at maturity.
206
207   The majority of dogs seen at Banfield are on a wellness plan (a set of pre-paid services
208 meant to enhance the provision of preventive care services). Wellness plan data is structured
209 within the system so the wellness plan variable was converted to a binary variable encoding if a
210 dog ever was on a wellness plan. Dogs whose wellness plan started at or after OA diagnosis were
211 treated as "never" having been on a plan.
212
213

214 **Table 1**: Body condition score conversion methodology

Point scale	Original BCS value	New BCS value	Classified as
5-point & 9-point	Body Condition Score - Thin 1	1	Too thin
5-point	Body Condition Score - Underweight 2	2	Too thin
3-point	Nutrition/Weight Control Advice - Thin 1	2	Too thin
9-point	Body Condition Score - Too Thin 2	2	Too thin
9-point	Body Condition Score - Too Thin 3	3	Too thin
9-point	Body Condition Score - Ideal 4	4	Ideal
3-point	Nutrition/Weight Control Advice - Normal 2	4	Ideal
5-point	Body Condition Score - Ideal 3	5	Ideal
9-point	Body Condition Score - Ideal 5	5	Ideal
9-point	Body Condition Score - Above Ideal 6	6	Overweight
5-point	Body Condition Score - Overweight 4	7	Overweight
9-point	Body Condition Score - Overweight 7	7	Overweight
3-point	Nutrition/Weight Control Advice - Heavy 3	7	Overweight
5-point	Body Condition Score - Obese 5	8	Obese
9-point	Body Condition Score - Obese 8	8	Obese
9-point	Body Condition Score - Obese 9	9	Obese

215
216
Statistical methods

Descriptive statistics

Mean, standard deviation (SD), median, 25th and 75th percentiles, and ranges were
calculated for continuous measures: age at baseline, mature and maximum weight, age at OA or
censorship/death and duration of follow up (years). Observation counts and proportions were
calculated for categorical variables: sex (Female or Male), mature and maximum BCS (1-3=Too
thin; 4-5=Ideal; 6-7=Overweight, 8-9=Obese), gonadectomy status (desexed or intact), age at
desexing (<6 months, 6-12 months, 12-24 months, 24-60 months), and ever had a Banfield
wellness plan (Yes or No). For all variables, the number of missing values were reported. All
descriptive statistics are reported for the total cohort as well as stratified by OA diagnosis.
227
Cox proportional hazards

Survival analysis was used to characterize the incidence of OA in companion dogs. The
analyses were conducted using duration of follow-up as the measure of survival time, calculated
as the years between a dog's first visit and their OA onset or death. This choice of follow-up
time as the survival time scale enables easier estimation of age as a risk factor for OA onset.
Dogs were right-censored at death.
234
A Cox proportional hazards model[34] was used to test joint associations of OA risk
factors. In particular, age at baseline, max weight, max BCS, desexing status and sex were used
as risk-factors. The Cox model was stratified, separating dogs on (76.3%) and not on (23.7%)
wellness plans (**Table 2**). As the dogs on wellness plans were more closely observed on average,
the likelihood of OA diagnosis in these dogs was higher (**Supplemental Figure 1**), mostly
likely due potential detection bias, and for this reason a stratified Cox proportional hazards

model was used. This stratified model accounts for this wellness plan-associated surveillance
bias by allowing the baseline hazard of dogs in these groups to be different.(35) On average,
dogs on wellness plans visited Banfield 25.06 times (SD 18.54) while dogs not on a wellness
plan visited only 4.67 times (SD 6.85).

After fitting the Cox proportional hazards model, using the *coxph* function from the
survival package(36) in R, the proportional hazards assumption was assessed using *cox.zph* and
visualization of residuals. The assumption being met, hazard ratios (HR) and their 95%
confidence intervals (CI) were estimated to test the magnitude and direction of each risk-factor
on time to OA diagnosis. To account for the risk-factors having varying scales of values,
standardized HRs were calculated for continuous risk factors (e.g., age at baseline) by
standardizing the risk factor (mean centering and dividing by the standard deviation). For all
models, both the natural scale and standardized HRs and 95% CIs are reported, along with
concordance index and number of observations.

The marginal effects of each risk-factor on OA-free survival time were plotted to better
understand the magnitude of effect sizes. Marginal effects reflect the average effect of a given
risk factor, holding all other covariates in the model constant. To visualize effects of continuous
risk factors, predicted OA-free survival curves were estimated based on quartile values or
clinical relevance. To visualize marginal effects of categorical variables, each level was used to
predict OA-free survival curves. For risk factors not being directly visualized, continuous values
were assigned as cohort median values (e.g., median weight) and categorical variables were
assigned as a proportion (e.g., proportion male). As the majority of dogs were on wellness plans,
strata were assigned to being on a wellness plan. For all interaction models, median survival
times and their 95% CIs are also reported.

Breed-specific analyses

For breeds with at least 500 observations, binomial tests of proportion were used to
compare breed-specific OA incidence rates against the cohort-wide incidence rate, using
Bonferroni adjusted p-value to adjust for multiple comparisons. Given the diversity and size of
the cohort, risk factors of OA were also evaluated in 12 representative breeds, chosen based on
breed weight and sample size. First, the maximum body weight of all dogs in the cohort was split
into quartiles and then breeds were assigned to a weight quartile based on the mean maximum
weight of all dogs of that breed. Then, the top three breeds within each quartile with the most
samples were chosen for further analysis. This ensures representation of a wide range of dog
weights while not including breeds lacking sufficient sample size to make substantive
conclusions. The same stratified Cox regression model used to test primary risk factors of OA
was applied to each of these 12 breeds.

All analyses were performed in R version 4.1.2. Type I error was set to $\alpha = 0.05$ for all
statistical tests.

Results

Descriptive statistics

Descriptive statistics of the final cohort resulting from the data processing steps outlined
are reported in **Table 2.** The final cohort included 131,140 dogs, 31,365 (23.9%) of which were
diagnosed with OA during the course of the study. The cohort of dogs had an average age
entering the study of 5.12 years (SD = 4.33), average mature weight of 41.8 lbs (SD=31.0), and
63,541 (48.5%) dogs were female. There were 318 unique breeds present in the study, with
Labrador Retrievers (n=11,718), Chihuahuas (n=10,147) and Yorkshire Terriers (n=6,570) as the
most frequent breeds observed.
293
Not all variables generated during the data processing steps outlined above contain
observable data. The relatively high proportion of missing data in mature BCS are due to strict
construct definitions to generate this variable. Mature BCS was defined as the median BCS score
between 1.5 to 2.5 years – many dogs did not have visits during this window in which BCS was
assessed. Similarly, age at desexing could only be quantified for dogs who were gonadectomized
at a Banfield hospital, which could be confidently matched to a visit date. Approximately 20% of
dogs in this cohort were desexed at Banfield, of which 1% were desexed after receiving an OA
diagnosis and 9.4% were desexed > 5 years of age. To preserve temporality and capture patterns
of voluntary desexing, age at desexing > 5 years of age or after OA were set to missing and are
therefore not included in age at desexing analyses.
304
305   In order to fully leverage all available data, only observed data of relevant variables were
306   included in statistical analyses.
307
308   **Table 2.** Cohort descriptive statistics

| | | | Osteoarthritis | |
	Total **(N=131,140)**	**Diagnosed** **(N=31,365)**	**Never diagnosed** **(N=99,775)**
Age (years) at baseline			
Mean (SD)	5.12 (4.33)	4.61 (4.04)	5.29 (4.41)
Median (Q1, Q3)	4.41 (0.906, 8.24)	4.00 (0.523, 7.77)	4.73 (0.999, 8.79)
Range	0.00274 - 21.0	0.00274 - 20.0	0.00274 - 21.0
Sex			
Female	63,541 (48.5%)	15,461 (49.3%)	48,080 (48.2%)
Male	67,574 (51.5%)	15,904 (50.7%)	51,670 (51.8%)
Missing	25 (0.0%)	0 (0%)	25 (0.0%)
Mature[1] weight (lbs)			
Mean (SD)	41.8 (31.0)	50.9 (30.7)	38.5 (30.5)
Median (Q1, Q3)	35.8 (14.0, 64.6)	52.7 (21.4, 71.8)	26.7 (12.6, 61.1)
Range	2.10 - 224	2.20 - 194	2.10 - 224
Missing	85,898 (65.5%)	19,419 (61.9%)	66,479 (66.6%)
Max[2] weight (lbs)			
Mean (SD)	42.0 (32.4)	52.3 (33.0)	38.8 (31.5)
Median (Q1, Q3)	29.6 (14.6, 66.6)	52.8 (20.8, 76.6)	24.8 (13.5, 62.0)
Range	2.00 - 262	2.44 - 230	2.00 - 262
Mature[1] BCS			

Too thin (1-3)	429 (0.3%)	79 (0.3%)	350 (0.4%)
Ideal (4-5)	8,626 (6.6%)	480 (1.5%)	8,146 (8.2%)
Overweight (6-7)	4,672 (3.6%)	1,223 (3.9%)	3,449 (3.5%)
Obese (8-9)	219 (0.2%)	47 (0.1%)	172 (0.2%)
Missing	117,194 (89.4%)	29,536 (94.2%)	87,658 (87.9%)
Max[2.] BCS			
Too thin (1-3)	2,279 (1.7%)	153 (0.5%)	2,126 (2.1%)
Ideal (4-5)	60,734 (46.3%)	13,564 (43.2%)	47,170 (47.3%)
Overweight (6-7)	52,596 (40.1%)	15,947 (50.8%)	36,649 (36.7%)
Obese (8-9)	5,433 (4.1%)	1,658 (5.3%)	3,775 (3.8%)
Missing	10,098 (7.7%)	43 (0.1%)	10,055 (10.1%)
Desexed			
Yes	112,704 (85.9%)	29,046 (92.6%)	83,658 (83.8%)
No	18,411 (14.0%)	2,319 (7.4%)	16,092 (16.1%)
Missing	25 (0.0%)	0 (0%)	25 (0.0%)
Age at desexing grouped			
<6 mos	13,178 (10.0%)	4,060 (12.9%)	9,118 (9.1%)
6-12 mos	7,956 (6.1%)	2,029 (6.5%)	5,927 (5.9%)
12-24 mos	1,590 (1.2%)	342 (1.1%)	1,248 (1.3%)
24-60 mos	1,862 (1.4%)	387 (1.2%)	1,475 (1.5%)
Missing	106,554 (81.3%)	24,547 (78.3%)	82,007 (82.2%)
Age (years) at OA or death			
Mean (SD)	11.2 (3.49)	10.7 (2.91)	11.4 (3.64)
Median (Q1, Q3)	11.5 (9.05, 13.7)	10.9 (8.88, 12.8)	11.7 (9.12, 13.9)
Range	0.353 - 22.0	0.353 - 20.5	1.34 - 22.0
Duration of follow up (years)			
Mean (SD)	6.10 (4.07)	6.11 (3.82)	6.10 (4.15)
Median (Q1, Q3)	5.79 (2.62, 9.27)	6.03 (2.83, 9.07)	5.71 (2.55, 9.34)
Range	0.00274 - 20.0	0.00274 - 18.5	0.00274 - 20.0
Had wellness plan			
Yes	100,086 (76.3%)	28,389 (90.5%)	71,697 (71.9%)
No	31,054 (23.7%)	2,976 (9.5%)	28,078 (28.1%)

309 [1.] 75th percentile of observations > 2 years of age
310 [2.] Median of observations within 1.5 and 2.5 years of age where available, if not, observation at desexing used
311 SD=Standard deviation; Q1=25th percentile; Q3=75th percentile
312
Primary risk factors

To explore if effects of previously reported risk factors were recapitulated in this
expansive cohort, we fit stratified Cox proportional hazards model using age at baseline, desexed
status (desexed/intact), sex (male/female), adult weight and adult BCS as our primary risk
factors, and treated wellness plan as a stratifying variable. Stratification along the presence of
being on a wellness plan was supported by evidence of a surveillance (or detection) bias in dogs
who were on wellness plans **(Supplemental Figure 1)**.
320
321   **Table 3** and **Figure 1** show that older age (Std. HR=3.70 (3.64-3.76)), higher adult body
322 weight (Std. HR=1.83 (1.81-1.85)), and desexing (Std. HR=1.37 (1.31-1.43)) are statistically
323 significantly associated with higher risks of OA in this cohort. While hazard ratios for adult BCS
324 and sex were both statistically significant, these effect sizes are minimal (**Figure 1**).

Table 3. Stratified Cox proportional hazard model testing the associations of primary risk factors on OA-free survival.

Variable	HR (95% CI)	Std. HR (95% CI)	p-value	Concordance	N obs
Age at baseline	1.35 (1.35-1.36)	3.70 (3.64-3.76)	<0.001***	0.79	121,035
Max weight	1.02 (1.02-1.02)	1.83 (1.81-1.85)	<0.001***		
Max BCS	1.02 (1.01-1.03)	1.02 (1.01-1.03)	<0.001***		
Desexed	1.37 (1.31-1.43)	1.37 (1.31-1.43)	<0.001***		
Male	0.91 (0.89-0.93)	0.91 (0.89-0.93)	<0.001***		

*Std. HR= Standardized continuous variables (mean centered, scaled by standard deviation).
Wellness plan treated as stratifying variable.

Figure 1. Predicted marginal OA-free survival curves of primary risk factors of OA.
Assignments for generating predicted survival curves: median values for continuous covariates
not being directly tested, proportions for binary variables, and assuming presence of a wellness
plan. Quantiles were used for continuous variable groupings being visualized, except for BCS,
which were determined based on clinical relevance.

Breed-specific risk factors: Are risk factors consistent within breeds?

To explore if the effects of these risk factors were consistent across breeds, the same
primary risk factor model was fit to a subset of breeds. The top three most frequent breeds within
each weight quartile were selected as an analysis subset (see **Statistical Methods: Breed-**
Specific Analyses section for more information). **Supplemental Table 1, Supplemental Figure**
2 and **Supplemental Figure 3** the 12 representative breeds as well as breed-specific OA
incidence tested against the overall cohort OA incidence rate.
346

Figure 2 summarizes these results using a dot-and-whisker plot of the standardized HR
and 95% CI for each of the risk factors within each breed. Higher weight and older age are
consistently associated with higher risks of OA across breeds. However, the effect of desexing
on OA incidence shows variability across breeds, where desexing does not appear to be a risk
factor for Yorkshire Terriers, Maltese, Shih Tzus, Pugs, Dachshunds and Beagles. The effects of
BCS also differ across breeds, where effects are more pronounced in smaller breeds. HRs and
95% CIs reported in **Supplemental Table 2.**
354

355
Figure 2. Dot-and-whisker plot results of breed-specific analysis of primary risk factors of OA.
Dots and whiskers reflect the standardized HR and 95% CIs associated with each risk factor,
respectively. Each panel reflects the model fit to an individual breed, where panels are grouped
by weight groups. Color assignments designate statistical significance of the HR.

Does desexing moderate the effect of max body weight on OA?

Both weight and desexing are independent risk factors of OA. However, it has been
reported that dogs who are desexed may be more likely to develop obesity,(21,22,37) which may
increase the risk of OA. To test if the influence of higher body weight on OA-free survival
differed by desexing status, the interaction between adult weight and desexing status was tested.

There was a statistically significant interaction between desexing and adult weight (Std.
HR=1.07 (1.03, 1.11), **Table 4**), suggesting that the increased risk of OA in larger dogs is greater
in those that are desexed compared to intact dogs. **Figure 3** shows the predicted survival curves
as well as the predicted median survival times for each weight class and desexing group. Both
desexed and intact dogs show that higher body weight is associated with higher OA risk;
however, this risk appears slightly larger in desexed dogs compared to intact dogs.

Table 4. Stratified Cox proportional hazard model testing the interaction between desexing and
maximum adult weight and BCS.

Model	Variable	HR (95% CI)	Std. HR (95% CI)	p-value	Concordance	N obs
Desexing x Weight	Age at baseline	1.35 (1.35-1.36)	3.70 (3.64-3.76)	<0.001***	0.79	121,035
	Max weight	1.02 (1.02-1.02)	1.73 (1.67-1.79)	<0.001***		
	Max BCS	1.02 (1.01-1.03)	1.02 (1.01-1.03)	<0.001***		
	Desexed	1.23 (1.15-1.32)	1.34 (1.29-1.40)	<0.001***		
	Male	0.91 (0.89-0.93)	0.91 (0.89-0.93)	<0.001***		
	Desexed x max weight	1.00 (1.00-1.00)	1.07 (1.03-1.11)	<0.001***		
Desexing x BCS	Age at baseline	1.35 (1.35-1.36)	3.70 (3.64-3.76)	<0.001***	0.79	121,035
	Max weight	1.02 (1.02-1.02)	1.83 (1.81-1.85)	<0.001***		
	Max BCS	1.02 (0.99-1.06)	1.03 (0.99-1.07)	0.186		
	Desexed	1.42 (1.14-1.76)	1.37 (1.31-1.43)	0.002**		
	Male	0.91 (0.89-0.93)	0.91 (0.89-0.93)	<0.001***		
	Desexed x max BCS	0.99 (0.96-1.03)	0.99 (0.95-1.04)	0.754		

*Std. HR= Standardized continuous variables (mean centered, scaled by standard deviation).
Wellness plan treated as stratifying variable.

Figure 3. Predicted marginal OA-free survival curves to visualize interaction effects between
weight and desexing status. Color indicates maximum weight values. Median survival times and
95% CIs are reported. To generate predictions, median values were used for continuous
covariates, proportions for binary variables, and the presence of wellness plan were used.

Does weight gain increase risk of developing OA?

Limitations of the modeling approach used in **Table 2** and **Figure 1** include the lack of
temporal precedence – specifically, the inability to test subsequent weight gain after desexing
and its influence on OA incidence. To test this hypothesis, percent changes in maximum weight
within desexed dogs (calculated as 100*[maximum weight - mature weight]/mature weight) was
tested as a risk factor for OA.

Both main effects of percent weight change as well as the interaction with mature body
weight were tested in the full cohort and in a sensitivity analysis restricting to dogs whose age at
desexing was $\leq$ 2 years of age. Mature body weight was treated as a covariate as larger-breed
dogs are less likely to have larger percent changes in body weight compared to smaller-breed
dogs. To determine if weight changes were independent of desexing status, these analyses were
performed in intact dogs as well, where percent changes in weight reflected overall changes from
mature body weight to maximum body weight.

Results in **Table 5** show that positive percent change in body weight was associated with
higher risk of OA (full cohort Std. HR=1.12 (1.09-1.15); age desexing $\leq$ 2 Std. HR=1.12 (1.09-
1.15)). There was also a significant interaction effect between percent change in body weight
after desexing and mature body size (Std. HR=1.09 (1.05-1.12)), which held in the dogs desexed
$\leq$ 2 years of age. These results suggest that percent increases in body weight after desexing may
have stronger deleterious effects in larger dogs, compared to smaller dogs.

In order to illustrate these effects in more interpretable units, **Figure 4** shows the main
effects of percent changes in body weight after desexing, where every 50% increase in body
weight after desexing was associated with an approximately 1 year decrease in median time to
OA. **Figure 5** visualizes the finding that percent increases in body weight after desexing was

associated with higher OA risks in larger dogs compared to smaller dogs. It is important to note that 50% increases in body weight become less likely as mature body size increases and the use of 50% increments are used only to facilitate interpretation of the direction and magnitude of effects.

Table 5. Stratified Cox proportional hazard model testing both the main effects and interaction effects of percent change in body weight in desexed dogs, dogs desexed ≤ 2 years of age, and intact dogs, respectively. For dogs who were desexed, percent change in body weight is based on weight change exclusively after desexing.

Subset	Model	Variable	HR (95% CI)	Std. HR (95% CI)	p-value	Concordance	N obs
All desexed dogs	Main effect	Age at baseline	1.20 (1.15-1.26)	2.23 (1.82-2.74)	<0.001***	0.69	23,099
		Male	0.86 (0.82-0.90)	0.86 (0.82-0.90)	<0.001***		
		% change in weight after desexing	1.01 (1.01-1.01)	1.12 (1.09-1.15)	<0.001***		
		Mature weight	1.02 (1.02-1.02)	1.95 (1.91-2.00)	<0.001***		
	Interaction	Age at baseline	1.21 (1.15-1.26)	2.25 (1.83-2.76)	<0.001***	0.69	23,099
		Male	0.86 (0.82-0.91)	0.86 (0.82-0.91)	<0.001***		
		% change in weight after desexing	1.00 (1.00-1.01)	1.05 (1.01-1.09)	0.027*		
		Mature weight	1.02 (1.02-1.02)	1.86 (1.80-1.92)	<0.001***		
		% change in weight after desexing x mature weight	1.00 (1.00-1.00)	1.09 (1.05-1.12)	<0.001***		
Dogs desexed ≤ 2 years old	Main effect	Age at baseline	1.12 (0.98-1.28)	1.02 (1.00-1.05)	0.108	0.69	21,296
		Male	0.86 (0.82-0.90)	0.86 (0.82-0.90)	<0.001***		
		% change in weight after desexing	1.01 (1.01-1.01)	1.12 (1.09-1.15)	<0.001***		
		Mature weight	1.02 (1.02-1.02)	2.00 (1.95-2.05)	<0.001***		
	Interaction	Age at baseline	1.12 (0.98-1.28)	1.02 (1.00-1.05)	0.097	0.69	21,296
		Male	0.86 (0.82-0.91)	0.86 (0.82-0.91)	<0.001***		
		% change in weight after desexing	1.00 (1.00-1.01)	1.04 (1.00-1.08)	0.057		
		Mature weight	1.02 (1.02-1.02)	1.89 (1.83-1.96)	<0.001***		
		% change in weight after desexing x mature weight	1.00 (1.00-1.00)	1.09 (1.05-1.13)	<0.001***		
Intact Dogs	Main effect	Age at baseline	1.44 (1.30-1.59)	4.84 (3.12-7.50)	<0.001***	0.69	5,519
		Male	1.23 (1.04-1.44)	1.23 (1.04-1.44)	0.014*		
		% change in weight	1.01 (1.01-1.01)	1.19 (1.14-1.25)	<0.001***		
		Mature weight	1.02 (1.02-1.02)	1.70 (1.59-1.81)	<0.001***		
	Interaction	Age at baseline	1.45 (1.31-1.60)	4.97 (3.20-7.71)	<0.001***	0.69	5,519
		Male	1.26 (1.07-1.48)	1.26 (1.07-1.48)	0.006**		
		% change in weight	1.01 (1.00-1.01)	1.12 (1.05-1.20)	0.001**		
		Mature weight	1.01 (1.01-1.02)	1.57 (1.45-1.71)	<0.001***		
		% change in weight x mature weight	1.00 (1.00-1.00)	1.13 (1.05-1.22)	0.001**		

*Std. HR= Standardized continuous variables (mean centered, scaled by standard deviation).
Wellness plan treated as stratifying variable.

423
424 **Figure 4**. Predicted marginal OA-free survival curves illustrating the main effect of percent
425 weight changes after desexing. Color indicates percent change values. To generate predictions,
426 median values were used for continuous covariates, proportions for binary variables, and the
427 presence of wellness plan were used.
428

429
Figure 5. Predicted marginal OA-free survival curves illustrating the interaction effect between
percent weight change after desexing and mature body weight. Color indicates percent change
values. Median survival times and 95% CIs are reported. To generate predictions, median values
were used for continuous covariates, proportions for binary variables, and the presence of
wellness plan were used.

Does increased BCS influence OA risks?

As BCS is a distinct construct from weight or body size, percent changes in body
condition score after desexing was also tested as a risk factor for OA. Paralleling the approach
used in the percent change in weight models, both main effects of percent BCS change as well as
the interaction with mature BCS were tested in the full cohort and in a sensitivity analysis
restricting to dogs whose age at desexing was ≤ 2 years of age. Mature body BCS was treated as
a covariate as dogs with higher BCS at maturity are less likely to have larger BCS changes in
BCS compared to dogs with lower BCS at maturity. To determine if weight changes were
independent of desexing status, these analyses were performed in intact dogs as well, where
percent changes in BCS reflected overall changes from mature BCS to maximum BCS.

Results indicate increases in percent change in BCS are associated with higher risk of OA
(Std. HR=1.12 (1.05-1.20), **Table 6**), and that these effects may be stronger in dogs with higher
mature BCS (Std. HR=1.16 (1.04-1.29), **Table 6**). However, limited variability in BCS in this
cohort makes determining the clinical significance of these effects difficult. As seen in **Figure 6**

and **Figure 7**, standard errors from model predictions show significant overlap, except in dogs
who have higher mature BCS, where effects begin to appear strongest. However, it is important
to note that due to there being a ceiling in the BCS tool, dogs with mature BCS of 7 can only
experience at most a percent change increase of 28% (28%=100*(9-7)/7).
455
456   These analyses were also performed in intact dogs, where percent changes in BCS reflect
457 overall changes from mature BCS to maximum BCS to identify any changes across desexing
458 status, where percent changes in BCS were not statistically significantly associated with OA
459 incidence.
460
461

462 **Table 6**. Stratified Cox proportional hazard model testing both the main effects and interaction
463 effects of percent change in BCS in desexed dogs, dogs desexed ≤ 2 years of age, and intact
464 dogs, respectively. For dogs who were desexed, percent change in BCS is based on BCS change
465 exclusively after desexing.

Subset	Model	Variable	HR (95% CI)	Std. HR (95% CI)	p-value	Concordance	N obs
All desexed dogs	Main effect	Age at baseline	1.25 (1.12-1.39)	2.60 (1.64-4.12)	<0.001***	0.67	7,484
		Male	0.92 (0.82-1.04)	0.92 (0.82-1.04)	0.186		
		% change in BCS after desexing	1.00 (1.00-1.01)	1.12 (1.05-1.20)	<0.001***		
		Mature weight	1.02 (1.02-1.02)	1.69 (1.61-1.78)	<0.001***		
		Mature BCS	1.16 (1.08-1.24)	1.18 (1.09-1.29)	<0.001***		
	Interaction	Age at baseline	1.24 (1.12-1.38)	2.56 (1.61-4.05)	<0.001***	0.67	7,484
		Male	0.92 (0.82-1.04)	0.92 (0.82-1.04)	0.166		
		% change in BCS after desexing	1.00 (1.00-1.00)	1.01 (0.90-1.13)	0.831		
		Mature weight	1.02 (1.02-1.02)	1.69 (1.60-1.78)	<0.001***		
		Mature BCS	1.16 (1.08-1.25)	1.19 (1.09-1.29)	<0.001***		
		% change in BCS after desexing x mature BCS	1.00 (1.00-1.00)	1.16 (1.04-1.29)	0.008**		
Dogs desexed ≤ 2 years old	Main effect	Age at baseline	1.16 (0.80-1.66)	1.03 (0.96-1.11)	0.432	0.67	6,921
		Male	0.91 (0.81-1.03)	0.91 (0.81-1.03)	0.135		
		% change in BCS after desexing	1.00 (1.00-1.01)	1.14 (1.06-1.21)	<0.001***		
		Mature weight	1.02 (1.02-1.02)	1.72 (1.63-1.81)	<0.001***		
		Mature BCS	1.15 (1.07-1.24)	1.17 (1.08-1.27)	<0.001***		
	Interaction	Age at baseline	1.14 (0.80-1.64)	1.03 (0.95-1.11)	0.471	0.67	6,921
		Male	0.91 (0.81-1.03)	0.91 (0.81-1.03)	0.124		
		% change in BCS after desexing	1.00 (1.00-1.01)	1.03 (0.92-1.14)	0.647		
		Mature weight	1.02 (1.02-1.02)	1.71 (1.62-1.81)	<0.001***		
		Mature BCS	1.16 (1.07-1.25)	1.18 (1.08-1.28)	<0.001***		
		% change in BCS after desexing x mature BCS	1.00 (1.00-1.00)	1.16 (1.05-1.29)	0.005**		
Intact dogs	Main effect	Age at baseline	2.07 (1.52-2.82)	23.46 (6.15-89.40)	<0.001***	0.67	2,153
		Male	1.21 (0.73-2.01)	1.21 (0.73-2.01)	0.467		
		% change in BCS	1.00 (1.00-1.01)	1.13 (0.98-1.30)	0.085		
		Mature weight	1.02 (1.01-1.02)	1.67 (1.39-2.00)	<0.001***		
		Mature BCS	1.00 (0.82-1.21)	1.00 (0.80-1.25)	0.981		
	Interaction	Age at baseline	2.07 (1.51-2.82)	23.16 (6.02-89.13)	<0.001***	0.67	2,153
		Male	1.21 (0.73-2.02)	1.21 (0.73-2.02)	0.463		
		% change in BCS	1.00 (1.00-1.01)	1.12 (0.92-1.36)	0.267		
		Mature weight	1.02 (1.01-1.02)	1.66 (1.39-2.00)	<0.001***		
		Mature BCS	1.00 (0.81-1.24)	1.00 (0.79-1.28)	0.971		
		% change in BCS x mature BCS	1.00 (1.00-1.00)	1.02 (0.76-1.39)	0.876		

466 *Std. HR= Standardized continuous variables (mean centered, scaled by standard deviation).
467 Wellness plan treated as stratifying variable.
468
469

470
471
472 **Figure 6**. Predicted marginal OA-free survival curves illustrating the main effect of percent BCS
473 after desexing. Color indicates percent change values. To generate predictions, median values
474 were used for continuous covariates, proportions for binary variables, and the presence of
475 wellness plan were used.

476
Figure 7. Predicted marginal OA-free survival curves illustrating the interaction effect between
percent BCS after desexing and mature BCS. Color indicates percent change values. Median
survival times and 95% CIs are reported. To generate predictions, median values were used for
continuous covariates, proportions for binary variables, and the presence of wellness plan were
used.

Does age at desexing influence OA risks?

To test the hypothesis that age at desexing is a risk factor for OA, age at desexing was
added into the primary risk factor model. Age at desexing was tested as a continuous variable as
well as a categorical variable (<6 months, 6-12 months, 12-24 months, and 24-60 months). Age
at desexing was significantly inversely associated with OA risk (Std. HR=0.90 (0.86-0.94) **Table
7**). Categorical analysis shows that relative to dogs desexed before 6 months of age, dogs
desexed 6-12 months are 13% less likely to develop OA, while dogs who are desexed 12-24 or
24-60 months are 28% and 32% less likely to develop OA, respectively. This categorical model
suggests a potential non-linear relationship between age at desexing and OA risk, such that the
effects of age at desexing diminish as dogs get older. To test this, a sensitivity analysis was
performed adding a quadratic term in the continuous age at desexing model. The quadratic term
for age at desexing was statistically significant (Std. HR=1.35 (1.22-1.49) **Table 7**)). Predictions
from all models are presented in **Figure 8**. Results from this quadratic model more closely relate
to the categorical model, providing additional evidence that the magnitude of the OA-related risk
associated with age at desexing diminishes with age.
499
500   **Table 7**. Stratified Cox proportional hazard model testing the role of age at desexing as a risk
501   factor for OA. Age at desexing is treated as a continuous measure and as a categorical variable.

Desexing variable	Variable	HR (95% CI)	Std. HR (95% CI)	p-value	Concordance	N obs
Continuous	Age at baseline	1.42 (1.32-1.53)	1.42 (1.32-1.53)	<0.001***	0.70	24,159
	Max weight	1.02 (1.02-1.02)	1.83 (1.79-1.87)	<0.001***		
	Max BCS	1.05 (1.02-1.07)	1.05 (1.03-1.08)	<0.001***		
	Male	0.88 (0.84-0.93)	0.88 (0.84-0.93)	<0.001***		
	Age at desexing	0.87 (0.82-0.92)	0.90 (0.86-0.94)	<0.001***		
Quadratic	Age at baseline	1.42 (1.32-1.52)	1.42 (1.32-1.52)	<0.001***	0.70	24,159
	Max weight	1.02 (1.02-1.02)	1.82 (1.79-1.86)	<0.001***		
	Max BCS	1.05 (1.02-1.07)	1.06 (1.03-1.08)	<0.001***		
	Male	0.89 (0.84-0.93)	0.89 (0.84-0.93)	<0.001***		
	Age at desexing	0.61 (0.53-0.69)	0.67 (0.61-0.75)	<0.001***		
	Age at desexing^2	1.09 (1.06-1.13)	1.35 (1.22-1.49)	<0.001***		
Grouped	Age at baseline	1.41 (1.31-1.51)	4.39 (3.26-5.90)	<0.001***	0.70	24,159
	Max weight	1.02 (1.02-1.02)	1.82 (1.78-1.86)	<0.001***		
	Max BCS	1.05 (1.02-1.07)	1.06 (1.03-1.08)	<0.001***		
	Male	0.89 (0.85-0.93)	0.89 (0.85-0.93)	<0.001***		
	Desexed <6 mos	*Reference*				
	Desexed 6-12 mos	0.87 (0.82-0.92)	0.87 (0.82-0.92)	<0.001***		
	Desexed 12-24 mos	0.72 (0.64-0.80)	0.72 (0.64-0.80)	<0.001***		
	Desexed 24-60 mos	0.68 (0.58-0.79)	0.68 (0.58-0.79)	<0.001***		

502   *Std. HR= Standardized continuous variables (mean centered, scaled by standard deviation).
503   Wellness plan treated as stratifying variable.
504

Figure 8. Predicted marginal OA-free survival curves illustrating the effect of age at desexing on OA incidence, treating age as a linear continuous variable, a categorical variable, or quadratic variable. Color indicates ages at desexing. Median survival times and 95% CIs are reported. To generate predictions, median values were used for continuous covariates, proportions for binary variables, and the presence of wellness plan were used.

Discussion

The proportion of dogs in this population that developed OA was 24.9%. It is difficult to compare this to occurrence data from other studies due to differences in study populations, case definitions, and other methodological factors. Previous analyses of primary practice databases in the UK, for example, reported a one-year period prevalence of OA in dogs of any age of 2.34%[1] and 2.5%.[4] An owner survey, also in the UK, reported an overall prevalence of 2.23%.[38]

In contrast, a proprietary industry survey of 200 veterinarians in the U.S. reported an overall prevalence of 20%[5] in dogs over 1 year of age, a figure frequently cited in reviews and other articles discussing the condition. A more recent study involving dogs 5-12 years of age and

over 11 kg in weight presenting for dental prophylaxis found 68% of dogs had radiographic evidence of OA in one or more joints.(39) However, owner surveys from a subset of the population reported no clinical signs in 71% of the dogs, and these would likely not have received a diagnosis of OA in routine clinical practice. Such variability creates significant uncertainty about the true rate of occurrence of OA in dogs.

In this study, the incidence of 24.9% represents the clinical diagnosis of OA at any age and by any criteria in dogs from a primary care population who died in 2019, with no fixed interval or duration of follow up. Diagnoses could be based on clinical impression, history and physical examination, radiographic interpretations, or other diagnostic procedures. Given the large size and broad geographic distribution of the study population, this should be reasonably representative of the occurrence of OA as routinely identified by general practice clinicians in companion dogs in the U.S.

Detection bias in retrospective cohort studies can influence both the apparent prevalence of a condition and the apparent impact of risk factors for the disease.(40) Owners of dogs in this population were offered various wellness plans, which consist of a prepaid packages of services that can reduce the cost of clinic visits and some diagnostic tests, including radiographs. Enrollment in such a wellness plan has previously been suggested to be a risk factor for detection bias,(37) and in this study it was strongly associated with OA diagnosis. Dogs with a wellness plan had far more clinic visits and were more likely to have radiographs than dogs not enrolled in such a plan (data not shown), likely leading to a higher rate of detection of OA. This factor was controlled for by stratification in the analysis of other putative risk factors.

Age was by far the strongest risk factor for OA diagnosis, consistent with the established understanding of OA as a disease of aging. This effect was consistent across breeds of different sizes and with different rates of OA occurrence (**Figure 2** and **Supplemental Table 2**). Diagnosis of OA in individual patients is more likely as time passes, both because it is an incurable condition and so will be present and available for diagnosis longer, and because it is progressive and so more likely to become clinically apparent as a dog ages.

In previous studies, sex has been inconsistently identified as a risk factor for OA.(6) In this study, males were at lower risk, though the size of this effect in the overall population was small. Within breeds, males were at lower risk in some breeds but not others, independent of body weight and BCS (**Figure 2** and **Supplemental Table 2**).

All types of OA were included in the case definition for this study, regardless of which joint was affected or whether a predisposing condition, such as joint dysplasia or cranial cruciate ligament rupture, was identified. Previous evaluations of sex as a risk factor for OA have focused more narrowly on particular joints, predisposing conditions, or breeds, and it is likely that the role of sex as a risk factor may vary among these.(4) The small overall risk difference between males and females suggests that diagnostic and preventative measures should be directed equally at dogs of both sexes.

Previous studies have shown that OA risk differs markedly between breeds.(6) Breed is a complex risk factor involving differences in genetic makeup, body size and conformation, and

likely also lifestyle variables influenced by owners, such as feeding practices and the type and intensity of activity. Because of the enormous phenotypic variability among dogs, evaluation of the role of body weight in OA risk can easily confound the effects of body size and breed with those of overweight.

In general, larger body size is often identified as a risk factor for OA in dogs. This pattern was confirmed in this study, with increased mature body weight positively associated with OA risk. When OA risk was compared among breeds, larger breeds had higher risk, consistent with previous findings **(Supplemental Figures 2** and **3** and **Supplemental Table 2**).

In an effort to disentangle the effects of body size and overweight to some extent, we examined the interaction between the percentage increase above mature body weight and the risk for OA. This analysis showed that weight gain after maturity, likely representing excess adipose mass, increases risk of OA. This effect is greater in larger dogs, possibly due to an exacerbation of the already elevated risk posed by larger body size, or because the absolute mechanical burden of weight gained after maturity is greater in larger dogs.

Ideally, body condition score should help to isolate overweight and obesity as a risk factor independent of mature body size, so it was expected that BCS would be a strong predictor of OA risk. In this study population, 40.1% of dogs were classed as overweight (BCS 6-7/9) and 4.1% as obese (BCS 8-9/9), and the proportion of overweight and obesity was higher in the dogs that developed OA (overweight= 50.8%, obese= 5.3%) than in those that did not (overweight= 36.7%, obese= 3.8%). These figures are broadly consistent with previous reports, though the estimated prevalence of overweight and obesity vary widely depending on the population, time period, assessment methods, and category definitions.(21,41–43)

While BCS was significantly correlated with OA risk, the size of this effect was very small. This relationship was stronger in smaller dogs, unlike the effect of weight gain after maturity. Overall, BCS was not a powerful predictor of OA diagnosis. A possible explanation for this unexpected result may be practical challenges in the implementation of BCS scoring. There are several scoring systems in use by veterinarians, and the validation data for these are limited, involving relatively small numbers of dogs, breeds, and clinicians.(44–47) Some studies have reported that BCS scoring correlates well with objective measures of body fat.(46) However, the strength of this correlation varies significantly between breeds. The consistency and reliability of BCS scoring among veterinarians scoring dogs of many different breeds and conformations in general practice is uncertain.

Various assessment tools for body condition were used at different times in evaluation of this population, and scores from a 3-point and a 5-point scale were mapped onto a 9-point scale for analysis. In principle, a system with more gradations should allow for a more accurate distinction between degrees of overweight among dogs and help to identify more clearly the relationship between BCS and clinical conditions such as OA. In practice, however, the distribution of BCS scores in this data set was clearly discontinuous, with the vast majority of dogs scored as 5/9 (45.4%) or 7/9 (35%). This suggests clinicians tended to lump dogs into broad "normal" and "overweight" categories, rather than utilizing BCS scoring as a more finely graded, continuous measure.

Though this was true across all sizes of dogs, there was greater use of the full range of
scores, particularly those below 5/9, in smaller dogs (**Supplemental Figure 4**). This suggests
clinicians may find it easier to distinguish gradations of body condition in smaller breeds. The
fact that BCS was a stronger predictor of OA in smaller dogs, among which BCS scores were
more granular, supports the possibility that the lumping of many dogs into either "normal" or
"overweight" obscured the overall relationship between BCS and OA.

Further evaluation of the reliability of extant BCS scoring systems in heterogenous
general practice populations, and the development and validation of alternative assessment tools
that obviate the limitations of BCS, would be useful in clarifying the role of overweight and
obesity as risk factors for disease.

Neuter status has been consistently identified as a risk for OA, with intact individuals at
lower risk than neutered dogs. There is also evidence that neutering before 12 months of age is
associated with an increased risk of several of the most important conditions predisposing to OA
(hip and elbow dysplasia and cranial cruciate ligament rupture), though this effect may only
appear in larger dogs (over approximately 43 lbs in one recent analysis).(18)

This study is consistent with previous findings that the risk of OA is increased following
desexing in most medium and large breed dogs (over 30 lbs) and in some smaller breeds. While
predisposing conditions were not evaluated, the impact of neutering on OA risk overall increased
with body size, and neutering was less frequently associated with OA risk in smaller breeds,
supporting the apparent greater significance of neutering as a risk factor in larger dogs.

Neutering is also an established risk factor for obesity,(21,22,37) and it is possible that
one way in which neutering increases OA risk is by increasing the propensity for overweight and
obesity. Because of challenges in the interpretation of BCS scores, it was difficult to compare the
occurrence of overweight and obesity between intact and neutered dogs. The effect of body
weight gain after maturity on OA risk was not different between intact and neutered dogs, so
overweight and obesity are clearly important risk factors in all dogs regardless of neuter status.
While increasing the incidence of overweight and obesity may be one mediator of the impact of
neutering on OA risk, this hypothesis could not be directly evaluated in this study.

In this study, age at neutering was inversely associated with OA risk in the total
population, indicating that risk decreased progressively with delayed neutering up to 2 years of
age. This effect was the same in dogs of different body sizes, which is not consistent with
previous reports suggesting earlier neutering (before 12 months of age) increased the risk of
predisposing conditions, such as hip and elbow dysplasia and cruciate ligament rupture, in larger
dogs more than smaller dogs.(18) In this large a cohort, the power to detect such a difference in
risk associated with size should be high, so these data suggest the OA risk associated with age at
neutering would apply to all dogs in similar, heterogenous primary care populations. Differences
between this study and others in the appearance of a size effect for this risk factor may be
explained by differences in detection strategies, case definitions, or study populations.

This study serves to confirm the importance of several key risk factors for OA in dogs, including age, body weight, and neutering. Chronological age itself is not, of course, modifiable, but the potential to delay the onset of age-associated health conditions, including OA, has been demonstrated in dogs and numerous other species.(27,28) The most successful strategy for achieving this in research animals, severe caloric restriction, is not practical in companion dogs. Such a strategy requires precisely formulated diets and extensive monitoring to prevent malnutrition, and it can actually reduce lifespan in some animals due to nutrition/gene interactions.(48,49) Feeding is also a critical element of the human-animal bond, and asking dog owners to drastically underfeed their canine companions is unlikely to be accepted by many. However, caloric restriction research has done much to elucidate the general mechanisms of aging, raising the possibility that pragmatic therapies to delay aging-associated diseases may be developed.

Body weight is a function of both breed and body condition. Breed risks for specific health conditions, such as OA, can potentially be modified in populations by selective breeding and in individuals by nutritional interventions and, theoretically, the use of drugs targeting the physiologic characteristics of specific breeds that increase disease risk.

Body condition can clearly be altered by nutrition and exercise. Despite the unexpectedly small impact of BCS score on OA risk in this study, OA incidence was greater in overweight and obese dogs than in dogs scored as normal weight. Weight gain after maturity, which likely reflects accumulation of excess adipose, was also a significant risk factor for OA. These findings support the role of obesity as a risk factor for OA and the importance of maintaining optimal body condition in dogs through diet and exercise to minimize OA risk. These interventions are particularly important in larger dogs, who are at greatest risk of obesity and most significantly impacted by weight gain and other risk factors, such as neutering.

Neutering is an important modifiable risk factor for OA. Overall, gonadectomy appears to increase the risk of OA, particularly in larger dogs. Earlier neutering adds to this effect in both large and small dogs. The pathogenesis of OA, of course, is complex and incompletely elucidated, and the role of neutering should be understood in this context, as one of many risk interacting risk factors. Given this complexity, it is not likely that the decision whether or not to neuter an individual dog will be the primary determinant of whether or not that individual develops OA. However, recognizing that an association exists on a population level between neutering and OA risk can help clinicians identify individuals who may be at elevated risk and utilize more proactive surveillance to ensure early diagnosis and appropriate preventative and therapeutic interventions.

Any potential increase in OA risk associated with neutering must also be considered in the context of other risks and benefits associated with neutering on both an individual and population level. Changes in neutering practices intended to mitigate one health risk may entail unpredictable increases in other risks not yet identified, so judicious integration of such information with the larger context and the circumstances and needs of individual patients is critical.

704   Future research in such large, primary-care cohorts can further clarify the relationship
705   between putative risk factors and common aging-associated diseases such as OA. This will
706   support efforts to reduce the burden of these conditions on companion dogs and their caregivers
707   by informing appropriate preventative intervention.
708

Limitations

As a retrospective cohort study, this analysis relied on electronic medical records,
collected across thousands of clinics and veterinarians. While these data provide longitudinal
insights into risk factors associated with incident diseases, the data themselves were not collected
to test hypotheses surrounding risk factors for incident osteoarthritis. The study population (all
dogs seen between 1998-2019 and dying in 2019) was large and expected to be representative of
the patient population at Banfield during this period, but it may not be representative of the
overall U.S. companion dog population.
717

Electronic health records rely on real-time recording of interactions between clinicians,
clients, and patients. Information provided by clients, such as age, breed, and prior medical
history, may be subject to misclassification or recall bias. Information provided by veterinarians,
such as diagnosis of OA and scoring of BCS, may be subject to misclassification bias and may
vary over time and between clinicians. For example, a diagnosis of OA could be based on
clinical impression, history and physical examination, radiographic interpretations, or other
factors, and no standardized diagnostic criteria were in place.
725

Overall incidence rates may differ from true population-level incidence, and should be
interpreted with caution. Additionally, percentiles were used to summarize adult body weight
and BCS across the follow-up window, which fails to capture duration of time over or
underweight. Finally, age at desexing was only known for dogs being desexed at a Banfield Pet
Hospital clinic, which does not account for desexing occurring outside of Banfield clinics.

Conflict of Interest

Author Contributions
JG conducted the data analyses and wrote the Materials and Methods and the Results sections.
BM wrote the Introduction and Discussion sections. All authors contributed to manuscript
revisions and have read and approved the final manuscript.

Acknowledgements
The authors would like to thank Alexander Naka for significant contributions to the initial
conception and design of this project, Zane Koch for assistance with data processing and
analysis, Banfield Pet Hospital for sharing the data set analyzed in this project, and the

veterinarians and clients of Banfield Pet Hospital for generating this invaluable information
resource.

Contribution to the Field
Osteoarthritis, also called degenerative joint disease, is the most common joint disorder in dogs.
It is an incurable, progressive condition that can cause significant pain and disability. Identifying
risk factors for the development of osteoarthritis can help veterinarians recognize individual dogs
at high risk and make appropriate efforts to diagnose and treat the condition as early as possible.
Identifying those risk factors that can be changed can help dog owners and veterinarians prevent
or mitigate the development of this condition in some dogs. This study analyzed a very large
body of data about companion dogs examined by veterinarians in many general practice
veterinary clinics throughout the United States. This analysis confirmed the importance of
several previously suspected risk factors for the development of osteoarthritis, including age and
body weight. The study also added to existing information about the relationship between the
risk of developing osteoarthritis and several factors, including body weight, desexing, and age at
desexing. This research will be useful both to clinicians treating canine patients and to other
researchers studying the factors that lead to the development of osteoarthritis in dogs.

References

1. O'Neill DG, James H, Brodbelt DC, Church DB, Pegram C. Prevalence of commonly diagnosed
disorders in UK dogs under primary veterinary care: results and applications. BMC Vet Res.
2021 Dec;17(1):1–14.

2. Robinson NJ, Dean RS, Cobb M, Brennan ML. Investigating common clinical presentations in
first opinion small animal consultations using direct observation. Vet Rec. 2015
May;176(18):463–463.
3. Wolf S, Selinger J, Ward MP, Santos-Smith P, Awad M, Fawcett A. Incidence of presenting
complaints and diagnoses in insured Australian dogs. Aust Vet J. 2020 Jul;98(7):326–32.
4. Anderson KL, O'Neill DG, Brodbelt DC, Church DB, Meeson RL, Sargan D, et al. Prevalence,
duration and risk factors for appendicular osteoarthritis in a UK dog population under primary
veterinary care. Sci Rep 2018 81. 2018 Apr;8(1):1–12.
5. Johnston SA. Osteoarthritis. Joint anatomy, physiology, and pathobiology. Vet Clin North Am
Small Anim Pract. 1997;27(4):699–723.
6. Anderson KL, Zulch H, O'Neill DG, Meeson RL, Collins LM. Risk Factors for Canine
Osteoarthritis and Its Predisposing Arthropathies: A Systematic Review. Front Vet Sci.
2020;7:220–220.
7. Belshaw Z, Dean R, Asher L. "You can be blind because of loving them so much": The impact
on owners in the United Kingdom of living with a dog with osteoarthritis. BMC Vet Res. 2020
Jun;16(1):1–10.
8. Pegram C, Gray C, Packer RMA, Richards Y, Church DB, Brodbelt DC, et al. Proportion and
risk factors for death by euthanasia in dogs in the UK. Sci Rep. 2021 Dec;11(1).

9. O'Neill DG, Church DB, McGreevy PD, Thomson PC, Brodbelt DC. Longevity and mortality of owned dogs in England. Vet J Lond Engl 1997. 2013 Dec;198(3):638–43.

10. Urfer SR, Kaeberlein M, Promislow DEL, Creevy KE. Lifespan of companion dogs seen in three independent primary care veterinary clinics in the United States. Canine Med Genet. 2020;7:7.

11. Bonnett BN, Egenvall A. Age patterns of disease and death in insured Swedish dogs, cats and horses. J Comp Pathol. 2010 Jan;142 Suppl 1:S33-38.

12. Clements DN, Carter SD, Innes JF, Ollier WER. Genetic basis of secondary osteoarthritis in dogs with joint dysplasia. Am J Vet Res. 2006 May;67(5):909–18.

13. Sanderson S. The epidemic of canine obesity and its role in osteoarthritis. Isr J Vet Med. 2012;67:195–202.

14. Shahid M, Manchi G, Slunsky P, Naseer O, Fatima A, Leo B, et al. A systemic review of existing serological possibilities to diagnose canine osteoarthritis with a particular focus on extracellular matrix proteoglycans and protein. Pol J Vet Sci. 2017 Mar;20(1):189–201.

15. Smith GK, Mayhew PD, Kapatkin AS, McKelvie PJ, Shofer FS, Gregor TP. Evaluation of risk factors for degenerative joint disease associated with hip dysplasia in German Shepherd Dogs, Golden Retrievers, Labrador Retrievers, and Rottweilers. J Am Vet Med Assoc. 2001 Dec;219(12):1719–24.

16. Torres de la Riva G, Hart BL, Farver TB, Oberbauer AM, Messam LLM, Willits N, et al. Neutering dogs: effects on joint disorders and cancers in golden retrievers. PloS One. 2013;8(2):e55937.

17. Hart BL, Hart LA, Thigpen AP, Willits NH. Long-term health effects of neutering dogs: comparison of Labrador Retrievers with Golden Retrievers. PLoS One. 2014 Jul;9(7):e102241–e102241.

18. Hart BL, Hart LA, Thigpen AP, Willits NH. Assisting Decision-Making on Age of Neutering for 35 Breeds of Dogs: Associated Joint Disorders, Cancers, and Urinary Incontinence. Front Vet Sci. 2020 Jul;7:388–388.

19. McKenzie B. Evaluating the benefits and risks of neutering dogs and cats. CABI Rev. 2010 Jan;2010:1–18.

20. Lefebvre SL, Yang M, Wang M, Elliott DA, Buff PR, Lund EM. Effect of age at gonadectomy on the probability of dogs becoming overweight. J Am Vet Med Assoc. 2013 Jul;243(2):236–43.

21. Lund EM, Armstrong PJ, Kirk CA, Klausner JS. Prevalence and risk factors for obesity in adult dogs from private US veterinary practices. Int J Appl Res Vet Med. 4(2):177–86.

22. Bjørnvad CR, Gloor S, Johansen SS, Sandøe P, Lund TB. Neutering increases the risk of obesity in male dogs but not in bitches — A cross-sectional study of dog- and owner-related risk factors for obesity in Danish companion dogs. Prev Vet Med. 2019 Oct;170:104730–104730.

23. Spain CV, Scarlett JM, Houpt KA. Long-term risks and benefits of early-age gonadectomy in dogs. J Am Vet Med Assoc. 2004 Feb 1;224(3):380–7.

24. Kutzler MA. Possible Relationship between Long-Term Adverse Health Effects of Gonad-Removing Surgical Sterilization and Luteinizing Hormone in Dogs. Anim Open Access J MDPI [Internet]. 2020 Apr;10(4). Available from: /pmc/articles/PMC7222805/

25. Grandi FC, Bhutani N. Epigenetic Therapies for Osteoarthritis. Trends Pharmacol Sci. 2020 Aug;41(8):557–69.

26. Shen J, Abu-Amer Y, O'Keefe RJ, McAlinden A. Inflammation and epigenetic regulation in osteoarthritis. https://doi.org/101080/0300820720161208655. 2016 Jan;58(1):49–63.

27. Kealy RD, Lawler DF, Ballam JM, Mantz SL, Biery DN, Greeley EH, et al. Effects of diet restriction on life span and age-related changes in dogs. J Am Vet Med Assoc. 2002 May 1;220(9):1315–20.

28. Kealy RD, Lawler DF, Ballam JM, Lust G, Smith GK, Biery DN, et al. Five-year longitudinal study on limited food consumption and development of osteoarthritis in coxofemoral joints of dogs. J Am Vet Med Assoc. 1997 Jan 15;210(2):222–5.

29. Huck JL, Biery DN, Lawler DF, Gregor TP, Runge JJ, Evans RH, et al. A Longitudinal Study of the Influence of Lifetime Food Restriction on Development of Osteoarthritis in the Canine Elbow. Vet Surg. 2009 Feb;38(2):192–8.

30. Runge JJ, Biery DN, Lawler DF, Gregor TP, Evans RH, Kealy RD, et al. The effects of lifetime food restriction on the development of osteoarthritis in the canine shoulder. Vet Surg VS. 2008 Jan;37(1):102–7.

31. Hazewinkel H, Goedegebuure SA, Poulos P, Wolvekamp W. Influences of chronic calcium excess on the skeletal development of growing Great Danes. J Am Anim Hosp Assoc. 1985;

32. Hedhammar A, Wu F, Krook L, Schryver H, Lahunta A de, Whalen J, et al. Overnutrition and skeletal disease. An experimental study in growing Great Dane dogs. Cornell Vet. 1974;

33. Manfredi S, Di Ianni F, Di Girolamo N, Canello S, Gnudi G, Guidetti G, et al. Effect of a commercially available fish-based dog food enriched with nutraceuticals on hip and elbow dysplasia in growing Labrador retrievers.

34. Cox DR. Regression Models and Life-Tables. J R Stat Soc Ser B Methodol. 1972;34(2):187–220.

35. Therneau TM, Grambsch PM. Modeling Survival Data: Extending the Cox Model [Internet]. New York, NY: Springer; 2000 [cited 2023 Jul 18]. (Dietz K, Gail M, Krickeberg K, Samet J, Tsiatis A, editors. Statistics for Biology and Health). Available from: http://link.springer.com/10.1007/978-1-4757-3294-8

36. Therneau TM, Grambsch PM. A Package for Survival Analysis in S, version 2.38. Model Surviv Data Extending Cox Model [Internet]. 2000; Available from: http://cran.r-project.org/package=survival

37. Benka VA, Scarlett JM, Sahrmann J, Rieke K, Briggs JR, Ruple A, et al. Age at gonadectomy, sex, and breed size affect risk of canine overweight and obese outcomes: a retrospective cohort study using data from United States primary care veterinary clinics. J Am Vet Med Assoc. 2023 May 19;1–10.

38. Wiles BM, Llewellyn-Zaidi AM, Evans KM, O'Neill DG, Lewis TW. Large-scale survey to estimate the prevalence of disorders for 192 Kennel Club registered breeds. Canine Genet Epidemiol. 2017;4:8.

39. Millis D, Tichenor MG, Hecht S, Hunt T. Prevalence of Osteoarthritis in Dogs Undergoing Routine Dental Prophylaxis. In: World Small Animal Veterinary Association World Congress Proceedings. Cape Town, South Africa; 2014.

40. Combelles L, Corbiere F, Calavas D, Bronner A, Hénaux V, Vergne T. Impact of imperfect disease detection on the identification of risk factors in veterinary epidemiology. Front Vet Sci. 2019;6(MAR):66–66.

41. Association for the Prevention of Pet Obesity. State of U.S. Pet Obesity 2022. 222AD.

42. Chiang CF, Villaverde C, Chang WC, Fascetti AJ, Larsen JA. Prevalence, Risk Factors, and Disease Associations of Overweight and Obesity in Dogs that Visited the Veterinary Medical Teaching Hospital at the University of California, Davis from January 2006 to December 2015. Top Companion Anim Med. 2022 May;48:100640–100640.

43. McGreevy PD, Thomson PC, Pride C, Fawcett A, Grassi T, Jones B. Prevalence of obesity in dogs examined by Australian veterinary practices and the risk factors involved. Vet Rec. 2005 May;156(22):695–702.

44. Laflamme D. Development and Validation of a Body Condition Score System for Dogs. Canine Pract. 1997;22(1):10–5.

45. Burkholder WJ. Use of body condition scores in clinical assessment of the provision of optimal nutrition. J Am Vet Med Assoc. 2000;217(5):650–4.

46. Mawby DI, Bartges JW, d'Avignon A, Laflamme DP, Moyers TD, Cottrell T. Comparison of Various Methods for Estimating Body Fat in Dogs. J Am Anim Hosp Assoc. 2004 Mar;40(2):109–14.

47. German AJ, Holden SL, Moxham GL, Holmes KL, Hackett RM, Rawlings JM. A Simple, Reliable Tool for Owners to Assess the Body Condition of Their Dog or Cat. J Nutr. 2006 Jul;136(7):2031S-2033S.

48. Lee MB, Hill CM, Bitto A, Kaeberlein M. Antiaging diets: Separating fact from fiction. Science. 2021 Nov 19;374(6570):eabe7365.

49. Nikolai S, Pallauf K, Huebbe P, Rimbach G. Energy restriction and potential energy restriction mimetics. Nutr Res Rev. 2015 Dec;28(2):100–20.