1

A special role for Anterior Cingulate Cortex, but not Orbitofrontal Cortex or Basolateral Amygdala, in choices involving information

Valeria V. González[1], Sonya A. Ashikyan[1], Yifan Zhang[5], Anne Rickard[1], Ibrahim Yassine[1], Juan Luis Romero-Sosa[1], Aaron P. Blaisdell[1-3], & Alicia Izquierdo[1-4]

[1]Department of Psychology, University of California-Los Angeles, Los Angeles, California, USA.

[2]The Brain Research Institute, University of California-Los Angeles, Los Angeles, California, USA.

[3]Integrative Center for Learning and Memory, University of California-Los Angeles, Los Angeles, California, USA.

[4]Integrative Center for Addictions, University of California-Los Angeles, Los Angeles, California, USA.

[5]Department of Computer Science, University of Southern California, Los Angeles, California, USA.

This work was supported by R01 DA047870 (Izquierdo), BCS-1844144 (Blaisdell) and UC's Chancellor Postdoctoral Fellowship (González). The authors have no competing interests to declare that are relevant to the content of this article.

Correspondence concerning this article should be addressed to Dr. Valeria V. González, UCLA Department of Psychology, 1285 Franz Hall, Los Angeles, CA 90095-1563. E-mail: vgonzalez@psych.ucla.edu

Abstract

Humans and other animals make decisions under uncertainty. Choosing an option that provides
information can improve decision making. However, subjects often choose information that does
not increase the chances of obtaining reward. In a procedure that promotes such paradoxical
choice, animals choose between two alternatives: The richer option is followed by a cue that is
rewarded 50% of the time (No-info) and the leaner option is followed by one of two cues, one
always rewarded (100%), and the other never rewarded, 0% (Info). Since decisions involve
comparing the subjective value of options after integrating all their features perhaps including
information value, preference for information may rely on cortico-amygdalar circuitry. To test
this, male and female Long-Evans rats were prepared with bilateral inhibitory DREADDs in the
anterior cingulate cortex (ACC), orbitofrontal cortex (OFC), basolateral amygdala (BLA), or null
virus infusions as a control. Using a counterbalanced design, we inhibited these regions after
stable preference was acquired and during learning of new Info and No-info cues. We found that
inhibition of ACC, but not OFC or BLA, selectively destabilized choice preference in female rats
without affecting latency to choose or the response rate to cues. A logistic regression fit revealed
that the previous choice strongly predicted preference in control animals, but not in female rats
following ACC inhibition. BLA inhibition tended to decrease the learning of new cues that signaled
the Info option, but had no effect on preference. The results reveal a causal, sex-dependent role
for ACC in decisions involving information.

Keywords: chemogenetics, frontal cortex, DREADDs, information, choice, paradoxical, curiosity,

explore-exploit, information-seeking, value-based decision making

The environment is full of unpredictable events, and information that reduces

uncertainty about such events allows an organism to better predict and prepare for the future.

However, several psychiatric conditions are characterized by a strong preference for

information, or an Intolerance of Uncertainty, including Autism Spectrum Disorder, Substance

Use Disorders, Attention-Deficit Hyperactivity Disorder, Generalized Anxiety Disorder, and

Obsessive-Compulsive Disorder (Boulter et al., 2014; Dugas et al., 2004; Jenkinson et al., 2020;

Mandali et al., 2021; Tolin et al., 2003). Intolerance of uncertainty has been linked to

'pathological doubt' during which the preference for information is dramatically increased

(Tolin et al., 2003).

Obtaining information can be crucial for survival. However, if information cannot be

used to modify action, a bias toward information can be considered paradoxical or even

suboptimal because organisms should not invest resources to obtain information that does not

affect the outcome of a choice. Imagine a situation in which an organism chooses between two

sources of delayed reward; if rewards are signaled before each choice, it will be worth investing

in that information to choose the best option. In contrast, if the outcomes are signaled *after* the

choice, information is useless, since the organism cannot change its choice. The latter has been

broadly studied at the behavioral level. In the so-called paradoxical or suboptimal choice task

(Stagner & Zentall, 2010), animals are presented with two alternatives, one providing a lower

rate of reinforcement with different stimuli indicating the presence (S+) or absence (S-) of

delayed food (i.e. Info), and another one (S3) providing a higher rate of reinforcement but with

non-differential stimuli signaling food (i.e. No-Info) (See **Figure 1A**). Animals often prefer the

leaner but informative option despite the difference in reinforcement rates between

alternatives (Fortes et al., 2016; Macias et al., 2021; Stagner & Zentall, 2010) .

The neural substrates of reinforcement uncertainty have been broadly researched, with

evidence pointing to a distributed network that involves prefrontal cortex (PFC) (Rushworth &

Behrens, 2008), striatum, hippocampus, basolateral amygdala (BLA) and mediodorsal thalamus

(Soltani & Izquierdo, 2019; Winstanley & Floresco, 2016). Nevertheless, to our knowledge there

is no investigation of the specific brain regions using this behaviorally well-documented

paradoxical choice procedure. Additionally, compared to primates, there is a paucity of rodent

studies on the value of non-instrumental information and its neural substrates (Iigaya et al.,

2019; Iigaya et al., 2016). When human subjects are assessed on choices between two cued

alternatives- an informative vs. a non-informative one (but with no difference in overall

reinforcement rate), subjects reliably prefer advanced information and Blood-oxygen-level-

dependent (BOLD) signal in ventromedial PFC tracks the value of the anticipation of reward

(Iigaya et al., 2019; Iigaya et al., 2016). Neural correlates of uncertainty have been found in

different subregions of the PFC in several species, among them, orbitofrontal cortex (OFC) and

anterior cingulate cortex (ACC) (Rushworth & Behrens, 2008). Electrophysiological recording

studies in rats and monkeys demonstrate that activity in OFC is associated with stimulus (cue)

value and expected uncertainty, or probabilistic risk (Jenni et al., 2023; Jo & Jung, 2016;

Namboodiri et al., 2019; Riceberg & Shapiro, 2017). Similarly, studies have shown that ACC

neurons signal value, uncertainty of predictions about rewards or punishments (Jezzini et al.,

2021; Monosov, 2017), and track trial-by-trial outcomes of choice (Procyk et al., 2000; Shidara

& Richmond, 2002). On the other hand, inactivation of BLA decreases the likelihood of choosing

uncertain options in probabilistic-discounting tasks (Ghods-Sharifi et al., 2009; St Onge et al.,

2012; Stopper & Floresco, 2011). This suggest the contribution of BLA may be biasing choice

towards larger rewards, especially when the delivery of these rewards is uncertain.

Furthermore, selectively disrupting PFC-to-BLA connections increases choice of the larger yet

increasingly uncertain reward, implying that communication between these two regions serves

to modify choice biases (St Onge et al., 2012). Finally, lesions to either OFC or BLA (or their

connection), results in slower learning about which option has the better payout, suggesting

that these areas form a functional circuit for the adaptation of reward-maximizing strategies

(Zeeb & Winstanley, 2011, 2013).

In this experiment, we examined the specific contributions of ACC, OFC, and BLA to this

seemingly 'suboptimal' choice phenomenon via chemogenetic manipulation. On the additional

evidence that these regions participate in decision confidence under uncertainty (Lak et al.,

2014; Stolyarova, Rakhshan, Hart, O'Dell, et al., 2019), we hypothesize that they may play vital,

yet dissociable roles in decisions involving information value. To study this, we inactivated

these regions during stable preference and during the learning of new informative and non-

informative cues. We found that inhibition of ACC, but not OFC or BLA, destabilized choices

involving information in females and reduced their ability to use previous choices to guide

current decisions.

Materials and Methods

Animals.

Sixty-six Long Evans rats (*Rattus Norvegicus*), 36 females and 30 males, acquired from Envigo

served as subjects. Subjects were between aged post-natal days (PND) 90 and 140 at the start

of the experiment. Subjects were pair-housed before and single-housed after surgeries in

transparent plastic tubs with wood shaving bedding in a vivarium maintained on a reverse 12-hr

light cycle. Experiments were conducted during the dark portion of the cycle at a minimum of

five days per week. A progressive food restriction schedule was imposed prior to the beginning

of the experiment to maintain rats at 85% of their initial free-feeding weights. Water was

always available in their home-cages. The procedures used in this experiment were conducted

under approval and following the guidelines established by the Chancellor's Animal Research

Committee at UCLA.

Figure 1. Task structure and experimental timeline. **(A)** Rats choose between two levers (left or right). After pressing once, both levers retract and an auditory cue commences. If the rat chooses the 'Info' alternative, on 20% of the trials, tone S+ plays for 60s always ending with the delivery of one sugar pellet; the other 80% of the trials sound S- plays for 60s always ending without food. If the rat chooses the 'No-Info' option, a third sound S3 plays for 60s ending with the delivery of one sugar pellet in 50% of the trials. **(B)** CNO or VEH was administered 10 min before starting the task. Each session consisted of 54 trials in which rats received 18 choice trials (both levers available to choose), 18 forced Info trials (only the lever associated with the Informative alternative was available) and 18 forced No-Info trials (only the No-Info lever was available). Animals were required to complete the session in 2 hrs. **(C)** Rats were trained on the task over 15 sessions before they underwent bilateral viral infusion of inhibitory hM4Di DREADDs or null virus enhanced Green Fluorescent Protein (eGFP) in ACC, BLA, or OFC. After two weeks of recovery, baseline performance was reestablished for 5 sessions before administrating CNO and VEH (order counterbalanced) for three consecutive sessions with a wash-out day between drugs. Then, the three originally trained auditory cues were replaced with three new auditory cues, while maintaining the rest of the structure as before. Rats received CNO or VEH for 15 to 20 sessions before receiving 5 final sessions of the task without drug.

Viral Constructs.

To express DREADDs on putative projection neurons in ACC, OFC, or BLA, an adeno-

associated virus AAV8 driving the hM4Di-mCherry sequence under the CaMKIIa promoter was

used (AAV8-CaMKIIa-hM4D(Gi)-mCherry, packaged by Addgene, viral prep #50477-AAV8). A

virus without the hM4Di DREADD gene but containing the fluorescent tag enhanced Green

Fluorescent Protein, eGFP (AAV8-CaMKIIa-EGFP, packaged by Addgene, viral prep #50469-

AAV8) was infused into ACC, OFC, or BLA as a null virus control. This null virus allowed us to

control for non-specific effects of surgical procedures (i.e., craniotomy, anesthesia), exposure to

AAV8, and non-specific effects of drug or injections.

Behavioral apparatus.

This experiment was conducted using operant testing chambers, measuring 30 x 25 x 20

cm (L x W x H). Each chamber was housed in separate sound- and light- attenuating

environmental isolation chests (ENV-008, Med Associates, Georgia, VT). The front and back

walls and ceiling of the chambers were constructed of clear Plexiglas, the side walls were made

of aluminum, and the floors were built of stainless-steel rods measuring 0.5 cm in diameter,

spaced 1.5 cm center-to-center.

Each chamber had a pellet-dispenser (ENV-203-45, Med Associates) and a cup type

pellet receptacle (ENV-200R1M, Med Associates). When activated, one sucrose pellet was

delivered into the cup. The opening of the cup was equipped with an infrared beam and

photodetector to record entries into the food niche. A 3.5 cm wide operant lever was

positioned one cm to the left of the food niche on the metal wall.

A speaker (ENV-224DM) on the ceiling of the chamber delivered a siren (cycling

between 1500 and 1900 Hz at a 0.5 s rate), a 1000-Hz tone and a white noise, all were 8 dB

above background to serve as the initial S+, S- and S3; and another siren (cycling between 4000

and 3500 Hz at a 0.5 s rate), a 3000-Hz tone and a click train (4/s) 8 dB above background

served as a second set of S+, S- and S3 cues, counterbalanced across subjects. A diffuse

incandescent light (ENV-227M, Med Associates) was located on the bottom panel of the right-

side chamber wall, 6 cm from the ceiling.

Surgical procedures.

After completing the training phase, rats were anesthetized with isoflurane for bilateral

infusion of ACC, OFC or BLA inhibitory (Gi) DREADDS (AAV8-CaMKIIα-hM4D(Gi)-mCherry,

Addgene, Cambridge, MA, viral prep #50477-AAV8) or eGFP (AAV8-CaMKIIa-EGFP, Addgene,

Cambridge, MA, viral prep #50469-AAV8). Craniotomies were created, and a 26-gauge guide

cannula (PlasticsOne, Roanoke, VA) with a dummy injector was lowered, after which the

dummy injector was replaced with a 33-gauge internal injector (PlasticsOne, Roanoke, VA) was

inserted. Animals were infused with two bilateral sites of injections in BLA (0.2 µL at 0.1 µL/min

in AP: -2.5, ML: ±5.0, DV: -8.6, and 0.1 µL at 0.1 µL/min in AP: -2.5, ML: ±5.0, DV: -8.3; total

volume per side = 0.3 µL), two bilateral sites in OFC (0.15 µL at 0.1 µL/min in AP: 4.0, ML: ±2.5,

DV: -4.4, and 0.2 µL at 0.1 µL/min in AP: 3.7, ML: ±2.5, DV: -4.6; total volume per side = 0.35 µL)

and one bilateral site in ACC (0.3 µL at 0.1 µL/min AP: +3.7, ML: ±0.8, DV: -2.4 for a total volume

of 0.3 µL per side). All measurements were taken from bregma. After infusion, the cannula was

left in place for ten additional minutes to allow diffusion.

Drug treatment.

Before testing, rats were given intraperitoneal (i.p.) injections of vehicle (VEH: 95%

saline + 5% DMSO) or clozapine-N-oxide, CNO (3 mg/kg CNO in 95% saline + 5% DMSO) 10 min

prior to beginning the behavioral task. The time was shorter than some other work (Hart et al.,

2020; Ye et al., 2023) to account for the longer duration of behavioral testing sessions as in a

previous experiment (Stolyarova, Rakhshan, Hart, O'Dell, et al., 2019). CNO and VEH were

administered during *Stable Info vs No Info* condition in a within-subject design (**Figure 1B**),

where animals received three sessions of CNO (or VEH) followed by a washout day with no

injection or training, and then three sessions of VEH (or CNO) followed by another washout day,

such that the order of drug administration was counterbalanced across animals. Following this,

CNO or VEH was administered during the a*cquisition of new cues* in a between-subject design

(drug condition counterbalanced) for 15-20 sessions.

Behavioral procedure.

Pretraining. Before training, ten sucrose pellets were given in the rat home cage to

avoid food neophobia. On the first day, rats were trained to eat pellets from the pellet tray by

delivering one pellet every 20 ± 15 s in the chamber (actual intertrial interval (ITI) values = 5, 10,

15, 20, 25, 30, and 35 s) for a total of 40 pellets. On days 2 and 3, rats were trained in an

autoshaping procedure to lever press the left and right lever (lever presented in alternate

order). Reinforcements were delivered following each lever press (i.e., a continuous

reinforcement schedule) for a maximum of 40 pellets within a session, or after 30 min had

elapsed.

Training of the Paradoxical choice task. The training stage of the task was comprised of

two types of trials: choice and forced trials (**Figure 1C**). In choice trials, rats were required to

choose between two levers available simultaneously. A choice trial started with the

simultaneous insertion of both levers and the houselight turning on. A choice was made by

pressing a lever one time. After the choice was made, if the animal chose the Info alternative,

both levers retracted and the houselight turned off. If the lever associated with the Info option

was pressed, 20% of the time an auditory cue (S+) was presented for 60 s always ending with

food; the other 80% of the time another auditory cue (S-) was on for 60 s never ending with

food. The total percentage of reinforced trials was 20%. If the rat chose the No-Info alternative,

a third auditory cue (S3) was presented for 60 s and ended with food on half of the trials. The

percentage of reinforcement on this alternative was 50%. On forced trials, only one lever was

presented at a time, following the same contingencies described above. All trials were

separated by a 10-s intertrial-interval (ITI) during which all stimuli were off. A single session

constituted 54 trials, 18 choice trials, and 36 forced trials. The maximum duration of a session

was set to 120 min. The assignment of sounds to S+, S- or S3 and the lever side for each option

was counterbalanced across animals but remained the same for individual animals throughout

training. This condition lasted 15 sessions.

Stable preference. Approximately two weeks after surgery, rats received five training

sessions as described above as a reminder of the task, used as a measure of baseline preference

(Baseline Info vs No Info, **Figure 1B**). After this, rats received three sessions of the task with an

i.p. injection of CNO and three sessions with an injection of VEH (order counterbalanced across

animals). After the three-session round of injections, animals underwent a washout day in

which they were not tested on the task. This condition lasted six sessions (eight days).

New cues preference. Three new tones were introduced to replace the auditory cues

used in previous conditions as S+, S- and S3. This change was used to evaluate the inhibition

effect o of BLA, OFC, or ACC in learning about new informative cues. Aside from the auditory

cues, all else remained identical. Animals received a minimum of 15 sessions of training and

stability criteria (no more than 20% change over the last three sessions) with a maximum of 20

sessions (acquisition of new cues, **Figure 1B**). Finally, animals received five sessions of the same

procedure without drug to establish baseline performance (baseline new cues, **Figure 1B**)

Histology.

At the conclusion of the experiment, rats were euthanized by an overdose of sodium

pentobarbital (Euthasol, 0.8 mL, i.p.; Virbac, Fort Worth, TX) and transcardially perfused with

phosphate buffered saline (PBS) followed by 10% buffered formalin acetate. Brains were

extracted and post-fixed in this solution for 24 hours followed by 30% sucrose cryoprotection.

Tissue was sectioned in 40-μM thick slices and cover slipped with DAPI mounting medium

(Prolong gold, Invitrogen, Carlsbad, CA) visualized using a BZ-X710 microscope (Keyence, Itasca,

IL), and analyzed with BZ-X Viewer software (**Figure 2A**).

We have previously validated the efficacy of our CNO-activated DREADDs ex-vivo in slice

(Aguirre et al., 2023; Hart et al., 2020) and behaviorally (Aguirre et al., 2023; Stolyarova,

Rakhshan, Hart, O'Dell, et al., 2019) in OFC, ACC, and BLA. A subset of 40 μm coronal sections

containing ACC were also stained for c-Fos, following an adapted Abcam protocol for dry-

mounted slides (Schneider Gasser et al., 2006). In this protocol, mounted tissue was marked

with a hydrophobic pen, any medium was added with a pippette and removed using a

vacuuum. The tissue was incubated for 22-24 h at 4 °C in a solution of primary antibody (1:2000

rabbit polyclonal to c-Fos, Abcam, Cambridge, MA) with 5% normal goat serum (Abcam,

Cambridge, MA), and 0.1% TritonX-100 (Sigma, St. Louis, MO) in PBS. After the incubation, the

tissue was washed with PBS 3 times during a 5-min period, then the brain slides incubated in a

secondary antibody for 90 min protected from light at room temperature (0.1% TritonX-100,

5% normal goat serum, PBS solution with 1:500 goat anti-rabbit Alexa 488; Abcam, Cambridge,

MA). Slides were washed again as described above. Then, tissue was incubated during 3 min

with quenching reagent (1:1 ratio) to reduce background. Slides were washed with PBS and

then cover-slipped with fluoroshield DAPI mounting medium (Abcam, Cambridge, MA).

Figure 2. Inhibitory DREADDs in ACC, BLA and OFC; Validation of ACC inhibition via c-fos immunohistochemistry. **(A)** Representative placement of inhibitory hM4Di DREADDs and eGFP null virus at Anterior-Posterior (AP) level +3.7 for ACC and OFC, and -2.8 for BLA relative to Bregma. Note, there were no eGFP for OFC. **(B)** Reconstructions of placement of inhibitory hM4Di DREADDs (pink) and eGFP null virus (*green*) at AP level +4.2, +3.7 and +3.2 for ACC, OFC, and AP -2.56, -2.8 and -3.14 for BLA relative to Bregma. **(C)** Representative image showing DAPI (blue), hM4Di-mCherry (red), c-fos immunoreactivity (green) and their overlap after injections of CNO in ACC at 10x and 20x. **(D)** Mean cell count of 3 images per condition, hM4Di+CNO, DAPI+CNO for ACC. **(E)** Mean spread of viral expression across brain regions, pixel quantification was done using ImageJ software. ns= nonsignificant, *p<0.05, Mann-Whitney U test.

c-Fos inmunoreactivity quantification images were visualized with a 20× objective with a

724 µm by 543 µm field of view using a confocal microscope (Model LSM 900, Zeiss, Germany).

For each region, three images were taken from two or three separate coronal sections from

both hemispheres at the same approximate AP coordinate (ACC +3.7 mm). To verify DREADD-

mediated inhibition of neurons in ACC after CNO administration we compared the number of c-

fos–positive cells in the hM4Di-expressing regions to the number of c-fos–positive cells in

neighboring (non-hM4Di-expressing, DAPI) areas. Cell counts were conducted using ImageJ

software (**Figure 2C**). We found greater overlap number of c-fos positive cells in hM4Di+ than

DAPI+ cell areas (Mann-Whitney U test: W= 0, p = 0.04) (**Figure 2D**). DREADDs and eGFP

expression were determined by matching histological sections to a standard rat brain atlas

(Paxinos and Watson, 2007) and quantifying fluorescence using ImageJ (Rueden et al., 2017)

where two independent raters measured the area of fluorescent pixels for each animal per

hemisphere (**Figure 2E**).

An independent sample t-test compared viral spread for ACC hM4Di (Mean$_{ACC\ hM4Di}$=

88542.87) against ACC eGFP (Mean$_{ACC\ eGFP}$= 75795.5), the same analysis compared BLA hM4Di

(Mean$_{BLA\ hM4Di}$= 44399.54) and BLA eGFP (Mean$_{BLA\ eGFP}$= 56519.33). No comparison with eGFP

was conducted for OFC hM4Di (Mean$_{OFC\ hM4Di}$= 100132). However, there were no significant

differences between ACC and OFC hM4Di expression (p=0.26), and we show below that control

groups were no different from each other and could be collapsed. We found no significant

differences in the spread between DREADDs hM4Di and control eGFP for ACC (t(18.52)= -1.77,

p = 0.093, -95% CI [27850, 2355]) or for BLA t(2.46)= 0.56, p = 0.624, 95% CI [-66729, 90969]).

Data Analysis.

All analyses were performed via custom-written code in MATLAB (MathWorks, Inc.,

Natick, MA). There were 3 main conditions in our analyses: 1) preference 2) latency to choose,

and 3) response rate during the 60-s cue duration.

Learning and performance data were analyzed with a series of mixed-effects General

Linear Models (GLMs) (*fitglme* function; Statistics and Machine Learning Toolbox) first in

omnibus analyses that included all factors (drug, virus and sex), with all brain regions (ACC, BLA

and OFC). Analyses were further pursued pending significant predictors and interactions in the

full model. All post-hoc tests were corrected for the number of comparisons (Bonferroni).

Figure 3. Preference was stable, and response latency and response rate (RR) exhibited the typical phenotype. (A) Mean change in preference for the info alternative between the last three sessions of training and baseline after surgery, with individual rats represented as scatter plots. Note that change in preference scale (y axis) ranges between -1 to 1. **(B)** Density of response latencies on the last three sessions of training are shown for each trial type. Dashed lines indicate the median latency. A marginally significant difference was found between trial types, where responses in choice trials were faster than in Forced (Info and No Info) trials. **(C)** Violin plots of the distribution of total RR during the 60s cue presentations are presented for each cue during the last three sessions of training. Dots indicate median RR for each cue. A significant effect of cue was found. **p<0.01, +p=0.06, ns = nonsignificant.

We also fit a logistic regression, a type of GLM for binary classification, to each rat's data

that belonged to either an experimental group or a control condition. We aimed to predict the

probability of choice using 3 features: 1) Latency to choose, 2) Previous choice, and 3) Previous

reward. We used only free-choice data for the outcome variable, and the previous choice and

reward as either from free-choice or forced-choice data. The full model corresponds to *y ~ [1 +*

PrevChoice + PrevReward + PrevLatency], with a binomial distribution.

Statistical significance for all analyses was noted when p-values were less than 0.05, and

p-values between 0.05 and 0.06 were noted as trending toward significance.

Results

Control group. Our control group was created by combining the animals that were

infused with the eGFP virus, but also animals that were originally assigned to the eGFP group

but wherein the histological analysis revealed either no-expression (n=9) or unilateral

expression (n=4) compared with bilateral expression of eGFP (n=9). To assess if these groups

could be collapsed in further analyses, we performed a GLM for each phase of the experiment.

Using the mean of the last 3 sessions of training, a GLM was conducted for mean preference

using virus, sex, and control group (eGFP, unilateral, or no-expression) as between-subject

factors, and individual rat as a random factor (full model: $y \sim [1 + sex * control\ group + (1\ |\ rat)]$).

We found no significant effect of control group (p= 0.54), sex (p = .335), or control group * sex

interaction (p= .955). Similarly, a GLM was performed on *preference change* during the stable

preference phase of the experiment, adding drug as a within-subject factor (full model: $y \sim [1 +$

*drug * sex*control group + (1 + drug| rat)*]). We found no significant predictors or interactions.

Based on these results, the animals in the control groups were collapsed and treated as a single

group for subsequent analysis, and added to the 'virus' factor as a fourth group (ACC hM4Di,

BLA hM4Di, OFC hM4Di, and control).

Training: Rats developed a consistent preference, responded quickly during choice trials and

responded more to the informative cue (S+).

294   *Choice*. Training data for preference for the info option were first analyzed across all

295 sessions, to evaluate learning of preference on the task. A GLM was conducted for preference

296 for the info alternative using sex as between-subject factor, session as within-subjects factor,

297 and individual rat as a random factor using the following formula: *γ ~ [1 + sex*session +*

298 *(1+session| rat)*]. We found a main effect of session (GLM: $\beta_{session}$= -0.008, t(868)=-2.09, *p*

299 =0.03) and sex (GLM: β_{sex}= -0.12, t(868)=-2.34, *p* =0.02) but no significant interaction (GLM:

300 $\beta_{session*sex}$= 0.006, t(868)=1.13, *p* =0.26) (see **Supplement 1A**).

301   To evaluate wether the observed preference remained consistent within each animal,

302 we compared preference in no-drug conditions before any drug experience. To do so, the last 3

303 sessions of preference data for training were averaged, and compared to the preference in the

304 last three sessions of the baseline after surgery. A GLM comparing the averaged preference

305 during training and baseline phase was performed, in which phase (training and baseline) was a

306 within-subject factor, and sex was a between-subject factor (full model: *γ ~ [1 + phase * sex (1 +*

307 *phase| rat)*]. No significant effects (p_{sex}= 0.93, p_{phase}= 0.26) or interactions ($p_{sex*phase}$= 0.73) were

308 found, suggesting that preference remained stable across conditions (See **Figure 3A**).

309   *Latencies.* A GLM was conducted on median latencies in the last three sessions of

310 training using trial type (forced info, forced no-info, and choice) as within-subject factors, sex as

311 a between-subject factor, and individual rat as a random factor using the following formula: *γ ~*

312 *[1 + trial type * sex + (1 + trial type| rat)*]. The results yielded a trend for an effect of trial type

313 (GLM: $\beta_{trial\ type}$= 6.16, t(179)=1.85, *p* =0.06) wherein rats tended to be faster during choice trials

314 (Median$_{Choice}$= 10.6 s, SEM±1.69) than either of the forced trials (Median$_{Forced\ Info}$=21.8

s,SEM±3.97, and Median$_{\text{Forced No-Info}}$= 18.1 s, SEM±3.81) (See **Figure 3B**). This result replicates

what previous literature has shown in this task in pigeons and starlings (González et al., 2023).

Response Rate. The median response rate (RR) during the 60 s cue was analyzed for the

last 3 sessions of training. A GLM was conducted for median RR using sex as between-subject

factor, and individual rat and cue (S+, S- and S3) as random factors using the following formula:

γ ~ [1 + sex + (1|rat:cue)]. Cue was introduced as a random factor to account for the fact that

rats were presented with different frequencies of cues given the programmed contingencies

(e.g., S+ present only 20% of all Info trials) but this difference also depended on the rat's choice.

For instance, an animal that chooses the No-Info alternative exclusively during choice trials

would have more presentations of cue S3 than any other cue. We found a main effect of cue

(GLM: β_{cue}= -2.11, t(185)=-2.89, *p* = 0.004), indicating that RR was greater for the cue predicting

food (Median$_{\text{S+}}$= 11 presses/minute, SEM±0.34) than to the no-info cue (Median$_{\text{S3}}$= 6

presses/minute ,SEM±0.14), but the lowest RR was to the cue predicting absence of food

(Median$_{\text{S-}}$= 3 presses/minute , SEM±0.06). This also is in line with previous work in which the

lowest RR was reported for S-, and typically a greater RR for S+ than S3 (Gonzalez & Blaisdell,

2021; Hinnenkamp et al., 2017).

Info Preference: Inhibition of ACC destabilized preference in female but not male rats.

Choice. Stable preference was analyzed by averaging the three sessions of baseline, and

then comparing this choice preference with the 3 sessions of CNO and three sessions of VEH for

each subject. We compared changes in stable preference by calculating the absolute difference

between preference in baseline-to-CNO and baseline-to-VEH conditions (See **Figure 4A**). A GLM

was conducted for *absolute preference change* using drug as a within-subject factor, virus and

sex as between-subject factors, and individual rat as random factor using the following formula:

γ ~ [1 + virus * drug * sex + (1 + drug| rat)]. We found a significant interaction of drug*virus

(GLM: $\beta_{drug*virus}$= 0.065, t(124)=4.88, *p* = 3.24^{e-06}) and sex*drug*virus (GLM: $\beta_{sex*drug*virus}$ = -0.065,

t(124)=-3.15, *p* = 0.002). Thus, we were justified to conduct follow-up analyses with individual

group comparisons: *γ ~ [1 + virus * drug * sex + (1 + drug| rat)]*. For the comparison of ACC

hM4Di vs Control, we found a significant interaction of drug*virus (GLM: $\beta_{drug*virus}$= 0.074,

t(68)=5.58, *p* = 4.61^{e-07}) and drug*virus*sex interaction (GLM: $\beta_{sex*virus*drug}$ = -0.077, t(68)= -3.70,

p = 0.0004). Comparisons of OFC hM4Di vs. Control and BLA hM4Di vs. Control were not

significant. Post-hoc analysis using Bonferroni correction and sex as a covariate for the ACC

hM4Di group resulted in a significant effect of drug (t(29)= 3.66, *p* =0.002). In contrast, there

was no significant effect of drug in the control group (t(41)= 1.59, *p* = 0.239). Overall, these

results found that ACC inhibition causes a desestabilization on preference for the female rats.

Latencies. Latency to choose was analyzed by calculating the median for CNO and VEH

sessions (See **Figure 4B**). A GLM was conducted for median latency using drug as within-subject

factor, virus and sex as between-subject factors, and individual rat as a random factor using the

following formula: *γ ~ [1 + trial type * virus * drug * sex + (1 + trial type + drug| rat)]*. We found

a main effect of trial type (GLM: $\beta_{trial\ type}$= 5.61, t(380)=2.19, *p* =0.028) indicating that rats were

faster in choosing during choice trials (Median$_{Choice}$= 8.4 s, SEM±0.4) than forced trials

(Median$_{Forced\ Info}$=14.8 s,SEM±0.66, and Median$_{Forced\ No-Info}$= 15.1 s, SEM±0.62). Note that we did

not find any difference by sex or virus, suggesting that the change in preference observed in the

female rats following ACC inhibition did not affect decision speed.

Response rate. RR during the 60s cue duration was also analyzed by calculating the

median during CNO and VEH sessions (See **Figure 4C**). A GLM was conducted for median RR

using drug as within-subject factor, virus, and sex as between-subject factors, and individual rat

Figure 4. Inhibition of ACC destabilized info preference in female but not male rats, while latencies and response rate (RR) were unaltered by inhibition. **(A)** Baseline Info preference for females (top) and males (bottom) over two baselines. The first baseline (BL1) was obtained after surgery and the second baseline (BL2) was obtained at the end of the experiment. Note the lack of dramatic changes within individual animals. **(B)** Mean change in preference for the info alternative between the last three sessions of baseline after surgery minus the preference during VEH and CNO administration for females (top) and males (bottom). Note that a positive change indicates a shift in preference towards the non-info option, whereas a negative change indicates a preference shift toward the info option. Values around 0 indicate no change in preference. **(C)** Density of latencies to choose across drug conditions is shown for each trial type. Dashed lines indicate the median latency. A significant difference between trial types was found, where choice trials were faster than Forced (Info and No Info) trials. **(D)** Violin plots of the distribution of total RR during the 60s cue presentations are presented for each cue for all drug conditions for females (left) and males (right). Dots indicate median response rate for each cue. Females responded more than males and a significant effect of cue was found. **p<0.01, *p=0.06.

and cue (S+, S- and S3) as random factors using the following formula for the full model: $y \sim [1 +$

*cue * virus * drug * sex + (1 + drug| rat:cue)*]. As in training, cue was defined as a random factor

given that there were different probabilities of experiencing each cue based on an individual

rat's choices. We found a main effect of sex (GLM: β_{sex}= -7.87, t(379)=-2.17, *p* = 0.03), with

females showing higher response rates (Median$_{Female}$= 7 presses/min , SEM±0.06) than males

(Median$_{Male}$= 4 presses/min , SEM±0.46).

Learning of new cues: Inhibition of BLA tended to decrease learning of new informative cues.

Choice. Preference for the info alternative was analyzed across all sessions to evaluate

learning about the informative and non-informative cues for rats administered CNO or VEH.

Similar to above, a GLM was conducted for preference using session as within-subject factor,

virus, sex, brain and drug as between-subject factors, and individual rat as random factors using

the full model formula: *γ ~ [1 + session* virus * drug * sex + (1 + session| rat)]*. We found a main

effect of drug*sex (GLM: $\beta_{drug*sex}$= -0.54, t(978)= -3.00, *p* = 0.003), sex*virus (GLM: $\beta_{sex*virus}$= -

0.17, t(978)= -2.28, *p* = 0.02), and drug*sex*virus (GLM: $\beta_{drug*sex*virus}$= 0.39, t(978)=-3.96, *p* =

7.89^{e-05}). Given this result, we excluded session as a factor, and sex was added as a covariate

according to the formula: *γ ~ [1 + sex + virus * drug + (1| rat)]*. We found a main effect of virus

(GLM: β_{virus}= -0.077, t(989)=-2.03, *p* = 0.04), and drug*virus interaction (GLM: $\beta_{drug* virus}$= 0.13,

t(989)=-2.77, *p* = 0.005).

The interaction above justified a GLM analysis on preference separated by brain region

vs. the control group using the formula: *γ ~ [1 + virus * drug + (1| rat)]*. Separate GLMs for OFC

hM4Di and for ACC hM4Di compared to the control group resulted in no significant predictors

or interactions. Only a GLM for BLA hM4Di vs control resulted in a main effect of virus (GLM:

β_{virus}= -0.13, t(533)=-2.75, *p* = 0.006) and marginal interaction of virus*drug (GLM: $\beta_{drug*virus}$=

0.14, t(533)=1.87, *p* = 0.06). However, post-hocs tests using Bonferroni correction did

not result in a significant effect of drug in either BLA hM4Di (p = .127) or control (p = .528)

groups (**Figure 5**).

Figure 5. Effect of inhibition on info preference change during acquisition of new cues. Average change in preference for the info alternative, between the first and last three sessions of new cues acquisition for BLA, OFC, ACC and control animals during VEH or CNO administration. Note that a positive change indicates a shift in preference towards the non-info option, whereas a negative change indicates a preference shift toward the info option. Values around 0 indicate no change in preference. There was a trend for a decrease in learning the new info option following BLA inhibition. p=0.06 (drug x virus mixed-effects GLM).

Latencies. We performed GLM analysis on median latencies separated by brain region

and compared against the control group: $γ \sim [1 + session*sex * virus*drug*trial type + (1+trial$

$type*session | rat)]$. Though we highlight several 3- and 4-way interactions in **Table 1A-C**, there

was no significant interaction of drug * virus.

RR. We performed GLM analysis on median RR separated by brain region and compared

against the control group: *γ ~ [1 + session*sex* virus*drug*trial type + (1+trial type*session|*

rat)]. No significant effects were found for BLA vs control. RR for ACC vs control found a

significant effect of session*drug (*p* = 0.48) and session*cue*drug (p = 0.035). RR for OFC vs

control found only an effect of session*cue*drug (*p* = 0.041). However, there was no significant

interaction of drug * virus in any group (**Supplement 1B**).

Previous choices are not predictive of current information choices in ACC-inhibited females

To further understand the nature of the effect of ACC inhibition on choice behavior, we

fit a logistic regression, a type of GLM for binary classification, to each rat's data that belonged

Figure 6. *Female rats do not use previous choices to guide choices involving information during ACC inhibition.* Each panel represents a feature of the model (Previous Choice, Previous Reward, and Latency) for female animals following CNO administration in four different groups: ACC hM4Di, BLA hM4Di, OFC hM4Di and Control. The y-axis is the odds ratio of the predictive feature. Box, error bars, and line represent 80% confidence interval, 95% confidence interval, and mean odds ratio, respectively. Mean model R^2 for control group = 0.4613, ACC group = 0.4189, OFC group = 0.4613, BLA group = 0.5181.

to either the test group (female ACC hM4Di-CNO) or control (female control) condition. Using

the three features described in the analysis section we aimed to predict the probability of

choice. We computed the odds ratio and 95%, 80% CI for each condition's regression fit: If an

odds ratio was greater than 1, it represented an increase in the odds of the outcome happening

given a one-unit increase in the predictor. If the odds ratio was less than 1, it represented a

decrease in the odds of the outcome happening given a one-unit increase in the predictor. And

finally, an odds ratio of exactly 1 signified that the value of the predictor did not affect the

probability of the outcome.

The model resulted in a significant effect of Previous Choice (t(651)= 11.19, p = 4.38^{e-29})

and Previous Reward (t(651)= -3.07, p = 0.002), but no effect of Latency (t(651)= -0.84, p =

0.40). We found that in the control group, previous choice was a strong predictor for the next

choice (n= 95.45%). Similarly, most BLA hM4Di animals (n = 91%) showed that previous choice

was a good predictor of future choices. Interestingly, a similar result to ACC inhibition occurs

with the OFC hM4Di group in which previous choices do not predict future behavior well.

Finally, both control and BLA hM4Di groups used previous reward as a predictor (86.36% of rats

showed a significant effect). In contrast, only 20% of ACC-inhibited animals showed good

prediction of choice from previous reward, suggesting that inhibition of ACC changes choice

patterns uniquely (**Figure 6**).

These findings reveal dissociable patterns of choice prediction across different brain

areas but also suggest that previous choices have the most predictive impact on subsequent

behavior for all conditions. Conversely, previous rewards reduce the odds of the event of

interest across all conditions, and latency instead has no predictive impact (Odds Ratio close to

1).

Discussion

429    Despite the well-characterized roles of OFC and ACC in decision-making (Bromberg-

430 Martin & Monosov, 2020; Izquierdo, 2017; Sosa et al., 2021), few studies find a clear

431 dissociation between the role of both regions in rodents. For example, both OFC and ACC are

432 important in confidence report as measured by temporal wagering (Stolyarova et al. 2019; Lak

433 et al. 2014) and are also involved in stimulus-based reversal learning (Ye et al. 2023). In the

434 present study, we found a clear dissociation between ACC and OFC that may shed light on their

435 individual contributions in decision-making about non-instrumental information, and point to a

436 hierarchy of functions within rodent frontal cortex. Specifically, we found that ACC inhibition

437 rendered animals' decisions about information more stochastic, also corroborated by the

438 logistic regression model which showed that previous choice was not a good predictor of future

439 choices when ACC was offline. The pattern of results on latencies and response rate also

440 indicates the effect of inhibition on decision-making is not due to performance decrements (i.e.,

441 we found unchanged latencies and response rates). Interestingly, the logistic regression

442 revealed a similar finding of reduced use of previous choice following OFC inhibition, but this

443 did not translate into a significant change in behavior. Thus, it could be that "monitoring" is a

444 general feature of rodent frontal cortex, with ACC exerting a more prominent role than OFC.

445 Indeed, ACC has been linked more to the representation of reward opportunities across the

446 environment at different timescales while also keeping track of animals' previous actions

447 (Kolling et al., 2016; Spitmaan et al., 2020; Wittmann et al., 2016), suggestive of a more

448 metacognitive or performance monitoring role (Kane et al., 2022; Stolyarova, Rakhshan, Hart,

449 O'Dell, et al., 2019; Takeuchi et al., 2022; van Veen et al., 2004). Instead, BLA inhibition did not

450 alter stable performance but it resulted in a tendency to affect learning of new informative

cues, supporting the role of BLA in biasing choices towards larger rewards in uncertain

environments (Ghods-Sharifi et al., 2009; St Onge et al., 2012).

Our results presented here also replicate the vast literature investigating the behavior

behind the preference for the informative, yet "suboptimal" alternative. As other researchers

have previously reported, the preference for the informative option is more variable in rats

than pigeons or starlings when the informative option results in less food (Stagner & Zentall,

2010; Vasconcelos et al., 2015). This indicates that perhaps there are different sensitivities to

information and/or reinforcement history between species. Regarding latencies to make a

choice, we replicate what has been observed in several studies: animals respond faster on

"true" choice trials than forced-choice trials. Ecologists have discussed this counterintuitive

result by suggesting that animals did not evolve to make simultaneous but rather sequential

choices in nature, such that latencies during forced choice trials can be used to predict

preference (Kacelnik et al., 2010). Finally, response rate during the cue presentation followed

the typical pattern reported previously: animals respond more in the presence of the always

and partially-reinforced cue, and do not respond to the non-reinforced cue (Gonzalez &

Blaisdell, 2021; Hinnenkamp et al., 2017). It is important to highlight that animals are not

required to respond during the duration of the cue; however, they typically do, suggesting a

Pavlovian association established between the cue and the outcome. This association we

believe is central to the preference, in which a stable preference emerges when an association

between the reinforced cue (S+) and the outcome and the non-reinforced cue (S-) and the

outcome is established (González et al., 2023).

472   A similar variation of the task presented in this study was used to assess preference for

473 non-instrumental information in humans and monkeys. In those studies, subjects are given a

474 choice between two alternatives: one that provides informative cues that indicate the trial's

475 outcome, and another provides non-informative cues that do not indicate the trial's outcome.

476 Similar to our task, the information provided by the cues does not influence or change the

477 outcome. However, humans and monkeys were also informed about the quantity (money for

478 humans and juice for monkeys) of the outcome in a given trial. The results showed that

479 macaque monkeys and humans prefer information, and they were willing to sacrifice

480 water/money to obtain immediate information about the outcomes. One group previously

481 found that OFC neurons encode variables that are relevant in learning and decision-making but

482 do not integrate these variables into a single value (Blanchard et al., 2015). This result is

483 consistent with our findings in which OFC inactivation resulted in an altered pattern of using

484 previous choice information revealed by the logistic regression, suggesting problems in

485 retrieving the memory of previous choice, but this did not translate into a real change in

486 preference. ACC may have a higher-level role in sustaining information-seeking (Blanchard et

487 al., 2015). For example, activity ramps up in this region in anticipation of information becoming

488 available to resolve uncertainty before reward delivery (White et al., 2019). Our results support

489 the monitoring role of ACC in decisions about information, where inhibition altered the

490 integration of value, destabilizing preference.

491   We found that ACC inhibition desestabilized preference, but only in female animals. We

492 did not find changes in latency to choose or response rates, indicating that the observed

493 differences were not due to motor impairment or due to changes in the association between

cues and outcomes. Therefore, the change in preference with ACC inhibition is unlikely due to

problems in accessing overall value of each option. Previous studies have determined that ACC

does not support simple effort or the hedonic value of a given alternative (i.e., one option).

Instead, it computes the value *across* options, that is, it tracks what is the overall "best" choice

(Hart et al., 2020; Hart et al., 2017). Here, we found that the change in preference following

ACC inhibition was not in any particular direction. If animals with ACC offline had issues in

accessing the overall value of the best option, in this case the no-info alternative, we would

have observed an increase in preference for the info alternative. In contrast, we found that

preference became more stochastic. The preference change may also indicate a reduced ability

to link previous actions (in this case, the previous lever press) to the current trial. However, this

is also unlikely because the motor response to lever press did not change (latencies were

unaffected), nor did the assigned value of the cues (response rate to cues also did not change).

We propose instead that ACC inhibition results in an impairment in accessing the value of

information. Previous research using this paradigm indicates that the contrast between both

informative cues (i.e., the difference in information between the S+ and the S-) is essential for

the development of a preference for information. Previous research has already suggested that

ACC is important for information-seeking behavior; however, previous studies reported this

following electrophysiological recording and neuroimaging in monkeys and humans,

respectively (Blanchard et al., 2015; Bromberg-Martin & Hikosaka, 2009, 2011; Bromberg-

Martin & Monosov, 2020; Wallis, 2012). Our study provides the first causal evidence of the

importance of ACC in decision-making involving non-instrumental information.

Sex differences in the involvement of ACC in value-based decision making have been

reported before. In a recent study, Cox et al. (2023) found that ACC inhibition disrupted the

relationship between the value of each alternative and motivation to engage in the task in

female mice. However, they did not find changes in preference as we found in our study. This

group also reported that the ACC-to-dorsomedial striatal neuron pathway represents negative

outcomes more strongly in female than males rats, suggesting different sensitivity to negative

feedback (Cox et al., 2023). Similarly, in tasks involving decision-making under risk, it has been

determined that female rats are more risk-averse (Orsini et al., 2016), indicating that female

and male rats use different strategies to make decisions (Orsini et al., 2021). In our task, similar

to Cox et al. (2023), we did not find differences in preference between female and male rats

before inhibition. However, the destabilization of preference might indicate that females rely

more strongly on ACC to keep track of reward statistics. Indeed, our model of trial-by-trial

choice indicated that when ACC was inhibited, previous choice was not a good predictor of

preference, whereas it remained so in other conditions.

Similar to the results in rodents, groups studying human subjects have uncovered sex

differences in strategy in decision-making (Chen et al., 2021; Chowdhury et al., 2019), usually in

tasks involving risk (van den Bos et al., 2013). These sex differences are important to

understand because, on the one hand, there is a higher prevalence of depression or anxiety-

related disorders in women (Cyranowski et al., 2000); and on the other hand, there is increasing

evidence that neuropsychiatric conditions such as ADHD, bipolar disorder, and autism show

different onset, symptom severity, and prognosis dependent on sex (Grissom & Reyes, 2019;

Hwang et al., 2020). This evidence suggests differential involvement of circuits involved in

decision-making by sex, and our study contributes to the effort in finding the neural

mechanisms behind these potential differences.

540   **Author contributions**

ABP: Resources, funding, writing- review & editing. **AI:** Conceptualization, methodology, formal

analysis, data curation, resources, writing-original draft, writing- review & editing, software,

supervision, and funding. **AR:** Investigation and data curation. **IY:** Investigation and data

curation. **JLR:** Investigation, writing- review & editing. **SA:** Investigation, writing- original draft,

and data curation. **VVG**: Conceptualization, methodology, formal analysis, investigation, data

curation, writing- original draft preparation, writing- review & editing, software, visualization,

and funding. **YZ:** Software and formal analysis.

References

Aguirre, C. G., Woo, J. H., Sosa, J. L. R., Munier, J. J., Perez, J., Goldfarb, M., Das, K., Gomez, M., Ye, T., Pannu, J., Evans, K., Oa Neill, P. R., Spigelman, I., Soltani, A., & Izquierdo, A. (2023). Sex-dependent contributions of ventrolateral orbitofrontal cortex and basolateral amygdala to learning under uncertainty. *bioRxiv*. https://doi.org/10.1101/2023.04.03.535471

Blanchard, T. C., Hayden, B. Y., & Bromberg-Martin, E. S. (2015). Orbitofrontal cortex uses distinct codes for different choice attributes in decisions motivated by curiosity. *Neuron*, *85*(3), 602-614. https://doi.org/10.1016/j.neuron.2014.12.050

Boulter, C., Freeston, M., South, M., & Rodgers, J. (2014). Intolerance of uncertainty as a framework for understanding anxiety in children and adolescents with autism spectrum disorders. *J Autism Dev Disord*, *44*(6), 1391-1402. https://doi.org/10.1007/s10803-013-2001-x

Bromberg-Martin, E. S., & Hikosaka, O. (2009). Midbrain dopamine neurons signal preference for advance information about upcoming rewards. *Neuron*, *63*(1), 119-126. https://doi.org/10.1016/j.neuron.2009.06.009

Bromberg-Martin, E. S., & Hikosaka, O. (2011). Lateral habenula neurons signal errors in the prediction of reward information. *Nat Neurosci*, *14*(9), 1209-1216. https://doi.org/10.1038/nn.2902

Bromberg-Martin, E. S., & Monosov, I. E. (2020). Neural circuitry of information seeking. *Curr Opin Behav Sci*, *35*, 62-70. https://doi.org/10.1016/j.cobeha.2020.07.006

Chen, C. S., Knep, E., Han, A., Ebitz, R. B., & Grissom, N. M. (2021). Sex differences in learning from exploration. *Elife*, *10*, e69748. https://doi.org/10.7554/eLife.69748

Chowdhury, T. G., Wallin-Miller, K. G., Rear, A. A., Park, J., Diaz, V., Simon, N. W., & Moghaddam, B. (2019). Sex differences in reward- and punishment-guided actions. *Cogn Affect Behav Neurosci*, *19*(6), 1404-1417. https://doi.org/10.3758/s13415-019-00736-w

Cox, J., Minerva, A. R., Fleming, W. T., Zimmerman, C. A., Hayes, C., Zorowitz, S., Bandi, A., Ornelas, S., McMannon, B., Parker, N. F., & Witten, I. B. (2023). A neural substrate of sex-dependent modulation of motivation. *Nat Neurosci*, *26*(2), 274-284. https://doi.org/10.1038/s41593-022-01229-9

Cyranowski, J. M., Frank, E., Young, E., & Shear, M. K. (2000). Adolescent onset of the gender difference in lifetime rates of major depression: a theoretical model. *Arch Gen Psychiatry*, *57*(1), 21-27. https://doi.org/10.1001/archpsyc.57.1.21

Dugas, M. J., Buhr, K., & Ladouceur, R. (2004). The Role of Intolerance of Uncertainty in Etiology and Maintenance. In *Generalized anxiety disorder: Advances in research and practice.* (pp. 143-163). The Guilford Press.

Fortes, I., Vasconcelos, M., & Machado, A. (2016). Testing the boundaries of "paradoxical" predictions: Pigeons do disregard bad news. *J Exp Psychol Anim Learn Cogn*, *42*(4), 336-346. https://doi.org/10.1037/xan0000114

Ghods-Sharifi, S., St Onge, J. R., & Floresco, S. B. (2009). Fundamental contribution by the basolateral amygdala to different forms of decision making. *J Neurosci*, *29*(16), 5251-5259. https://doi.org/10.1523/JNEUROSCI.0315-09.2009

Gonzalez, V. V., & Blaisdell, A. P. (2021). The role of inhibition in the suboptimal choice task. *J Exp Psychol Anim Learn Cogn*, *47*(4), 429-444. https://doi.org/10.1037/xan0000300

González, V. V., Izquierdo, A., & Blaisdell, A. P. (2023). Theoretical Mechanisms of Paradoxical Choices Involving Information. *Comparative Cognition & Behavior Reviews*, *18*, 11-31. https://doi.org/10.3819/ccbr.2023.180002

Grissom, N. M., & Reyes, T. M. (2019). Let's call the whole thing off: evaluating gender and sex differences in executive function. *Neuropsychopharmacology*, *44*(1), 86-96. https://doi.org/10.1038/s41386-018-0179-5

Hart, E. E., Blair, G. J., O'Dell, T. J., Blair, H. T., & Izquierdo, A. (2020). Chemogenetic Modulation and Single-Photon Calcium Imaging in Anterior Cingulate Cortex Reveal a Mechanism for Effort-Based Decisions. *J Neurosci*, *40*(29), 5628-5643. https://doi.org/10.1523/JNEUROSCI.2548-19.2020

Hart, E. E., Gerson, J. O., Zoken, Y., Garcia, M., & Izquierdo, A. (2017). Anterior cingulate cortex supports effort allocation towards a qualitatively preferred option. *Eur J Neurosci*, *46*(1), 1682-1688. https://doi.org/10.1111/ejn.13608

Hinnenkamp, J. E., Shahan, T. A., & Madden, G. J. (2017). How suboptimal is suboptimal choice? *J Exp Anal Behav*, *107*(1), 136-150. https://doi.org/10.1002/jeab.239

Hwang, Y. I. J., Arnold, S., Srasuebkul, P., & Trollor, J. (2020). Understanding anxiety in adults on the autism spectrum: An investigation of its relationship with intolerance of uncertainty, sensory sensitivities and repetitive behaviours. *Autism*, *24*(2), 411-422. https://doi.org/10.1177/1362361319868907

Iigaya, K., Hauser, T. U., Kurth-Nelson, Z., O'Doherty, J. P., Dayan, P., & Dolan, R. J. (2019). Hippocampal-midbrain circuit enhances the pleasure of anticipation in the prefrontal cortex. -. https://doi.org/10.1101/588699

Iigaya, K., Story, G. W., Kurth-Nelson, Z., Dolan, R. J., & Dayan, P. (2016). The modulation of savouring by prediction error and its effects on choice. *Elife*, *5*(APRIL2016). https://doi.org/10.7554/eLife.13747

Izquierdo, A. (2017). Functional Heterogeneity within Rat Orbitofrontal Cortex in Reward Learning and Decision Making. *J Neurosci*, *37*(44), 10529-10540. https://doi.org/10.1523/JNEUROSCI.1678-17.2017

Jenkinson, R., Milne, E., & Thompson, A. (2020). The relationship between intolerance of uncertainty and anxiety in autism: A systematic literature review and meta-analysis. *Autism*, *24*(8), 1933-1944. https://doi.org/10.1177/1362361320932437

Jenni, N. L., Symonds, N., & Floresco, S. B. (2023). Medial orbitofrontal cortical regulation of different aspects of Pavlovian and instrumental reward seeking. *Psychopharmacology (Berl)*, *240*(3), 441-459. https://doi.org/10.1007/s00213-022-06265-8

Jezzini, A., Bromberg-Martin, E. S., Trambaiolli, L. R., Haber, S. N., & Monosov, I. E. (2021). A prefrontal network integrates preferences for advance information about uncertain rewards and punishments. *Neuron*, *109*(14), 2339-2352.e2335. https://doi.org/https://doi.org/10.1016/j.neuron.2021.05.013

Jo, S., & Jung, M. W. (2016). Differential coding of uncertain reward in rat insular and orbitofrontal cortex. *Sci Rep*, *6*, 24085. https://doi.org/10.1038/srep24085

Kacelnik, A., Vasconcelos, M., Monteiro, T., & Aw, J. (2010). Darwin's "tug-of-war" vs. starlings' "horse-racing": how adaptations for sequential encounters drive simultaneous choice. *Behavioral Ecology and Sociobiology*, *65*(3), 547-558. https://doi.org/10.1007/s00265-010-1101-2

Kane, G. A., James, M. H., Shenhav, A., Daw, N. D., Cohen, J. D., & Aston-Jones, G. (2022). Rat Anterior Cingulate Cortex Continuously Signals Decision Variables in a Patch Foraging Task. *J Neurosci*, *42*(29), 5730-5744. https://doi.org/10.1523/JNEUROSCI.1940-21.2022

Kolling, N., Behrens, T., Wittmann, M. K., & Rushworth, M. (2016). Multiple signals in anterior cingulate cortex. *Curr Opin Neurobiol*, *37*, 36-43. https://doi.org/10.1016/j.conb.2015.12.007

Lak, A., Costa, G. M., Romberg, E., Koulakov, A. A., Mainen, Z. F., & Kepecs, A. (2014). Orbitofrontal cortex is required for optimal waiting based on decision confidence. *Neuron*, *84*(1), 190-201. https://doi.org/10.1016/j.neuron.2014.08.039

Macias, A., Gonzalez, V. V., Machado, A., & Vasconcelos, M. (2021). The functional equivalence of two variants of the suboptimal choice task: choice proportion and response latency as measures of value. *Anim Cogn*, *24*(1), 85-98. https://doi.org/10.1007/s10071-020-01418-8

Mandali, A., Sethi, A., Cercignani, M., Harrison, N. A., & Voon, V. (2021). Shifting uncertainty intolerance: methylphenidate and attention-deficit hyperactivity disorder. *Transl Psychiatry*, *11*(1), 12. https://doi.org/10.1038/s41398-020-01118-4

Monosov, I. E. (2017). Anterior cingulate is a source of valence-specific information about value and uncertainty. *Nat Commun*, *8*(1), 134. https://doi.org/10.1038/s41467-017-00072-y

Namboodiri, V. M. K., Otis, J. M., van Heeswijk, K., Voets, E. S., Alghorazi, R. A., Rodriguez-Romaguera, J., Mihalas, S., & Stuber, G. D. (2019). Single-cell activity tracking reveals that orbitofrontal neurons acquire and maintain a long-term memory to guide behavioral adaptation. *Nat Neurosci*, *22*(7), 1110-1121. https://doi.org/10.1038/s41593-019-0408-1

Orsini, C. A., Blaes, S. L., Hernandez, C. M., Betzhold, S. M., Perera, H., Wheeler, A. R., Ten Eyck, T. W., Garman, T. S., Bizon, J. L., & Setlow, B. (2021). Regulation of risky decision making by gonadal hormones in males and females. *Neuropsychopharmacology*, *46*(3), 603-613. https://doi.org/10.1038/s41386-020-00827-0

Orsini, C. A., Willis, M. L., Gilbert, R. J., Bizon, J. L., & Setlow, B. (2016). Sex differences in a rat model of risky decision making. *Behav Neurosci*, *130*(1), 50-61. https://doi.org/10.1037/bne0000111

Procyk, E., Tanaka, Y. L., & Joseph, J. P. (2000). Anterior cingulate activity during routine and non-routine sequential behaviors in macaques. *Nat Neurosci*, *3*(5), 502-508. https://doi.org/10.1038/74880

Riceberg, J. S., & Shapiro, M. L. (2017). Orbitofrontal Cortex Signals Expected Outcomes with Predictive Codes When Stable Contingencies Promote the Integration of Reward History. *J Neurosci*, *37*(8), 2010-2021. https://doi.org/10.1523/JNEUROSCI.2951-16.2016

Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T., & Eliceiri, K. W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*, *18*(1), 529. https://doi.org/10.1186/s12859-017-1934-z

Rushworth, M. F. S., & Behrens, T. E. J. (2008). Choice, uncertainty and value in prefrontal and cingulate cortex. In *Nature Neuroscience* (Vol. 11, pp. 389-397): Nature Publishing Group.

Schneider Gasser, E. M., Straub, C. J., Panzanelli, P., Weinmann, O., Sassoe-Pognetto, M., & Fritschy, J. M. (2006). Immunofluorescence in brain sections: simultaneous detection of presynaptic and postsynaptic proteins in identified neurons. *Nat Protoc*, *1*(4), 1887-1897. https://doi.org/10.1038/nprot.2006.265

Shidara, M., & Richmond, B. J. (2002). Anterior cingulate: Single neuronal signals related to degree of reward expectancy. *Science*, *296*(5573), 1709-1711. https://doi.org/10.1126/science.1069504

Soltani, A., & Izquierdo, A. (2019). Adaptive learning under expected and unexpected uncertainty. In *Nature Reviews Neuroscience* (Vol. 20, pp. 635-644): Nature Publishing Group.

Sosa, J. L. R., Buonomano, D., & Izquierdo, A. (2021). The orbitofrontal cortex in temporal cognition. *Behav Neurosci*, *135*(2), 154-164. https://doi.org/10.1037/bne0000430

Spitmaan, M., Seo, H., Lee, D., & Soltani, A. (2020). Multiple timescales of neural dynamics and integration of task-relevant signals across cortex. *Proc Natl Acad Sci U S A*, *117*(36), 22522-22531. https://doi.org/10.1073/pnas.2005993117

St Onge, J. R., Stopper, C. M., Zahm, D. S., & Floresco, S. B. (2012). Separate prefrontal-subcortical circuits mediate different components of risk-based decision making. *J Neurosci*, *32*(8), 2886-2899. https://doi.org/10.1523/JNEUROSCI.5625-11.2012

Stagner, J. P., & Zentall, T. R. (2010). Suboptimal choice behavior by pigeons. *Psychon Bull Rev*, *17*(3), 412-416. https://doi.org/10.3758/PBR.17.3.412

Stolyarova, A., Rakhshan, M., Hart, E. E., O'Dell, T. J., Peters, M. A. K., Lau, H., Soltani, A., & Izquierdo, A.
(2019). Contributions of anterior cingulate cortex and basolateral amygdala to decision
confidence and learning under uncertainty. *Nat Commun*, *10*(1), 4704.
https://doi.org/10.1038/s41467-019-12725-1
Stolyarova, A., Rakhshan, M., Hart, E. E., O'Dell, T. J., Peters, M. A. K., Lau, H., Soltani, A., & Izquierdo, A.
(2019). Dissociable roles for Anterior Cingulate Cortex and Basolateral Amygdala in Decision
Confidence and Learning under Uncertainty. *000*. https://doi.org/10.1101/655860
Stopper, C. M., & Floresco, S. B. (2011). Contributions of the nucleus accumbens and its subregions to
different aspects of risk-based decision making. *Cogn Affect Behav Neurosci*, *11*(1), 97-112.
https://doi.org/10.3758/s13415-010-0015-9
Takeuchi, D., Roy, D., Muralidhar, S., Kawai, T., Bari, A., Lovett, C., Sullivan, H. A., Wickersham, I. R., &
Tonegawa, S. (2022). Cingulate-motor circuits update rule representations for sequential choice
decisions. *Nat Commun*, *13*(1), 4545. https://doi.org/10.1038/s41467-022-32142-1
Tolin, D. F., Abramowitz, J. S., Brigidi, B. D., & Foa, E. B. (2003). Intolerance of uncertainty in obsessive-
compulsive disorder. *Journal of Anxiety Disorders*, *17*(2), 233-242.
https://doi.org/https://doi.org/10.1016/S0887-6185(02)00182-2
van den Bos, R., Homberg, J., & de Visser, L. (2013). A critical review of sex differences in decision-
making tasks: focus on the Iowa Gambling Task. *Behav Brain Res*, *238*, 95-108.
https://doi.org/10.1016/j.bbr.2012.10.002
van Veen, V., Holroyd, C. B., Cohen, J. D., Stenger, V. A., & Carter, C. S. (2004). Errors without conflict:
implications for performance monitoring theories of anterior cingulate cortex. *Brain Cogn*, *56*(2),
267-276. https://doi.org/10.1016/j.bandc.2004.06.007
Vasconcelos, M., Monteiro, T., & Kacelnik, A. (2015). Irrational choice and the value of information. *Sci
Rep*, *5*, 13874. https://doi.org/10.1038/srep13874
Wallis, J. D. (2012). Cross-species studies of orbitofrontal cortex and value-based decision-making. In
Nature Neuroscience (Vol. 15, pp. 13-19): NIH Public Access.
White, J. K., Bromberg-Martin, E. S., Heilbronner, S. R., Zhang, K., Pai, J., Haber, S. N., & Monosov, I. E.
(2019). A neural network for information seeking. *Nat Commun*, *10*(1), 5168.
https://doi.org/10.1038/s41467-019-13135-z
Winstanley, C. A., & Floresco, S. B. (2016). Deciphering Decision Making: Variation in Animal Models of
Effort- and Uncertainty-Based Choice Reveals Distinct Neural Circuitries Underlying Core
Cognitive Processes. *J Neurosci*, *36*(48), 12069-12079. https://doi.org/10.1523/JNEUROSCI.1713-
16.2016
Wittmann, M. K., Kolling, N., Akaishi, R., Chau, B. K., Brown, J. W., Nelissen, N., & Rushworth, M. F.
(2016). Predictive decision making driven by multiple time-linked reward representations in the
anterior cingulate cortex. *Nat Commun*, *7*(1), 12327. https://doi.org/10.1038/ncomms12327
Ye, T., Romero-Sosa, J. L., Rickard, A., Aguirre, C. G., Wikenheiser, A. M., Blair, H. T., & Izquierdo, A.
(2023). Theta oscillations in anterior cingulate cortex and orbitofrontal cortex differentially
modulate accuracy and speed in flexible reward learning. *Oxf Open Neurosci*, *2*, kvad005.
https://doi.org/10.1093/oons/kvad005
Zeeb, F. D., & Winstanley, C. A. (2011). Lesions of the basolateral amygdala and orbitofrontal cortex
differentially affect acquisition and performance of a rodent gambling task. *J Neurosci*, *31*(6),
2197-2204. https://doi.org/10.1523/JNEUROSCI.5597-10.2011
Zeeb, F. D., & Winstanley, C. A. (2013). Functional disconnection of the orbitofrontal cortex and
basolateral amygdala impairs acquisition of a rat gambling task and disrupts animals' ability to
alter decision-making behavior after reinforcer devaluation. *J Neurosci*, *33*(15), 6434-6443.
https://doi.org/10.1523/JNEUROSCI.3971-12.2013