A transmission bottleneck for malaria? Quantification of sporozoite expelling from laboratory and natural *P. falciparum* infections

Chiara Andolina*[1], Wouter Graumans*[1], Moussa Guelbeogo[2], Geert Jan van Gemert[1], Jordache Ramjith[1], Soré Harouna[2], Zongo Soumanaba[2], Rianne Stoter[1], Marga Vegte-Bolmer[1], Martina Pangos[3], Photini Sinnis[4], Katharine Collins[1], Sarah G Staedke[5], Alfred B Tiono[2], Chris Drakeley[6], Kjerstin Lanke[1] & Teun Bousema[1,6].

Author affiliation:

[1]Department of Medical Microbiology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; [2]Centre National de Recherche et de Formation sur le Paludisme, Ouagadougou, Burkina Faso; [3]Department of Plastic and Reconstructive Surgery, Azienda Ospedaliero Universitaria Giuliano Isontina Trieste, Italy; [4]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA; [5]Liverpool School of Tropical Medicine, Liverpool, UK; [6]Department of Immunology and Infection, London School of Hygiene and Tropical Medicine, London, UK.

Abstract

It is currently unknown whether all *Plasmodium falciparum* infected mosquitoes are equally infectious. We assessed sporogonic development using cultured gametocytes in the Netherlands and natural infections in Burkina Faso. We quantified the number of sporozoites expelled into artificial skin in relation to intact oocysts, ruptured oocysts, and residual salivary gland sporozoites. Sporozoites were quantified by highly sensitive qPCR; intact and ruptured oocysts by fluorescence microscopy following anti-circumsporozoite antibody staining. In laboratory conditions, higher total sporozoite burden in mosquitoes was associated with a shorter duration of sporogony (p<0.001). Overall, 53% (116/216) of *P. falciparum* infected *An. stephensi* mosquitoes expelled sporozoites into artificial skin. The geometric means of expelled and residual salivary gland sporozoites were 116 (interquartile range (IQR: 33-501) and 21,016 (IQR: 9127-78,380), respectively. There was a strong correlation between ruptured oocyst number and salivary gland sporozoite load (ρ=0.74; p<0.0001) and a weaker positive correlation between salivary gland sporozoite load and the number of sporozoites expelled (ρ=0.35; p=0.0002). In Burkina Faso, *An. coluzzii* mosquitoes were infected by natural gametocyte carriers. Among mosquitoes that were salivary gland sporozoite positive, 97.2% (36/37) expelled sporozoites with a geometric mean of 420 expelled sporozoites (IQR: 116-2,779) and harbored a geometric mean of 35,149 residual salivary gland sporozoites (IQR: 20,310-164,900). Again, we observed a strong correlation between ruptured oocyst number and salivary gland sporozoite load (ρ=0.84; p<0.0001) and a positive correlation between salivary gland sporozoite load and the number of sporozoites expelled (ρ=0.68; p=0.0003). Whilst sporozoite expelling was regularly observed from mosquitoes with low infection burdens, our findings indicate that mosquito infection burden is associated with the number of expelled sporozoites and may need to be considered in estimations of transmission potential.

Corresponding author: Teun Bousema. Teun.Bousema@radboudumc.nl

Funding: European Research Council.

Keywords: malaria; *P. falciparum*; sporozoites quantification; expelling; oocysts.

Introduction

Malaria transmission to mosquitoes depends on the presence of mature gametocytes in human peripheral blood that are ingested by a mosquito during blood feeding. Ingested parasites undergo several developmental transformations in a process called sporogony. After ingestion gametocytes transform into male and female gametes that fuse to form a zygote. The zygote differentiates into a motile ookinete that penetrates the midgut epithelium to form an oocyst. Multiple rounds of mitotic replication result in the formation of sporozoites inside an oocyst. Upon oocyst rupture, sporozoites are released into the haemocoel and invade the salivary glands [1]. These sporozoites penetrate the distal portion of the two lateral and medial lobes of salivary glands and accumulate extracellularly inside secretory cavities, before entering the salivary ducts [2]. Despite the large number of sporozoites in the cavities, only a small number can pass through the proximal part of the lobes, where the ducts become narrow [3], prior to being inoculated with saliva into the skin when a mosquito takes a blood meal. Developmental bottlenecks during sporogony, as well as the size of the sporozoite inoculum, remain incompletely understood [4].

The density of gametocytes in human peripheral blood is an important determinant of human to mosquito transmission. While gametocyte densities that are below the microscopic threshold for detection can infect mosquitoes [5,6], the likelihood of infecting mosquitoes and the oocyst density that is achieved in infected mosquitoes are low and mosquito infection prevalence and infection burden increase with increasing gametocyte density [7]. Because of the abundance of low-density gametocyte carriers among infected populations, these are considered important drivers of malaria transmission [8-10]. Importantly, this conclusion is based on the assumption that all infected mosquitoes are equally infectious regardless of oocyst densities.

In apparent support of this assumption, a single oocyst can indeed result in thousands of salivary gland sporozoites [11,12] and only tens or low hundreds of sporozoites are assumed to be inoculated per mosquito bite [4], in part due to the physiology of salivary glands [7]. A positive correlation between oocyst densities and salivary gland sporozoites was previously observed in *P. falciparum* [13-15] as well as in *P. vivax* [16], suggesting that on average 1250 sporozoites reach the salivary glands per single ruptured oocyst. The few studies that quantified sporozoite inoculum by allowing mosquitoes to salivating into capillary tubes containing mineral oil, sucrose solution, or blood [17-19], estimated median inocula ranging between 8 to 39 sporozoites with a minority of mosquitoes expelling >100 sporozoites (reviewed in [4]).

While these studies provide some insights into sporozoites expelling and transmission dynamics, they do not reflect natural feeding conditions and the microscopy techniques used for sporozoite quantification may have underestimated the number of sporozoites [20]. Subsequent studies with rodent *Plasmodium* models allowed natural mosquito probing through skin and showed a positive association between sporozoite density in salivary glands and expelled sporozoites [20,21]. A recent study using a *P. yoelii* malaria model showed that mosquitoes with over 10,000 salivary gland sporozoites were 7.5 times more likely to initiate a malaria infection in mice [21]. This finding has not been replicated for human malarias but is broadly consistent with post-hoc analysis of infection likelihood in controlled human malaria infections where only mosquitoes that had >1000 *P. falciparum* salivary gland sporozoites remaining after probing (the residual sporozoite load), were capable of establishing an infection in malaria-naïve volunteers [22].

If low oocyst/low sporozoite densities in mosquitoes are unlikely to initiate infections in humans, this may have profound consequences for our understanding of transmission[23]. If mosquitoes with low infection burdens have limited transmission potential then the rationale for targetting low-density infections in humans that give rise to low infection burdens in mosquitoes [7] may be diminished.

Here, we examined the progression of sporozoite development and the number of sporozoites expelled into artificial skin by individual *Anopheles* mosquitoes infected either with *P. falciparum* gametocyte cultures or natural gametocyte infections. We directly assessed the association between oocyst burden, salivary gland infection intensity and sporozoites expelled.

Results

Low numbers of *P. falciparum* sporozoites are quantifiable by qPCR

Multicopy mitochondrial COX-1 and 18S rRNA gene targets were analysed in octuplicate on serial dilutions of sporozoites to assess qPCR sensitivity in order to select the gene target that achieved highest sensitivity and consistent detection. COX-1 outperformed 18S in detecting sporozoites (**Supplementary Figure 1**), the LOD and LOQ for COX-1 was determined at 20 sporozoites per sample (8/8 sample positivity with a coefficient of variation <2) (**Figure 1A**). Next, we confirmed the qPCR performance in combination with the matrix that was used for expelling experiments by spotting serial dilutions of sporozoites in whole-blood on INTEGRA® dermal substitute artificial skin [24] prior to nucleic acid extraction. The matrix had no apparent impact on sporozoite detectability and quantification (**Supplementary Figure 2**).

Immunolabeling allows quantification of ruptured oocysts

Subsequently, mosquito feeding assays were performed by offering diluted *in vitro* cultured gametocytes to mosquitoes to obtain a broad range of oocyst densities. The association between $\log_{10}$ oocyst intensity and infection prevalence in mosquitoes was assessed using a logistic regression model (using data from 457 mosquitoes, **Figure 1B**). Mosquito infection prevalence was strongly associated with oocyst intensity, corroborating earlier work[25] ,with a strong positive sigmoidal association and a 14.68 (95% CI: 8.18-26.35, p<0.0001) times higher odds of infection prevalence associated with a 10-fold higher oocyst density. In this analysis, oocysts were enumerated microscopically following standard mercurochrome staining. We previously used 3SP2-Alexa 488 anti-circumsporozoite (CSP) immunostaining to visualize ruptured and intact oocysts [26]. The concordance between oocyst prevalence by standard oocyst mercurochrome staining (day 8 post infection) and anti-CSP immunostaining on day 18 post infection (**Figure 1C**) was investigated. For this, oocyst density distributions by both methods were compared within batches of mosquitoes that were fed from the same standard membrane feeding assay (**Figure 1D**). We observed no statistically significant difference in oocyst densities determined by day 8 mercurochrome staining (median 5 oocysts, IQR: 2-20, N=252) and day 18 immunolabelling (median 6 oocysts, IQR: 2-14, N=167; Student t-test on log densities, p=0.944).

Highly infected mosquitoes become salivary gland sporozoite positive earlier

Following assay validation, the extrinsic incubation period (EIP) was compared between mosquitoes with low and high oocyst densities. Batches of high and low infected mosquitoes were generated using standard membrane feeding assay standard concentrations of cultured gametocytes of culture material that was 5- or 10-fold diluted (**Figure 1B**). On day 8 post infection, 20 mosquitoes were dissected and batches that had ≥70% oocyst infection prevalence and means of ≤5 or >20 oocysts were selected for subsequent dissections (**Figure 1B**). On days 9, 10 and 11, salivary glands and remaining mosquito body were collected separately and analysed for sporozoite density by COX-1 qPCR. Mosquitoes were then binned into four categories of sporozoite infection intensity, defined as the sum of mosquito body and salivary gland sporozoite density (**Figure 2A**). This total sporozoite density was examined in relation with the likelihood of being salivary gland sporozoite positive and thus having completed sporogonic development.

On day 9 post-infection, 54.3 % of highly infected mosquitoes (>50,000 sporozoites) were salivary glands sporozoite positive (**Figure 2A**) and had 3.17 (CI 95%: 0.7-14.4, p=0.4278) times odds of being salivary gland positive compared to low infected mosquitoes with <1000 sporozoites (27.3% salivary gland sporozoite positive) (**Supplementary Table S1**). By day 10 post-infection, 82.2 % of mosquitoes with 10,000-50,000 (10k-50k) sporozoites were salivary glands positive and had 11.56 (CI 95%: 1.83, 73.25, p=0.0449) times odds of being salivary glands positive compared to low infected mosquitoes with <1000 sporozoites (28.6 % salivary gland sporozoite positive). On day 11 post-infection, all 15 highly infected mosquitoes (>50,000 sporozoites) were salivary gland sporozoite positive and meaningful odds ratios and 95% CIs could not be determined. When considering the entire period over which EIP experiments were conducted, mosquitoes with > 50,000 sporozoites had a 13.44 times higher odds of being salivary glands positive compared to low infected mosquitoes (<1000 sporozoites; 95% CI: 4.02-44.88, p<0.0001) (**Supplementary Table S2**). Mosquitoes harbouring 10,000-50,000 sporozoites had 5.98 times higher odds of being salivary glands positive when compared to low infected mosquitoes(CI 95%: 1.88-19.07, p=0.0119). These data demonstrate that EIP is shorter in high infected compared to low infected mosquitoes in a temperature and humidity controlled environment.

Sporozoite densities increase with oocyst age

To quantify the number of sporozoites per oocyst, individual oocysts were isolated from midguts stained with 1% mercurochrome on days 9 and 10 post infection [27]. The median sporozoite density was 10,485 (IQR: 9171.3-12,322.5; 12 examined mosquitoes) per oocyst for day 9 and 15,390 (IQR: 10,600-20,887, 19 examined mosquitoes,) for day 10 (**Figure 2B**, p=0.04995, by Welsch two sample t-test on $\log_{10}$-transformed data).

Oocyst density, salivary gland density, and the size of the sporozoite inoculum are positively associated in mosquitoes infected with cultured gametocytes

We performed artificial skin feeding experiments with individual mosquitoes on day 15 post infection to assess sporozoite expelling. To avoid interference of residual blood with oocyst immunolabeling, we assessed mosquito oocyst density (ruptured and intact oocysts) and sporozoite density in the salivary glands on day 18, allowing 3 days for bloodmeal digestion. This approach allowed us to determine the density of intact and ruptured oocysts and associate this to sporozoite density in the same mosquito. It was noted that a minority of oocysts failed to rupture during this time span; 5% (93/1854) of all oocysts were visually intact and 88.3% (166/188) of examined mosquitoes had at least one unruptured oocyst on day 18. While we observed good concordance between oocyst densities by mercurochrome staining on day 8 and immunostaining on day 18 post infection (**Figure 1D**), oocysts sporadically did not take up the 3SP2-Alexa488 anti-CSP antibody labelling (**Supplementary Figure 4**). In 54% (12/22) of mosquitoes without evidence of ruptured oocysts, we observed salivary gland sporozoites. Nevertheless, there was a strong positive association observed between ruptured oocysts and salivary gland sporozoite load (ρ=0.80, p<0.0001; N=185) (**Figure 3B**). When intact oocysts were also included, this association was nearly identical (ρ=0.80, p<0.0001; N=185) (**Supplementary Figure 5A**). We estimated a median of 4951 (IQR: 3016-8318) salivary gland sporozoites per ruptured oocyst in *An. stephensi*. Next, the association between sporozoite density and the number of expelled sporozoites (inoculum size) was determined. For this, individual mosquitoes in miniature cages were allowed to probe for a maximum of 8 minutes on blood-soaked artificial skin (**Supplementary Figure 3**). The skin on the feeder (surface feeding area 201 mm^2) was used as input for DNA extraction with excess skin discarded, after we confirmed that there was no evidence for sporozoite migration outside this area (**Supplementary information S1**).

Among all mosquitoes used in skin feeding experiments, 53% (116/216) expelled sporozoites at any parasite density, and 45% (97/216) expelled sporozoites above our threshold for reliable detection and quantification of 20 sporozoites/skin (**Supplementary Figure 6**). In line with previous work with rodent *P. berghei* [28], sporozoite expelling was observed in mosquitoes that did not take a bloodmeal; 33% (5/15) of mosquitoes that probed but failed to take a blood meal expelled sporozoites (sporozoite range 5-1802). To examine sporozoite expelling in relation to infection burden, mosquitoes were binned into four categories of salivary gland infection intensity that was estimated by combining the residual sporozoite load in the salivary glands and the sporozoites successfully expelled into the skin. In this way also heavily infected mosquitoes that expelled the majority of sporozoites were categorized as heavily infected. We observed a strong positive association between oocyst sheets and total salivary gland sporozoite load (Spearman's correlation coefficient (ρ) = 0.80, p<0.0001; N=111; **Figure 3B**). We observed no statistically significant association between salivary gland infection intensity and the prevalence of expelling sporozoites (**Figure 3A;** p=0.1880). Among mosquitoes that expelled sporozoites, the geometric mean number of expelled sporozoites was 126 (IQR: 30-501) while the highest number of sporozoites detected in skin was 4166. We observed a weak but statistically significant positive association between total sporozoite load and the number of expelled sporozoites (ρ=0.35, p=0.0002; N=112; **Figure 3C**). We observed no evidence for a sharp increase in sporozoite expelling at sporozoite densities ≥10,000, as was previously described in rodent malaria models [21]; 28% (53/186) of our mosquitoes harbored sporozoites below this density. Among these low-infected mosquitoes, 64% (34/53) expelled sporozoites and the median number of expelled sporozoites was 67 (IQR: 13-128).

Natural infected mosquitoes by gametocyte carriers in Burkina Faso show comparable correlation between oocyst density, salivary gland density and sporozoite inoculum

Seven gametocyte donors (age 5-15 years) were recruited in Balonghin, Burkina Faso. Their blood was offered to locally reared *An. coluzzii* via membrane at the gametocyte density observed and following gametocyte enrichment by magnetic activated cell-sorting whereby gametocyte concentration was increased ~3-4-fold (**Supplementary Table S3**). Five donors infected mosquitoes in at least one of these two conditions (**Figure 4A**); as expected mosquito infection

rates and oocyst densities were significantly increased after gametocyte enrichment (**Supplementary Table S4**). Mosquito batches with ≥50% infection prevalence were used for artificial skin feeding experiments described above. From a total of 53 mosquitoes, 31 mosquito midguts were available for immunolabelling of which 67.7% (23/31) had oocyst sheets detected by immunolabelling on day 19; salivary gland sporozoites were detected in 69% (37/53) of mosquitoes on the same day. Two mosquito midguts were negative by immunolabeling while their residual salivary gland sporozoite loads were 7958 and 14,750 respectively, suggesting oocyst staining failure or rupture followed by complete oocyst disappearance. 87.8% (370/421) of all detected oocysts ruptured while 39% (9/23) of oocyst positive mosquitoes harbored at least one intact oocyst (range 1-19). Failure to rupture was uncommon in low infected mosquitoes (≤5 oocysts, N=80) where only 2 midguts had 1 intact oocyst. Among 53 mosquitoes used for artificial skin feeding on day 16 post-infection, 89% (33/37) salivary gland sporozoite positive mosquitoes with a median of 45,100 residual salivary gland sporozoites (IQR: 20,310-164,900) expelled a median of 1035 sporozoites (IQR: 171-2969). We estimated a median of 6350 (IQR: 4225-8475) salivary gland sporozoites per ruptured oocyst in these experiments. Three skin samples were positive by COX-1 qPCR (range 1-64) while salivary glands were negative, suggesting either all sporozoites were expelled or a technical failure in DNA extraction from the salivary glands. During the artificial skin feeding, 30% (16/53) of probing mosquitoes did not ingest blood of which 68% (11/16) expelled sporozoites (range 1-11,970). There appeared to be a trend towards higher prevalence of expelling with increasing sporozoite density (**Figure 4B**). There was a strong association between ruptured oocyst density and total salivary gland sporozoite density (ρ=0.84, p<0.0001) (**Figure 4C**); when intact oocysts were also included the association was very similar (ρ=0.86, p<0.0001; N=30) (**Supplementary Figure 5B**). There was also a strong positive association between total sporozoite load and the number of sporozoites expelled (ρ=0.68, p=0.0003) (**Figure 4D**).

Discussion

We examined *P. falciparum* sporogony in different mosquito species by employing sensitive molecular and staining techniques in conjunction with mosquito probing experiments on artificial skin. Through the visualization of ruptured oocysts and the simultaneous quantification of sporozoites that are expelled by individual mosquitoes, we observed that 95% of oocysts rupture

to release sporozoites and that higher salivary gland sporozoite load is associated with shorter time to infectivity and larger inoculum size.

The proportion infectious mosquitoes is a central component of malariometric indices, both in terms of quantifying the force of infection and the human infectious reservoir. The entomological inoculation rate (EIR) is defined as the number of infectious bites per person per time-unit and is the product of human biting rate and the proportion of sporozoite-positive mosquitoes [29,30]. While EIR is a common measure of human malaria exposure, mosquito infection prevalence is used in this calculation and thus assumes all sporozoite-positive mosquitoes are equally infectious [31]. Similarly, assessments of the human infectious reservoir for malaria typically take the number of oocyst-positive mosquitoes as measure of transmission [10,32] and thereby not only assume that all oocysts will lead to salivary gland sporozoites but also that all oocyst positive mosquitoes have equal transmission potential. Recent work with a rodent malaria model challenged this central assumption[21]; we provide the first direct evidence for *P. falciparum* that sporozoite burden may indeed be a relevant determinant in efficient sporozoite expelling.

In the current study, we assessed sporozoite expelling by mosquitoes carrying low and high infection burdens. Our findings confirm that the vast majority (~95%) of oocysts rupture to release sporozoites. This estimate is higher than a previous study with cultured gametocytes (~72%) [15] that did not provide a second bloodmeal that may accelerate oocyst maturation [33]; a second bloodmeal also better mimics natural feeding habits where multiple bloodmeals are taken within the period required for sporogony. Moreover, we observed a strong positive association between sporozoite salivary gland load and ruptured oocyst density in mosquitoes infected with both cultured and natural gametocytes, with similar median numbers of 4951 (IQR: 3016-8318) and 6350 (IQR: 4225-8475) salivary gland sporozoites per ruptured oocyst in *An. stephensi* and *An. coluzzii* respectively. Previous studies have shown that despite the substantial release of sporozoites per oocyst, only a proportion of sporozoites successfully reach the salivary glands [11,34]. In line with this, we observed considerably higher sporozoite estimates in intact oocysts (10,485-15,390 sporozoites per oocyst). Whilst these sporozoite estimates are higher than commonly reported [4], an early study from the 1960s reported microscopy-detected sporozoite densities above 9000 sporozoites per oocyst [12] and a recent study using qPCR similarly observed up to 12,583 sporozoites from a single oocyst [35]. These high sporozoite numbers per oocyst also mean

that, different from *P. yoellii* [21] and *P. berghei* [22], only a minority of infected mosquitoes had sporozoite densities below the threshold values previously reported in relation to a very low likelihood of achieving secondary infections. Even among infected *An. stephensi* mosquitoes with only 1 ruptured oocyst, 92.9% (26/28) had >1000 and 35.7% (10/28) had >10,000 sporozoites in their salivary glands. In our experiments in natural gametocyte carriers in Burkina Faso, only 2 mosquitoes were observed with single oocysts and both had >10,000 salivary gland sporozoites. Although *P. falciparum* sporozoite densities below these lower thresholds were thus uncommon, we observed that 39% of sporozoite-positive mosquitoes across a wide range of infection densities failed to expel sporozoites upon probing. This finding broadly aligns with an earlier study of Medica and Sinnis that reported that 22% of *P. yoelii* infected mosquitoes failed to expel sporozoites [20]. For highly infected mosquitoes, this inefficient expelling has been related to a decrease of apyrase in the mosquito saliva [36,37].

Importantly, we observed a positive association between salivary gland sporozoite density and the number of expelled sporozoites. For unknown reasons, this association was markedly stronger in experiments where *An. coluzzii* mosquitoes were infected by natural gametocyte carriers compared to *An. stephensi* mosquitoes infected with cultured gametocytes. In our experiments with natural gametocyte carriers, sporozoite density appeared associated with both the likelihood of expelling any sporozoites and the inoculum size. The most heavily infected mosquitoes expelled ~14-fold higher oocyst numbers compared to the lowest quartile. In addition, we found evidence that the extrinsic incubation period (EIP) – the period required for a mosquito to become infectious – is shorter for heavily infected mosquitoes. Previous studies have not found consistent effects of parasite burden on EIP [38,39] but, unlike our experiments, were also not specifically designed to examine this association across a broad range of infection intensities. Our associations of a shorter EIP and larger sporozoite inoculum size make mosquito infection intensity a plausible factor in determining onward transmission potential to humans. Heavily infected mosquitoes may be infectious sooner and be more infectious. On the other side of the infection spectrum, it is conceivable that submicroscopic gametocyte carriers that typically result in low oocyst burdens in mosquitoes give rise to infected mosquitoes with a reduced transmission potential. This would greatly reduce their importance for sustaining malaria transmission and make it less important to identify individuals with low parasite densities for malaria elimination purposes [40]. At the same time, we occasionally observed high sporozoite inocula from mosquitoes with low infection

intensity. This argues against a clear threshold sporozoite or oocyst density below which mosquitoes are truly irrelevant for transmission. Moreover, the high number of sporozoites per oocyst make it plausible that even the low oocyst burdens that are typically observed from asymptomatic parasite carriers (in the range of 1-5 oocysts/infected gut [8,32]) would be sufficient to result in salivary gland sporozoite loads that are sufficient to result in secondary infections.

Our study leaves a number of questions and has several limitations. While the use of two mosquito and gametocyte sources was a relevant strength of our study; an uncertainty relates to the choice of artificial skin that has a realistic 1.33 mm thickness but is arguably less natural than microvascularized skin with all the natural cues for mosquito probing. We initially considered filter paper cards to study expelling [41] (**Supplementary Figure 9**) but this approach was abandoned because of a marked loss in sensitivity due to incomplete DNA recovery (~10-fold signal reduction) [42] and since it is definitely not a natural skin mimic. Whilst genuine skin might have improved natural feeding behaviour, probing and blood-feeding were highly efficient in our model and we see no reasons to assume bias in the comparison between high- and low-infected mosquitoes. In addition, our quantification of sporozoite inoculum size is informative for comparisons between groups of high and low-infected mosquitoes but does not provide conclusive evidence on the likelihood of achieving secondary infections. Given striking differences in sporozoite burden between different *Plasmodium* species – low sporozoite densities appear considerably more common in mosquitoes infected with *P. yoelli* and *P. berghei* [4,22,43] , the association between sporozoite inoculum and the likelihood of achieving secondary infections may be best examined in controlled human infection studies. Since conventional controlled infection studies typically maximize infection burden in mosquitoes [4], dedicated studies that examine a range of mosquito salivary gland burdens may be required.

In conclusion, we observed that the majority of oocysts rupture and contribute to salivary gland infection load. We further observe that this sporozoite load is highly variable and an important determinant of the number of sporozoites that is expelled into the skin upon probing.

Methods

P. falciparum in vitro culture and mosquito infection

Plasmodium falciparum gametocytes, NF54 (West-Africa) and NF135 (Cambodia) [44,45] were cultured in an automated culture system [46] and maintained as previously described at Radboudumc, Nijmegen [47]. *An. stephensi* mosquitoes, Nijmegen Sind-Kasur strain [48], were reared at 30°C and 70-80% humidity with a 12-hour reverse day/night cycle. To have a range of infection intensities in mosquitoes, undiluted and diluted cultured gametocytes (0.3%-0.5% gametocytes) were generated in heparin blood. 100-150, 1-3-days old female mosquitoes were fed using glass membrane mini-feeders [49].

P. falciparum isolates from natural gametocyte carriers and mosquitoes feeding

Gametocyte donors were recruited at schools in Saponé Health District, 45 kilometres southwest of Ouagadougou. Following informed consent, finger prick blood was examined for gametocytes by counting against 500-1000 white blood cells (WBCs) in thick blood films. The gametocyte counts were done by two independent microscopists and expressed as density/µL by assuming 8,000 WBCs per µL of blood (**Supplementary Table S3**). If gametocyte densities were above 16 /µL, 2-5mL of venous blood was drawn by venepuncture in Lithium heparin tubes (BD Vacutainers, ref. 368496) and transported to Centre National de Recherche et de Formation sur le Paludisme (CNRFP) insectary in Ouagadougou in thermos flasks filled with water at 35.5°C [50] (**Supplementary Table S4**).

700 µL of whole blood was used for immediate feeding, that was performed as described elsewhere, using 3 to 5 day-old *Anopheles coluzzii* mosquitoes per glass mini feeder (Coelen Glastechniek, The Netherlands) that was attached to a circulating water bath set up at 38°C (Isotemp®, Fisher Scientific) [51,52]. 50 mosquitoes per cup (starved for 12 hours) were allowed to feed in the dark for 15-20 minutes through a Parafilm membrane. To increase mosquito infection prevalence and intensity, 1mL of blood was used to enrich gametocytes with magnetic cells sorting columns (MACS), as described by Graumans et al. [53] (**Supplementary Figure 10**). In this process, a pre-warmed 23G hypodermic needle (Becton Dickinson, Franklin Lakes, NJ) was attached to an LC column on a QuadroMACS™ separator (MiltenyiBiotech, UK) that was placed inside a temperature-monitored incubator (37°C). The flow-through was collected in a 15ml Falcon tube. Following hydration with 1ml RPMI, 1ml donor blood (Lithium Heparin) was added to MACS

column. The column was rinsed with 2ml warm RPMI. The 15mL Falcon tube was replaced with a new tube, and the needle was removed. Enriched (bound) gametocytes were washed off the column with 4ml of warm RPMI, the plunger was used to press the last 1mL of medium through the column. The two 15ml Falcon tubes, one with the blood mixed to RPMI and the second with the gametocyte suspension were spun down at 2000rpm for 5 minutes in a temperature-controlled centrifuge at 37°C (Eppendorf 5702 R). RPMI supernatant was removed from both tubes with 3ml disposable Pasteur pipettes. To prepare the mosquito blood meal, the small visible pellet of concentrated gametocytes (~30µl), was resuspended with 150µl of prewarmed malaria naïve serum (Sanquin Bloodbank, Nijmegen, The Netherlands) with a blunt needle and 200µl of the patients packed red cells were added and mixed. About 350µl gametocyte enriched blood meal was added to a water-jacket glass feeder as described above.

In both insectaries, at Radboudumc (The Netherlands) and at CNRFP (Burkina Faso), following membrane feeding unfed mosquitoes were immediately removed from cups with an aspirator.

On day 4-6 post infection mosquitoes were given a second blood meal to synchronize the oocyst development. Mosquitoes were kept at 27–29°C in the insectaries on 5-10% glucose and dissected 7-8 days to assess infection prevalence. 20 mosquito midguts were stained with 1% mercurochrome and oocysts were examined and confirmed by two independent microscopists at 400X under an optic microscope (CX 40 Olympus). If oocysts prevalence was above 40%, infected mosquitoes were transferred to the bio-secure insectary in Nijmegen, whilst in Ouagadougou cups with infected mosquitoes were placed into secured metal cages (30x30x30cm) and kept in a temperature and relative humidity controlled environment (27–29°C and 70-80% HR) with double screened doors to prevent escaping sporozoite positive mosquitoes.

Extrinsic incubation period

The extrinsic incubation period (EIP) defined as duration of sporogony was estimated in *An. stephensi* mosquitoes in order to assess i) the best day to perform the expelling experiment, and ii) if EIP differed in low vs high infected mosquitoes. Cages containing 150 mosquitoes were fed with *P. falciparum* NF54 and NF135 gametocytes, and 20 midguts were dissected on day 6-8 post infection (PI) to assess oocyst infection intensity. Groups of low (mean of 5 oocysts) and high (mean of 20 oocysts) infected mosquitoes (prevalence of infection >70%) were maintained in a secure insectary until salivary glands dissections were performed on day 9-10-11. Salivary glands for individual mosquitoes were collected in 1,5mL Eppendorf tubes containing 180µl oocyst lysis

buffer (NaCl 0.1M: EDTA 25mM: TRIS-HCl 10mM), and stored at -20°C for further molecular analysis. Mosquito bodies were first homogenised by beat-beating as previously described [54], in 200μl PBS.

Sporozoite expelling experiments

15 and 16 days post feeding, at Radboudumc and CNRFP respectively, infected mosquitoes were used to quantify the number of expelled sporozoites. To prevent contamination, all instruments and equipment were cleaned from nucleic acids by 30-min exposure to sodium hypochlorite (10% in H2O), rinsed with water and paper dried on the day before the experiment. New gloves were used each time the experiments were performed. Integra dermal substitute (Dermal Regeneration Template, single layer 20x25cm, ref number 68101), hereafter referred to as artificial skin, was cut into 3.5cm squares. Squares were transferred to Petri dishes filled with sterile nuclease-free water (VWR, E476), and left overnight at room temperature (RT). Mosquitoes were individually collected in small Perspex cages (5x5x7cm, covered with netting material on the top and bottom sides) (**Supplementary Figure 3C**). Mosquitoes were starved 14-16 hours prior to feeding. On the day of the expelling experiment, artificial skin was transferred with gloves to an inverted positioned glass membrane mini-feeder (convex bottom, 15mm diameter, ref 70172000) connected to a heated circulating water bath (CORIO C-B5, Julabo) set to 39°C. A rubber band was wrapped around the feeder to secure the artificial skin. Paper tissue was gently pressed on the skin four times to absorb water. 100 μL of naïve donor blood (EDTA, BD Vacutainers, ref. 367525) was pipetted on the circular artificial skin area and spread evenly across the surface with the horizontal side of the tip. The feeder was then turned around and placed on top of the cage, without touching the netting, for a maximum of ten minutes. Following mosquito probing, a scalpel (Dalhausen präzisa plus, no 11) was used to cut the artificial skin above the rubber band around the entire feeder. The artificial skin was transferred with tweezers to a 1,5mL Eppendorf tube containing 180μl oocyst lysis buffer, and stored at -20°C. After feeding, mosquitoes were transferred to metal cages. Mosquitoes were kept at 27-29°C in the insectaries on 5% glucose.

Immunolabelling of intact and ruptured oocysts

After allowing for blood meal digestion that interferes with immunolabelling, mosquito salivary glands and midguts were dissected on day 18 PI for *An. stephensi* (Nijmegen), and day 19 PI for

An. coluzzii (Burkina Faso). Salivary glands were collected in 1,5mL Eppendorf tubes containing 180µl oocyst lysis buffer (NaCl 0.1M: EDTA 25mM: TRIS-HCl 10mM), and stored at -20°C for further molecular analysis. Midguts were dissected in 20 µl phosphate buffered saline (PBS, pH 7.2) without mercurochrome. For experiments performed in Nijmegen, *An. stephensi* midguts were transferred to a fresh drop of (1:400) 3SP2-Alexa488 anti-CSP antibodies and incubated for 30 minutes at RT in a slide humidity incubation box. Following staining, midguts were washed twice with 10µl of PBS for 10 minutes. Midguts were transferred to glass slides and secured with a cover slip. Intact/degenerated and ruptured oocysts were counted using a using an incident light fluorescence microscope GFP filter at 400X. Due to lack of fluorescence microscopes at CNRFP, we combined a formalin fixation method with immunostaining. *An. coluzzii* midguts were transferred into individual screw cap tubes (Eppendorf) filled with 400µl of 4% formalin and stored at 4°C until shipped to the Netherlands. Midguts stored in 4% formalin were collected by using a p1000 Gilson pipette with the point of the tip cut and placed on a slide. Midguts were rinsed from formalin 3 times in PBS 1X-Tween 0.05% by moving the midgut with a needle from drop to drop. They were then transferred to a fresh drop of 3SP2-Alexa488 anti-CSP antibodies (1:400) and incubated for 20 minutes. Midguts were rinsed in PBS and examined as described above (**Supplementary Figure 7**).

Sample extraction and sporozoite quantification by qPCR

Serial dilutions of *P. falciparum* sporozoites were generated to generate standard curves for qPCR. Therefore pooled salivary glands from highly infected mosquitoes were collected in a glass pestle grinder that contained 500µl PBS. The sample was homogenised and subsequently diluted 100 times in PBS. Sporozoites were transferred to a hemocytometer and counted under a phase contrast light microscopy (400x magnification), by two independent microscopists. Serial dilutions were prepared in PBS, using glass test tubes and low binding tips. For each concentration, 100µl was filled out in a 1,5ml Eppendorf tube. Eppendorf tubes were stored at -70°C for at least one day before sample processing. Prior to DNA extraction, 30µl proteinase K (Qiagen, cat no. 19133) was added to Eppendorf tubes containing collected artificial skin and salivary glands samples. PBS was added to all samples to have an equal volume (410µL) for all samples before incubation overnight at 56°C. The following day, total nucleic acids (NA) were extracted with the automated MagNA Pure LC instrument (Roche) using the MagNA Pure LC DNA Isolation Kit – High performance (Roche, product no. 03310515001), and eluted in 50µL. Samples were used immediately or stored

at -20°C. *P. falciparum* sporozoites were quantified by qPCR, targeting the mitochondrial gene COX-1. A previously published primer set [55] was modified to improve template annealing, forward primer 5'-CATCAGGAATGTTATTGCTAACAC-3' and reverse primer 5'-GGATCTCCTGCAAATGTTGGGTC-3', resulting in an amplicon length of 112bp. A probe was designed for amplicon detection 6FAM-ACCGGTTTTAACTGGAGGAGTA-BHQ1. qPCR reactions were prepared with TaqMan Fast Advanced Master Mix (Applied Biosystems, ref 4444557). For each reaction was used 12.5µl mix, 0.4µl primers (stock 50µM, final concentration 800nM), 0.1µl probe (stock 100µM, final concentration 400nM) 7µl PCR grade water and 5µl template DNA. In each run, standard curves and negative controls (water) were included. Melt curves were visually inspected. Samples were run on a Bio-Rad CFX 96 real time System at 95°C for 15 sec, followed by 30 cycles of 95°C for 15 sec, 60°C for 60 seconds.

Statistical analysis

Statistical analyses were performed in R, version 3.1.12 [56]. Associations between $\log_{10}$ oocyst intensity and infection prevalence was modelled using a logistic regression model (using N=457, **Figure 1B**). The difference in mean $\log_{10}$ oocyst densities between staining types were compared using a t-test (t = 0.070075, df = 405, p-value = 0.9442, N=406, **Figure 1D**). The association between total sporozoite density and experiment day with the prevalence of salivary gland sporozoite was modelled using a mixed effects logistic regression with a random intercept for the different experiments (using N=266, **Figure 2A**). Welsch t-test was used to compare sporozoite density between days 9 and 10 using a (t = -2.0467, df = 28.66, p=0.04995, N=31, **Figure 2B**). The association between total sporozoite load and sporozoite expelling prevalence was modelled using a logistic regression (using N=186, **Figure 3A**).

Spearman's correlation coefficient (ρ) was used to assess the association between oocyst sheets and salivary gland sporozoite load (one outlier not included), (ρ=0.80, p<0.0001; N=111, **Figure 3B**); the association between total sporozoite density and the number of sporozoites that was expelled into the artificial skin (ρ=0.35, p=0.0002; N=112, **Figure 3C**) (one outlier not included); the associations between ruptured oocyst density and total sporozoite load, and between total sporozoite load and skin expelling (ρ=0.84, p<0.0001; N=25, **Figure 4C**) and (ρ=0.68, p<0.0003; N=25, **Figure 4D**).

Ethics declarations

The study protocol in Burkina Faso was approved by the London School of Hygiene and Tropical Medicine ethics committee (Review number: 14724), the Centre National de Recherche et de Formation sur le Paludisme institutional review board (Deliberation N° 2018/000,002/MS/SG/CNRFP/CIB) and the Ethics Committee for Health Research in Burkina Faso (Deliberation N° 2018–01-010). Experiments with in vitro cultured parasites and *An. stephensi* mosquitoes at Radboud university medical center were conducted following approval from the Radboud University Experimental Animal Ethical Committee (RUDEC 2009-019, RUDEC 2009-225).

Consent for publication

All authors have given their consent for this publication.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

We thank all the study participants for their willingness to support the study and donate blood. We also thank all the lab staff, drivers and procurement for their dedication in the study. We would like to thank Jacqueline Kuhnen, Laura Pelser-Posthumus, Astrid Pouwelsen, and Jolanda Klaassen for all mosquito breeding.

Additional information

Funding

Funding was provided by the fellowship from the European Research Council to TB (ERC-CoG 864180; QUANTUM) and an AMMODO Science Award (2019) to TB.

Author information

These authors contributed equally: Chiara Andolina, Wouter Graumans

Contributions

Conceptualization and study design were done by C.A., W.G., P.S., K,L. and T.B. Formal analysis, visualization, data curation were done by J.R. and T.B. Methodology was performed by C.A., W.G, K.L., G.J. S.H; Z.S and G.M performed samples collection and sporozoite expelling in Burkina Faso. Manuscript drafting, review and editing was by C.A.,W.G, J.R, K.L. and T.B. Supervision and project administration was done by K.L, T.B, G.G, M.G, A.B.T. Resources, funding acquisition by T.B. All authors reviewed the manuscript.

References

Mueller, A. K., Kohlhepp, F., Hammerschmidt, C. & Michel, K. Invasion of mosquito salivary glands by malaria parasites: prerequisites and defense strategies. *Int J Parasitol* **40**, 1229-1235, doi:10.1016/j.ijpara.2010.05.005 (2010).
Wells, M. B. & Andrew, D. J. Anopheles Salivary Gland Architecture Shapes Plasmodium Sporozoite Availability for Transmission. *mBio* **10**, doi:10.1128/mBio.01238-19 (2019).
Frischknecht, F. *et al.* Imaging movement of malaria parasites during transmission by Anopheles mosquitoes. *Cell Microbiol* **6**, 687-694, doi:10.1111/j.1462-5822.2004.00395.x (2004).
Graumans, W., Jacobs, E., Bousema, T. & Sinnis, P. When Is a Plasmodium-Infected Mosquito an Infectious Mosquito? *Trends Parasitol* **36**, 705-716, doi:10.1016/j.pt.2020.05.011 (2020).
Schneider, P. *et al.* Submicroscopic Plasmodium falciparum gametocyte densities frequently result in mosquito infection. *Am J Trop Med Hyg* **76**, 470-474 (2007).
Bousema, J. T. *et al.* Moderate effect of artemisinin-based combination therapy on transmission of Plasmodium falciparum. *J Infect Dis* **193**, 1151-1159, doi:10.1086/503051 (2006).
Bradley, J. *et al.* Predicting the likelihood and intensity of mosquito infection from sex specific Plasmodium falciparum gametocyte density. *Elife* **7**, doi:10.7554/eLife.34463 (2018).
Andolina, C. *et al.* Sources of persistent malaria transmission in a setting with effective malaria control in eastern Uganda: a longitudinal, observational cohort study. *The Lancet. Infectious diseases* **21**, 1568-1578, doi:10.1016/s1473-3099(21)00072-4 (2021).
Slater, H. C. *et al.* The temporal dynamics and infectiousness of subpatent Plasmodium falciparum infections in relation to parasite density. *Nat Commun* **10**, 1433, doi:10.1038/s41467-019-09441-1 (2019).
Ouedraogo, A. L. *et al.* Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding Assays and Ultrasensitive Malaria Diagnosis in Burkina Faso. *J Infect Dis* **213**, 90-99, doi:10.1093/infdis/jiv370 (2016).
Rosenberg, R. & Rungsiwongse, J. The number of sporozoites produced by individual malaria oocysts. *Am J Trop Med Hyg* **45**, 574-577 (1991).
Pringle, G. A count of the sporozoites in an oöcyst of plasmodium falciparum. *Transactions of The Royal Society of Tropical Medicine and Hygiene* **59**, 289-290, doi:10.1016/0035-9203(65)90009-x (1965).
Vaughan, J. A., Noden, B. H. & Beier, J. C. Population dynamics of Plasmodium falciparum sporogony in laboratory-infected Anopheles gambiae. *J Parasitol* **78**, 716-724 (1992).
Miura, K. *et al.* Strong concordance between percent inhibition in oocyst and sporozoite intensities in a Plasmodium falciparum standard membrane-feeding assay. *Parasit Vectors* **12**, 206, doi:10.1186/s13071-019-3470-3 (2019).
Stone, W. J. *et al.* The relevance and applicability of oocyst prevalence as a read-out for mosquito feeding assays. *Sci Rep* **3**, 3418, doi:10.1038/srep03418 (2013).
Solarte, Y. *et al.* Plasmodium vivax sporozoite production in Anopheles albimanus mosquitoes for vaccine clinical trials. *Am J Trop Med Hyg* **84**, 28-34, doi:10.4269/ajtmh.2011.09-0499 (2011).
Beier, J. C., Davis, J. R., Vaughan, J. A., Noden, B. H. & Beier, M. S. Quantitation of Plasmodium falciparum sporozoites transmitted in vitro by experimentally infected Anopheles gambiae and Anopheles stephensi. *Am J Trop Med Hyg* **44**, 564-570, doi:10.4269/ajtmh.1991.44.564 (1991).
Beier, J. C. *et al.* Quantitation of malaria sporozoites transmitted in vitro during salivation by wild Afrotropical Anopheles. *Med Vet Entomol* **5**, 71-79, doi:10.1111/j.1365-2915.1991.tb00523.x (1991).
Rosenberg, R., Wirtz, R. A., Schneider, I. & Burge, R. An estimation of the number of malaria sporozoites ejected by a feeding mosquito. *Trans R Soc Trop Med Hyg* **84**, 209-212, doi:10.1016/0035-9203(90)90258-g (1990).
Medica, D. L. & Sinnis, P. Quantitative dynamics of Plasmodium yoelii sporozoite transmission by infected anopheline mosquitoes. *Infect Immun* **73**, 4363-4369, doi:10.1128/IAI.73.7.4363-4369.2005 (2005).

Aleshnick, M., Ganusov, V. V., Nasir, G., Yenokyan, G. & Sinnis, P. Experimental determination of the force of malaria infection reveals a non-linear relationship to mosquito sporozoite loads. *PLoS Pathog* **16**, e1008181, doi:10.1371/journal.ppat.1008181 (2020).

Churcher, T. S. *et al.* Probability of Transmission of Malaria from Mosquito to Human Is Regulated by Mosquito Parasite Density in Naive and Vaccinated Hosts. *PLoS Pathog* **13**, e1006108, doi:10.1371/journal.ppat.1006108 (2017).

WHO. *Meeting report of the WHO Evidence Review Group on Low-Density Malaria Infections*, <https://www.who.int/publications/m/item/WHO-HTM-GMP-MPAC-2017.10> (2017).

Agostinis, C. *et al.* Differential Capability of Clinically Employed Dermal Regeneration Scaffolds to Support Vascularization for Tissue Bioengineering. *Biomedicines* **9**, doi:10.3390/biomedicines9101458 (2021).

Churcher, T. S. *et al.* Measuring the blockade of malaria transmission--an analysis of the Standard Membrane Feeding Assay. *Int J Parasitol* **42**, 1037-1044, doi:10.1016/j.ijpara.2012.09.002 (2012).

Stone, W. J. R. *et al.* The relevance and applicability of oocyst prevalence as a read-out for mosquito feeding assays. *Scientific reports* **3**, 3418, doi:10.1038/srep03418 (2013).

Soontarawirat, I. *et al.* Plasmodium vivax genetic diversity and heterozygosity in blood samples and resulting oocysts at the Thai–Myanmar border. *Malaria Journal* **16**, 355, doi:10.1186/s12936-017-2002-x (2017).

Matsuoka, H., Yoshida, S., Hirai, M. & Ishii, A. A rodent malaria, Plasmodium berghei, is experimentally transmitted to mice by merely probing of infective mosquito, Anopheles stephensi. *Parasitol Int* **51**, 17-23, doi:10.1016/s1383-5769(01)00095-2 (2002).

Tusting, L. S., Bousema, T., Smith, D. L. & Drakeley, C. Measuring changes in Plasmodium falciparum transmission: precision, accuracy and costs of metrics. *Adv Parasitol* **84**, 151-208, doi:10.1016/B978-0-12-800099-1.00003-X (2014).

Shaukat, A. M., Breman, J. G. & McKenzie, F. E. Using the entomological inoculation rate to assess the impact of vector control on malaria parasite transmission and elimination. *Malar J* **9**, 122, doi:10.1186/1475-2875-9-122 (2010).

Smith, D. L. *et al.* Ross, macdonald, and a theory for the dynamics and control of mosquito-transmitted pathogens. *PLoS Pathog* **8**, e1002588, doi:10.1371/journal.ppat.1002588 (2012).

Goncalves, B. P. *et al.* Examining the human infectious reservoir for Plasmodium falciparum malaria in areas of differing transmission intensity. *Nat Commun* **8**, 1133, doi:10.1038/s41467-017-01270-4 (2017).

Shaw, W. R. *et al.* Multiple blood feeding in mosquitoes shortens the Plasmodium falciparum incubation period and increases malaria transmission potential. *PLoS Pathog* **16**, e1009131, doi:10.1371/journal.ppat.1009131 (2020).

Hillyer, J. F., Barreau, C. & Vernick, K. D. Efficiency of salivary gland invasion by malaria sporozoites is controlled by rapid sporozoite destruction in the mosquito haemocoel. *Int J Parasitol* **37**, 673-681, doi:10.1016/j.ijpara.2006.12.007 (2007).

Wang, C. Y. T. *et al.* Assessing Plasmodium falciparum transmission in mosquito-feeding assays using quantitative PCR. *Malar J* **17**, 249, doi:10.1186/s12936-018-2382-6 (2018).

Rossignol, P. A., Ribeiro, J. M. & Spielman, A. Increased intradermal probing time in sporozoite-infected mosquitoes. *Am J Trop Med Hyg* **33**, 17-20, doi:10.4269/ajtmh.1984.33.17 (1984).

Thievent, K., Zilio, G., Hauser, G. & Koella, J. C. Malaria load affects the activity of mosquito salivary apyrase. *J Insect Physiol* **116**, 10-16, doi:10.1016/j.jinsphys.2019.04.003 (2019).

Guissou, E. *et al.* Intervention reducing malaria parasite load in vector mosquitoes: No impact on Plasmodium falciparum extrinsic incubation period and the survival of Anopheles gambiae. *PLoS Pathog* **19**, e1011084, doi:10.1371/journal.ppat.1011084 (2023).

Stopard, I. J., Churcher, T. S. & Lambert, B. Estimating the extrinsic incubation period of malaria using a mechanistic model of sporogony. *PLoS Comput Biol* **17**, e1008658, doi:10.1371/journal.pcbi.1008658 (2021).

Slater, H. C. *et al.* Assessing the impact of next-generation rapid diagnostic tests on Plasmodium falciparum malaria elimination strategies. *Nature* **528**, S94-101, doi:10.1038/nature16040 (2015).

Brugman, V. A. *et al.* Detection of malaria sporozoites expelled during mosquito sugar feeding. *Sci Rep* **8**, 7545, doi:10.1038/s41598-018-26010-6 (2018).

Holzschuh, A. & Koepfli, C. Tenfold difference in DNA recovery rate: systematic comparison of whole blood vs. dried blood spot sample collection for malaria molecular surveillance. *Malar J* **21**, 88, doi:10.1186/s12936-022-04122-9 (2022).

Ahmed, A. *et al.* Invasive Malaria Vector Anopheles stephensi Mosquitoes in Sudan, 2016-2018. *Emerging infectious diseases* **27**, 2952-2954, doi:10.3201/eid2711.210040 (2021).

Conway, D. J. *et al.* Origin of Plasmodium falciparum malaria is traced by mitochondrial DNA. *Mol Biochem Parasitol* **111**, 163-171, doi:10.1016/s0166-6851(00)00313-3 (2000).

Teirlinck, A. C. *et al.* NF135.C10: a new Plasmodium falciparum clone for controlled human malaria infections. *J Infect Dis* **207**, 656-660, doi:10.1093/infdis/jis725 (2013).

Ponnudurai, T., Lensen, A. H., Leeuwenberg, A. D. & Meuwissen, J. H. Cultivation of fertile Plasmodium falciparum gametocytes in semi-automated systems. 1. Static cultures. *Trans R Soc Trop Med Hyg* **76**, 812-818, doi:10.1016/0035-9203(82)90116-x (1982).

van de Vegte-Bolmer, M. *et al.* A portfolio of geographically distinct laboratory-adapted Plasmodium falciparum clones with consistent infection rates in Anopheles mosquitoes. *Malar J* **20**, 381, doi:10.1186/s12936-021-03912-x (2021).

Feldmann, A. M. & Ponnudurai, T. Selection of Anopheles stephensi for refractoriness and susceptibility to Plasmodium falciparum. *Med Vet Entomol* **3**, 41-52, doi:10.1111/j.1365-2915.1989.tb00473.x (1989).

Ponnudurai, T. *et al.* Infectivity of cultured Plasmodium falciparum gametocytes to mosquitoes. *Parasitology* **98 Pt 2**, 165-173 (1989).

Soumare, H. M. *et al.* Maintaining Plasmodium falciparum gametocyte infectivity during blood collection and transport for mosquito feeding assays in the field. *Malar J* **20**, 191, doi:10.1186/s12936-021-03725-y (2021).

Ouédraogo, A. L. *et al.* A protocol for membrane feeding assays to determine the infectiousness of P. falciparum naturally infected individuals to Anopheles gambiae. (2013).

Musiime, A. K. *et al.* Is that a real oocyst? Insectary establishment and identification of Plasmodium falciparum oocysts in midguts of Anopheles mosquitoes fed on infected human blood in Tororo, Uganda. *Malar J* **18**, 287, doi:10.1186/s12936-019-2922-8 (2019).

Graumans, W. *et al.* Plasmodium falciparum Gametocyte Enrichment in Peripheral Blood Samples by Magnetic Fractionation: Gametocyte Yields and Possibilities to Reuse Columns. *Am J Trop Med Hyg* **100**, 572-577, doi:10.4269/ajtmh.18-0773 (2019).

Graumans, W. *et al.* Semi-high-throughput detection of Plasmodium falciparum and Plasmodium vivax oocysts in mosquitoes using bead-beating followed by circumsporozoite ELISA and quantitative PCR. *Malar J* **16**, 356, doi:10.1186/s12936-017-2011-9 (2017).

Boissiere, A. *et al.* Application of a qPCR assay in the investigation of susceptibility to malaria infection of the M and S molecular forms of An. gambiae s.s. in Cameroon. *PLoS One* **8**, e54820, doi:10.1371/journal.pone.0054820 (2013).

Team, R. C. *The R Project for Statistical Computing* <https://www.r-project.org/.> (2019).

Figure. 1 The detectability of sporozoites by molecular methods and oocysts by immunolabeling. A. qPCR performance for *P. falciparum* sporozoites**.** Serial dilutions of sporozoites (x-axis) were prepared in PBS and assessed in octuplicate on a single plate to assesses qPCR COX-1 limit of detection and quantification. Dots represent sample cycle threshold (left y-axis), and bars coefficient of variation (right y-axis). For each serial dilution Ct sample positivity is shown as the percentage (%) of total tested. **B.** The relationship between oocyst density versus infection prevalence from 457 membrane feeding experiments using cultured gametocytes. Colours represent regular feeds (red) and those selected for performing experiments on the extrinsic incubation period (EIP; green) or sporozoite expelling experiments (blue). **C**. Immunofluorescence staining with 3SP2-Alexa 488 anti-CSP. Empty sheet and intact oocyst. **D.** Violin plots of oocysts staining - day 8 post infection by mercurochrome (purple) and - day 18 by 3SP2-Alexa 488 anti-CSP immunostaining (yellow) for cultured parasites. Box plots show interquartile range, whiskers show the 95% intervals.

Figure. 2 Extrinsic Incubation Period in high versus low infected mosquitoes. A. Total sporozoites (SPZ) per mosquito in body plus salivary glands (x-axis) were binned by infection load <1k; 1k-10k; 10k-50k; >50k and plotted against the proportion of mosquitoes (%) that were sporozoite positive (y-axis). 120, 120, 40 mosquitoes were dissected on day 9 (blue), 10 (dark green), 11 (light green) respectively. Error bars show the 95% confidence intervals. **B.** Violin plots of sporozoite density (SPZ) in single oocysts dissected on day 9 (purple) and day 10 (green). The box indicates the interquartile range (IQR) (Q1 and Q3 quartiles) and the median. Lines extending Q1 and Q3 quartiles indicate the range of the data within 1.5 times IQR.

Figure 3. Sporozoite expelling in relation to infection burden in *Anopheles stephensi* mosquitoes infected with cultured gametocytes. A. Binning of mosquitoes by total sporozoite load and expelling prevalence (N=186). **B.** The number of ruptured oocysts stained by 3SP2-Alexa 488 anti-CSP and fluorescent microscopy (X-axis) in relation to total salivary gland sporozoite density (Y-axis), assessed by COX-1 qPCR. The red dot indicates a mosquito which had 9 ruptured oocysts but only 126 residual salivary glands sporozoites while expelled 1567 sporozoites. Considering the high number of ruptured oocysts in the midgut it is possible that some lobes of salivary glands were missed during dissection and sporozoite load was underestimated by qPCR. **C.** Total sporozoite density (residual salivary gland sporozoites + sporozoites expelled, X-axis) in relation to the number of expelled sporozoites (Y-axis) by COX-1 qPCR. The dotted line on the x- axis shows the threshold of qPCR detection of 20 sporozoites.

Figure 4. Sporozoite expelling in relation to infection load in *Anopheles coluzzii* mosquitoes infected by naturally acquired gametocyte infections in Burkina Faso. (**A**) Direct feeding (blue) vs magnetic-activated cell sorting (MACS; green). Bars show the infection prevalence for each of the 7 gametocyte carriers. Scatter plots with median lines show the midgut oocyst density as result of direct feeding (blue) and MACS (green). (**B**) Binning of total sporozoite load and expelling prevalence (N=25).(**C**) Scatter plot of absolute numbers of ruptured oocyst (sheet) density assessed by fluorescent microscopy vs total salivary gland sporozoite density assessed by COX-I qPCR (orange). The line represents the fitted linear regression line and the grey shaded area is the 95% CI. (**D**) Scatter plot of absolute numbers of total sporozoite density (residual salivary gland sporozoites + sporozoites expelled) and sporozoites expelled into the artificial skin assessed by COX-I qPCR (purple). The line represents the fitted linear regression line and the grey shaded area is the 95% CI.