DsrMKJOP is the terminal reductase complex in anaerobic sulfate respiration

Ana C. C. Barbosa[‡], Sofia S. Venceslau[‡]*, Inês A. C. Pereira*

Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, 2780-156 Oeiras, Portugal

[‡]These authors contributed equally to this work

*Corresponding authors: ipereira@itqb.unl.pt, sofiasvenceslau@gmail.com

A.C.C.B.: ORCID ID: 0000-0003-3577-2927

S.S.V.: ORCID ID: 0000-0001-8774-1351

I.A.C.P.: ORCID ID: 0000-0003-3283-4520

KEYWORDS

Dissimilatory sulfur metabolism, sulfate reducers, DsrAB/C, DsrMKJOP

The authors have no competing interests.

ABSTRACT

Microbial dissimilatory sulfate reduction is a key process in the Earth biogeochemical sulfur cycle. In spite of its importance to the sulfur and carbon cycles, industrial processes and human health, it is still not clear how reduction of sulfate to sulfide is coupled to energy conservation. A central step in the pathway is the reduction of sulfite by the DsrAB dissimilatory sulfite reductase, which leads to the production of a DsrC-trisulfide. The involvement of the membrane-bound DsrMKJOP complex in the production of sulfide has been inferred from sequence analysis, but its precise function was so far not determined. Here, we present biochemical studies revealing that the DsrMKJOP complex of *Archaeoglobus fulgidus* works as a menadiol:DsrC-trisulfide oxidoreductase. Our results confirm a close interaction between the DsrC-trisulfide and the DsrMKJOP complex and show that electrons flow from menadiol to DsrM hemes *b*, to DsrK noncubane $[4Fe-4S]^{3+/2+}$ catalytic center, and finally to the DsrC-trisulfide with concomitant release of sulfide. These results clarify the role of this widespread respiratory membrane complex and indicate that DsrMKJOP can provide the missing link to energy conservation by generating a *proton motive force* across the membrane in the last step of dissimilatory sulfate reduction.

INTRODUCTION

Sulfur is a key element for life, mainly due to its chemical versatility and ability to switch between several biologically relevant oxidation states, ranging from -2, as in hydrogen sulfide (H_2S), up to +6, as in sulfate (SO_4^{2-}). A critically important microbial process in the biogeochemical sulfur cycle is dissimilatory sulfate reduction (DSR) driven by anaerobic bacteria and archaea, in which the reduction of sulfate to hydrogen sulfide is linked to energy conservation [1, 2]. This process has particular impact in marine sediments, accounting for up to 50% of carbon mineralization in the sea floor [3], and preventing methane emissions through its contribution to anaerobic methane oxidation [4, 5]. Despite being a very ancient bioenergetic pathway [6], it is still not clear how sulfate reducers are able to conserve energy during sulfate respiration. In DSR, a central step is the reduction of sulfite, performed by the dissimilatory sulfite reductase DsrAB and its physiological partner DsrC, which mediate a four-electron reduction of sulfite to zero-valent sulfur in the form of a trisulfide formed between two conserved cysteines of DsrC [7, 8]. The reduction of the DsrC-trisulfide to hydrogen sulfide is thought to derive electrons from the quinone pool via the membrane-bound DsrMKJOP complex, pointing to a critical role of this complex in energy conservation [7, 9, 10]. The Dsr proteins, namely DsrAB, DsrC and DsrMKJOP, are widespread in the microbial world, being also present in phototrophic and chemotrophic sulfur-oxidizing organisms where they are involved in sulfite oxidation [11, 12], and in many organisms capable of sulfur, thiosulfate, sulfite or organosulfonates reduction, and in sulfur disproportionators [13, 14]. However, there is still no concrete evidence demonstrating the physiological function of the DsrMKJOP complex.

The DsrMKJOP complex was first isolated and characterized from *Archaeoglobus fulgidus* [15], then from *Desulfovibrio desulfuricans* [10], and later from *Allochromatium vinosum* [16], where this complex has been shown to be essential for sulfur globule oxidation [17]. This transmembrane complex contains two periplasmic subunits (DsrJ and DsrO), two integral membrane subunits (DsrM and DsrP) and a cytoplasmic subunit (DsrK), and appears to be composed of a combination of two modules, DsrMK and DsrJOP [18].

The MK module is homologous to the membrane-bound HdrED heterodisulfide reductase (Hdr) from

methanogens [19–21], comprising a quinone-interacting membrane diheme cytochrome *b* DsrM, with

both hemes displaying His/His coordination, and an iron-sulfur subunit DsrK predicted to have two

canonical $[4Fe-4S]^{2+/1+}$ clusters, and one noncubane $[4Fe-4S]^{3+/2+}$ cluster [10]. The noncubane cluster is

predicted on the basis of a conserved five-cysteine motif $CX_nCCGX_nCX_2C$ (the CCG motif) [22], which was

identified to bind the catalytic noncubane cluster for heterodisulfide reduction in the homologous HdrD

[15]. The DsrJOP module contains DsrP, an integral membrane subunit with ten predicted transmembrane

helices, and DsrO, an iron-sulfur subunit predicted to coordinate three to four $[4Fe-4S]^{2+/1+}$ clusters, which

together resemble the quinone-interacting NrfCD module widespread in bacterial oxidoreductases [23].

DsrJ is a membrane-anchored triheme cytochrome *c*, where each heme is proposed to have distinct axial

coordination: His/His, His/Met and His/Cys [10, 24]. The His/Cys heme *c* coordination is very unusual but

is also found in SoxXA [25], TdsA [26] and PufC [27], which are involved in thiosulfate oxidation and/or

electron transfer, suggesting that DsrJ could have a catalytic role in sulfur redox chemistry. Notably, the

DsrJOP module is absent in many Gram-positive (Firmicutes) and archaeal sulfate reducers [18], and so

does not seem to be essential. In contrast, the DsrMK proteins are present in all organisms containing

DsrAB and DsrC, often encoded in the same operon and sometimes with multiple copies in the genome,

which suggests this is the minimum functional module of this respiratory complex [18]. Based on its

similarity to HdrED, the DsrMK module was proposed to catalyze the reduction of DsrC-trisulfide to

hydrogen sulfide through the oxidation of the quinone pool [7, 9, 10, 28]. However, the presence of the

DsrJOP module suggests that there may be a more elaborate mechanism [7]. Both DsrM and DsrP are

homologous to quinone-interacting proteins, and the presence of two putative quinone-binding sites led

to the suggestion that a quinone cycle between the DsrMK and DsrJOP modules may be present [9].

The X-ray structure of the soluble HdrABC-MvhAGD complex from the methanogenic archaeon

Methanothermococcus thermolithotrophicus, comprising the MvhAGD hydrogenase and the HdrABC

heterodisulfide reductase, provided the first structural insights into the two noncubane [4Fe-4S]$^{3+/2+}$

clusters present in the active site of HdrB, and how they mediate the reduction of CoM-S-S-CoB

heterodisulfide [29]. Each noncubane cluster is ligated by five cysteines from the predicted CCG motif.

There are two such motifs in both HdrB and HdrD, the catalytic subunits of the two different types of Hdrs

in methanogens. Cleavage of the CoM-S-S-CoB disulfide bond has been suggested to involve the two

noncubane clusters in a homolytic mechanism, along with binding of each substrate to one of the clusters

[29]. Remarkably, only one CCG motif is present in DsrK, raising the question of how this protein may

operate with a single noncubane cluster. Recently, studies of the noncubane [4Fe-4S]$^{3+/2+}$ cluster in the

HdrB protein from *Desulfovibrio vulgaris* Hildenborough revealed that this cluster can adopt "closed" and

"open" conformations corresponding to the substrate-free and substrate-bound forms, respectively [30],

suggesting that cluster flexibility may be important for binding the DsrC-trisulfide in the active site of its

homolog, DsrK.

Overall, unraveling the role of the DsrMKJOP complex and its link to DsrC-trisulfide reduction and

corresponding energy conservation mechanism is a major challenge in understanding dissimilatory sulfur

metabolism. In the present study, we provide evidence for the activity of the DsrMKJOP complex as a

menadiol:DsrC-trisulfide oxidoreductase, clarifying the functional role of this membrane complex in the

final step of the sulfate respiratory chain, where it is likely coupled to energy conservation.

MATERIALS AND METHODS

Purification of *A. fulgidus* DsrMKJOP

A. fulgidus VC-16 (DSM 4304) was grown in sulfate-thiosulfate-lactate medium in 300 L fermenter at 83

°C, as previously described [31], without addition of dithionite. Frozen cells were resuspended in 50 mM

potassium phosphate buffer (KPi) pH 7.0, 10% (v/v) glycerol, homogenized and disrupted in an APV Model

2000 Homogenizer at 80 MPa in the presence of DNase (Sigma-Aldrich) and cOmplete^{TM} protease inhibitor

cocktail (Roche). The cell lysate was centrifuged at 7930 x g for 20 min at 4 °C, followed by

ultracentrifugation at 138000 x g for 2 h at 4 °C. The supernatant (soluble extract) was used to isolate *A.*

fulgidus DsrAB (see below). The pellet (membrane extract) was solubilized with gentle stirring overnight

at 4 °C in 50 mM potassium phosphate buffer pH 7.0, 10% (v/v) glycerol, and 2% (w/v) n-dodecyl β-D-

maltoside (DDM, Glycon Biochemicals GmbH). The solubilized membrane proteins were separated by

ultracentrifugation at 138000 x g for 2 h at 4 °C, and the membrane pellet was used for a second

solubilization for 5 h with gentle stirring at 4 °C, followed by a third ultracentrifugation step under the

same conditions. The pooled solubilized membrane proteins were loaded into a Q-Sepharose high

performance column (2.6 x 10.0 cm, Amersham Pharmacia Biotech) equilibrated with buffer A (50 mM

KPi pH 7.0, 10% (v/v) glycerol, 0.1% (w/v) DDM, and a cOmplete™ protease inhibitor tablet/L). A stepwise

gradient of increasing concentrations of NaCl (from 0 to 1 M NaCl) was applied and the fractions eluted

at 0.45 M and 0.5 M NaCl were pooled, concentrated and the ionic strength of the solution was lowered

by dilution with buffer A and ultrafiltration (50 kDa cutoff, Amicon, Millipore). The concentrate was

applied to a Resource Q column (1.6 x 3.0 cm, Amersham Pharmacia Biotech) equilibrated with buffer A.

A stepwise gradient of increasing concentrations of NaCl (from 0 to 1 M NaCl) was performed and the

fractions eluted at 0.3 M and 0.35 M NaCl were pooled, concentrated and the ionic strength lowered by

dilution and concentration steps as above. All the chromatographic steps were performed at 4 °C, at air,

and monitored by UV-visible spectroscopy. The purity of DsrMKJOP was analyzed by a 10% Tricine-SDS-

PAGE gel and a Blue Native (BN)-PAGE gel, followed by Coomassie Blue staining and heme-staining [32],

and the protein concentration was determined using the previously determined absorption coefficient (ε)

of 132.4 mM^{-1}.cm^{-1} at 555 nm in the reduced state [33].

UV-visible Spectroscopy

UV–visible spectra were acquired in a Shimadzu UV-1800 spectrophotometer inside a Coy anaerobic

chamber (98 % N_2, 2 % H_2). Spectra were obtained at 60 °C, using as-purified DsrMKJOP, prepared in 20

mM Tris-HCl pH 7.5, then incubated with menadiol and an excess of sodium dithionite to allow full

reduction. Menadiol was prepared by reduction of menadione with sodium dithionite [34], stored in dry

state at -20 °C under anaerobic conditions, and solubilized in ethanol prior to use. A control spectrum of

the as-purified DsrMKJOP complex was obtained by addition of the same volume of ethanol as for

menadiol incubation.

Electron Paramagnetic Resonance Spectroscopy

X-band electron paramagnetic resonance (EPR) spectra were obtained on a Bruker EMX spectrometer

(Billerica, MA) equipped with an Oxford Instruments ESR-900 continuous flow helium cryostat (Abingdon,

UK). A menadiol-treated sample of the DsrMKJOP complex, in 20 mM Tris-HCl pH 7.5 at a concentration

of 25 µM, was prepared inside a Coy anaerobic chamber (98 % N_2, 2 % H_2) by incubation for 2h at 60 °C

with a solution of menadiol in ethanol in a 1:500 ratio. As control an as-purified DsrMKJOP sample was

also incubated for 2 h with the same volume of ethanol as was used in the menadiol experiment. Samples

were transferred to EPR tubes within the anaerobic chamber, capped, and immediately frozen in liquid

nitrogen upon removal from the chamber. All spectra were recorded using a microwave frequency of 9.38

GHz, a field modulation frequency of 100 kHz, a microwave power of 20.1 mW, a modulation amplitude

of 1.0 mT, and a temperature of 23 K.

Purification of *A. fulgidus* DsrAB

The purification protocol of the soluble extract from *A. fulgidus* cells followed two consecutive ion

exchange chromatography steps at 4 °C under air, as described in detail in Santos *et al.* [7]. The purity of

DsrAB was analyzed in a 10% Tricine-SDS-PAGE gel, and the protein concentration was determined by UV-

visible spectroscopy using an ε of 60 $mM^{-1}.cm^{-1}$ at 593 nm determined previously [35]. The purified DsrAB

was stored under anaerobic conditions to prevent prolonged contact with O_2, which leads to gradual

activity loss.

Purification of Recombinant *A. fulgidus* DsrC

A. fulgidus DsrC with the two structural Cys (Cys[77] and Cys[85]) modified to Ala was expressed in *Escherichia*

coli and purified by an affinity exchange chromatography step on a HiTrap[TM] immobilized metal-ion affinity

chromatography (IMAC) high performance column (GE Healthcare), as described in detail in Ferreira *et al.*

[36]. The purity of DsrC was analyzed on a 10% Tricine-SDS-PAGE gel, and the protein concentration was

determined by bicinchoninic acid (BCA) assay (Merck Millipore) and by UV-visible spectroscopy using an ε

of 24 $mM^{-1} \cdot cm^{-1}$ at 280 nm. Before its use in activity assays, DsrC was treated with 5 mM dithiothreitol

(DTT, Sigma-Aldrich) during 30 min at 37 °C, to ensure its Cys were in the reduced state, and excess

reductant was removed with a HiTrap[TM] Desalting column (GE Healthcare) [7]. Reduced DsrC was

concentrated, if necessary, by ultrafiltration (10 kDa cutoff, Amicon Millipore) and kept under anaerobic

conditions until use. The redox state of the DsrC (5 μg) was analyzed by a gel-shift assay using MalPEG

(methoxy-polyethylene glycol maleimide, Fluka), as described in Venceslau *et al.* [37].

Production and Analysis of DsrC-trisulfide

The enzymatic production of DsrC-trisulfide was performed inside a Coy anaerobic chamber (98% N_2, 2%

H_2) at 60 °C using Zn-reduced methyl viologen (Sigma-Aldrich) as the electron donor [7]. The activity assays

were run in 50 mM KPi pH 7.0, 1 mM Zn-reduced methyl viologen (MV^+), 200 nM of DsrAB, 0.5 mM sodium

sulfite (Sigma-Aldrich), with a range from 10 μM to 100 μM of reduced DsrC. After a 5 min incubation of

DsrAB in the reaction buffer, followed by a 1 min incubation with DsrC, the reaction was started by

addition of sodium sulfite. The oxidation of MV^+ was monitored at 732 nm (ε = 3.15 $mM^{-1} \cdot cm^{-1}$) in a

spectrophotometer. A prolonged incubation time (from 15 to 60 min depending on the DsrC-trisulfide

concentration) was used to try to drive the enzymatic reaction to completion and have DsrC as much as

possible in the trisulfide form. The enzymatic solution was diluted and concentrated by ultrafiltration (10

kDa cutoff, Amicon Millipore) to remove the methyl viologen. The redox state of DsrC was monitored with

the MalPEG gel-shift assay as previously described [37].

Determination of DsrC Thiol Content by DTNB

DsrC in different redox states was subject to a DTNB (5,5-dithio-bis-(2-nitrobenzoic acid) – Ellman's

reagent) assay to quantify the sulfhydryl groups present. DsrC (0.8 – 3 nmol) was incubated in 100 mM

KPi pH 8.0, 1 mM EDTA, and 0.2 mM DTNB (Sigma-Aldrich) for 15 min at room temperature. The

production of 2-nitro-5-thiobenzoate anion (TNB^{2-}) was quantified by measuring the absorbance at 412

nm using the ε of 14150 $M^{-1} \cdot cm^{-1}$ [38].

Surface Plasmon Resonance Analysis

The surface plasmon resonance (SPR) experiments were performed at 25 °C on a BIAcore 2000 instrument

(GE Healthcare). DsrC-trisulfide was immobilized on a NTA sensor chip (GE Healthcare) by the His-tag tail

present at the N terminus. All the assays were performed with 10 mM HEPES pH 7.4, 150 mM NaCl, 50

µM EDTA, and 0.005% (v/v) Tween 20 running buffer. DsrC-trisulfide (45 nM) was bound to a previously

activated flow cell at a flow rate of 5 µL/min in the presence of 500 µM $NiCl_2$ in running buffer in order to

have an immobilization of around 350 resonance units (RU). Another activated flow cell was similarly

treated with running buffer in the absence of DsrC-trisulfide (control cell). Interaction experiments with

DsrMKJOP complex were performed with a 3 min injection of increasing concentrations of DsrMKJOP

complex (range from 8.9 nM to 568 nM) at a flow rate of 30 µL/min. At the end of sample injection, the

NTA sensor chip was regenerated with 350 mM EDTA in running buffer for 1 min at a flow rate of 10

µL/min before a new cycle of surface activation and immobilization. Sensorgrams were obtained by

subtracting the unspecific binding to the control flow cell in order to remove buffer artifacts and

normalizing to the baseline injection timepoint. The equilibrium dissociation constant (K_D) was

determined from duplicate experiments and according to $RU_{steady\text{-}state} = (RU_{max} \times [analyte])/K_D + [analyte])$.

DsrMKJOP Activity Assays

Menadiol:DsrC-trisulfide oxidoreductase activity assays by DsrMKJOP were performed inside a Coy

anaerobic chamber (98% N_2, 2% H_2) at 60 °C using menadiol as electron donor. The activity assays were

run in 50 mM KPi pH 7.0, 0.5 mM menadiol, and 100 nM DsrMKJOP in the presence of DsrC-trisulfide (10

µM). The reaction was started by addition of DsrC-trisulfide after a 5 min incubation of DsrMKJOP in the

reaction buffer. The oxidation of menadiol was monitored at 270 nm (using $\varepsilon_{DMN} = 16$ mM^{-1}·cm^{-1} [39]) in

a spectrophotometer. For peptide mass fingerprinting of DsrC, the enzymatic reactions were

concentrated and dialyzed by ultrafiltration (10 kDa cutoff, Amicon Millipore). To monitor the redox state

of DsrC cysteines, the MalPEG gel-shift assay was used [37].

Peptide Mass Fingerprinting

The redox state of DsrC cysteines was analyzed by peptide mass fingerprinting using a matrix-assisted

laser desorption/ionization-time-of-flight (MALDI-TOF/TOF) mass spectrometer, as described in Santos *et*

al. [7]. Alkylation was performed inside an anaerobic chamber by incubation of DsrC with 50 mM

iodoacetamide (IA, Sigma-Aldrich) in 50 mM ammonium bicarbonate pH 7.8 during 30 min at 55 °C and

run on a 10% Tricine-SDS-PAGE gel (15 µg) under non-reducing conditions. The protein bands were

excised, destained and digested with Asp-N (Roche, 20 ng/µL) overnight at 37 °C. This endoproteinase

generates a peptide comprising both conserved C-terminal cysteine residues, [101]DA**C**RIAGLPKPTG**C**V[115],

with predicted mass of 1500.8 Daltons (Da). The digested peptides were desalted and concentrated using

POROS C18 (Empore, 3M) and eluted directly onto the MALDI plate using 1 µL of 5 mg/mL alpha-cyano-4-

hydroxycinnamic acid (CHCA, Sigma-Aldrich) in 50% (v/v) acetonitrile and 5% (v/v) formic acid.

The data were acquired in positive reflector MS mode using a 5800 MALDI-TOF/TOF (AB Sciex) mass

spectrometer and TOF/TOF Series Explorer Software v.4.1.0 (Applied Biosystems). External calibration was

performed using CalMix5 (Protea). Mass Spectrometry (MS) data were analyzed using Data Explorer

Software v.4.11 (AB Sciex) and obtained at the Mass Spectrometry Unit (UniMS), ITQB/iBET, Oeiras,

Portugal.

Determination of Hydrogen Sulfide by HPLC

Hydrogen sulfide was quantified using monobromobimane (mBBr) derivatization [40]. mBBr (Thermo

Fisher Scientific) was diluted in HPLC ultrapure acetonitrile to a final concentration of 50 mM. For

derivatization, 10 µL of sample from the enzymatic reaction were reacted with 3 µL of mBBr in 50 mM

EPPS   (4-(2-hydroxyethyl)piperazine-1-propanesulfonic   acid)   pH   8.0   and   5   mM   DTPA

(diethylenetriaminepentaacetic acid) in a final volume of 86 µL, inside an anaerobic chamber. The

reactions were incubated in the dark at room temperature for 10 min, after which 4 µL of 5 M

methanesulfonic acid was added to stop the reaction.

Derivatized hydrogen sulfide (sulfide dibimane) was measured by high performance liquid

chromatography (HPLC, Waters Alliance 2695) with a fluorescence detector (Waters 2475) with an

excitation/emission wavelength of 380 and 480 nm, respectively, using a reverse phase Ultrasphere® ODS

column (4.6 x 250 mm, 5 µm, Hichrom) run at a flow rate of 1.2 mL·min^{-1} and 35 °C, as described in detail

in Santos *et al.* [7]. The retention time was 15.5 min for sulfide dibimane. A calibration curve was

performed with freshly prepared solutions of sodium sulfide nonahydrate (Sigma-Aldrich) in 50 mM KPi

pH 7.0, reacted with excess mBBr.

RESULTS AND DISCUSSION

DsrMKJOP isolation and reduction with Menadiol

The DsrMKJOP transmembrane complex was purified with DDM as solubilizing detergent from

membranes of the thermophilic archaeon *A. fulgidus,* by adapting the protocol previously reported by

Mander *et al.* [15]. The DsrMKJOP complex is the only heme-containing membrane complex in *A. fulgidus*

[18], which facilitates its purification compared with other sulfate reducers, such as *D. desulfuricans* [10].

In *A. fulgidus*, the predicted molecular mass for each subunit of the complex is 38.4 kDa (DsrM, locus tag:

AF0501), 64.4 kDa (DsrK, AF0502), 16.7 kDa (DsrJ, AF0503), 30.5 kDa (DsrO, AF0499), and 44.2 kDa (DsrP,

AF0500). The purified complex displayed four bands on a Tricine-SDS-PAGE gel (**Figure S1**) with apparent

molecular masses of 61, 31, 28, and 17 kDa, in line with the report by Mander *et al.* [15]. The 17 kDa band

was identified as a cytochrome *c* (DsrJ) by heme-staining, and the other bands were assigned as DsrK (61

kDa), DsrM (31 kDa), DsrO (28 kDa), based on previous identifications [10, 15]. As reported before, DsrP

is not detected by SDS-PAGE [10, 15] because it is a highly hydrophobic subunit (ten predicted

transmembrane helices), which hinders its migration in the gel.

The UV-visible spectrum of the as-purified DsrMKJOP (oxidized state) is dominated by absorption maxima

at 280, 410 and 527 nm (**Figure 1A**), confirming the presence of hemes. Upon reduction, the Soret peak

shifts to 420 nm, and the β- and α-bands are observed at 524 and 557 nm, respectively, which is in line

with previously reported spectra of DsrMKJOP from *A. fulgidus* [15] and *D. desulfuricans* [10]. The α-band

is quite broad supporting the presence of hemes *b* and *c*. Addition of menadiol reduced 38% of the total

hemes as compared with full reduction by dithionite. This corresponds to approximately two hemes

reduced, and since hemes *b* typically have higher redox potentials than hemes *c,* this likely corresponds

to the two predicted hemes *b* in the quinone-interacting DsrM membrane subunit. A similar result was

reported by Pires *et al.* [10], where 40% of hemes belonging to the *D. desulfuricans* DsrMKJOP were also

reduced upon incubation with menadiol.

The electron paramagnetic resonance (EPR) spectrum of the as-purified *A. fulgidus* DsrMKJOP shows an

intense rhombic signal characteristic of a noncubane $[4Fe-4S]^{3+/2+}$ cluster in the oxidized state, with *g*-

values of 2.025, 1.995 and 1.947 (**Figure 1B**). This EPR signature was first observed in heterodisulfide

reductases from methanogens (upon addition of HS-CoM or HS-CoB), with *g*-values of 2.013, 1.991 and

1.938 for HdrABC from *Methanothermobacter marburgensis* and 2.012, 1.993 and 1.946 for HdrED from

Methanosarcina barkeri [41, 42], both of which possess two noncubane $[4Fe-4S]^{3+/2+}$ catalytic centers

involved in mediating heterodisulfide (CoM-S-S-CoB) reduction in two one-electron steps [21, 43]. The

previously reported *g*-values of *A. fulgidus* DsrMKJOP were 2.031, 1.994 and 1.941 [15], while in *D.*

desulfuricans DsrMKJOP the *g*-values were 2.027, 1.991 and 1.943 [10]. Given the similarity of DsrK to

HdrD, the noncubane $[4Fe-4S]^{3+/2+}$ cluster present in DsrK is likely involved in thiol/disulfide chemistry,

suggesting that DsrK is the catalytic subunit of the DsrMKJOP complex. Upon incubation with menadiol

under anaerobic conditions, the intensity of the rhombic signal decreased by 85%, revealing a pathway

for electron transfer from the hemes *b* of DsrM to the noncubane $[4Fe-4S]^{3+/2+}$ cluster of DsrK. The redox

potential of the noncubane center in *A. fulgidus* DsrK was previously determined to be +90 mV [15], which

agrees with its reduction by menadiol (midpoint potential of approx. -70 mV).

The DsrC-trisulfide interacts strongly with DsrMKJOP

In *A. vinosum*, coelution upon protein purification suggested that DsrK interacts with DsrC [16]. Following

our identification of the DsrC-trisulfide as the product of sulfite reduction by DsrAB, we proposed that this

is the actual substrate for the DsrMKJOP complex [7]. To assess if there is a protein-protein interaction

between the DsrC-trisulfide and the DsrMKJOP complex, surface plasmon resonance (SPR) was initially

used. SPR is a highly sensitive technique for studying protein-protein interactions, without the need for

labeling. The *A. fulgidus* DsrC-trisulfide was produced by *in vitro* enzymatic reaction using DsrAB from the

same organism, as described in Santos *et al.* [7]. The DsrC-trisulfide thus produced was immobilized on a

nitrilotriacetic acid (NTA) sensor chip by the N-terminal His-tag, and the interaction was studied with

increasing concentrations of DsrMKJOP complex (**Figure 2**). A strong interaction was observed between

the two proteins, which could be detected even at low concentrations of the membrane complex. Steady-

state conditions could not be achieved (even by increasing the association time), which means that no

kinetic model could fit the experimental data, and so the association and dissociation rate constants could

not be obtained. Nevertheless, an estimate for the equilibrium dissociation constant (K_D) of 200 ± 27 nM

was derived from duplicate sensorgrams using the maximum response reached. This K_D confirms a strong

interaction between the DsrC-trisulfide and the DsrMKJOP complex (strong interactions are in the

nanomolar range, or even lower as in the case of high-affinity antibodies), supporting the DsrC-trisulfide

as a possible substrate for the membrane-bound DsrMKJOP complex. The SPR data were obtained at 25

°C, whereas a higher temperature (as used in the activity assays) should favor the association and

dissociation rates of DsrC-trisulfide to DsrMKJOP complex since *A. fulgidus* is a thermophile, suggesting

that the actual K_D at physiological temperatures may be somewhat different, and possibly even lower.

The DsrC-trisulfide is reduced by DsrMKJOP

Given that menadiol can reduce the noncubane [4Fe-4S]$^{3+/2+}$ catalytic center of DsrK, and that the DsrC-

trisulfide interacts with DsrMKJOP complex, we tested whether the DsrC-trisulfide can serve as substrate

and electron acceptor for the menadiol-reduced DsrMKJOP complex. Menadiol:DsrC-trisulfide

oxidoreductase activity by the DsrMKJOP complex was thus measured. These experiments confirmed a

fast oxidation of menadiol (measured by menadione formation at 270 nm) in the presence of the DsrC-

trisulfide with a specific activity of 85.5 ± 5.4 mU/mg, assuming DsrMKJOP as a 5-subunit monomer (200

kDa) (**Figure 3A**). In control reactions without DsrMKJOP, menadiol or DsrC-trisulfide no activity was

detected. In addition, sequential additions of DsrC-trisulfide, after the reaction has stopped, trigger a new

phase of menadiol oxidation confirming it had stopped due to substrate depletion (**Figure 3B**).

The redox state of DsrC, before and after the reaction, was analyzed by a gel-shift assay using MalPEG,

which selectively labels cysteine thiol groups [7, 37]. In this assay, the DsrC-trisulfide presents no shift in

the gel as it cannot bind MalPEG, whereas the reduced form binds one or two MalPEG molecules, as the

last Cys_A is more accessible for labeling than Cys_B (Cys^{114} and Cys^{103} in *A. fulgidus*, respectively) [7, 37]. At

time zero (T_0), most DsrC shows no shift, as expected for the trisulfide, but a small amount of reduced

species is present, as revealed by labeling for one MalPEG, due to incomplete production of DsrC-trisulfide

by DsrAB (**Figure 3C**). After the enzymatic reaction (T_f), most DsrC is now in the reduced form, as revealed

by the shift for one (Cys_A) and two labeled cysteines (Cys_A and Cys_B) (**Figure 3C**). In the control experiments,

without DsrMKJOP or menadiol, DsrC remains in the oxidized DsrC-trisulfide state. Using the MalPEG assay

to follow the reaction in time, the decrease in the DsrC-trisulfide band and increase in the band for

reduced, doubly labeled DsrC, is clearly seen (**Figure 3D**). Increasing concentrations of menadiol for the

same reaction time also increase the extent of reaction (**Figure 3E**). As a control, reduced DsrC was tested

as a substrate for DsrMKJOP under the same conditions. In this case, no activity is observed (**Figure S2A**),

as expected, and the MalPEG assay confirms the reduced state of DsrC in the beginning and end of the

experiment (**Figure S2B).**

Finally, the redox state of DsrC cysteines was also characterized by peptide mass fingerprinting after

protein digestion with Asp-N endoproteinase, which gives rise to a small peptide containing both Cys_A and

Cys_B, followed by alkylation with iodoacetamide (IA), which adds 57 Da per reduced cysteine [7]. In

reduced DsrC, the C-terminal peptide ^{101}DA**C**RIAGLPKPTG**C**V^{115} has a mass of 1498.7 Da [7], and upon

alkylation, a peptide with 1614.9 Da is detected (**Figure 4A**). In the DsrC-trisulfide formed by reaction with

DsrAB, the C-terminal peptide shows a mass increase of 32 Da, 1530.7 Da, which remains unaltered by

alkylation with IA (**Figure 4B**). A small amount of unreacted IA-labelled DsrC is still detected in this sample

(1614.8 Da), as well as a species that does not react with IA (1498.8) and likely corresponds to DsrC with

a disulfide bond between the two Cys. It should be noted that the peak intensities are not directly

correlated with relative amounts, as they depend on the different ionization properties of each peptide.

As expected, analysis of the DsrC peptide after the reaction with DsrMKJOP, reveals the disappearance of

the DsrC-trisulfide peak at 1530.7 Da, and the increase of the 1614.8 Da peak characteristic of reduced

DsrC, where two cysteines react with IA (**Figure 4C**).

Sulfide is the product of DsrMKJOP catalysis

The products of DsrC-trisulfide reduction by the DsrMKJOP complex should be sulfide and reduced DsrC

(**Figure 5A**). To quantify the sulfide released during DsrMKJOP catalysis, the reaction was performed with

increasing concentrations of DsrC-trisulfide. These kinetic studies showed an increase in the length of the

reaction with increasing substrate concentrations (**Figure 5B**), but not an increase in reaction rate. This

indicates that the concentrations used are already saturating and that the Michaelis-Menten constant

(K_M) for the DsrC-trisulfide is probably in the nM range, in agreement with the measured K_D. The reaction

rates slow down as the substrate becomes exhausted. The product sulfide, quantified by HPLC as the

monobromobimane fluorescent derivative, increases linearly with increasing DsrC-trisulfide

concentrations (**Figure 5C**), although in sub-stoichiometric amounts (40 to 50%). This can be explained by

two factors: the partial escape of hydrogen sulfide to the gas phase at pH 7.0, made worse by the high

temperature required for the DsrMKJOP assay, and also by a degree of uncertainty associated with

quantification of the DsrC-trisulfide. This quantification was performed indirectly using the BCA (for

protein quantification) and DTNB (for thiol quantification) assays, by the amount of reduced DsrC

remaining in samples of the DsrC-trisulfide. However, we observed by mass spectrometry (MS) that upon

production of the DsrC-trisulfide some part of DsrC is in the disulfide state and is not alkylated by IA (peak

1498.8 Da in **Figure 4B and 4C**) and thus is also not quantified by DTNB derivatization, generating an

uncertainty in the actual level of DsrC-trisulfide present in the samples. Nevertheless, taken globally these

results show clearly that DsrMKJOP can catalyze full reduction of the DsrC-trisulfide to sulfide using

menadiol, finally clarifying the functional role of this respiratory complex in dissimilatory sulfite reduction.

Insights into the catalytic mechanism of DsrMKJOP

A catalytic mechanism for heterodisulfide reduction by Hdrs has been proposed based on the structure of

HdrABC with substrate bound, where the CoM-S-S-CoB heterodisulfide binds between the two noncubane

$[4Fe-4S]^{3+/2+}$ clusters and is homolytically cleaved [29]. The thyil radicals formed may be stabilized by

coordination with the reduced $[4Fe-4S]^{2+}$ clusters, leading to the binding of the CoM and CoB sulfurs to

the two $[4Fe-4S]^{3+}$ clusters, resulting in the "open" conformation of the clusters [30]. Two consecutive

single-electron reduction steps release CoB-SH and CoM-SH [29, 30]. However, the catalytic mechanism

of the DsrMKJOP complex is likely different since it coordinates a single noncubane $[4Fe-4S]^{3+/2+}$ cluster in

DsrK instead of the two found in HdrD and HdrB (**Figure S3**). Interestingly, in many DsrK proteins from

different sulfate reducers the last cysteine of the CCG motif for binding this cluster is replaced by a

conserved aspartate (**Figure S4**). In *A. fulgidus*, the DsrK protein of the DsrMKJOP complex (AF0502) and

another homolog (AF0543) have this aspartate, whereas two other DsrK homologs (AF0544 and AF0547),

encoded in *dsrMK* arrangements, have a cysteine, suggesting that both residues may be functional. In

addition, one of the conserved cysteines from the missing noncubane cluster (Cys[329] in *A. fulgidus*

numbering) is strictly conserved in DsrK proteins and is likely involved in the catalytic mechanism of DsrC-

trisulfide reduction. This cysteine is part of a conserved GECGH sequence motif that is highly conserved in

DsrK from several organisms (**Figure S4**) and is one of the cysteines from the second CCG signature motif

in HdrD and HdrB (**Figure S3**). A similar motif (ECGLH), present in APS and PAPS reductases, includes a

redox-active catalytic cysteine [44]. Other proteins from sulfate reducers, homologous to DsrK and HdrD,

such as HmcF from the HmcABCDEF complex [18, 23, 45] and TmcB from the TmcABCD complex [46] also

contain only one CCG motif and this conserved cysteine in a T/GECGH motif (**Figure S3**), suggesting that

this cysteine is probably essential for disulfide reduction catalysis. The Hmc and Tmc complexes are both

transmembrane complexes with a similar architecture to the DsrMKJOP complex, but they are not present

in *A. fulgidus* [18]. Interestingly, the characteristic EPR signal of the noncubane $[4Fe-4S]^{3+}$ cluster was

initially reported for the methanogenic HdrB and HdrD only in the presence of HS-CoM or HS-CoB [41–

43], which bind to the cluster, whereas for DsrK and TmcB proteins this signal is observed in the as-isolated

protein, in the absence of any added substrate [10, 15, 46]. This could suggest that this conserved cysteine

might be binding the cluster and generating the open $S = ½$ conformation. However, for the recombinant

HdrB protein produced in *E. coli* the signal for the noncubane cluster was visible already in the absence of

substrate [22].

In methanogens, the redox potential of the CoM-S-S-CoB heterodisulfide has been determined to be $E^{0'}$=-

143 ± 10 mV vs. SHE [47], which is more positive than that of typical disulfides such as glutathione and

cysteine. It is also more positive than the potential of methanophenazine ($E^{0'}$=-165 ± 6 mV), the

membrane-associated quinone-like redox carrier that is the electron donor to HdrED [47]. The redox

potential of the DsrC-trisulfide is unknown, so it is not clear whether its reduction by menaquinol ($E^{0'}$ ~ -

70 mV) is thermodynamically favorable. Protein disulfide bonds are known to span a wide range of

potentials. For example, *A. fulgidus* proteins from the thioredoxin superfamily have potentials ranging

from -32 mV to -309 mV (pH 7.0) [48]. A value of -346 mV (pH 7.0, vs. SHE) has been reported for several

organic water-soluble trisulfides [49], but we know of no reports for redox potentials of protein-bound

trisulfides, so it is very difficult to estimate the redox potential of the DsrC trisulfide. We tested the

chemical reduction of this species by different chemical reducing agents, and surprisingly none of them,

not even those with very low redox potential, could reduce the DsrC-trisulfide as analyzed with the

MalPEG shift assay (**Figure S5**). Moreover, MS confirmed the intact state of the DsrC-trisulfide after

reduction with dithionite, TCEP and DTT (**Table S1**). This unexpected result suggests that the DsrC-

trisulfide can only be reduced by the DsrMKJOP complex. The fact that it cannot be chemically reduced

could suggest that it has a low redox potential, which could increase upon interaction with the DsrMKJOP

complex, but it is also possible that the reduction of the trisulfide is not thermodynamically, but kinetically

prevented, i.e. that the reaction has a high activation energy and occurs only when it is enzymatically

catalyzed. This sort of argument has been put forward by Flohé and coworkers in discussing the

glutathione redox potential and its physiological significance [50, 51]. Such kinetic impediment for

reduction of the DsrC-trisulfide, which may be based on steric inaccessibility of the trisulfide, makes sense

considering that this species is likely to face a considerable intracellular sulfide concentration.

Nevertheless, the full reduction of the DsrC-trisulfide, as observed by MS, and the increasing production

of sulfide upon reaction of DsrMKJOP and menadiol with increasing concentrations of DsrC-trisulfide

confirm that its reduction by this menaquinol analog does occur. This reaction could, in principle, only

involve the DsrMK subunits in analogy to HdrED, with DsrM as the menaquinol-oxidizing subunit that

transfers electrons to DsrK for trisulfide reduction, and in fact, in many organisms only these two subunits

are present [2, 18]. This leaves the function of the DsrJOP subunits as an enigma.

It has been suggested that the DsrJ cytochrome could react with a periplasmic sulfur substrate [10, 15,

24], due to the presence of the unusual His/Cys heme coordination that is also present in SoxXA involved

in thiosulfate oxidation [52] or in TsdA thiosulfate dehydrogenase [26]. We tested both thiosulfate and

sulfite as electron donors to the complex and could not detect any activity with menadione and/or DsrC-

trisulfide as electron acceptors, and sulfite actually decreased the rate of DsrC-trisulfide reduction by

DsrMKJOP (**Figure S6**). Since the DsrJOP module is not essential for some organisms, a hypothesis is that

its role is to increase the energetic efficiency of the DsrMKJOP complex vs. the DsrMK version. We have

previously suggested that the DsrMK and DsrJOP modules could be involved in some form of proton-

translocating quinone cycling, as present in the bc_1 complex, possibly involving quinone-based electron

bifurcation [9]. Alternatively, the DsrJOP module may allow for proton pumping upon DsrC-trisulfide

reduction. Notably, the DsrOP proteins form a quinone-interacting redox module that is widespread in

bacterial oxidoreductases [23, 53], and some proteins of the DsrP family may be involved in proton

pumping, such as the alternative complex III [54, 55] and the reductive dehalogenase complex [56].

CONCLUSION

In conclusion, our results are a major step forward in the molecular understanding of dissimilatory sulfur

metabolism by identifying the function of the DsrMKJOP complex, showing that it interacts with the DsrC-

trisulfide and acts as a menadiol:DsrC-trisulfide oxidoreductase that produces sulfide. The architecture of

the DsrMKJOP complex indicates that it is quite an elaborate machine and clearly more detailed

biochemical and structural studies will be essential to understand its molecular mechanism and how it is

coupled to energy conservation.

ACKNOWLEDGEMENTS

We thank João Carita and Cristina Leitão from ITQB NOVA for the growth of *A. fulgidus* and HPLC analysis,

respectively. This work was funded by the Fundação para a Ciência e Tecnologia (Portugal) through

Fellowships PD/BD/135488/2018 and COVID/BD/152504/2022 (A.C.C.B.), Grants PTDC/BIA-

MIC/6512/2014 and PTDC/BIA-BQM/29118/2017, Research Unit Molecular, Structural and Cellular

Microbiology (MOSTMICRO-ITQB) (UIDB/04612/2020 and UIDP/04612/2020), and Associate Laboratory

Life Sciences for a Healthy and Sustainable Future (LS4FUTURE) (LA/P/0087/2020).

AUTHOR CONTRIBUTIONS

S.S.V. and I.A.C.P. designed research; A.C.C.B. and S.S.V. performed research; A.C.C.B., S.S.V. and I.A.C.P.

analyzed data; A.C.C.B., S.S.V. and I.A.C.P. wrote the paper.

The authors have no competing interests.

Data Availability Statement:

The raw data is available from the authors upon request.

REFERENCES

1.   Muyzer G, Stams AJM. The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev*

Microbiol 2008; **6**: 441–454.

2.   Rabus R, Venceslau SS, Wöhlbrand L, Voordouw G, Wall JD, Pereira IAC. A Post-Genomic View of

the Ecophysiology, Catabolism and Biotechnological Relevance of Sulphate-Reducing Prokaryotes.

Adv Microb Physiol 2015; **66**: 55–321.

3.   Jørgensen BB, Findlay AJ, Pellerin A. The biogeochemical sulfur cycle of marine sediments. *Front*

Microbiol 2019; **10**: 1–27.

4.   McGlynn SE, Chadwick GL, Kempes CP, Orphan VJ. Single cell activity reveals direct electron

transfer in methanotrophic consortia. *Nature* 2015; **526**: 531–535.

5.   Wegener G, Krukenberg V, Riedel D, Tegetmeyer HE, Boetius A. Intercellular wiring enables

electron transfer between methanotrophic archaea and bacteria. *Nature* 2015; **526**: 587–590.

6.   Canfield DE, Rosing MT, Bjerrum C. Early anaerobic metabolisms. *Philos Trans R Soc B Biol Sci* 2006;

361: 1819–1836.

7.   Santos AA, Venceslau SS, Grein F, Leavitt WD, Dahl C, Johnston DT, et al. A protein trisulfide couples

dissimilatory sulfate reduction to energy conservation. *Science* 2015; **350**: 1541–1545.

8.   Oliveira TF, Vonrhein C, Matias PM, Venceslau SS, Pereira IAC, Archer M. The crystal structure of

Desulfovibrio vulgaris dissimilatory sulfite reductase bound to DsrC provides novel insights into the

mechanism of sulfate respiration. *J Biol Chem* 2008; **283**: 34141–34149.

9.   Venceslau SS, Stockdreher Y, Dahl C, Pereira IAC. The 'bacterial heterodisulfide' DsrC is a key

protein in dissimilatory sulfur metabolism. *Biochim Biophys Acta - Bioenerg* 2014; **1837**: 1148–

1164.

10.   Pires RH, Venceslau SS, Morais F, Teixeira M, Xavier A V., Pereira IAC. Characterization of the

Desulfovibrio desulfuricans ATCC 27774 DsrMKJOP complex - A membrane-bound redox complex

involved in the sulfate respiratory pathway. *Biochemistry* 2006; **45**: 249–262.

11.   Pott AS, Dahl C. Sirohaem sulfite reductase and other proteins encoded by genes at the dsr locus

of *Chromatium vinosum* are involved in the oxidation of intracellular sulfur. *Microbiology* 1998;

144: 1881–1894.

12.   Dahl C. A biochemical view on the biological sulfur cycle. In: Piet N. L. Lens (ed.). Environmental

Technologies to Treat Sulfur Pollution: Principles and Engineering. IWA Publishing, 2020; pp 55–

96.

13.   Simon J, Kroneck PMH. Microbial Sulfite Respiration. *Adv Microb Physiol* 2013; **62**: 45–117.

14.   Wasmund K, Mußmann M, Loy A. The life sulfuric: microbial ecology of sulfur cycling in marine

sediments. *Environ Microbiol Rep* 2017; **9**: 323–344.

15.   Mander GJ, Duin EC, Linder D, Stetter KO, Hedderich R. Purification and characterization of a

membrane-bound enzyme complex from the sulfate-reducing archaeon *Archaeoglobus fulgidus*

related to heterodisulfide reductase from methanogenic archaea. *Eur J Biochem* 2002; **269**: 1895–

1904.

16.   Grein F, Pereira IAC, Dahl C. Biochemical characterization of individual components of the

Allochromatium vinosum DsrMKJOP transmembrane complex aids understanding of complex

function in vivo. *J Bacteriol* 2010; **192**: 6369–6377.

17.   Sander J, Engels-Schwarzlose S, Dahl C. Importance of the DsrMKJOP complex for sulfur oxidation

in *Allochromatium vinosum* and phylogenetic analysis of related complexes in other prokaryotes.

Arch Microbiol 2006; **186**: 357–366.

18.   Pereira IAC, Ramos AR, Grein F, Marques MC, da Silva SM, Venceslau SS. A comparative genomic

analysis of energy metabolism in sulfate reducing bacteria and archaea. *Front Microbiol* 2011; **2**:

25–36.

19.   Heiden S, Hedderich R, Setzke E, Thauer RK. Purification of a two-subunit cytochrome-*b*-containing

heterodisulfide reductase from methanol-grown *Methanosarcina barkeri*. *Eur J Biochem* 1994;

221: 855–861.

20.   Künkel A, Vaupel M, Heim S, Thauer RK, Hedderich R. Heterodisulfide reductase from methanol-

grown cells of *Methanosarcina barkeri* is not a flavoenzyme. *Eur J Biochem* 1997; **244**: 226–234.

21.   Hedderich R, Hamann N, Bennati M. Heterodisulfide reductase from methanogenic archaea: A new

catalytic role for iron-sulfur cluster. *Biol Chem* 2005; **386**: 961–970

22.   Hamann N, Mander GJ, Shokes JE, Scott RA, Bennati M, Hedderich R. A cysteine-rich CCG domain

contains a novel [4Fe-4S] cluster binding motif as deduced from studies with subunit B of

heterodisulfide reductase from Methanothermobacter marburgensis. *Biochemistry* 2007; **46**:

12875–12885.

23.   Duarte AG, Barbosa ACC, Ferreira D, Manteigas G, Domingos RM, Pereira IAC. Redox loops in

anaerobic respiration - The role of the widespread NrfD protein family and associated dimeric

redox module. *Biochim Biophys Acta - Bioenerg* 2021; **1862**: 148416.

24.   Grein F, Venceslau SS, Schneider L, Hildebrandt P, Todorovic S, Pereira IAC, et al. DsrJ, an essential

part of the DsrMKJOP transmembrane complex in the purple sulfur bacterium *Allochromatium*

vinosum, is an unusual triheme cytochrome c. *Biochemistry* 2010; **49**: 8290–8299.

25.   Cheesman MR, Little PJ, Berks BC. Novel heme ligation in a *c*-type cytochrome involved in

thiosulfate oxidation: EPR and MCD of SoxAX from *Rhodovulum sulfidophilum*. *Biochemistry* 2001;

40: 10562–10569.

26.   Denkmann K, Grein F, Zigann R, Siemen A, Bergmann J, van Helmont S, et al. Thiosulfate

dehydrogenase: A widespread unusual acidophilic *c*-type cytochrome. *Environ Microbiol* 2012; **14**:

2673–2688.

27.   Alric J, Tsukatani Y, Yoshida M, Matsuura K, Shimada K, Hienerwadel R, et al. Structural and

functional characterization of the unusual triheme cytochrome bound to the reaction center of

Rhodovulum sulfidophilum. *J Biol Chem* 2004; **279**: 26090–26097.

28. Grein F, Ramos AR, Venceslau SS, Pereira IAC. Unifying concepts in anaerobic respiration: Insights

from dissimilatory sulfur metabolism. *Biochim Biophys Acta - Bioenerg* . 2013; **1827**: 145–160

29. Wagner T, Koch J, Ermler U, Shima S. Methanogenic heterodisulfide reductase (HdrABC-MvhAGD)

uses two noncubane [4Fe-4S] clusters for reduction. *Science* 2017; **357**: 699–703.

30. Pelmenschikov V, Ferreira D, Venceslau SS, Hildebrandt P, Pereira IAC, Todorovic S. Substrate-

Dependent Conformational Switch of the Noncubane [4Fe-4S] Cluster in Heterodisulfide Reductase

HdrB. *J Am Chem Soc* 2023; **145**: 7–11.

31. Stetter KO, Lauerer G, Thomm M, Neuner A. Isolation of Extremely Thermophilic Sulfate Reducers:

Evidence for a Novel Branch of Archaebacteria. *Science* 1987; **236**: 822–824.

32. Francis RT, Becker RR. Specific indication of hemoproteins in polyacrylamide gels using a double-

staining process. *Anal Biochem* 1984; **136**: 509–514.

33. Venceslau SS. Electron transfer chains in sulfate reducing bacteria. Instituto de Tecnologia Química

e Biológica António Xavier, Universidade Nova de Lisboa. Ph.D. Thesis Dissertation 2011.

34. Fieser LF. Convenient Procedures for the Preparation of Antihemorrhagic Compounds. *J Biol Chem*

1940; **133**: 391–396.

35. Dahl C, Kredich NM, Deutzmann R, Truper HG. Dissimilatory sulphite reductase from

Archaeoglobus fulgidus: Physico-chemical properties of the enzyme and cloning, sequencing and

analysis of the reductase genes. *J Gen Microbiol* 1993; **139**: 1817–1828.

36. Ferreira D, Barbosa ACC, Oliveira GP, Catarino T, Venceslau SS, Pereira IAC. The DsrD functional

marker protein is an allosteric activator of the DsrAB dissimilatory sulfite reductase. *Proc Natl Acad*

Sci USA 2022; **119**.

37. Venceslau SS, Cort JR, Baker ES, Chu RK, Robinson EW, Dahl C, et al. Redox states of *Desulfovibrio*

vulgaris DsrC, a key protein in dissimilatory sulfite reduction. *Biochem Biophys Res Commun* 2013;

441: 732–736.

38.   Riddles PW, Blakeley RL, Zerner B. Reassessment of Ellman's Reagent. *Methods Enzymol* 1983; **91**:

564   49–60.

565   39.   Weiner JH, Cammack R, Cole ST, Condon C, Honoré N, Lemire BD, et al. A mutant of *Escherichia*

566   *coli* fumarate reductase decoupled from electron transport. *Proc Natl Acad Sci* 1986; **83**: 2056–

567   2060.

568   40.   Fahey RC, Newton GL. Determination of Low-Molecular-Weight Thiols using Monobromobimane

569   Fluorescent Labeling and High-Performance Liquid Chromatography. *Methods Enzymol* 1987; **143**:

570   85–96.

571   41.   Madadi-Kahkesh S, Duin EC, Heim S, Albracht SPJ, Johnson MK, Hedderich R. A paramagnetic

572   species with unique EPR characteristics in the active site of heterodisulfide reductase from

573   methanogenic archaea. *Eur J Biochem* 2001; **268**: 2566–2577.

574   42.   Duin EC, Bauer C, Jaun B, Hedderich R. Coenzyme M binds to a [4Fe-4S] cluster in the active site of

575   heterodisulfide reductase as deduced from EPR studies with the [^{33}S]coenzyme M-treated enzyme.

576   *FEBS Lett* 2003; **538**: 81–84.

577   43.   Duin EC, Madadi-Kahkesh S, Hedderich R, Clay MD, Johnson MK. Heterodisulfide reductase from

578   *Methanothermobacter marburgensis* contains an active-site [4Fe-4S] cluster that is directly

579   involved in mediating heterodisulfide reduction. *FEBS Lett* 2002; **512**: 263–268.

580   44.   Bhave DP, Hong JA, Keller RL, Krebs C, Carroll KS. Iron-sulfur cluster engineering provides insight

581   into the evolution of substrate specificity among sulfonucleotide reductases. *ACS Chem Biol* 2012;

582   **7**: 306–315.

583   45.   Rossi M, Pollock WBR, Reij MW, Keon RG, Fu R, Voordouw G. The hmc operon of *Desulfovibrio*

584   *vulgaris* subsp. *vulgaris* Hildenborough encodes a potential transmembrane redox protein

585   complex. *J Bacteriol* 1993; **175**: 4699–4711.

46. Pereira PM, Teixeira M, Xavier A V., Louro RO, Pereira IAC. The Tmc complex from *Desulfovibrio*

vulgaris Hildenborough is involved in transmembrane electron transfer from periplasmic hydrogen

oxidation. *Biochemistry* 2006; **45**: 10359–10367.

47. Tietze M, Beuchle A, Lamla I, Orth N, Dehler M, Greiner G, et al. Redox potentials of

methanophenazine and CoB-S-S-CoM, factors involved in electron transport in methanogenic

archaea. *ChemBioChem* 2003; **4**: 333–335.

48. Chobot SE, Hernandez HH, Drennan CL, Elliott SJ. Direct Electrochemical Characterization of

Archaeal Thioredoxins. *Angew Chemie Int Ed* 2007; **46**: 4145–4147.

49. Zöphel A, Kennedy MC, Beinert H, Kroneck PMH. Investigations on microbial sulfur respiration.

Arch Microbiol 1988; **150**: 72–77.

50. Flohé L. The fairytale of the GSSG/GSH redox potential. *Biochim Biophys Acta - Gen Subj* 2013;

1830: 3139–3142.

51. Berndt C, Lillig CH, Flohé L. Redox regulation by glutathione needs enzymes. *Front Pharmacol* 2014;

5.

52. Bamford VA, Bruno S, Rasmussen T, Appia-Ayme C, Cheesman MR, Berks BC, et al. Structural basis

for the oxidation of thiosulfate by a sulfur cycle enzyme. *EMBO J* 2002; **21**: 5599–5610.

53. Duarte AG, Catarino T, White GF, Lousa D, Neukirchen S, Soares CM, et al. An electrogenic redox

loop in sulfate reduction reveals a likely widespread mechanism of energy conservation. *Nat*

Commun 2018; **9**: 5448.

54. Sousa JS, Calisto F, Langer JD, Mills DJ, Refojo PN, Teixeira M, et al. Structural basis for energy

transduction by respiratory alternative complex III. *Nat Commun* 2018; **9**: 1728.

55. Sun C, Benlekbir S, Venkatakrishnan P, Wang Y, Hong S, Hosler J, et al. Structure of the alternative

complex III in a supercomplex with cytochrome oxidase. *Nature* 2018; **557**: 123–126.

56. Kublik A, Deobald D, Hartwig S, Schiffmann CL, Andrades A, von Bergen M, et al. Identification of a

multi-protein reductive dehalogenase complex in *Dehalococcoides mccartyi* strain CBDB1 suggests

a protein-dependent respiratory electron transport chain obviating quinone involvement. *Environ*

Microbiol 2016; **18**: 3044–3056.

Figure Legends

Figure 1 – UV-visible and EPR spectroscopy of the *A. fulgidus* DsrMKJOP complex. (A) UV-visible spectra of as-purified DsrMKJOP (black), menadiol-reduced (dark grey), and dithionite-reduced (light grey). (B) EPR spectra of the $[4Fe-4S]^{3+/2+}$ cluster of the as-purified DsrMKJOP (black), and upon menadiol reduction (grey). *g*-values are marked in the spectrum of as-purified DsrMKJOP. All menadiol reduction spectra were obtained in a 1:500 protein to menadiol ratio.

Figure 2 – Interaction assays between DsrC-trisulfide and DsrMKJOP complex. Sensorgrams of DsrC-trisulfide interaction with increasing concentrations of DsrMKJOP (8.9 nM to 568 nM) using SPR. Sensorgrams were measured in duplicate.

Figure 3 – Kinetic assays of the effect of DsrC-trisulfide on DsrMKJOP activity. (A) Menadiol:DsrC-trisulfide oxidoreductase activity catalyzed by *A. fulgidus* DsrMKJOP (black), and in the absence of DsrC-trisulfide (light grey), absence of DsrMKJOP (dashed line), and absence of menadiol (dotted line). (B) Menadiol:DsrC-trisulfide oxidoreductase activity catalyzed by *A. fulgidus* DsrMKJOP upon successive additions of DsrC-trisulfide (arrow indicates addition of 5 µM DsrC-trisulfide). (C and D) Gel-shift analysis of MalPEG-labeled DsrC from (A) and after reaction with DsrMKJOP and 0.5 mM menadiol over time (at 0, 1, 2, 3, 5, 10, 15, 20 and 30 min), respectively. MalPEG causes a shift of approximately 10 kDa per each labeled cysteine. (E) Gel-shift analysis of MalPEG-labeled DsrC with increasing concentrations of menadiol (0, 0.8, 4, 20, 100 and 500 µM), reacted for 30 mins. (M, molecular marker; T_0, time zero; T_f, time at the end of reaction).

Figure 4 – Peptide mass fingerprinting spectra of DsrC C-terminal peptides after alkylation with IA. (A) DTT-reduced DsrC (before the reaction with *A. fulgidus* DsrAB and sulfite). (B) DsrC-trisulfide (produced in

the *in vitro* enzymatic reaction with *A. fulgidus* DsrAB and sulfite). (C) DsrC recovered after the *in vitro*

enzymatic reaction with *A. fulgidus* DsrMKJOP complex and menadiol. The C-terminal peptide of DsrC-

trisulfide (1530.7 Da) is marked with an arrow in (C) showing its disappearance from (B) to (C). (IA,

iodoacetamide; DTT, dithiothreitol).

Figure 5 – Kinetic assays of the effect of DsrC-trisulfide on DsrMKJOP activity and sulfide production. (A)

Schematic representation of menadiol:DsrC-trisulfide oxidoreductase activity catalyzed by *A. fulgidus*

DsrMKJOP. (B) Menadiol:DsrC-trisulfide oxidoreductase activity catalyzed by *A. fulgidus* DsrMKJOP with

increasing concentrations of DsrC-trisulfide (0 μM to 10 μM). (C) Hydrogen sulfide measured by HPLC with

increasing concentrations of DsrC-trisulfide (0 μM to 10 μM) after 30 min of enzymatic reaction.

648
649

A

B

C