Embryonic life histories in annual killifish: adapted to what?

Tom JM Van Dooren

team VPA Phenotypic Variability and Adaptation

CNRS, Institute for Ecology and Environmental Sciences iEES Paris

Sorbonne University

4, Place Jussieu, 75005 Paris France

tvdooren@gmail.com

Abstract: 218 words

Main text: 8102 words

Five figures and one table

Supplementary Material with two figures

Acknowledgements

This study was supported by action Climat-AmSud 21-CLIMAT-06, funded by CNRS France, CONICYT

Chile and ANII Uruguay.

Abstract (218)

Adaptation requires an evolving strategy and an environment. Given an environment, we predict or

estimate which strategies are adapted. Given a strategy, we want to know in which environments it

might be adapted. Example calculations aiming to determine such environments, named

evolutionarily singular environments ESE, are presented using lab data on embryonic life histories of

Austrolebias annual killifish. Annual killifish embryos can arrest development and survive desiccation

of temporary ponds in the soil. They might implement diversified bet-hedging, generally seen as an

adaptation to uncertain environments. Using parameter estimates and parsimonious assumptions, a

population dynamical model is constructed with explicit developmental stages. Using invasion

fitness gradients of rates of development and hatching probabilities, it is investigated whether these

could be adapted to pond filling regimes with gradual filling and drying and deterministic within-year

variation only. The life history as a whole is not adapted to the regular within-year annual cycles

investigated, with one or two periods where reproduction can occur. Faster development rates were

always favoured, just as in constant environments. Only for hatching probabilities, pond filling

regimes were found which made their invasion fitness sensitivities zero. However, the observed trait

values did not have long-term evolutionary stability in these ESE. Therefore, neither the

developmental rates nor the hatching strategy seem adapted to within-year patterns of

environmental change.

Keywords: Adaptation, environmental change, annual killifish, diapause, pond life cycle, eco-evo-

devo, cumulative hazards, longitudinal studies, Adaptive Dynamics.

Introduction

Evolutionary adaptation requires traits which can evolve and an environment. Part of the environment, the "external driver" (e.g., Metz et al. 2008), is usually seen as given and not affected by changes in an evolving population. Effects of environmental variation on adaptation are assessed by the population structuring they generate (Southwood 1977), and through the means and variances of demographic parameters (Cohen 1966, Simons 2011, Starrfelt and Kokko 2012). In modelling, the actual environment often figures by means of examples congruent with assumptions on demographic parameters (e.g., ten Brink et al. 2020). Similarly, calls for better predictions of adaptation to climate change urge more elaborate demographic and trait models (Urban et al. 2023) but it is unclear whether we are grasping the environments themselves and their effects on demography well. In evolutionary conservation biology, adaptive population management (e.g., Zhan et al. 2015) includes controlling the environment, which then isn't external nor a given.

Evolutionary anthropologists have discussed the "environment of evolutionary adaptedness" (Bowlby 1973), the environment in which a population evolved and presumably became adapted. Historical aspects of this definition were problematic (Irons 1998). We currently favour a rather a-historical view of adaptation (Reeve & Sherman 1993), where an adapted suite of traits confers highest fitness among a set of alternatives in a given environment. For a given organismal strategy, observed in a single or across different environments, we often want to know in which environments it would be adapted. It would be the most common strategy in them (Reeve & Sherman 1993) or one of the strategies maintained in a protected polymorphism (Geritz et al. 1998). In these environments, there is no long-term directional selection giving rare alternative strategies a systematic advantage (e.g., Ellner 1997, Coulson 2021). For adaptive management, conservation, and to investigate whether we have understood the environment sufficiently, knowing such environments seems useful.

Here, an example is presented of a search for environments to which an observed strategy could be

adapted. This is done for embryonic life history traits in an annual killifish (Van Dooren & Varela-

Lasheras 2018). Annual killifish are oviparous cyprinodontiform fish species from the suborder

Aplocheiloidei which evolved "annualism" repeatedly (Furness et al. 2015a, Helmstetter et al. 2016,

Thompson et al. 2021). They use embryonic diapauses to persist in temporary ponds in Africa and

South America which dry out seasonally (Berois et al. 2015) and where the start and end of a wet

period generally vary between years. Many annual fish species are threatened (Volcan & Lanés

2018). Their habitats are restricted in space and managing them seems feasible. We could aim to

keep annual fish populations adapted by preserving or changing their environments, not by seeing

these as given.

Data patterns in annual killifish embryonic life histories have been interpreted as in agreement with

bet-hedging adaptations to environmental uncertainty (Furness et al. 2015b, Pinceel et al. 2015,

Polačik et al. 2017, Van Dooren & Varela-Lasheras 2018), but also to predictable properties of the

environments. Furness et al. (2015b) proposed that a diapause occurs less or not at high

temperatures because these predict the probability that another generation can be completed

before the cool dry season starts. Similarly, Polačik et al. (2017) proposed that eggs from younger

mothers develop faster to be ready when an extra opportunity to hatch and reproduce within a

season presents itself. However, in *Austrolebias* annual fish (Alonso et al. 2023) for example,

embryos will not hatch and grow when they have not experienced a dry period yet. The environment

proposed where this would be adaptive therefore needs to be an environmental regime with two

periods within a year where reproduction can occur, separated by at least partial drying of a pond.

Complete drying would avoid competition between cohorts hatched in the two periods. A

demographic model and fitness calculations across the entire life cycle (Childs et al. 2010, de Jong et

al. 2011, Van Dooren & Varela-Lasheras 2018) in different seasonal patterns of wet and dry periods

are required to investigate whether the observed patterns would be adapted to them.

Such a model is presented here and it is used to investigate whether developmental rates and

hatching probabilities might be adaptations to deterministic within-year environmental patterns.

The model makes two important assumptions: it is assumed that a developmental strategy can only

be adapted when there is no persistent directional selection on it. Second, data observed on

individual development in lab conditions are assumed to be representative of what individuals do in

the field. The model structure is tailored to specifics of annual killifish life histories and is integrating

estimated parameters mostly from *A. bellottii*, the Argentinian pearl killifish (Steindachner 1881, Van

Dooren & Varela-Lasheras 2018). The results demonstrate that the traits of observed embryonic life

histories are not adapted to environments changing deterministically within years. Faster

developmental rates are always favoured. For hatching probabilities, however, directional selection

vanishes in some environments investigated. Based on the results, expectations are formulated for

the adaptiveness of patterns when between-year variation occurs, deterministic or not.

Methods

In the model detailed below, annual fish individuals can find themselves in different pond

compartments and environments (Figure one). Environmental and population states are projected

forward in time on a day-to-day basis.

Embryonic life histories

The strategy for which we want to find environments where it would be adapted, consists of rates of

development and hatching probabilities estimated for a group of embryos of the Argentinian pearl

killifish *A. bellottii* (Van Dooren & Varela-Lasheras 2018). The data used to parameterize the

embryonic life history were collected at 19.5C and 20.5C, while the average annual temperature for

their site of origin was 17.9C (Ingeniero Maschwitz Argentina, Worldclim value used in Helmstetter

et al. 2020). In this section, next to detailing the given strategy, we also include the description of

mortality rates in different conditions, so that this section can be seen as the description of life

history events in the egg bank compartment of the model. Events occur with different rates when

embryos are in (1) wet conditions (e.g., their location covered by water and conditions hypoxic),

when they are in (2) dry conditions (oxygenated with limited desiccation) and haven't been rewetted

yet (first dry period) and (3) in subsequent dry periods. In dry conditions, mortality rates differ

between developmental stages.

The cohort from which development rates were estimated for "dry" conditions started incubation in

wet oxygenated conditions. After some time, it was further incubated in air with 100% relative

humidity (Supplementary Material). Five distinct developmental stages were considered (Figure one,

Van Dooren & Varela-Lasheras 2018, simplified from Wourms 1972a), each with an age-dependent

pattern of the risk to enter the next developmental stage in the first four developmental stages.

Embryos were assigned to stage one at age zero. In stage two, a neural keel and somites have

appeared. In stage three, optic cups were visibly present. In stage four, the eyes had become

pigmented. In the fifth stage the embryo completely surrounded the yolk sac. In the conditions

where the strategy has been determined, diapause has been observed in *Austrolebias bellottii* in

stages one (DI) and five (DIII, Van Dooren & Varela-Lasheras 2018) as age intervals with negligible

risk to enter the next stage. From stage five, embryos can hatch when they are rewetted. Hatching

probabilities were estimated using logistic regressions (Van Dooren & Varela-Lasheras 2018). It was

noted that after some time, the rate of events (deaths or developmental transitions) decreased to

almost zero. A similar observation was made by Polačik & Vrtilek (2023). This slowdown is more

prominent even after the first rewetting of eggs and we incorporated that into the life history model

in an ad-hoc manner as explained below.

For the first dry period embryos experience, estimated cumulative hazards were used to obtain

transition probabilities between developmental stages or from any developmental stage to the dead

state (Supplementary Material). Fifteen different transitions (from each developmental stage to all

more advanced ones and to the dead state) can be observed across a time interval, some more often

than others. It is only age given in the number of days in dry conditions which is assumed to affect

developmental progress, therefore we use this integer as the individual age variable. Matrices were

derived with transition probabilities per day using age-dependent multi-state modelling with

competing risks (de Wreede et al. 2011). Van Dooren & Varela-Lasheras (2018) found that many

embryos don't resume development after a first rewetting. To integrate this in the model in a

straightforward manner, a 1/0 state variable was added to the individual state denoting whether an

embryo would be in a state where development proceeds (1), or remains in a generally arrested

state (0). Each time an embryo experiences wet conditions after the first dry period, it enters this

non-advancing state with probability p_d which is maintained during the next dry period. The

probability of survival per day $e^{-\mu_d}$ in the non-advancing state, which was equal in all stages, and

the proportion entering the advancing state p_d were estimated based on the data in Van Dooren &

Varela-Lasheras (2018) and are given in Table one.

When stage five embryos are rewetted (i.e., covered by standing water again), they can hatch on

that day only in the model. This occurs with probabilities which differ between the first and later

rewettings and with an age effect at first rewetting (Van Dooren & Varela-Lasheras 2018). Three

hatching parameters β_1 , β_2 , β_3 required to model this were estimated and are given in Table one.

Upon first rewetting, embryos in stage five hatch with probability $g(\beta_1 + \beta_2 a)$, with g the inverse

logit function and a the number of days an embryo has been dry. First rewetting is therefore an 1/0

state variable. At later rewettings, embryos hatch with probability $g(\beta_3)$. Eggs which don't hatch

remain in the soil at the same location. In wet soil conditions, embryonic development does not

proceed (Van Goor 2009, Polačik et al. 2021, 2023), presumably due to lack of oxygen. In a pilot

study on the related species *Acantholebias luteoflammulatus* (van Goor 2009), individual survival

was followed in the field. The probability per day that an egg survives in wet conditions $e^{-\mu_w}$ was

computed from the results and used across all stages (Table one). The value used is lower than an

estimate (0.997) based on the data in Polačik et al. (2023). There are no stage-specific survival

probabilities in wet soil estimated yet. Fitness effects of developmental rates per stage in dry

conditions and in the advancing state were predicted as detailed below, and fitness effects of

hatching parameters β_1 and β_3. The probabilities of individual switching between 1/0 states and of

the age plasticity of hatching were not treated as traits which can evolve. The parameter values for

these transitions were based on observations at few censuses and considered too much ad-hoc in

comparison to other embryonic life history parameters.

Evolutionarily Singular Environments

Given a strategy, we want to know in which environments it might be adapted. When we start from

a vector of scalar trait values **x** representing a strategy (for function-valued **x**, see Dieckmann et al.

2006), we first need to embed it into a population dynamical model with population regulation and a

mode of genetic inheritance, and in an "external" environment **E**. This allows calculation or

simulation of "resident" stationary populations where all individuals have the same genotype and

strategy **x**, and following the dynamics of rare strategies with a different genotype and phenotype **x'**.

For the sake of easiness of presentation, we assume that there is a single such attractor of the

population dynamics Att(**x**, **E**). Using this model and an Att(**x**, **E**), we can determine invasion fitnesses

of rare mutants **x'**. We assume that a mutation occurs at a single locus and that it has phenotype **x'**

in heterozygotes. When a mutant allele is rare, the frequency of mutant homozygotes is negligible.

We can then write invasion fitnesses as functions s(**x'**, Att(**x**, **E**), **E**). For a fixed **x**, we can call an

environment **E** an evolutionarily singular environment ESE when

$\left. \dfrac{\partial}{\partial x'} s(\boldsymbol{x'}, Att(\boldsymbol{x}, \boldsymbol{E}), \boldsymbol{E}) \right|_{x'=x} = \boldsymbol{0}$ (Eqn. 1)

This is a vector of partial fitness derivatives towards each of the phenotypic traits in mutants,

evaluated at the resident trait values x. When this vector is zero, no mutant with phenotype x'

slightly differing from x in one or several traits can invade.

This definition does not suggest that E evolves. E does not have to be the environment in which an

organism supposedly became adapted. It is just singular. If multiple attractors of the population

dynamics occur for an x, the definition needs to be checked in each of them. Adaptive dynamics

(Metz et al. 1996, Geritz et al. 1998) usually assumes that x evolves in a given E. Trait vector x^* is

called an evolutionary singularity or evolutionarily singular strategy ESS, when Eqn.1 holds for it, for

a given E. Eqn (1) is a first condition for E to be an environment where x is adapted and is expected

to stay adapted. It can then be verified using adaptive dynamics in a given E whether resident

strategies which are slightly different from x would be replaced by strategies more similar to it and

whether evolution halts when x is the or a common strategy in the population (Geritz et al. 1998).

The environment

The temporary pond is assumed to be a disc with a fixed center. The maximum radius which can be

covered by water is r_{max} (Table 1). The state of a pond is characterized by how much of its extent is

covered by water. All non-negative radiuses within [0, r_{max}] are allowed. The radius covered by water

$r_{pond}(t)$ can change gradually and deterministically per day, where t can be any date. When $r_{pond}(t) =$

0, it is in the dry state. In a wet state with positive $r_{pond}(t)$, water is present and fish can hatch, grow

and reproduce. When they reproduce, adults lay their eggs in the part of the pond soil which is wet.

Depending on the location of an egg within the pond extent, dry conditions start at different times.

Location is therefore a state variable of each individual egg. Location states were rounded down to

arrive at a limited number of discrete location classes to model (Supplementary Material).

Pond radius profiles over time $r_{pond}(t)$ were kept relatively simple, with up to two wet periods within

a year and with linear radius increases and decreases within a wet period. In longer simulations,

patterns within a year were repeated across years. In leap years, $r_{pond}(t)$ of the previous year was

copied into January 1 to December 30 and December 31 of the previous year was duplicated. To

gain a first insight in the effects of pond regime on population dynamics, a small number of seasonal

scenarios were explored with a single wet period allowing reproduction (see Figure two for

examples). In part of these scenarios, this period was preceded by a wet period too short to allow

reproduction. In the simplest scenario (Y0), the pond had water starting on April 11 (day 101),

increasing to maximum size r_{max} on July 19 (day 200) and became dry again on October 28 (day 301).

In scenario Y0, the pond covers $0.3r_{max}$ on day 130 and $0.7r_{max}$ on day 170. From this scenario the

period from April 11 to July 19 was then modified, making the period where most of the pond extent

is covered by water smaller or larger. Deceleration scenarios were constructed where there was an

initial phase of fast filling where $0.7r_{max}$ would be reached on days 163 (YD1), 150 (YD2) and 145

(YD3), respectively. This was followed by a second phase of slow filling with a time span of 37 (YD1),

50 (YD2) and 55 (YD3) days respectively to reach the maximum size r_{max} by 19 July. Scenarios with

accelerated filling were constructed where $0.3r_{max}$ would be reached after a period of slow filling on

days 136 (YA1), 148 (YA2) and 163 (YA3), followed by faster filling, with r_{max} still reached on July 19.

In a second category of modifications of Y0, a short wet period was inserted from days 71 to 90, with

the maximum pond radius within that short period reached on day 80 at values mr_{max}. Scaling

parameter m varied from 0.2 to 1 in steps of 0.2, yielding scenarios YS1-YS5. Refilling of a partly

desiccated pond was not considered for now and neither was between-year variability. Other

environmental state variables than $r_{pond}(t)$ were not modelled. The model does not contain explicit

temperature effects yet. Diurnal variation is prevalent in annual fish habitats (Žák & Reichard 2020)

but not explicitly modelled here.

Fish life histories

Fish are assumed to move freely within the available water body. Parameter estimates are based on

Helmstetter & Van Dooren (2019) and Van Dooren & Varela-Lasheras (2018). Individual fish have

their age since hatching and sex as state variables. The number of fish per state and their produced

eggs are modelled on a day-to-day basis. Different events which can happen on the same day are

treated sequentially. The sex ratio among the hatchlings on each day is equal. Juveniles mature at

age since hatching α. A simple model of population regulation is adopted. The number of fish within

a pond is limited by its surface. It is assumed that each square meter of wet surface can contain a

maximum of K adults. When more than the maximum number of allowed adults is present on a day

(K times the area of the pond), surplus adults are chosen at random and erased from the population

state. There is no other source of daily mortality among adults during a wet period, senescence is

not expected for the wet season lengths modelled (Liu and Walford 1990). When the number of

adults N(t) is below the maximum which a pond allows, juveniles can survive with a survival

probability per day equal to the proportion of surface which is not "occupied" on that day $[1 - N(t) /$

$(\pi K \, r(t)^2)]_{\geq 0}$ times a density-independent survival probability $e^{-\mu_J}$. Reproduction occurs each day

after density-dependence. Adult females have a fixed individual fecundity per day of eggs f which

are all fertilized when there are males present in the pond on that day after competition and the sex

ratio (ratio numbers of females/males) is within the interval $[S^{-1}, S]$. Otherwise, fecundity is set to

zero because of presumed sperm limitation in males or male aggression. Eggs are deposited at a

random location within the pond (Supplementary Material). At the start of a dry period, the

remaining adults die. Parameter estimates α, f, μ_J (Table 1) are based on Helmstetter & Van Dooren

(2019) and Van Dooren & Varela-Lasheras (2018).

Simulations of population dynamics

As the extent of the wet pond area changes continually, numbers of individuals are projected

forward, not their densities per unit of area (Supplementary Material). Given an initial population

state, an egg bank with a number of embryos or a fish population can be projected forward over a

series of successive years, where each year repeats the pond cycle of the previous year, to find the

attractors of the population dynamics. On these attractors, population states change within a year,

but on Poincaré sections at a specific day within the year, the population states can have fixed-point

or non-equilibrium attractors. Only fixed-point attractors were observed.

The model for the pond environment, population dynamics and invasion fitness is coded in R (R Core

Team 2020) and the scripts are available on https://doi.org/10.5281/zenodo.10992414.

For the purpose of illustration, forward projection was done for example profiles Y0, YA, YD and YS

and the stationary dynamics over one year was then simulated on a day-to-day basis.

Mutational effects

To implement the check of adaptation proposed by Eqn. 1, we need a vector of phenotypic trait

parameters $\boldsymbol{x}$ and the effects of mutations. The embryonic annual fish strategy considered consists

of four stage-specific developmental transition parameters and two hatching parameters. The rates

are all age-dependent strategies, which would make them four function-valued strategies to assess

(Kirkpatrick & Heckman 1989). I opted for a simplification. Four stage-specific trait parameters

represent multiplicative changes to the cumulative integrated stage-specific rates (Supplementary

Material). The trait values are all one for resident individuals and can be smaller or larger than that

for mutants, which then develop slower or faster out of the stages where their trait value differs

from value one. For hatching, multiplicative changes to hatching parameters β_1 and β_3 of the linear

predictor in the logistic regression are the traits. When the trait values are all fixed at value one, the

transition probabilities and hatching probabilities determined from the data are unchanged and

represent the assumed common strategy in a resident population of this species. The Supplementary

Material details how modified rates affect the transition probabilities between stages.

Invasion analysis

It is assumed that there is no genetic variation in the resident population with phenotypic effects.

For a mutation which is relatively rare, the number of mutant alleles equals the number of mutant

individuals because these are approximately all heterozygotes at a single locus of a mutant and a

resident allele. Offspring numbers are determined by female fecundity and the availability of adult

males. Mutant adult females contribute the mutant allele to half of their eggs. Male adult mutant

individuals contribute their mutant allele to half of the eggs they fertilize each day, which is assumed

to equal the total number of eggs produced on that day (f times the number of resident adult

females present), divided by the resident males present.

To estimate invasion fitness of a specific mutation with phenotype x', the population dynamics of

the data-based resident strategy x was simulated to obtain so-called resident stationary regimes

(Supplementary Material). Then the mutant genotype was introduced and simulated competing

against the resident strategy over eight complete dry-wet period cycles. The final number of mutant

individuals was divided by the initial number at the start of this time interval and raised to the power

one over the interval length (in days) and the result log transformed. Invasion fitness is thus the log

geometric average population growth rate per day of this mutant population given an environmental

regime and a resident data-based population strategy.

Fitness sensitivities

To assess the effects of mutants with specific trait changes on invasion fitness in a given

environmental regime, mutants with trait values between 0.9 and 1.1 were generated for each of

the separate traits and their invasion fitnesses were estimated in an established population with the

data-based strategy. This was done for each of the pond filling scenarios Y0, YA, YD and YS. Per

environmental regime, the invasion fitnesses for mutations in each trait with value 1.1 were used as

a measure of invasion fitness sensitivity (Caswell 2019) and plotted to see how environmental

properties might affect strengths of selection and change the potential adaptedness of strategies.

For the five stage-specific mortalities, invasion fitnesses were also calculated using the same

approach as for rates of development. This allowed a comparison of strengths of selection between

developmental and mortality rates. To compare these strengths of selection more easily between

environments, we assigned parameter values to the scenarios Y0, YD and YA, and to Y0 and YS, such

that invasion fitnesses of a specific trait change could be plotted as functions of environmental

parameters. YD1-3 were given parameter values -1, -2 and -3, Y0 value zero and YA1-3 the values 1,

2 and 3. To compare effects of adding a short wet period where reproduction is impossible, Y0 was

again assigned parameter value zero, and YS1-5 the corresponding values of m.

Evolutionarily Singular Environments

A more detailed parametric specification of pond filling regimes (Supplementary Figure one) was

investigated to search for ESE. Here, a parameter vector ρ contained ten parameters ρ_i with values

between zero and one, affecting dry and wet interval lengths and maximum pond radiuses within

intervals. A search algorithm on vectors ρ was implemented to investigate whether the data-based

strategy might be adapted to one or some of these. The Euclidian norm of the vector of invasion

fitnesses of mutants with trait values 1.1 in the four developmental traits and in the two hatching

probabilities was calculated for each environmental regime visited by the algorithm as a measure of

the selection gradient. These gradients were minimized as a function of ρ by means of the L-BFGS-B

method (Byrd et al. 1995), programmed in the optim function for R (R Core team 2020). In an ESE

environment ρ^* where the data-based strategy is adapted, this norm would become zero.

Optimizations were started from different initial pond regimes with one or two wet periods, also

with two wet periods long enough for reproduction to occur. In all cases explored, the algorithm

converged to a local minimum of the gradient of invasion fitness. Its initial and final values at these

local minima were calculated. Environmental regimes with the smallest gradients found are plotted,

their resident population dynamics simulated and shown, as well as the pattern of invasion fitnesses

of mutants in these environments.

Results

Population Dynamics

Pond regime differences within years can change the expected number of individuals in the egg bank

more than tenfold (Figure two). Fish are always a minority among the individuals present in the

population (Fig. 2). Different speeds of pond filling make periods with many reproducing adults

shorter or longer. Coexistence of adults and juveniles can occur over extended periods (Fig. 2). Their

numbers vary with pond extent, but the number of adults is not always matching the profile changes

closely. It seems that the strong competition effects of adults on juveniles prevent this. During dry

periods all embryonic developmental stages are usually present but stage five is generally most

abundant (Fig. 2). At different distances from the pond center, the number and distribution over

stages of embryos change predictably, with more advanced stages found sooner at larger distances

from the pond center (Supplementary Figure two). At the start of a dry period after reproduction,

earlier stages are the most abundant near the pond center and more advanced developmental

stages dominate near the pond edge (Supplementary Figure two).

The largest egg bank changes occur within wet periods, due to hatching, overall lower survival, and

the addition of new eggs. Variable speeds of pond filling have clear effects. With faster initial and

slower later filling as the maximum radius is approached, the overall size of the egg bank increases at

all times because more individuals reproduce (Fig. 2A). For comparison, in pond regimes with step

functions between no water and a full pond, populations go extinct when the wet season is shorter

than 189 days (results not shown). The increase in number of embryos accelerates or decelerates as

the pond radius does (Fig. 2A), but there cannot be a causal relationship, as a deceleration also

occurs for other pond regimes where a short wet period precedes the one with reproduction (Fig.

2B). Accelerating egg number changes with increasing pond size might be indicative of a lack of

competition between mature fish. With an additional short pre-period of pond filling covering the

pond radius to some extent (Fig. 2B), stage five embryos hatch in that period without any chance of

reproducing and the embryos which experienced their first rewetting have their hatching state

variable changed. In the main wet periods (Fig. 2B), we can see more intense competition: the

fraction of the juvenile population contributing to the pool of reproducing adults is smaller, while

overall, there are many more new eggs produced. If we need to guess which among these

environments might be the most adapted, based on simple principles such as the optimization of

population size (Charlesworth 1994) or the pessimization of the environment (Metz et al. 2008), it

would be the pond filling regime where the egg bank is largest and where competition is most

intense, so YS1 or YS2. A combination of decelerated pond filling and a short wet period before a

long one might be the environment where the data based strategy is the most adapted, if only

within year variation would occur.

Invasion fitness patterns

When invasion fitnesses are estimated for changes in each trait in each of the example scenarios, it

can be observed that invasion fitness changes are largest for mortality and development in stage

one, and for early hatching (Figure three). Note that for mortalities in other stages than stage one,

the slope of invasion fitness on mutational trait change is small. This can be explained by the already

high survival rates in each of these stages. For the development rates, there seems only weak

selection to increase developmental speed in most stages (Fig. 3). However, selection was always

towards acceleration. Invasion fitness slopes changed when environmental parameters changed, in

particular for the traits under stronger selection (Figure four). For the range of changes explored in

(Fig. 4), invasion fitness slopes did not change sign except for hatching at the first rewetting, when

short wet periods without reproduction were added to the reference scenario Y0. In these YS pond

filling regimes, parameter m was varied gradually to find an ESE but it was not present, at value m =

0.1, selection changed direction abruptly.

Evolutionarily Singular Environments

It was found that the minimization algorithm could make jumps to environmental regimes where dry

periods would disappear and the population went extinct. These iterations were restarted with a

constraint which imposed that a dry period would exist for of at least 10% of the year. In general,

initial conditions with two periods where reproduction could occur were replaced by pond regimes

with a single wet period with reproduction. Therefore, the annual fish strategy here is not adapted

to regular pond regimes with two wet periods within a year where reproduction occurs. When pond

regimes were changed iteratively such that the invasion fitness gradient for the developmental

speeds and hatching traits became minimized, the output pond regimes to which the algorithm

converged were local minima which had filling regimes with a subset of parameters modified. Figure

five shows two representative initial and final pond filling regimes. In general, the length of the

invasion fitness gradient vector would become at least halved, but vectors of zero length were not

obtained. Therefore, we did not find evolutionarily singular environments with within-year changes

and variation only. The only trait which ever became really adapted in one of the environments, was

late hatching (Fig. 5C, lower panel). Minimization of the invasion fitness sensitivity in early hatching,

a trait which contributed much to the reduction of the gradient, could co-occur with an increase in

invasion fitness sensitivities for other developmental speeds (Fig. 5C). The attractors of the

population dynamics in each of the regimes at local minima of the invasion fitness gradient, had

large maximum numbers of eggs in the egg bank and showed that many juveniles did not become

reproducing adults. Note that the output example in the first row has a larger number of eggs in the

egg bank than any of the other pond filling regimes investigated. At the start of the first wet period

in the year, the two scenarios differed in number of eggs in the egg bank by a factor of six. In all,

these simulations also found that faster developmental transitions were always favoured which

replicates the standard conclusion for "constant" environments.

For the output pond filling regime in the second row in Fig. 5, the evolutionary stability of hatching

at later rewettings β_3 was investigated and it was found to be an evolutionary repeller (Geritz et al.

1998). For the output example in the first row in Fig. 5, pond filling regimes were further optimized

as a function of the invasion fitness sensitivity on the hatching at first rewetting trait β_1 alone. A

pond filling regime was found (with ρ_8 = 0.68) where that sensitivity vanished, but it was found that

the parameter value in the data corresponded to an evolutionary repellor (Geritz et al. 1998).

Therefore, even while ESE can occur for observed hatching probabilities, these have no long-term

evolutionary stability in these environments. For values of m above 0.1 in the YS pond filling

scenarios, an exploratory analysis was carried out. It was investigated to which evolutionarily

singular value evolution in β_1 would converge, which turned out to be values which corresponded to

very low probabilities of hatching at first rewetting. When this was done for β_3 it could either

converge to higher or lower values, indicating for each m that there was an evolutionary repeller

nearby the data value for this trait. There is therefore no evidence that the intermediate hatching

probabilities are adapted to within-year pond filling variation, this analysis suggests that all-or-none

hatching probabilities would be adapted.

Discussion

For an embryonic life history strategy of annual fish (Van Dooren & Varela-Lasheras 2018), it is

investigated to which environments it might be adapted. For this, the observed strategy is

complemented with a model for the environment and the annual fish population dynamics in it. This

allows searching for Evolutionarily Singular Environments ESE where the derivatives of invasion

fitness with respect to the strategy trait parameters are all zero at the observed strategy. This is a

first condition for being an evolutionarily stable strategy ESS in a given environment, and can

therefore be used to single out environments to which the observed strategy could be adapted. The

analysis did not find an ESE among the regimes considered in which all six traits would be adapted.

Only for the two hatching probabilities, ESE pond filling regimes without any between-year

variability were found. However, the hatching probabilities did not have evolutionary stability there

and might remain some time unchanged by selection, but not so I the long term. Given model

assumptions, an environment with regular within year variation and one or two periods where

reproduction can occur does not seem to be the environment to which the embryonic rates of

development and hatching probabilities of these annual fish are adapted.

Adaptation to deterministic pond size variation within years

In a sense, within-year stochastic variability in demographic parameters is present. Random egg

deposition within a pond combined with the arrival of a wet period too short for reproduction

reduce the correlation between individual fitnesses of randomly chosen pairs of individuals (Frank &

Slatkin 1990, Starrfelt & Kokko 2012) but also increases their variance in reproductive success.

Hatching probabilities could be bet-hedging traits adapted to this within-year uncertainty, Selection

for smaller hatching probabilities at first rewetting than the observed one was advantageous in

cases where the wet period where reproduction could occur was preceded by a shorter period

during which juveniles could not mature. In one such case, by changing the extent of the pond which

was covered by water during this period, an environmental regime was found which minimized the

effect of this hatching probability on invasion fitness. In this ESE and a different one for hatching at

later rewettings, the hatching probability traits were evolutionarily unstable. The suggestion by

Pinceel et al. (2015) that observed hatching probabilities would be adapted to short-term

environmental uncertainty could not be confirmed for this system. An exploratory analysis rather

suggested that the evolutionary dynamics in a given environment would converge to a combination

of very low and high hatching probabilities and therefore no diversified bet-hedging.

In the study of adaptation to environmental variability, most attention has been given to uncertain

environments (Tuljapurkar 1990) and not to deterministic patterns of change as modelled here. The

reference environment contrasted with uncertain environments is usually called "constant" (Stearns

2000, Rose et al. 2002). If this refers to demographic parameters being invariable across years,

deterministic seasonal patterns can be implicitly present and even randomness at the individual

level. The boundaries of seasons often define the moment of census (Venable 2007). It would be

better to focus on models where the underlying demographic processes and the observation process

can be uncoupled, such that we can predict population states at any census point, to be chosen

freely within a year and keeping seasons explicit (e.g., Cooch et al. 2003). Phenological changes

which shift seasonal boundaries now occur often due to climate change (Park & Post 2022) and this

then generates between-year variability (ten Brink et al. 2020). This can occur without differences in

instantaneous rates, just by duration variation.

Not all possible within-year deterministic changes were explored, such as refilling of a pond after a

partial desiccation. This is postponed until data on competition between juveniles and adults and on

survival of different stages in wet conditions become available, which both are expected to affect

patterns of selection. When periods where a partially desiccated pond refills occur, different stage-

specific mortality rates in wet hypoxic conditions might switch the direction of selection on

development such that stages with increased mortality in wet hypoxic conditions are avoided. A

state switch such as between advancing/non-advancing states might be of use then. Note that low

rates of development in stages two and three and diapause II, claimed to be the hallmark of

annualism (Furness et al. 2015a, Furness 2016), were not observed in the data (Van Dooren &

Varela-Lasheras 2018). Assuming that this absence is adaptive, the presence of diapause I suggests

that mortality in wet hypoxic conditions might increase for this species in stages two or three where

diapause II occurs, rather than decrease (Podrabsky et al. 2007).

In some of the simulation steps aiming to minimize the invasion fitness gradient, pond regime

vectors ρ were chosen without a completely dry period. These resulted in population extinctions,

but might be useful to explore the adaptedness of developmental patterns in non-annual killifish

(Varela-Lasheras & Van Dooren 2014) and why killifish clades which evolved annualism never seem

to have lost it. On the other hand, regimes with two periods where reproduction could occur always

had larger fitness gradients than regimes with a single such period, such that no support for

adaptation to a second chance to hatch and reproduce within a year (Furness et al. 2015, Polačik et

al. 2017) was found.

Adaptation to variation between years

In models for annual plants, dormancy hedging seed germination over years could substitute for a

within-year bet-hedging trait (ten Brink et al. 2020). Therefore, bet-hedging in hatching probabilities

might still not be adaptive when years differ. A common conclusion has been that risk avoidance by

bet-hedging in one trait, would pre-empt the scope for bet-hedging in other traits (e.g. Koons et al.

2008, ten Brink et al. 2020). This is however, only partly supported (Snyder 2006). Many of these

conclusions derive from models where individuals could hatch or germinate any time within a year

or every year (Metcalf et al. 2015, Poethke et al. 2016, ten Brink et al. 2020), which is different from

how annual fish interact with their environment. In annual fish, hatching is provoked by the arrival of

water. In dry years, hatching is impossible. If long-term survival in the pre-hatching stage is reduced

relative to earlier stages, individuals should become spread out over years earlier in development

and stage-specific survival would constrain hatching to spreading individuals over within-year

occasions only.

Delaying and bet-hedging can be selected against in any constant environment or with recurring

within-year deterministic environmental changes, but become advantageous with different

population fluctuations. This could be stochastic variation, when random samples of annual pond

regimes are concatenated, or rather deterministic time series with population size fluctuations

(Bulmer 1984, Van Dooren & Metz 1998). Between-year variation in pond filling patterns would only

render delaying adaptive if it manages to dampen fluctuations in the densities of embryos in the egg

bank.

Evolutionary ecology has a long tradition of studying adaptation to variable uncertain environments

(Cohen 1966, Ellner 1997, Stearns 2000, Evans & Dennehy, 2005), i.e., environments with

unpredictable random changes. Stochastic environmental variability in pond filling regimes between

years will render changes in numbers between years random, even without stochastic variation in

instantaneous embryonic life history rates in wet and dry soil. The body of theoretical results on life

in uncertain environments then becomes applicable to this system (Tuljapurkar 1990). As this model

has explicit links between environment and demography, it would permit relating trait variation to

magnitudes of environmental uncertainty (Simons 2011, e.g., Pinceel et al. 2017) or to variances in

life history statistics or demographic parameters such as survival or fecundity (Simons 2011, e.g.,

Venable 2007). In modelling, however, the environmental variation itself has been kept relatively

implicit, and the focus has been on random 'environmental' life history variation (Rose et al. 2002,

Ellner 1997). Between different within-year pond filling patterns shown, the stationary population

size in the egg bank can differ by a factor hundred (Figs. 2 and 4) such that there seems scope for

pond filling regimes which vary between years to generate egg number fluctuations and for

adaptation to dampen these via the adjustment of embryonic life history rates.

Unfortunately, with the current trends in environmental change, it is unclear whether we can still

expect real populations to be near attractors of their eco-evolutionary dynamics for much of the

time. For example, the recently described *Austrolebias wichi* was at that time known from a single

pond in a region with large changes in agricultural practise (Alonso et al. 2018), but can be now

found in nearby man-made roadside ditches as well. Similar to the search for environments

rendering embryonic life histories adapted as implemented here, simulations of anthropogenic

disturbances and rearrangements of the natural habitats could explore which ones would render

patterns of development adapted and would minimize adaptational lags (Van Dooren 2022) across a

wider range of projected climate change patterns. Such managed novel environmental regimes

which are ESE could aid in the conservation of the several hundred annual fish species occurring in

South-America and Africa, often at risk of extinction and with currently small population sizes.

Model validation and falsification

Different types of data can be collected to test model predictions and assumptions. First of all,

snapshot distributions of embryos in the egg bank across developmental stages as in Polacik et al.

(2023) can be compared to model predictions. The model predicts on many days that a single stage

will dominate in the egg bank (Fig. 2 and Supp. Fig. 2), therefore, large sample sizes will be required

to reject the model on the basis of the relative occurrences of the less abundant stages. Predictions

can also be made for transfers of embryos between lab and field (Polacik et al. 2023) and simulations

can be used to optimize power of such experiments. Several model assumptions were made based

on the lack of more detailed data, each attempting to avoid introducing more than a minimal

amount of parameters. Lacunes in the data call for a field embryology with longitudinal studies in

time sequences of natural conditions and also for studies on competition between fish. The field

studies of Polacik et al. (2021, 2023) collected data on batches of eggs, not individuals. They could

not detect any developmental arrest in the field, but did report data which allows an estimate of

survival of embryos in wet soil which would imply that one third of them would be able to survive

one year in wet soil. I used a lower estimate of survival based on unpublished individual data, from

eggs placed in wet soil one day after they were laid. Estimates of survival in wet soil conditions in

other stages than the first are needed and a longitudinal follow-up of embryos after their first

rewetting. More data are needed on hatching probabilities after the first rewetting, to estimate age

and other effects there. Detailed data on pond filling regimes are needed as well, to investigate what

the speeds of emptying and filling actually are, how these vary within a year and what the

magnitude is of between-year variation.

Resources, rates and sizes

Searches for an annual pond filling cycle where embryonic life histories would be adapted halted at

environmental parameters where the invasion fitness gradient was at a local minimum. There,

invasion fitness sensitivities of developmental rates and hatching probabilities were smaller than in

the initial environments but never zero for rates of development. It is necessary to consider that

trade-offs might nevertheless make life histories adapted: if mortality rates and developmental rates

are coupled by trade-offs provoked by allocation of resources, negative and positive invasion fitness

changes might balance due to within-year variation alone. Rates within and between stages might

be coupled. For example, Van Dooren & Lasheras (2018) found that when parental pairs were

compared, low mortality in stage one was associated with faster development from stage 4 into the

prehatching stage. Also constraints might render the observed strategy adaptive. It would then be

on a boundary of the allowed trait space. In recent years there have been efforts to understand

whether and how plasticity and maternal effects evolve given uncertainty (de Jong & Behera 2002,

Dey et al. 2016, King & Hadfield 2019, Van Dooren 2022). Decisions on the allocation of resources

might be plastic, and the adaptiveness of amounts of plasticity was not investigated here. We know

that desiccation plasticity is limited in the annual fish studied (Van Dooren & Varela-Lasheras 2018),

such that it might be little predictive of the future. Insight in temperature plasticity is lacking. In

African *Nothobranchius* annual fish, Furness et al. (2015b) demonstrated temperature plasticity in

the entrance into diapause. However, these authors defined developmental duration as time to

hatching or death, such that separate rates of development were not estimated.

Here, competition between juvenile and adult fish was modelled according a very simple density-

dependence. It ignored size-structured competition to a large extent (e.g., de Roos et al. 2003). Size

effects can extend to the embryonic stages, if egg size and hatchling size correlate. Vrtilek et al.

(2020) found no correlation between them in *Austrolebias bellottii*, while in a different species

hatchling size depended in a hypo-allometric manner on egg size. Including more size structure

where possible would permit investigating coexistence and co-adaptation of annual fish species

(Helmstetter et al. 2020).

Eco-evo-devo modelling

Developmental life histories were embedded into a demography across the entire life cycle, bringing

us closer to an eco-evo-devo which is predictive and can start to relate environmental regimes to

adaptation. Four of the six traits modelled summarize effects of mutations across all ages on

function-valued development rates. This might not correspond to how specific genes would change

phenotypes, but at least most possible events in the embryonic life histories are included in the

assessment of adaptedness. The multi-state model used for the embryonic life history has

cumulative hazards which depends on the number of days in dry conditions, leading to a Markovian

model (de Wreede et al. 2010). This might not be the model which integrates developmental

mechanisms easily. Van Dooren & Varela-Lasheras (2018) modelled development with age-within

stage as timescales, which seems reasonable when a minimum time is required to complete each

developmental stage. With such timescales or coexisting other timescales, the only way to arrive at

predictions of distributions of individuals over states is forward simulation of all competing

processes (de Wreede et al. 2010). This is individual-based modelling and recalls conclusions of

Hogeweg and others in the classical school of bioinformatics (Hogeweg 2012), dealing with the

information exchange of organisms and their environments and the incompressibility of this process.

Modelling the dynamics of pools of physiological resources and state variables explicitly is an

alternative approach which others have implemented to keep track of processes with different

timescales (DEB modelling, Kooijman 2010), and this should be explored as well. Embryos within an

egg have well-circumscribed resource pools available to them. In the end, multiple such efforts, even

for the same empirical model, seem required to understand evolutionary constraints on

development and to be able to project adaptation and evolution as processes fuelled by mutations

forward into the future. Fortunately, for annual fish, rather mechanistic models are available for

processes surrounding diapause I (Reig et al. 2017, Montenegro-Rojas et al. 2023). However, an

approach which is customary, even in evo-devo, is to separate the processes generating phenotypic

variation from selective processes such as mortality (e.g., Salazar-Ciudad & Cano-Fernández 2023).

This should be abandoned. Individual phenotypes are not composed on a conveyor belt moving

individuals to a station where their quality is controlled. They can change the speed of the belt, they

can fall off, jump off and back on again. At any point in time there are competing risks of advancing

their phenotype and of dying which can vary between genotypes.

References

Alonso F, Terán GE, Calviño P, García I, Cardoso Y, García G. 2018. An endangered new species of

seasonal killifish of the genus *Austrolebias* (Cyprinodontiformes: Aplocheiloidei) from the Bermejo

river basin in the Western Chacoan Region. *PLoS One* 13: e0196261.

Alonso F, Terán GE, Serra Alanís WS, Calviño P, Montes MM, García ID, Barneche JA, Almirón A,

Ciotek L, Giorgis P, Casciotta J. 2023. From the mud to the tree: phylogeny of *Austrolebias* killifishes,

new generic structure and description of a new species (Cyprinodontiformes: Rivulidae), *Zoological*

Journal of the Linnean Society 2023: zlad032.

Berois N, García G, de Sá RO. 2015. *Annual fishes. Life history strategy, diversity, and evolution*. CRC

Press, Boca Raton, FL.

Bowlby J. 1973. *Attachment and Loss. Vol. 2: Separation: Anxiety and anger*. New York:

Basic Books.

Byrd RH, Lu P, Nocedal J, Zhu C. 1995. A limited memory algorithm for bound constrained

optimization. *SIAM Journal on Scientific Computing* 16: 1190-1208.

Bulmer MG. 1984. Delayed germination of seeds: Cohen's model revisited. *Theoretical Population*

Biology 26: 367-377.

Caswell H. 2019. *Sensitivity analysis: matrix methods in demography and ecology*. Springer Nature.

Charlesworth B. 1994. *Evolution in Age-Structured Populations. 2nd ed., Cambridge Studies in*

Mathematical Biology. Cambridge: Cambridge University Press.

Childs DZ, Metcalf CJE, Rees M. 2010. Evolutionary bet-hedging in the real world: empirical evidence

and challenges revealed by plants. *Proceedings of the Royal Society B: Biological Sciences* 277: 3055–

3064.

Cohen D. 1966. Optimizing reproduction in a randomly varying environment. *Journal of Theoretical*

Biology 12: 119-129,

Cooch EG, Gauthier G, Rockwell RF. 2003. Apparent differences in stochastic growth rates based on

timing of census: a cautionary note. *Ecological Modelling* 209: 133-143.

Coulson T. 2021. Environmental perturbations and transitions between ecological and evolutionary

equilibria: an eco-evolutionary feedback framework. *Peer Community Journal* 1: e6.

de Jong G, Behera, N. 2002. The influence of life-history differences on the evolution of reaction

norms. *Evolutionary Ecology Research* 4: 1-25.

de Jong IG, Haccou P, Kuipers OP. 2011. Bet hedging or not? A guide to proper classification of

microbial survival strategies. *Bioessays* 33: 215– 223.

De Roos AM, Persson L, McCauley E. 2003. The influence of size-dependent life-history traits on the

structure and dynamics of populations and communities. *Ecology Letters* 6: 473-487.

de Wreede LC, Fiocco M, Putter H. 2011. mstate: An R Package for the analysis of competing risks

and multi-state models. *Journal of Statistical Software* 38: 1– 30.

Dey S, Proulx SR, Teotónio H. 2016. Adaptation to temporally fluctuating environments by the

evolution of maternal effects. *PLoS Biology* 14: e1002388.

Dieckmann U, Heino M, Parvinen K. 2006. The adaptive dynamics of function-valued traits. *Journal of

Theoretical Biology* 241: 370–389.

Ellner S. 1997. You bet your life: life-history strategies in fluctuating environments. In: *Case Studies in

Mathematical Modeling: Ecology, Physiology, and Cell Biology*. Eds. Othmer HG, Adler FR, Lewis MA,

Dallon JC. Prentice-Hall, New Jersey, pp. 3-24.

Evans ME, Dennehy JJ. 2005. Germ banking: bet-hedging and variable release from egg and seed

dormancy. *Quarterly Review of Biology* 80: 431-51.

Frank S, Slatkin M. 1990. Evolution in a Variable Environment. *The American Naturalist* 136: 244–

260.

Furness AI. 2016. The evolution of an annual life cycle in killifish: adaptation to ephemeral aquatic

environments through embryonic diapause. *Biological Reviews* 91: 796-812.

Furness AI, Lee K, Reznick DN. 2015a. Adaptation in a variable environment: Phenotypic plasticity

and bet-hedging during egg diapause and hatching in an annual killifish. *Evolution* 69: 1461– 1475.

Furness AI, Reznick DN, Springer MS, Meredith RW. 2015b. Convergent evolution of alternative

developmental trajectories associated with diapause in African and South American killifish.

Proceedings of the Royal Society B: Biological Sciences 282: 20142189.

Geritz SAH, Meszéna G, Metz JAJ. 1998. Evolutionarily singular strategies and the adaptive growth

and branching of the evolutionary tree. *Evolutionary Ecology* 12: 35-57.

Helmstetter AJ, Van Dooren TJM. 2019. Determinants of growth and body size in *Austrolebias* South-

American annual killifish. *BioRxiv* 2019.12.31.891648. https://doi.org/10.1101/2019.12.31.891648

Helmstetter AJ, Papadopulos AS, Igea J, Van Dooren TJM, Leroi AM, Savolainen V. 2016. Viviparity

stimulates diversification in an order of fish. *Nature Communications* 7: 11271.

Helmstetter AJ, Papadopulos, AST, Igea, J, Van Dooren, TJM. 2020. Trait evolution and historical

biogeography shape assemblages of annual killifish. *Journal of Biogeography* 47: 1955– 1965.

Hogeweg P. 2012. Toward a theory of multilevel evolution: long-term information integration shapes

the mutational landscape and enhances evolvability. *Advances in Experimental Medicine and Biology*

751: 195-224.

Irons W. 1998. Adaptively relevant environments versus the environment of evolutionary

adaptedness. *Evolutionary Anthropology* 6: 194-204.

King JG, Hadfield JD. 2019. The evolution of phenotypic plasticity when environments fluctuate in

time and space. *Evolution Letters* 3: 15–27.

Kirkpatrick M, Heckman N. 1989. A quantitative genetic model for growth, shape, reaction norms,

and other infinite-dimensional characters. *Journal of Mathematical Biology* 27: 429-50.

Kooijman SALM. 2010. *Dynamic Energy Budget Theory for Metabolic Organisation*. Cambridge

University Press, Cambridge.

Liu RK, Walford RL. 1970. Observations on the lifespans of several species of annual fishes and of the

world's smallest fishes. *Experimental Gerontology* 5: 241-6.

Koons D, Metcalf C, Tuljapurkar S. 2008. Evolution of delayed reproduction in uncertain

environments: a life-history perspective. *The American Naturalist* 172: 797–805.

Metcalf CJE, Burghardt LT, Koons DN. 2015. Avoiding the crowds: the evolution of plastic responses

to seasonal cues in a density-dependent world. *Journal of Ecology* 103: 819-828.

Metz JAJ, Geritz SAH, Meszena G, Jacobs FJA, van Heerwaarden JS. 1996. Adaptive dynamics: A

geometrical study of the consequences of nearly faithful reproduction. In*: Stochastic and Spatial*

Structures of Dynamical Systems. Eds. Van Strien SJ, Verduyn Lunel SM. North-Holland, Amsterdam,

pp. 183-231.

Metz JAJ, Mylius SD, Diekmann O. 2008. When does evolution optimize? *Evolutionary Ecology*

Research 10: 629–654

Montenegro-Rojas I, Yañez G, Skog E, Guerrero-Calvo O, Andaur-Lobos M, Dolfi L, Cellerino A, Cerda

M, Concha ML, Bertocchi C, Rojas NO, Ravasio Andrea, Rudge TJ. 2023. A computational framework

for testing hypotheses of the minimal mechanical requirements for cell aggregation using early

annual killifish embryogenesis as a model. *Frontiers in Cell and Developmental Biology* 11: 959611.

Park JS, Post E. 2022. Seasonal timing on a cyclical Earth: Towards a theoretical framework for the

evolution of phenology. *PLoS Biology* 27: e3001952.

Pinceel T, Vanschoenwinkel B, Deckers P, Grégoir A, Ver Eecke T, Brendonck L. 2015. Early and late

developmental arrest as complementary embryonic bet-hedging strategies in African killifish.

Biological Journal of Linnean Society 114: 941–948.

Pinceel, T., Vanschoenwinkel, B., Hawinkel, W. Tuyens K, Brendonck L. 2017. Aridity promotes bet

hedging via delayed hatching: a case study with two temporary pond crustaceans along a latitudinal

gradient. *Oecologia* 184: 161–170.

Podrabsky JE, Lopez JP, Fan TWM, Higashi R, Somero GN. 2007. Extreme anoxia tolerance in embryos

of the annual killifish *Austrofundulus limnaeus*: insights from a metabolomics analysis. Journal of

Experimental Biology 210: 2253–2266.

Poethke HJ, Hovestadt T, Mitesser O. 2016. The evolution of optimal emergence times: bet hedging

and the quest for an ideal free temporal distribution of individuals. *Oikos* 125: 1647-1656.

Polačik M, Vrtílek M. 2023. Cryptic stasis during the development of Nothobranchius furzeri suggests

new stages of dormancy outside of the typical three diapauses of annual killifishes. *Environmental*

Biology of Fishes 106: 575-583.

Polačik M, Smith C, Reichard M. 2017. Maternal source of variability in the embryo development of

an annual killifish. *Journal of Evolutionary Biology* 30: 738– 749.

Polačik M, Vrtílek M, Reichard M, Žák J, Blažek R, Podrabsky J. 2021. Embryo ecology: developmental

synchrony and asynchrony in the embryonic development of wild annual fish populations. *Ecology*

and Evolution 11: 4945-4956.

Polačik M., García D., Arezo MJ, Papa N, Schlueb H, Blanco D, Podrabsky JE, Vrtílek M. 2023.

Embryonic development of natural annual killifish populations of the genus *Austrolebias*:

Evolutionary parallelism and the role of environment. *Freshwater Biology* 00: 1–13

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/

Reeve HK, Sherman PW. 1993. Adaptation and the goals of evolutionary research. *The Quarterly*

Review of Biology 68: 1–32.

Reig G, Cerda M, Sepúlveda N, Flores D, Castaneda V, Tada M, Härtel S, Concha ML. 2017. Extra-

embryonic tissue spreading directs early embryo morphogenesis in killifish. *Nature communications*

8: 15431.

Rose KE, Rees M, Grubb PJ. 2002. Evolution in the real world: stochastic variation and the

determinants of fitness in *Carlina vulgaris*. *Evolution* 56: 1416–1430.

Salazar-Ciudad I, Cano-Fernández H. 2023. Evo-devo beyond development: Generalizing evo-devo to

all levels of the phenotypic evolution. *Bioessays* 45: 20232200205.

Simons AM. 2011. Modes of response to environmental change and the elusive empirical evidence

for bet hedging. *Proceedings of the Royal Society B: Biological Sciences* 278: 1601-9.

Snyder RE. 2006. Multiple risk reduction mechanisms: can dormancy substitute for dispersal?

Ecology Letters 9: 1106-1114.

Southwood TRE. 1977. Habitat, the templet for ecological strategies? *Journal of Animal Ecology 46:*

337–365.

Starrfelt J, Kokko H. 2012. Bet-hedging--a triple trade-off between means, variances and

correlations. *Biological reviews of the Cambridge Philosophical Society* 87: 742-55.

Stearns S. 2000. Life history evolution: successes, limitations, and prospects. *Naturwissenschaften*

87: 476–486.

Steindachner F. 1881. Beiträge zur Kenntniss der Flussfische Südamerika's (III) und Ichthyologische

Beiträge (XI). *Anzeiger der Akademie der Wissenschaften in Wien, Mathematisch-*

Naturwissenschaftliche Classe 18: 97– 100.

ten Brink H, Gremer JR, Kokko H. 2020. Optimal germination timing in unpredictable environments:

the importance of dormancy for both among- and within-season variation. *Ecology Letters* 23: 620-

630.

Thompson AW, Black AC, Huang Y, Shi Q, Furness AI, Braasch I, Hoffmann FG, Ortí G. 2021.

Deterministic shifts in molecular evolution correlate with convergence to annualism in killifishes.

BioRxiv 2021.08.09.455723. https://doi.org/10.1101/2021.08.09.455723

Tuljapurkar S. 1990. *Population Dynamics in Variable Environments*. Lecture Notes in

Biomathematics 85, Springer Verlag.

Urban MC, Swaegers J, Stoks R, Snook RR, Otto SP, Noble DWA, Moiron M, Hällfors MH, Gómez-

Llano M, Fior S, Cote J, Charmantier A, Bestion E, Berger D, Baur J, Alexander JM, Saastamoinen M,

Edelsparre AH, Teplitsky C. 2023. When and how can we predict adaptive responses to climate

change? Evolution Letters 29: 172-187.

Van Dooren TJM. 2022. Adaptational lags during periods of environmental change. *BioRxiv*:

742916v2 https://doi.org/10.1101/742916

Van Dooren TJM, Metz JAJ. 1998. Delayed maturation in temporally structured populations with

non-equilibrium dynamics. *Journal of Evolutionary Biology* 11: 41-62.

Van Dooren TJM, Varela-Lasheras I. 2018. Embryonal life histories: Desiccation plasticity and

diapause in the Argentinean pearlfish *Austrolebias bellottii*. *Ecology & Evolution* 8: 11246– 11260.

van Goor HA. 2009. Innovative Design: A killifish embryo's life behind glass. Unpublished Msc thesis,

Leiden University, The Netherlands.

Varela-Lasheras I, Van Dooren TJM. 2014. Desiccation plasticity in the embryonic life histories of

non-annual rivulid species. *Evodevo* 29: 5-16.

Venable DL. 2007. Bet Hedging in a Guild of Desert Annuals. *Ecology* 88: 1086–1090.

Volcan MV, Krause Lanés LE. 2018. Brazilian killifishes risk extinction. *Science* 361: 340-341.

Vrtílek M, Van Dooren TJM, Beaudard M. 2020. Egg size does not universally predict embryonic

resources and hatchling size across annual killifish species. *Comparative Biochemistry and Physiology*

Part A: Molecular & Integrative Physiology 249: 110769.

Wourms JP. 1972a. Developmental biology of annual fishes. I. Stages in the normal development of

Austrofundulus myersi Dahl. *Journal of Experimental Zoology* 182: 143– 167.

Žák J, Reichard M. 2020. Fluctuating temperatures extend median lifespan, improve reproduction

and reduce growth in turquoise killifish. *Experimental Gerontology* 140: 111073.

Zhan J, Thrall PH, Papaïx J, Xie L, Burdon JJ. 2015. Playing on a pathogen's weakness: using evolution

to guide sustainable plant disease control strategies. *Annual Reviews Phytopathology* 53: 19-43.

Figure Legends

Figure one. Schematic depiction of the annual killifish life history interacting with the pond
environment.

Above. Main processes over time in water and soil compartments and exchanges between them are
listed. The egg bank is located in the soil compartment and fish in the water compartment. Dry and
wet periods alternate: during a wet period at least part of the pond is filled (blue). Embryos in the
soil egg bank cay be in dry (pale drab) or wet conditions (drab), depending on the extent of the pond
covered by water. Individuals can hatch from stage five embryos when water reaches the location
where they are buried (thick black line), survive and mature into adults. Reproducing adults
contribute eggs to the soil egg bank at locations where water is still covering the soil. Below. A
depiction of the developmental strategy modelled. Five developmental stages are considered, each
with stage-specific mortality and development rates in dry conditions. Stage five embryos can hatch
according a sub-model for hatching probabilities. Stages four and five are easy to discern due to
their pigmented eyes. In stage two and three the body axis can be seen already, in stage three with a
clear head (white arrows). Photographs of embryos reproduced from Van Dooren & Varela-Lasheras
(2018) with permission.

Figure two. Population dynamics of embryos and fish from day to day in different pond filling
regimes.

The environmental pond filling regimes are shown above each panel, with the pond radius varying
between zero and r_{max} = 10. The population dynamics was first iterated for twenty years, then it was
checked that it had become stationary. The stationary result was iterated from day to day over one
year and the numbers of individuals in different states are plotted. The numbers of individual embryos
in the egg bank are plotted (increasing downward) and fish numbers (increasing upward). Note that
fish and embryos are drawn with different scales. The bars are as high for one fish as for ten embryos.
(A) Effects of changes in the speed of pond filling within a wet period. From left to right, the scenario
with accelerating filling YA3, the reference scenario Y0 and the scenario with decelerating filling YD3.
(B) Effects of the main wet period where reproduction can occur being preceded by a short wet period
which does not allow reproduction. From left to right, scenarios YS1, YS3 and YS5 where this short
period covers m = 0.2, 0.6, and the full extent of the pond. Note the additional differences in scale
between rows (A) and (B).
836

Figure three. Invasion fitnesses of phenotypic mutational changes in each of the embryonic life history
traits, in the simplest pond filling regime Y0.

Invasion fitnesses are shown for invasions of mutant phenotypes in a resident population which has
become stationary in scenario Y0. The resident trait value for each trait is at value one. Invasion
fitnesses are estimated at five equidistant points and connected by a line.

Figure four. Sensitivities of invasion fitness to pond regime changes seem limited for many traits.

Patterns of invasion fitness for a fixed difference between mutant phenotype and resident phenotype
are shown. Invasion fitnesses of mutants increasing cumulative hazards and logistic regression
parameters with a multiplication factor of 1.1 plotted as a function of an environmental parameter.
This gives a qualitative impression of strengths of selection in different environmental regimes on each
of the traits. Labels of rates next to panels are ranked according values. (A) Decelerating and
accelerating pond filling. At zero on the x-axis, the pond filling regime is the one of the middle panel
in row A of Figure two. (B) Short wet periods with increasing fractions m of the egg bank covered. For
hatching at first rewetting, it was determined in a stepwise manner for which value of m the sign of
the sensitivity changed. The change was abrupt.

Figure five. Minimization of invasion fitness gradients by changing pond filling regimes results in
limited reductions of selection strengths.

Results of optimizing environmental pond filling regimes such that the data-based strategy becomes
as adapted as possible, by a parameter search. (A) Two initial pond filling regimes from which
optimization started (purple), the upper one with two wet periods, the lower profile with a single one.
Resulting pond filling regimes after minimization of the invasion fitness gradient are shown in black.
(B) Attractors of the resident population dynamics in each of the two pond regimes to which the
algorithm converged. Note that fish and embryos are drawn with different scales. The bars are as high
for one fish as for ten embryos. Both rows additionally have different scales. The population sizes are
much larger for the first row. (C) The patterns of invasion fitnesses per trait in the regimes obtained
after minimization. The invasion fitnesses in the initial pond regimes are shown as hairlines for
comparison. Traits with very similar invasion fitness patterns are given the same colour. Labels of rates
next to panels are ranked according their values. It can be seen that for some traits, the invasion fitness
slope increased while the multi-trait fitness gradient became minimized.

870
871

872   Table 1. Model parameters fixed across simulations

Model Parameter	Compartments	Value	Description
r_{max}	All	10	Maximum pond radius
$e^{-\mu_w}$	Egg bank (wet location)	0.975	Probability of survival per day[#]
β_1	Egg bank (wet location)	2.693	Hatching parameter first rewetting[*]
β_2	Egg bank (wet location)	-0.034	Hatching parameter age effect[*]
β_3	Egg bank (wet location)	-0.080	Hatching parameter later rewettings[*]
p_d	Egg bank (dry location)	0.498	Proportion arresting after rewetting
$e^{-\mu_d}$	Egg bank (dry location)	0.994	Survival per day, those arrested after rewetting
f	Water body	10	Female fecundity per day
$e^{-\mu_J}$	Water body	0.989	Juvenile survival probability per day[$]
a	Water body	60	Age at maturity (days)
K	Water body	10	Maximum density of adults per m^2
S	Water body	5	Maximum number of adults of one sex relative to the other allowing successful reproduction

#: according Van Goor 2008.

*: according Van Dooren & Varela-Lasheras 2018.

$: according Helmstetter & Van Dooren 2020.

Supplementary Material

Embryonic life histories in annual killifish: adapted to what?

Transition probabilities

Cumulative hazards are measures of the summed probabilities across all observation times that

individuals have made a specific transition by a specific time (de Wreede et al. 2011) and are

represented using time-dependent matrices $A(a)$, here of age a, with a ranging from zero to the

maximum age observed/considered. $A(0)$ equals zero. Each entry A_{ij}, with i denoting the row and j

the column of each matrix element, represents the cumulative hazard at that time for the transition

from state i to state j. Note that this is different from standard practise in matrix population

dynamics models, where columns denote departure states and rows arrival states (Caswell 2000). Six

states can occur in an embryonic life history and are indexed as the five developmental stages ($i, j =$

1…5) and the dead state (i or j = 6). All sub-diagonal elements A_{ij} are zero. In a statistical analysis,

non-parametric estimates of matrix function A can only change at ages where individuals have been

observed. If we give observation times a_τ an index τ, then the estimated changes in matrices A at

ages a_τ are given by matrices $dA(a_\tau)$. Transition probabilities from age a_1 to a_2 are calculated as

(Eqn. 1, de Wreede et al. 2011)

$$\boldsymbol{P}_{a_1,a_2} = \prod_{a_\tau\ in\ (a_1,a_2]}(\boldsymbol{I} + d\boldsymbol{A}(a_\tau)) \hspace{4cm} \text{(Eqn. A.1.1)}$$

where I use interval limits as in de Wreede et al. (2011). Using the data-based transition probability

matrices (Eqn.1), a set of matrices $\boldsymbol{P}_{a,a+1}$ is constructed projecting numbers of embryos in different

stages and sharing their age state variable one day forward, for any age $a \in [0, \infty[$. For all ages

above the maximum a_{max} observed (which was 169 days), it is assumed that $\boldsymbol{P}$ remains equal to

$\boldsymbol{P}_{a_{max-1},a_{max}}$. To project numbers of embryos forward in time, we only need to project the numbers

of individuals alive. Therefore the 5 by 5 minorized matrices M_{66} are used, with the sixth row and

column of P removed. The life history transitions in this environmental compartment imply that

embryos need, next to their location, their age and stage as state variables. However, as it is only

age given in the number of days in dry conditions which is assumed to determine developmental

progress, we use this number as the individual age variable.

Population Dynamics

The equations are given for the population dynamics of a single mutant genotype and phenotype

across wet and dry periods. The equations include the feedback from the resident population. It is

explained how to obtain the population dynamics of individuals with the resident genotype and

phenotype using similar equations. In principle, development and mortality are processes occurring

or modelled on a continuous timescale, but the periodicity in environmental conditions within a day

is strong (Jacobs et al. 2008) and a motivation to use days as the discrete time step to carry out the

modelling instead.

Embryos have different individual state variables. Numbers of embryos N_{ijkl} are indexed i for stage (i =

1...5), j for their location, i.e., their distance from the pond center in meter ($j \geq 0$), k for the first or

later rewetting variable (hatching state k = 1, 0) and l for the advancing (l = 1) and non-advancing

states (l = 0). Embryos are not differentiated according their sex.

A location j is dry when $r_{pond}(t) < j$ or $r_{pond}(t) = 0$. In dry conditions at their location on day t,

embryos are projected forward into the next day according

$$N_{j,k,1}(a + 1, t + 1) = M_{66}^T(a)N_{j,k,1}(a, t)$$ (Eqn. A2.1a)

$$N_{i,j,k,0}(a + 1, t + 1) = e^{-\mu_d}N_{i,j,k,0}(a, t)$$ (Eqn. A2.1b)

where $\boldsymbol{N}_{j,k,1}$ is the vector of numbers $N_{i,j,k,1}$ of embryos in the five developmental stages. A location

j is wet when $r_{pond}(t) \geq j$. On the first day of wet conditions at a location j, embryos in the pre-

hatching stage can hatch. To determine the number of individuals hatching $H_{j,s}(0,t)$ at time t, with

sex s and with their age as fish zero, from embryos of all ages and per location j receiving water, we

need to use the hatching probability functions and account for the fact that half of the individuals

are assumed to be females, half males. Index s equals f for females and m for males. When a

location is already wet for at least one day $H_{j,s}(0,t)$ is zero. For a category j which is wetted

$H_{j,s}(0,t) = \frac{1}{2}\sum_{a,l} g(\beta_1 + \beta_2 a)N_{5,j,1,l}(a,t) + g(\beta_3)\,N_{5,j,0,l}(a,t)$ (Eqn. A2.2)

The individuals which did not hatch at j are

$N_{5,j,0,l}(0,t)' = \sum_{a,l}(1 - g(\beta_1 + \beta_2 a))N_{5,j,1,l}(a,t) + (1 - g(\beta_3))N_{5,j,0,l}(a,t)$ (Eqn. A2.3)

Since these individuals all experienced a rewetting now, the index k is set to zero for all of them and

$N_{5,j,1,l}(0,t)'$ equals zero for both l. Then on the same day, the advancing and non-advancing states

are reset and the age state variable, the number of days in dry conditions, is reset to zero. For stages

one to four this yields

$N_{i,j,k,0}(0,t)'' = p_d \sum_{a,l} N_{i,j,k,l}(a,t)$ (Eqn. A2.4.a)

$N_{i,j,k,1}(0,t)'' = (1 - p_d) \sum_{a,l} N_{i,j,k,l}(a,t)$ (Eqn. A2.4.b)

with $N_{i,j,k,l}(0,t)''$ the number after resetting at location j. For stage five, this occurs in the

individuals remaining in the egg bank after they decided not to hatch

$$N_{5,j,0,0}(0,t)'' = p_d\big(N_{5,j,0}(0,t)' + N_{5,j,1}(0,t)'\big)$$ (Eqn. A2.5.a)

$$N_{5,j,0,1}(0,t)'' = (1 - p_d)\big(N_{5,j,0}(0,t)' + N_{5,j,1}(0,t)'\big)$$ (Eqn. A2.5.b)

The reset individuals N'' are then advancing to the next day according the survival probability in wet

conditions

$$N_{i,j,k,l}(0,t+1) = e^{-\mu_w}N_{i,j,k,l}(0,t)''$$ (Eqn. A2.6)

and on subsequent wet days at location j, individuals are updated according

$$N_{i,j,k,l}(0,t+1) = e^{-\mu_w}N_{i,j,k,l}(0,t)$$ (Eqn. A2.7)

Fish can be present when $r_{pond}(t) \geq 0$. The fish population is initiated from hatchlings.

The number of juvenile fish of age one are the summed individuals that decided to hatch on the

previous day. For older ages (a below α), they are the juveniles which survived from the previous

day, including the effects of density dependence from the resident population

$$N_{J,s}(1,t) = \sum_j H_{j,s}(0,t-1)$$ (Eqn. A2.8.a)

$$N_{J,s}(a,t) = e^{-\mu_J} \left[\frac{KS_{pond}(t-1) - N_A^R(t-1)}{KS_{pond}(t-1)} \right]_{\geq 0} N_{J,s}(a-1, t-1)$$ (Eqn. A2.8.b)

where $N_A^R(t-1)$ is the number of adult resident individuals present at the start of day $t-1$ and

$S_{pond}(t-1)$ the pond wet surface area on that day. Factor $\left[\frac{KS_{pond}(t-1) - N_A^R(t-1)}{KS_{pond}(t-1)} \right]_{\geq 0}$ is non-negative.

Juveniles become adults at age α. Adults only die because of density dependence, or when the pond

radius has become zero.

$$N_{A,s}(a,t) = \left[\frac{KS_{pond}(t-1)}{N_A^R(t-1)} \right]_{\leq 1} N_{A,s}(a-1, t-1)$$ (Eqn. A2.9)

The density dependent factor $\left[\frac{KS_{pond}(t-1)}{N_A^R(t-1)} \right]_{\leq 1}$ cannot be larger than one. Adult females which will

survive until the next day t produce new eggs every day throughout their lives, which are fertilized

by males, which will also survive until the next day. Reproduction requires that resident individuals

of both sexes are present and it is also set at zero for very biased resident sex ratios (ratio resident

females to males outside of the interval $[S^{-1}, S]$). If all conditions are fulfilled, the number of new

eggs with a mutant genotype appearing on day t equals

$$M(0,t) = \frac{1}{2} f \sum_a N_{A,f}(a,t) + \frac{1}{2} \frac{\sum_a N_{A,m}(a,t)}{\sum_a N_{A,m}^R(a,t)} f \sum_a N_{A,f}^R(a,t)$$ (Eqn. A2.10)

where the factors $\frac{1}{2}$ account for the fact that mutant individuals are heterozygotes and only transfer

the mutant allele to half of their offspring. $N_{A,m}^R(a,t)$ is the resident number of adult males and

$N_{A,f}^R(a,t)$ the number of resident adult females of a specific age present at time t. New eggs and

their embryos have experienced zero dry days, they are put in wet conditions in the soil. These eggs

are distributed at random over the pond bottom which adults can reach. This implies that location

category j receives eggs proportional to its surface (which is $\pi(2j+1)$), giving the $M_j(0,t)$ per

location category.

On the first day with dry conditions at location j, all new eggs deposited in that location in the

preceding wet period $M_j(0,t)$ are added to the stage one embryos from older wet periods. The

buffer with new eggs is emptied.

1000

1001 $N_{1,j,1,1}(0,t) = e^{-\mu_w}N_{i,j,1,1}(0,t-1) + \sum_a e^{-\mu_w a}M_j(0,t-a)$ (Eqn. A2.11)

1002

Resident individuals have the homozygous genotype and the phenotype of a homozygote. To

determine their numbers we can use mostly the same equations as above where resident individuals

are now counted. Only in the number of new eggs produced (Eqn. A2.10), a modification has to be

made to make it represent the number of new eggs with resident genotype. Only the route via

females needs to be accounted for and resident genotypes are transmitted with probability one, not

one half as all males will contribute the resident allele when mutants are rare.

1009

1010   Mutational effects on transition probabilities

1011   Some more detail is required to understand what a mutation change in one of the rates will actually

1012   change to a transition matrix. If we make the elements of a minorized transition matrix $\boldsymbol{M}_{66}$ from one

1013   day to the next explicit, we can write that matrix as consisting of transition probabilities $p_{i \to j}$ from

1014   state i to state j, where each state is a developmental stage (i, j in 1,…, 5).

1015

1016 $$M_{66} = \begin{matrix} p_{1\to1} & p_{1\to2} & p_{1\to3} & p_{1\to4} & p_{1\to5} \\ 0 & p_{2\to2} & p_{2\to3} & p_{2\to4} & p_{2\to5} \\ 0 & 0 & p_{3\to3} & p_{3\to4} & p_{3\to5} \\ 0 & 0 & 0 & p_{4\to4} & p_{4\to5} \\ 0 & 0 & 0 & 0 & p_{5\to5} \end{matrix}$$ (Eqn. A.3.1)

1017

Transitions $p_{i \to dead}$ can be inferred from the other transitions as $p_{i \to dead} = 1 - \sum_{j \neq dead} p_{i \to j}$.

Diagonal elements in M_{66} can be decomposed as $p_{i \to i} = 1 - \sum_{j \neq i} p_{i \to j} - p_{i \to dead}$. The probabilities

of developmental transitions are not conditional on not dying. All transitions are treated on an equal

footing. One needs to be careful when changing hazards to keep all elements within the bounds [0,

1] and to preserve the sum constraint on the probabilities out of a state. Therefore, when for a

particular day, no events have been observed in the data (due to lack of events or of observations),

the matrix for that day is not changed by trait changes. Also, when one of the probabilities on a row

is one or the diagonal is zero, trait changes in the involved cumulative hazards are not affecting that

row. Only diagonal elements $p_{i \to i}$ of M_{66} (Eqn. A.3.1) with values within the interval]0, 1[are

affected by trait changes to the cumulative hazards for death rates. It is assumed that trait changes

in stage-specific rates of development, affect all cumulative hazards for changes out of that stage

equally. For example, a multiplicative increase c in the rate of development out of stage one affects

four cumulative hazards, changing all off-diagonal probabilities in]0, 1[in the first row of M_{66} (Eqn.

A.3.1) with a factor c and therefore reduces the diagonal with $(c - 1) \sum_{j=2,3,4,5} p_{1 \to j}$. Trait changes

studied were limited to the interval [0.9, 1.1] for which it was checked that all probabilities in

transition matrices M_{66} remain between zero and one.

1034

Invasion fitness

To determine invasion fitness and adaptedness, it is necessary to assume stationarity of the

environment in the long run (Metz et al. 1992, Van Dooren 2022). If we want to assess whether

organisms which are prevalent in a given environment are adapted to it, then we should determine

in the first place that organisms with genotypes determining different but similar phenotypic

strategies would not be able to increase in frequency when they would appear. This is the idea of

adaptation bringing populations to a final stop of phenotypic and genetic evolution (Hammerstein

1996), which became formalized in adaptive dynamics approximations based on invasion fitness

(Metz et al. 1992, Metz et al. 1996, Geritz et al. 1998). At such endpoints of evolution, invasion

fitnesses of rare (mutant) phenotypic strategies slightly differing from the prevailing or so-called

resident strategy should not indicate that these can invade. At adapted population states, the

invasion fitness as a function of mutational changes is at an optimum and the first condition for that

is that the linear approximation of such a function equals zero for small mutational changes, i.e., all

partial derivatives (slopes) of invasion fitness with respect to a mutational change in a single trait are

zero (Metz et al. 1996). When invasion fitness is estimated for mutants appearing in a resident

population with stationary but non-constant numbers or densities, it is best to calculate initial and

final numbers of mutants over a period which is a multiple of the periodicity in the resident

population dynamics. Respecting this periodicity minimizes errors in estimation (Van Dooren & Metz

1998). All pond regimes investigated have a duration after which the same pattern of wet and dry

periods repeats itself, and the resident attractors follow this periodicity. Invasion fitnesses were

calculated from numbers at the start and end of a multiple of the total duration of the resident

pattern.

1057

1058 As further supplementary material, two figures are provided.

Supplementary figure one. Parameterization of pond filling time profiles.

Radiuses $r_{pond}(t)$ are piecewise linear continuous functions (when time is given as a real number). Differences between a large number of piecewise linear pond filling profiles can be specified by a change in one or several of ten parameters. These parameters, all constrained between zero and one, are involved in the specification of each pond filling regime, for a fixed parameter r_{max}, the maximum radius the pond can have within a year. As they are all proportions, they are indicated in the figure by bars, which show the intervals on which each parameter works, and by arrows, showing the actual value they took to produce the pond profile shown. Parameter ρ_1 is the proportion of the entire year taken up by a first pond filling and drying period. Parameter ρ_2 is the proportion of that period which is wet. Parameter ρ_3 if the proportion of that wet period at which

maximum radius is reached. Parameter ρ_4 is the proportion of the first wet period where the radius
increases, at which the speed of filling changes. Parameters ρ_5 to ρ_7 are the corresponding
parameters for the second pond filling and drying period. Parameter ρ_8 is the proportion of r_{max}
which is the maximum reached within the first period. Parameter ρ_9 determines the radius at which
the speed of filling changes in the first wet period and parameter ρ_{10} determines this radius for the
second wet period.

1076

1077

1078

1079

Supplementary figure two. Distributions of embryos in different stages at different time points
within a year.

Distributions over stages at the end of each period in a pond filling regime drawn above the stacked
barcharts are shown, as a function of distance to the pond center. The pond filling regime has a
short period where reproductions cannot occur and a main longer one where new eggs are added to
the egg bank. By the end of the wet period, on day 300, all eggs at nine to ten meter from the pond
center have been dry since day 212.

1087

1088

1089   References

1090

Caswell H. 2000. *Matrix population models*. Sinauer, Sunderland, MA.

Hammerstein P. 1996. Darwinian adaptation, population genetics and the streetcar theory of

evolution. *Journal of Mathematical Biology* 34: 511–532.

Jacobs AF, Heusinkveld BG, Kraai A, Paaijmans KP. 2008. Diurnal temperature fluctuations in an

artificial small shallow water body. *International Journal of Biometeorology* 52: 271-80.

Metz JAJ, Nisbet RM, Geritz SAH. 1992. How should we define 'fitness' for general ecological

scenarios? *Trends in Ecology and Evolution* 7: 198-202.

1098

1099

1100
1101

1102

1103

1104