Correlative single-cell hard X-ray tomography and X-ray fluorescence imaging

Zihan Lin[1,6], Xiao Zhang[1,6], Purbasha Nandi[1], Yuewei Lin[2], Liguo Wang[3], Yong Chu[4], Timothy Paape[1,5], Yang Yang[4,*], Xianghui Xiao[4,*], Qun Liu[1,4,*]

[1]Biology Department, [2]Computational Science Initiative, [3]LBMS, [4]NSLS-II, Brookhaven National Laboratory, Upton, New York, USA. [5]U.S. Department of Agriculture's Agricultural Research Service at Children's Nutrition Research Center, Houston, Texas, USA.

[6]These authors contributed equally to the work: Z. L., X. Z.

*Correspondence to: Y.Y.: yyang@bnl.gov, X.X.: xiao@bnl.gov, or Q.L.: qunliu@bnl.gov

Abstract:

X-ray tomography (XRT) and X-ray fluorescence (XRF) imaging are two non-invasive imaging techniques to study cellular structures and chemical element distributions, respectively. However, correlative X-ray tomography and fluorescence imaging for the same cell have yet to be routinely realized due to challenges in sample preparation and X-ray radiation damage. Here we report an integrated experimental and computational workflow for achieving correlative multi-modality X-ray imaging of a single cell. The method consists of the preparation of radiation-resistant single-cell samples using live-cell imaging-assisted chemical fixation and freeze-drying procedures, targeting and labeling cells for correlative XRT and XRF measurement, and computational reconstruction of the correlative and multi-modality images. With XRT, cellular structures including the overall structure and intracellular organelles are visualized, while XRF imaging reveals the distribution of multiple chemical elements within the same cell. Our correlative method demonstrates the feasibility and broad applicability of using X-rays to understand cellular structures and the roles of chemical elements and related proteins in signaling and other biological processes.

Main:

Cells are the building blocks of living organisms. X-rays have deep-penetration power and are ideal for non-invasive imaging of cells at nanometer resolutions [1,2]. Transmission X-ray microscopy tomography (XRT) and X-ray scanning fluorescence (XRF) imaging are two nanometer-resolution imaging techniques capable of studying cellular structure and function in high spatial resolution. XRT is analog in its configuration to optical bright-field microscopy, utilizing a condenser to focus X-rays onto the sample and employing a zone plate as an X-ray objective lens to magnify images for full-field tomographic imaging [3,4]. Meanwhile, XRF imaging operates as a spectroscopic technique based on scanning, utilizing element-specific X-ray fluorescence spectra to map multiple chemical elements [5-7]. The two methods complement each other, where XRT provides the overall cellular and subcellular structures while XRF imaging generates elemental maps crucial for understanding element-related uptake, metabolism, and cellular localization.

Both X-ray imaging modalities have found extensive applications in biological research [3,5,8]. XRT utilizes either absorption or phase contrast for imaging biological cells. Soft X-rays at energy within the water window (183-530 eV) exhibit a larger mass absorption cross-section difference for carbon and nitrogen relative to water, thus creating absorption contrast for tomographic imaging of hydrated cells [1,9]. However, these soft X-rays are also associated with high parasitic X-ray absorption and radiation damage, necessitating the use of cryofixation and data collection at cryogenic temperatures [9]. Soft X-rays are also limited to a short focal length and samples thinner than 20 μm and thus, large cells such as some plant and animal cells may be challenging for soft X-ray tomography. In comparison, hard X-rays with an energy of 5 keV or higher possess a smaller cross-section but generate much less radiation damage while penetrating to great deeper, e.g. 1 mm for 10 keV X-rays [10]. Hard X-rays have been utilized to study cellular structures of unstained MC3 preosteoblasts [11] and fission yeast cells [12]. Due to the small cross-section difference between carbon, nitrogen, and water, the absorption contrast is very low for imaging biological cells. Hence, previous researchers preferred to use phase contrast mechanisms by a Zernike phase plate to create a phase delay

between absorbed and scattered X-rays [11,12]. To enhance the contrast for hard X-ray

imaging, staining cells using heavy metals such as osmium, lead, and uranyl is

generally required, but such a method limited the resolution and caused stain-related

artifacts [12].

XRF is highly sensitive and capable of detecting elemental distributions in a sample at

sub-ppm (Parts Per Million) levels. Unlike other methods, XRF does not require the use

of external markers or tags, enabling simultaneous detection of multiple elements with

exceptional resolution using a highly focused X-ray beam [5,6]. This characteristic makes

XRF suitable to probe and differentiate compositions of both metal and non-metal

elements inside cells, thereby aiding in the understanding of their roles in signaling,

homeostasis, and other cellular functions. As hard X-rays can detect biological elements

with an atomic number of 15 (phosphorus) or higher and are insensitive to water, both

hydrated and dehydrated samples have been employed for XRF [13-15]. XRF has been

utilized to study cellular functions of metals, for example, zinc in leaf cells [16], copper in

neuronal cells [17], and calcium, phosphorus, and iron in algae [18].

Although it is attractive to image the same cells using multiple X-ray modalities [19,20],

challenges exist in developing suitable sample preparation protocols to optimize

contrast and mitigate the radiation damage associated with the measurement of the

same cells, as well as data correlation between the two imaging modalities. Addressing

these challenges is crucial to facilitate the routine application of these correlative

techniques for precise imaging at cellular and subcellular levels. Here we developed a

single-cell X-ray imaging workflow that encompasses sample preparation, targeting,

multi-modality data acquisition, and computational alignments of multi-modality image

data at nanometer resolutions. Our method offers a reliable and practical approach for

obtaining the structure and chemical information of a single cell with nanometer-level

resolution. The results of our research pave the way for the broad application of hard X-

rays for correlative characterization of cell structure and function, opening up new

possibilities for advanced biological investigations.

Results

A single-cell multi-modality X-ray imaging workflow

HEK293 cells were utilized in this study to demonstrate the workflow. Cells are fragile
and their manipulation and targeting require the use of compatible sample holders and
experimental setups across various imaging modalities. **Fig. 1** depicts the devised
workflow for achieving single-cell-level correlative multi-modal imaging. To begin, an X-
ray transparent Si_3N_4 membrane is used for growing cells (**Fig. 1a**). Subsequently, the
grown cells are chemically fixed using paraformaldehyde (PFA) to preserve cellular
structure and morphology (**Fig. 1b**). Fixed cells are manually blotted using filter paper
and rapidly plunge-freeze into liquid ethane using a Vitrobot (Thermo Fisher Scientific
Inc.), and later are transferred into liquid nitrogen (**Fig. 1c**). The vitrified cells were then
freeze-dried [21] to retain cell structure while gradually depleting water through
sublimation. (**Fig. 1d**). Dried cells are imaged using a confocal light microscope for
targeting and labeling cells (**Fig. 1e**). The labeled cells are measured first with XRT and
then with XRF (**Fig. 1f**). XRT and XRF data are processed separately to obtain
respective 3D tomograms and 2D chemical maps (**Fig. 1g**). Two sets of images are
then computationally aligned to generate the final integrated images (**Fig. 1h**). This
integrated workflow provides a comprehensive and detailed visualization of the cellular
structure and the distribution of chemical elements within the same cell.

Single-cell sample preparation and labeling

Freeze drying plays a critical role in preparing cell samples for XRF experiments [22,23].
However, XRF imaging is limited to detecting fluorescence-emitting chemicals such as
Zn and Ca while does not provide information about the carbon-, nitrogen-, and oxygen-
rich cellular structure. Consequently, fixation conditions optimized for XRF imaging may
not be optimal for the correlated XRT, which aims to reveal the three-dimensional
cellular structure. To address this issue, the fixation conditions were optimized using 4%
paraformaldehyde (PFA), a routinely employed fixative for immunolabeling and x-ray
fluorescence imaging [23,24]. The goal was to strike a balance between maintaining the
integrity of the cellular structure for XRT and preserving chemical elements for
subsequent XRF imaging. Upon the addition of 4% PFA to the cultured cells, the entire
fixation process was continuously monitored using a Nanolive imaging system at 30-
second intervals over two hours (**Supplementary Video 1**). The snapshots before and
after fixation are shown in **Supplementary Fig. 1**. No significant changes were

observed in the fixed cells compared to the initial time point (0 min). Furthermore, we didn't observe membrane blebbing or loss of cytoplasmic content during the entire two-hour fixation period [25].

Chemical fixation of cells is essential to maintain cellular structures for XRT analysis. Insufficient PFA fixation (0 minutes and 15 minutes) resulted in bubbles and the loss of internal features during XRT analysis (**Supplementary Fig. 2a,b**). Without fixation (0 minutes), the freeze-dried cell collapsed and was flatted on the Si_3N_4 membrane (**Supplementary Fig. 2a**). Sixty-minute fixation did not result in bubbles, but the internal cellular features were mostly lost compared to those of the 120-minute fixation (**Supplementary Fig. 2c and Fig. 2c-d**).

To perform correlative XRT and XRF imaging on a single cell, the cell density on the Si_3N_4 membrane was optimized to ensure sufficient separation between cells. Additionally, a confocal microscope was used to capture both the overall view and zoomed-in images of cells on the Si_3N_4 membrane (**Supplementary Fig. 3**). The cells were labeled and the labeled locations were utilized to target the same cells in XRT and XRF experiments. This multi-resolution imaging approach can be employed to label and target any cells on a Si_3N_4 membrane.

Hard X-ray tomography

XRT experiments were conducted at the Full-field X-ray Imaging (FXI) beamline at the National Synchrotron Light Source II (NSLS-II) at Brookhaven National Laboratory (BNL). The FXI beamline is equipped with a Transmission X-ray Microscope (TXM) capable of rapid measurements at 30 nm spatial resolution in less than one minute [26]. To minimize X-ray radiation damage to the samples, an optical telescope, coaligned with the X-ray beam, was utilized for target cell identification and pre-alignment. This telescope ensured that the samples were not exposed to the X-ray beam until the actual XRT data collection started. Lower energy X-rays increase intensity and contrast. With the optics set up at the FXI beamline at NSLS-II, 7 keV is the lowest X-ray energy we can reach. We conducted all the tomography experiments at 7 keV.

The tomograms of the target cells were reconstructed using TXM sandbox [27] and

visualized with IMOD [28]. **Fig. 2a,b** displays the projection of a cell's tomogram, revealing

a well-defined plasma membrane encompassing rich cellular structures. The shape of

the cell matches the confocal microscope image depicted in **Supplementary Fig. 3**.

Within the cell, two elongated organelles of about 500-800 nm in size are visible close

to the plasma membrane (**Fig. 2c,d**). Based on their appearance, these organelles are

likely bilayer endosome/lysosome (**Fig. 2c**) or multilayer mitochondrion (**Fig. 2d**). In

addition, we observed double membrane structures (**Fig. 2d**), i.e. plasma membrane

and intracellular membrane. By considering their length and smooth curvature of the

internal membrane, we assigned it as the nucleus membrane. Upon viewing the cell

from the top of the Si_3N_4 membrane, a few small vesicles of 300 nm can be identified

(**Fig. 2e**). Based on the size and high x-ray contrast, we assigned these vesicles as lipid

droplets. These lipid droplets exhibit spherical shapes, contrasting with the surrounding

air. Another compartment membrane structure is displayed between the plasma

membrane and intracellular membrane (**Fig. 2e**). However, we were not able to

rationally assign the membrane structure to any known organelle.   Based on the

features of the nucleus membrane and high contrast from chromatin, we were able to

segment the nucleus manually inside the cell (**Fig. 2f,g**). The 2D views of the 3D

volume along three orthogonal planes demonstrate the positioning of the segmented

nucleus within the cell (**Fig. 2g**). More structure details of this cell can be visualized in

Supplementary Fig. 4 and **Supplementary Video 2**.

X-ray fluorescence imaging

XRT is an effective method for visualizing cellular structures. Nonetheless, due to low

abundance, it cannot reveal the chemical element of trace micronutrient metals such as

Zn, Fe, Cu, and Ca, essential for proper cell function. To investigate the elemental

distributions within the same cell, we performed XRF measurement at the Submicron

Resolution X-ray Spectroscopy (SRX) beamline at NSLS-II. In order to precisely locate

the same cell, we utilized an in-line optical microscope at SRX to identify and center the

cell for the collection of raster scanning x-ray fluorescence data. To achieve high spatial

resolution, we utilized a beam size of approximately 100 nm and a step size of 100 nm

at an x-ray energy of 12 keV.

Figure 3 presents the chemical mapping of six common chemical elements: Ca, Cl, Cu,

Fe, P, and Zn. Among these, zinc was predominantly localized in the center of the cell,

i.e. nucleus, with its concentration gradually decreasing but remaining within the

nucleus region. Phosphorus is mostly enriched in the chromatin/nucleus, consistent with

its co-localization with zinc, which is known to bind to transcription factors to regulate

gene transcription in the nucleus [29]. Interestingly, calcium appears to be distributed in

separate regions of various sizes of about 1 µm or smaller in diameter. Given calcium's

cytotoxic nature, it is typically confined to calcium stores [30]. Therefore, it is likely that

these calcium-rich regions correspond to calcium stores associated with ER or

mitochondria. Copper seems to be attached to the outside of the phosphorus and zinc

map but extends beyond the zinc map (**Supplementary Fig. 5**). The chloride

distribution appears to resemble that of zinc and phosphorus, suggesting its localization

to the nucleus. Unlike other elements, which remain immobile during the fixation

process, chloride is soluble within the cell. It remains unclear why it remained in the

nucleus despite the PFA fixation.

Correlation of X-ray tomography and fluorescence data

We established a correlation between the 3D tomography image and the 2D

fluorescence data obtained from the same cell, displaying chemical element localization

within the cell. To achieve accurate alignment between both images, additional XRF

data were acquired by performing rotations at -15°, 25°, and 55° along the z-axis,

consistent with the rotation axis employed at the FXI. To generate correlation maps, the

3D tomogram was appropriately rotated to match the orientation of the 2D fluorescence

maps. Subsequently, we performed the total projections of the segmented cell plasma

membrane and nucleus onto their respective 2D fluorescence maps (**Fig. 4 and**

Supplementary Fig. 6). These multi-orientation XRF data facilitated the precise overlay

of fluorescence signals onto the corresponding cellular structures, enabling the

visualization and examination of chemical element distribution within the cell. For

instance, the four orientations utilized for the calcium maps indicate that these calcium
stores are situated close to, yet outside of the nucleus, suggesting their association with
the rough ER or mitochondria. Interestingly, copper is accumulated at one side of the
cell and is also close to, but outside of, the nucleus. The elongated shape of the Cu
map along the nucleus membrane, suggests its localization to rough ER or Golgi
apparatus. Copper and calcium have been reported to be involved in the function of
both mitochondria and the ER, raising intriguing questions for future studies [31,32]. The
correlative and non-invasive multi-modality imaging of the same cell provides a tool to
further characterize element distribution inside a cell which may not be possible using
other imaging tools.

Radiation damage

To assess the X-ray-caused radiation damage, a series of six consecutive XRT scans
were performed on the same cell, employing different experimental conditions such as
exposure time, rotation speed, and X-ray energy. Subsequently, a 7[th] scan was
conducted with the same parameters as the initial one. The resulting tomograms were
reconstructed using TXM sandbox and IMOD, employing the same reconstruction
parameters and post-processing filters. The first and the last tomograms were chosen
for comparative analysis (**Fig. 5**). The results indicate that the repeated tomography
data collection induced the cell-volume shrinkage by approximately 5% along each
dimension (**Supplementary Fig. 7**). In addition, high-resolution features seen in the first
scan were blurred or even disappeared in the 7[th] scan. For example, two small white
objects (1 and 2) were blurred in the last scan. One tiny object (4) and two elongated
structures (2 and 3) disappeared in the last scan (**Fig. 5**). The estimated X-ray dose for
each scan was about 5 MGy determined by using the program RADDOSE3D [33]. To
quantify the progressive change in cell volume with respect to repeated X-ray scans and
radiation damage, we measured the alternation in cell dimension as shown in
Supplementary Fig. 7. Notably, significant changes in cell volume were observed after
the first three scans, equivalent to an X-ray dose of 15 MGy. We suggest that for single-
cell X-ray tomography, the total dose should be limited to 15 MGy or lower.

Discussion

In this work, we developed an experimental and computational workflow that allows for the correlation of two distinct bioimaging modalities: nano-XRT and nano-XRF imaging. This approach enables the investigation of intracellular structures and chemical-element distributions within the same cells. The fast XRT serves as an effective screening tool to identify cells with interesting structural features. The selected cells are then scanned with XRF to obtain elemental maps, providing complementary information about the distribution of specific elements within the cell. In addition, the tomogram of a cell serves as a spatial framework, allowing for the precise localization of elemental maps within the context of cellular structure. The correlation between X-ray tomography and fluorescence provides valuable insights into cellular compositions (**Fig. 4**). For instance, phosphorus, a key component of DNA and chromatin, enriches the central region of the cell, corresponding to the segmented nucleus. Interestingly, zinc, which exhibits stronger fluorescence signals compared to phosphorus, co-localized with phosphorus in the same cellular region. This suggests a potential involvement of zinc in DNA-related biological activities, supported by previous research [34].

The abundance of each chemical element within a cell has a wide range of distributions, resulting in XRF signals with varying intensities. Further optimization of detector sensitivity and data collection conditions may increase the signal-to-noise ratios for better-detecting low-concentration trace elements such as Fe, Mn, Co, and Ni, which are essential to support proper cell function [35]. During our cell culture and sample preparation, no additional chemical elements were introduced. The observed distribution and localization of chemical elements within the HEK293 cell likely reflect their physiological concentrations and localization. In addition to correlating two X-ray imaging modalities, further correlation with fluorescence optical microscopy will provide additional insights into the function of transporter proteins associated with these elements. For instance, overexpression or knockdown of zinc transporters [36,37] in the plasma membrane and intracellular membranes of a cell can shed light on their role in regulating zinc homeostasis and their physiological and pathological implications.

Radiation damage presents a significant challenge in achieving high-resolution imaging

of biological samples [38,39]. Chemical fixation has proven to be an effective method for

preserving biological samples and mitigating radiation damage [23,40]. By employing

chemical fixation, it was possible to perform hard X-ray imaging on cells at room

temperature while preserving the integrity of intracellular features [13,25,41,42]. To prepare

radiation-resistant cell samples, we employed an extended fixation time of 2 hours using

PFA followed by freeze-drying to remove water. Live-cell imaging provides an ideal

approach to monitor the fixation process and identify optimal fixation time for multi-

modality X-ray experiments (**Supplementary Movie 1**). We hypothesize that the longer

fixation duration allowed for better penetration of PFA into intracellular organelles

including the nucleus for a more complete fixation of cellular contents. Our sample

preparation method was developed specifically for HEK293 cells, but it is generalizable

and can be applied to identify optimal sample preparation conditions for other cell types.

In our sample preparation, we deliberately avoided the use of contrast-enhancing

metals, such as osmium, lead, and uranyl, as they have the potential to introduce

additional artifacts to the cellular structure and interfere with the XRF analysis.

In other techniques such as cryogenic X-ray crystallography and cryo-electron

microscopy (cryo-EM), the recommended radiation doses for biological samples are 20

MGy for frozen-hydrated samples and 0.38 MGy for room-temperature hydrated

samples [41,43]. With hydrated algae *Chlorella*, doses of 0.0015 MGy changed the

ultrastructural organization. With hydrated living Chinese hamster ovarian (CHO) cells,

doses of about 0.1 MGy caused morphological changes [44]. Dehydrated cells are more

resistant to X-ray radiation damage than hydrated cells. It has been reported in several

studies that dose damage is significantly more visible in wet samples than in dehydrated

samples [44,45]. For chemically fixed hydrated samples, doses of about 1 MGy were

shown to cause cell shrinkage and mass loss[46]. In contrast, for dehydrated samples,

doses of up to 10 MGy did not lead to appreciable mass loss [46]. It seems the

combination of dehydration and chemical fixation will increase the dose damage

resistance. In our study, we investigated the dose damage effects in dehydrated and

chemically fixed cells. We believe that the dose damage delivered by one XRT (5 MGy)

is consistent with the nanometer resolution features without observing any visible

structural changes. Remarkably, we observed no significant overall cell volume shrinkage even after three scans, resulting in an accumulated dose of 15 MGy, which is significantly higher than the tolerable dose (1 MGy) for hydrated cells. In our time-dependent fixation test, we found that 2-hour fixation (**Fig. 2**) is better than 15-minute and 60-minute fixation (**Supplementary Fig. 2**) in terms of preserving structural features characterized by XRT analyses. A long fixation time in the sample preparation is thus recommended for XRT measurements to alleviate the radiation damage challenge. Exploring the precise relationship between fixation time and X-ray damage could be an intriguing avenue for future studies.

In our multimodal experiments, we conducted XRF measurements after performing seven scans for XRT data collection. However, we didn't consider the cumulative dose prior to the XRF experiments. Consequently, we observed a slight discrepancy between the XRF maps obtained after the seventh scan and the X-ray tomography data from the initial scan, attributed to cell shrinkage and radiation damage (**Fig. 5 and supplementary Fig. 7**). XRF imaging primarily detects chemical elements bound to proteins and nucleic acids, making it less sensitive to radiation damage compared to fine cellular structures. Retrospectively, a more rational experimental strategy would have been to allocate the total dose of 15 MGy between XRT and XRF data acquisition for better results. Nevertheless, our analysis of radiation damage underscores the significance of careful planning and the selection of appropriate experimental conditions for correlative multimodal X-ray experiments.

Methods

Visualization of the chemically fixed cell using live-cell imaging

Adherent HEK293 cells (ATCC Cat# CRL-1573) were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented by 10% (v/v) fetal bovine serum and 100 units/ml of penicillin-streptomycin (Cat# 15140122, Thermo Fisher). Cell subculture was performed twice a week to maintain cell viability and proliferation. To visualize the cell growth, 1 ml of suspension of cells with a concentration of 1×10^5 cells/ml was seeded

onto a 35 mm Ibidi μ-Dish (Cat#80136, Ibidi) and allowed cells to grow overnight. Cell growth was visually inspected by a light microscope before chemical fixation.

To visualize the chemical fixation process, a 4% (v/v) paraformaldehyde (PFA) solution was diluted from 16% stock (Cat# 50-980-487, Electron Microscopy Sciences) by 1X Gibco™ DPBS (Cat# 14190-136, Thermo Fisher). Firstly, cell culture media was discarded carefully followed by two washes, each with 1 ml of 1X Gibco™ DPBS. Subsequently, the fixation solution was gently added from the side of the well to cover the middle area of the dish. A transparent lid was placed on the dish, and the dish was placed in stage top CO_2 incubator on a 3D Cell Explorer-fluo light microscope (Nanolive). The incubator was operated at 37°C, 5% CO_2, and 95-100% humidity. The microscope was operated to perform real-time imaging for 2 hours, with images captured at 30-second intervals.

Cell culture on Si_3N_4 membrane

Seeding cells on Si_3N_4 membranes were adapted from the reported procedure [21]. Si_3N_4 membranes (Cat# NX5200E, Norcada) were glow-discharged using a PELCO easiGlow Glow Discharge Cleaning System (Tedpella) operated at 30 mA for 30 seconds. Subsequently, they were treated with 10 μl of poly-L-lysine (Cat# 25988-63-0, Sigma) and incubated in a cell-culture incubator (37 °C and 8% CO_2) for 25 min. Afterward, the membranes were washed 3 times for 10 seconds each with sterile MilliQ water and vacuum-dried overnight. Then, the membranes were seeded with 10 μl of 1×10^6 cells/ml in a Falcon 24-well cell culture plate (Cat# 353047, Fisher Scientific) and the plate was incubated in the cell-culture incubator (37 °C and 8% CO_2) for 25 minutes, resulting in roughly 250 cells/mm^2 on the Si_3N_4 after freeze-drying. Furthermore, 500 μl of the cell culture media was gently added from the side of the well to ensure the coverage of the entire membranes. The plate was then stored in the incubator for overnight cell culture.

Chemical fixation

The chemical fixation procedure was modified from [21,25]. Cell culture media was removed from the wells containing the Si_3N_4 membrane with attached cells.

351 Subsequently, the Si_3N_4 membranes were washed with 1 ml DPBS twice and fixed for 2

352 hours with 1 ml of 4% PFA as previously described. After fixation, membranes were

353 washed twice with 1 ml of ammonium acetate (150 mM, pH 7.4) to remove residual

354 fixatives.

Plunge-freezing and freeze-drying

To prepare the samples for freeze-drying, samples were blotted manually by using filter

paper to remove liquid on the frame and the back of the Si_3N_4 window. The plunge-

freezing was performed using a Mark IV Vitrobot (Thermo Fisher). Frozen samples were

then transferred into a pre-cooled 2-ml screw cap tube (Cat# HS10060, Millipore Sigma)

for freeze-drying using a FreeZone Console Freeze Dryer (Cat# 711211110, Labconco).

After the freeze dryer reached –80°C with a vacuum level of 0.05 mbar, the sample

tubes were quickly transferred into a freeze-drying container preloaded with liquid

nitrogen. The container was swiftly connected to the freeze-drying machine while the

vacuum valve was turned from off to on. The samples were subjected to freeze-drying

for 2 days to ensure complete removal of water.

Cell imaging and labeling for correlated multimodality data acquisition

To collect correlative data at multiple X-ray beamlines, it is necessary to perform cell

imaging and labeling. We used a Leica TCS SP5 (Leica Microsystems Inc., Deerfield,

IL, USA) to provide an overall view of cells on the Si_3N_4 membranes as well as to label

cells for correlated X-ray imaging data acquisition. The freeze-dried sample was gently

put upside down on the center of a µ-Dish before imaging. A 10x objective lens was

employed to survey the cells present on the membranes. Once a specific region of

interest was identified, the objective lens was switched to 50x magnification to capture

high-resolution images for cell labeling, serving as a reference for subsequent X-ray

data collection.

X-ray tomography data collection

The samples were manually mounted onto the FXI beamline data collection stage, and

the target cell was identified using a telescope that precisely aligned with the X-ray

beam. The XRT data were acquired using X-rays at 7 KeV. Two different exposure

schemes were employed: 0.1 sec exposure at a speed of 3°/ sec and 0.2 sec exposure

at a speed of 1.5°/sec. For each scheme, three data sets were collected with a defocus

of 0, 50, and 100 μm. To assess the extent of radiation damage, an additional data set

was collected with the same condition as used for the first data set.

Correlated XRF data collection

The same sample was used at the SRX beamline [47] for XRF measurements following

the XRT experiments. At SRX, a pair of rhodium-coated KB (Kirkpatrick–Baez) mirrors

was used to focus the X-ray beam down to 190 nm (H) x 230 nm (V) nm. By employing

an in-line optical light microscope (VLM), we were able to locate the same cell for

correlative chemical mapping. The focused beam was used to raster scan the cell,

allowing for the measurement of elements of biological significance. Cells were fly

scanned with 100 nm step size with 0.1 sec dwell time, with the photon energy at 12

keV and flux of about 4.5×10^{11} ph/sec. The estimated dose is 14 MGy per pixel. These

XRF maps are scanned in both horizontal and vertical directions with the same steps.

For the 0-degree scan, the step size was 100 nm, with 270 x 301 pixels. For the other

angles, the step size was 300 nm, with 141 x101 pixels. A Vortex 7-element silicon drift

detector (Hitachi, Ltd.) was located at an angle of 75° with respect to the incident beam

to capture the fluorescence emission. The obtained spectra were subsequently

analyzed using the open software PyXRF [48] to generate elemental distribution maps for

each element.

X-ray imaging data analysis

X-ray tomograms were reconstructed using TXM-Sandbox [27] and IMOD [28]. As described

in TXM-sandbox, the rotation center position was first identified in TXM-sandbox's "Trial

Center" mode in the raw images. Three filters from the TXM-sandbox were used to

enhance the signal-to-noise ratio (SNR). Initially, a stripe removal filter with a pixel size

range of 21-81 was used to minimize ring artifacts. Subsequently, a phase retrieval filter

and a median filter with a kernel size of 5x5 were applied to further improve the SNR.

The choice of filter parameters was guided by the SNR of the reconstructed image.

Then a full image reconstruction was performed with "Gridrec" algorithm under "Volume

Reconstruction" mode. For the reconstruction, the voxel size is 21 nm and the field of

view is 52 µm.

The cell membrane was segmented semiautomatically using FIJI [49]

(https://imagej.net/software/fiji/) and IMOD. First, a Canny edge detector was employed

to enhance the boundaries within a predefined region of interest in the image, followed

by the application of thresholding to convert these enhanced boundaries into binary

masks. The threshold value was chosen to retain only the cell membrane boundaries

while eliminating the nucleus membrane boundaries. Afterward, image binary opening

and closing operations were conducted to eliminate smaller noise objects. First, a

Canny edge detector was used to enhance the cell membrane region and then

thresholding was used to convert the enhanced membrane region to binary masks. The

resultant membrane masks were converted to a 3D IMOD model using "imodmop" and

"imodauto" commands. The cell nucleus was segmented manually using IMOD. The

resultant 3D model was visualized with IMOD and Napari [50] (http://narpari.org).

Correlation of X-ray tomography and fluorescence images

The reconstructed 3D X-ray tomograms were first projected into 2D projection maps

along the 4 different angles (-15°, 0°, 25°, 55°). Then 2D rigid image registration

(translation + rotation) was performed with FIJI [49] (https://imagej.net/software/fiji/) on 2D

X-ray projection image and 2D fluorescent image separately for four different angles.

The registered images from two imaging modalities were overlaid and visualized with

Napari.

References

1   Le Gros, M. A., McDermott, G. & Larabell, C. A. X-ray tomography of whole cells.
Curr. Opin. Struc. Biol. 15, 593-600, (2005).
2   Ou, X., Chen, X., Xu, X., Xie, L., Chen, X., Hong, Z., Bai, H., Liu, X., Chen, Q., Li,
L. & Yang, H. Recent development in X-Ray imaging technology: future and
challenges. *Research* 2021, 9892152, (2021).
3   Andrews, J. C., Meirer, F., Liu, Y. J., Mester, Z. & Pianetta, P. Transmission X-
Ray microscopy for full-field nano imaging of biomaterials. *Microsc. Res. Tech.*
74, 671-681, (2011).

4   Spence, S., Lee, W. K., Lin, F. & Xiao, X. H. Transmission x-ray microscopy and
its applications in battery material research-a short review. *Nanotechnology* 32,
442003, (2021).
5   Pushie, M. J., Sylvain, N. J., Hou, H., Hackett, M. J., Kelly, M. E. & Webb, S. M.
X-ray fluorescence microscopy methods for biological tissues. *Metallomics* 14,
mfac032, (2022).
6   Paunesku, T., Vogt, S., Maser, J., Lai, B. & Woloschak, G. X-ray fluorescence
microprobe imaging in biology and medicine. *J. Cell Biochem.* 99, 1489-1502,
(2006).
7   Decelle, J., Veronesi, G., Gallet, B., Stryhanyuk, H., Benettoni, P., Schmidt, M.,
Tucoulou, R., Passarelli, M., Bohic, S., Clode, P. & Musat, N. Subcellular
chemical imaging: new avenues in cell biology. *Trends Cell Biol.* 30, 173-188,
(2020).
8   Chen, J. H., Vanslembrouck, B., Loconte, V., Ekman, A., Cortese, M.,
Bartenschlager, R., McDermott, G., Larabell, C. A., Le Gros, M. A. & Weinhardt,
V. A protocol for full-rotation soft X-ray tomography of single cells. *STAR Protoc.*
3, 101176, (2022).
9   Larabell, C. A. & Nugent, K. A. Imaging cellular architecture with X-rays. *Curr.*
Opin. Struc. Biol. 20, 623-631, (2010).
10   Hémonnot, C. Y. & Ko□ster, S. Imaging of biological materials and cells by X-ray
scattering and diffraction. *ACS nano* 11, 8542-8559, (2017).
11   Stampanoni, M., Mokso, R., Marone, F., Vila-Comamala, J., Gorelick, S., Trtik,
P., Jefimovs, K. & David, C. Phase-contrast tomography at the nanoscale using
hard x rays. *Phys. Rev. B* 81, 140105, (2010).
12   Yang, Y., Li, W., Liu, G., Zhang, X., Chen, J., Wu, W., Guan, Y., Xiong, Y., Tian,
Y. & Wu, Z. 3D visualization of subcellular structures of Schizosaccharomyces
pombe by hard X-ray tomography. *J. Microsc.* 240, 14-20, (2010).
13   Jin, Q., Paunesku, T., Lai, B., Gleber, S. C., Chen, S. I., Finney, L., Vine, D.,
Vogt, S., Woloschak, G. & Jacobsen, C. Preserving elemental content in
adherent mammalian cells for analysis by synchrotron-based x-ray fluorescence
microscopy. *J. Microsc.* 265, 81-93, (2017).
14   Matsuyama, S., Shimura, M., Fujii, M., Maeshima, K., Yumoto, H., Mimura, H.,
Sano, Y., Yabashi, M., Nishino, Y., Tamasaku, K., Ishizaka, Y., Ishikawa, T. &
Yamauchi, K. Elemental mapping of frozen-hydrated cells with cryo-scanning X-
ray fluorescence microscopy. *X-Ray Spectrom.* 39, 260-266, (2010).
15   Gramaccioni, C., Yang, Y., Pacureanu, A., Vigano, N., Procopio, A., Valenti, P.,
Rosa, L., Berlutti, F., Bohic, S. & Cloetens, P. Cryo-nanoimaging of single human
macrophage cells: 3D structural and chemical quantification. *Anal. Chem.* 92,
4814-4819, (2020).
16   Vigani, G., Bohic, S., Faoro, F., Vekemans, B., Vincze, L. & Terzano, R. Cellular
fractionation and nanoscopic X-ray fluorescence imaging analyses reveal
changes of zinc distribution in leaf cells of iron-deficient plants. *Front. Plant Sci.*
9, 1112, (2018).
17   Dodani, S. C., Domaille, D. W., Nam, C. I., Miller, E. W., Finney, L. A., Vogt, S. &
Chang, C. J. Calcium-dependent copper redistributions in neuronal cells revealed

by a fluorescent copper sensor and X-ray fluorescence microscopy. ***Proc. Natl.***
Acad. Sci. USA 108, 5980-5985, (2011).

18 Diaz, J., Ingall, E., Vogt, S., de Jonge, M. D., Paterson, D., Rau, C. & Brandes, J.
A. Characterization of phosphorus, calcium, iron, and other elements in
organisms at sub-micron resolution using X-ray fluorescence spectromicroscopy.
Limnol. Oceanogr-Meth. 7, 42-51, (2009).

19 Yang, Y., Fus, F., Pacureanu, A., da Silva, J. C., De Nolf, W., Biot, C., Bohic, S.
& Cloetens, P. Three-dimensional correlative imaging of a malaria-infected cell
with a hard x-ray nanoprobe. ***Anal. Chem.*** 91, 6549-6554, (2019).

20 Deng, J. J., Vine, D. J., Chen, S., Nashed, Y. S. G., Jin, Q. L., Phillips, N. W.,
Peterka, T., Rossc, R., Vogt, S. & Jacobsen, C. J. Simultaneous cryo X-ray
ptychographic and fluorescence microscopy of green algae. ***Proc. Natl. Acad.***
Sci. USA 112, 2314-2319, (2015).

21 Bissardon, C., Reymond, S., Salome, M., Andre, L., Bayat, S., Cloetens, P. &
Bohic, S. Cell culture on silicon nitride membranes and cryopreparation for
synchrotron X-ray fluorescence nano-analysis. ***J. Vis. Exp.***, 31885377, (2019).

22 Finney, L. A. & Jin, Q. Preparing adherent cells for X-ray fluorescence imaging
by chemical fixation. ***J. Vis. Exp.***, 25867691, (2015).

23 Li, Y., Almassalha, L. M., Chandler, J. E., Zhou, X., Stypula-Cyrus, Y. E., Hujsak,
K. A., Roth, E. W., Bleher, R., Subramanian, H., Szleifer, I., Dravid, V. P. &
Backman, V. The effects of chemical fixation on the cellular nanostructure. ***Exp.***
Cell Res. 358, 253-259, (2017).

24 Zu, Y., Tong, X. J., Wang, Z. X., Liu, D., Pan, R. C., Li, Z., Hu, Y. Y., Luo, Z.,
Huang, P., Wu, Q., Zhu, Z. Y., Zhang, B. & Lin, S. TALEN-mediated precise
genome modification by homologous recombination in zebrafish. ***Nat. Methods***
10, 329-+, (2013).

25 Huebinger, J., Spindler, J., Holl, K. J. & Koos, B. Quantification of protein mobility
and associated reshuffling of cytoplasm during chemical fixation. ***Sci. Rep.*** 8,
17756, (2018).

26 Lee, W. K., Reininger, R., Loo, W., Gambella, R., O'Hara, S., Chu, Y. S., Zhong,
Z. & Wang, J. FXI: a full-field imaging beamline at NSLS-II. ***Proc. Spie.*** 9592,
959209, (2015).

27 Xiao, X., Xu, Z., Lin, F. & Lee, W. K. TXM-Sandbox: an open-source software for
transmission X-ray microscopy data analysis. ***J Synchrotron Radiat.*** 29, 266-
275, (2022).

28 Mastronarde, D. N. & Held, S. R. Automated tilt series alignment and
tomographic reconstruction in IMOD. ***J. Struct. Biol.*** 197, 102-113, (2017).

29 Eide, D. J. Transcription factors and transporters in zinc homeostasis: lessons
learned from fungi. ***Crit. Rev. Biochem. Mol. Biol.*** 55, 88-110, (2020).

30 Sammels, E., Parys, J. B., Missiaen, L., De Smedt, H. & Bultynck, G. Intracellular
Ca2+ storage in health and disease: a dynamic equilibrium. ***Cell Calcium*** 47,
297-314, (2010).

31 Ruiz, L. M., Libedinsky, A. & Elorza, A. A. Role of copper on mitochondrial
function and metabolism. ***Front. Mol. Biosci.*** 8, 711227, (2021).

32 Ashby, M. C. & Tepikin, A. V. ER calcium and the functions of intracellular
organelles. ***Semin. Cell Dev. Biol.*** 12, 11-17, (2001).

Zeldin, O. B., Gerstel, M. & Garman, E. F. RADDOSE-3D: time- and space-resolved modelling of dose in macromolecular crystallography. *J. Appl. Crystallogr.* 46, 1225-1230, (2013).

Beyersmann, D. & Haase, H. Functions of zinc in signaling, proliferation and differentiation of mammalian cells. *Biometals* 14, 331-341, (2001).

Zoroddu, M. A., Aaseth, J., Crisponi, G., Medici, S., Peana, M. & Nurchi, V. M. The essential metals for humans: a brief overview. *J. Inorg. Biochem.* 195, 120-129, (2019).

Pratt, E. P. S., Damon, L. J., Anson, K. J. & Palmer, A. E. Tools and techniques for illuminating the cell biology of zinc. *Biochim. Biophys. Acta Mol. Cell Res.* 1868, (2021).

Kambe, T., Taylor, K. M. & Fu, D. Zinc transporters and their functional integration in mammalian cells. *J. Biol. Chem.* 296, 100320, (2021).

Baksh, P., Odstrcil, M., Parsons, A., Bailey, J., Deinhardt, K., Chad, J. E., Brocklesby, W. S. & Frey, J. G. Quantitative evaluation of hard x-ray damage to biological samples using EUV ptychography. *J. Phys.: Conf. Ser.* 849, 012034, (2017).

Howells, M. R., Beetz, T., Chapman, H. N., Cui, C., Holton, J., Jacobsen, C., Kirz, J., Lima, E., Marchesini, S. & Miao, H. An assessment of the resolution limitation due to radiation-damage in x-ray diffraction microscopy. *J. Electron Spectros. Relat. Phenomena* 170, 4-12, (2009).

Hopwood, D. Fixatives and fixation: a review. *Histochem. J.* 1, 323-360, (1969).

de la Mora, E., Coquelle, N., Bury, C. S., Rosenthal, M., Holton, J. M., Carmichael, I., Garman, E. F., Burghammer, M., Colletier, J. P. & Weik, M. Radiation damage and dose limits in serial synchrotron crystallography at cryo- and room temperatures. *Proc. Natl. Acad. Sci. USA* 117, 4142-4151, (2020).

Murk, J. L., Posthuma, G., Koster, A. J., Geuze, H. J., Verkleij, A. J., Kleijmeer, M. J. & Humbel, B. M. Influence of aldehyde fixation on the morphology of endosomes and lysosomes: quantitative analysis and electron tomography. *J. Microsc.* 212, 81-90, (2003).

Henderson, R. Cryoprotection of Protein Crystals against Radiation-Damage in Electron and X-Ray-Diffraction. *Proc. R. Soc. Lond. B Biol. Sci.* 241, 6-8, (1990).

Kirz, J., Jacobsen, C. & Howells, M. Soft-X-ray microscopes and their biological applications. *Q. Rev. Biophys.* 28, 33-130, (1995).

Beetz, T. & Jacobsen, C. Soft X-ray radiation-damage studies in PMMA using a cryo-STXM. *J. Synchrotron Radiat.* 10, 280-283, (2003).

Williams, S., Zhang, X., Jacobsen, C., Kirz, J., Lindaas, S., Vanthof, J. & Lamm, S. S. Measurements of wet metaphase chromosomes in the scanning-transmission X-ray microscope. *Journal of Microscopy* 170, 155-165, (1993).

Nazaretski, E., Coburn, D. S., Xu, W., Ma, J., Xu, H., Smith, R., Huang, X., Yang, Y., Huang, L., Idir, M., Kiss, A. & Chu, Y. S. A new Kirkpatrick-Baez-based scanning microscope for the Submicron Resolution X-ray Spectroscopy (SRX) beamline at NSLS-II. *J Synchrotron Radiat.* 29, 1284-1291, (2022).

48   Li, L., Yan, H., Xu, W., Yu, D. T., Heroux, A., Lee, W. K., Campbell, S. I. & Chu,
Y. S. PyXRF: Python-based X-ray fluorescence analysis package. ***Proc. Spie.***
10389, (2017).
49   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D.
J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A. Fiji: an open-source
platform for biological-image analysis. ***Nat. Methods*** 9, 676-682, (2012).
50   Contributors, N. napari: a multi-dimensional image viewer for Python. ***Zenodo***
https://doi. org/10.5281/zenodo 3555620, (2019).

Acknowledgments This work was supported by Laboratory Directed Research and

Development Program LDRD21-013 and the U.S. Department of Energy (DOE), Office

of Biological and Environmental Research (KP1601011). The work used NSLS-II which

is supported in part by the U.S. Department of Energy, Office of Science, Office of Basic

Energy Sciences Program under contract number DESC0012704 (KC0401040).

Author contributions Y.Y., X. X., and Q.L. designed the study and experiments. Z. L.,

X. Z., Y.Y., and X. X. performed the experiments. All authors analyzed the data. Z. L.

Y.Y, X. X. and Q.L. wrote the manuscript with help from other coauthors.

Competing interests Authors declare no competing interests.

Correspondence and requests for materials should be addressed to Y.Y., X.X, or Q.L.

Figures and captions

Cell culture | Chemical fixation | Plunge freeze | Freeze drying

FXI tomography | 3D reconstruction | Images correlation | 2D reconstruction | SRX Fluorescence | Cell labeling

Fig. 1. Overall workflow of single-cell correlative X-ray imaging. **a**, HEK293T cells are grown adherently on the flat surface of a poly-lysine-treated Si_3N_4 membrane. **b**, Adherent cells are washed twice with PBS and a chemical fixation solution is added to fix the cell. **c**, Si_3N_4 membrane is washed twice with 150 mM ammonium acetate and then plunge-freeze using a Vitrobot. The membrane is stored in a screw-cap microcentrifuge tube. **d**, The tube with the sample is transferred to a freeze-dryer device for freezing-drying. **e**, Cells are labeled using a confocal microscope for multiple beamlines data collection. **f**, Single-cell data are collected using x-ray tomography at the FXI beamline and x-ray fluorescence microscopy at the SRX beamline, respectively. **g**, Reconstruction of 3D tomograms and 2D fluorescence map. **h**, Correlation of multimodality images.

Fig. 2. Single-cell X-ray tomography. Total projection of the reconstructed X-ray tomogram in XY (**a**) and YZ (**b**) directions. **c-e**, cellular structure features including mitochondrion (**c**), nucleus membrane (**d**), and lipid droplets (**e**). The diameter for lysosome (L, yellow), mitochondria (Mito, red), and lipid droplets (LD, green) are 500 nm, 800 nm, and 350 nm, respectively. **f**, Orthoslices of the cell along three directions. The whole cell is visualized in 3D (XY, YZ, and XZ) views. (**g**) View of f superimposed with the segmentation for nucleus (cyan) and plasma membrane (orange). Scale bar is 5 µm for **a** and **b, and** 2 µm for **c-g**.

Fig. 3. X-ray fluorescence mapping of multiple chemical elements. The X-ray emission energy is 12 keV. Intensities are normalized for each component. The sizes of the three calcium-rich regions (labeled with numbers 1-3) are 1.0 μm,1.1 μm, 0.9 μm, respectively. Scale bar is 5 μm.

Fig. 4. Correlation of multi-modality data. Superimposition of 2D XRF and 3D XRT images along 4 different view angles. The cell was rotated along the y-axis by -15°, 0°, 25°, and 55°, respectively. Gray scale maps show the max projection of the 3D X-ray tomogram of the cells. Red masks show a segmented nucleus from the tomogram. Green maps show the corresponding localization of multiple elements. Scale bar is 5 µm.

629
Fig. 5. Radiation damage in X-ray tomographic imaging. Representative slices of
the reconstruction results of **a** from the first x-ray tomography scan and **b** from the last
scan (7th scan). The same experimental conditions are used for the first the last scans.
Due to radiation damage, features indicated by color arrows in **a** were weakened (1,5)
or disappeared (2-4) in **b**. Scale bars are 5 µm.

Supplementary information

0 minutes

2 hours

Supplementary Fig. 1. Nanolive live-cell imaging to monitor the PFA fixation
process. The first and the last images of fixation imaging were selected to investigate
the feature changes caused by the fixative. Some minor changes were observed such
as the plasma membrane shrinkage on the top of the cell and the faded nucleus
membrane. Although the fixative damage occurred, the overall cell structure didn't have
a significant change in the 2-hour fixation.

Supplementary Fig. 2. XRT reconstructions of insufficiently fixed cells. The projections of XRT tomograms at two orthogonal views. Arrows indicate bubbles induced by X-ray exposure. Scale bar is 5 μm. **a,** no fixation (0 minutes)**. b,** 15 minutes of fixation. **c,** 60 minutes of fixation.

650 **Supplementary Fig. 3. Multi-resolution confocal images for targeting and labeling**

651 **cells for correlative multimodality X-ray imaging data collection.** Both images

652 played important roles in displaying the target cell location that provided a reference for

653 X-ray imaging. A red arrow indicates the cell used for this study.

Supplementary Fig. 4. Representative slice views of intracellular organelles.
Lysosome (**a**, yellow), mitochondria (**b**, red), nucleus membrane (**b**, blue), and lipid
droplets (**c**, green) were observed in the reconstructed tomogram. Scale bars are 2 μm.
Numbers at the bottom left of each slice indicate the slice number in the 3D
reconstructed tomogram. Each slice is 21.7 nm in thickness. Distances between the first
slice and fourth slice for **a**, **b,** and **c** are 0.26 μm, 0.33 μm, and 0.46 μm, respectively.

Supplementary Fig. 5. X-ray fluorescence imaging indicates relative localization of Ca, Cu, and Zn. Intensities are normalized for each component to showcase the relative amount in the cell. Scale bar is 5 µm.

673
Supplementary Fig. 6. Correlation of multi-modality data. Superimposition of single-
cell 2D XRF with the first (**a**) and last (**b**) scans of the 3D XRT images along 4 different
angles. The cell was rotated along the y-axis by -15°, 0°, 25°, and 55° to collect angular
XRF data. Gray scale maps show the max projection of 3D XRT of the cells. Red masks
show segmented nucleus from the 3D XRT images. Green maps show the
corresponding localization of indicated chemical elements.

Supplementary Fig. 7. Change of cell size caused by radiation damage. The size of

the cell was measured in three orthogonal directions, namely, the Z, X, and Y axes,

using the maximum length as the basis with respect to the number of XRT scans.

Panels **a-c** illustrate the changes in size across seven different scans along the Z, X,

and Y axes.

Supplementary Video 1. Live cell imaging to monitor the PFA fixation process.

Supplementary Video 2. Sliced view of X-ray tomography (XRT) from top of the cell to the bottom membrane support.