Highly conserved, positively charged tandem repeats mediate multimerization of ice nucleation proteins

Thomas Hansen[1], Jocelyn C. Lee[1], Naama Reicher[2], Gil Ovadia[3], Shuaiqi Guo[4], Wangbiao Guo[4], Jun Liu[4], Ido Braslavsky[3], Yinon Rudich[2], Peter L. Davies[1*]

[1]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON Canada K7L 3N6

[2]Department of Earth and Planetary Sciences, The Weizmann Institute of Science, Rehovot 7610001, Israel

[3]The Robert H. Smith Faculty of Agriculture, Food and Environment, Institute of Biochemistry, Food Science, and Nutrition, The Hebrew University of Jerusalem, Rehovot 7610001, Israel

[4]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT 06536

[*]Corresponding author: Peter L. Davies, peter.davies@queensu.ca

Abstract

Ice nucleating bacteria like *Pseudomonas borealis* (*Pb*) can trigger frost formation at temperatures as high as −2 °C. The efficiency of ice formation by these bacteria comes from the ability of their >100 kDa ice nucleation proteins (INPs) to form ~10 MDa multimers. The INP monomers contain an array of tandem repeats, each of which is predicted to form a single coil in a β-solenoid. The 12 coils at the C-terminal end of the INP differ from the other 53 coils in lacking water organizing motifs and by having arginines on one side of the solenoid in place of acidic residues. Here we show that these arginine-containing coils (R-coils) are a distinct subdomain that, along with the adjacent putative 41-residue capping structure, facilitate INP multimerization. Bioinformatic analyses show that the cap structure is highly conserved, as is the number of R-coils. Indeed, the loss of just a few R-coils eliminated ice-nucleation activity, as did mutations designed to spoil the fold of the capping structure. Movement of the R-coils to the N-terminal end or to the middle of the β-solenoid caused a large decrease in ice nucleation temperature. Replacing the arginines with acidic residues decreased the nucleation temperature.

Activity was restored when these residues were in turn replaced by lysines. A role for electrostatic interactions in INP self-assembly was suggested by the effect of pH on ice nucleation activity in bacterial lysates. This activity declined by 3-6 °C at pH values below 5.0. When the INP-producing *E. coli* were examined by cryo-electron tomography, the cell cytoplasm contained clusters of 100 to 200 nm-long and ~5 nm-wide fibres that were absent from control bacteria. Each fibre is wide enough to have up to two INPs in cross-section and is long enough to span several INPs end to end. The ice nucleation activity in the lysates was remarkably resistant to heat treatment up to ~70 °C, above which there was a slight decline with increasing temperature. Even after heating to 99 °C, the lysate ice nucleation activity only lost ~8°C. We suggest that the R-coils play a crucial role in INP fibre assembly and that a bundle of these fibers could amass a sufficient number of ice-like water molecules to initiate ice nucleation at high sub-zero temperatures.

Introduction

Ice crystals grow from ice embryos, which are crystalline aggregates of water molecules that spontaneously form (homogeneous nucleation) in pure H_2O at approximately -38 °C (1). Ice can arise in nature at much warmer temperatures because various surfaces act as stabilizers of ice embryos (heterogeneous nucleation). Only once an ice embryo reaches a critical number of organized water molecules will it become stable enough to spontaneously grow at elevated temperatures, a process called ice nucleation (2). The most active heterogeneous ice nucleators are bacterial ice nucleation proteins (INPs), which can stabilize an ice embryo at temperatures as warm as −2 °C (3).

INPs are large proteins (up to ~150 kDa) that are thought to form multimers on the surface of the bacteria that express them (4, 5). AlphaFold predictions have provided some insight into the INP monomer structure (**Fig. 1A**). For the INP from *Pseudomonas borealis* (*Pb*INP) AlphaFold predicted a folded domain of ~100 residues at the N terminus followed by a flexible linker of ~50 residues, a repetitive domain composed of 65 16-residue tandem repeats, and a small 41-residue C-terminal capping structure (6). The predicted fold of the repetitive domain agrees with some previous homology-based models in which each 16-residue repeat forms a single coil of a β-solenoid structure (7, 8).

Most coils of the β-solenoid contain putative water-organizing motifs like Thr-Xaa-Thr (TXT) that occupy the same position in each coil to form long parallel arrays, and where Xaa is an inward pointing amino acid residue (**Fig. 1B**). Shorter versions of similar arrays have convergently evolved in insects to form the ice-binding sites of several hyperactive antifreeze proteins (AFPs) (9-11). According to the anchored clathrate water hypothesis, the ability of these arrays to position ice-like water molecules facilitates AFP adsorption to the ice crystal surface (12). In the ~10 times longer INPs, which additionally form multimers, the organizing effect on nearby water molecules is thought to increase to the point where ice embryos can be stabilized enough to lead to the spontaneous growth of ice at high sub-zero temperatures. Recently, we demonstrated that interruptions in the continuity of these water-organizing motifs decrease the ice nucleation temperature as much as extensive deletions of the water-organizing coils (6).

Previously we showed that the final 12 C-terminal coils lack the water organizing motifs and that deleting these coils resulted in a near total loss of activity (6). Interestingly, the necessity for these C-terminal coils was demonstrated by Green and Warren in 1986 in the first publication of an INP sequence (13) but was not further investigated. While the water-organizing coils (WO-coils) are characterized by their conserved TXT, SLT, and Y motifs, the defining feature of the C-terminal-most coils, other than the lack of these motifs, is that position 12 of the 16-residue coil is typically occupied by arginine. Thus, we refer to these non-WO-coils as R-coils. Since both coil types maintain the same predicted fold but serve different functions, we consider them subdomains of the same β-solenoid (14). In the WO-coils, position 12 is usually occupied by residues of the opposite charge, Asp and Glu. This charge inversion is noteworthy as it has been shown that electrostatic interactions contribute to the formation of INP multimers (15, 16). It has also been shown that INP activity is affected by pH, which is consistent with a role for electrostatic interactions (17). We and others have suggested that INPs may assemble through salt-bridging of the sidechains in these positions of the coil (6, 18).

Multimerization of INPs was shown in 1988 by Govindarajan and Lindow, who used radiation inactivation analysis to suggest a multimer size of 19 MDa (>100 monomers) (4). Computational estimates indicate that up to a 5-MDa (34 INPs) structure of laterally assembled INPs is

necessary for optimal activity, which puts the multimer size in the same order of magnitude as that determined experimentally (19). The tendency to form such large structures is one of many factors that makes these proteins difficult to work with (3) and may be part of why, despite many attempts, very little is known about them at a molecular level (20). The size of these structures does, however, make them amenable to size-based separation from most proteins (21, 22). INP multimers are also large enough to be visible using negative stain transmission electron microscopy (TEM) on enriched samples, revealing a fibril-like morphology (22, 23).

In nature, these multimers form on the surface of bacteria, anchored to the outer membrane by the N-terminal domain (24), which has been used extensively for the surface display of many proteins (25, 26). Remarkably, INPs with N-terminal truncations are only slightly less active, suggesting that INPs can multimerize in the cytoplasm whether anchored to the inner surface of the plasma membrane or free in solution (27) (28).

Here, we have studied the role of the R-coils in INPs through a series of mutations and rearrangements. Additionally, using cryo-focused-ion-beam (cryo-FIB) milling and cryo-electron tomography (cryo-ET), we have observed the fibrillar morphology of INP multimers within cells recombinantly expressing INPs. The R-coils' length, location, and sequence are critical for INP multimerization and hence INP activity. Although we report results using *Pb*INP, the bioinformatic analysis presented here indicates that these findings are universally applicable to the INP family, including the more commonly studied InaZ from *Pseudomonas syringae*.

Results

Bioinformatic analysis reveals conservation in the number of R-coils across all INPs

In *Pb*INP, there are 53 WO-coils and 12 R-coils (**Fig. 1A**), each composed of 16 residues (**Fig. 1B**). To determine whether this ratio of coil types is consistent across known INPs, we analyzed INP and INP-like solenoid sequences in the *nr* (NCBI) database. The long tandem arrays of coils in INPs make them prone to mis-assembly when using short-read DNA sequencing (29) so we opted to limit our dataset to sequences obtained by long-read technologies (Oxford Nanopore and Pacific Biosciences SMRT sequencing). From this bioinformatics study, it is apparent that

the number of WO-coils varies considerably from over 70 coils to around 30, with a median

length of 58 coils (**Fig. 1C**). In contrast, the length of the R-coil region is much less variable

across sequences, with 107 of 120 sequences containing either 10 or 12 R-coils (**Fig. 1C**). The

stark difference in length variation between the numbers of WO-coils and the R-coils supports

the hypothesis that these two regions have different functions.

The differences observed between the *Pb*INP WO-coil consensus sequence and the R-coil

consensus sequence, which include the loss of putative WO motifs in the former and the

appearance of basic residues at position 12 in the latter, are consistent across the entire dataset

(**Fig. 1D**). This is apparent from the sequence logos comparing them. Also worth noting is the

similarity of the sequence logos for WO-coils and R-coils in *Pb*INP (6) with those based on 120

sequences from the database.

Figure 1. Classification of unique INP tandem arrays into either WO-coil or R-coil subdomains. A) AlphaFold 2 model of *Pb*INP coloured by domain. Purple: N-terminal domain, pink: flexible linker, green: water-organizing (WO) coils, blue: arginine-rich (R) coils, yellow: C-terminal cap. The inset shows a cross section through the solenoid coil. **B)** 16-residue tandem repeat forming one coil from the β-solenoid with positions numbered from N- to C-terminus. **C)** The number of repeats in the WO-coils and R-coils for each unique sequence. *Pb*INP is included and indicated in orange. n = 121. **D)** Sequence logos constructed from each 16-residue repeat present in the dataset.

Incremental replacement of R-coils with WO-coils severely diminishes ice nucleating activity

Given the remarkable conservation of the R-coil count compared to the variability of the WO-coil numbers, we measured the functional impact of shortening the R-coil region. We designed mutants in which the R-coils were incrementally replaced with WO-coils, shortening the R-coil subdomain from 12 to 10, 8, 6, 4, or 1 coil(s), while retaining the same overall length as wild-type *Pb*INP (**Fig. 2A**). To avoid disrupting any potential interaction between the C-terminal cap structure and the R-coils, one R-coil was left in place to produce the 1 R-coil mutant.

Ice nucleation assays were performed to assess the activity of the incremental replacement mutants. In theory, replacement of R-coils by WO-coils could result in a gain of function as the water-organizing surface increased in area. However, as the number of R-coils was reduced, the nucleation temperature decreased (**Fig. 2B**). Replacing two or four R-coils to leave ten or eight in place resulted in a slight loss of activity compared to the wild-type protein (T_{50} = -9.1 °C and -10.2 °C, respectively, where T_{50} is the temperature at which 50% of droplets have frozen). Reducing the R-coil count to six dramatically decreased the activity (T_{50} = -35.7 °C). The construct with only four R-coils in place showed only the slightest amount of activity, and activity was entirely lost in the construct containing only one R-coil. Evidently, small decreases in the R-coil region length produce disproportionately large decreases in activity. Halving the length of the R-coils by replacing just six coils reduced ice nucleation activity by 26.9 °C, whereas reducing the WO-coil length in half decreased the T_{50} by less than 2 °C (6). Since the R-coils mostly lack the motifs required for water-organizing, we attribute the observed changes in nucleation temperature to changes in INP multimer formation.

Figure 2. Ice nucleation activity of mutant INPs in which R-coils are incrementally replaced with WO-coils. A) Diagram indicating the domain map of *Pb*INP and from it, the design of the INP mutants. Sections of the WO-coils used in the replacement of the R-coils are indicated. wt: wild type. **B)** Ice nucleation temperatures measured by WISDOM for *E. coli* cells expressing *Pb*INP and mutants with different numbers of R-coils. The temperature at which fifty percent of the droplets froze (T_{50}) is indicated with a hollow circle, and its corresponding value written nearby. The shaded region indicates standard deviation. **C)** The same constructs assayed for ice nucleation using a nanoliter osmometer. The grey box indicates temperatures beyond the lower limit of the NLO apparatus for detecting ice nucleation in this experiment.

The location of the R-coil subdomain is crucial

In addition to its length, we investigated whether the location of the R-coil subdomain is important for ice nucleation activity. We produced constructs where 11 of the 12 R-coils were relocated to either the N-terminal end of the solenoid or the approximate midpoint of the solenoid (**Fig. 3A**). As before, the C-terminal R-coil was left in place adjacent to the cap structure. We also produced an R-coil deletion construct, where the same 11 R-coils were deleted entirely from the protein.

The N-terminal relocation construct displayed markedly lower activity with a T_{50} = -22.4 °C compared to wild-type *Pb*INP, and the midpoint relocation construct displayed almost no activity (T_{50} = -36.1 °C), and was indistinguishable in activity from the construct where the R-coils were deleted (**Fig. 3B**).

Figure 3. Ice nucleation activity of mutant INPs with entirely relocated or deleted R-coil subdomain. a) Diagram indicating the design of the constructs. 11 of the 12 R-coil repeats were either moved within the construct or deleted. **b)** Freezing curves with T_{50} and number of unfrozen droplets indicated where applicable.

Targeted mutations reveal that positively charged residues are important for R-coil function

Having established the importance of R-coil position and length for high activity, we next investigated the features of this subdomain that are required for its activity. Looking at the charge distribution along the solenoid from N terminus to C terminus (**Fig. 4A**), we noted a switch at the start of the R-coils from an abundance of acidic residues to their replacement by basic residues. To probe the significance of this observation, we mutated all basic residues (R/K/H) in the R-coils to match those found in the same repeat positions of the WO-coils (D/G/E for positions 11 and 12, and S for position 14). In total, 17 basic residues – 10 Arginine (R), 4 Lysine (K), 3 Histidine (H) – were replaced in the R-coils to generate the RKH replacement mutant. The side chains at these positions are predicted by the AlphaFold model to point outward from the solenoid, so these mutations are unlikely to compromise the stability of the solenoid core.

There was a 10.3 °C drop in T_{50} from wild-type activity after RKH replacement (**Fig. 4B**) ($T_{50} =$ -19.1 °C). Although this is a large decrease in activity, it was not as deleterious as the relocation or deletion mutations (**Fig. 3**). The prominent, entirely conserved tyrosine in position 3 of the WO-coils is only present in the first three R-coils and is missing from the following nine coils, making it another candidate for mutation. Upon extending this "tyrosine ladder" through the R-coils (**Fig. 4A**), there was a 1.3-°C loss in activity. However, when combining the RKH replacement with the tyrosine ladder extension, an almost total loss of activity was observed (T_{50} = -36.1 °C on WISDOM) (**Fig. 4B**).

In the final mutated *Pb*INP construct in this series, all arginines in the R-coils were replaced by lysines (K-coils). This mutant nucleated ice formation at essentially the same temperature as the wild type (**Fig. 4b**) (p = 0.89), suggesting that positive charges in these locations are more important than side chain geometry.

Figure 4. Site-specific mutagenesis of noteworthy motifs in the R-coil subdomain. A) Diagram indicating the design of the constructs. Translucent bars indicate continuity of three conserved motifs along the length of the central repetitive domain. Neg. charge: Negative residues present in positions 11, 12, and 14 of the repeat. TxT: Thr-Xaa-Thr motif in positions 6-8 of the repeat. Y-ladder: An entirely conserved Tyr is position 3 of the motif. Three mutants were created in which these motifs were extended into the R-coils. K-coils: All arginines in the R-coils were replaced with lysines. **B)** Freezing curves with T_{50} and number of unfrozen droplets indicated where applicable.

Droplet freezing assays show recombinant cell lysate supernatant has ice nucleation activity that is affected by pH

The experiments described above were performed using whole recombinant bacteria rather than extracted INPs. In *E. coli*, the vast majority of the expressed INP is intracellular (27). Indeed, with our GFP-tagged constructs, we observe intense green fluorescence in the cytoplasm. To study the effects of pH on *Pb*INP activity, it was necessary to lyse the *E. coli* to change the pH surrounding the INP multimers. After centrifuging the sonicate to remove cell debris and passing the supernatant through a 0.2-µm filter to remove any unbroken cells, the extracts were tested to see how ice nucleation activity is affected by pH between 2.0 to 11.0. The activity of the filtered

supernatant was only a few degrees lower than that of whole bacteria (T_{50} = -9.6 °C) (**Fig. 5A**),

which agrees with the results of Kassmannhuber *et al.* (28). This indicates that large INP

structures are present within the bacterial cytoplasm.

The effect of pH on Snomax activity has been previously reported (16). However, Snomax is

comprised of freeze-dried *P. syringae* cells in which the INPs are thought to be membrane

bound. Our assays on bacterial lysate tested free, cytoplasmic *Pb*INP complexes, producing a

similar trend regarding the effect of pH but with somewhat greater loss of activity on the lower

end of the optimal range (**Fig. 5B**). Ice nucleation activity decreased by a few degrees below pH

5.0, and by ~8 °C at pH 2.0. The loss of activity in the alkaline buffers up to pH 11.0 was

minimal. Similar to the findings of Chao *et al.* (30), we did not observe a major change in

activity (i.e. ΔT_{50} > 10 °C) even at the extremes of pH 2.0 and 11.0, suggesting that the

mechanism of ice nucleation is not pH-dependent.

Figure 5. A) A comparison of the nucleation temperatures of *Pb*INP when assayed using intact *E. coli* cells and when assayed with filtered supernatant. **B)** A box and whisker plot showing the nucleation temperatures of filtered supernatant containing *Pb*INP under different pH conditions. Boxes and bars indicate quartiles, with medians indicated by a centre line. Outliers are indicated by diamonds.

INP activity is remarkably heat resistant

Having access to lysate also provided an opportunity to examine the heat stability of the INP complexes. The filtered lysate was heat-treated to 60, 70, 80, 90, or 99 °C for 10 min in sealed tubes before being chilled and assayed for ice nucleation activity (**Fig. 6A**). The activity of the 60 °C sample (T_{50} = -9.9 °C) was nearly identical to the non-treated wild-type control (T_{50} = -9.6 °C), and the 70 °C sample only displayed a minor loss of activity (T_{50} = -10.2 °C). From 80 °C to 99 °C the activity incrementally decreased (T_{50} = -11.3 °C, -12.7 °C, -14.6 °C, respectively), but the activity loss never exceeded 6 °C. Indeed, the heat resistance of the INP complex is remarkable. The C-terminal GFP tag provided an internal control for the effectiveness of heat treatment, as GFP denatures at around 73 °C (31). The green colour of the bacteria was robust at 65 °C and with very few exceptions, gone at 75 °C (**Fig. 6B**). There was no fluorescence at 90 °C.

We also assayed the lysate of a construct from our previous study in which repeats 16-47 (residues 411-923) of the solenoid had been deleted, leaving 32 coils (6). The overall freezing profile remained the same, indicating that this construct is also extremely resistant to heat denaturation, with each temperature sample freezing at slightly lower temperatures than their full-length counterparts (Fig. 6A). While the Δ411-923 construct had slightly lower overall activity, its heat resistance was not affected by the truncation.

Figure 6. A) Measured freezing temperatures of heat-treated droplets containing either *Pb*INP or *Pb*INP with the first 32 repeats (counting from the N-terminal end) deleted. Wild-type *Pb*INP freezing without heat treatment (kept at roughly 10 °C) is indicated- on the left. **B)** Fluorescent microscopy images of recombinant *E. coli* cells expressing *Pb*INP tagged with GFP viewed under bright-field (BF) or fluorescent excitatory (GFP) light. Representative images are shown (n = 3). Note: cells that retain their fluorescence after 75 °C treatment are rarely observed.

The β-solenoid of INPs is stabilized by a capping structure at the C terminus, but not at the N terminus

There is a clear C-terminal capping structure in the AlphaFold model (**Fig. 1A**), but a possible N-terminal cap was more nebulous. Most protein solenoids are N- and/or C-terminally capped to help maintain the fold and/or prevent end-to-end associations (32). Looking at the N-terminal sequence, we tested if any part of the extended linker region serves as an N-terminal capping motif. To investigate this, we made a series of incremental N-terminal deletions starting at residues Asp150 (Truncation 1), Gln159 (Truncation 2), and Gln175 (Truncation 3) (**Fig. 7A**). Truncation 1 lacked most of the N-terminal domain, leaving the last few residues of the unstructured linker. Truncation 2 removed those linker residues so that the putative cap (a single β-strand) was located at the very N-terminal end of the protein. Truncation 3 removed the β-strand along with the rest of the first coil of the solenoid. When tested, there was no difference between the activities of the three truncations and the wild type (p = 0.82 by one-way ANOVA) (**Fig. 7D**). This result is in line with those from Kassmannhuber *et al.* (28), which showed that deletion of the N-terminal domain does not significantly affect ice nucleation activity.

Previously, we demonstrated that the C-terminal cap is essential for ice nucleation activity (6). Bioinformatic analysis showed a high degree of conservation in the C-terminal cap residues (**Fig. 7B**). Rather than deleting the cap, we made targeted mutations: F1204D, D1208L, and Y1230D, to disrupt the structure predicted by AlphaFold. These mutations were chosen based on the putative role of the residues in the AlphaFold model. F1204 sits atop the final R-coil to cover its hydrophobic core. D1208 helps to maintain a tight loop through strategic hydrogen bonds, and Y1230 fills a gap in the surface of the cap (**Fig. 7C**). When comparing these selections to the aligned C-terminal cap sequences, we see that all three residues are highly conserved. The resulting triple mutant displayed greatly reduced activity (T_{50} = -27.8 °C).

Figure 7. A) Sites of N-terminal truncations to *Pb*INP, indicating the location of the starting residue in the shortened construct. **B)** Alignment of representative INP C-terminal domains from the genus *Pseudomonas*. Mutated residues and their one-letter codes are indicated above. Symbols at the bottom indicate consensus (* for fully conserved, : for conservation of strongly similar chemical properties, . for conservation of weakly similar chemical properties). **C)** Predicted location of mutated residues in the *Pb*INP C-terminal cap with sidechains shown and predicted H-bonds for D1208 shown as dashed lines. **D)** The ice nucleation curves for the N- and C-terminal cap mutants.

Cryo-electron tomography reveals INPs multimers form bundled fibres in recombinant cells

E. coli cells recombinantly overexpressing INPs were plunge-frozen and milled into ~150-nm thick lamella using cryo-FIB (**Fig. 8A**). Grids containing lamellae were transferred into either a 200- or a 300-kV transmission electron microscope for imaging under cryogenic conditions. Many *E. coli* cells were observed within the low-magnification cryo-TEM overview image of the lamella (**Fig. 8B**). Tilt series were collected near individual *E. coli* cells, and 3-D tomograms were reconstructed to reveal cellular and extracellular features. Strikingly, *E. coli* cells overproducing wild-type INPs appear to be lysed after 3 days of cold acclimation at 4 °C and contain clusters of fibres in the cytoplasm (**Fig. 8 C, D, E**, tomograms in **Movies S2** and **S3**). Individual fibres are up to a few hundred nanometers in length but only a few nanometers in width. Intriguingly, these fibre clusters were not observed in those *E. coli* that overexpress INP mutants lacking R-coils and the cell envelopes stay integral after being cold acclimated over the same period as those of wild-type INP-producing *E. coli*. (**Fig. S5 A, B, C, D**).

Figure 8. Fibrous bundles observed by cryo-FIB and cryo-ET in in *E. coli* cells expressing INP. A) Ion-beam image of a thin lamella containing *E. coli* cells expressing INP obtained from cryo-FIB milling. **B)** Zoomed-in view of a cryo-TEM image of the lamella in A). Boxes with dashed-lines indicate areas where tilt series were collected. **C) and D)** Snapshots from 3-D cryo-tomograms reconstructed from tilt series collected in the boxed regions in B) showing striking fibrous bundles (yellow arrowheads). The *E. coli* cell envelopes are indicated with thick dash-lines. **E)** Further examples of the fibrous bundles produced by INP-expressing *E. coli*. Size markers in A) is 10 µm, in B) is 2 µm and in C), D) and E) are 100 nm, respectively.

Discussion

Previously, we showed that the PbINP solenoid domain is made up of two subdomains: the larger N-terminal region of WO-coils accounting for 80-90% of the total length; and the smaller C-terminal R-coil region accounting for the remaining 10-20% (6). The length of the WO-coil region and the continuity of the water-organizing motifs were shown to directly affect ice nucleation temperature. Although the R-coil region lacks water-organizing motifs, its presence was critical for ice nucleation activity, which led us to propose a key role for this region in INP

multimer formation. Here we have characterized the R-coil subdomain in terms of the attributes it needs to support INP multimerization and have shown by cryo-ET the first *in situ* view of what these multimers look like. In addition, we have advanced a working model for the INP multimer structure that is compatible with all of the known INP properties.

Removal of the front or middle halves of the PbINP solenoid (leaving 33 coils) only dropped the ice nucleation activity by ~2 °C. In these experiments, the R-coils were not shortened, so these deletions caused a disproportional loss of WO-coils from 53 to 21. Here we have confirmed this tolerance of WO-coil count variation through bioinformatic analysis of natural INPs. The majority of bacterial INPs have WO-coil counts ranging from ~30 to 70. *Pb*INP is average in this respect with 53 WO-coils.

It seems counterintuitive that these bacteria have not been uniformly selected for the highest WO-coil count, which might give them an advantage in causing frost damage to plants at the highest possible temperature (33). However, it is clear that INPs are not functioning as monomers but rather as large multimers. Thus, any loss of length can potentially be compensated for by adding more monomers to the multimer. The ability to form these superstructures is a key property of the INPs and centers on the R-coil subdomain. This was shown in the same bioinformatic analysis where there is remarkably little variation to the R-coil length of 10-12 coils. The importance of a minimal R-coil length is supported here by experiments. Whereas over 50% of the WO-coils can be removed with minimal loss of activity, when six of the 12 *Pb*INP R-coils were replaced by WO-coils there was a catastrophic loss of ice nucleation activity, and no activity at all with further shortening of the R-coils. We postulate that at least eight R-coils are required for efficient multimer formation and that the ice nucleation activity of a monomer is inconsequential in the natural environment.

In the absence of detailed structural information, we have probed the properties of the multimers to help develop feasible models for their structure and assembly. The location of the R-coils at the C-terminal end of the solenoid next to the highly conserved cap structure is critical, as they do not function in the middle of the WO-coil region, and only poorly at the N-terminal end. These R-coils have a strong positive charge from the Arg and Lys residues, whereas the WO-coils are negatively charged, and their interaction potentially provides an electrostatic component to the fibre assembly. In this regard, the negative charges of the WO-coils are consistent

throughout their length, which offers no clue as to where on the WO-coils the R-coils might

interact. One possible advantage of this uniformity is that multimer assembly could still happen

if the WO-coil length is appreciably shortened.

As expected, changing the charge on the R-coils from positive to negative caused a large loss of

ice nucleation activity of ~9 °C, consistent with charge repulsion between these two solenoid

regions. The minimal effects of pH on native INP activity are reminiscent of the insensitivity of

antifreeze activity to pH (30, 34). The ice-binding sites of AFPs are typically devoid of charged

residues and there should be no effect of pH on the ability of these sites to organize ice-like

waters. The same can be said for the water-organizing motifs in INPs. Any effect of pH on ice-

binding or ice-nucleating activity is likely due to changes to the protein fold or INP multimer.

The small effect of pH on INP activity encouraged us to consider other possible interactions in

the assembly of multimers. We noted the extraordinary heat stability of INP multimers. Even

after heating to 99 °C for 10 min the bacterial extracts only lost 5 °C of ice nucleation activity,

whereas the heat-stable internal GFP control was denatured at 75 °C. We cannot rule out the

possibility that the INP multimers were also denatured by heat treatment but could reassemble on

cooling.

Figure 9. Filamentous multimer model for bacterial INPs. A) Antiparallel INP dimers are joined end to end by forming electrostatic interactions between negatively (red) and positively (blue) charged surfaces. The tyrosine ladder (yellow strip) is proposed as a dimerization interface. The C-terminal cap is shown in orange, Light green is the solenoid surface displaying the TxT WO-motifs and the darker green is the solenoid surface displaying the SxT motifs. The N and C termini of the INP solenoids are labeled N and C, respectively. **B)** The same dimerization interface in a parallel orientation. Highlighted in red is an apparent steric clash at the C-terminal end. **C)** Illustration of a possible assembly where the sheet in A is curled around to form a second set of electrostatic interactions to produce a fibre with an approximate diameter of 5-6 nm. **D**) Cross section looking down the axis of the fibre. **E**) A version of the model where 32 WO-coils have been removed.

Working model of the INP multimer.

The fundamental unit of the INP multimer in this hypothetical model is a dimer (**Fig. 9**). The dimerization interface involves an interaction of the stacked tyrosine ladders (coloured yellow in **Fig. 9A**) from the two INP monomers as previously suggested (8, 22). However, in this model the INPs are aligned antiparallel to each other (**Fig. 9B**). This orientation is more likely than a parallel alignment since the R-coils and C-terminal cap structure appear to clash when modelled (**Movie S4**). The antiparallel dimer would not be a rigid flat sheet but could hinge at the tyrosine ladder. Another advantage of the antiparallel arrangement is that the two dimer termini are identical allowing end-to-end linking to form a long fibre.

The end-to-end dimer associations involve electrostatic interactions between the basic side of the R-coils and the acidic side of the WO-coils (**Fig. 9A**). If these interactions also form a hinge, it

should be possible to roll the end-linked dimers into a compact fibre (**Fig. 9C**) with a diameter

close to that seen by cryo-ET (**Fig. 8**). At the dimer ends where the R-coils form an electrostatic

interaction with the far end of the WO-coils there are two sets of interacting surfaces involved in

making the rolled up fibre. The antiparallel arrangement of the dimers gives a sidedness to the

multimer where TxT motifs (light green) face outwards and inwards in an alternating pattern

with SxT motifs (dark green) in the opposite phase. A cross-sectional view of the INP fibre (**Fig.**

9D) shows the alternating water-organizing surfaces, the interactions between the negatively

(red) and positively (blue) charged regions, and the openness of the structure to organize ice-like

water molecules inside and outside of the fibre.

In addition, the additional stabilizing interfaces offered by this assembly would further stabilize

the multimer into a rigid fiber and also help to explain the apparent heat resistance of INP

samples.

Working model of the INP multimer is consistent with the properties of INPs and their multimers.

We previously showed that the length of the WO-coil region can be shortened by ~60% with

only a few °C decrease in ice nucleation temperature (6). The working model can accommodate

these huge deletions simply by closing the gap between the dimers. For example, the deletion of

32 WO-coils, leaving just 21 and the 12 R-coils (**Fig. 9E**) retains all the molecular interactions

seen in the longer fibre but with fewer stacked tyrosine interactions. This can help explain the

heat stability of the INP multimers and the minimal difference (2-3 °C) in activity loss between

full-length *Pb*INP with 65 coils and the truncated version with 33 coils (**Fig. 6**). Similarly, longer

WO-coil regions can be accommodated by lengthening the gap. This can explain the wide range

of WO-coil lengths seen in nature (**Fig. 2**). They all fit in the same model.

Our model also shows how the interaction between the R-coils and the WO-coils of the adjacent

dimers supports fibre formation. Any shortening of the R-coil subdomain jeopardizes the ability

to link up the dimers. The catastrophic loss of ice nucleation activity seen below 8 R-coils is

because the interacting length of R-coils and WO-coils has too few electrostatic and other

interactions to bridge the dimers together. The importance of electrostatic interaction has been

illustrated in this study in two ways. When the R-coil basic residues were replaced by acidic

residues, the ice nucleation activity was severely compromised but was fully restored when the mutated residues were all converted to lysines. Additionally, in cell-free extracts of lysed INP-producing *E. coli* ice nucleation activity decreased by a few degrees Celsius at low pH values where the charge on acidic residue side chains was reduced or eliminated. When the carboxyl groups of aspartate and glutamate involved in electrostatic pairing lose their negative charges at low pH they can still form hydrogen bonds with basic amino acid partners, which can explain why the lowering of pH was not as disruptive as reversing the charge on these residues.

The observation that low, variable levels of ice nucleation activity remained in the construct where the R-coil basic residues were replaced by acidic residues, suggests that there are additional binding interactions between the dimers other than electrostatic ones. Here we suggest the involvement of the highly conserved C-terminal capping structure. When three mutations designed to disrupt the cap fold were introduced all ice nucleation activity was lost. Also of note is the disruptive effect of extending the tyrosine ladder further into the R-coil sub-domain in the mutant where the acidic residues replaced the basic one. The subtle details of the R-coil region will require detailed structural analysis for their elucidation.

The relocation of the R-coils to the N-terminal end of the solenoid caused a loss of just over 50% activity and it is possible to accommodate such a change in the model while retaining a charge interaction between the R-coils and WO-coils. However, the separation from the cap structure might account for some of the activity loss. Movement of the R-coils to the centre of the WO-coil region is not compatible with the model and sure enough this construct was devoid of ice nucleation activity (**Fig. 3B**).

Other features supporting the model are that the dimer's C- and N-terminal ends are exposed and can accommodate tags and extensions without disrupting the fibre. Thus, the addition of a C-terminal GFP tag has no detrimental effect in ice nucleation activity. Nor is there any difference in activity if the N-terminal INP domain and linker region are present or not (**Fig. 7**) (ref). Even the incorporation of bulky mRuby into the WO-coils (6) can be accommodated because the fibre is too thin for the solenoids to be packed together in a bundle.

Electron microscopy of newly synthesized INPs in a cell-free system shows them as thin molecules of dimensions 4-6 nm in diameter by a few hundred nm in length (23). Negatively

stained images of recombinantly produced INP multimers isolated by centrifugation and

chromatography show an elongated structure ~5-7 times longer than a monomer but not much

wider (22). The fibres seen *in situ* in INP-expressing *E. coli* (**Fig. 8**) are similarly long but

slightly thinner, consistent with the absence of negative staining. The model in **Fig. 9** is the

thinnest structure we can project for a fibrillar multimer.

Solving the structure of the INP fibres at atomic detail will be the key to understanding the

remarkable ability of biological ice nucleators to start the freezing process at high sub-zero

temperatures.

Methods

AlphaFold prediction

442 The AlphaFold model for *Pb*INP was generated by Forbes *et al.* as described (6).

Bioinformatic analysis of INPs

NCBI's BLAST was accessed using the BioPython library v1.81 (35). The consensus sequence

for the 16-residue coil 'AGYGSTQTAGEDSSLT' was used as the query agsinst the non-

redundant (nr) database. The PAM30 scoring matrix was used due to the short query.

Quality control (QC) was performed using custom Python scripts, making use of BioPython's

Entrez module to fetch information on the protein, BioProject, and assembly method for each

BLAST result (**Fig. S1**). Custom Python scripts were used to automatically identify the tandem

repeats and classify them as WO-coils or R-coils.

Sequence logos were made using the Logomaker package v0.8 (36). Alignment of C-terminal

cap sequences was performed using JalView software v2.11.2.6 (37).

Synthesis of *Pb*INP genes

456 Experiments for this project used a synthetic *Pb*INP gene previously developed by our group.

457 This codon-optimized gene encodes the *P. borealis* INP gene (GenBank accession: EU573998).

458 Additionally, the DNA sequence for enhanced green fluorescent protein (eGFP) (GenBank

459 accession: AAB02572) was fused to the 3′-end of the *Pb*INP gene using a hexanucleotide

encoding two linker residues (Asn-Ser). More details about the *Pb*INP-eGFP sequence are

provided in Forbes *et al*. (6).

All mutants for this study were designed by modifying the aforementioned synthetic gene.

GenScript performed all (Piscataway, NJ, USA) gene syntheses, which we subsequently cloned

into the pET-24a expression vector.

Five *Pb*INP mutants were designed to test the effect of replacing the R-coils. The R-coils were

incrementally replaced with the sequences of WO-coils adjacent to the R-coil region as

indicated, resulting in constructs containing 10, 8, 6, 4, and 1 R-coil (**Fig. 2A**). Replacements

were designed such that they maintained the periodicity of the tandem repeats. The C-terminal

R-coil was left untouched to avoid disturbing possible interactions with the putative C-terminal

cap structure.

The R-coils were either relocated within the protein or deleted (**Fig. 3A**), while again leaving the

N- and C-terminal coils untouched to avoid interactions with adjacent domains.

Targeted mutations were introduced to the R-coil region gene to produce four additional

constructs (**Fig. 4A**). For the first construct (RKH replacement), any positively charged residues

(Arg, Lys, His) in positions 11, 12, or 14 in the R-coils were replaced with residues commonly

found in those locations in the WO-coils (Asp, Glu, and Gly for positions 11 and 12, Ser for

position 14). The second construct Y extend in **Fig. 4A** extends the stacked tyrosine ladder

present at position 3 of the coils through 7 additional coils toward the C terminus of the solenoid.

The third construct (RKH replacement + Y extend in **Fig. 4A** is a combination of both mutants.

The fourth construct (K-coils in **Fig. 4A**) converted every Arg residue in the R-coil section to a

Lys residue.

Protein expression in *E. coli*

Each *Pb*INP construct was transformed into the ArcticExpress strain of *E. coli*, since its

expression of two cold-adapted chaperones, Cpn10 and Cpn60, promotes the correct folding of

proteins at low temperatures (38). Transformation and induction with IPTG were performed

according to the supplier's instructions (Agilent Technologies, Catalog #230192). Cells

expressed at 10 °C for 24 h post-induction. The eGFP tag allowed expression to be confirmed

using fluorescence microscopy (6).

Ice nucleation assays by WISDOM

Constructs were assayed on WISDOM (WeIzmann Supercooled Droplets Observation on a

Microarray) (39) in a similar way as described in Forbes *et al.* (6).

Ice nucleation assays by nanoliter osmometer

Ice nucleation activity was quantified using a droplet freezing assay protocol (Lee *et al*.,

submitted) that makes use of a LabVIEW-operated nanoliter osmometer (Micro-Ice, Israel) (40).

Briefly: Following induction and cold incubation, nanoliter-sized droplets of liquid cultures were

pipetted into oil-filled wells resting on a cold stage. The temperature of the cold stage was

lowered at a rate of 1 °C per minute while a video recording was taken of the sample grid.

Freezing was characterized by a distinct change in droplet appearance. After assay completion,

the videos were analyzed to record the temperatures of all freezing events. The fraction of frozen

droplets (f_{ice}) as a function of temperature was plotted, generating ice nucleation curves for each

sample. This apparatus could not reach temperatures as low as those achieved on WISDOM, but

results are in agreement between the two approaches (**Fig. 2B**).

Heat treatment and pH

To obtain cell lysates, *E. coli* cultures were centrifuged at $3,200 \times g$ for 30 min post-induction.

Cell pellets were then resuspended in a lysis buffer of 50 mM Tris-HCl, 150 mM NaCl,

containing Pierce Protease Inhibitor (Thermo Scientific, Canada) before sonication at 70%

amplitude for 30-s rounds. Lysate was centrifuged at $31,000 \times g$ and the resulting supernatant

was passed through a 0.2 μm filter.

For heat treatment, filtered lysate in sealed Eppendorf tubes was heated at 60 °C, 70 °C, 80 °C,

90 °C, or 99 °C for 10 min in a thermocycler and then quenched on ice prior to being assayed for

activity.

For the pH experiments, aliquots of filtered lysate were diluted 50-fold in pH-adjusted buffer of

100 mM sodium citrate, 100 mM sodium phosphate, and 100 mM sodium borate following the

protocol by Chao *et al*. (30). Before assaying, we verified using universal indicator strips that

addition of lysate to the buffer mixtures did not meaningfully affect the pH of the final mixtures.

Preparation of the cryo-EM grids

After confirming eGFP-INP expression, the *E. coli* cultures were incubated at 4 $°C$ for an

additional 3 days. The *E. coli* cells were spun down and resuspended in PBS to an $OD_{600\ nm}$ of ~ 3.

These concentrated *E. coli* samples were deposited onto freshly glow-discharged QUANTIFOIL

holey carbon grids (Electron Microscopy Sciences). The grids were then blotted from the back

side with the filter paper for ~5 s before plunge-frozen in liquid ethane, using a manual plunger-

freezing apparatus as described previously (41, 42).

Cryo-FIB milling

The plunge-frozen grids with *E. coli* cells were clipped into cryo-FIB AutoGrids and mounted into

the specimen shuttle under liquid nitrogen. An Aquilos2 cryo-FIB system (Thermo Fisher

Scientific) was used to mill the thick bacterial samples into lamellae of < 200 nm in thickness. The

milling process was completed using a protocol as previously described (43).

Cryo-ET data acquisition and tomogram reconstruction

Grids containing the lamellae obtained from cryo-FIB milling were loaded into either a 300-kV

Titan Krios electron microscope (Thermo Fisher Scientific) equipped with a Direct Electron

Detector and energy filter (Gatan) or a 200-kV Glacios Electron Microscope at Yale University.

The FastTOMO script was used with the SerialEM software to collect tilt series with defocus

values of approximately −6 μm (44), and a cumulative dose of ~70 e^-/Å covering angles from −48°

to 48° (3° tilt step). Images were acquired at 42,000 × magnification with an effective pixel size

of 2.148 Å. All recorded images were first drift corrected by MotionCor2 (45), stacked by the

software package IMOD (46), and then aligned by IMOD using Pt particles as fiducial markers.

TOMO3D was used to generate tomograms by simultaneous iterative reconstruction technique

(SIRT) (47). In total, 10 tomograms were reconstructed with TOMO3D for the WT INP while 5

tomograms were produced for the R-coil mutant.

Contributions

T.H., J.C.L., and P.L.D. planned these experiments based on the identification of the R-coil subdomain by T.H. and P.L.D. in a prior work. T.H., J.C.L., and P.L.D. wrote the original manuscript for peer review. T.H. and P.L.D. designed the model for multimerization. T.H. coded the pipeline and performed the bioinformatic analysis of INPs.

T.H. and J.C.L. designed all mutants and prepared all samples except where otherwise noted.

J.C.L. prepared the buffers and samples for the pH experiments, and performed all measurements of ice nucleation activity on the nanoliter osmometer. T.H. and J.C.L. performed the heat treatment experiments.

G.O. and I.B. prepared samples whose activity was assayed by N.R. and Y.R. on WISDOM.

S.G., W.G. and J.L. performed cryo-FIB and cryo-ET experiments. S.G. prepared Fig. 8 and Movies S2 and S3. T.H. prepared all other figures.

Acknowledgements

This work was supported by CIHR Foundation Grant FRN 148422 to P.L.D., who holds the Canada Research Chair in Protein Engineering, and by an Israel Science Foundation grant to I.B. YR acknowledges support by a research grant from the Yotam project and the Weizmann institute sustainability and energy research initiative. S.G. was supported by a CIHR Post-Doctoral Fellowship and an NIH RO1 grant (R01AI087946) of J.L. W.G. was also supported by the NIH RO1 grant (R01AI087946) of J.L. We thank Virginia K. Walker for the gift of the *Pseudomonas borealis* strain.

References

1. Hoose, C., and Möhler, O. (2012) Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments Atmospheric Chemistry and Physics **12**, 9817-9854 http://dx.doi.org/10.5194/acp-12-9817-2012

2. Vali, G., DeMott, P. J., Möhler, O., and Whale, T. F. (2015) Technical Note: A proposal for ice nucleation terminology Atmospheric Chemistry and Physics **15**, 10263-10270 http://dx.doi.org/10.5194/acp-15-10263-2015

3. Lukas, M., Schwidetzky, R., Eufemio, R. J., Bonn, M., and Meister, K. (2022) Toward Understanding Bacterial Ice Nucleation J Phys Chem B **126**, 1861-1867 http://dx.doi.org/10.1021/acs.jpcb.1c09342

4. Govindarajan, A. G., and Lindow, S. E. (1988) Size of bacterial ice-nucleation sites measured in situ by radiation inactivation analysis Proc Natl Acad Sci U S A **85**, 1334-1338 http://dx.doi.org/10.1073/pnas.85.5.1334

5. Lindow, S. E., Lahue, E., Govindarajan, A. G., Panopoulos, N. J., and Gies, D. (1989) Localization of ice nucleation activity and the iceC gene product in Pseudomonas syringae and Escherichia coli Mol Plant Microbe Interact **2**, 262-272 http://dx.doi.org/10.1094/mpmi-2-262

6. Forbes, J., Bissoyi, A., Eickhoff, L., Reicher, N., Hansen, T., Bon, C. G. *et al.* (2022) Water-organizing motif continuity is critical for potent ice nucleation protein activity Nat Commun **13**, 5019 http://dx.doi.org/10.1038/s41467-022-32469-9

7. Graether, S. P., and Jia, Z. (2001) Modeling Pseudomonas syringae ice-nucleation protein as a beta-helical protein Biophys J **80**, 1169-1173 http://dx.doi.org/10.1016/S0006-3495(01)76093-6

8. Garnham, C. P., Campbell, R. L., Walker, V. K., and Davies, P. L. (2011) Novel dimeric beta-helical model of an ice nucleation protein with bridged active sites BMC Struct Biol **11**, 36 http://dx.doi.org/10.1186/1472-6807-11-36

9. Graether, S. P., Kuiper, M. J., Gagne, S. M., Walker, V. K., Jia, Z., Sykes, B. D. *et al.* (2000) Beta-helix structure and ice-binding properties of a hyperactive antifreeze protein from an insect Nature **406**, 325-328 http://dx.doi.org/10.1038/35018610

10. Graham, L. A., Qin, W., Lougheed, S. C., Davies, P. L., and Walker, V. K. (2007) Evolution of hyperactive, repetitive antifreeze proteins in beetles J Mol Evol **64**, 387-398 http://dx.doi.org/10.1007/s00239-005-0256-3

11. Kristiansen, E., Ramlov, H., Hojrup, P., Pedersen, S. A., Hagen, L., and Zachariassen, K. E. (2011) Structural characteristics of a novel antifreeze protein from the longhorn beetle Rhagium inquisitor Insect Biochem Mol Biol **41**, 109-117 http://dx.doi.org/10.1016/j.ibmb.2010.11.002

12. Garnham, C. P., Campbell, R. L., and Davies, P. L. (2011) Anchored clathrate waters bind antifreeze proteins to ice Proc Natl Acad Sci U S A **108**, 7363-7367 http://dx.doi.org/10.1073/pnas.1100429108

13. Green, R. L., and Warren, G. J. (1985) Physical and functional repetition in a bacterial ice nucleation gene Nature **317**, 645-648 http://dx.doi.org/10.1038/317645a0

14. Kajava, A. V., and Steven, A. C. (2006) The turn of the screw: variations of the abundant beta-solenoid motif in passenger domains of Type V secretory proteins J Struct Biol **155**, 306-315 http://dx.doi.org/10.1016/j.jsb.2006.01.015

15. Madzharova, F., Bregnhoj, M., Chatterley, A. S., Lovschall, K. B., Drace, T., Andersen Dreyer, L. S. *et al.* (2022) Electrostatics Trigger Interfacial Self-Assembly of Bacterial Ice Nucleators Biomacromolecules **23**, 505-512 http://dx.doi.org/10.1021/acs.biomac.1c01217

16. Lukas, M., Schwidetzky, R., Kunert, A. T., Poschl, U., Frohlich-Nowoisky, J., Bonn, M. *et al.* (2020) Electrostatic Interactions Control the Functionality of Bacterial Ice Nucleators J Am Chem Soc **142**, 6842-6846 http://dx.doi.org/10.1021/jacs.9b13069

17. Schwidetzky, R., Lukas, M., YazdanYar, A., Kunert, A. T., Poschl, U., Domke, K. F. *et al.* (2021) Specific Ion-Protein Interactions Influence Bacterial Ice Nucleation Chemistry **27**, 7402-7407 http://dx.doi.org/10.1002/chem.202004630

18. Juurakko, C. L., diCenzo, G. C., and Walker, V. K. (2022) Brachypodium Antifreeze Protein Gene Products Inhibit Ice Recrystallisation, Attenuate Ice Nucleation, and Reduce Immune Response Plants (Basel) **11**, http://dx.doi.org/10.3390/plants11111475

19. Qiu, Y., Hudait, A., and Molinero, V. (2019) How Size and Aggregation of Ice-Binding Proteins Control Their Ice Nucleation Efficiency J Am Chem Soc **141**, 7439-7452 http://dx.doi.org/10.1021/jacs.9b01854

20. Roeters, S. J., Golbek, T. W., Bregnhoj, M., Drace, T., Alamdari, S., Roseboom, W. *et al.* (2021) Ice-nucleating proteins are activated by low temperatures to control the structure of interfacial water Nat Commun **12**, 1183 http://dx.doi.org/10.1038/s41467-021-21349-3

21. Schmid, D., Pridmore, D., Capitani, G., Battistutta, R., Neeser, J.-R., and Jann, A. (1997) Molecular organisation of the ice nucleation protein InaV from Pseudomonas syringae FEBS Letters **414**, 590-594 http://dx.doi.org/10.1016/s0014-5793(97)01079-x

22. Hartmann, S., Ling, M., Dreyer, L. S. A., Zipori, A., Finster, K., Grawe, S. *et al.* (2022) Structure and Protein-Protein Interactions of Ice Nucleation Proteins Drive Their Activity Front Microbiol **13**, 872306 http://dx.doi.org/10.3389/fmicb.2022.872306

23. Novikova, I. V., China, S., and Evans, J. E. (2018) Overcoming bottlenecks for in vitro synthesis and initial structural insight of ice nucleating protein InaZ bioRxiv 334987 http://dx.doi.org/10.1101/334987

24. Li, Q., Yan, Q., Chen, J., He, Y., Wang, J., Zhang, H. *et al.* (2012) Molecular characterization of an ice nucleation protein variant (inaQ) from Pseudomonas syringae and the analysis of its transmembrane transport activity in Escherichia coli Int J Biol Sci **8**, 1097-1108 http://dx.doi.org/10.7150/ijbs.4524

25. Li, L., Kang, D. G., and Cha, H. J. (2004) Functional display of foreign protein on surface of Escherichia coli using N-terminal domain of ice nucleation protein Biotechnol Bioeng **85**, 214-221 http://dx.doi.org/10.1002/bit.10892

26. Nicchi, S., Giuliani, M., Giusti, F., Pancotto, L., Maione, D., Delany, I. *et al.* (2021) Decorating the surface of Escherichia coli with bacterial lipoproteins: a comparative analysis of different display systems Microb Cell Fact **20**, 33 http://dx.doi.org/10.1186/s12934-021-01528-z

27. Kassmannhuber, J., Rauscher, M., Schoner, L., Witte, A., and Lubitz, W. (2017) Functional display of ice nucleation protein InaZ on the surface of bacterial ghosts Bioengineered **8**, 488-500 http://dx.doi.org/10.1080/21655979.2017.1284712

28. Kassmannhuber, J., Mauri, S., Rauscher, M., Brait, N., Schoner, L., Witte, A. *et al.* (2020) Freezing from the inside: Ice nucleation in Escherichia coli and Escherichia coli ghosts by inner membrane bound ice nucleation protein InaZ Biointerphases **15**, 031003 http://dx.doi.org/10.1116/1.5142174

29. Wick, R. R., Judd, L. M., Gorrie, C. L., and Holt, K. E. (2017) Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads PLoS Comput Biol **13**, e1005595 http://dx.doi.org/10.1371/journal.pcbi.1005595

30. Chao, H., Sonnichsen, F. D., DeLuca, C. I., Sykes, B. D., and Davies, P. L. (1994) Structure-function relationship in the globular type III antifreeze protein: identification of a cluster of surface residues required for binding to ice Protein Sci **3**, 1760-1769 http://dx.doi.org/10.1002/pro.5560031016

31.   Melnik, T., Povarnitsyna, T., Solonenko, H., and Melnik, B. (2011) Studies of irreversible heat
denaturation of green fluorescent protein by differential scanning microcalorimetry
Thermochimica Acta **512**, 71-75 http://dx.doi.org/10.1016/j.tca.2010.09.002

32.   Kajava, A. V., and Steven, A. C. (2006) Beta-rolls, beta-helices, and other beta-solenoid proteins
Adv Protein Chem **73**, 55-96 http://dx.doi.org/10.1016/S0065-3233(06)73003-0

33.   Lindow, S. E., Arny, D. C., and Upper, C. D. (1982) Bacterial ice nucleation: a factor in frost injury
to plants Plant Physiol **70**, 1084-1089 http://dx.doi.org/10.1104/pp.70.4.1084

34.   Liu, K., Wang, C., Ma, J., Shi, G., Yao, X., Fang, H. *et al.* (2016) Janus effect of antifreeze proteins
on ice nucleation Proc Natl Acad Sci U S A **113**, 14739-14744
http://dx.doi.org/10.1073/pnas.1614379114

35.   Cock, P. J., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A. *et al.* (2009) Biopython:
freely available Python tools for computational molecular biology and bioinformatics
Bioinformatics **25**, 1422-1423 http://dx.doi.org/10.1093/bioinformatics/btp163

36.   Tareen, A., and Kinney, J. B. (2019) Logomaker: Beautiful sequence logos in python bioRxiv
635029 http://dx.doi.org/10.1101/635029

37.   Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M., and Barton, G. J. (2009) Jalview
Version 2--a multiple sequence alignment editor and analysis workbench Bioinformatics **25**,
1189-1191 http://dx.doi.org/10.1093/bioinformatics/btp033

38.   Belval, L., Marquette, A., Mestre, P., Piron, M. C., Demangeat, G., Merdinoglu, D. *et al.* (2015) A
fast and simple method to eliminate Cpn60 from functional recombinant proteins produced by
E. coli Arctic Express Protein Expr Purif **109**, 29-34 http://dx.doi.org/10.1016/j.pep.2015.01.009

39.   Reicher, N., Segev, L., and Rudich, Y. (2018) The Weizmann Supercooled Droplets Observation
on a Microarray (WISDOM) and application for ambient dust Atmospheric Measurement
Techniques **11**, 233-248 http://dx.doi.org/10.5194/amt-11-233-2018

40.   Braslavsky, I., and Drori, R. (2013) LabVIEW-operated novel nanoliter osmometer for ice binding
protein investigations J Vis Exp e4189 http://dx.doi.org/10.3791/4189

41.   Liu, J., Lin, T., Botkin, D. J., McCrum, E., Winkler, H., and Norris, S. J. (2009) Intact flagellar motor
of Borrelia burgdorferi revealed by cryo-electron tomography: evidence for stator ring curvature
and rotor/C-ring assembly flexion J Bacteriol **191**, 5026-5036
http://dx.doi.org/10.1128/JB.00340-09

42.   Zhao, X., Zhang, K., Boquoi, T., Hu, B., Motaleb, M. A., Miller, K. A. *et al.* (2013) Cryoelectron
tomography reveals the sequential assembly of bacterial flagella in Borrelia burgdorferi Proc
Natl Acad Sci U S A **110**, 14390-14395 http://dx.doi.org/10.1073/pnas.1308306110

43.   Xiang, Y., Surovtsev, I. V., Chang, Y., Govers, S. K., Parry, B. R., Liu, J. *et al.* (2021) Interconnecting
solvent quality, transcription, and chromosome folding in Escherichia coli Cell **184**, 3626-3642
e3614 http://dx.doi.org/10.1016/j.cell.2021.05.037

44.   Xu, A., and Xu, C. (2021) FastTomo: A SerialEM Script for Collecting Electron Tomography Data
bioRxiv 2021.2003.2016.435675 http://dx.doi.org/10.1101/2021.03.16.435675

45.   Zheng, S. Q., Palovcak, E., Armache, J. P., Verba, K. A., Cheng, Y., and Agard, D. A. (2017)
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy Nat Methods **14**, 331-332 http://dx.doi.org/10.1038/nmeth.4193

46.   Kremer, J. R., Mastronarde, D. N., and McIntosh, J. R. (1996) Computer visualization of three-
dimensional image data using IMOD J Struct Biol **116**, 71-76
http://dx.doi.org/10.1006/jsbi.1996.0013

47.   Agulleiro, J. I., and Fernandez, J. J. (2015) Tomo3D 2.0--exploitation of advanced vector
extensions (AVX) for 3D reconstruction J Struct Biol **189**, 147-152
http://dx.doi.org/10.1016/j.jsb.2014.11.009

Supplemental information

Figure S1. Flowchart and quality control steps in sequence selection for bioinformatic analysis. Ten known INPs from literature were used to generate a consensus sequence for WO-coils which was then used as a query in a BLAST against the *nr* database to identify INPs. NCBI E-Utils were used to generate a data set using only genes from long-read DNA sequencing data.

Figure S5. *E. coli* **expressing INP mutant lacking R-coils shows no fibre clusters as observed in those cells overexpressing WT INP. A-D**) Representative snapshots from 3-D cryo-tomograms showing cytoplasmic and extracellular features of various *E. coli* cells overexpressing an INP mutant in which all but the C-terminal R-coil have been replaced by WO-coils. All four images are in the same scale and the scale bar represents 100 nm.

713